

Dear reviewer and editors,

The authors appreciate the helpful comments from the reviewer very much, which help to improve the manuscript. According to these comments, the manuscript has been thoroughly revised. The responses to the reviewer's comments are listed below.

“Page 2, Line 5: What kind of ‘climate change’ are you talking about? “

The previous wording of “climate change” has been revised to “The global climate changed induced by anthropogenic activities”

*“Page 4, Lines 5-6: “The marine coccolithophore *Emiliana huxleyi* (morphotype A, strain NIWA1108) was isolated from the Chatham Rise, east of New Zealand by Dr. H. Chang in 2009, as detailed in Feng et al. (2017).*

*I have checked Feng et al. (2017) and Feng (2014) (PhD thesis of the corresponding author of this manuscript that shows original study of this manuscript), however, could not find detailed information of the strain in the papers. Please refer an appropriate paper that describes detailed information of the strain, or provide following information in this manuscript instead; sampling locality (latitude and longitude), sampling method, depth of seawater sample that yielded NIWA1108, hydrographic condition (temperature, salinity, etc.) of seawater at the sampling time, date of sampling, isolation method, and information of sampling cruise and cruise report. There is a possibility that culture strains originate from different hydrographic condition have different environmental preferences, as authors mention “we can speculate that the diverse *E. huxleyi* strains growing in different temperature regions might have different requirements for nitrogen” in the page 14. I think the detailed information of origin culture strain will help discussion of this study, and comparison of results from this study with result of other studies.”*

The detailed information of the *Emiliana huxleyi* strain NIWA 1108 has been added in the revised manuscript as “The marine coccolithophore *Emiliana huxleyi* (morphotype A, strain NIWA1108) was isolated from the surface water (depth of 5-6 m, salinity of 34.78) at 41°35.8'S 175°41.5'E, east of New Zealand by Dr. H. Chang aboard the *RV Tangaroa* on the research voyage TAN0909 in November 2009.

The water temperature was 12.1°C at the sampling site”.

“Page 4, Line 14; Please describe how did you know the culture strain is in exponential growth phase.”

The text “determined by the growth curve” has been added in order to clarify how the exponential growth phase was determined.

“Page 5, Line 18: How long did you keep your subsample in dark 4°C condition before counting?”

The text has been revised to “...and stored in dark at 4°C for no more than 5 days before counting”.

“Page 11, Line 15: Put ‘*E. huxleyi*’ in italics.”

The font has been corrected.

“Page 13, Lines 14-16: “In our 15 accompanying study, the growth, photosynthetic, and calcification rates all increased with rising temperature until the optimal temperature was reached (Feng et al., 2017)”;

Please mention the optimal temperature of the strain NIWA 1108 reported by Feng et al. (2017), and then discuss whether the optimal temperature of the strain in laboratory experiment is same/close to the temperature of sampling locality.”

The optimal temperatures for growth, photosynthetic and calcification rates are now included and discussed in the revised manuscript, as “...the growth, photosynthetic, and calcification rates all increased with rising temperature until the optimal temperature was reached at 25°C, 24°C, and 20°C respectively (Feng et al., 2017), which were all higher than the stock culture growth temperature or the temperature at the isolation site of *E. huxleyi* strain NIWA 1108”.

Thank you again for your help in the review process of the manuscript. Look forward to hearing back from you soon.

Sincerely yours,

Yuanyuan Feng and the co-authors

Environmental controls on the elemental composition of a Southern Hemisphere strain of the coccolithophore *Emiliana huxleyi*

Yuanyuan Feng^{1, 2, 3, 4}, Michael Y. Roleda^{2, 5}, Evelyn Armstrong⁶, Cliff S. Law^{6, 7}, Philip W. Boyd^{6, 7, 8}, Catriona L. Hurd^{2, 8}

5 ¹ College of Marine and Environmental Sciences, Tianjin University of Science and Technology, Tianjin 300457, China

² Department of Botany, University of Otago, PO Box 56, Dunedin 9054, New Zealand

³ Tianjin Key Laboratory of Marine Resources and Chemistry, Tianjin 300457, China

10 ⁴ Tianjin Marine Environmental Protection and Restoration Technology Engineering Center, Tianjin 300457, China

⁵ Norwegian Institute of Bioeconomy Research, Bodø 8027, Norway

⁶ NIWA/University of Otago Research Centre for Oceanography, Department of Chemistry, University of Otago, PO Box 56, Dunedin 9054, New Zealand

⁷ NIWA, Greta Point, Wellington 6023, New Zealand

15 ⁸ Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, 7005, Tasmania, Australia

Correspondence to: Yuanyuan Feng (yfeng@tust.edu.cn)

Abstract. A series of semi-continuous incubation experiments were conducted with the coccolithophore *Emiliana huxleyi* strain NIWA1108 (Southern Ocean isolate) to examine the effects of five environmental drivers (nitrate concentration, phosphate concentration, irradiance, temperature, and $p\text{CO}_2$) on both the physiological rates and elemental composition of the coccolithophore. Here, we report the alteration of the elemental composition of *E. huxleyi* in response to the changes in these environmental drivers. A series of dose response curves for the cellular elemental composition of *E. huxleyi* were fitted for each of the five drivers across an environmentally-representative gradient. The importance of each driver in regulating the elemental composition of *E. huxleyi* was ranked using a semi-quantitative approach. The percentage variations in elemental composition arising from the change in each driver between present day and model-projected conditions for the year 2100 were calculated. Temperature was the most important driver controlling both cellular particulate organic and inorganic carbon content, whereas nutrient concentrations were the most important regulator of cellular particulate nitrogen and phosphorus of *E. huxleyi*. In contrast, elevated $p\text{CO}_2$ had the greatest influence on cellular particulate inorganic carbon to organic carbon ratio, resulting in a decrease in the ratio. Our results

20
25
30

indicate that the different environmental drivers play specific roles in regulating the elemental composition of *E. huxleyi* with wide-reaching implications for coccolithophore-related marine biogeochemical cycles, as a consequence of the regulation of *E. huxleyi* physiological processes.

1 Introduction

5 The global climate change induced by anthropogenic activities is causing a wide range of alterations to the marine environment including ocean acidification (OA), rising sea surface temperature (SST), and intensified stratification due to increased density gradients between surface and subsurface waters, with associated shifts in mean irradiance levels and nutrient availability in the upper water column (Boyd and Doney, 2002; Rost and Riebesell, 2004; Stocker, 2013). All these global changes in environmental variables will affect the physiology and ecology of phytoplankton, both individually and interactively, in a complex way (Boyd and Hutchins, 2012; Boyd et al., 2010; Boyd et al., 2016; Feng et al., 2017).

15 Phytoplankton elemental composition is an important cellular property that reflects the metabolic rates of phytoplankton (Raven and Geider, 1988). Elemental composition is strongly influenced by environmental conditions and by phytoplankton adaptations to these conditions (Sterner and Elser, 2002) which in turn influences marine foodweb structure, particulate carbon export to the deep ocean, and ultimately marine biogeochemistry (Finkel et al., 2010 and references therein). The widely-recognized average molar elemental ratio of C:N:P is 106:16:1 for marine phytoplankton assemblages - the Redfield ratio (Redfield et al., 1963). However, individual phytoplankton species may have elemental ratios deviating, on short timescales (days to months) from Redfield depending on the environmental conditions they encounter. Such deviations subsequently influence the accumulation of these elements in the upper food web and also marine biogeochemistry (Finkel et al., 2010; Ho et al., 2003; Sardans et al., 2012).

25 Different environmental drivers may play a range of roles in regulating the stoichiometry of marine phytoplankton. Nutrient availability (Hecky et al., 1993; Perry, 1976) has been proven to affect phytoplankton stoichiometry directly. Irradiance provides the energy source for nutrient assimilation in the cells (Goldman, 1986). In addition, temperature changes, which mainly alter metabolic rates, can

also influence the diffusive uptake of nutrients into cells (Raven and Geider, 1988; Roleda et al., 2013). Increased levels of dissolved CO₂ during cell growth may result in higher cellular C:N and C:P ratios, due to increased CO₂ availability as a substrate for photosynthesis (Beardall et al., 2009; Feng et al., 2008; Fu et al., 2007; Fu et al., 2008). However, the dependency of C:N or C:P ratios on CO₂ availability can be species-specific (Burkhardt and Riebesell, 1997; Burkhardt et al., 1999). The effect of rising pCO₂ on the N:P ratio is still unclear, due to large variations observed in previous environmental manipulation studies (Sardans et al., 2012). For example, the N:P ratio of *Synechococcus* increased with elevated CO₂ concentration, but remained unchanged for *Prochlorococcus* (Fu et al., 2007) and *Emiliana huxleyi* (Feng et al., 2008).

Marine coccolithophores are responsible for almost half of the global marine calcium carbonate production, and are important in the marine carbon cycle through both the organic carbon pump and the inorganic carbon counter pump (Rost and Riebesell, 2004). *Emiliana huxleyi* is the most widely distributed coccolithophore species (Balch et al., 1991; Holligan et al., 1993; Holligan et al., 1983), and has been selected as a model phytoplankton species in the context of the marine carbon cycle (Westbroek et al., 1993). A wide range of environmental drivers, such as CO₂ concentration, nutrient concentration, irradiance, and temperature influence the growth, photosynthesis, and calcification of *E. huxleyi* both individually and interactively (Feng et al., 2017; Raven and Crawford, 2012; Zondervan, 2007). Changes in these physiological processes may in turn alter the elemental stoichiometry and composition of coccolithophores. Knowledge of how different environmental drivers will affect the elemental composition of *E. huxleyi* is important for a more complete understanding of the physiological responses of this species to the changing environment and the consequent effects on biogeochemical cycles. In addition, the magnitude of change of each environmental driver will be different with the future climate change, depending on location and scenario; hence a systematic study across a gradient of each driver is required.

This study advances previous findings by relating the change in elemental composition of *E. huxleyi* cells, in response to environmental forcing, to the physiological rate responses presented in the study of Feng et al. (2017). The major objective of the present study is to investigate and rank the importance of the environmental drivers, including the nitrate and phosphate concentrations, irradiance, temperature,

and $p\text{CO}_2$, on setting the elemental composition of a Southern Ocean strain of *E. huxleyi*. The combined results of this study and Feng et al. (2017) provide new insights into how environmental changes will impact the marine biogeochemical cycles related to *E. huxleyi*.

2 Materials and methods

5 2.1 Experimental setup

The marine coccolithophore *Emiliana huxleyi* (morphotype A, strain NIWA1108) was isolated from the [surface water \(depth of 5-6 m, salinity of 34.78\) at Chatham Rise, 41°35.8'S 175°41.5'E](#), east of New Zealand by Dr. H. Chang [aboard the *R/V Tangaroa* on the research voyage TAN0909 in November 2009, as detailed in Feng et al. \(2017\)](#). The water temperature was 12.1°C at the sampling site. The stock culture was maintained in the laboratory at 14°C and an irradiance of $\sim 140 \mu\text{mol m}^{-2} \text{s}^{-1}$, under a light/dark cycle of 12 h/12 h. The medium used for maintaining the stock culture was seawater obtained from Otago Harbour, New Zealand (nutrient concentrations: phosphate 0.3-0.6 μM , nitrate 3-6 μM), filtered using 0.2 μm pore size filtration cartridge (WhatmanTM) and supplemented with nutrient stock solutions to give final concentrations of nitrate 96 μM and phosphate 6 μM , without silicate addition. Trace metal and vitamin stock solutions were added according to the f/20 recipe for both stock culture and the manipulation experiments (10 times dilution of f/2 level; Guillard and Ryther, 1962).

For the manipulation experiments, *E. huxleyi* cells, in exponential growth phase [determined by the growth curve](#), were transferred into acid-cleaned 500-mL polycarbonate bottles with screw caps and subjected to a series of semi-continuous incubation experiments under different nutrient, irradiance, temperature, and $p\text{CO}_2$ conditions (Feng et al., 2017). Only one environmental driver was manipulated at a time for each incubation experiment, with the other environmental drivers remaining the same as the stock culture conditions. The manipulation of each of the different drivers was carefully selected to cover a broad range of conditions observed in the natural environment and those commonly employed for laboratory incubations (Feng et al., 2017). Initial cell abundances were $\sim 10^4 \text{ cell mL}^{-1}$ and *in vivo* chlorophyll *a* (Chl-*a*) fluorescence readings were monitored daily as indicators of Chl-*a* biomass and cell growth. After 2-3 days of acclimation under the experimental conditions daily dilution was started

带格式的: 字体: 倾斜

by adding freshly made seawater medium into each incubation bottle to adjust the biomass to the level of the previous day. All the incubation experiments were carried out in walk-in growth chambers (Model 650, Contherm, New Zealand), with metal halide lamps (full spectrum) as the light source, under a light/dark cycle of 12 h/12 h. Irradiance levels inside the incubation bottles were measured using a quantum PAR sensor (2100 series, Biospherical Instruments Inc., USA). The temperature in each incubation experiment was monitored continuously using a HOBO Pendant[®] temperature/light data logger (Onset, Australia), with variation no more than $\pm 0.5^{\circ}\text{C}$.

The experimental conditions for each environmental driver used for the manipulation experiments are presented in Table 1, and described in Feng et al. (2017). All the treatments were conducted in triplicate. For each culture, the final sampling was performed after the daily monitored growth rate remained relatively constant (daily variations $<10\%$) for more than seven generations (Feng et al., 2008). This yielded total acclimation of the cultures to the experimental conditions for ~ 20 days. Samples were collected for cell counts, Chl-*a* biomass, and elemental components, including particulate organic carbon (POC), particulate inorganic carbon (PIC), particulate organic nitrogen (PON), and particulate organic phosphorus (POP), starting 2 hours after the beginning of the light incubation phase and finishing within 2 hours for all the experimental treatments.

2.2 Sample analyses

2.2.1 Cell counts and Chl-*a*

One mL subsamples for cell counts were taken from each incubation bottle, preserved by adding 6 μL modified Lugol's solution, prepared by dissolving 10 g KI and 5 g iodine crystals in 20 mL Milli-Q water, then adding 50 mL of 5% anhydrous sodium acetate solution, and stored in dark at 4°C [for no more than 5 days before counting](#). Cell abundance was determined with a nanoplankton counting chamber (PhycoTech, USA) using a Zeiss microscope (Axiostar plus, Germany). *In vitro* Chl-*a* concentration was analysed using a Turner 10-AU fluorometer (Turner design, USA) with 90% acetone extracted samples, as detailed in Welschmeyer (1994).

2.2.2 Elemental composition

Subsamples from each incubation bottle for PON, total particulate carbon (TPC), and POC measurement were filtered onto pre-combusted GF/F glass fibre filters (Whatman™), and analysed using an elemental combustion system (Elementar vario EL III, Germany). Filters for POC analysis, 5 were fumed with saturated HCl for 3 hours in order to remove all the inorganic carbon (Zondervan et al., 2002). The PIC content was calculated by subtracting POC from TPC values. Subsamples for particulate organic phosphate (POP) measurements were filtered onto pre-combusted GF/F filters (Whatman™), and analysed following the molybdate colorimetric method of Solórzano and Sharp (1980). The particulate inorganic nitrogen (PIN) and particulate inorganic phosphorus (PIP) contents 10 were both neglected due to their relatively low values for laboratory cultures (Feng et al., 2008).

2.2.3 Data analyses

The effects of different environmental drivers on the elemental composition of *E. huxleyi* and their stoichiometric ratios were identified with one-way analysis of variance (ANOVA) using the commercial statistical software package SigmaStat (Version 3.5; Jandel Scientific, San Rafael, CA, US). Differences 15 between treatments were considered significant when $p < 0.05$. Post-hoc comparisons using the Student-Neuman-Keuls (SNK) test were conducted to determine any differences between particular treatments.

For the environmental drivers that had significant effects on the elemental composition and ratios of *E. huxleyi* within the examined range, the response curves to the drivers were fitted using the models listed in Table S1. All of the curve-fitting was performed using least square fit with Prism software 20 (version 5.0; GraphPad Prism Software, US) with all the triplicate data for each of the experimental treatments.

The same approach as used in Feng et al. (2017) was performed to rank the relative importance of each environmental driver, that was found to have significant effects using the one-way ANOVA analyses, on the elemental composition of *E. huxleyi*. Firstly, the two values for the elemental 25 composition at the average present-day conditions and the projected conditions for the year 2100 were derived from each fitted dose-response curve for the environmental driver that had significant effects.

The environmental conditions were projected using Coupled Model Intercomparison Project phase 5 (CMIP5) models (Boyd and Law, 2011; Law et al., 2016; Rickard et al., 2016), which suggested 33% decreases in both nitrate and phosphate concentrations, 2°C warming, a 25% increase in irradiance, and an increase in $p\text{CO}_2$ from 39 to 75 Pa in the Chatham Rise area for the year 2100 compared to present day conditions (Feng et al., 2017). The magnitude and direction of percentage change in the elemental composition under projected future conditions relative to the present day condition for each environmental driver was then calculated. The ranking was finally determined by comparing the absolute values of the calculated percentage changes of the physiological metrics caused by each driver. The driver that caused the largest percentage change was selected as the most important controlling driver.

3 Results

3.1 Changes in cellular POC content in response to environmental drivers

Cellular POC content was significantly affected by alteration of irradiance, temperature, and $p\text{CO}_2$ (Fig. 1). Increasing irradiance from 14 to 80 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ increased the cellular POC content by around two-fold from 8.20 ± 2.39 to 14.07 ± 1.17 pg cell^{-1} ($p < 0.05$). POC content decreased at the two highest irradiance levels (350 and 650 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, Fig. 1c). A trend of decreased *E. huxleyi* cellular POC content with elevated temperature was evident from the temperature manipulation experiment (Fig. 1d). The cellular POC content (28.85 ± 6.98 pg cell^{-1}) was significantly higher than all the other treatments ($p < 0.05$) at the lowest temperature of 4°C and significantly reduced by ~ 70% at both 20°C and 25°C ($p < 0.05$). Raising $p\text{CO}_2$ from 8 to 15 Pa significantly increased the cellular POC content from 9.63 ± 1.67 to 12.93 ± 1.84 pg cell^{-1} (Fig. 1e), with cellular POC content being relatively uniform from 15 to 109 Pa.

3.2 Alteration of cellular PIC content in response to environmental drivers

Temperature was the only driver that significantly altered the cellular PIC content (Fig. 2). There was a general trend of decreased cellular PIC content of *E. huxleyi* with warming from 11°C to 20°C (Fig. 2d). The cellular PIC content was significantly lower at 20°C and 25°C compared to the other four

temperature treatments ($p < 0.05$). More than a 50% decrease in cellular PIC content was observed at the two highest temperature conditions, relative to the 7°C treatment. However, there were no significant differences in cellular PIC content between the other temperature treatments. The fitted Q_e value (the plateau for one phase decay) was 6.94 ± 0.93 pg cell⁻¹, close to the average value at the two highest temperatures (Table S1).

3.3 Changes in the cellular PIC:POC ratio in response to environmental drivers

As for POC, the cellular ratio of PIC:POC was mainly affected by changes in irradiance, temperature, and $p\text{CO}_2$ (Fig. 3). The highest cellular PIC:POC ratio of 1.20 ± 0.09 was observed at the lowest irradiance ($19 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$; $p < 0.05$, compared to all other irradiance treatments). The ratio then decreased with increasing irradiance to 0.72 ± 0.10 at $190 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ and slightly increased again at the two highest irradiances ($p < 0.05$ between 190 and $650 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, Fig. 3c). In the temperature manipulation experiment, the PIC:POC ratio was significantly lower ($p < 0.05$) at the lowest temperature (4°C) than any other treatment, with a value of 0.45 ± 0.03 pg cell⁻¹. The PIC:POC value then leveled off between the range of 7°C to 25°C, with the average value more than double that at 4°C (Fig. 3d). With the variation of $p\text{CO}_2$ levels, the cellular PIC:POC ratio decreased by more than 40% from 1.46 ± 0.02 pg cell⁻¹ at 8 Pa to 0.90 ± 0.15 at 39 Pa and stayed similar between the range of 39 and 109 Pa ($p < 0.05$) (Fig. 3e), mainly due to the increased cellular POC quota with rising $p\text{CO}_2$.

3.4 Alteration of cellular PON content in response to environmental drivers

The cellular PON content increased with increasing nitrate concentration. The content at the two lowest nitrate concentrations of 3.7 and 6.0 μM was less than half of the average value (2.06 ± 0.36 pg cell⁻¹) of the three highest nitrate treatments (Fig. 4a). Warming from 4°C to 25°C decreased the cellular PON content ($p < 0.05$). The value of 4.07 ± 0.00 pg cell⁻¹ at 4°C was double that at 15°C (1.93 ± 0.10 pg cell⁻¹) and three-fold greater than the PON content of 1.31 ± 0.24 pg cell⁻¹ at 25°C (Fig. 4d).

3.5 Changes in cellular POP content in response to environmental drivers

The cellular POP content of *E. huxleyi* was significantly altered by nitrate, phosphate, temperature, and $p\text{CO}_2$. POP content was slightly less at the three low nitrate concentrations (3.7, 6.0, and 12 μM), compared to those at 96 and 200 μM ($p < 0.05$; Fig. 5a). Cellular POP content significantly increased with rising phosphate concentration (Fig. 5b), with the highest POP content observed at 20 μM phosphate. As observed for cellular POC and cellular PON contents, warming greatly decreased the cellular POP content (Fig. 5d), with a reduction of 65% from $1.08 \pm 0.14 \text{ pg cell}^{-1}$ at 4°C to $0.38 \pm 0.04 \text{ pg cell}^{-1}$ at 11°C , but then only a further decrease of $\sim 0.1 \text{ pg cell}^{-1}$ from 15°C to 25°C . Significant differences in POP content were detected between the two lowest temperature treatments compared to all others. Conversely, with rising $p\text{CO}_2$ level there was a trend of increased cellular POP content (Fig. 5e), which almost doubled from $0.20 \pm 0.04 \text{ pg cell}^{-1}$ at 8 Pa to $0.38 \pm 0.02 \text{ pg cell}^{-1}$ at 109 Pa ($p < 0.05$).

3.6 Alteration of cellular C to Chl-*a* ratio in response to environmental drivers

Alteration of all the five environmental drivers greatly affected the cellular ratio of POC to Chl-*a* content (C:Chl-*a*, g:g) ($p < 0.05$) (Fig. 6). C:Chl-*a* decreased exponentially with increased nitrate concentration up to 50 μM , but stabilised between 50 and 200 μM (Fig. 6a). The highest ratio of 422.36 ± 74.28 was observed at the lowest nitrate concentration of 3.7 μM , significantly higher than all other treatments ($p < 0.05$). The ratio then decreased by 87% at 200 μM . An increase in phosphate concentration, however, only slightly decreased the C:Chl-*a* ratio (Fig. 6b). Compared to the ratios at the two lowest concentrations, a significant decrease ($p < 0.05$) at 6.0 μM and 20 μM was observed (by $\sim 20\%$ each). Increased irradiance increased the C:Chl-*a* ratio linearly, with more than a doubling at 650 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ compared to the ratio of 47.45 ± 12.58 at 14 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (Fig. 6c). The C:Chl-*a* ratio dramatically decreased with warming, especially between 4°C and 7°C (Fig. 6d). The ratio of 131.26 ± 42.96 observed at 4°C was significantly higher than all the other temperatures ($p < 0.05$). Significantly lower C:Chl-*a* ratios were observed at the two lowest $p\text{CO}_2$ levels of 8 and 15 Pa compared with the other treatments ($p < 0.05$, Fig. 6e), with the ratio increasing by 42% from low to high $p\text{CO}_2$.

3.7 Shifts in cellular elemental molar ratios in response to environmental drivers

The PON to POP (N:P) ratio was significantly lower ($p < 0.05$) at the two lowest nitrate treatments compared to the others (Table 2). In contrast, the POC to PON (C:N) ratio was significantly higher ($p < 0.05$) at the two lowest nitrate concentrations. There was no significant difference in C:N ratios across the other four nitrate treatments ($p > 0.05$). Changes in nitrate concentration did not significantly affect the POC to POP (C:P) ratio.

The N:P ratio of *E. huxleyi* increased at low phosphate concentrations (0.4 and 2 μM), with highest value in the 0.4 μM phosphate treatment ($p < 0.05$). There was a significant increase in the C:P ratio ($p < 0.05$) at the two lowest phosphate concentrations compared to the others. The highest C:P ratio, recorded at the lowest phosphate concentration (0.4 μM), was almost double the value at 2 μM and more than three times the average ratio of the other treatments (Table 2). In contrast, there were no significant differences in the calculated C:N ratio across the phosphate treatments ($p > 0.05$).

Decreased C:N ratios were observed for low irradiances; the value at 14 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ being significantly lower than the three highest irradiances ($p < 0.05$). Similarly, a decreased C:P ratio was found at low irradiance, with a significantly lower value at 14 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ compared to the three highest irradiances. Warming significantly increased the N:P ratio from 4°C to 20°C (Table 2).

3.8 Ranking the importance of environmental drivers in altering *Emiliana huxleyi* elemental composition

Ranking the response of the Southern Ocean *E. huxleyi* isolated to projected future changes in oceanic properties revealed differential responses between drivers and processes (Table 3, Fig. 7). Cellular POC and cellular PIC:POC ratio were both significantly influenced by CO_2 and temperature, with temperature affecting cellular POC content the most, while CO_2 was the most important factor regulating PIC:POC. However, only one driver (temperature) significantly regulated cellular PIC, with a 4°C warming causing a 14.2 % decrease. The cellular PON content was significantly affected by future nitrate concentration and temperature, with nitrate ranking the most important. Four (phosphate, temperature, CO_2 , and nitrate) out the five environmental drivers, under end of the century conditions, significantly affected cellular POP content, with future phosphate concentration playing the most

important role. The rankings associated with statistically non-significant differences among the treatment intervals, as marked in Table 3 and Fig. 7, need to be considered with caution (see Feng et al. (2017)).

4 Discussion

5 This is the first detailed study of the individual effect of five environmental drivers (nitrate concentration, phosphate concentration, irradiance, temperature, and $p\text{CO}_2$) on the cellular elemental composition of the coccolithophore *E. huxleyi*. Moreover, it is the first to rank the importance of the predicted changes in these environmental drivers on *E. huxleyi* elemental stoichiometry for the year 2100 relative to the present-day conditions. Relating changes in elemental composition is an important
10 addition to the responses of growth, photosynthesis, and calcification rates (Feng et al., 2017), providing insights into the biogeochemical consequences of the physiological effects induced by change in the five essential environmental drivers.

4.1 Effects of nutrient concentration on the elemental stoichiometry of *Emiliana huxleyi*

The PON and POP cell quotas of *E. huxleyi* in the present study were mainly controlled by nitrate
15 and phosphate concentrations, respectively, as phytoplankton relies on seawater nutrient availability as the external elemental source (Hecky et al., 1993; Price, 2005; Sakshaug and Holmhansen, 1977). Nitrate concentration plays an important role in regulating the growth, photosynthetic, and calcification rates of *E. huxleyi* (Feng et al., 2017); however, the three lowest nitrate concentrations only resulted in slightly decreased cellular POP contents and had no significant effect on cellular POC or PIC content.
20 This indicates that the regulation of the nitrate concentration on the POC and PIC productivity in our study was mainly a consequence of decreased growth rate of the cells under nitrate limitation, as shown by Feng et al. (2017). This finding is in contrast to Paasche (1998) who observed higher *E. huxleyi* cellular PIC:POC ratios under nitrate limitation as a result of decreased cellular POC and increased coccolith abundance per cell in *E. huxleyi* strain BOF 92 isolated from the North Atlantic. Higher
25 PIC:POC ratios under nitrate limitation was alternatively attributed to increased calcite mass per lith of *E. huxleyi* strain CCMP 378 isolated from the Gulf of Maine (Fritz, 1999). In addition, phosphate

concentration did not significantly affect *E. huxleyi* cellular carbon content nor the PIC:POC ratio of cells in the present study. However, Paasche (1998) observed greatly increased PIC content of *E. huxleyi* (strain BOF 92) under phosphate limiting conditions, and Riegman et al. (2000) observed that a greater increase PIC quotas under phosphate limitation than nitrate limitation for *E. huxleyi* (strain L).

5 These discrepancies between studies in the nitrate or phosphate effects on cellular PIC:POC ratio are mainly due to the different nutrient concentrations in the culturing media. Paasche (1998) observed an increase in *E. huxleyi* PIC cell quota under the stationary phase of batch incubation, i.e. when cell division ceased as nitrate dropped to ≤ 0.2 μM and phosphate dropped to ≤ 0.03 μM . This supports the findings of both Riegman et al. (2000) and Fritz (1999) who conducted continuous incubations with
10 high cell densities of *E. huxleyi*. These studies observed an increased cellular PIC content when phosphate concentration fell below 0.4 nM (Riegman et al., 2000) or nitrate concentration was below the detection limit (Fritz, 1999). However, the present study used a semi-continuous incubation method with higher and relatively steady nutrient concentrations (with lowest nitrate and phosphate concentrations of 3.6 and 0.4 μM , respectively) and the cells were grown and sampled at a healthy
15 exponential growth phase. Similarly, Müller et al. (2008) only found higher *E. huxleyi* (strain CCMP371) cellular calcite content during the stationary but not the exponential growth phase under both nitrate and phosphate limitation, due to the different cell cycle phases during which the calcification and cell division occurred. The authors explained that calcification continued during the G1 phase of cell assimilation when cell division was restricted under nutrient limitation, and thus the
20 cellular PIC content was increased (Müller et al., 2008). Further studies at extremely low nutrient concentrations (< 0.1 μM) in a steady-state growth phase are still needed to understand the potential connection between carbon production and extreme nutrient limitation, given reports of areal expansion of oligotrophic waters in the world oceans with global climate change (Polovina et al., 2008).

4.2 Irradiance effects on the elemental stoichiometry of *Emiliana huxleyi*

25 In the present study, irradiance was the main environmental factor affecting cellular POC content which in turn altered the PIC:POC ratio. The increased PIC:POC cellular ratio at low irradiance indicates that calcification is less dependent on irradiance than organic carbon fixation, as discussed in

Feng et al. (2017). Although both processes require light as an energy source, calcification requires less energy (Anning et al. 1996) than photosynthesis (Paasche, 1965; Balch et al., 1992). Therefore, the calcification rate is generally saturated at lower irradiance levels than photosynthesis (Paasche, 1964; Zondervan, 2007). Feng et al. (2017) reported greatly reduced photosynthetic rates under the two lowest irradiance levels, while observing that this trend was less significant for the calcification rate. Hence, limiting irradiance will lead to less POC content in the cells compared to the cellular PIC quota, and thus a higher cellular PIC:POC ratio would be expected at low irradiance when growth and photosynthesis are light-limited (Raven and Crawford, 2012), as confirmed by the response of calcification:photosynthesis in Feng et al. (2017).

10 Increasing irradiance also elevated the C:Chl-*a* ratio linearly in the present study, due to the increase in POC quota and a decrease in Chl-*a* quota, as also reported for diatoms and dinoflagellates (Geider, 1987). The reduced cellular pigment quota under high irradiance helps to reduce the energy required for light harvesting in phytoplankton cells, which is a strategy to balance the energy demands for growth and POC production with photon harvesting (Kiefer, 1993). In addition, the present study revealed that
15 the C:N and C:P ratios of *E. huxleyi* both increased at high light levels, as a consequence of increased cellular POC content driven by increased irradiance but no significant change in cellular PON or POP quota, further suggesting that organic carbon content is more light dependent than the accumulation of cellular N or P (Geider et al., 1998).

4.3 Temperature effects on the elemental stoichiometry of *Emiliana huxleyi*

20 Temperature is important in regulating dissolved chemical diffusion and transport, non-enzymatic and enzymatic reactions, and the metabolic rates of phytoplankton (Raven and Geider, 1988). In our accompanying study, the growth, photosynthetic, and calcification rates all increased with rising temperature until the optimal temperature was reached at 25°C, 24°C, and 20°C respectively (Feng et al., 2017), which were all higher than the stock culture growth temperature or the temperature at the
25 isolation site of *E. huxleyi* strain NIWA 1108. In the present studyConversely, the cellular POC, PON, and POP content all reduced significantly as temperature increased in the present study. It has been proposed that reduced cell size is a universal strategy in response to increasing temperature for both

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

terrestrial and aquatic organisms (Gardner et al., 2011), following a hypothesis suggested by Atkinson et al. (2003). A study on the coccolithophores *E. huxleyi* (strain EH2) and *Gephyrocapsa oceanic* (strain GO1) observed decreased cell size and thinner coccospheres upon raising temperature from 10°C to 25°C, which was attributed to the relatively suppressed cell division at low temperature (Sorrosa et al., 5 2005). This decrease in cell volume (Fig. S1) could be the main cause of reduced cellular elemental components in the present study. Previous studies also reported that warming resulted in reduced cell volume of *E. huxleyi* (strain AC481, De Bodt et al., 2010; strain L, van Rijssel and Gieskes, 2002), and decreased cellular POC and PIC quotas of coccolithophore *Coccolithus pelagicus* when the temperature was raised from 10°C to 15°C (Gerecht et al., 2014). Similarly, warming significantly decreased the 10 cellular elemental contents to their lowest levels measured in the present study over the range from 4°C to 25°C, with a decrease in cell size at higher temperatures (Fig. S1), as growth rate increased (Feng et al, 2017).

However, contrary to the observed changes in POC, PON, and POP cell quota, the cellular PIC content of *E. huxleyi* only decreased when temperature was higher than 11°C in the present study, due to 15 the strongly reduced calcification and malformation at low temperatures of 4°C and 7°C (Feng et al., 2017). The reduced cell division rate (i.e. enlarged cell volume, Fig. S1) offset the reduced calcification rate at lower temperatures, and so there was no significant difference in PIC cell quota at temperatures below 11°C. Consequently, the cellular PIC:POC ratio was lower at 4°C and 7°C, consistent with the trend observed for the calcification: photosynthesis ratio (Feng et al., 2017), indicating suppression of 20 PIC formation relative to POC production at low temperature (Watabe and Wilbur, 1966). The PIC:POC ratio then decreased with warming from 11°C to 15°C and remained relatively steady afterwards, mainly due to the lower optimal temperature for calcification (20°C) compared to photosynthesis (24°C) as suggested in Feng et al. (2017).

Furthermore, warming from 4°C to 20°C significantly increased the *E. huxleyi* cellular N:P ratio in 25 the present study, in agreement with the recent model study on a natural phytoplankton community (Toseland et al., 2013). Toseland et al. (2013) found that with increasing temperature the rate of cellular protein synthesis in phytoplankton was higher, but with a lower number of phosphorus-rich ribosomes, thereby increasing the cellular N:P ratio. In the present study, the cellular N:P ratio of *E. huxleyi* at 20°C

increased by 74% from that at 4°C, in spite of both cellular PON and cellular POP decreasing with warming. Although this study presents results for a single strain of *E. huxleyi*, if the temperature dependency of cellular resource allocation is a universal trend for all the *E. huxleyi* genotypes, we can speculate that the diverse *E. huxleyi* strains growing in different temperature regions might have different requirements for nitrogen vs. phosphorus, and that the growth of *E. huxleyi* strains in the temperate to tropical regions might be more readily limited by nitrate than sub-polar strains. Similarly, Toseland et al. (2013) suggested that future warming might accentuate nitrate limitation in the oceans.

4.4 Effects of CO₂ on the elemental stoichiometry of *Emiliana huxleyi*

The photosynthesis of *E. huxleyi* was saturated at a higher $p\text{CO}_2$ than that for growth rate (Feng et al., 2017). In the present study, CO₂ plays the most important role in regulating the cellular PIC:POC ratio. The PIC:POC ratio was significantly higher at the lowest $p\text{CO}_2$ level, as a consequence of the lower cellular POC and higher cellular PIC at 8 Pa. In general, cell growth of *E. huxleyi* is less limited by low CO₂ concentrations than in other phytoplankton groups (Clark and Flynn, 2000; Paasche et al., 1996; Riebesell et al., 2000a; Rost et al., 2003). Moreover, recent studies suggest that *E. huxleyi* operates an active carbon concentrating mechanism (CCM) to utilize HCO₃⁻ through the enzyme carbonic anhydrase (CA; Reinfelder, 2011), and may have high affinity for CO₂ in photosynthesis (Stojkovic et al., 2013). However, the efficiency of CCMs in *E. huxleyi* (strain B92/11) is considered to be low as a consequence of the leakage of CO₂ from the cell (Rost et al., 2006), and so coccolithophore photosynthesis is more dependent than cell growth on CO₂ concentration (Rost and Riebesell, 2004). This discrepancy between growth and organic carbon fixation can lead to a decrease in cellular POC at low $p\text{CO}_2$. This difference in CO₂ requirements between the two processes may also have resulted in the lower cellular POP content at 8 Pa compared to other $p\text{CO}_2$ treatments.

The increasing trend observed for cellular POC and POP was not apparent for cellular PIC quota, as calcification rates significantly decreased with increasing $p\text{CO}_2$ level >40 Pa (Feng et al., 2017). Hence the cellular PIC:POC ratio was significantly higher at the two lowest $p\text{CO}_2$ levels, consistent with previous findings for CO₂ manipulations at saturating irradiances on *E. huxleyi* (strain PML B92/11A; Zondervan et al., 2002; Zondervan et al., 2001). No further significant change in cellular carbon content

or PIC:POC ratio occurred at higher $p\text{CO}_2$, in contrast to the linear decrease in the calcification:photosynthesis ratio with rising $p\text{CO}_2$ (Feng et al., 2017). This difference is noteworthy as both cellular PIC:POC and calcification:photosynthesis ratios are commonly used to examine the relative change of PIC and POC production in coccolithophores (Raven and Crawford, 2012). These changes have biogeochemical implications for the marine rain ratio in the carbon cycle (Klaas and Archer, 2002; Rost and Riebesell, 2004), which is the export ratio of calcite to organic carbon into the deep ocean. The ^{14}C -labelling technique used in this study (see Feng et al., 2017) to measure carbon fixation (photosynthesis and calcification) rates was conducted during the light period, thus the measured rate is an indicator of net carbon fixation, that does not account for the energy-consuming respiratory process or CO_2 leakage out of the cells (Bach et al., 2015, 2013; Rost et al., 2006). Conversely, the cellular carbon content indicates the gross accumulated carbon in the cells over longer period of growth (Engel et al., 2010; Fabry and Balch, 2010). The most compelling reason for the relatively higher PIC:POC ratio (~1.5) than calcification: photosynthesis ratio (~1.0) in the lowest $p\text{CO}_2$ treatment (8 Pa, 79 ppm) in our study may then be attributed to diffusive CO_2 loss limiting inorganic carbon active uptake from the substrate (Bach et al., 2013), resulting in less POC fixation into the cells relative to the PIC fixation by the calcification process.

The C:Chl-*a* ratio of *E. huxleyi* was lowest at $p\text{CO}_2$ of 8 Pa across all the $p\text{CO}_2$ treatments in the present study, mainly due to the decreased cellular POC at low $p\text{CO}_2$, rather than any change in cellular Chl-*a* content. However, increasing $p\text{CO}_2$ did not have significant effects on the C:N, N:P or C:P ratios in the present study. This is in accordance with a recent study on *E. huxleyi* (strain PML B92/11A), which also exhibited constant C:N:P ratios across a $p\text{CO}_2$ range of 18 to 75 Pa for cultures at steady growth phase under phosphate-limited continuous incubation (Engel et al., 2014).

4.5 Biogeochemical implications and future directions

The comparisons between present day conditions and those projected for year 2100 for the Chatham Rise area are summarized in the conceptual figure (Fig. 7). These results indicate that the 2°C warming will decrease both POC and PIC cellular quotas of *E. huxleyi*, but may slightly increase the PIC:POC ratio. Rising $p\text{CO}_2$ alone will result in decreased cellular PIC:POC ratio. Although the 33% decrease in

nitrate concentration is the major factor controlling the growth, photosynthetic, and calcification rates (Feng et al., 2017), change in nitrate concentration did not significantly affect the elemental stoichiometry except for the cellular PON contents of *E. huxleyi*. In addition, increasing temperature may increase the cellular N:P ratio, while rising $p\text{CO}_2$ will decrease the N:P and C:P ratios. These results provide a more detailed perspective that can improve our knowledge on how the model coccolithophore species, *E. huxleyi*, may respond to future environmental changes. For example, our results suggest that rising $p\text{CO}_2$ in the future oceanic environment will decrease the *E. huxleyi* cellular PIC:POC ratio by 5.4%; however, the projected warming and increase in irradiance level may offset this decreased PIC:POC by 2.4% and 0.3%, respectively. The changes in PIC:POC have implications for the marine “rain ratio” and so alter the marine carbon cycle (Rost and Riebesell, 2004). Similarly, the cellular N:P ratio will be decreased by rising $p\text{CO}_2$, although this trend may be canceled out by warming. The altered C:N:P stoichiometry will in turn affect the nutrient cycle at higher trophic levels (Jones and Flynn, 2005) and marine biogeochemical cycles (Beardall and Raven, 2004).

It is noteworthy that the research presented here only examined the physiological response norms of *E. huxleyi* to a single environmental driver when other drivers were all kept at the stock culture growth condition (i.e. a set of single dimensional space experiments). However, these responses (such as the shape of the curves and the optimal conditions) may be different when the other background conditions are changed. For example, Sett et al. (2014) observed the dose-response curves of calcification of *E. huxleyi* PML B9/11 to CO_2 concentration was regulated by temperature. Therefore, in order to comprehensively understand how *E. huxleyi* physiology will respond to multiple environmental drivers and fill this knowledge gap, future research on a full environmental matrix is still necessary. These experiments will not only help to further explore the potential interactions (i.e. synergistic or agnostic effects) between environmental drivers, but also provide a better understanding of the underlying mechanisms of these interactive effects. In addition, the present study is only based on a single strain of southern hemisphere *E. huxleyi*. Due to the wide distribution of this species in the natural marine environment, *E. huxleyi* presents high variability in terms of genetic, morphological, and physiological characteristics (Cook et al., 2011; Read et al., 2013; Young et al., 2014). Therefore, the physiology of different *E. huxleyi* strains isolated from different geographic locations might respond differently to

changing environmental drivers. For example, within the context of OA research, extensive previous studies suggest a strain-specificity of *E. huxleyi* in response to changes in seawater carbonate chemistry (Langer et al., 2009; Raven and Crawford, 2012; Blanco-Ameijerías et al., 2016). It has also been observed that different *E. huxleyi* ecotypes/morphotypes responded differently to OA (Müller et al., 5 2015), which is likely a consequence of their genetic variation (Cook et al., 2011). The present study and Feng et al. (2017) demonstrate the important roles of different environmental drivers in controlling the physiology of *E. huxleyi* strain NIWA1108, and so further work is required to determine if the findings apply to other strains.

In summary, this study, in combination with Feng et al. (2017), have a number of implications for 10 research into the response of *E. huxleyi* to ocean acidification and global climate change. In addition to seawater carbonate chemistry (Riebesell et al., 2010), it is necessary to report the experimental conditions of all the environmental drivers carefully. The predictions presented will provide useful information for biogeochemical models, such as that of Bopp et al. (2001), of how the elemental stoichiometry of *E. huxleyi* will respond to the alteration of these environmental conditions individually, 15 in order to predict the future changes in the marine biogeochemical cycles. In addition, multiple environmental drivers tend to change simultaneously in the future global climate change scenario (Boyd and Hutchins, 2012), and so future studies should also investigate the interactions between these multiple drivers on phytoplankton physiology. The predicted future changes in marine physical properties (such as sea surface temperature (SST) and mixed layer depth) will vary from one oceanic 20 region to another (Boyd and Doney, 2002). The dose response curves from our study suggest that the range of alteration in environmental drivers may control the outcome of the effects of environmental perturbation on *E. huxleyi* physiology and biogeochemistry. For future multi-factorial manipulation experimental designs, our results suggest that the magnitudes of change in each environmental driver need to be determined/decided cautiously and should have environmental relevance in order to make 25 more accurate predictions, and the understanding of interactive effects of multiple environmental drivers and the underlying mechanisms should be further explored.

5 Acknowledgements

We would like to thank Hoe Chang at National Institute of Water and Atmospheric Research Ltd. (NIWA) Wellington for providing the stock culture of *Emiliana huxleyi* NIWA1108. We also thank Kim Currie at NIWA, Department of Chemistry, University of Otago for helping with the DIC and alkalinity analysis and Graham Rickard for generating the future estimates of environmental variables for New Zealand waters from the CMIP5 models. This work was supported by New Zealand Marsden grant (09-UOO-175) to CLH and National Natural Science Foundation of China (No. 41306118 and No. 41676160) to YF.

10 References

- Anning, T., Nimer, N., Merrett, M. J., and Brownlee, C.: Costs and benefits of calcification in coccolithophorids, *J. Mar. Syst.*, 9, 45-56, doi:10.1016/0924-7963(96)00015-2, 1996.
- Atkinson, D., Ciotti, B. J., and Montagnes, D. J. S.: Protists decrease in size linearly with temperature: ca. 2.5% degrees C⁻¹, *Proc. R. Soc. Lond. [Boil.]*, 270, 2605-2611, doi:10.1098/rspb.2003.2538, 2003.
- 15 Bach, L.T., Mackinder, L.C.M., Schulz, K.G., Wheeler, G., Schroeder, D.C., Brownlee, C., Riebesell, U.: Dissecting the impact of CO₂ and pH on the mechanisms of photosynthesis and calcification in the coccolithophore *Emiliana huxleyi*, *New Phytol.*, 199, 121-134, doi:10.1111/nph.12225, 2013.
- Bach, L. T., Riebesell, U., Gutowska, M. A., Federwisch, L., Schulz, K. G.: A unifying concept of coccolithophore sensitivity to changing carbonate chemistry embedded in an ecological framework,
- 20 *Prog. Oceanogr.*, 135, 125-138, doi:10.1016/j.pocean.2015.04.012, 2015.
- Balch, W. M., Holligan, P. M., Ackleson, S. G., and Voss, K. J.: Biological and optical properties of mesoscale coccolithophore blooms in the Gulf of Maine, *Limnol. Oceanogr.*, 36, 629-643, 1991.
- Balch, W. M., Holligan, P. M., and Kilpatrick, K. A.: Calcification, photosynthesis and growth of the bloom-forming coccolithophore, *Emiliana huxleyi*, *Cont. Shelf Res.*, 12, 1353-1374, doi:10.1016/0278-
- 25 4343(92)90059-s, 1992.

- Beardall, J., and Raven, J. A.: The potential effects of global climate change on microalgal photosynthesis, growth and ecology, *Phycologia*, 43, 26-40, doi:10.2216/i0031-8884-43-1-26.1, 2004.
- Beardall, J., Allen, D., Bragg, J., Finkel, Z. V., Flynn, K. J., Quigg, A., Rees, T. A. V., Richardson, A., and Raven, J. A.: Allometry and stoichiometry of unicellular, colonial and multicellular phytoplankton,
5 *New Phytol.*, 181, 295-309, doi:10.1111/j.1469-8137.2008.02660.x, 2009.
- Blanco-Ameijeiras, S., Lebrato, M., Stoll, H.M., Iglesias-Rodriguez, D., Müller, M.N., Méndez-Vicente, A., Oschlies, A.: Phenotypic variability in the Coccolithophore *Emiliana huxleyi*. *PLoS ONE*, 11(6): e0157697, doi:10.1371/journal.pone.0157697, 2016.
- Bopp, L., Monfray, P., Aumont, O., Dufresne, J. L., Le Treut, H., Madec, G., Terray, L., and Orr, J. C.:
10 Potential impact of climate change on marine export production, *Global Biogeochem. Cy.*, 15, 81-99, doi:10.1029/1999gb001256, 2001.
- Boyd, P. W., and Doney, S. C.: Modelling regional responses by marine pelagic ecosystems to global climate change, *Geophys. Res. Lett.*, 29, 1806, 10.1029/2001gl014130, 2002.
- Boyd, P. W., Strzepek, R., Fu, F. X., and Hutchins, D. A.: Environmental control of open-ocean
15 phytoplankton groups: Now and in the future, *Limnol. Oceanogr.*, 55, 1353-1376, doi:10.4319/lo.2010.55.3.1353, 2010.
- Boyd, P. W., and Law, C. S.: A climate change atlas for the New Zealand exclusive economic zone, National Institute of Water and Atmosphere (NIWA) Information Series, Wellington, New Zealand, 22 pp., 2011.
- 20 Boyd, P. W., and Hutchins, D. A.: Understanding the responses of ocean biota to a complex matrix of cumulative anthropogenic change, *Mar. Ecol. Prog. Ser.*, 470, 125-135, doi:10.3354/meps10121, 2012.
- Boyd, P. W., Dillingham, P. W., McGraw, C. M., Armstrong, E. A., Cornwall, C. E., Feng, Y., Hurd, C. L., Gault-Ringold, M., Roleda, M. Y. Timmins-Schiffman, E., Nunn, B. L.: Physiological responses of a Southern Ocean diatom to complex future ocean conditions, *Nat. Clim. Chang.*, 6, 207-213, 2016.
- 25 Burkhardt, S., and Riebesell, U.: CO₂ availability affects elemental composition (C:N:P) of the marine diatom *Skeletonema costatum*, *Mar. Ecol. Prog. Ser.*, 155, 67-76, doi:10.3354/meps155067, 1997.
- Burkhardt, S., Zondervan, I., and Riebesell, U.: Effect of CO₂ concentration on C : N : P ratio in marine phytoplankton: a species comparison, *Limnol. Oceanogr.*, 44, 683-690, 1999.

- Clark, D. R., and Flynn, K. J.: The relationship between the dissolved inorganic carbon concentration and growth rate in marine phytoplankton, *Proc. R. Soc. Lond. [Boil.]*, 267, 953-959, 2000.
- Cook, S. S., Whitlock, L., Wright, S. W., and Hallegraeff, G. M.: Photosynthetic pigment and genetic differences between two southern ocean morphotypes of *Emiliana huxleyi* (Haptophyta), *J. Phycol.*, 47, 615-626, doi:10.1111/j.1529-8817.2011.00992.x, 2011.
- 5 De Bodt, C., Van Oostende, N., Harlay, J., Sabbe, K., and Chou, L.: Individual and interacting effects of $p\text{CO}_2$ and temperature on *Emiliana huxleyi* calcification: study of the calcite production, the coccolith morphology and the coccosphere size, *Biogeosciences*, 7, 1401-1412, doi:10.5194/bg-7-1401-2010, 2010.
- 10 Engel, A., Barcelos e Ramos, J., Geider, R., Hutchins, D. A., Lee, C., Rost, B., Röttgers, R., and Thingstad, F.: Production and export of organic matter, in: *Guide to Best Practices for Ocean Acidification Research and Data Reporting*, edited by: Riebesell, U., Fabry, V. J., Hansson, L., and Gattuso, J. P., Publications Office of the European Union, Luxembourg, 181-200, 2010.
- Engel, A., Novoa, C. C., Wurst, M., Endres, S., Tang, T., Schartau, M., and Lee, C.: No detectable effect of CO_2 on elemental stoichiometry of *Emiliana huxleyi* in nutrient-limited, acclimated continuous cultures, *Mar. Ecol. Prog. Ser.*, 507, 15-30, doi:10.3354/meps10824, 2014.
- 15 Fabry, V. L., and Balch, W. M.: Direct measurements of calcification rates in planktonic organisms, in: *Guide for Best Practices in Ocean Acidification Research and Data Reporting*, edited by: Riebesell, U., Fabry, V. L., Hansson, L., and Gattuso, J.P., Publications Office of the European Union, Luxembourg, 201-212, 2010.
- 20 Feng, Y., Warner, M. E., Zhang, Y., Sun, J., Fu, F. X., Rose, J. M., and Hutchins, D. A.: Interactive effects of increased $p\text{CO}_2$, temperature and irradiance on the marine coccolithophore *Emiliana huxleyi* (Prymnesiophyceae), *Eur. J. Phycol.*, 43, 87-98, doi:10.1080/09670260701664674, 2008.
- Feng, Y., Roleda, M.Y., Armstrong, E.A., Boyd, P.W. and Hurd, C.L.: Environmental controls on the growth, photosynthetic and calcification rates of a Southern Hemisphere strain of the coccolithophore *Emiliana huxleyi*, *Limnol. Oceanogr.*, 62: 519-540, doi:10.1002/lno.10442, 2017.
- 25

- Finkel, Z. V., Beardall, J., Flynn, K. J., Quigg, A., Rees, T. A. V., and Raven, J. A.: Phytoplankton in a changing world: cell size and elemental stoichiometry, *J. Plankton Res.*, 32, 119-137, doi:10.1093/plankt/fbp098, 2010.
- Fritz, J. J.: Carbon fixation and coccolith detachment in the coccolithophore *Emiliania huxleyi* in nitrate-limited cyclostats, *Mar. Biol.*, 133, 509-518, doi:10.1007/s002270050491, 1999.
- 5 Fu, F. X., Warner, M. E., Zhang, Y. H., Feng, Y. Y., and Hutchins, D. A.: Effects of increased temperature and CO₂ on photosynthesis, growth, and elemental ratios in marine *Synechococcus* and *Prochlorococcus* (Cyanobacteria), *J. Phycol.*, 43, 485-496, doi:10.1111/j.1529-8817.2007.00355.x, 2007.
- 10 Fu, F. X., Zhang, Y. H., Warner, M. E., Feng, Y. Y., Sun, J., and Hutchins, D. A.: A comparison of future increased CO₂ and temperature effects on sympatric *Heterosigma akashiwo* and *Prorocentrum minimum*, *Harmful Algae*, 7, 76-90, doi:10.1016/j.hal.2007.05.006, 2008.
- Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L., and Heinsohn, R.: Declining body size: a third universal response to warming?, *Trends Ecol. Evol.*, 26, 285-291, doi:10.1016/j.tree.2011.03.005, 2011.
- 15 Geider, R. J.: Light and temperature-dependence of the carbon to chlorophyll-*a* ratio in microalgae and cyanobacteria - implications for physiology and growth of phytoplankton, *New Phytol.*, 106, 1-34, doi:10.1111/j.1469-8137.1987.tb04788.x, 1987.
- Geider, R. J., MacIntyre, H. L., and Kana, T. M.: A dynamic regulatory model of phytoplanktonic acclimation to light, nutrients, and temperature, *Limnol. Oceanogr.*, 43, 679-694, 1998.
- 20 Gerecht, A. C., Supraha, L., Edvardsen, B., Probert, I., and Henderiks, J.: High temperature decreases the PIC/POC ratio and increases phosphorus requirements in *Coccolithus pelagicus* (Haptophyta), *Biogeosciences*, 11, 3531-3545, doi:10.5194/bg-11-3531-2014, 2014.
- Goldman, J. C.: On phytoplankton growth-rates and particulate C:N:P ratios at low light, *Limnol. Oceanogr.*, 31, 1358-1363, 1986.
- 25 Guillard, R. R., and Ryther, J. H.: Studies of marine planktonic diatoms.1. *Cyclotella nana* (Hustet), and *Detonula confervacea* (Cleve Gran). *Can. J. Microbiol.*, 8, 229-239, 1962.
- Hecky, R. E., Campbell, P., and Hendzel, L. L.: The stoichiometry of carbon, nitrogen, and phosphorus in particulate matter of lakes and oceans, *Limnol. Oceanogr.*, 38, 709-724, 1993.

- Ho, T. Y., Quigg, A., Finkel, Z. V., Milligan, A. J., Wyman, K., Falkowski, P. G., and Morel, F. M. M.: The elemental composition of some marine phytoplankton, *J. Phycol.*, 39, 1145-1159, doi:10.1111/j.0022-3646.2003.03-090.x, 2003.
- Holligan, P. M., Viollier, M., Harbour, D. S., Camus, P., and Champagnephippe, M.: Satellite and ship studies of coccolithophore production along a continental-shelf edge, *Nature*, 304, 339-342, doi:10.1038/304339a0, 1983.
- Holligan, P. M., Fernandez, E., Aiken, J., Balch, W. M., Boyd, P., Burkill, P. H., Finch, M., Groom, S. B., Malin, G., Muller, K., Purdie, D. A., Robinson, C., Trees, C. C., Turner, S. M., and Vanderwal, P.: A biogeochemical study of the coccolithophore, *Emiliana huxleyi*, in the North Atlantic, *Global Biogeochem. Cy.*, 7, 879-900, doi:10.1029/93gb01731, 1993.
- Jones, R. H., and Flynn, K. J.: Nutritional status and diet composition affect the value of diatoms as copepod prey, *Science*, 307, 1457-1459, doi: 10.1126/science.1107767, 2005.
- Kiefer, D. A.: Modelling growth and light absorption in the marine diatom *Skeletonema costatum*, in: *Towards a Model of Ocean Biogeochemical Processes*, edited by: Evans, G. T. and Fasham, M. J. R., 93-118, Berlin: Springer-Verlag, 1993.
- Klaas, C., and Archer, D. E.: Association of sinking organic matter with various types of mineral ballast in the deep sea: Implications for the rain ratio, *Global Biogeochem. Cy.*, 16, doi:10.1029/2001gb001765, 2002.
- Langer, G., Nehrke, G., Probert, I., Ly, J., and Ziveri, P.: Strain-specific responses of *Emiliana huxleyi* to changing seawater carbonate chemistry, *Biogeosciences*, 6, 2637-2646, 2009.
- Law, C.S., Rickard, G.J., Mikaloff-Fletcher, S.E., Pinkerton, M.H., Gorman, R., Behrens, E., Chiswell, S.M., Bostock, H.C., Anderson, O. and Currie, K.: The New Zealand EEZ and South West Pacific. Synthesis Report RA2, Marine Case Study, in: *Climate Changes, Impacts and Implications (CCII) for New Zealand to 2100*, MBIE contract C01X1225, 40pp, 2016.
- Müller, M. N., Antia, A. N., and LaRoche, J.: Influence of cell cycle phase on calcification in the coccolithophore *Emiliana huxleyi*, *Limnol. Oceanogr.*, 53, 506-512, doi:10.4319/lo.2008.53.2.0506, 2008.

- Müller, M. N., Trull, T. W., and Hallegraeff, G. M.: Differing responses of three Southern Ocean *Emiliana huxleyi* ecotypes to changing seawater carbonate chemistry, *Mar. Ecol. Prog. Ser.*, 531, 81–90, doi:10.3354/meps11309, 2015.
- Paasche, E.: A tracer study of the inorganic carbon uptake during coccolith formation and photosynthesis in the coccolithophorid *Coccolithus huxleyi*, *Physiol. Plant. Suppl.*, 3, 1-82, 1964.
- 5 Paasche, E.: Effect of 3-(p-chlorophenyl)-1,1-dimethylurea (cmu) on photosynthesis and light-dependent coccolith formation in *Coccolithus huxleyi*, *Physiol. Plant.*, 18, 138-145, doi:10.1111/j.1399-3054.1965.tb06876.x, 1965.
- Paasche, E., Brubak, S., Skattebol, S., Young, J. R., and Green, J. C.: Growth and calcification in the coccolithophorid *Emiliana huxleyi* (Haptophyceae) at low salinities, *Phycologia*, 35, 394-403, doi:10.2216/i0031-8884-35-5-394.1, 1996.
- 10 Paasche, E.: Roles of nitrogen and phosphorus in coccolith formation in *Emiliana huxleyi* (Prymnesiophyceae), *Eur. J. Phycol.*, 33, 33-42, doi:10.1017/s0967026297001480, 1998.
- Perry, M. J.: Phosphate utilization by an oceanic diatom in phosphorus-limited chemostat culture and in oligotrophic waters of central North Pacific, *Limnol. Oceanogr.*, 21, 88-107, 1976.
- 15 Polovina, J. J., Howell, E. A., and Abecassis, M.: Ocean's least productive waters are expanding, *Geophys. Res. Lett.*, 35, L03618, doi:10.1029/2007GL031745, 2008.
- Price, N. M.: The elemental stoichiometry and composition of an iron-limited diatom, *Limnol. Oceanogr.*, 50, 1159-1171, 2005.
- 20 Raven, J. A., and Geider, R. J.: Temperature and algal growth, *New Phytol.*, 110, 441-461, doi:10.1111/j.1469-8137.1988.tb00282.x, 1988.
- Raven, J. A., and Crawford, K.: Environmental controls on coccolithophore calcification, *Mar. Ecol. Prog. Ser.*, 470, 137-166, doi:10.3354/meps09993, 2012.
- Read, B. A., Kegel, J., Klute, M. J., Kuo, A., Lefebvre, S. C., Maumus, F., Mayer, C., Miller, J., Monier, A., Salamov, A., Young, J., Aguilar, M., Claverie, J. M., Frickenhaus, S., Gonzalez, K., Herman, E. K., Lin, Y. C., Napier, J., Ogata, H., Sarno, A. F., Shmutz, J., Schroeder, D., de Vargas, C., Verret, F., von Dassow, P., Valentin, K., Van de Peer, Y., Wheeler, G., Dacks, J. B., Delwiche, C. F., Dyhrman, S. T., Glockner, G., John, U., Richards, T., Worden, A. Z., Zhang, X. Y., and Grigoriev, I. V.: Pan genome of

- the phytoplankton *Emiliana* underpins its global distribution, *Nature*, 499, 209-213, doi:10.1038/nature12221, 2013.
- Redfield, A. C., Ketchum, B. H., and Richards, F. A.: The influence of organisms on the composition of sea-water, in: *The Sea. Ideas and observations on progress in the study of the seas. v2. The composition of sea-water. Comparative and descriptive oceanography.*, edited by: Hill, M. N., Interscience, 26-77, 1963.
- Reinfelder, J. R.: Carbon Concentrating Mechanisms in Eukaryotic Marine Phytoplankton, in: *Annual Review of Marine Science*, Vol 3, edited by: Carlson, C. A., and Giovannoni, S. J., *Annu. Rev. Mar. Sci.*, 291-315, 2011.
- 10 Rickard, G. J., Behrens, E., and Chiswell, S. M.: CMIP5 earth system models with biogeochemistry: An assessment for the southwest Pacific Ocean, *J. Geophys. Res. Oceans*, 121, doi:10.1002/2016JC011736, 2016.
- Riebesell, U., Revill, A. T., Holdsworth, D. G., and Volkman, J. K.: The effects of varying CO₂ concentration on lipid composition and carbon isotope fractionation in *Emiliana huxleyi*, *Geochim. Cosmochim. Acta*, 64, 4179-4192, doi:10.1016/s0016-7037(00)00474-9, 2000.
- 15 Riebesell, U., Fabry, V. J., Hansson, L., and Gattuso, J.-P.: Guide to best practices for ocean acidification research and data reporting, Publications Office of the European Union, Luxembourg, 2010.
- Riegman, R., Stolte, W., Noordeloos, A. A. M., and Slezak, D.: Nutrient uptake, and alkaline phosphate (EC 3 : 1 : 3 : 1) activity of *Emiliana huxleyi* (Prymnesiophyceae) during growth under N and P limitation in continuous cultures, *J. Phycol.*, 36, 87-96, doi:10.1046/j.1529-8817.2000.99023.x, 2000.
- Roleda, M. Y., Slocombe, S. P., Leakey, R. J. G., Bell, E. M., and Stanley, M. S.: Effects of temperature and nutrient regimes on biomass and lipid production by six oleaginous microalga in batch culture employing a two-phase cultivation strategy, *Bioresour. Technol.*, 129, 439-449, 25 doi:10.1016/j.biortech.2012.11.043, 2013.
- Rost, B., Riebesell, U., Burkhardt, S., and Sültemeyer, D.: Carbon acquisition of bloom-forming marine phytoplankton, *Limnol. Oceanogr.*, 48, 55-67, 2003.

- Rost, B., and Riebesell, U.: Coccolithophores and the biological pump: Responses to environmental changes, in: Coccolithophores: From Molecular Processes to Global Impact, edited by: Thierstein, H. R., and Young, J. R., 99-125, 2004.
- Rost, B., Riebesell, U., and Sultemeyer, D.: Carbon acquisition of marine phytoplankton: Effect of photoperiod length, *Limnol. Oceanogr.*, 51, 12-20, 2006.
- 5 Sakshaug, E., and Holmhansen, O.: Chemical composition of *Skeletonema costatum* (grev) cleve and pavlova (monochrysis) lutheri (droop) green as a function of nitrate-limited, phosphate-limited, and iron-limited growth, *J. Exp. Mar. Biol. Ecol.*, 29, 1-34, doi:10.1016/0022-0981(77)90118-6, 1977.
- Sardans, J., Rivas-Ubach, A., and Penuelas, J.: The C:N:P stoichiometry of organisms and ecosystems in a changing world: A review and perspectives, *Perspect. Plant Ecol.*, 14, 33-47, doi:10.1016/j.ppees.2011.08.002, 2012.
- 10 Sett, S., Bach, L. T., Schulz, K. G., Koch-Klavsen, S., Lebrato, M., and Riebesell, U.: Temperature modulates coccolithophorid sensitivity of growth, photosynthesis and calcification to increasing seawater $p\text{CO}_2$, *PLoS One*, 9, e88308, doi:10.1371/journal.pone.0088308, 2014.
- 15 Solórzano, L., and Sharp, J. H.: Determination of total dissolved phosphorus and particulate phosphorus in natural waters, *Limnol. Oceanogr.*, 25, 756-760, 1980.
- Sorrosa, J. M., Satoh, M., and Shiraiwa, Y.: Low temperature stimulates cell enlargement and intracellular calcification of Coccolithophorids, *Mar. Biotechnol.*, 7, 128-133, doi:10.1007/s10126-004-0478-1, 2005.
- 20 Sterner, R. W., and Elser, J. J.: *Ecological Stoichiometry: The Biology of Elements from Molecules to Biosphere*, Princeton University Press, Princeton, 2002.
- Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M.: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 25 Stojkovic, S., Beardall, J., and Matear, R.: CO_2 concentrating mechanisms in three southern hemisphere strains of *Emiliania huxleyi*, *J. Phycol.*, 49, 670-679, 2013.

- Toseland, A., Daines, S. J., Clark, J. R., Kirkham, A., Strauss, J., Uhlig, C., Lenton, T. M., Valentin, K., Pearson, G. A., Moulton, V., and Mock, T.: The impact of temperature on marine phytoplankton resource allocation and metabolism, *Nat. Clim. Chang.*, 3, 979-984, doi:10.1038/nclimate1989, 2013.
- van Rijssel, M., and Gieskes, W. W. C.: Temperature, light, and the dimethylsulfoniopropionate (DMSP) content of *Emiliana huxleyi* (Prymnesiophyceae), *J. Sea Res.*, 48, 17-27, doi:10.1016/s1385-1101(02)00134-x, 2002.
- Watabe, N., and Wilbur, K. M.: Effects of temperature on growth calcification and coccolith form in *Coccolithus huxleyi* (coccolithineae), *Limnol. Oceanogr.*, 11, 567-575, 1966.
- Westbroek, P., Brown, C. W., Vanbleijswijk, J., Brownlee, C., Brummer, G. J., Conte, M., Egge, J., Fernandez, E., Jordan, R., Knappertsbusch, M., Stefels, J., Veldhuis, M., Vanderwal, P., and Young, J.: A model system approach to biological climate forcing - the example of *Emiliana huxleyi*, *Global Planet. Change*, 8, 27-46, doi:10.1016/0921-8181(93)90061-r, 1993.
- Welschmeyer, N. A. Fluorometric analysis of chlorophyll *a* in the presence of chlorophyll *b* and pheopigments. *Limnol. Oceanogr.* 39, 1985-1992, 1994.
- Young, J. R., Poulton, A. J., and Tyrrell, T.: Morphology of *Emiliana huxleyi* coccoliths on the North West European shelf - is there an influence of carbonate chemistry?, *Biogeosciences Discussions*, 11, 4531-4561, 2014.
- Zondervan, I., Zeebe, R. E., Rost, B., and Riebesell, U.: Decreasing marine biogenic calcification: A negative feedback on rising atmospheric $p\text{CO}_2$, *Global Biogeochem. Cy.*, 15, 507-516, doi:10.1029/2000gb001321, 2001.
- Zondervan, I., Rost, B., and Riebesell, U.: Effect of CO_2 concentration on the PIC/POC ratio in the coccolithophore *Emiliana huxleyi* grown under light-limiting conditions and different daylengths, *J. Exp. Mar. Biol. Ecol.*, 272, 55-70, doi:10.1016/s0022-0981(02)00037-0, 2002.
- Zondervan, I.: The effects of light, macronutrients, trace metals and CO_2 on the production of calcium carbonate and organic carbon in coccolithophores - a review, *Deep-Sea Res. PT II*, 54, 521-537, doi:10.1016/j.dsr2.2006.12.004, 2007.




Figure 1. Changes in *Emiliania huxleyi* cellular particulate organic carbon (POC) content in response to different environmental drivers: a) cellular POC vs. nitrate concentration; b) cellular POC vs. phosphate concentration; c) cellular POC vs. irradiance; d) cellular POC vs. temperature; and e) cellular POC vs. pCO₂. Error bars represent standard deviations (n=3).

5 For Figures 1-5: The dashed lines represent the fitted dose-response curves. “Φ” represent the fitted values for the present day conditions in the Chatham Rise area, and “X” represent the fitted values for the predicted future conditions (2100) in the Chatham Rise area. Error bars represent standard deviation (n=3).




Figure 2. Changes in *Emiliania huxleyi* cellular particulate inorganic carbon (PIC) content in response to different environmental drivers: a) cellular PIC vs. nitrate concentration; b) cellular PIC vs. phosphate concentration; c) cellular PIC vs. irradiance; d) cellular PIC vs. temperature; and e) cellular PIC vs. $p\text{CO}_2$.




Figure 3. Changes in the ratio of *Emiliana huxleyi* cellular particulate inorganic carbon content to particulate organic carbon content (PIC:POC) in response to different environmental drivers: a) PIC:POC ratio vs. nitrate concentration; b) PIC:POC ratio vs. phosphate concentration; c) PIC:POC ratio vs. irradiance; d) PIC:POC ratio vs. temperature; and e) PIC:POC ratio vs. $p\text{CO}_2$.




Figure 4. Changes in *Emiliana huxleyi* cellular particulate organic nitrogen (PON) content in response to different environmental drivers: a) cellular PON vs. nitrate concentration; b) cellular PON vs. phosphate concentration; c) cellular PON vs. irradiance; d) cellular PON vs. temperature; and e) cellular PON vs. $p\text{CO}_2$.




Figure 5. Changes in *Emiliania huxleyi* cellular particulate organic phosphorus (POP) content in response to different environmental drivers: a) cellular POP vs. nitrate concentration; b) cellular POP vs. phosphate concentration; c) cellular POP vs. irradiance; d) cellular POP vs. temperature; and e) cellular POP vs. $p\text{CO}_2$.




Figure 6. Changes in the ratio of *Emiliana huxleyi* cellular particulate organic carbon content to chlorophyll *a* content (C:Chl-*a*) in response to different environmental drivers: a) C:Chl-*a* ratio vs. nitrate concentration; b) C:Chl-*a* ratio vs. phosphate concentration; c) C:Chl-*a* ratio vs. irradiance; d) C:Chl-*a* ratio vs. temperature; and e) C:Chl-*a* ratio vs. $p\text{CO}_2$. Error bars represent standard deviations ($n=3$).

5




Figure 7. Conceptual figure of the specific effects of each the five environmental drivers, under the projected future conditions (year 2100), on the elemental composition of *Emiliana huxleyi*. Q represents the cellular quota of each element of *Emiliana huxleyi*.

5 ***The box denotes the *E. huxleyi* cell. Solid blue arrows indicate positive effects of the future environmental changes; and dash red arrows indicate negative effects of the future environmental changes. Arrows in bold indicate the environmental drivers that play the most important role regulating the connected physiological metrics under the predicted environmental conditions for the year 2100.

Table 1. Treatment conditions for each environmental driver used in the manipulation experiments.

Environmental driver	Treatments
Nitrate (μM)	3.7; 6; 12; 48; 96; 200
Phosphate (μM)	0.4; 2; 6; 10; 20
Irradiance ($\mu\text{mol photons m}^{-2} \text{s}^{-1}$)	14; 40; 80; 190; 350; 650
Temperature ($^{\circ}\text{C}$)	4; 7; 11; 15; 20; 25
$p\text{CO}_2$ (Pa)	8; 15; 39; 58; 74; 109

Table 2. Elemental molar ratios of N:P, C:N and C:P of *Emiliana huxleyi* from the five single-factorial manipulation experiments. The errors are standard deviations around the mean (n=3). The values in bold are significantly different compared to other treatments.

Environmental driver	Treatment	N:P (mol:mol)	C:N (mol:mol)	C:P (mol:mol)
Nitrate (μM)	3.7	9.09±2.48	15.90±4.09	137.69±8.27
	6	9.07±1.55	13.16±1.13	118.45±10.06
	12	15.90±1.38	9.00±0.07	143.08±11.28
	48	14.46±1.39	7.01±0.64	107.76±11.93
	96	16.16±3.45	6.56±1.07	103.56±6.54
	200	14.31±0.40	7.32±0.47	104.82±8.34
Phosphate (μM)	0.4	41.57±4.14	6.32±0.42	261.67±17.30
	2	21.47±0.28	9.27±5.08	137.57±11.31
	6	14.17±3.59	6.16±1.24	85.00±4.51
	10	17.06±2.06	5.24±0.06	89.28±9.80
	20	13.99±0.89	5.22±0.08	73.04±3.48
Irradiance ($\mu\text{mol photons m}^{-2} \text{ s}^{-1}$)	14	22.03±7.27	4.26±0.29	73.82±6.48
	40	14.27**	4.38**	62.50**
	80	16.03±2.90	5.24±0.69	83.96±17.05
	190	18.71±1.74	5.99±0.23	112.39±14.60
	350	17.15±0.83	6.47±0.54	110.63±3.98
	650	16.11±1.77	5.70±0.31	91.49±6.66
Temperature ($^{\circ}\text{C}$)	4	8.92±1.29	8.67±2.64	71.34±27.92
	7	10.46±2.05	7.60±1.32	78.70±14.78
	11	13.58±1.91	6.21±0.32	86.56±11.92
	15	14.12±0.66	6.70±0.31	94.47±3.84
	20	15.53±1.06	5.98±0.15	92.88±6.33
	25	13.67±2.99	7.08±1.39	93.96±4.87
$p\text{CO}_2$ (Pa)	8	19.39±2.41	6.81±1.09	122.61±9.97
	15	24.01±6.80	5.80±0.71	137.50±33.51
	39	16.96±3.62	7.41±0.06	155.64±31.42
	58	17.89±0.80	6.55±1.16	116.67±16.15
	74	18.22±2.45	6.42±0.43	116.93±17.60
	109	13.56±2.78	7.41±1.20	99.25±17.70

** Sample loss during analysis resulted in only single values at this irradiance.

Table 3. Comparison of cellular particulate organic carbon (POC) contents, particulate inorganic carbon (PIC) contents, PIC:POC ratios, particulate organic nitrogen (PON) contents and particulate organic phosphorus (POP) contents of *Emiliania huxleyi* between projected (year 2100) and present day Chatham Rise conditions, with rankings of the importance of the environmental drivers which caused significant effects on each physiological parameter. The numbers of the ranking scheme represent the gradient of the most (1) to least (4) important effects. Effect of “+” represents an increase and “-“ represents a decrease in the elemental composition/ratio in the future, respectively.

Physiological parameter	Environmental driver	Fitted values at different conditions of environmental drivers*		Future vs. present day comparisons		
		Present day	Future	Change (%)**	Effects (+/-)	Ranking
Cellular POC content (pg cell ⁻¹)	Temperature	10.798	9.713	10.0	-	1*
	CO ₂	14.632	15.436	5.5	+	2
	Irradiance	14.774	14.827	0.3	+	3
	Nitrate					n.s.
	Phosphate					n.s.
Cellular PIC content (pg cell ⁻¹)	Temperature	10.206	8.753	14.2	-	1
	Nitrate					n.s.
	Phosphate					n.s.
	Irradiance					n.s.
	CO ₂					n.s.
PIC:POC	CO ₂	0.868	0.821	5.4	-	1
	Temperature	1.017	1.042	2.4	+	2
	Irradiance	0.777	0.780	0.3	+	3
	Nitrate					n.s.
	Phosphate					n.s.
Cellular PON content (pg cell ⁻¹)	Nitrate	1.380	1.162	15.8	-	1
	Temperature	2.013	1.819	9.6	-	2
	Phosphate					n.s.
	Irradiance					n.s.
	CO ₂					n.s.
Cellular POP content (pg cell ⁻¹)	Phosphate	0.106	0.078	25.9	-	1
	Temperature	0.304	0.269	11.6	-	2
	CO ₂	0.312	0.342	9.6	+	3
	Nitrate	0.249	0.227	8.9	-	4
	Irradiance					n.s.

*The fitted values for “present day” and “future” were extracted from the fitted dose-response curves (Fig.1 - 5) at the stock culture growing conditions, average present day conditions in the Chatham Rise area, and the predicted future conditions (2100) of Chatham Rise, respectively.

** The percentage changes were calculated as the changes caused by each environmental driver under the future predicted condition relative to that under the present day condition.

*Numbers in bold indicate statistically significant difference between the range of present day and future conditions (nitrate treatments: 6.0 and 12.0 μM; phosphate treatments: 0.4 and 2 μM; irradiance treatments: 80 and 190 μmol photons m⁻² s⁻¹; temperature treatments: 11°C, 15°C and 20°C) based on the one-way ANOVA. “n.s.” indicates non-significant difference (one-way ANOVA) among all the treatments used for the fitting.