

We would like to thank the editor for the critical comments, which we think helped to improve the quality and clarity of this manuscript. We hope our responses and adaptations are adequate to accept this manuscript for publication in Biogeosciences. Please find our detailed responses below.

Associate Editor Comment (21. September 2017)

Dear Authors,

I went through your new version of manuscript and appreciate the modifications you made in order to clarify presentation of your ideas, moderate some statements and delimit limits of your study. However, I would ask you to make an effort to synthesize your discussions by shortening the sections that are viewed as speculative by the two referees. This important effort will avoid a dilution of the main results of your study in speculative discussions using the conditional form.

Regards,

Sébastien Fontaine

Authors Reply: We agree with the editor/reviewers that the discussion was, in some parts, too repetitive and speculative. In order to follow your suggestions, following adaptations were made:

- 1. We drastically shortened the first part of the discussion (4.1.) to reduce the amount of repetition from the results and the amount of unwarranted speculation.**
- 2. In the discussion part 4.3, we deleted the paragraph about our calculation of deep methanogenesis, which was criticized of being too speculative. As we did not actively measure deep methanogenesis in the present study, we agree that this part might go above the scope of our research focus. We further deleted comparisons with other environments (including Table 2) in this chapter, to stay focused on the study site.**
- 3. In part 4.3, we were able to add proof to our hypothesis of close coupling between AOM and methanogenesis in surface sediments, which decreases the degree of speculation of this statement. This cryptic methane cycling has been recently demonstrated in labeling incubations with surface sediments from the close-by Aarhus Bay, Denmark (Xiao et al, 2017), and thus could very likely occur in sediments from Eckernförde Bay.**
- 4. We shortened the summary part to decrease unnecessary repetition.**
- 5. We further decided to provide a more concise definition of "surface methanogenesis" by introducing the term "SRZ methanogenesis" (i.e. methanogenesis in the sulfate reduction zone) opposite to ("deep") methanogenesis below the sulfate methane transition zone (SMTZ). We think these terms are more scientifically correct and avoid confusion with the term "sediment surface" in general.**

1 Microbial methanogenesis in the sulfate-reducing zone ~~in~~from ~~of~~in the sediments
2 from in the Eckernförde Bay, SW Baltic Sea

3 Johanna Maltby^{a,b*}, Lea Steinle^{c,a}, Carolin R. Löscher^{d,a}, Hermann W. Bange^a, Martin A. Fischer^e, Mark
4 Schmidt^a, Tina Treude^{f,g*}

5 ^a *GEOMAR Helmholtz Centre for Ocean Research Kiel, Department of Marine Biogeochemistry, 24148*
6 *Kiel, Germany*

7 ^b *Present Address: Natural Sciences Department, Saint Joseph's College, Standish, Maine 04084, USA*

8 ^c *Department of Environmental Sciences, University of Basel, 4056 Basel, Switzerland*

9 ^d *Nordic Center for Earth Evolution, University of Southern Denmark, 5230 Odense, Denmark*

10 ^e *Institute of Microbiology, Christian-Albrecht-University Kiel, 24118 Kiel, Germany*

11 ^f *Department of Earth, Planetary, and Space Sciences, University of California Los Angeles (UCLA), Los*
12 *Angeles, California 90095-1567, USA*

13 ^g *Department of Atmospheric and Oceanic Sciences, University of California Los Angeles (UCLA), Los*
14 *Angeles, California 90095-1567, USA*

15
16 *Correspondence: jmaltby@sjcme.edu, ttreude@g.ucla.edu

27 Abstract

28 Benthic microbial methanogenesis is a known source of methane in marine systems. In most
29 sediments, the majority of methanogenesis is located below the sulfate-reducing zone, as sulfate
30 reducers outcompete methanogens for the major substrates hydrogen and acetate. Coexistence of
31 methanogenesis and sulfate reduction has been shown before and is possible by usage of non-
32 competitive substrates by the methanogens such as methanol or methylated amines. However, the
33 knowledge about magnitude, seasonality and environmental controls on this non-competitive
34 methane production is sparse. In the present study, the presence of [surface-methanogenesis \(0-30](#)
35 [centimeters below seafloor, cmbsf\)](#), here defined as methanogenesis within the [sulfate-rich sulfate](#)
36 [reduction zone \(SRZ methanogenesis\)](#), was investigated in sediments [\(0-30 centimeters below](#)
37 [seafloor, cmbsf\)](#) of the seasonally hypoxic Eckernförde Bay, southwestern Baltic Sea. Water column
38 parameters ~~like such as~~ oxygen, temperature, and salinity together with porewater geochemistry and
39 benthic methanogenesis rates were determined in the sampling area “Boknis Eck” quarterly from
40 March 2013 to September 2014, to investigate the effect of seasonal environmental changes on the
41 rate and distribution of [surface-SRZ](#) methanogenesis, to estimate its potential contribution to benthic
42 methane emissions, and to identify potential methanogenic groups responsible for [surface-methane](#)
43 [production-SRZ methanogenesis](#). The metabolic pathway of methanogenesis in the presence or
44 absence of sulfate reducers and after the addition of a non-competitive substrate was studied in four
45 experimental setups: 1) unaltered sediment batch incubations (net methanogenesis), 2) ¹⁴C-
46 bicarbonate labeling experiments (hydrogenotrophic methanogenesis), 3) manipulated experiments
47 with addition of either molybdate (sulfate reducer inhibitor), 2-bromoethane-sulfonate (methanogen
48 inhibitor), or methanol (non-competitive substrate, potential methanogenesis), 4) addition of ¹³C-
49 labeled methanol (potential methylotrophic methanogenesis). After incubation with methanol,
50 molecular analyses were conducted to identify key functional methanogenic groups during
51 methylotrophic methanogenesis. To also compare magnitudes of [surface-SRZ](#) methanogenesis with
52 [deep](#)-methanogenesis below the sulfate reduction zone (> 30 cmbsf), hydrogenotrophic
53 methanogenesis was determined by ¹⁴C-bicarbonate radiotracer incubation in samples collected in
54 September 2013.

55 [Surface-SRZ](#) methanogenesis changed seasonally in the upper 30 cmbsf with rates increasing from
56 March (0.2 nmol cm⁻³ d⁻¹) to November (1.3 nmol cm⁻³ d⁻¹) 2013 and March (0.2 nmol cm⁻³ d⁻¹) to
57 September (0.4 nmol cm⁻³ d⁻¹) 2014, respectively. Its magnitude and distribution appeared to be
58 controlled by organic matter availability, C/N, temperature, and oxygen in the water column,
59 revealing higher rates in warm, stratified, hypoxic seasons (September/November) compared to
60 colder, oxygenated seasons (March/June) of each year. The majority of [surface-SRZ](#) methanogenesis
61 was likely driven by the usage of non-competitive substrates (e.g., methanol and methylated

62 compounds), to avoid competition with sulfate reducers, as it was indicated by the 1000-3000-fold
63 increase in potential methanogenesis activity observed after methanol addition. Accordingly,
64 competitive hydrogenotrophic methanogenesis increased in the sediment only below the depth of
65 sulfate penetration (> 30 cmbsf). Members of the family *Methanosarcinaceae*, which are known for
66 methylotrophic methanogenesis, were detected by PCR using *Methanosarcinaceae*-specific primers
67 and are likely to be responsible for the observed [surfaceSRZ](#) methanogenesis.
68 The present study indicates that [surfaceSRZ](#) methanogenesis is an important component of the
69 benthic methane budget and carbon cycling in Eckernförde Bay. Although its contribution to
70 methane emissions from the sediment into the water column are probably minor, [surfaceSRZ](#)
71 methanogenesis could directly feed into methane oxidation above the sulfate-methane transition
72 zone.

73 1. Introduction

74 After water vapor and carbon dioxide, methane is the most abundant greenhouse gas in the
75 atmosphere (e.g. Hartmann et al., 2013; Denman et al., 2007). Its atmospheric concentration
76 increased more than 150 % since preindustrial times, mainly through increased human activities such
77 as fossil fuel usage and livestock breeding (Hartmann et al., 2013; Wuebbles & Hayhoe, 2002;
78 Denman et al., 2007). Determining the natural and anthropogenic sources of methane is one of the
79 major goals for oceanic, terrestrial and atmospheric scientists to be able to predict further impacts
80 on the world's climate. The ocean is considered to be a modest natural source for atmospheric
81 methane (Wuebbles & Hayhoe, 2002; Reeburgh, 2007; EPA, 2010). However, research is still sparse
82 on the origin of the observed oceanic methane, which automatically leads to uncertainties in current
83 ocean flux estimations (Bange et al., 1994; Naqvi et al., 2010; Bakker et al., 2014).

84 Within the marine environment, the coastal areas (including estuaries and shelf regions) are
85 considered the major source for atmospheric methane, contributing up to 75 % to the global ocean
86 methane production (Bange et al., 1994). The major part of the coastal methane is produced during
87 microbial methanogenesis in the sediment, with probably only a minor part originating from
88 methane production within the water column (Bakker et al., 2014). However, the knowledge on
89 magnitude, seasonality and environmental controls of benthic methanogenesis is still limited.
90 In marine sediments, methanogenesis activity is mostly restricted to the sediment layers below
91 sulfate reduction, due to the successful competition of sulfate reducers with methanogens for the
92 mutual substrates acetate and hydrogen (H₂) (Oremland & Polcin, 1982; Crill & Martens, 1986;
93 Jørgensen, 2006). Methanogens produce methane mainly from using acetate (acetoclastic
94 methanogenesis) or H₂ and carbon dioxide (CO₂) (hydrogenotrophic methanogenesis). Competition
95 with sulfate reducers can be relieved through usage of non-competitive substrates (e.g. methanol or

96 methylated compounds, methylotrophic methanogenesis) (Cicerone & Oremland, 1988; Oremland &
97 Polcin, 1982). Coexistence of sulfate reduction and methanogenesis has been detected in a few
98 studies from organic-rich sediments, e.g., salt-marsh sediments (Oremland et al., 1982; Buckley et al.,
99 2008), coastal sediments (Holmer & Kristensen, 1994; Jørgensen & Parkes, 2010) or sediments in
100 upwelling regions (Pimenov et al., 1993; Ferdelman et al., 1997; Maltby et al., 2016), indicating the
101 importance of these environments for [surface-methanogenesis within the sulfate reduction zone](#)
102 [\(SRZ methanogenesis\)](#). So far, however, environmental controls of [surfaceSRZ](#) methanogenesis
103 remain elusive.

104 The coastal inlet Eckernförde Bay (southwestern Baltic Sea) is an excellent model environment to
105 study seasonal and environmental controls of benthic [surfaceSRZ](#) methanogenesis. Here, the muddy
106 sediments are characterized by high organic loading and high sedimentation rates (Whiticar, 2002),
107 which lead to anoxic conditions within the uppermost 0.1-0.2 centimeter below seafloor (cmbsf)
108 (Preisler et al., 2007). Seasonally hypoxic (dissolved oxygen < 63 µM) and anoxic (dissolved oxygen =
109 0 µM) events in the bottom water of Eckernförde Bay (Lennartz et al., 2014) provide ideal conditions
110 for anaerobic processes at the sediment surface.

111 Sulfate reduction is the dominant pathway of organic carbon degradation in Eckernförde Bay
112 sediments in the upper 30 cmbsf, followed by methanogenesis in deeper sediment layers where
113 sulfate is depleted (> 30 cmbsf) (Whiticar 2002; Treude et al. 2005; Martens et al. 1998) (Fig. 1). This
114 [deep-methanogenesis below the sulfate-methane transition zone \(SMTZ\)](#) can be intense and often
115 leads to methane oversaturation in the porewater below 50 cm sediment depth, resulting in gas
116 bubble formation (Abegg & Anderson, 1997; Whiticar, 2002; Thießen et al., 2006). Thus, methane is
117 transported from the methanogenic zone (> 30 cmbsf) to the surface sediment by both molecular
118 diffusion and advection via rising gas bubbles (Wever et al., 1998; Treude et al., 2005a). Although
119 upward diffusing methane is mostly retained by anaerobic oxidation of methane (AOM) (Treude et
120 al. 2005), a major part is reaching the sediment-water interface through gas bubble transport
121 (Treude et al. 2005; Jackson et al. 1998), resulting in a supersaturation of the water column with
122 respect to atmospheric methane concentrations (Bange et al., 2010). The Time Series Station “Boknis
123 Eck” in the Eckernförde Bay is a known site of methane emissions into the atmosphere throughout
124 the year due to this supersaturation of the water column (Bange et al., 2010).

125 The source for benthic and water column methane was seen in [deep-methanogenesis below the](#)
126 [SMTZ \(> 30 cmbsf\) below the penetration of sulfate](#) (Whiticar, 2002), however, coexistence of
127 sulfate reduction and methanogenesis has been postulated (Whiticar, 2002; Treude et al., 2005a).
128 Still, the magnitude and environmental controls of [surfaceSRZ](#) methanogenesis is poorly understood,
129 even though it may make a measurable contribution to benthic methane emissions given its short
130 diffusion distance to the sediment-water interface (Knittel & Boetius, 2009). Production of methane

131 within the sulfate reduction zone of Eckernförde Bay [surface](#)-sediments could further explain peaks
132 of methane oxidation observed in top sediment layers, which was previously attributed to methane
133 transported to the [sediment](#) surface via rising gas bubbles (Treude et al., 2005a).
134 In the present study, we investigated [surface](#)-sediments [from within](#) (< 30 cmbsf, on a seasonal
135 basis), ~~deep sediment~~ [and below the sulfate reduction zone](#) (> 30 cmbsf, on one occasion), and the
136 water column (on a seasonal basis) at the Time Series Station "Boknis Eck" in Eckernförde Bay, to
137 validate the existence of [surfaceSRZ](#) methanogenesis and its potential contribution to benthic
138 methane emissions. Water column parameters like oxygen, temperature, and salinity together with
139 porewater geochemistry and benthic methanogenesis were measured over a course of 2 years. In
140 addition to seasonal rate measurements, inhibition and stimulation experiments, stable isotope
141 probing, and molecular analysis were carried out to find out if [surfaceSRZ](#) methanogenesis 1) is
142 controlled by environmental parameters, 2) shows seasonal variability, 3) is based on non-
143 competitive substrates with a special focus on methylotrophic methanogens.

144 2. Material and Methods

145 2.1 Study site

146 Samples were taken at the Time Series Station "Boknis Eck" (BE, 54°31.15 N, 10°02.18 E;
147 www.bokniseck.de) located at the entrance of Eckernförde Bay in the southwestern Baltic Sea with a
148 water depth of about 28 m (map of sampling site can be found in e.g. Hansen et al., (1999)). From
149 mid of March until mid of September the water column is strongly stratified due to the inflow of
150 saltier North Sea water and a warmer and fresher surface water (Bange et al., 2011). Organic matter
151 degradation in the deep layers causes pronounced hypoxia (March-Sept) or even anoxia
152 (August/September) (Smetacek, 1985; Smetacek et al., 1984). The source of organic material is
153 phytoplankton blooms that occur regularly in spring (February-March) and fall (September-
154 November) and are followed by pronounced sedimentation of organic matter (Bange et al., 2011). To
155 a lesser extent, phytoplankton blooms and sedimentation are also observed during the summer
156 months (July/August) (Smetacek et al., 1984). Sediments at BE are generally classified as soft, fine-
157 grained muds (< 40 µm) with a carbon content of 3 to 5 wt% (Balzer et al., 1986). The bulk of organic
158 matter in Eckernförde Bay sediments originates from marine plankton and macroalgal sources (Orsi
159 et al., 1996), and its degradation leads to production of free methane gas (Wever & Fiedler, 1995;
160 Abegg & Anderson, 1997; Wever et al., 1998). The oxygen penetration depth is limited to the upper
161 few millimeters when bottom waters are oxic (Preisler et al., 2007). Reducing conditions within the
162 sulfate reduction zone lead to a dark grey/black sediments color with a strong hydrogen sulfur odor
163 in the upper meter of the sediment and dark olive-green color the deeper sediment layers (> 1 m)
164 (Abegg & Anderson, 1997).

165 **2.2 Water column and sediment sampling**

166 Sampling was done on a seasonal basis during the years of 2013 and 2014. One-Day field trips with
167 either F.S. Alkor (cruise no. AL410), F.K. Littorina or F.B. Polarfuchs were conducted in March, June,
168 and September of each year. In 2013, additional sampling was conducted in November. At each
169 sampling month, water profiles of temperature, salinity, and oxygen concentration (optical sensor,
170 RINKO III, detection limit= 2 μ M) were measured with a CTD (Hydro-Bios). In addition, water samples
171 for methane concentration measurements were taken at 25 m water depth with a 6-Niskin bottle (4
172 Liter each) rosette attached to the CTD (Table 1). Complementary samples for water column
173 chlorophyll were taken at 25 m water depth with the CTD-rosette within the same months during
174 standardized monthly sampling cruises to Boknis Eck organized by GEOMAR.

175 Sediment cores were taken with a miniature multicorer (MUC, K.U.M. Kiel), holding 4 core liners
176 (length= 60 cm, diameter= 10 cm) at once. The cores had an average length of \sim 30 cm and were
177 stored at 10°C in a cold room (GEOMAR) until further processing (normally within 1-3 days after
178 sampling).

179 In September 2013, a gravity core was taken in addition to the MUC cores. The gravity core was
180 equipped with an inner plastic bag (polyethylene; diameter: 13 cm). After core recovery (330 cm
181 total length), the polyethylene bag was cut open at 12 different sampling depths resulting in intervals
182 of 30 cm and sampled directly on board for sediment porewater geochemistry (see Sect. 2.4),
183 sediment methane (see Sect. 2.5), sediment solid phase geochemistry (see Sect. 2.6), and microbial
184 rate measurements for hydrogenotrophic methanogenesis as described in section 2.8.

185 **2.3 Water column parameters**

186 At each sampling month, water samples for methane concentration measurements were taken at 25
187 m water depth in triplicates. Therefore, three 25 ml glass vials were filled bubble free directly after
188 CTD-rosette recovery and closed with butyl rubber stoppers. Biological activity in samples was
189 stopped by adding -saturated mercury chloride solution, followed by storage- at room temperature
190 until further treatment.

191 Concentrations of dissolved methane (CH_4) were determined by headspace gas chromatography as
192 described in Bange et al. (2010). Calibration for CH_4 was done by a two-point calibration with known
193 methane concentrations before the measurement of headspace gas samples, resulting in an error of
194 < 5 %.

195 Water samples for chlorophyll concentration were taken by transferring the complete water volume
196 (from 25 m water depth) from one water sampler into a 4.5 L Nalgene bottle, from which then
197 approximately 0.7-1 L (depending on the plankton content) were filtrated back in the GEOMAR
198 laboratory using GF/F filter (Whatman, 25 mm diameter, 8 μ M pores size). Dissolved chlorophyll a

199 concentrations were determined using the fluorometric method by Welschmeyer (1994) with an
200 error < 10 %.

201 **2.4 Sediment porewater geochemistry**

202 Porewater was extracted from sediment within 24 hours after core retrieval using nitrogen (N₂) pre-
203 flushed rhizons (0.2 µm, Rhizosphere Research Products, Seeberg-Elverfeldt et al., 2005). In MUC
204 cores, rhizons were inserted into the sediment in 2 cm intervals through pre-drilled holes in the core
205 liner. In the gravity core, rhizons were inserted into the sediment in 30 cm intervals directly after
206 retrieval.

207 Extracted porewater from MUC and gravity cores was immediately analyzed for sulfide using
208 standardized photometric methods (Grasshoff et al., 1999).

209 Sulfate concentrations were determined using ion chromatography (Methrom 761). Analytical
210 precision was < 1 % based on repeated analysis of IAPSO seawater standards (dilution series) with an
211 absolute detection limit of 1 µM corresponding to a detection limit of 30 µM for the undiluted
212 sample.

213 For analysis of dissolved inorganic carbon (DIC), 1.8 ml of porewater was transferred into a 2 ml glass
214 vial, fixed with 10 µl saturated HgCl₂ solution and crimp sealed. DIC concentration was determined
215 as CO₂ with a multi N/C 2100 analyzer (Analytik Jena) following the manufacturer's instructions.
216 Therefore, the sample was acidified with phosphoric acid and the outgassing CO₂ was measured. The
217 detection limit was 20 µM with a precision of 2-3 %.

218 **2.5 Sediment methane concentrations**

219 In March 2013, June 2013 and March 2014, one MUC core was sliced in 1 cm intervals until 6 cmbsf,
220 followed by 2 cm intervals until the end of the core. At the other sampling months, the MUC core
221 was sliced in 1 cm intervals until 6 cmbsf, followed by 2 cm intervals until 10 cmbsf and 5 cm intervals
222 until the end of the core.

223 Per sediment depth (in MUC and gravity cores), 2 cm⁻³ of sediment were transferred into a 10 ml-
224 glass vial containing 5 ml NaOH (2.5 %) for determination of sediment methane concentration per
225 volume of sediment. The vial was quickly closed with a butyl septum, crimp-sealed and shaken
226 thoroughly. The vials were stored upside down at room temperature until measurement via gas
227 chromatography. Therefore, 100 µl of headspace was removed from the gas vials and injected into a
228 Shimadzu gas chromatograph (GC-2014) equipped with a packed Haysep-D column and a flame
229 ionization detector. The column temperature was 80°C and the helium flow was set to 12 ml min⁻¹.

230 CH₄ concentrations were calibrated against CH₄ standards (Scotty gases). The detection limit was 0.1
231 ppm with a precision of 2 %.

232 **2.6 Sediment solid phase geochemistry**

233 Following the sampling for CH₄, the same cores described under section 2.5 were used for the
234 determination of the sediment solid phase geochemistry, i.e. porosity, particulate organic carbon
235 (POC) and particulate organic nitrogen (PON).

236 Sediment porosity of each sampled sediment section was determined by the weight difference of 5
237 cm³ wet sediment after freeze-drying for 24 hours. Dried sediment samples were then used for
238 analysis of particulate organic carbon (POC) and particulate organic nitrogen (PON) with a Carlo-Erba
239 element analyzer (NA 1500). The detection limit for C and N analysis was < 0.1 dry weight percent (%)
240 with a precision of < 2 %.

241 **2.7 Sediment methanogenesis**

242 **2.7.1 Methanogenesis in MUC cores**

243 At each sampling month, three MUC cores were sliced in 1 cm intervals until 6 cmbsf, in 2 cm
244 intervals until 10 cmbsf, and in 5 cm intervals until the bottom of the core. Every sediment layer was
245 transferred to a separate beaker and quickly homogenized before sub-sampling. The exposure time
246 with air, i.e. oxygen, was kept to a minimum. Sediment layers were then sampled for determination
247 of net methanogenesis (defined as the sum of total methane production and consumption, including
248 all available methanogenic substrates in the sediment), hydrogenotrophic methanogenesis
249 (methanogenesis based on the substrates CO₂/H₂), and potential methanogenesis (methanogenesis
250 at ideal conditions, i.e. no lack of nutrients) as described in the following sections.

251 *2.7.1.1 Net methanogenesis*

252 Net methanogenesis was determined with sediment slurry experiments by measuring the headspace
253 methane concentration over time. Per sediment layer, triplicates of 5 cm³ of sediment were
254 transferred into N₂-flushed sterile glass vials (30 ml) and mixed with 5 ml filtered bottom water. The
255 slurry was repeatedly flushed with N₂ to remove residual methane and to ensure complete anoxia.
256 Slurries were incubated in the dark at in-situ temperature, which varied at each sampling date (Table
257 1). Headspace samples (0.1 ml) were taken out every 3-4 days over a time period of 4 weeks and
258 analyzed on a Shimadzu GC-2104 gas chromatograph (see Sect. 2.5). Net methanogenesis rates were
259 determined by the linear increase of the methane concentration over time (minimum of 6 time
260 points, see also Fig. S1).

261 *2.7.1.2 Hydrogenotrophic methanogenesis*

262 To determine if hydrogenotrophic methanogenesis, i.e., methanogenesis based on the competitive
263 substrates H₂, is present in the sulfate-reducing zone, radioactive sodium bicarbonate (NaH¹⁴CO₃)
264 was added to the sediment.

265 Per sediment layer, sediment was sampled in triplicates with glass tubes (5 mL) which were closed
266 with butyl rubber stoppers on both ends according to (Treude et al. 2005). Through the stopper,
267 $\text{NaH}^{14}\text{CO}_3$ (dissolved in water, injection volume 6 μl , activity 222 kBq, specific activity = 1.85-2.22
268 GBq/mmol) was injected into each sample and incubated for three days in the dark at in-situ
269 temperature (Table 1). To stop bacterial activity, sediment was transferred into 50 ml glass-vials filled
270 with 20 ml sodium hydroxide (2.5 % w/w), closed quickly with rubber stoppers and shaken
271 thoroughly. Five controls were produced from various sediment depths by injecting the radiotracer
272 directly into the NaOH with sediment.

273 The production of ^{14}C -methane was determined with the slightly modified method by Treude et al.,
274 (2005) used for the determination of anaerobic oxidation of methane. The method was identical,
275 except no unlabeled methane was determined by gas chromatography. Instead, DIC values were
276 used to calculate hydrogenotrophic methane production.

277 2.7.1.3 Potential methanogenesis in manipulated experiments

278 To examine the interaction between sulfate reduction and methanogenesis, inhibition and
279 stimulation experiments were carried out. Therefore, every other sediment layer was sampled
280 resulting in the following examined six sediment layers: 0-1 cm, 2-3 cm, 4-5 cm, 6-8 cm, 10-15 cm
281 and 20-25 cm. From each layer, sediment slurries were prepared by mixing 5 ml sediment in a 1:1
282 ratio with adapted artificial seawater medium (salinity 24, Widdel & Bak, 1992) in N_2 -flushed, sterile
283 glass vials before further manipulations.

284 In total, four different treatments, each in triplicates, were prepared per depth: 1) with sulfate
285 addition (17 mM), 2) with sulfate (17 mM) and molybdate (22 mM) addition, 3) with sulfate (17 mM)
286 and 2-bromoethane-sulfonate (BES, 60 mM) addition, and 4) with sulfate (17 mM) and methanol (10
287 mM) addition. From here on, the following names are used to describe the different treatments,
288 respectively: 1) control treatment, 2) molybdate treatment, 3) BES treatment, and 4) methanol
289 treatment. Control treatments feature the natural sulfate concentrations occurring in [surface](#)
290 [sediments of the sulfate reduction zone of-at](#) the sampling site. Molybdate was used as an enzymatic
291 inhibitor for sulfate reduction (Oremland & Capone, 1988) and BES was used as an inhibitor for
292 methanogenic archaea (Hoehler et al., 1994). Methanol is a known non-competitive substrate, which
293 is used by methanogens but not by sulfate reducers (Oremland & Polcin, 1982), thus it is suitable to
294 examine non-competitive methanogenesis. Treatments were incubated similar to net
295 methanogenesis (2.7.1.1) by incubating sediment slurries at the respective in-situ temperature (Table
296 1) in the dark for a time period of 4 weeks. Headspace samples (0.1 ml) were taken every 3-5 days
297 over a time period of 4 weeks and potential methanogenesis rates were determined by the linear
298 increase of methane concentration over time (minimum of 6 time points).

300 2.7.1.4 Potential methylotrophic methanogenesis from methanol using stable isotope probing

301 One additional experiment was conducted with sediments from September 2014 by adding ¹³C-
302 labelled methanol to investigate the production of ¹³C-labelled methane. Three cores were stored at
303 1°C after the September 2014 cruise until further processing ~ 3.5 months later. The low storage
304 temperature together with the expected oxygen depletion in the enclosed supernatant water after
305 retrieval of the cores likely led to slowed anaerobic microbial activity during storage time and
306 preserved the sediments for potential methanogenesis measurements.

307 Sediment cores were sliced in 2 cm intervals and the upper 0-2 cmbsf sediment layer of all three
308 cores was combined in a beaker and homogenized. Then, sediment slurries were prepared by mixing
309 5 cm³ of sediment with 5 ml of artificial seawater medium in N₂-flushed, sterile glass vials (30 ml).

310 After this, methanol was added to the slurry with a final concentration of 10 mM (see 2.7.1.3).

311 Methanol was enriched with ¹³C-labelled methanol in a ratio of 1:1000 between ¹³C-labelled (99.9 %
312 ¹³C) and non-labelled methanol mostly consisting of ¹²C (manufacturer: Roth). In total, 54 vials were
313 prepared for nine different sampling time points during a total incubation time of 37 days. All vials
314 were incubated at 13°C (in situ temperature in September 2014) in the dark. At each sampling point,
315 six vials were stopped: one set of triplicates were used for headspace methane and carbon dioxide
316 determination and a second set of triplicates were used for porewater analysis.

317 Headspace methane and carbon dioxide concentrations (volume 100 µl) were determined on a
318 Shimadzu gas chromatograph (GC-2014) equipped with a packed Haysep-D column a flame ionization
319 detector and a methanizer. The methanizer (reduced nickel) reduces carbon dioxide with hydrogen
320 to methane at a temperature of 400°C. The column temperature was 80°C and the helium flow was
321 set to 12 ml min⁻¹. Methane concentrations (including reduced CO₂) were calibrated against methane
322 standards (Scotty gases). The detection limit was 0.1 ppm with a precision of 2 %.

323 Analyses of ¹³C/¹²C-ratios of methane and carbon dioxide were conducted after headspace
324 concentration measurements by using a continuous flow combustion gas chromatograph (Trace
325 Ultra, Thermo Scientific), which was coupled to an isotope ratio mass spectrometer (MAT253,
326 Thermo Scientific). The isotope ratios of methane and carbon dioxide given in the common delta-
327 notation (δ ¹³C in permill) are reported relative to Vienna Pee Dee Belemnite (VPDB) standard.

328 Isotope precision was +/- 0.5 ‰, when measuring near the detection limit of 10 ppm.

329 For porewater analysis of methanol concentration and isotope composition, each sediment slurry of
330 the triplicates was transferred into argon-flushed 15 ml centrifuge tubes and centrifuged for 6
331 minutes at 4500 rpm. Then 1 ml filtered (0.2 µm) porewater was transferred into N₂-flushed 2 ml
332 glass vials for methanol analysis, crimp sealed and immediately frozen at -20 °C. Methanol

333 concentrations and isotope composition were determined via high performance liquid
334 chromatography-ion ratio mass spectrometry (HPLC-IRMS, Thermo Fisher Scientific) at the MPI
335 Marburg. The detection limit was 50 μM with a precision of 0.3‰.

336 2.7.2 Methanogenesis in the gravity core

337 Ex situ hydrogenotrophic methanogenesis was determined in a gravity core taken in September 2013.
338 The pathway is thought to be the main methanogenic pathway in the ~~deep-sediment layers~~ (below
339 ~~sulfate penetration~~) the SMTZ in Eckernförde Bay (Whiticar, 2002). Hydrogenotrophic
340 methanogenesis was determined using radioactive sodium bicarbonate ($\text{NaH}^{14}\text{CO}_3$). At every
341 sampled sediment depth (12 depths in 30 cm intervals), triplicate glass tubes (5 mL) were inserted
342 directly into the sediment. Tubes were filled bubble-free with sediment and closed with butyl rubber
343 stoppers on both ends according to (Treude et al. 2005). Methods following sampling were identical
344 as described in 2.7.1.2.

345 2.8 Molecular analysis

346 During the non-labeled methanol treatment of the 0-1 cmbsf horizon from the September 2014
347 sampling (see 2.7.1.3), additional samples were prepared to detect and quantify the presence of
348 methanogens in the sediment. Therefore, additional 15 vials were prepared with addition of
349 methanol as described in 2.7.1.3 for five different time points (day 1 (= t_0), day 8, day 16, day 22, and
350 day 36) and stopped at each time point by transferring sediment from the triplicate slurries into
351 whirl-packs (Nasco), which then were immediately frozen at -20°C . DNA was extracted from ~ 500 mg
352 of sediment using the FastDNA[®] SPIN Kit for Soil (Biomedical). Quantitative real-time polymerase
353 chain reaction (qPCR) technique using TaqMan probes and TaqMan chemistry (Life Technologies) was
354 used for the detection of methanogens on a ViiA7 qPCR machine (Life Technologies). Primer and
355 Probe sets as originally published by Yu et al. (2005) were applied to quantify the orders
356 *Methanobacteriales*, *Methanosarcinales* and *Methanomicrobiales* along with the two families
357 *Methanosarcinaceae* and *Methanosaetaceae* within the order *Methanosarcinales*. In addition, a
358 universal primer set for detection of the domain *Archaea* was used (Yu et al. 2005).
359 Absolut quantification of the 16S rDNA from the groups mentioned above was performed with
360 standard dilution series. The standard concentration reached from 10^8 to 10^1 copies per μL .
361 Quantification of the standards and samples was performed in duplicates. Reaction was performed in
362 a final volume of 12.5 μL containing 0.5 μL of each Primer ($10\text{pmol } \mu\text{L}^{-1}$, MWG), 0.25 μL of the
363 respective probe ($10\text{ pmol } \mu\text{L}^{-1}$, Life Technologies), 4 μL H_2O (Roth), 6.25 μL TaqMan Universal Master
364 Mix II (Life Technologies) and 1 μL of sample or standard. Cycling conditions started with initial
365 denaturation and activation step for 10 min at 95°C , followed by 45 cycles of 95°C for 15 sec, 56°C

366 for 30 sec and 60°C for 60 sec. Non-template controls were run in duplicates with water instead of
367 DNA for all primer and probe sets, and remained without any detectable signal after 45 cycles.

368 2.9 Statistical Analysis

369 To determine possible environmental control parameters of [surfaceSRZ](#) methanogenesis, a Principle
370 Component Analysis (PCA) was applied according to the approach described in Gier et al. (2016).

371 Prior to PCA, the dataset was transformed into ranks to assure the same data dimension.

372 In total, two PCAs were conducted. The first PCA was used to test the relation of parameters in the
373 surface sediment (integrated methanogenesis (0-5 cm, $\text{mmol m}^{-2} \text{d}^{-1}$), POC content (average value
374 from 0-5 cmbsf, wt %), C/N (average value from 0-5 cmbsf, molar) and the bottom water (25 m water
375 depth) (oxygen (μM), temperature ($^{\circ}\text{C}$), salinity (PSU), chlorophyll ($\mu\text{g L}^{-1}$), methane (nM)). The
376 second PCA was applied on depth profiles of sediment [surfaceSRZ](#) methanogenesis ($\text{nmol cm}^{-3} \text{d}^{-1}$),
377 sediment depth (cm), sediment POC content (wt%), sediment C/N ratio (molar), and sampling month
378 (one value per depth profile at a specific month, the later in the year the higher the value).

379 For each PCA, biplots were produced to view data from different angles and to graphically determine
380 a potential positive, negative or zero correlation between methanogenesis rates and the tested
381 variables.

382 3. Results

383 3.1 Water column parameters

384 From March 2013 to September 2014, the water column had a pronounced temporal and spatial
385 variability of temperature, salinity, and oxygen (Fig. 2 and 3). In 2013, temperature of the upper
386 water column increased from March (1°C) to September (16°C), but decreased again in November
387 (11°C). The temperature of the lower water column increased from March 2013 (2°C) to November
388 2013 (12°C). In 2014, lowest temperatures of the upper and lower water column were reached in
389 March (4°C). Warmer temperatures of the upper water column were observed in June and
390 September (around 17°C), while the lower water column peaked in September (13°C).
391 Salinity increased over time during 2013, showing the highest salinity of the upper and lower water
392 column in November (18 and 23 PSU, respectively). In 2014, salinity of the upper water column was
393 highest in March and September (both 17 PSU), and lowest in June (13 PSU). The salinity of the lower
394 water column increased from March 2014 (21 PSU) to September 2014 (25 PSU).

395 In both years, June and September showed the most pronounced vertical gradient of temperature
396 and salinity, featuring a pycnocline at around ~ 14 m water depth.

397 Summer stratification was also seen in the O_2 profiles, which showed O_2 depleted conditions ($\text{O}_2 <$
398 $150 \mu\text{M}$) in the lower water column from June to September in both years, reaching concentrations

399 below 1- 2 μM (detection limit of CTD sensor) in September of both years (Fig. 2 and 3). The water
400 column was completely ventilated, i.e. homogenized, in March of both years with O_2 concentrations
401 of 300-400 μM down to the sea floor at about 28 m.

402

403 3.2 Sediment geochemistry in MUC cores

404 Sediment porewater and solid phase geochemistry results for the years 2013 and 2014 are shown in
405 Fig. 2 and 3, respectively.

406 Sulfate concentrations at the sediment surface ranged between 15-20 mM. Concentration decreased
407 with depth at all sampling months but was never fully depleted until the bottom of the core (18-29
408 cmbsf, between 2 and 7 mM sulfate). November 2013 showed the strongest decrease from ~ 20 mM
409 at the top to ~ 2 mM at the bottom of the core (27 cmbsf).

410 Opposite to sulfate, methane concentration increased with sediment depth in all sampling months
411 (Fig. 2 and 3). Over the course of a year (i.e. March to November in 2013, and March to September in
412 2014), maximum methane concentration increased, reaching the highest concentration in November
413 2013 (~ 1 mM at 26 cmbsf) and September 2014 (0.2 mM at 23 cmbsf), respectively. Simultaneously,
414 methane profiles became steeper, revealing higher methane concentrations at shallower sediment
415 depth late in the year. Magnitudes of methane concentrations were similar in the respective months
416 of 2013 and 2014.

417 In all sampling months, sulfide concentration increased with sediment depth (Fig. 2 and 3). Similar to
418 methane, sulfide profiles revealed higher sulfide concentrations at shallower sediment depth
419 together with higher peak concentrations over the course the sampled months in each sampling
420 year. Accordingly, November 2013 (10.5 mM at 15 cmbsf) and September 2014 (2.8 mM at 15
421 cmbsf) revealed the highest sulfide concentrations, respectively. September 2014 was the only
422 sampling month showing a pronounced decrease in sulfide concentration from 15 cmbsf to 21 cmbsf
423 of over 50 %.

424 DIC concentrations increased with increasing sediment depth at all sampling months. Concomitant
425 with highest sulfide concentrations, highest DIC concentration was detected in November 2013 (26
426 mM at 27 cmbsf). At the surface, DIC concentrations ranged between 2-3 mM at all sampling
427 months. In June of both years, DIC concentrations were lowest at the deepest sampled depth
428 compared to the other sampling months (16 mM in 2013, 13 mM in 2014).

429 At all sampling months, POC profiles scattered around 5 ± 0.9 wt % with depth. Only in November
430 2013, June 2014 and September 2014, POC content exceeded 5 wt % in the upper 0-1 cmbsf (5.9, 5.2
431 and 5.3 wt %, respectively) with the highest POC content in November 2013. Also in November 2013,
432 surface C/N ratio (0-1 cmbsf) of the particulate organic matter was lowest of all sampling months

433 (8.6). In general, C/N ratio increased with depth in both years with values around 9 at the surface and
434 values around 10-11 at the deepest sampled sediment depths.

435 3.3 Sediment geochemistry in gravity cores

436 Results from sediment porewater and solid phase geochemistry in the gravity core from September
437 2013 are shown in Fig. 4. Please note that the sediment depth of the gravity core was corrected by
438 comparing the sulfate concentrations at 0 cmbsf in the gravity core with the corresponding sulfate
439 concentration and depth in the MUC core from September 2013 (Fig. 2). The soft surface sediment is
440 often lost during the gravity coring procedure. Through this ~~correction~~ correction, the topmost layer
441 of the gravity core was set at a depth of 14 cmbsf.

442 Porewater sulfate concentration in the gravity core decreased with depth (i.e. below 0.1 mM at 107
443 cmbsf) and stayed below 0.1 mM until 324 cmbsf. Sulfate increased slightly (1.9 mM) at the bottom
444 of the core (345 cmbsf). In concert with sulfate, also methane, sulfide, DIC, POC and C/N profiles
445 showed distinct alteration in the profile at 345 cmbsf (see below, Fig. 4). As fluid seepage has not
446 been observed at the Boknis Eck station (Schlüter et al., 2000), these alterations could either indicate
447 a change in sediment properties or result from a sampling artifact from the penetration of seawater
448 through the core catcher into the deepest sediment layer. The latter process is, however, not
449 expected to considerably affect sediment solid phase properties (POC and C/N), and we therefore
450 dismissed this hypothesis.

451 Methane concentration increased steeply with depth reaching a maximum of 4.8 mM at 76 cmbsf.
452 Concentration stayed around 4.7 mM until 262 cmbsf, followed by a slight decrease until 324 cmbsf
453 (2.8 mM). From 324 cmbsf to 345 cmbsf methane increased again (3.4 mM).

454 Both sulfide and DIC concentrations increased with depth, showing a maximum at 45 cmbsf (~ 5mM)
455 and 345 cmbsf (~ 1mM), respectively. While sulfide decreased after 45 cmbsf to a minimum of ~ 300
456 μM at 324 cmbsf, it slightly increased again to ~1 mM at 345 cmbsf. In accordance, DIC
457 concentrations showed a distinct decrease between 324 cmbsf to 345 cmbsf (from 45 mM to 39
458 mM).

459 While POC contents varied around 5 wt % throughout the core, C/N ratio slightly increased with
460 depth, revealing the lowest ratio at the surface (~3) and the highest ratio at the bottom of the core
461 (~13). However, both POC and C/N showed a distinct increase from 324 cmbsf to 345 cmbsf.

462

463 3.4 Methanogenesis activity in MUC cores

464 3.4.1 Net methanogenesis

465 Net methanogenesis activity (calculated by the linear increase of methane over time, see Fig. S1) was
466 detected throughout the cores at all sampling months (Fig. 2 and 3). Activity measured in MUC cores

467 increased over the course of the year in 2013 and 2014 (that is: March to November in 2013 and
468 March to September in 2014) with lower rates mostly $< 0.1 \text{ nmol cm}^{-3} \text{ d}^{-1}$ in March and higher rates $>$
469 $0.2 \text{ nmol cm}^{-3} \text{ d}^{-1}$ in November 2013 and September 2014, respectively. In general, November 2013
470 revealed highest net methanogenesis rates ($1.3 \text{ nmol cm}^{-3} \text{ d}^{-1}$ at 1-2 cmbsf). Peak rates were
471 detected at the sediment surface (0-1 cmbsf) at all sampling months except for September 2013
472 where the maximum rates were situated between 10-15 cmbsf. In addition to the surface peaks, net
473 methanogenesis showed subsurface (= below 1 cmbsf until 30 cmbsf) maxima at all sampling
474 months, but with alternating depths (between 10 and 25 cmbsf).
475 Comparison of integrated net methanogenesis rates (0-25 cmbsf) revealed highest rates in
476 September and November 2013 ($0.09 \text{ mmol m}^{-2} \text{ d}^{-1}$ and $0.08 \text{ mmol m}^{-2} \text{ d}^{-1}$, respectively) and lowest
477 rates in March 2014 ($0.01 \text{ mmol m}^{-2} \text{ d}^{-1}$) (Fig. 5). A trend of increasing areal net methanogenesis rates
478 from March to September was observed in both years.

479 **3.4.2 Hydrogenotrophic methanogenesis**

480 Hydrogenotrophic methanogenesis activity determined by ^{14}C -bicarbonate incubations of MUC cores
481 is shown in Fig. 2 and 3. In 2013, maximum activity ranged between 0.01 - $0.2 \text{ nmol cm}^{-3} \text{ d}^{-1}$, while in
482 2014 maxima ranged only between 0.01 and $0.05 \text{ nmol cm}^{-3} \text{ d}^{-1}$. In comparison, maximum
483 hydrogenotrophic methanogenesis was up to two orders of magnitude lower compared to net
484 methanogenesis. Only in March 2013 both activities reached a similar range.

485 Overall, hydrogenotrophic methanogenesis increased with depth in March, September, and
486 November 2013 and in March, June, and September 2014. In June 2013, activity decreased with
487 depth, showing the highest rates in the upper 0-5 cmbsf and the lowest at the deepest sampled
488 depth.

489 Concomitant with integrated net methanogenesis, integrated hydrogenotrophic methanogenesis
490 rates (0-25 cmbsf) were high in September 2013, with slightly higher rates in March 2013 (Fig. 5).
491 Lowest areal rates of hydrogenotrophic methanogenesis were seen in June of both years.

492 Hydrogenotrophic methanogenesis activity in the gravity core is shown in Fig. 4. Highest activity (\sim
493 $0.7 \text{ nmol cm}^{-3} \text{ d}^{-1}$) was measured at 45 cmbsf and 138 cmbsf, followed by a decrease with increasing
494 sediment depth reaching $0.01 \text{ nmol cm}^{-3} \text{ d}^{-1}$ at the deepest sampled depth (345 cmbsf).

495 **3.4.3 Potential methanogenesis in manipulated experiments**

496 Potential methanogenesis rates in manipulated experiments included either the addition of
497 inhibitors (molybdate for inhibition of sulfate reduction or BES for inhibition of methanogenesis) or
498 the addition of a non-competitive substrate (methanol). Control treatments were run with neither
499 the addition of inhibitors nor the addition of methanol.

500 *Controls.* Potential methanogenesis activity in the control treatments was below $0.5 \text{ nmol cm}^{-3} \text{ d}^{-1}$
501 from March 2014 to September 2014 (Fig. 6). Only in November 2013, control rates exceeded 0.5
502 $\text{nmol cm}^{-3} \text{ d}^{-1}$ below 6 cmbsf. While rates increased with depth in November 2013 and June 2014,
503 they decreased with depth at the other two sampling months.

504 *Molybdate.* Peak potential methanogenesis rates in the molybdate treatments were found in the
505 uppermost sediment interval (0-1 cmbsf) at almost every sampling month with rates being 3-30
506 times higher compared to the control treatments ($< 0.5 \text{ nmol cm}^{-3} \text{ d}^{-1}$). In November 2013, potential
507 methanogenesis showed two maxima (0-1 and 10-15 cmbsf). Highest measured rates were found in
508 September 2014 ($\sim 6 \text{ nmol cm}^{-3} \text{ d}^{-1}$), followed by November 2013 ($\sim 5 \text{ nmol cm}^{-3} \text{ d}^{-1}$).

509 *BES.* Profiles of potential methanogenesis in the BES treatments were similar to the controls mostly
510 in the lower range $< 0.5 \text{ nmol cm}^{-3} \text{ d}^{-1}$. Only in November 2013 rates exceeded $0.5 \text{ nmol cm}^{-3} \text{ d}^{-1}$.
511 Rates increased with depth at all sampling months, except for September 2014, where highest rates
512 were found at the sediment surface (0-1 cmbsf).

513 *Methanol.* At all sampling months, potential rates in the methanol treatments were three orders of
514 magnitude higher compared to the control treatments ($< 0.5 \text{ nmol cm}^{-3} \text{ d}^{-1}$). Except for November
515 2013, potential methanogenesis rates in the methanol treatments were highest in the upper 0-5
516 cmbsf and decreased with depth. In November 2013, highest rates were detected at the deepest
517 sampled depth (20-25 cmbsf).

518

519 **3.4.4 Potential methanogenesis followed by ^{13}C -methanol labeling**

520 The concentration of total methanol concentrations (labeled and unlabeled) in the sediment
521 decreased sharply in the first 2 weeks from $\sim 8 \text{ mM}$ at day 1 to 0.5 mM at day 13 (Fig. 7). At day 17,
522 methanol was below the detection limit. In the first 2 weeks, residual methanol was enriched with
523 ^{13}C , reaching $\sim 200 \text{ ‰}$ at day 13.

524 Over the same time period, the methane content in the headspace increased from 2 ppmv at day 1
525 to $\sim 66,000 \text{ ppmv}$ at day 17 and stayed around that value until the end of the total incubation time
526 (until day 37) (Fig. 7). The carbon isotopic signature of methane ($\delta^{13}\text{C}_{\text{CH}_4}$) showed a clear enrichment
527 of the heavier isotope ^{13}C (Table 3) from day 9 to 17 (no methane was detectable at day 1). After day
528 17, $\delta^{13}\text{C}_{\text{CH}_4}$ stayed around 13 ‰ until the end of the incubation. The content of CO_2 in the headspace
529 increased from $\sim 8900 \text{ ppmv}$ at day 1 to $\sim 29,000 \text{ ppmv}$ at day 20 and stayed around $30,000 \text{ ppmv}$
530 until the end of the incubation (Fig. 7). Please note, that the major part of CO_2 was dissolved in the
531 porewater, thus the CO_2 content in the headspace does not show the total CO_2 abundance in the
532 system. CO_2 in the headspace was enriched with ^{13}C during the first 2 weeks (from -16.2 to -7.3 ‰)
533 but then stayed around -11 ‰ until the end of the incubation.

534 3.5 Molecular analysis of benthic methanogens

535 In September 2014, additional samples were run during the methanol treatment (see Sect. 2.7.) for
536 the detection of benthic methanogens via qPCR. The qPCR results are shown in Fig. 8. For a better
537 comparison, the microbial abundances are plotted together with the sediment methane
538 concentrations from the methanol treatment, from which the rate calculation for the methanol-
539 methanogenesis at 0-1 cmbsf was done (shown in Fig. 6).
540 Sediment methane concentrations increased over time revealing a slow increase in the first ~10 days,
541 followed by a steep increase between day 13 and day 20 and ending in a stationary phase.
542 A similar increase was seen in the abundance of total and methanogenic archaea. Total archaea
543 abundances increased sharply in the second week of the incubation reaching a maximum at day 16
544 ($\sim 5000 \cdot 10^6$ copies g^{-1}) and stayed around $3000 \cdot 10^6$ - $4000 \cdot 10^6$ copies g^{-1} over the course of the
545 incubation. Similarly, methanogenic archaea, namely the order *Methanosarcinales* and within this
546 order the family *Methanosarcinaceae*, showed a sharp increase in the first 2 weeks as well with the
547 highest abundances at day 16 ($\sim 6 \cdot 10^8$ copies g^{-1} and $\sim 1 \cdot 10^6$ copies g^{-1} , respectively). Until the end of
548 the incubation, the abundances of *Methanosarcinales* and *Methanosarcinaceae* decreased to about a
549 third of their maximum abundances ($\sim 2 \cdot 10^8$ copies g^{-1} and $\sim 0.4 \cdot 10^6$ copies g^{-1} , respectively).

550 3.6 Statistical Analysis

551 The PCA of integrated [surfaceSRZ](#) methanogenesis (0-5 cmbsf) (Fig. 10) showed a positive correlation
552 with bottom water temperature (Fig. 10a), bottom water salinity (Fig. 10a), bottom water methane
553 (Fig. 10b), surface sediment POC content ([0-5 cmbsf](#), Fig. 10c), and surface sediment C/N ([0-5 cmbsf](#),
554 Fig. 10b). A negative correlation was found with bottom water oxygen concentration (Fig. 10b). No
555 correlation was found with bottom water chlorophyll.

556 The PCA of methanogenesis depth profiles showed positive correlations with sediment depth (Fig.
557 11a) and C/N (Fig. 11b), and showed negative correlations with POC (Fig. 11a).

558

559 4. Discussion

560 4.1 Methanogenesis in the sulfate-reducing zone

561 On the basis of the results presented in Fig. 2 and 3, it is evident that methanogenesis and sulfate
562 reduction were concurrently active in the [surface-sediments/sulfate reduction zone](#) (0-30 cmbsf) at
563 Boknis Eck. Even though sulfate reduction activity was not directly determined, the decrease in
564 sulfate concentrations with a concomitant increase in sulfide within the upper 30 cmbsf clearly
565 indicated its presence (Fig. 2 and 3). Several previous studies confirmed the high activity of sulfate
566 reduction in the surface sediment of Eckernförde Bay, revealing rates up to 100 - $10,000$ $nmol\ cm^{-3}\ d^{-1}$

567 in the upper 25 cmbsf (Treude et al., 2005a; Bertics et al., 2013; Dale et al., 2013). Microbial
568 fermentation of organic matter was probably high in the organic-rich sediments of Eckernförde Bay
569 (POC contents of around 5 %, Fig. 2 and 3), providing high substrate availability and variety for
570 methanogenesis.

571

572 The results of this study further identified methylotrophy to be a potentially important non-
573 competitive methanogenic pathway in the sulfate-reducing zone. The pathway utilizes alternative
574 substrates, such as methanol, to bypass competition with sulfate reducers for H₂ and acetate. A
575 potential for methylotrophic methanogenesis within the sulfate-reducing zone was supported by the
576 following observations; ~~which will be discussed in more detail in the subsequent chapters:~~

577 1) ~~1)~~ Hydrogenotrophic methanogenesis was up to two orders of magnitude lower compared to
578 net methanogenesis, resulting in insufficient rates to explain the observed net
579 methanogenesis in the upper 0-30 cmbsf (Fig. 2 and 3). This points towards ~~pointing to~~ the
580 presence of alternative methanogenic processes in the ~~sediment surface layers~~ sulfate
581 reduction zone, such as methylotrophic methanogenesis (Fig. 2 and 3).

582 2) ~~2)~~ Methanogenesis increased when sulfate reduction was inhibited by molybdate,
583 confirming the inhibitory effect of sulfate reduction on methanogenesis with competitive
584 substrates (H₂ and acetate (Oremland & Polcin, 1982; King et al., 1983)) (Fig. 6),
585 Consequently, usage of non-competitive substrates was preferred in ~~surface sediments~~
586 where sulfate reduction was active zone (especially in the upper 0-1 cmbsf, Fig. 6).
587 Accordingly, hydrogenotrophic methanogenesis increased at depths where sulfate was
588 depleted and thus the competitive situation was relieved (Fig. 4).

589 3) ~~3)~~ The addition of BES did not result in the inhibition of methanogenesis, indicating the
590 presence of unconventional methanogenic groups using non-competitive substrates (Fig. 7).
591 The unsuccessful inhibition by BES can be explained either by incomplete inhibition or by the
592 fact that the methanogens were insensitive to BES (Hoehler et al., 1994; Smith & Mah, 1981;
593 Santoro & Konisky, 1987). The BES concentration applied in the present study (60 mM) has
594 been shown to result in successful inhibition of methanogens in previous studies (Hoehler et
595 al., 1994). Therefore, the presence of methanogens that are insensitive to BES is more likely.
596 The insensitivity to BES in methanogens is explained by heritable changes in BES permeability
597 or formation of BES-resistant enzymes (Smith & Mah, 1981; Santoro & Konisky, 1987). Such
598 BES resistance was found in *Methanosarcina* mutants (Smith & Mah, 1981; Santoro &
599 Konisky, 1987). This genus was successfully detected in our samples (for more details see
600 point 5), and is known for mediating the methylotrophic pathway (Keltjens & Vogels, 1993).
601 supporting our hypothesis on the utilization of non-competitive substrates by methanogens.

Formatted: Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

602 4) -4) The addition of methanol to sulfate-rich sediments increased methanogenesis rates up
603 to three orders of magnitude, confirming the potential of the methanogenic community to
604 utilize non-competitive substrates especially in the 0-5 cmbsf sediment horizon (Fig. 67). At
605 this sediment depth either the availability of non-competitive substrates, including
606 methanol, was highest (derived from fresh organic matter), or the usage of non-competitive
607 substrates was increased due to the high competitive situation as sulfate reduction is most
608 abundant active in the 0-5 cmbsf layer (Treude et al., 2005a; Bertics et al., 2013). It should be
609 noted that even though methanogenesis rates were calculated assuming a linear increase in
610 methane concentration over the entire incubation to make a better comparison between
611 different treatments, the methanol treatments generally showed a delayed response in
612 methane development (Fig. 8, Supplement, Fig. S2). We suggest that this delayed response
613 was a reflection of cell growth by methanogens utilizing the surplus methanol. We are
614 therefore unable to decipher whether methanol plays a major role as a substrate in the
615 Eckernförde Bay sediments compared to possible alternatives, as its concentration is
616 relatively low in the natural setting (1.05 μ M in the 0-1 cmbsf layer, \sim 1.2 μ M at between 0-1
617 and -25 cmbsf, June 2014 sampling, G.-C. Zhuang unpubl. data). It is conceivable that other
618 non-competitive substrates, such as methylated sulfides (e.g., dimethyl sulfide or
619 methanethiol), are more relevant for the support of surface SRZ methanogenesis.
620 5) -5) Methylotrophic methanogens of the order *Methanosarcinales* were detected in the
621 methanol-treatment (Fig. 8), confirming the presence of methanogens that utilize non-
622 competitive substrates in the natural environment (Boone et al., 1993), (Fig. 8). The delay in
623 growth of *Methanosarcinales* moreover hints towards the predominant usage of other non-
624 competitive substrates over methanol (see also point 4).
625 6) -6) Stable isotope probing revealed highly 13 C-enriched methane produced from
626 13 C-labelled methanol, furthermore confirming the potential of the methanogenic
627 community to utilize non-competitive substrates (Fig. 7). The production of both methane
628 and CO₂ from methanol has been shown previously in different strains of methylotrophic
629 methanogens (Penger et al., 2012). The fast conversion of methanol to methane and CO₂
630 (methanol was consumed completely in 17 days) is hinting towards the presence of
631 methylotrophic methanogens (e.g. members of the family *Methanosarcinaceae*, which is
632 known for the methylotrophic pathway (Keltjens & Vogels, 1993)). Please note, however,
633 that the storage of the cores (3.5 months) prior to sampling could have led to shifts in the
634 microbial community and thus might not reflect in-situ conditions of the original microbial
635 community in September 2014. The delay in methane production also seen in the stable
636 isotope experiment was, however, only slightly different (methane developed earlier,

Formatted: Not Highlight

637 between day 8 and 12, data not shown) from the non-labeled methanol treatment (between
638 day 10 to 16, Fig. S2), which leads us to the assumption that the storage time at 1°C did not
639 dramatically affect the methanogen community. Similar, in a previous study with arctic
640 sediments, addition of substrates had no stimulatory effect on the rate of methanogenesis or
641 on the methanogen community structure at low temperatures (5°C, (Blake et al., 2015)).

643 **4.1.1 Hydrogenotrophic methanogenesis**

644 We demonstrated that hydrogenotrophic methanogenesis was insufficient to explain the observed
645 net methanogenesis, pointing to the presence of alternative pathways that utilize substrates other
646 than H₂. One exemption was detected in the March 2013 incubation, where rates of
647 hydrogenotrophic methanogenesis exceeded net methanogenesis in discrete depths (5–6 cmbsf and
648 25–30 cmbsf). It is possible that additional carbon sources led to increased local fermentation
649 processes, for instance from the deposition of macro-algae detritus, which is produced during winter
650 storms and can be transported into deeper sediment layers by bioturbation, where it is digested and
651 released as fecal pellets (Meyer-Reil, 1983; Bertics et al., 2013). Such additional carbon sources from
652 fresh material could lead to the local accumulation of excess hydrogen through fermentation and
653 reduce the competition for H₂ between sulfate reducers and methanogens (Treude et al., 2009). C/N
654 ratios in March 2013 were more scattered compared to other months in 2013 and 2014, indicating
655 the transport of labile material into the sediment. Eckernförde Bay sediments are known for
656 bioturbation especially during early spring by mollusks and polychaetes in the upper 10 cm of the
657 sediment (D'Andrea et al., 1996; Orsi et al., 1996; Bertics et al., 2013; Dale et al., 2013), and empty
658 mollusk shells were observed even at depth of ~ 20 cmbsf during sampling in the present study
659 (personal observation).

660 Hydrogenotrophic methanogenesis was also detected in the gravity core in September 2013.
661 Maximum rates were found at 45 cmbsf and 138 cmbsf, indicating a higher usage of H₂ at depths >
662 40 cmbsf, where sulfate was depleted and thus the competition between sulfate reducers and
663 methanogens was relieved. It should be noted, however, that the peak in at 45 cmbsf could also be a
664 result of tracer (H¹⁴CO₃⁻) back flux associated with AOM (Holler et al., 2011), as this peak is situated
665 directly at the SMTZ (Fig. 4)

666 **4.1.2 Inhibition of sulfate reducers**

667 Supposedly the competition between methanogens and sulfate reducers within the upper 30 cmbsf
668 led to the predominant utilization of non-competitive substrates by methanogenesis, as indicated by
669 lower hydrogenotrophic vs. higher net methanogenesis rates (see discussion above). After the
670 addition of the sulfate reducer inhibitor molybdate, competitive substrates (H₂ and acetate

671 (Oremland & Polcin, 1982; King et al., 1983) were available for methanogenesis resulting in the (up to
672 30 times) increase in potential activity (Fig. 6 and 7). Notably, highest rates in the molybdate
673 treatment were measured at the shallowest sediment depth at most sampling months (except
674 November 2013), pointing towards the strongest competition between sulfate reducers and
675 methanogens directly at the top 0–1 cmbsf. Accordingly, maximum sulfate reduction activity was
676 detected in this depth layer in earlier studies (Bertics et al. 2013; Treude et al. 2005). In conclusion,
677 findings from the molybdate addition experiment highlight that the methanogenic community is
678 subject to a strong competition with sulfate reducers in the surface sediments and that the majority
679 of the observed methane production under sulfate-reducing conditions can be attributed to the
680 utilization of non-competitive substrates.

681 4.1.3 Inhibition of methanogenesis by BES

682 BES acts as a specific inhibitor of methanogens, because it is a structural analogue of (coenzyme M),
683 an enzyme only found in methanogens (Gunsalus et al., 1978; Hoehler et al., 1994). Addition of BES
684 did not result in the expected inhibition of potential methanogenesis; instead rates were in the same
685 range as the control treatment (Fig. 7). Consequently, either the inhibition of BES was incomplete, or
686 the methanogens were insensitive to BES (Hoehler et al., 1994; Smith & Mah, 1981; Santoro &
687 Konisky, 1987). The BES concentration applied in the present study (60 mM) has been shown to
688 result in successful inhibition of methanogens in previous studies (Hoehler et al., 1994). Therefore,
689 the presence of methanogens that are insensitive to BES is more likely. The insensitivity to BES in
690 methanogens is explained by heritable changes in BES permeability or formation of BES-resistant
691 enzymes (Smith & Mah, 1981; Santoro & Konisky, 1987). Such BES resistance was found in
692 *Methanosarcina* mutants (Smith & Mah, 1981; Santoro & Konisky, 1987). This genus was successfully
693 detected in our samples (for more details see 4.1.5), and is known for mediating the methylotrophic
694 pathway (Keltjens & Vogels, 1993), supporting our hypothesis on the utilization of non-competitive
695 substrates by methanogens.

696 4.1.4 Methanol addition

697 High potential methanogenesis rates observed after the addition of the non-competitive substrate
698 methanol (Fig. 6) leads to the assumption that methylotrophic methanogens are present in surface
699 sediments of Eckernförde Bay. Except for November 2013, highest rates in the methanol treatment
700 were detected in the upper 0–5 cmbsf and decreased with depth. This observation can be interpreted
701 twofold: (1) The availability of non-competitive substrates, including methanol, was most likely
702 highest at the sediment surface, as those substrates are derived from fresh organic matter, such as
703 pectin or betaine and dimethylpropiothetin (both osmoprotectants) (Zinder, 1993). Hence, the
704 methanol-utilizing methanogenic community had its highest abundance in this zone. (2) Sulfate

705 reduction is most dominant in the 0-5 cmbsf (Treude et al., 2005a; Bertics et al., 2013), which
706 probably leads prevalent methanogens to an increased usage of non-competitive substrates.
707 It should be noted that even though methanogenesis rates were calculated assuming a linear
708 increase in methane concentration over the entire incubation to make a better comparison between
709 different treatments, the methanol treatments generally showed a delayed response in methane
710 development (Fig. 8, Supplement, Fig. S2). We suggest that this delayed response was a reflection of
711 cell growth by methanogens utilizing the surplus methanol. We are therefore unable to decipher
712 whether methanol plays a major role as a substrate in the Eckernförde Bay sediments compared to
713 possible alternatives, as its concentration is relatively low in the natural setting (1.05 μM in the 0-1
714 cmbsf layer, $\sim 1.2 \mu\text{M}$ at 1-25 cmbsf, June 2014 sampling, G. C. Zhuang unpubl. data). It is conceivable
715 that other non-competitive substrates, such as methylated sulfides (e.g., dimethyl sulfide or
716 methanethiol), are more relevant for the support of surface methanogenesis. In the marine
717 environment, dimethyl sulfide mainly originate from the algae osmoregulatory compound
718 dimethylsulfoniopropionate (DMSP) (Van Der Maarel & Hansen, 1997), which could have
719 accumulated in Eckernförde Bay sediments, due to intense sedimentation of algae blooms (Bange et
720 al., 2011). (Maltby et al., 2016) detected a similar delay in methane production in organic rich surface
721 sediments sampled off Peru after the addition of methanol, and suggested the predominant use of
722 methylated sulfides. Certain *Methanosarcina* species have been shown to use DMSP as a substrate
723 (Sieburth et al., 1993; Van Der Maarel & Hansen, 1997), a genus, which has been detected in our
724 samples (see 4.1.5 for more details).

725 4.1.5 Presence of methylotrophic methanogens

726 Simultaneously with the increase in methane concentration after methanol addition in the surface
727 layer (0-1 cmbsf) in September 2014, the DNA counts for the order *Methanosarcinales* and the family
728 *Methanosarcinaceae* within the order *Methanosarcinales* increased 10^2 to 10^6 times, respectively,
729 compared to the respective DNA abundances at the start of the incubation (Fig. 8). The successful
730 enrichment of *Methanosarcinaceae* indicates that this family is present in the natural environment
731 and thus could in part be responsible for the observed surface methanogenesis. As the members of
732 the family *Methanosarcinaceae* are known for utilization of methylated substrates (Boone et al.,
733 1993), our hypothesis for the presence of methylotrophic methanogenesis is supported. The delay in
734 growth of *Methanosarcinales* and *Methanosarcinaceae*, moreover hints towards the predominant
735 usage of other non-competitive substrates over methanol (see also 4.1.4).

736 4.1.6 Stable isotope experiment

737 Samples taken in September 2014 for the labeling experiment (^{13}C -enriched methanol, initial isotopic
738 signature: +26 ‰) showed that methanol was completely consumed after 17 days and converted to

739 methane and CO₂, as both revealed a concomitant enrichment in ¹³C. The production of both
740 methane and CO₂ from methanol has been shown previously in different strains of methylotrophic
741 methanogens (Penger et al., 2012).
742 Isotopic fractionation factors of methylotrophic methanogenesis from methanol to methane have
743 been found to be 1.07–1.08 (Heyer et al., 1976; Krzycki et al., 1987). This fractionation leads to a
744 progressive enrichment of ¹³C in the residual methanol until all methanol is consumed. Accordingly,
745 methanol was enriched in ¹³C in the first 13 days, as the consumption of ¹²C-methanol was preferred
746 by the microbes. The fast conversion of methanol to methane is hinting towards the presence of
747 methylotrophic methanogens (e.g. members of the family *Methanosarcinaceae*, which is known for
748 the methylotrophic pathway (Keltjens & Vogels, 1993)). Please note, however, that the storage of the
749 cores (3.5 months) prior to sampling could have led to shifts in the microbial community and thus
750 might not reflect in-situ conditions of the original microbial community in September 2014. The delay
751 in methane production also seen in the stable isotope experiment was, however, only slightly
752 different (methane developed earlier, between day 8 and 12, data not shown) from the non-labeled
753 methanol treatment (between day 10 to 16, Fig. S2), which leads us to the assumption that the
754 storage time at 1 °C did not dramatically affect the methanogen community. Similar, in a previous
755 study with arctic sediments, addition of substrates had no stimulatory effect on the rate of
756 methanogenesis or on the methanogen community structure at low temperatures (5 °C, (Blake et al.,
757 2015).

758 4.2 Environmental control of [surface-methanogenesis in the sulfate reduction zone](#)

759 [SurfaceSRZ](#) methanogenesis in Eckernförde Bay sediments showed variations throughout the
760 sampling period, which may be influenced by variable environmental factors such as temperature,
761 salinity, oxygen, and organic carbon. In the following, we will discuss the potential impact of those
762 factors on the magnitude and distribution of [surfaceSRZ](#) methanogenesis.

763 4.2.1 Temperature

764 During the sampling period, bottom water temperatures increased over the course of the year from
765 late winter (March, 3–4 °C) to autumn (November, 12 °C, Fig. 2 and 3). The PCA revealed a positive
766 correlation between bottom water temperature and integrated [surfaceSRZ](#) methanogenesis (0–5
767 cmbsf). A temperature experiment conducted with sediment from ~75 cmbsf in September 2014
768 within a parallel study revealed a mesophilic temperature optimum of methanogenesis (20 °C, data
769 not shown). Whether methanogenesis in [the surface sediments sulfate reduction zone](#) (0–30 cm) has
770 the same physiology remains speculative. However, AOM organisms, which are closely related to
771 methanogens (Knittel & Boetius, 2009), studied in [surface sediments the sulfate reduction zone](#) from
772 the same site were confirmed to have a mesophilic physiology, too (Treude et al. 2005). The sum of

773 these aspects lead us to the conceivable conclusion that [surfaceSRZ](#) methanogenesis activity in the
774 Eckernförde Bay is positively impacted by temperature increases. Such a correlation between benthic
775 methanogenesis and temperature has been found in several previous studies from different
776 environments ((Sansone & Martens, 1981; Crill & Martens, 1983; Martens & Klump, 1984).

778 **4.2.2 Salinity and oxygen**

779 From March 2013 to November 2013, and from March 2014 to September 2014, salinity increased in
780 the bottom-near water (25 m) from 19 to 23 PSU and from 22 to 25 PSU (Fig. 2 and 3), respectively,
781 due the pronounced summer stratification in the water column between saline North Sea water and
782 less saline Baltic Sea water (Bange et al., 2011). The PCA detected a positive correlation between
783 integrated [surfaceSRZ](#) methanogenesis (0-5 cmbsf) and salinity in the bottom-near water (Fig. 10a).
784 This correlation can hardly be explained by salinity alone, as methanogens feature a broad salinity
785 range from freshwater to hypersaline (Zinder, 1993). More likely, ~~methanogenesis was affected by~~
786 ~~variations in water column sulfate concentrations, which change alongside salinity (Pattnaik et al.,~~
787 ~~2000), providing either more (high salinity) or less (low salinity) of the electron acceptor for the~~
788 ~~degradation of organic matter by the sulfate-reducing bacteria in the sediment. Alternatively,~~ salinity
789 ~~may also serves~~ as an indicator of water-column stratification, which is often correlated with low O₂
790 concentrations in the Eckernförde Bay (Anon, n.d.)(Fig. S3, Bange et al., 2011; Bertics et al., 2013).
791 Methanogenesis is sensitive to O₂ (Oremland, 1988; Zinder, 1993), and hence conditions might be
792 more favorable during hypoxic or anoxic events, particular in the sediment closest to the sediment-
793 water interface, but potentially also in deeper sediment layers due to the absence of bioturbating
794 and bioirrigating infauna (Dale et al., 2013; Bertics et al., 2013), which could introduce O₂ beyond
795 diffusive transport. Accordingly, the PCA revealed a negative correlation between O₂ concentration
796 close to the seafloor and [surfaceSRZ](#) methanogenesis.

798 **4.2.4 Particulate organic carbon**

799 The supply of particulate organic carbon (POC) is one of the most important factors controlling
800 benthic heterotrophic processes, as it determines substrate availability and variety (Jørgensen,
801 2006). In Eckernförde Bay, the organic material reaching the seafloor originates mainly from
802 phytoplankton blooms in spring, summer and autumn (Bange et al., 2011). It has been estimated that
803 >50 % in spring (February/March), <25 % in summer (July/August) and >75 % in autumn
804 (September/October) of these blooms is reaching the seafloor (Smetacek et al., 1984), resulting in a
805 overall high organic carbon content of the sediment (5 wt %), which leads to high benthic microbial
806 degradation rates including sulfate reduction and methanogenesis (Whiticar, 2002; Treude et al.,
807 2005a; Bertics et al., 2013). Previous studies revealed that high organic matter availability can relieve

808 competition between sulfate reducers and methanogens in sulfate-containing, marine sediments
809 (Oremland et al., 1982; Holmer & Kristensen, 1994; Treude et al., 2009; Maltby et al., 2016).
810 To determine the effect of POC concentration and C/N ratio (the latter as a negative indicator for the
811 freshness of POC) on [surfaceSRZ](#) methanogenesis, two PCAs were conducted with a) the focus on the
812 upper 0-5 cmbsf, which is directly influenced by freshly sedimented organic material from the water
813 column (Fig. 10), and b) the focus on the depth profiles throughout the sediment cores (up to 30
814 cmbsf) (Fig. 11).

815 **a) Effect of POC and C/N ratio in the upper 0-5 cmbsf**

816 For the upper 0-5 cmbsf in the sediment, a positive correlation was found between [surfaceSRZ](#)
817 methanogenesis (integrated) and POC content (averaged) (Fig. 10c), indicating that POC content is an
818 important controlling factor for methanogenesis in this layer. In support, highest bottom-near water
819 chlorophyll concentrations coincided with highest bottom-near water methane concentrations and
820 high integrated [surfaceSRZ](#) methanogenesis (0-5 cmbsf) in September 2013, probably as a result of
821 the sedimentation of the summer phytoplankton bloom (Fig. 9). Indeed, the PCA revealed a -positive
822 correlation between integrated [surfaceSRZ](#) methanogenesis rates and bottom-near water methane
823 concentrations (Fig. 10b), when viewed over all investigated months. However, no correlation was
824 found between bottom water chlorophyll and integrated [surfaceSRZ](#) methanogenesis rates (Fig. 10).
825 As seen in Fig. 9, bottom-near high chlorophyll concentrations did not coincide with high bottom-
826 near methane concentration in June/September 2014. We explain this result by a time lag between
827 primary production in the water column and the export of the produced organic material to the
828 seafloor, which was probably even more delayed during stratification. Such a delay was observed in a
829 previous study (Bange et al., 2010), revealing enhanced water methane concentration close to the
830 seafloor approximately one month after the chlorophyll maximum. The C/N ratio (averaged over 0-5
831 cmbsf) also showed no -correlation with integrated [surface](#)-methanogenesis [from the same depth](#)
832 [layer](#) (0-5 cmbsf), which is surprising as we expected that a higher C/N ratio, indicative for less labile
833 organic carbon, should have a negative effect on non-competitive methanogenesis. However,
834 methanogens are not able to directly use most of the labile organic matter due their inability to
835 process large molecules (more than two C-C bondings) (Zinder, 1993). Methanogens are dependent
836 on other microbial groups to degrade large organic compounds (e.g. amino acids) for them (Zinder,
837 1993). Because of this substrate speciation and dependence, a delay between the sedimentation of
838 fresh, labile organic matter and the increase in methanogenesis can be expected, which would not be
839 captured by the applied PCA.

840 **b) Effect of POC and C/N ratio over 0-30 cmbsf**

841 In the PCA for the [surface](#)-sediment profiles [from the sulfate reduction zone](#) (0-30 cmbsf), POC
842 showed a negative correlation with methanogenesis and sediment depth, while C/N ratio showed a

843 positive correlation with methanogenesis and sediment depth (Fig 11.). Given that POC remained
844 basically unchanged over the top 30 cmbsf, with the exemption of the topmost sediment layer, its
845 negative correlation with methanogenesis is probably solely explained by the increase of
846 methanogenesis with sediment depth, and can therefore be excluded as a major controlling factor.
847 As sulfate in this zone was likely never depleted to levels that are critically limiting sulfate reduction
848 (lowest concentration 1300 μM , compare e.g. with Treude et al., 2014) we do not expect a significant
849 change in the competition between methanogens and sulfate reducers. It is therefore more likely
850 that the progressive degradation of labile POC into dissolvable methanogenic substrates over depth
851 and time had a positive impact on methanogenesis. The C/N ratio indicates such a trend as the labile
852 fraction of POC decreased with depth.

853 4.3 Relevance of ~~surface~~ methanogenesis in the sulfate reduction zone of Eckernförde Bay sediments

854 The time series station Boknis Eck in Eckernförde Bay is known for being a methane source to the
855 atmosphere throughout the year due to supersaturated waters, which result from significant benthic
856 methanogenesis and emission (Bange et al., 2010). The benthic methane formation is thought to take
857 place mainly in ~~the deeper, sulfate depleted~~ sediments below the SMTZ layers (Treude et al., 2005a;
858 Whiticar, 2002).

859 In the present study, we show that surfaceSRZ methanogenesis within the sulfate zone is present
860 despite sulfate concentrations > 1 mM, a limit above which methanogenesis has been thought to be
861 negligible (Alperin et al., 1994; Hoehler et al., 1994; Burdige, 2006), and thus could contribute to
862 benthic methane emissions. In support of this hypothesis, high dissolved methane concentration in
863 the water column occurred with concomitant high surfaceSRZ methanogenesis activity (Fig. 9).

864 ~~If~~ However, whether the this observed water-column methane in the water column originated solely
865 from surfaceSRZ methanogenesis or also from gas ebullition from caused by deep methanogenesis
866 below the SMTZ, or a mixture of both, -remains speculative.

867 In fact, surface methanogenesis in the Eckernförde Bay could even increase in the future, as
868 temperature and oxygen, two important controlling factors identified for surface methanogenesis
869 (Maltby et al., 2016) and this study), are predicted to increase and decrease, respectively (Lennartz et
870 al., 2014). We will therefore have a closer look at the magnitude and potential relevance of this
871 process for the benthic methane budget and carbon cycling of Eckernförde Bay.

872 Surface methanogenesis rates determined in the present study are in a similar range of other sulfate-
873 containing, organic rich surface sediments (e.g. salt marsh sediments, sediments from the upwelling
874 region off Chile and Peru, or coastal sediments from Limfjorden, North Sea), (Table 2, References
875 herein). In comparison with methanogenesis rates below the SMTZ of organic rich sediments (i.e.,
876 coastal and upwelling systems), rates were mainly lower (2-5 times) (Table 2), which is explained by
877 the competition relief below the SMTZ, which makes more substrates available for methanogenesis.

878 ~~However, absolute surface methanogenesis rates in Eckernförde Bay sediments are in the same~~
879 ~~magnitude as deep methane production in other organic-rich sediments from the North Sea (0.076~~
880 ~~mmol m⁻² d⁻¹, Jørgensen & Parkes, 2010), or from the upwelling region off Chile (0.068–0.13 mmol m⁻²~~
881 ~~d⁻¹, Treude et al., 2005b), indicating the general importance of this process. Compared to these other~~
882 ~~sites, Eckernförde Bay features extremely high methanogenesis activity below the SMTZ, resulting in~~
883 ~~gas bubble formation and ebullition (Abegg & Anderson, 1997; Jackson et al., 1998; Treude et al.,~~
884 ~~2005a).~~
885 ~~We also performed a comparison between surface (0–30 cmbsf) and deep (below the SMTZ) net~~
886 ~~methanogenesis for the present study site to investigate the relevance of surface methanogenesis in~~
887 ~~Eckernförde Bay sediments for the overall benthic methane budget. In the gravity core of September~~
888 ~~2013, the SMTZ was situated between 45 and 76 cmbsf (Fig. 4). The methane flux was estimated~~
889 ~~according to Iversen & Jørgensen, (1993) using a sediment methane diffusion coefficient of $D_s =$~~
890 ~~$1.64 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$. The sediment diffusion coefficient was derived from the seawater methane-~~
891 ~~diffusion coefficient at 10 °C (Schulz, 2006), which was corrected by porosity according to Iversen &~~
892 ~~Jørgensen, (1992). The calculated deep methane production (1.55 mmol m⁻² d⁻¹) was similar to earlier~~
893 ~~calculated deep methanogenesis in Eckernförde Bay (0.66–1.88 mmol m⁻² d⁻¹; Treude et al., 2005a).~~
894 ~~However, integrated hydrogenotrophic methanogenesis measured in the presented study below 45~~
895 ~~cmbsf (determined by interpolation, $0.5 \pm 0.2 \text{ mmol m}^{-2} \text{ d}^{-1}$) was up to 3 times lower compared to the~~
896 ~~calculated deep methanogenesis, indicating that the interpolation missed hot spots of~~
897 ~~hydrogenotrophic methanogenesis, as alternative pathways are not predicted for this zone given the~~
898 ~~isotopic signature of methane (Whiticar, 2002). Surface methanogenesis in September 2013~~
899 ~~represented 3–8 % of deep methanogenesis. While this percentage seems low, absolute surface~~
900 ~~methanogenesis rates in Eckernförde Bay sediments are in the same magnitude as deep methane~~
901 ~~production in other organic-rich sediments from the North Sea (0.076 mmol m⁻² d⁻¹, Jørgensen &~~
902 ~~Parkes, 2010), or from the upwelling region off Chile (0.068–0.13 mmol m⁻² d⁻¹, Treude et al., 2005b),~~
903 ~~indicating the general importance of this process. Compared to these other sites, Eckernförde Bay~~
904 ~~features extremely high methanogenesis activity below the SMTZ, resulting in gas bubble formation~~
905 ~~and ebullition (Abegg & Anderson, 1997; Jackson et al., 1998; Treude et al., 2005a).~~
906 How much of ~~the~~ methane produced in the surface sediment is ~~ultimately~~ emitted into the water
907 column depends on the rate of methane consumption, i.e., aerobic and anaerobic oxidation of
908 methane in the sediment (Knittel & Boetius, 2009) (Fig. 1). In organic-rich sediments, such as in the
909 presented study, the ~~oxygenated~~ sediment layer is often only mm-thick, due to the high ~~rates~~ O_2
910 ~~demand of of~~ microorganisms during ~~bial~~ organic matter degradation, ~~which rapidly consumes~~
911 ~~oxygen~~ (Jørgensen, 2006; Preisler et al., 2007). Thus the anaerobic oxidation of methane (AOM)
912 might play a more important role ~~for methane consumption~~ in the ~~studied present study~~ Eckernförde

Formatted: Subscript

913 [Bay sediments](#). In an earlier study from [Eckernförde Bay](#) [this site](#) (Treude et al., 2005a), AOM activity
914 was detected ~~between~~ [throughout the top 0-25 cmbsf](#), ~~which~~ [which](#) included zones that were well
915 [above the actual SMTZ](#) (Treude et al., 2005a) ~~was above the expected steepest increase in methane~~
916 [concentration](#). Hence, ~~a part of the AOM zone could have been missed during sampling~~. But the
917 authors concluded that ~~the activity found~~ [methane oxidation](#) was ~~entirely~~ [completely](#) fueled by ~~deep~~
918 methanogenesis [from below sulfate penetration](#), as ~~the~~ integrated AOM rates ($0.8\text{-}1.5\text{ mmol m}^{-2}\text{ d}^{-1}$)
919 were in the same range as the predicted ~~deep~~ methane flux ($0.66\text{-}1.88\text{ mmol m}^{-2}\text{ d}^{-1}$) ~~from below into~~
920 the SMTZ.

921 Together with the data set presented here we postulate that AOM above the SMTZ ($0.8\text{ mmol m}^{-2}\text{ d}^{-1}$,
922 Treude et al., (2005a) could be partially or entirely fueled by [surfaceSRZ](#) methanogenesis. ~~In fact,~~
923 ~~such a similar close coupling between methane oxidation and methanogenesis without in the~~
924 ~~absence of definite methane profiles has been shown~~ [was recently proposed from isotopic labeling](#)
925 [experiments in with surface sediments from the sulfate reduction zone of the close-by Aarhus Bay,](#)
926 [Denmark](#) (Xiao et al., 2017). ~~Therefore it is therefore~~ [very likely that such a cryptic methane cycling](#)
927 [also occurred in the presented](#) [occurs in the sulfate reduction zone of sediments in the from](#)
928 [Eckernföerde Bay](#). If, in ~~the an~~ extreme scenario, [surfaceSRZ](#) methanogenesis would represent the
929 only methane source for AOM above the SMTZ, then [maximum surfaceSRZ](#) methanogenesis ~~is more~~
930 ~~likely could be~~ in the [range order](#) of $1.6\text{-}9\text{ mmol m}^{-2}\text{ d}^{-1}$ ($1.5\text{ mmol m}^{-2}\text{ d}^{-1}$ AOM + $0.09\text{ mmol m}^{-2}\text{ d}^{-1}$
931 net [surfaceSRZ](#) methanogenesis).

932 Even though the contribution of [surfaceSRZ](#) methanogenesis to ~~surface~~ AOM [above the SMTZ](#)
933 remains speculative, it leads to the assumption that [surfaceSRZ](#) methanogenesis could play a much
934 bigger role for benthic carbon cycling in the Eckernförde Bay than previously thought. Whether
935 [surfaceSRZ](#) methanogenesis at Eckernförde Bay has the potential for the direct emission of methane
936 into the water column goes beyond the scope of this study and should be tested in the future. ~~In fact,~~
937 ~~surface methanogenesis was found to correlate with methane concentrations in the water column~~
938 ~~near the seafloor, but at the same time this could be related to gas ebullition from below the SMTZ,~~
939 ~~which is likely a more potent methane source to the water column (Fig. 1).~~

940 5. Summary

941 The present study demonstrated that methanogenesis and sulfate reduction were concurrently
942 active within the sulfate-reducing zone in sediments at Boknis Eck (Eckernförde Bay, SW Baltic Sea).
943 The observed methanogenesis was probably based on non-competitive substrates due to the
944 competition with sulfate reducers for the substrates H_2 and acetate. Accordingly, members of the
945 family *Methanosarcinaceae*, which are known for methylotrophic methanogenesis, were found in the
946 [surface sulfate reduction zone of the](#) sediments and are likely to be responsible for the observed

947 methanogenesis ~~potentially with the potential use of using~~ non-competitive substrates such as the
948 ~~substrates~~ methanol, methylamines or methylated sulfides.
949 ~~Important Potential~~ environmental factors controlling surfaceSRZ methanogenesis ~~could beare~~
950 ~~identified as~~ POC content, C/N ratio, oxygen, and temperature, resulting in highest methanogenesis
951 ~~activity during the warm, stratified, and hypoxic months after the~~ ~~summer and autumn~~late summer
952 ~~phytoplankton blooms.~~
953 ~~This study provides new insights into the presence and seasonality of~~ surfaceSRZ methanogenesis in
954 ~~coastal sediments, and was able to showdemonstrate that surfacethe process~~ methanogenesis
955 ~~mightcould~~ play an important role ~~infor the benthic~~ methane budget and carbon cycling ~~inof~~
956 ~~Eckernfoerde Bay~~ sediments, e.g., by directly fueling AOM above the SMTZ.

957
958 ~~An important factor controlling surface methanogenesis in the upper 0-5 cmbsf was the POC content,~~
959 ~~resulting in highest methanogenesis activity after summer and autumn phytoplankton blooms.~~
960 ~~Increased stratification (indicated by salinity) was also found to be beneficial for surface~~
961 ~~methanogenesis, as it leads to the decline of oxygen below the pycnocline. Accordingly, oxygen~~
962 ~~depletion during later summer showed a positive correlation with surface methanogenesis, enabling~~
963 ~~more organic matter to reach the seafloor and providing a larger habitable anoxic zone for~~
964 ~~methanogens in the surface sediment.~~
965 ~~With increasing sediment depth (0-30 cmbsf), methanogenesis rates revealed a weak positive~~
966 ~~correlation with C/N ratio, indicating that a progressive mobilization of dissolved methanogenic~~
967 ~~substrates from fermentation of organic material at greater sediment depth plays an important role~~
968 ~~for controlling non-competitive methanogenesis.~~
969 ~~Even though surface methanogenesis was low compared to methanogenesis below the SMTZ, it may~~
970 ~~play an underestimated role in the carbon cycling at Boknis Eck, e.g., by directly fueling AOM above~~
971 ~~the SMTZ.~~

972 Author Contribution

973 J.M. and T.T. designed the experiments. J.M. carried out all experiments. H.W.B. coordinated
974 measurements of water column methane and chlorophyll. C.R.L. and M.A.F. conducted molecular
975 analysis. M.S. coordinated ¹³C-Isotope measurements. J.M. prepared the manuscript with
976 contributions from all co-authors.

977 Data Availability

978 Research data for the present study can be accessed via the public data repository PANGEA
979 (doi:10.1594/PANGAEA.873185).

980 **Acknowledgements**

981 We thank the captain and crew of F.S. Alkor, F.K. Littorina and F.B. Polarfuchs for field assistance. We
982 thank G. Schüssler, F. Wulff, P. Wefers, A. Petersen, M. Lange, and F. Evers for field and laboratory
983 assistance. For the geochemical analysis we want to thank B. Domeyer, A. Bleyer, U. Lomnitz, R.
984 Suhrberg, and V. Thoenissen. We thank F. Malien, X. Ma, A. Kock and T. Baustian for the O₂, CH₄, and
985 chlorophyll measurements from the regular monthly Boknis Eck sampling cruises. Further we thank
986 R. Conrad and P. Claus at the MPI Marburg for the ¹³C-Methanol measurements. This study received
987 financial support through the Cluster of Excellence “The Future Ocean” funded by the German
988 Research Foundation, through the Sonderforschungsbereich (SFB) 754, and through a D-A-CH project
989 funded by the Swiss National Science Foundation and German Research foundation (grant no.
990 200021L_138057, 200020_159878/1). Further support was provided through the EU COST Action
991 PERGAMON (ESSEM 0902), through the BMBF project BioPara (grant no. 03SF0421B) and through
992 the EU’s H2020 program (Marie Curie grant NITROX # 704272 to CRL).

993

994 **References**

- 995 Abegg, F. & Anderson, A.L. (1997). The acoustic turbid layer in muddy sediments of Eckernförde Bay
996 , Western Baltic : methane concentration , saturation and bubble characteristics. *Marine*
997 *Geology*. 137. pp. 137–147.
- 998 Alperin, M.J., Albert, D.B. & Martens, C.S. (1994). Seasonal variations in production and consumption
999 rates of dissolved organic carbon in an organic-rich coastal sediment. *Geochimica et*
1000 *Cosmochimica Acta*. 58 (22). pp. 4909–4930.
- 1001 Anon (n.d.). *Bange 1994-Methane on continental shelves.pdf*.
- 1002 Bakker, D.E., Bange, H.W., Gruber, N., Johannessen, T., Upstill-Goddard, R.C., Borges, A.V., Delille, B.,
1003 Löscher, C.R., Naqvi, S.W.A., Omar, A.M. & Santana-Casiano-J.M. (2014). Air-sea interactions of
1004 natural long-lived greenhouse gases (CO₂, N₂O, CH₄) in a changing climate. In: P. S. Liss & M. T.
1005 Johnson (eds.). *Ocean-Atmosphere Interactions of Gases and Particles*. Heidelberg: Springer-
1006 Verlag, pp. 113–169.
- 1007 Balzer, W., Pollehne, F. & Erlenkeuser, H. (1986). Cycling of Organic Carbon in a Marine Coastal
1008 System. In: P. G. Sly (ed.). *Sediments and Water Interactions*. New York, NY: Springer New York,
1009 pp. 325–330.
- 1010 Bange, H.W., Bartell, U.H., Rapsomanikis, S. & Andreae, M.O. (1994). Methane in the Baltic and North
1011 Seas and a reassessment of the marine emissions of methane. *Global Biogeochemical Cycles*. 8
1012 (4). pp. 465–480.
- 1013 Bange, H.W., Bergmann, K., Hansen, H.P., Kock, A., Koppe, R., Malien, F. & Ostrau, C. (2010).
1014 Dissolved methane during hypoxic events at the Boknis Eck time series station (Eckernförde
1015 Bay , SW Baltic Sea). *Biogeosciences*. 7. pp. 1279–1284.
- 1016 Bange, H.W., Hansen, H.P., Malien, F., Laß, K., Karstensen, J., Petereit, C., Friedrichs, G. & Dale, A.
1017 (2011). Boknis Eck Time Series Station (SW Baltic Sea): Measurements from 1957 to 2010.
1018 *LOICZ-Affiliated Activities*. Inprint 20. pp. 16–22.
- 1019 Bertics, V.J., Löscher, C.R., Salonen, I., Dale, A.W., Gier, J., Schmitz, R.A. & Treude, T. (2013).

1020 Occurrence of benthic microbial nitrogen fixation coupled to sulfate reduction in the seasonally
1021 hypoxic Eckernförde Bay, Baltic Sea. *Biogeosciences*. 10 (3). pp. 1243–1258.

1022 Blake, L.I., Tveit, A., Øvreås, L., Head, I.M. & Gray, N.D. (2015). *Response of Methanogens in Arctic*
1023 *Sediments to Temperature and Methanogenic Substrate Availability*.

1024 Buckley, D.H., Baumgartner, L.K. & Visscher, P.T. (2008). Vertical distribution of methane metabolism
1025 in microbial mats of the Great Sippewissett Salt Marsh. *Environmental microbiology*. 10 (4). pp.
1026 967–77.

1027 Burdige, D.J. (2006). *Geochemistry of Marine Sediments*. New Jersey, U.S.A.: Princeton University
1028 Press.

1029 Cicerone, R.J. & Oremland, R.S. (1988). Biogeochemical aspects of atmospheric methane. *Global*
1030 *Biogeochemical Cycles*. 2 (4). pp. 299–327.

1031 Crill, P. & Martens, C. (1983). Spatial and temporal fluctuations of methane production in anoxic
1032 coastal marine sediments. *Limnology and Oceanography*. 28. pp. 1117–1130.

1033 Crill, P.M. & Martens, C.S. (1986). Methane production from bicarbonate and acetate in an anoxic
1034 marine sediment. *Geochimica et Cosmochimica Acta*. 50. pp. 2089–2097.

1035 Dale, a. W., Bertics, V.J., Treude, T., Sommer, S. & Wallmann, K. (2013). Modeling benthic–pelagic
1036 nutrient exchange processes and porewater distributions in a seasonally hypoxic sediment:
1037 evidence for massive phosphate release by Beggiatoa? *Biogeosciences*. 10 (2). pp. 629–651.

1038 Denman, K.L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P.M., Dickinson, R.E., Hauglustaine, D.,
1039 Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., da Silva Dias, P.L., Wofsy, S.C.
1040 & Zhang, X. (2007). Couplings Between Changes in the Climate System and Biogeochemistry. In:
1041 S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, & H. L. Miller
1042 (eds.). *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the*
1043 *Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge,
1044 United Kingdom and New York, NY, USA: Cambridge University Press.

1045 EPA (2010). *Methane and nitrous oxide emissions from natural sources*. Washington, DC, USA.

1046 Ferdelman, T.G., Lee, C., Pantoja, S., Harder, J., Bebout, B.M. & Fossing, H. (1997). Sulfate reduction
1047 and methanogenesis in a Thioploca-dominated sediment off the coast of Chile. *Geochimica et*
1048 *Cosmochimica Acta*. 61 (15). pp. 3065–3079.

1049 Gier, J., Sommer, S., Löscher, C.R., Dale, A.W., Schmitz, R.A. & Treude, T. (2016). Nitrogen fixation in
1050 sediments along a depth transect through the Peruvian oxygen minimum zone. *Biogeosciences*.
1051 13 (14). pp. 4065–4080.

1052 Grasshoff, K., Ehrhardt, M. & Kremmling, K. (1999). *Methods of Seawater Analysis*. Weinheim: Verlag
1053 Chemie.

1054 Hansen, H.-P., Giesenhausen, H.C. & Behrends, G. (1999). Seasonal and long-term control of bottom-
1055 water oxygen deficiency in a stratified shallow-water coastal system. *ICES Journal of Marine*
1056 *Science*. 56. pp. 65–71.

1057 Hartmann, D.L., Klein Tank, A.M.G., Rusticucci, M., Alexander, L.V., Brönnimann, S., Charabi, Y.,
1058 Dentener, F.J., Dlugokencky, D.R., Easterling, D.R., Kaplan, A., Soden, B.J., Thorne, P.W., Wild,
1059 M. & Zhai, P.M. (2013). Observations: Atmosphere and Surface. In: *Climate Change 2013: The*
1060 *Physical Science Basis. Contribution Group I to the Fifth Assessment Report of the*
1061 *Intergovernmental Panel on Climate Change*. United Kingdom and New York, NY, USA:
1062 Cambridge University Press.

1063 Hoehler, T.M., Alperin, M.J., Albert, D.B. & Martens, C.S. (1994). Field and laboratory studies of
1064 methane oxidation in an anoxic marine sediment: Evidence for a methanogen-sulfate reducer
1065 consortium. *Global Biogeochemical Cycles*. 8 (4). pp. 451–463.

- 1066 Holmer, M. & Kristensen, E. (1994). Coexistence of sulfate reduction and methane production in an
1067 organic-rich sediment. *Marine Ecology Progress Series*. 107. pp. 177–184.
- 1068 Jackson, D.R., Williams, K.L., Wever, T.F., Friedrichs, C.T. & Wright, L.D. (1998). Sonar evidence for
1069 methane ebullition in Eckernförde Bay. *Continental Shelf Research*. 18. pp. 1893–1915.
- 1070 Jørgensen, B.B. (2006). Bacteria and marine Biogeochemistry. In: H. D. Schulz & M. Zabel (eds.).
1071 *Marine Geochemistry*. Berlin/Heidelberg: Springer-Verlag, pp. 173–207.
- 1072 Jørgensen, B.B. & Parkes, R.J. (2010). Role of sulfate reduction and methane production by organic
1073 carbon degradation in eutrophic fjord sediments (Limfjorden, Denmark). *Limnology and*
1074 *Oceanography*. 55 (3). pp. 1338–1352.
- 1075 Keltjens, J.T. & Vogels, G.D. (1993). Conversion of methanol and methylamines to methane and
1076 carbon dioxide. In: J. G. Ferry (ed.). *Methanogenesis: Ecology, Physiology, Biochemistry &*
1077 *Genetics*. Chapman & Hall, pp. 253–303.
- 1078 King, G.M., Klug, M.J. & Lovley, D.R. (1983). Metabolism of acetate, methanol, and methylated
1079 amines in intertidal sediments of lowes cove, maine. *Applied and environmental microbiology*.
1080 45 (6). pp. 1848–1853.
- 1081 Knittel, K. & Boetius, A. (2009). Anaerobic oxidation of methane: progress with an unknown process.
1082 *Annual review of microbiology*. 63. pp. 311–34.
- 1083 Kuivila, K.M., Murray, J.W. & Devol, a. H. (1990). Methane production in the sulfate-depleted
1084 sediments of two marine basins. *Geochimica et Cosmochimica Acta*. 54. pp. 403–411.
- 1085 Lennartz, S.T., Lehmann, A., Herrford, J., Malien, F., Hansen, H.-P., Biester, H. & Bange, H.W. (2014).
1086 Long-term trends at the Boknis Eck time series station (Baltic Sea), 1957–2013: does climate
1087 change counteract the decline in eutrophication? *Biogeosciences*. 11 (22). pp. 6323–6339.
- 1088 Maltby, J., Sommer, S., Dale, A.W. & Treude, T. (2016). Microbial methanogenesis in the sulfate-
1089 reducing zone of surface sediments traversing the Peruvian margin. *Biogeosciences*. 13. pp.
1090 283–299.
- 1091 Martens, C.S., Albert, D.B. & Alperin, M.J. (1998). Biogeochemical processes controlling methane in
1092 gassy coastal sediments---Part 1 . A model coupling organic matter flux to gas production ,
1093 oxidation and transport. *Continental Shelf Research*. 18. pp. 14–15.
- 1094 Martens, C.S. & Klump, J. V (1984). Biogeochemical cycling in an organic-rich coastal marine basin 4.
1095 An organic carbon budget for sediments dominated by sulfate reduction and methanogenesis.
1096 *Geochimica et Cosmochimica Acta*. 48. pp. 1987–2004.
- 1097 Naqvi, S.W. a., Bange, H.W., Fariás, L., Monteiro, P.M.S., Scranton, M.I. & Zhang, J. (2010). Marine
1098 hypoxia/anoxia as a source of CH₄ and N₂O. *Biogeosciences*. 7 (7). pp. 2159–2190.
- 1099 Oremland, R.S. (1988). Biogeochemistry of methanogenic bacteria. In: A. J. B. Zehnder (ed.). *Biology*
1100 *of Anaerobic Microorganisms*. New York: J. Wiley & Sons, pp. 641–705.
- 1101 Oremland, R.S. & Capone, D.G. (1988). Use of specific inhibitors in biogeochemistry and microbial
1102 ecology. In: K. C. Marshall (ed.). *Advances in Microbial Ecology*. Advances in Microbial Ecology.
1103 Boston, MA: Springer US, pp. 285–383.
- 1104 Oremland, R.S., Marsh, L.M. & Polcin, S. (1982). Methane production and simultaneous sulfate
1105 reduction in anoxic, salt-marsh sediments. *Nature*. 286. pp. 143–145.
- 1106 Oremland, R.S. & Polcin, S. (1982). Methanogenesis and Sulfate Reduction : Competitive and
1107 Noncompetitive Substrates in Estuarine Sediments. *Applied and Environmental Microbiology*. 44
1108 (6). pp. 1270–1276.
- 1109 Orsi, T.H., Werner, F., Milkert, D., Anderson, a. L. & Bryant, W.R. (1996). Environmental overview of
1110 Eckernförde Bay, northern Germany. *Geo-Marine Letters*. 16 (3). pp. 140–147.
- 1111 Penger, J., Conrad, R. & Blaser, M. (2012). Stable carbon isotope fractionation by methylotrophic

- 1112 methanogenic archaea. *Applied and environmental microbiology*. [Online]. 78 (21). pp. 7596–
 1113 602. Available from:
 1114 <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3485729&tool=pmcentrez&render>
 1115 [type=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3485729&tool=pmcentrez&render). [Accessed: 13 October 2014].
- 1116 Pimenov, N., Davidova, I., Belyaev, S., Lein, A. & Ivanov, M. (1993). Microbiological processes in
 1117 marine sediments in the Zaire River Delta and the Benguela upwelling region. *Geomicrobiology*
 1118 *Journal*. 11 (3–4). pp. 157–174.
- 1119 Preisler, A., de Beer, D., Lichtschlag, A., Lavik, G., Boetius, A. & Jørgensen, B.B. (2007). Biological and
 1120 chemical sulfide oxidation in a Beggiatoa inhabited marine sediment. *The ISME journal*. 1 (4).
 1121 pp. 341–353.
- 1122 Reeburgh, W. (2007). Oceanic methane biogeochemistry. *Chemical Reviews*. pp. 486–513.
- 1123 Sansone, F.J. & Martens, C.S. (1981). Methane Production from Acetate and Associated Methane
 1124 Fluxes from Anoxic Coastal Sediments. *Science*. 211 (4483). pp. 707–709.
- 1125 Santoro, N. & Konisky, J. (1987). Characterization of bromoethanesulfonate-resistant mutants of
 1126 *Methanococcus voltae*: Evidence of a coenzyme M transport system. *Journal of Bacteriology*.
 1127 169 (2). pp. 660–665.
- 1128 Schlüter, M., Sauter, E., Hansen, H.-P. & Suess, E. (2000). Seasonal variations of bioirrigation in
 1129 coastal sediments: modelling of field data. *Geochimica et Cosmochimica Acta*. 64 (5). pp. 821–
 1130 834.
- 1131 Seeberg-Elverfeldt, J., Schlüter, M., Feseker, T. & Kolling, M. (2005). Rhizon sampling of porewaters
 1132 near the sediment-water interface of aquatic systems. *Limnology and Oceanography-Methods*.
 1133 3. pp. 361–371.
- 1134 Senior, E., Lindström, E.B., Banat, I.M. & Nedwell, D.B. (1982). Sulfate reduction and methanogenesis
 1135 in the sediment of a saltmarsh on the East coast of the United Kingdom. *Applied and*
 1136 *environmental microbiology*. 43 (5). pp. 987–996.
- 1137 Smetacek, V. (1985). The Annual Cycle of Kiel Bight Plankton: A Long-Term Analysis. *Estuaries*. 8
 1138 (June). pp. 145–157.
- 1139 Smetacek, V., von Bodungen, B., Knoppers, B., Peinert, R., Pollehne, F., Stegmann, P. & Zeitzschel, B.
 1140 (1984). Seasonal stages characterizing the annual cycle of an inshore pelagic system. *Rapports*
 1141 *et Proces-Verbaux des Reunions Conseil International pour l'Exploration de la Mer*. 186. pp.
 1142 126–135.
- 1143 Smith, M.R. & Mah, R. a. (1981). 2-Bromoethanesulfonate: A selective agent for isolating
 1144 resistant *Methanosarcina* mutants. *Current Microbiology*. 6 (5). pp. 321–326.
- 1145 Thießen, O., Schmidt, M., Theilen, F., Schmitt, M. & Klein, G. (2006). Methane formation and
 1146 distribution of acoustic turbidity in organic-rich surface sediments in the Arkona Basin, Baltic
 1147 Sea. *Continental Shelf Research*. 26 (19). pp. 2469–2483.
- 1148 Treude, T., Krause, S., Maltby, J., Dale, A.W., Coffin, R. & Hamdan, L.J. (2014). Sulfate reduction and
 1149 methane oxidation activity below the sulfate-methane transition zone in Alaskan Beaufort Sea
 1150 continental margin sediments: Implications for deep sulfur cycling. *Geochimica et*
 1151 *Cosmochimica Acta*. 144. pp. 217–237.
- 1152 Treude, T., Krüger, M., Boetius, A. & Jørgensen, B.B. (2005a). Environmental control on anaerobic
 1153 oxidation of methane in the gassy sediments of Eckernförde Bay (German Baltic). *Limnology*
 1154 *and Oceanography*. 50 (6). pp. 1771–1786.
- 1155 Treude, T., Niggemann, J., Kallmeyer, J., Wintersteller, P., Schubert, C.J., Boetius, A. & Jørgensen, B.B.
 1156 (2005b). Anaerobic oxidation of methane and sulfate reduction along the Chilean continental
 1157 margin. *Geochimica et Cosmochimica Acta*. 69 (11). pp. 2767–2779.

- 1158 Treude, T., Smith, C.R., Wenzhöfer, F., Carney, E., Bernardino, A.F., Hannides, A.K., Krgüer, M. &
1159 Boetius, A. (2009). Biogeochemistry of a deep-sea whale fall: Sulfate reduction, sulfide efflux
1160 and methanogenesis. *Marine Ecology Progress Series*. 382. pp. 1–21.
- 1161 Welschmeyer, N.A. (1994). Fluorometric analysis of chlorophyll a in the presence of chlorophyll b and
1162 pheopigments. *Limnology and Oceanography*. 39 (8). pp. 1985–1992.
- 1163 Wever, T.F., Abegg, F., Fiedler, H.M., Fechner, G. & Stender, I.H. (1998). Shallow gas in the muddy
1164 sediments of Eckernförde Bay, Germany. *Continental Shelf Research*. 18. pp. 1715–1739.
- 1165 Wever, T.F. & Fiedler, H.M. (1995). Variability of acoustic turbidity in Eckernförde Bay (southwest
1166 Baltic Sea) related to the annual temperature cycle. *Marine Geology*. 125. pp. 21–27.
- 1167 Whitticar, M.J. (2002). Diagenetic relationships of methanogenesis, nutrients, acoustic turbidity,
1168 pockmarks and freshwater seepages in Eckernförde Bay. *Marine Geology*. 182. pp. 29–53.
- 1169 Widdel, F. & Bak, F. (1992). Gram-Negative Mesophilic Sulfate-Reducing Bacteria. In: A. Balows, H. G.
1170 Trüper, M. Dworkin, W. Harder, & K.-H. Schleifer (eds.). *The Prokaryotes*. New York, NY:
1171 Springer New York, pp. 3352–3378.
- 1172 Wuebbles, D.J. & Hayhoe, K. (2002). Atmospheric methane and global change. *Earth-Science Reviews*.
1173 57 (3–4). pp. 177–210.
- 1174 Xiao, K.Q., Beulig, F., Kjeldsen, K.U., Jørgensen, B.B. & Risgaard-Petersen, N. (2017). Concurrent
1175 methane production and oxidation in surface sediment from Aarhus Bay, Denmark. *Frontiers in*
1176 *Microbiology*. pp. 1–12.
- 1177 Zinder, S.H. (1993). Physiological ecology of methanogens. In: J. G. Ferry (ed.). *Methanogenesis*. New
1178 York, NY: Chapman & Hall, pp. 128–206.

1179

1180 **Figure Captions**

1181 **Figure 1:** Overview of processes relevant for benthic methane production, consumption, and
1182 emission in the Eckernförde Bay. The thickness of arrows for emissions and coupling between surface
1183 processes indicates the strength of methane supply. Note that this figure combines existing
1184 knowledge with results from the present study. See discussion for more details.

1185 **Figure 2:** Parameters measured in the water column and sediment in the Eckernförde Bay at each
1186 sampling month in the year 2013. Net methanogenesis (MG) and hydrogenotrophic (hydr.)
1187 methanogenesis rates are shown in triplicates with mean (solid line).

1188 **Figure 3:** Parameters measured in the water column and sediment in the Eckernförde Bay at each
1189 sampling month in the year 2014. Net methanogenesis (MG) and hydrogenotrophic (hydr.)
1190 methanogenesis rates are shown in triplicates with mean (solid line).

1191 **Figure 4:** Parameters measured in the sediment gravity core taken in the Eckernförde Bay in
1192 September 2013. Hydrogenotrophic (hydr.) methanogenesis rates are shown in triplicates with mean
1193 (solid line).

1194 **Figure 5:** Integrated net methanogenesis (MG) rates (determined by net methane production) and
1195 hydrogenotrophic MG rates (determined by radiotracer incubation) in surface sediments (0-25
1196 cmbsf) of Eckernförde Bay for different sampled time points.

1197 **Figure 6:** Potential methanogenesis rates versus sediment depth in sediment sampled in November
1198 2013, March 2014, June 2014 and September 2014. Presented are four different types of incubations
1199 (treatments): *Control* (blue symbols) is describing the treatment with sediment plus artificial
1200 seawater containing natural salinity (24 PSU) and sulfate concentrations (17 mM), *molybdate* (green
1201 symbols) is the treatment with addition of molybdate (22 mM), *BES* (purple symbols) is the
1202 treatment with 60 mM BES addition, and *methanol* (red symbols) is the treatment with addition of 10
1203 mM methanol. Shown are triplicates per depth interval and the mean as a solid line. Please note the
1204 different x-axis for the methanol treatment (red).

1205 **Figure 7:** Development of headspace gas content and isotope composition of methane (CH₄) and
1206 carbon dioxide (CO₂), and porewater methanol (CH₃OH) concentration and isotope composition
1207 during the ¹³C-labeling experiment (with sediment from the 0-2 cmbsf horizon in September 2014)
1208 with addition of ¹³C-enriched methanol (¹³C:¹²C = 1:1000). *Figure above:* Concentrations of porewater
1209 methanol (CH₃OH) and headspace content of methane (CH₄) and carbon dioxide (CO₂) over time.
1210 *Figure below:* Isotope composition of porewater CH₃OH, headspace CH₄, and headspace CO₂ over
1211 time. Shown are means (from triplicates) with standard deviation.

1212 **Figure 8:** Sediment methane concentrations (with sediment from the 0-1 cmbsf in September 2014)
1213 over time in the treatment with addition of methanol (10 mM) are shown above. Shown are triplicate
1214 values per measurement. DNA copies of *Archaea*, *Methanosarcinales* and *Methanosarcinaceae* are
1215 shown below in duplicates per measurement. Please note the secondary y-axis for
1216 *Methanosarcinales* and *Methanosarcinaceae*. More data are available for methane (determined in
1217 the gas headspace) than from DNA samples (taken from the sediment) as sample volume for
1218 molecular analyzes was limited.

1219 **Figure 9:** Temporal development of integrated net surface methanogenesis (0-5 cmbsf) in the
1220 sediment and chlorophyll (green) and methane concentrations (orange) in the bottom water (25 m).
1221 Methanogenesis (MG) rates and methane concentrations are shown in means (from triplicates) with
1222 standard deviation.

1223 **Figure 10:** Principle component analysis (PCA) from three different angles of integrated surface
1224 methanogenesis (0-5 cmbsf) and surface particulate organic carbon averaged over 0-5 cmbsf (surface
1225 sediment POC), surface C/N ratio averaged over 0-5 cmbsf (surface sediment C/N), bottom water
1226 salinity, bottom water temperature (T), bottom water methane (CH₄), bottom water oxygen (O₂), and
1227 bottom water chlorophyll. Data were transformed into ranks before analysis. a) Correlation biplot of
1228 principle components 1 and 2, b) correlation biplot of principle components 1 and 3, c) correlation
1229 biplot of principle components 2 and 3. Correlation biplots are shown in a multidimensional space
1230 with parameters shown as green lines and samples shown as black dots. Parameters pointing into
1231 the same direction are positively related; parameters pointing in the opposite direction are
1232 negatively related.

1233
1234 **Figure 11:** Principle component analysis (PCA) from two different angles of surface methanogenesis
1235 depth profiles and sampling month (Month), sediment depth, depth profiles of particulate organic
1236 carbon (POC) and C/N ratio (C/N). Data was transformed into ranks before analysis. a) Correlation
1237 biplot of principle components 1 and 2, b) correlation biplot of principle components 1 and 3.
1238 Correlation biplots are shown in a multidimensional space with parameters shown as green lines and
1239 samples shown as black dots. Parameters pointing into the same direction are positively related;
1240 parameters pointing in the opposite direction are negatively related.

1241

1242

1243

1244 **Table 1:** Sampling months with bottom water (~ 2 m above seafloor) temperature (Temp.), dissolved
1245 oxygen (O₂) and dissolved methane (CH₄) concentration

Sampling Month	Date	Instrument	Temp. (°C)	O ₂ (μM)	CH ₄ (nM)	Type of Analysis
March 2013	13.03.2013	CTD	3	340	30	WC
		MUC				All
Juni 2013	27.06.2013	CTD	6	94	125	WC
		MUC				All
September 2013	25.09.2013	CTD	10	bdl	262*	WC
		MUC				All
		GC				GC-All
November 2013	08.11.2013	CTD	12	163	13	WC
		MUC				All
March 2014	13.03.2014	CTD	4	209	41*	WC
		MUC				All
June 2014	08.06.2014	CTD	7	47	61	WC
		MUC				All
September 2014	17.09.2014	CTD	13	bdl	234	WC
		MUC				All

1246 MUC = multicorer, GC= gravity corer, CTD = CTD/Rosette, bdl= below detection limit (5μM), All = methane gas
1247 analysis, porewater analysis, sediment geochemistry, net methanogenesis analysis, hydrogenotrophic
1248 methanogenesis analysis, GC-All= analysis for gravity cores including methane gas analysis, porewater analysis,
1249 sediment geochemistry, hydrogenotrophic methanogenesis analysis, WC= Water column analyses including
1250 methane analysis, chlorophyll analysis

1251 **Concentrations from the regular monthly Boknis Eck sampling cruises on 24.09.13 and 05.03. 14 (www.bokniseck.de)

1252

1253

1254

1255

1256

1257

1258

1259

1260 **Table 2:** Comparison of surface methanogenesis rates in shallow water marine sediments of different
 1261 geographical origin

Study site	Water depth (m)	Sediment depths (cm)	Rate (nmol cm ⁻² d ⁻¹)	Reference
<i>Sulfate-containing, organic-rich sediments</i>				
Eckernförde Bay (Baltic Sea)	28	0-25	0-1.3	Present study
Upwelling region off Peru (Pacific)	70-1025	0-25	0-1.5	(Maltby et al., 2016)
Upwelling region off Chile (Pacific)	87	0-6	0-0.6	(Ferdelman et al., 1997)
Limfjorden (North Sea)	7-10	0-100	0-0.05	(Jørgensen & Parkes, 2010)
Colne Point Saltmarsh (Essex, UK)	-	0-30	0-0.03	(Senior et al., 1982)
<i>Sulfate-depleted, organic-rich sediments (sediment depth marks the depth at which sulfate was depleted)</i>				
Eckernförde Bay (Baltic Sea)	28	>100	0.01-1.4	Present Study
Limfjorden (North Sea)	7-10	>100	0.01-3.1	(Jørgensen & Parkes, 2010)
Saanich Inlet (British Columbia, Canada)	225	>20	0.3-7.0	(Kuivila et al., 1990)
Upwelling region off Peru (Pacific)	78	>50	0-2.1	(Maltby et al., 2016)

1262

1263

1264

1265

1266


1267

1268

1269

1270 **Figures**

1271 **Figure 1**



1272

1273

1274

1275

1276

1277

1278

1279


1280


1281

1282

1283


2013





1288 **Figure 4**

1289



1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303


1304

1305

1306

1307 Figure 5

1308



1309

1310

1311

1312

1313

1314

1315

1316

1317

1318


1319

1320

1321

1322 **Figure 6**

1323



1324

1325

1326

1327

1328

1329



1330

1331

1332

1333

1334 Figure 7



1335

1336

1337

1338

1339

1340


1341

1342

1343

1344

1345 Figure 8



1346

1347

1348

1349

1350


1351

1352

1353

1354

1355 **Figure 9**



1356

1357

1358

1359

1360

1361

1362

1363

1364

1365


1366

1367


1368

1369

1370



1373 **Figure 11**



1374

1375

1376

1377

1378

1379

1380

1381