



High-frequency productivity estimates for a lake from free-water CO₂ concentration measurements

Maria Provenzale¹, Anne Ojala^{2,3}, Jouni Heiskanen⁴, Kukka-Maaria Erkkilä¹, Ivan Mammarella¹, Pertti Hari³, and Timo Vesala^{1,3}

¹Department of Physics, University of Helsinki, Helsinki, Finland

²Department of Environmental Sciences, University of Helsinki, Lahti, Finland

³Department of Forest Sciences, University of Helsinki, Helsinki, Finland

⁴ICOS ERIC Head Office, Helsinki, Finland

Correspondence to: Maria Provenzale (maria.provenzale@helsinki.fi)

Abstract. Lakes are important actors in biogeochemical cycles and a powerful natural source of CO₂. However, they are not yet fully integrated in carbon budgets, and the carbon cycle in the water is still poorly understood. In freshwater ecosystems, productivity studies have usually been carried out with traditional methods (bottle incubations, ¹⁴C technique), which are imprecise and have a poor temporal resolution. Consequently, our ability to quantify and predict the net ecosystem productivity (NEP) is limited: the estimates are prone to errors and the NEP cannot be parameterized from environmental variables. Here we expand the testing of a free-water method based on the direct measurement of the CO₂ concentration in the water. The approach was proposed already in 2008, but was tested on a very short data set (3 days) under specific conditions (autumn turnover); despite showing promising results, it has not been used ever since. We tested the method under different conditions (summer stratification, typical summer conditions for boreal dark-water lakes) and on a much longer data set (40 days), and quantitatively validated it comparing our data and productivity models. We were able to evaluate the NEP with a high temporal resolution (minutes) and found an excellent agreement with the models. We also estimated the parameters of the productivity-irradiance (PI) curves that allow the calculation of the NEP from irradiance and water temperature. Overall, our work shows that the approach is suitable for productivity studies under a wider range of conditions, and is an important step towards developing it so that it becomes even more general.

1 Introduction

Lakes are very important actors in the local and global carbon cycles (Battin et al., 2009; Tranvik et al., 2009). They both fix carbon, through the photosynthesis of the in-lake primary producers, and release it, through the respiration of all the aquatic organisms (primary producers, consumers and microbes), through photochemical reactions and by transmitting the received carbon from the catchment (lateral transport) back to the atmosphere in gaseous form (CO₂). Many lakes - especially the oligotrophic ones typical of high latitudes - are net heterotrophic systems where the rate of community respiration exceeds that of primary production (Cole et al., 1994; Sobek et al., 2003); this contributes to make lakes one of the most important natural sources of greenhouse gases (Raymond et al., 2013). However, they are not yet fully integrated in the local and global carbon



budgets, and the lacustrine carbon cycle is still poorly known (Cole et al., 2007).

In freshwater ecology, productivity studies have usually relied on the light and dark bottle method (Gaarder and Gran, 1927) and the ^{14}C labeling technique (Stemann Nielsen, 1951; Peterson, 1980; Bender et al., 1987; Søndergaard, 2000). The first provides estimates both of the gross and the net primary productivity, whereas the latter gives an estimate that is between the gross and the net productivity, depending on the incubation time. These traditional methods require time- and effort-demanding measurements and have a poor temporal resolution. Periods of high productivity are easily missed (Karl et al., 2003) and, because of the low temporal resolution, the non-linear relationship between photosynthetically active solar radiation (*PAR*) and photosynthesis cannot be properly investigated. As a consequence, carbon balances may be imprecise and for instance the net ecosystem productivity (*NEP*) cannot be parameterized robustly as a function of ambient variables. Moreover, communities enclosed in bottles experience light and nutrient conditions far from the natural ones, since the movement of water or of the organisms themselves is limited (Mallin and Paerl, 1992; Reynolds, 2006), and the results can be unrealistic. Thus, advances in the methodology are necessary, to better estimate freshwater ecosystems productivity and to expand our understanding of the carbon cycle in the water column.

In the last 15 years, free-water methods, not requiring sampling and incubation, have become more common. These methods, however, are usually based on the measurement of the O_2 concentration in the water, which is then used as a proxy for CO_2 (Hoellein et al., 2013; Solomon et al., 2013): this introduces uncertainties (Staehr et al., 2010). The respiratory quotient that has to be applied when transforming rates from O_2 to CO_2 has, in fact, large variations (Berggren et al., 2012).

To study the in-water photosynthesis and respiration, Hari et al. (2008) proposed a free-water method based on the direct measurement of the CO_2 concentration in the water with non-dispersive infra-red (NDIR) CO_2 probes, associated with a concomitant assessment of the CO_2 flux between the lake and the atmosphere. The probes are designed to measure the CO_2 concentration in the air, but by building a gas collection system the concentration in the water is obtained. Similar probes have been used also in Johnson et al. (2010), albeit not for productivity studies. The temporal resolution is of five seconds, more than a hundredfold improvement over the traditional approaches. A requirement of the method is the concomitant assessment of the CO_2 flux from the lake to the atmosphere. Information on the in-lake vertical CO_2 flux is also needed (and, ideally, on the lateral transport as well). If such data are missing the method can be applied under specific conditions (e.g. stable stratification); it still allows the parameterization of the *NEP* from *PAR* and water temperature, from which the *NEP* can then be calculated under different conditions.

In Hari et al. (2008), the method was tested on a small boreal lake in Finland over three days only, during the autumn turnover. A cross-comparison was carried out between different measurement methods, but the *NEP* was not mathematically parameterized and the method was not quantitatively verified. Despite the very short data set and the specific conditions, the results were promising: the relationship between *PAR* and *NEP* was clearly visible, the measured respiration rate was 16 times higher than with the bottle method and the measured productivity was 5 times higher than with the ^{14}C technique. The numbers are in line with previous studies: Pace and Prairie (2005) reported similar discrepancies between an oxygen-based free-water approach and the bottle method in small lakes in Michigan, and a tendency of the ^{14}C method to underestimate the productivity is well known (Howarth and Michaels, 2000). However, the method has been overlooked and has not been used for productivity



calculations since 2008, possibly because of the limited testing.

Here we tested the method of Hari et al. (2008) on a different boreal lake, under different conditions and on a much longer data set, quantitatively verifying it. We continuously collected data for four summers, and then we focused on the periods when the lake was stably stratified, i.e. summer conditions typical of boreal dark-water lakes, in order to rule out the lateral CO₂ flux and the CO₂ flux from the deeper layers of the lake. Overall, we analysed 40 days of data. We calculated the *NEP* using the equations that are typically used in forest ecology, where high-frequency measurements are more common, in an effort of harmonizing the procedures between different fields. Once we had the *NEP* with a high temporal resolution, we verified the relationship between the *NEP* and irradiance, using a saturating Michaelis-Menten model. We found an excellent agreement between the data and the model. From that, we could also estimate the parameters of the productivity-irradiance (PI) curves, specific to the in-lake communities. These parameters are very important because they allow the calculation of the *NEP* from *PAR* and water temperature.

Whilst our efforts were mainly focused on method testing and development, we also checked whether the parameters of the PI curves we estimated changed significantly between the years. Our goal was to gather information on how sensitive the parameters are to variations in the communities living in the lake or in the environmental conditions. We investigated whether their behaviour could be related to their main drivers, water temperature and irradiance.

2 Materials and procedures

2.1 Study site

The study site is the boreal lake Kuivajärvi, in southern Finland (61°50.743' N, 24°17.134' E). Lake Kuivajärvi is typical dark-water boreal lake. It is small and oblong, and it is surrounded by managed coniferous forests. Its surface area is 0.62 km² and its length is 2.6 km; its mean depth and maximum depth are 6.3 m and 13.2 m respectively. The lake is humic (surface median DOC concentration = 11.8 mg l⁻¹ in 2011) and mesotrophic (surface median annual total nitrogen concentration = 370 μg l⁻¹ and annual total phosphorus concentration = 14 μg l⁻¹ in 2011), with a chlorophyll *a* concentration in the surface layer usually between 3 and 5 μg l⁻¹ (median 4.8 μg l⁻¹ in 2011), with summer values that can reach 30 μg l⁻¹ (Miettinen et al., 2015). The lake is dark coloured: the Secchi depth ranges from 1.2 to 1.5 m (Heiskanen et al., 2015). The lake is dimictic and it is frozen for five months every year on average; the spring turnover occurs immediately after the ice out in late April or early May, and after the turnover a thermocline starts developing. The thermocline deepens until the autumn turnover, and finally the lake freezes over in late November or early December (Heiskanen et al., 2015; Mammarella et al., 2015). A map with the location and bathymetry of the lake is available in the supplemental information (Fig. S1 and S2).

2.2 Measurements

All the instruments were mounted on a raft, which was moored in the middle of the lake (see the supplemental information, Fig. S2, for the exact position of the raft on the lake). To measure the CO₂ concentration in the water, a closed system consisting of



a NDIR probe (CARBOCAP® GMP343, Vaisala Oyj, Vantaa, Finland) for the CO₂ concentration in the air, gas impermeable tubes (stainless steel and teflon) and a submerged gas permeable tube (silicone rubber, Rotilabo 9572.1, Carl Roth GmbH and Co. KG, Karlsruhe, Germany) was built; the air was circulated continuously in the system by a diaphragm pump (KNF Neuberger Micro gas pump, KNF Neuberger AB, Stockholm, Sweden). Analog voltage outputs were used, logged with a Nokeval
 5 RMD680 serial transmitter to a ASCII-file on a Windows-based computer. Since silicone rubber has an excellent permeability to CO₂ (Carignan, 1998; Hari et al., 2008), the concentration of CO₂ in the air circulating in the system equilibrated with that in the water around the submerged tube. Hence, the CO₂ concentration in the water could be obtained from that in the air using the dependence of CO₂ solubility on temperature and pressure. The CO₂ concentration in the water C_{CO_2} (dissolved CO₂), in $\mu\text{mol m}^{-3}$, was calculated as

$$10 \quad C_{\text{CO}_2} = \chi_{\text{CO}_2} P K_{\text{H}}, \quad (1)$$

where χ_{CO_2} is the CO₂ gas phase mole fraction in the tube measured by the probe (in $\mu\text{mol mol}^{-1}$), P is the total air pressure inside the system and K_{H} is Henry's law constant (temperature dependent). For more details on the setup see Hari et al. (2008), Heiskanen et al. (2014) and the supplemental information (Fig. S3). The CO₂ concentration in the water was measured at a depth of 0.2 m (determined by the depth of the submerged silicone tube). The system was operating continuously from May to
 15 September 2010-2014, but the data from year 2012 are not used here due to technical problems. The silicone tube was cleaned once a week to avoid biofouling and changed once a month. The CO₂ sensors were calibrated using span and zero gases. A thermistor chain of 16 Pt100 resistance thermometers (depths: 0.2, 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 6.0, 7.0, 8.0, 10.0, and 12.0 m) was deployed and a *PAR* sensor (LI-192, LI-COR Inc., Nebraska, USA) for photosynthetic photon flux density (*PPFD*) was submerged in the water at the same depth as the CO₂ measurement (0.2 m). An eddy covariance (EC)
 20 system (with ultrasonic anemometer USA-1, Metek GmbH, Germany, and closed-path infra-red gas analyzer LI-7000, LI-COR Inc., Nebraska, USA, replaced in 2011 by enclosed-path infra-red gas analyzer LI-7200, LI-COR Inc., Nebraska, USA) was used to detect the CO₂ flux between the lake and the atmosphere. The fluxes were calculated and quality screened according to the standard procedures, see Vesala et al. (2006), Mammarella et al. (2009), Mammarella et al. (2015) and the supplemental information (Sect. S2). All the instruments were powered by mains electricity.

25 2.3 Calculation of the net ecosystem productivity

The net ecosystem productivity (NEP , $\mu\text{mol}(\text{CO}_2)\text{m}^{-2}\text{s}^{-1}$), also called net ecosystem uptake, can be defined as

$$NEP = GPP - R_{\text{h}}, \quad (2)$$

where GPP (gross primary productivity) is the amount of carbon fixed by the primary producers through photosynthesis and R_{h} (ecosystem respiration) is the amount of carbon lost through respiration, both autotrophic and heterotrophic. The NEP is
 30 the opposite of the net ecosystem exchange (NEE), whose expression can be derived from the conservation of mass. Hence, considering the mass balance of CO₂ in the mixed layer of the lake, where most of the photosynthesis takes place, and assuming



that lateral transport of CO₂ is of no importance, the *NEP* can also be expressed as

$$NEP = -NEE = - \int_{-h_{\text{mix}}}^0 \frac{\partial C_{\text{CO}_2}(z, t)}{\partial t} dz - F_a + F_u; \quad (3)$$

see Fig. 1 for a schematic representation. In Eq. (3), C_{CO_2} is the CO₂ concentration in the water calculated from Eq. (1), F_a is the CO₂ flux between the lake and the atmosphere (positive if from the lake to the atmosphere), F_u is the CO₂ flux between the deeper and the mixed layer of the lake (positive if upwards), t is time and z is depth. The integration is computed between the mixing depth h_{mix} and the surface. The mixing depth is defined as the depth at which the water temperature starts decreasing faster than one degree per meter (Staehr et al., 2010); in our study the average value was $h_{\text{mix}} = 1.5$ m. Given the dark water colour and the resulting low light conditions in the lake, there was no benthic primary production in the profundal zone. While C_{CO_2} was measured by the probe and F_a by the EC system, we had no precise way of measuring F_u . This is the main reason behind our choice to limit the analysis to the summer days when the lake was stably stratified and it was safe to assume no gas was exchanged through the thermocline: $F_u = 0$. The periods of stable stratifications were chosen on the basis of temperature plots and of the Schmidt stability of the lake, calculated with the LakeAnalyzer program, according to Read et al. (2011). For all the chosen days, the stability (Sc) is $> 100 \text{ Jm}^{-2}$. However, not all days with $Sc > 100 \text{ Jm}^{-2}$ were used: days with strong winds or stable atmospheric stratification were discarded because of their impact on fluxes (for more detailed information, see the end of this section). For the time series of isotherms for the whole summers (from 1 June to 31 August), and the time series of isotherms, Schmidt stability, CO₂ concentration and *PAR* at 0.2 m and air temperature for the periods of stable stratification chosen for analysis each year see the supplemental information (Figg. S4-S14). Overall, we analyzed 40 days in 10 periods occurring between mid-June and the end of July of each year.

It is worth pointing out that Eq. (3) resembles the equation used in terrestrial ecology to estimate the *NEP*. In fact, e.g. in forest EC calculations (Foken et al., 2012), neglecting lateral transport, the *NEP* is

$$NEP = -NEE = - \int_0^{h_m} \rho_d \frac{\partial \chi_{\text{CO}_2}(z, t)}{\partial t} dz - \rho_d \overline{w' \chi'_{\text{CO}_2}}. \quad (4)$$

In Eq. (4), $\rho_d \chi_{\text{CO}_2}$ (ρ_d = dry air density, χ_{CO_2} = CO₂ mixing ratio) replaces C_{CO_2} as the CO₂ concentration in the air instead of in the water, and z is the height (with h_m = measuring height); $\rho_d \overline{w' \chi'_{\text{CO}_2}}$ is F_a , the CO₂ flux from the forest to the atmosphere, calculated as the covariance between the fluctuations of the vertical wind velocity and the gas mixing ratio. From the analogy, we can see that using high-frequency CO₂ concentration measurements and Eq. (3) in aquatic ecology reduces the gap with terrestrial ecology, where high-frequency measurements are common, and leads towards a greater uniformity in the equations and procedures.

Resuming our calculation of the *NEP* in aquatic ecosystems through Eq. (3), to increase the precision of the concentration data, half-hourly averages of C_{CO_2} from the raw 5-sec data were used. A 30-min resolution is enough to capture the variations caused by the biological activity and at the same time filter out the ones caused by the physical mixing of the water (Staehr et al., 2010). However, the EC data set, which also has a resolution of 30 minutes, had many gaps, due to inherent problems



of the EC technique (wind not blowing along the lake, stability or not fully developed turbulence resulting in quality criteria not met) and technical problems (instrument failures). Approximately 70% of the data points for the summers were rejected or missing, with occurrences of consecutive days having no acceptable data points at all. Hence, for our data set, a point by point calculation of F_a in (3) was not possible. Even though in general it would not be needed, we had to use a daytime and a nighttime average value for F_a ; we maintained the half-hourly calculation of the NEP to preserve the temporal resolution. The daytime and nighttime average F_a values were calculated separately for each year, combining all the studied periods of water stable stratification of the same summer. Before doing so, we checked that the environmental conditions (temperature and relative humidity cycles, incoming radiation, wind speed and direction, atmospheric stability) were similar for all the analysed days in the summer. In particular, since wind and atmospheric stability have the greatest influence on the fluxes (given that the lake water is thermally stratified), as verified in Heiskanen et al. (2014), we discarded any day with winds $> 5 \text{ m s}^{-1}$ or with stable atmospheric stratification. For the remaining days, the wind was always weak, with averages $< 2.5 \text{ m s}^{-1}$; at such low speeds, the influence of the wind on the flux is negligible (Cole and Caraco, 1998). Under these circumstances (i.e. warm and sunny summer days without strong wind events), the CO_2 flux is expected to have similar daily cycles across the analysed days. Day and night were defined on the basis of PAR . When using PAR , we are referring to the average PAR value in the mixed layer, obtained from the 0.2 m value through the lake light extinction coefficient (1.5). The threshold between day and night was set to $20 \mu\text{mol}(\text{ph})\text{m}^{-2}\text{s}^{-1}$ and it was chosen by calculating the average value of PAR at which the CO_2 concentration in the water started decreasing in the morning after accumulating during the night, or increasing again in the evening. Using this procedure, “day” represents the fraction of the time series when photosynthesis dominates over respiration, and not the times when photosynthesis takes place in absolute terms.

At this point, we were able to calculate the half-hourly values of NEP for each period.

2.4 Relationship between NEP and PAR

In humic lakes, the photosynthesis is strongly driven by PAR , and the relationship can be described for instance by the Michaelis-Menten equation (Caperon, 1967; Kiefer and Mitchell, 1983). Assuming that the daytime respiration rate equals the nighttime respiration rate and that they depend exponentially on temperature (Carignan et al., 2000), the NEP can be expressed as

$$NEP = GPP - R_h = \frac{p_{\max} PAR}{PAR + b} - r_0 Q_{10}^{T/10}. \quad (5)$$

In (5), T is the water temperature (in $^{\circ}\text{C}$) and Q_{10} is a non-dimensional temperature coefficient whose generally accepted value for freshwater communities is 2 (in the literature, values between 1.88 and 2.19 are reported: Reynolds (1984); Raven and Geider (1988); Davison (1991)); p_{\max} , b and r_0 represent the maximum potential photosynthetic rate, the half-saturation constant (i.e. the value of PAR at which the photosynthetic rate is half of the maximum rate) and the basal respiration rate, respectively. These parameters are important, since they allow the calculation of NEP from water temperature and PAR . After calculating the NEP , we plotted the NEP versus irradiance curves. We then fitted the NEP data to the model (Eq. (5))



with the least-squares fitting method, in order to check the agreement between the data and the model and in order to estimate p_{\max} , b and r_0 .

3 Assessment

The NEP had the same trend as the incoming radiation, as expected; it had bigger negative values during the night, when only respiration took place, and smaller negative values during the day, when photosynthesis contributed with an uptake of CO_2 . Anyhow, the net productivity values are almost always negative, meaning that the ecosystem, overall, is heterotrophic and a source of CO_2 . Figure 2 shows the CO_2 concentration change in time over the mixed layer (the first term in Eq. (3)), which is usually referred to as storage flux in forest ecology calculations, the NEP , the average daytime and nighttime values of the CO_2 flux (F_{aday} and F_{anight}) and PAR for a sample period of stable stratification in July 2010, representative of the analyzed periods. The nine-day period in Fig. 2 is the longest of the entire data set. Generally, stable stratification lasted from two to five days; its short duration is due to the oblong shape of the lake, that makes it sensitive to wind action: as soon as the wind increases the mixing is enhanced (although complete mixing takes place only in spring and autumn). Figure 3 displays the NEP versus PAR . We decided to draw a different plot for each year, instead of combining all the data points from all the years, since the conditions (PAR and water T) varied from year to year. The figure also displays the model curve, calculated using the average water T of the studied periods of each year. From the plots, we can see that for low values of PAR the NEP was strongly negative; then, as PAR increased, the NEP quickly increased as well; however, as already noted, the NEP always remained negative, indicating net heterotrophy. None of the years exhibited signs of photoinhibition: the NEP did not seem to decrease even at high ($> 500 \mu\text{mol}(\text{ph})\text{m}^{-2}\text{s}^{-1}$) values of PAR . In case of photoinhibition, the addition of a specific term accounting for it in (5) would have been needed, but here it was not necessary. Differences can be seen between the years, with 2014 showcasing the smallest values of NEP . Year 2014 was particularly hot, so the strongly negative NEP can be due to increased respiration rates; year 2010 though displays the highest values of NEP despite having an intermediate average water temperature. Figure 4 shows the same plots, but focuses on the dependence of the NEP on T . The model is calculated for different values of water T , ranging from the minimum to the maximum water temperatures recorded during the studied periods of each year. The NEP decreases with increasing temperature, due to higher respiration rates. Finally, Fig. 5 features 3D plots of the data and the curves, to visualize simultaneously the dependence of the NEP on PAR and water T . The curves in Figg. 2-5 have the expected trends, and this confirms that the data and the equation used are proper tools for estimating the NEP at a high temporal resolution. The results of the fittings of the NEP versus PAR and T are reported in Table 1 below. Considering the assumptions we had to adopt, there is a very good agreement between the model and the data: the R^2 values range from 0.71 to 0.84. This clearly indicates that the method used here allows the NEP to be accurately parameterized as a function of irradiance and water temperature.

An in-depth analysis of the values of the model parameters (reported in Table 1) and their inter-annual variability is beyond the scope of this paper. However, some comments are possible. The maximum photosynthetic rate p_{\max} ranged between 1.55 (2014) and 0.63 (2013) $\mu\text{mol}(\text{CO}_2)\text{m}^{-2}\text{s}^{-1}$, and it was higher in 2011 and 2014 than in 2010 and 2013. The half-saturation



constant b ranged between 22 (2010) and 33 (2013) $\mu\text{mol}(\text{ph})\text{m}^{-2}\text{s}^{-1}$, being higher in 2011, 2013 and 2014 than in 2010. The values of b are relatively small. It indicates that the phytoplankton communities were well adapted to the low light conditions (boreal area and dark-water lake) and were able to start photosynthesising even when the incoming radiation was small. The basal respiration r_0 ranged between 0.228 (2010) and 0.482 (2014) $\mu\text{mol}(\text{CO}_2)\text{m}^{-2}\text{s}^{-1}$, being higher in 2011 and 2014 than in 2010 and 2013, as was the case with p_{max} . The parameters however do not appear to be strictly correlated to each other, and a clear and uniform pattern in their behavior cannot be identified.

We checked whether the differences in the parameter values between the years are statistically significant. This also gives an indication whether the parameters should be re-assessed each year or can be considered lake-specific. We calculated the parameters difference and its confidence interval (calculated as the uncertainty of the difference, from the confidence intervals of the parameters themselves), and verified whether it overlapped 0. The value of b does not change significantly between any of the years: this means that the algal communities adapted to the light conditions in a similar way every year. The values of the other parameters change: p_{max} is comparable only between 2011 and 2014, and r_0 is never comparable. The difference in p_{max} and r_0 can be due to different total algal biomass in the lake. In general, we can say that there are statistically significant differences between the years. Variations in the environmental conditions might have led to changes in the communities living in the lake, or the communities might have responded differently to the environmental conditions; p_{max} and r_0 seem to be more sensitive to variations than b .

Finally, we investigated whether the changes of the model parameters can be explained in terms of changes, during the analyzed periods, of the ambient variables that act as *NEP* drivers: water temperature and irradiance. The model parameters and the average, minimum and maximum values of water T and PAR for each year are reported in Table 2 (only the 40 analyzed days are considered in these statistics). In 2010 and 2011 the surface water temperature had similar average values, 22.9 and 22.7 °C respectively. Year 2013 was slightly colder, with an average value of 21.5 °C, while year 2014 was warmer, with an average value of 25.6 °C. The minimum temperatures of the study periods were similar for 2010 and 2013 ($\approx 20^\circ\text{C}$), slightly higher for 2011 (20.7 °C) and notably higher for 2014 (23.2 °C). The maximum temperatures ranged between 23.5 (2013) and 28.3 (2014) °C. Overall, 2013 can be considered as a cold year, 2014 as a hot year, 2010 and 2011 as intermediate years. The temperature variation pattern between the years cannot be easily linked to the variations in b . Concerning p_{max} , even though the largest value of p_{max} is associated with the warmest year (2014), and the smallest value of p_{max} with the coldest year (2013), years 2010 and 2011 had different values of p_{max} despite having similar temperatures. Besides, p_{max} and b are expected to depend more strongly on PAR than on T . Conversely, r_0 can be expected to be larger when temperatures are higher. This happened in 2011 and 2014, but not in 2010, which still had relatively high temperatures. Possible explanations are changes in the Q_{10} value, or the influence of other environmental variables. We did not investigate further possible changes in the Q_{10} value, both because we did not have an independent way to estimate it, and because its range is narrow according to the literature (Reynolds, 1984; Raven and Geider, 1988; Davison, 1991). Concerning PAR , in the analyzed periods the average values in the mixed layer ranged from 162 (2013) to 227 (2014) $\mu\text{mol}(\text{ph})\text{m}^{-2}\text{s}^{-1}$, being higher in 2010 and 2011 than in 2013, and notably higher in 2014 than in all the other years. Remarkably also in 2014, despite the high values of PAR , the communities did not show signs of photoinhibition (a PAR_{max} value of 741 for the mixed layer corresponds to a surface value



of $\approx 1900 \mu\text{mol}(\text{ph})\text{m}^{-2}\text{s}^{-1}$, given the light extinction coefficient of the lake of 1.5). Higher average PAR values could be responsible for larger p_{max} values, as observed in 2011 and 2014, and partially in 2010. However, the average PAR values are very similar in 2010 and 2011, while p_{max} values are not. Still, the very low value of p_{max} in 2013 could be explained by the low PAR_{ave} value. The variations of b between the years, though, cannot be linked to the changes in PAR : 2013 and 5 2014, despite having very different PAR values, had similar b values. The trend in r_0 also cannot be associated with the trend in PAR between the years. From what said so far, the changes of PAR and water temperature cannot fully account for the changes in the model parameters. A more extensive analysis would require more information on the algal communities living in the lake each year, and as already stated is beyond the scope of this paper.

4 Discussion

10 Firstly, it is important to notice that we are working under the assumption that the NEE , which is what can be measured, is equal in magnitude to the NEP . This concept is widely accepted in the scientific community (Aubinet et al., 2012), for forests as well as for other environments such as lakes.

The lateral transport of CO_2 had to be ruled for the sake of the calculations; we are of course fully aware of the lake being a 3D dynamic system, and we hope in the future to further develop the method we used so that it will work under more general 15 conditions. Besides, since this study focuses on the summer periods when the lake was stably stratified and there were no high winds or rains, the lateral transport is not expected to play a significant role here. A similar challenge is encountered in forest ecology studies as well, where the lateral transport in the air (advection) is also usually neglected.

The daytime and nighttime average values of the CO_2 flux were always positive, albeit having lower values during the day than during the night. This is not surprising: many lakes, especially at high latitudes, are supersaturated with respect to CO_2 ; 20 as a result, the CO_2 flux is from the lake to the atmosphere also during the day, when the aquatic primary producers are photosynthesising and absorbing CO_2 . The NEP values were in fact also always negative, confirming heterotrophy. This is in agreement with (Cole et al., 1994; Sobek et al., 2003).

Regarding oligotrophic lakes, it has been suggested that diurnal patterns in the epilimnion stratification and water convective motions (causing nighttime upwelling of CO_2) are important drivers of the diurnal variation of the surface water CO_2 concentration (Åberg et al., 2009). Lake Kuivajärvi though is mesotrophic (chl a 5-30 $\mu\text{g l}^{-1}$ during summer) and the primary production can be assumed to be the main driver of the CO_2 concentration, as observed also in some other lakes with high chl a (Hanson et al., 2003; Huotari et al., 2009). Also, we implemented strict selection criteria of the analyzed periods to minimize the effect of upwelling CO_2 : the thermistor data indicate that the winds, despite being weak, were strong enough to keep the top 1.5 m of the water column well mixed both day and night, without however disrupting the thermocline. Thus, no sign of hypolimnetic 30 upwelling was detected. Under these conditions, diurnal stratification patterns and convective motions had a minor impact on the mixed layer of our lake. It is also important to note that the photochemical production of CO_2 is generally negligible in humic lakes (Jonsson et al., 2001); its maximum contribution to the flux for a lake with similar characteristics as the one in our study lake was $< 4\%$ over the whole growing season, and was detectable only in the top 10 cm of the water column (Vähätalo



et al., 2000; Ojala et al., 2011).

In aquatic sciences, other models for describing the dependence of photosynthesis on irradiance are more commonly used than the Michaelis-Menten equation. The Michaelis-Menten equation was chosen in an effort of harmonizing productivity studies between aquatic and forest sciences, in order to study the carbon cycle consistently in the forest-lake continuum. However, we checked whether other models provided a better fit to the data. We used the equations by Smith (1936) and by Jassby and Platt (1976). Even though they agreed well with the data, they did not perform significantly better than the Michaelis-Menten equation: the R^2 and $RMSE$ values of the fits were very similar. Hence, we decided to proceed with our first choice. The Smith (1936) and Jassby and Platt (1976) model equations and fit statistics are reported in the supplementary information (Sect. S3 and Table S1).

In this study, we could not clearly link the environmental variables to the changes in the Michaelis-Menten model parameters, and more information on the algal communities living in the lake would have been required in order to extend the analysis. However, it is important for us to stress that the simplicity of this method lies in the fact that to estimate the parameters, which can then be used to calculate the productivity for each year, such information is not needed. Besides, an extensive application of the method would also allow for a comparison of the parameters between different lakes and different times, and the understanding of the relationship between the parameters and the environmental conditions or the specific phytoplankton communities would improve.

Our analysis was hindered by the problematic EC data set: due to inherent EC limitations and technical problems, the data set had many gaps and average flux values had to be used. The calculations could be improved with a better EC data set or chamber measurements performed regularly.

For further development, it would also be good to have measurements for F_u , the CO_2 flux from the deeper layer to the surface layer of the lake, in order not to limit the analysis to isothermal or stable stratification conditions. For example, water column turbulence measurements could be added to the CO_2 concentration and temperature measurements.

5 Conclusions

The high-frequency direct CO_2 concentration measurement method suggested in Hari et al. (2008) and tested only on 3 days of data under autumn turnover conditions was tested more extensively here, on a dataset of 40 days and under stable stratification conditions typical of summer for dark-water lakes. The method proved to be suitable for lake productivity studies under isothermal (Hari et al., 2008) or stable stratification conditions: its high temporal resolution allowed us to calculate the net ecosystem productivity at a temporal scale of minutes. A quantitative comparison between the calculated NEP and the modeled NEP was also carried out for the first time, and it showed a very good agreement between the two, further validating the method. From that, we were able to accurately parameterize the net productivity as a function of the ambient variables, estimating the productivity parameters typical of the communities in the lake.

Overall, we believe that the method proposed in Hari et al. (2008) and further tested, verified and developed here, with the explicit parameterization of the NEP from PAR and water T , represents a great improvement over the traditional approaches



(bottle method and ^{14}C technique). At the present stage it is still very system specific, and assumptions about lateral and vertical CO_2 exchange and photo-oxidation had to be made (negligible lateral exchange and photo-oxidation, no in-lake vertical exchange). Still, our study is an important step towards testing and developing the approach so that it becomes more general, also given the scarcity or even lack of high-frequency direct CO_2 measurements for productivity studies. We are looking for further contributions by the research community and we think the method should be widely adopted, in order first to gather more information about its usability under different conditions and then also to have a broader network of productivity studies on lakes. This is all the more true given that it is also easy to set up and relatively inexpensive. Its only requirement is a concomitant estimation of the CO_2 flux from the lake to the atmosphere. In our case the EC technique was used, which is expensive and can be laborious in the data processing phase. However, chamber measurements for example are an equally good option. Additionally, the method also relies on equations that are typically adopted in terrestrial ecology studies for the calculation of the NEP , where high-frequency measurements are more commonplace than in aquatic research. Extensively applying the method would reduce the gap in the CO_2 exchange measurements between aquatic and terrestrial ecology, which is beneficial in the framework of integrating research in different ecosystems, for which purpose a common language between different disciplines is needed. It would also help us achieve a better understanding of the biological processes behind the CO_2 exchange. This, in turn, would expand our knowledge on the carbon cycle in the water, which is still limited, and would lead to a better integration of aquatic ecosystems in the local and global carbon budgets.

Code and data availability. The data sets and the codes used in this paper can be obtained from the authors upon request.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This study was funded by the University of Helsinki and the Finnish Cultural Foundation - Häme fund (Hämeen rahasto - Suomen Kulttuurirahasto). Support also came from the Academy of Finland, through the Academy Professor projects (1284701 and 1282842), the CarLAC project (281196), ICOS-Finland (1281255) and the Finnish center of Excellence in Atmospheric sciences, and from the EU project GHG-LAKE (612642).



References

- Aubinet, M., Vesala, T., and Papale, D.: Eddy Covariance A Practical Guide to Measurement and Data Analysis, Springer, Dordrecht, the Netherlands, 2012.
- Åberg, J., Jansson, M., and Jonsson, A.: Importance of water temperature and thermal stratification dynamics for temporal variation of surface water CO₂ in a boreal lake, *J. Geophys. Res.*, 115, G02024, 2009.
- 5 Battin, T. J., Luyssaert, S., Kaplan, L. A., Aufdenkampe, A. K., Richter, A., and Tranvik, L. J.: The boundless carbon cycle, *Nature Geoscience*, 2, 598-600, 2009.
- Bender, M., Grande, K., Johnson, K., and others: A comparison of four methods for determining planktonic community production, *Limnol. Oceanogr.*, 32, 1085-1098, 1987.
- 10 Berggren, M., Lapierre, J. F., and del Giorgio, P. A.: Magnitude and regulation of bacterioplankton respiratory quotient across freshwater environmental gradients, *ISME J*, 6, 984-993, 2012.
- Caperon, J.: Population growth in micro-organisms limited by food supply, *Ecology*, 48, 715-722, 1967.
- Carignan, R.: Automated determination of carbon dioxide oxygen and nitrogen partial pressures in surface waters, *Limnol. Oceanogr.*, 43, 969-975, 1998.
- 15 Carignan, R., Planas, D., and Vis, C.: Planktonic production and respiration in oligotrophic Shield lakes, *Limnol. Oceanogr.*, 45, 189-199, 2000.
- Cole, J. J., and Caraco, N. F.: Atmospheric exchange of carbon dioxide in a low-wind oligotrophic lake measured by the addition of SF₆, *Limnol. Oceanogr.*, 43, 647-656, 1998.
- Cole, J. J., Caraco, N. F., Kling, G. W., and Kratz, T. K.: Carbon dioxide supersaturation in the surface waters of lakes, *Science-AAAS-Weekly Paper Edition*, 265, 1568-1569, 1994.
- 20 Cole, J. J., Prairie, Y. T., Caraco, N. F., and others: Plumbing the global carbon cycle: Integrating inland waters into the terrestrial carbon budget, *Ecosystems*, 10, 171-184, 2007.
- Davison, I. R.: Environmental effects on algal photosynthesis: temperature, *J. Phycol.*, 27, 2-8, 1991.
- Foken, T., Aubinet, M., and Leuning, R.: The eddy covariance method, pp. 1-19. In Aubinet, M., and others [eds.], *Eddy Covariance: A Practical Guide to Measurement and Data Analysis*, Springer, Dordrecht, the Netherlands, 2012.
- 25 Gaarder, T., and Gran, H. H.: Investigations of the production of plankton in the Oslo Fjord, *Rapp Proc Verb Conseil permanent international pour l'exploration de la mer*, 42, 1-48, 1927.
- Hanson, P., Bade, D., Carpenter, S., and Kratz, T.: Lake metabolism: Relationships with dissolved organic carbon and phosphorus, *Limnol. Oceanogr.*, 48, 1112-1119, 2003.
- 30 Hari, P., Pumpanen, J., Huotari, J., Kolari, P., Grace, J., Vesala, T., and Ojala, A.: High frequency measurements of productivity of planktonic algae using rugged nondispersive infrared carbon dioxide probes, *Limnol. Oceanogr.: Methods*, 6, 347-354, 2008.
- Heiskanen, J., Mammarella, I., Haapanala, S., Pumpanen, J., Vesala, T., MacIntyre, S., and Ojala, A.: Effects of cooling and internal wave motions on gas transfer coefficients in a boreal lake, *Tellus B*, 66, 22827, 2014.
- Heiskanen, J., Mammarella, I., Ojala, A., and others: Effects of water clarity on lake stratification and lake-atmosphere heat exchange, *J. Geophys. Res. Atmos.*, 120, 7412-7428, 2015.
- 35 Hoellein, T. J., Bruesewitz, D. A., and Richardson, D. C.: Revisiting Odum (1956): a synthesis of aquatic ecosystem metabolism, *Limnol. Oceanogr.*, 58, 2089-2100, 2013.



- Howarth, R. W., and Michaels, A. F.: The measurement of primary production in aquatic ecosystems, pp 72-85. In Sala, O. E., and others [eds.], *Methods in Ecosystems Science*, Springer, New York, 2000.
- Huotari, J., Ojala, A., Peltomaa, E., Pumpanen, J., Hari, P., and Vesala, T.: Temporal variations in surface water CO₂ concentrations in a boreal humic lake based on high-frequency measurements, *Bor. Env. Res.*, 14 suppl. A, 48-60, 2009.
- 5 Jassby, A. D., and Platt, T.: Mathematical formulation of the relationship between photosynthesis and light for phytoplankton, *Limnol. Oceanogr.*, 21, 540-547, 1976.
- Johnson, M. S., Billett, M. F., Dinsmore, K. J., Wallin, M., Dyson, K. E., and Jassal, R. S.: Direct and continuous measurement of dissolved carbon dioxide in freshwater aquatic systems - method and applications, *Ecohydrology*, 3, 68-78, 2010.
- Jonsson, A., Meili, M., Bergström, A.-K., and Jansson, M.: Whole-lake mineralization of allochthonous and autochthonous organic carbon
10 in a large humic lake (Örträsket, N. Sweden), *Limnol. Oceanogr.*, 46, 1691-1700, 2001.
- Karl, D. M., Laws, E. A., Morris, P., Williams, P. J. leB., and Emerson, S.: Metabolic balance of the open sea, *Nature*, 426, 5, 2003.
- Kiefer, D. A., and Mitchell, B. G.: A simple steady state description of phytoplankton growth based on absorption cross section and quantum efficiency, *Limnol. Oceanogr.*, 28, 770-776, 1983.
- Mallin, M. A., and Paerl, H. W.: Effects of variable irradiance on phytoplankton productivity in shallow estuaries, *Limnol. Oceanogr.*, 37,
15 54-62, 1992.
- Mammarella, I., Launiainen, S., Grönholm, T., Keronen, P., Pumpanen, J., Rannik, Ü., and Vesala, T.: Relative humidity effect on the high-frequency attenuation of water vapor flux measured by a closed-path eddy covariance system, *J. Atmos. Ocean. Technol.*, 26, 1856-1866, 2009.
- Mammarella, I., Nordbo, A., Rannik, Ü, and others: Carbon dioxide and energy fluxes over a small boreal lake in Southern Finland, *J. Geophys. Res. Biogeosci.*, 120, 1296-1314, 2015.
- 20 Miettinen, H., Pumpanen, J., Heiskanen, J. J., Aaltonen, H., Mammarella, I., Ojala, A., Levula, J., and Rantakari, M.: Towards a more comprehensive understanding of lacustrine greenhouse gas dynamics - two-year measurements of concentrations and fluxes of CO₂ CH₄ and N₂O in a typical boreal lake surrounded by managed forests, *Bor. Env. Res.*, 20, 75-89, 2015.
- Odum, H. T.: Primary production in flowing waters, *Limnol. Oceanogr.*, 1, 102-117, 1956.
- 25 Ojala, A., Lopez Bellido, J., Tulonen, T., Kankaala, P., and Huotari, J.: Carbon gas fluxes from a brown-water and a clear-water lake in the boreal zone during a summer with extreme rain events, *Limnol. Oceanogr.*, 56, 61-76, 2011.
- Pace, M. L., and Prairie, Y. T.: Respiration in lakes, pp 103-121. In del Giorgio, P. A., and Williams, P. J. leB. [eds.], *Respiration in Aquatic Ecosystems*, Oxford Univ. Press, New York, 2005.
- Peterson, B. J.: Aquatic primary productivity and the ¹⁴C-CO₂ method: a history of the productivity problem, *Ann. Rev. Ecol. Sys.*, 11,
30 359-385, 1980.
- Raven, J. A., and Geider, R. J.: Temperature and algal growth, *New Phytologist*, 11, 441-461, 1988.
- Raymond, P. A., and others: Global carbon dioxide emissions from inland waters, *Nature*, 503, 355-359, 2013.
- Read, J. S., Hamilton, D. P., Jones, I. D., Muraoka, K., Winslow, L. A., Kroiss, R., Wu, C. H., and Gaiser, E.: Derivation of lake mixing and stratification indices from high-resolution lake buoy data, *Environ. Model. Softw.*, 26, 1325-1336, 2011.
- 35 Reynolds, C. S.: *The Ecology of Freshwater Phytoplankton*, Cambridge Univ. Press, Cambridge, 1984.
- Reynolds, C. S.: *The Ecology of Phytoplankton*, Cambridge Univ. Press, Cambridge, 2006.
- Smith, E. L.: Photosynthesis in relation to light and carbon dioxide, *Proceedings of the National Academy of Sciences*, 22, 504-511, 1936.



- Sobek, S., Algsten, G., Bergström, A.-K., Jansson, M., and Tranvik, L. J.: The catchment and climate regulation of pCO₂ in boreal lakes, *Global Change Biol.*, 9, 630-641, 2003.
- Solomon, C. T., Bruesewitz, D. A., Richardson, D. C., and others: Ecosystem respiration: Drivers of daily variability and background respiration in lakes around the globe, *Limnol. Oceanogr.*, 58, 849-866, 2013.
- 5 Staehr, P. A., Bade, D., Van de Bogert, M. C., Koch, G. R., Williamson, C., Hanson, P., Cole, J. J., and Kratz, T.: Lake metabolism and the diel oxygen technique: state of the science, *Limnol. Oceanogr.: Methods*, 8, 628-644, 2010.
- Søndergaard, M.: A biography of Einar Steeman Nielsen: the man and his science, pp 1-15. In Williams, P. J. leB., and others [eds.], *Phytoplankton Productivity: Carbon Assimilation in Marine and Freshwater Ecosystems*. Blackwell Science, Oxford, 2000.
- Steemann Nielsen, E.: Measurement of the production of organic matter in the sea by means of carbon-14, *Nature*, 167, 684-685, 1951.
- 10 Tranvik, L. J., Downing, J. A., Cotner, J. B., and others: Lakes and reservoirs as regulators of carbon cycling and climate, *Limnol. Oceanogr.*, 54, 2298-2314, 2009.
- Vähätalo, A. V., Salkinoja-Salonen, M., Taalas, P., and Salonen, K.: Spectrum of the quantum yield for photochemical mineralization of dissolved organic carbon in a humic lake, *Limnol. Oceanogr.*, 45, 664-676, 2000.
- Vesala, T., Huotari, J., Rannik, Ü., Suni, T., Smolander, S., Sogachev, A., Launiainen, S., and Ojala, A.: Eddy covariance measurements of
15 carbon exchange and latent and sensible heat fluxes over a boreal lake for a full open-water period, *J. Geophys. Res.*, 111, D11101, 2006.




Figure 1. A schematic representation of the mass balance in the mixed layer of the lake. C_{CO_2} is the CO_2 concentration in the water, F_a is the CO_2 flux between the lake and the atmosphere (positive if from the lake to the atmosphere), F_u is the CO_2 flux from the deeper to the mixed layer of the lake (positive if upwards), t is time, z is depth and h_{mix} is the mixed layer depth.




Figure 2. A sample period of stable stratification in July 2010, representative of the studied periods. In the upper panel the solid line (sto) is the first term of Eq. (3), the CO_2 concentration change in time over the mixed layer, which is usually referred to as storage flux in forest ecology calculations; the dashed horizontal lines are the daytime and nighttime average CO_2 fluxes from the lake to the atmosphere (F_a). In the central panel, the solid line is the NEP and the dotted line is the zero rate. In the lower panel the solid line is the PAR (photon flux density measured in the PAR wavelength range) average value in the mixed layer.




Figure 3. The *NEP* versus *PAR* plots for each year; the fitted curve shown is calculated using the average water *T* of the studied periods of the year.




Figure 4. The *NEP* versus *PAR* plots for each year. Data points are color-classified according to water temperature classes, and the curves are calculated for the different temperatures; note that the curves are not individual fits, but are the result of the year's 3D fit, evaluated for the different temperatures. Water *T* is in °C.




Figure 5. Data and fitted *NEP* versus *PAR* and water *T* 3D-curves for each year.



Table 1. Parameters of the *NEP* vs *PAR* and water *T* model (Eq. (5)), with 95% confidence intervals, and fit statistics.

| Year | p_{\max} [$\mu\text{mol}(\text{CO}_2)\text{m}^{-2}\text{s}^{-1}$] | b [$\mu\text{mol}(\text{ph})\text{m}^{-2}\text{s}^{-1}$] | r_0 [$\mu\text{mol}(\text{CO}_2)\text{m}^{-2}\text{s}^{-1}$] | R^2 | $RMSE$ [$\mu\text{mol}(\text{CO}_2)\text{m}^{-2}\text{s}^{-1}$] |
|------|--|---|---|-------|--|
| 2010 | 1.05 ± 0.05 | 22 ± 5 | 0.228 ± 0.008 | 0.73 | 0.23 |
| 2011 | 1.47 ± 0.06 | 29 ± 6 | 0.399 ± 0.009 | 0.84 | 0.25 |
| 2013 | 0.63 ± 0.04 | 33 ± 10 | 0.290 ± 0.007 | 0.71 | 0.14 |
| 2014 | 1.55 ± 0.10 | 31 ± 11 | 0.482 ± 0.013 | 0.74 | 0.33 |



Table 2. The model parameters as in Table 1 (p_{\max} and r_0 in $\mu\text{mol}(\text{CO}_2)\text{m}^{-2}\text{s}^{-1}$, b in $\mu\text{mol}(\text{ph})\text{m}^{-2}\text{s}^{-1}$), average, minimum and maximum values of water T ($^{\circ}\text{C}$) and PAR in the mixed layer ($\mu\text{mol}(\text{ph})\text{m}^{-2}\text{s}^{-1}$) for the studied periods of each year.

| Year | p_{\max} | b | r_0 | T_{ave} | T_{min} | T_{max} | PAR_{ave} | PAR_{max} |
|------|------------|-----|-------|------------------|------------------|------------------|--------------------|--------------------|
| 2010 | 1.05 | 22 | 0.228 | 22.9 | 19.9 | 26.2 | 195 | 634 |
| 2011 | 1.47 | 29 | 0.399 | 22.7 | 20.7 | 25.3 | 197 | 708 |
| 2013 | 0.63 | 33 | 0.290 | 21.5 | 20.0 | 23.5 | 162 | 699 |
| 2014 | 1.55 | 31 | 0.482 | 25.6 | 23.2 | 28.3 | 227 | 741 |