

# N<sub>2</sub> fixation as a dominant new N source in the Western Tropical South Pacific Ocean (OUTPACE cruise)

Mathieu Caffin<sup>1</sup>, Thierry Moutin<sup>1</sup>, Rachel Ann Foster<sup>2</sup>, Pascale Bouruet-Aubertot<sup>3</sup>, Andrea Michelangelo Doglioli<sup>1</sup>, Hugo Berthelot<sup>1,4</sup>, Cécile Guieu<sup>5,6</sup>, Olivier Grosso<sup>1</sup>, Sandra Helias-Nunige<sup>1</sup>,  
5 Nathalie Leblond<sup>7</sup>, Audrey Gimenez<sup>1</sup>, Anne Alexandra Petrenko<sup>1</sup>, Alain de Verneil<sup>1</sup> and Sophie Bonnet<sup>8</sup>.

<sup>1</sup>Aix Marseille Université, CNRS, Université de Toulon, IRD, OSU Pythéas, Mediterranean Institute of Oceanography (MIO), UM 110, 13288, Marseille, France

<sup>2</sup>Stockholm University, Department of Ecology, Environment and Plant Sciences. Stockholm, Sweden

10 <sup>3</sup>Sorbonne Universités, UPMC Univ. Paris 06, LOCEAN, France

<sup>4</sup>Laboratoire des sciences de l'environnement marin, IUEM, Université de Brest-UMR 6539 CNRS/UBO/IRD/Ifremer, Plouzané, France

<sup>5</sup>Sorbonne Universités, UPMC Univ Paris 06, CNRS, Laboratoire d'Océanographie de Villefranche (LOV), 06230 Villefranche-sur-Mer, France

15 <sup>6</sup>Center for Prototype Climate Modeling, New York University Abu Dhabi, P.O. Box 129188, Abu Dhabi, United Arab Emirates

<sup>7</sup>Observatoire Océanologique de Villefranche, Laboratoire d'Océanographie de Villefranche, UMR 7093, Villefranche-sur-mer, France

20 <sup>8</sup>Aix Marseille Université, CNRS, Université de Toulon, IRD, OSU Pythéas, Mediterranean Institute of Oceanography (MIO), UM 110, 98848, Nouméa, New Caledonia

*Correspondence to:* Mathieu Caffin ([mathieu.caffin@mio.osupytheas.fr](mailto:mathieu.caffin@mio.osupytheas.fr))

## Abstract.

We performed nitrogen (N) budgets in the photic layer of three contrasting stations representing different trophic conditions in the western tropical South Pacific (WTSP) Ocean during austral summer conditions (Feb. Mar. 2015). Using a Lagrangian  
25 strategy, we sampled the same water mass for the entire duration of each long duration (5 days) station, allowing us to consider only vertical exchanges for the budgets. We quantified all major vertical N fluxes both entering (N<sub>2</sub> fixation, nitrate turbulent diffusion, atmospheric deposition) and leaving the photic layer (particulate N export). The three stations were characterized by a strong nitracline and contrasted deep chlorophyll maximum depths, which were lower in the oligotrophic Melanesian archipelago (MA, stations LD A and LD B) than in the ultra-oligotrophic waters of the South Pacific gyre (SPG,  
30 station LD C). N<sub>2</sub> fixation rates were extremely high at both LD A ( $593 \pm 51 \mu\text{mol N m}^{-2} \text{d}^{-1}$ ) and LD B ( $706 \pm 302 \mu\text{mol N m}^{-2} \text{d}^{-1}$ ), and the diazotroph community was dominated by *Trichodesmium*. N<sub>2</sub> fixation rates were lower ( $59 \pm 16 \mu\text{mol N m}^{-2} \text{d}^{-1}$ ) at LD C, and the diazotroph community was dominated by unicellular N<sub>2</sub>-fixing cyanobacteria (UCYN). At all stations, N<sub>2</sub> fixation was the major source of new N (> 90 %) before atmospheric deposition and upward nitrate fluxes induced by turbulence. N<sub>2</sub> fixation contributed circa 13-18 % of primary production in the MA region and 3 % in the SPG water and  
35 sustained nearly all new primary production at all stations. The e-ratio (e-ratio = particulate carbon export / primary production) was maximum at LD A (9.7 %) and was higher than the e-ratio in most studied oligotrophic regions (<5 %),

indicating a high efficiency of the WTSP to export carbon relative to primary production. The direct export of diazotrophs assessed by qPCR of the *nifH* gene in sediment traps represented up to 30.6 % of the PC export at LD A, while their contribution was 5 and < 0.1 % at LD B and LD C, respectively. At the three studied stations, the sum of all N input to the photic layer exceeded the N output through organic matter export. This disequilibrium leading to N accumulation in the upper layer appears as a characteristic of the WTSP during the summer season.

## 1 Introduction

Biological nitrogen fixation, the reduction of atmospheric dinitrogen ( $N_2$ ) to ammonia, is performed by a diverse group of prokaryotic organisms, commonly called diazotrophs. It provides the major external source of bio-available nitrogen (N) to the ocean, before riverine and atmospheric inputs (Deutsch et al., 2007; Gruber, 2008; Gruber and Sarmiento, 1997). In the oligotrophic ocean, N availability often limits phytoplankton growth (e.g. Moore et al., 2013) and  $N_2$  fixation sustains a significant part of new primary production (PP, i.e. the production unrelated to internal recycling of organic matter in the photic layer) such as in the North (Karl et al., 1997) and South Pacific Ocean (Moutin et al., 2008), the western Mediterranean Sea (Garcia et al., 2006), or the tropical North Atlantic (Capone et al., 2005). New N input by  $N_2$  fixation has thus been recognized as a significant process influencing global oceanic productivity, and can eventually fuel  $CO_2$  sequestration through the  $N_2$ -primed prokaryotic carbon (C) pump (Karl et al., 2003).

Low  $\delta^{15}N$  signatures of particles from sediment traps in the tropical North Pacific (Karl et al., 1997, 2012, Scharek et al., 1999a, 1999b) and Atlantic (Altabet, 1988; Bourbonnais et al., 2009; Knapp et al., 2005; Mahaffey et al., 2003) suggest that at least part of the recently fixed N is ultimately exported out of the photic zone. Knapp et al. (2008) and Bourbonnais et al. (2009) also observed a low  $\delta^{15}N$  of  $NO_3^-$  (relative to  $\delta^{18}O-NO_3^-$ ) in surface waters in the western and eastern subtropical Atlantic Ocean, supporting the role of  $N_2$ -fixers in these regions. Böttjer et al. (2017) revealed that  $N_2$  fixation supports 26-47 % of particulate N (PN) export over a 9-year time series (2005-2013) period at station ALOHA (North Pacific Subtropical Gyre). Export efficiency may depend on the diazotroph community composition present in surface waters. Blooms of diatom-diazotroph associations (DDAs) systematically observed in late summer at station ALOHA are thought to be directly responsible for the concomitant pulses of particulate export (Karl et al., 2012). High export associated with DDAs has also been observed in the Amazon River plume (Subramaniam et al., 2008), suggesting a high direct export efficiency associated with DDAs. *Trichodesmium* is one of the main contributors to global  $N_2$  fixation (Mahaffey et al., 2005), but is rarely recovered in sediment traps (Chen et al., 2003; Walsby, 1992), suggesting a low direct export efficiency. However, the  $N_2$  fixed by *Trichodesmium* is efficiently transferred to large, non-diazotrophic phytoplankton, such as diatoms (Berthelot et al., 2016; Bonnet et al., 2016a), which can be subsequently exported (Nelson et al., 1995), suggesting a potential indirect export pathway. A recent mesocosm study performed in New Caledonia unexpectedly revealed that the production sustained by unicellular diazotrophic cyanobacteria (hereafter referred to as UCYN) was much more efficient at promoting particle export than the production sustained by DDAs (Berthelot et al., 2015). However, the export efficiency of UCYN has been

poorly studied (White et al., 2012) in the open ocean, despite the fact that they contribute as much as *Trichodesmium* to  $N_2$  fixation rates in many parts of the ocean (Bonnet et al., 2009; Martínez-Pérez et al., 2016; Moisander et al., 2010; Montoya et al., 2004). More studies are thus needed to further investigate the ability of different diazotroph communities to fuel direct or indirect particle export in the oligotrophic ocean.

5 Studying the impact of  $N_2$  fixation on PN export in the ocean and the relative role of each diazotroph group in this process is technically challenging. It requires the measurement of all major N fluxes both entering the photic layer ( $N_2$  fixation, nitrate ( $NO_3^-$ ) eddy diffusion, atmospheric deposition) and leaving the photic layer (PN export) with an adequate time frame (i.e. linking production and export). In addition, the sampling has to been performed under contrasting situations, for example when either *Trichodesmium* or UCYN dominate the diazotroph community, hence allowing to assess the potential role of  
10 each diazotroph group. Most importantly, such N budgets must be performed in the same water mass to ensure that the particulate matter recovered in the sediment traps corresponds to the production that occurred just above in the photic layer. This is what we did during the OUTPACE (Oligotrophy to UItTra-oligotrophy PACific Experiment) cruise in the Western Tropical South Pacific (WTSP) in summer 2015, during which we used a Lagrangian strategy.

The WTSP Ocean has recently been identified as a hotspot of  $N_2$  fixation, including  $N_2$  fixation rates  $>500 \mu\text{mol N m}^{-2} \text{ d}^{-1}$   
15 (Bonnet et al., 2017). The region covered by the OUTPACE cruise is characterized by trophic and  $N_2$  fixation gradients (Moutin et al., 2017). The region covered by the OUTPACE cruise encompasses contrasting trophic regimes characterized by strong differences in top nitracline depths, from 46 to 141 m (Moutin et al., this issue), and representing a large part of the oligotrophic gradient at the scale of the world Ocean (Moutin and Prieur., 2012; their Fig. 9). The westward oligotrophic waters are characterized by high  $N_2$  fixation rates ( $631 \pm 286 \mu\text{mol N m}^{-2} \text{ d}^{-1}$ ) mainly associated with *Trichodesmium* (i.e.  
20 within the hot spot around Melanesian archipelago waters, hereafter named MA), and the eastward ultra-oligotrophic waters (in the eastern border of the South Pacific gyre, hereafter named SPG waters) are characterized by low  $N_2$  fixation rates ( $85 \pm 79 \mu\text{mol N m}^{-2} \text{ d}^{-1}$ ), mainly associated with UCYN (Bonnet et al., this issue; Stenegren et al., this issue). This west to east  $N_2$  fixation gradient has been mainly attributed to a decrease of iron availability in SPG waters as compared to MA waters (Bonnet et al., this issue; Guieu et al., in rev.). This region therefore provides ideal conditions to study the potential role of  
25  $N_2$  fixation on particulate export under contrasting situations.

In the present study, we focus on (i) the contribution of  $N_2$  fixation to new N inputs in the WTSP during the summer season; (ii) the coupling between  $N_2$  fixation and export; and (iii) the equilibrium versus disequilibrium between  $N_2$  fixation and particulate N export in the WSTP.

## 2 Material and methods

### 30 2.1 Station sampling strategy

The OUTPACE cruise was carried out during austral summer conditions (18 Feb.-3 Apr. 2015) along a west-east 5500 km transect from New-Caledonia ( $22^\circ\text{S} - 166^\circ\text{E}$ ) to French Polynesia ( $17^\circ30'\text{S} - 149^\circ30'\text{W}$ ). We performed a N budget at three

stations, hereafter named long duration (LD), stations that were chosen according to three criteria: (1) local minima of surface current intensity, (2) different trophic regimes, i.e. oligotrophic vs. ultra-oligotrophic, and (3) different diazotroph communities, i.e. *Trichodesmium* vs. UCYN.

To locate these three stations, we used a Lagrangian strategy developed during previous cruises such as LATEX (Doglioli et al., 2013; Petrenko et al., 2017) and KEOPS2 (d'Ovidio et al., 2015). Briefly, the regions of interest along the vessel route were first characterized at large scale through the analysis of satellite data. The altimetry-derived currents were processed by SPASSO (Software Package for an Adaptive Satellite-based Sampling for Ocean campaigns; <http://www.mio.univ-amu.fr/SPASSO/>) to derive Eulerian and Lagrangian diagnostics of ocean circulation : Okubo-Weiss parameter, particle retention time and advection, Lagrangian Coherent Structures (d'Ovidio et al., 2015), together with maps of the sea surface temperature and chlorophyll-a (Chl *a*) concentrations. The satellite data were processed on land in near real time and transmitted to the ship together with a daily bulletin proposing LD station positions (the complete series of 43 bulletins is available on the OUTPACE website at [https://outpace.mio.univ-amu.fr/OUT\\_Figures/Bulletins/](https://outpace.mio.univ-amu.fr/OUT_Figures/Bulletins/)). We also performed onboard quantitative Polymerase Chain Reaction (qPCR) analyses on the *nifH* gene to measure the abundance of six groups of diazotrophs (Stenengren et al., this issue). Thus, we located the stations in regions where either *Trichodesmium* or UCYN dominated the diazotroph community. Then, the exact locations of the three LD stations were determined on board in real time from a rapid survey using a Moving Vessel Profiler (MVP), equipped with conductivity-temperature-depth (CTD) and fluorimeter sensors, accompanied by the hull-mounted thermosalinograph and acoustic Doppler current profiler (de Verneil et al., this issue). Finally, Surface Velocity Program (SVP) drifters were deployed in order to study the relative dispersion at the surface (de Verneil et al., this issue) during the station occupation.

By using this strategy, LD A station (19°12.8'S - 164°41.3'E, 25 Feb. - 2 Mar.) was positioned in MA waters in the western part of the transect (Figure 1) offshore from New Caledonia. LD B station (18°14.4'S - 170°51.5'W, 15-20 Mar.) was positioned in MA waters near Niue Island and LD C station (18°25.2'S - 165°56.4'W, 23-28 Mar.) was positioned in the eastern part of the transect, in the SPG near the Cook Islands.

Each LD station was investigated for 5 days. The sequence of operations was the following: a drifting array equipped with three PPS5 sediment traps, current meters, oxygen sensors and high frequency temperature sensors (see <https://outpace.mio.univ-amu.fr/spip.php?article75> for details) was deployed at each station on the first day. Then, a series of CTD (SBE 911+ SeaBird) casts (0-500 m) were performed every 3-4 h near the actual position of the drifting array to study the high frequency evolution of temperature, salinity, Photosynthetically Available Radiation (PAR) and chlorophyll *a* fluorescence during the station occupation. Small-scale turbulence was characterized in the first 800 m from microstructure measurements using a Vertical Microstructure Profiler (VMP1000) that was typically deployed prior to or following each CTD cast (Bouruet-Aubertot et al, this issue). Nutrient concentration measurements (0-200 m) were performed every day on the midday CTD casts (hereafter named 'nut. CTD'). In addition to the 0-500 m casts every 3-4 h, production casts (0-150 m) (hereafter named 'prod. CTD') were performed three times at each LD station (on day 1, 3 and 5) to quantify N<sub>2</sub> fixation and primary production rates. Incubations with tracers (<sup>14</sup>C and <sup>15</sup>N<sub>2</sub>, see below) to quantify N<sub>2</sub> fixation and primary

production were performed on an in situ drifting production line deployed for 24 h from sunrise to sunrise. The drifting array with the traps was recovered at the end of each LD station. Aerosol sampling was performed throughout the cruise transect. The inputs of new N to the photic layer were induced by three different sources: atmospheric deposition at the air-sea interface, N<sub>2</sub> fixation as an interior source and NO<sub>3</sub><sup>-</sup> input by vertical diffusion. The N output was driven by PN sedimentation. The methods for the determination of each parameter are given below.

## 2. 2 Experimental procedures

### Physico-chemical parameters, nutrient concentrations and C:N ratios

In situ Chl *a* concentrations were derived from fluorescence measurements performed with a AquaTraka III (Chelsea Technologies Group Ltd) sensor mounted on the CTD. The chlorophyll fluorescence sensor was calibrated prior to the cruise and post-calibration was conducted using all HPLC measurements undertaken during the cruise. PAR was measured on each CTD profile. Phosphate (PO<sub>4</sub><sup>3-</sup>) and NO<sub>3</sub><sup>-</sup> concentrations were measured daily at 12 depths from the surface to 200 m on each nutrient CTD cast using standard colorimetric procedures (Aminot and K  rouel, 2007) on a AA3 AutoAnalyzer (Seal-Analytical). After filtration, one sample was directly analyzed on-board and the other poisoned with 50  l HgCl<sub>2</sub> (20 g L<sup>-1</sup>) and stored for analysis after the cruise in the laboratory. The quantification limits were 0.05   mol L<sup>-1</sup> for PO<sub>4</sub><sup>3-</sup> and NO<sub>3</sub><sup>-</sup>.

### 15 Primary production and associated N uptake

PP was measured in triplicate using the <sup>14</sup>C tracer method (Moutin and Raimbault, 2002). Samples were incubated in 320 mL polycarbonate bottles on the in situ drifting production line (Aquamout: <https://outpace.mio.univ-amu.fr/spip.php?article75>) for 24-h from dusk-to-dusk at 9 depths (75 %, 54 %, 36 %, 19 %, 10 %, 3 %, 1 %, 0.3 % and 0.1 % of surface irradiance levels), corresponding to the sub-surface (5 m) down to 105 m, 80 m and 180 m for stations LD A, LD B and LD C, respectively. A N-derived PP (N-PP) was obtained at each depth by dividing PP by the classical C:N Redfield ratio (6.625). Integrated N-PP (iN-PP) over the studied layer (surface to 125m, 100 m and 200 m for LD A, LD B and LD C, respectively) was calculated by using the trapezoidal method, assuming that surface N-PP was identical to N-PP measured in subsurface (5 m) and considering that N-PP 20 m below the deepest sampled depth was zero (JGOFS, 1988).

### Atmospheric deposition

25 N atmospheric deposition (NO<sub>3</sub><sup>-</sup> and NO<sub>2</sub><sup>-</sup> (nitrite), hereafter called NO<sub>x</sub>) was quantified along the transect after dissolution of aerosols collected continuously during the transect, as described in Guieu et al. (in rev.). Briefly, the sampling device, designed to avoid ship contamination was installed at the look-out post in the front of the ship, collected aerosols at ~20 L min<sup>-1</sup> on onto polycarbonate, 47-mm diameter, 0.45   m porosity (previously acid-cleaned with a 2% solution of HCl (Merck, Ultrapur, Germany) and thoroughly rinsed with ultra-pure water and dried under a laminar flow bench and stored in acid-cleaned Petri dishes). Dissolution experiments to determine NO<sub>x</sub> released in surface seawater after deposition were

performed on board using acid-cleaned Sartorius filtration units (volume 0.250 L) and filtered surface (5m) seawater. Each sample was subjected to two contact times: the first contact was at one minute, and the second contact was at 24 hours. NO<sub>x</sub> was analysed using a 1-m long Liquid Waveguide Capillary Cells (LWCC) made of quartz capillary tubing, following the protocol described in Louis et al., 2015. An extrapolated NO<sub>x</sub> release from dry deposition was estimated on the basis of a deposition velocity of submicronic particles (0.4 m s<sup>-1</sup>; Vong et al., 2010).

### Nitrogen fixation rates

N<sub>2</sub> fixation rates were measured using the <sup>15</sup>N<sub>2</sub> tracer method (Montoya et al. (1996), modified (see below)), on day 1, 3 and 5 at each LD station (hereafter named in situ\_1, in situ\_2 and in situ\_3, respectively). Seawater was collected from the Niskin bottles in duplicate 4.5 L polycarbonate bottles at 9 depths (same depths as for PP). 5 mL of <sup>15</sup>N<sub>2</sub> gas (99 atom% <sup>15</sup>N, Eurisotop) were injected in each bottle through the septum cap using a gas-tight syringe. The purity of the <sup>15</sup>N<sub>2</sub> Cambridge isotopes stocks was previously checked by Dabundo et al. (2014) and more recently by Benavides et al. (2015) and Bonnet et al. (2016a). They were found to be lower than 2 x 10<sup>-8</sup> mol:mol of <sup>15</sup>N<sub>2</sub>, leading to a potential N<sub>2</sub> fixation rates overestimation of <1 %. All bottles were shaken 20 times to facilitate the <sup>15</sup>N<sub>2</sub> dissolution and incubated in situ on the production line at the same depth of sampling for 24 h from dusk to dusk (hereafter called ‘in situ incubation method’). It has been previously shown that the bubble method potentially underestimated N<sub>2</sub> fixation rates (Großkopf et al., 2012; Mohr et al., 2010a) compared to methods consisting in adding the <sup>15</sup>N<sub>2</sub> as dissolved in a subset of seawater previously N<sub>2</sub> degassed (Mohr et al., 2010a). This underestimation is due to incomplete equilibration of the <sup>15</sup>N<sub>2</sub> gas with surrounding seawater. However, other studies did not find any significant difference between the two methods (Bonnet et al., 2016b; Shiozaki et al., 2015). In the present study, we intentionally decided to use the bubble method due to the high risk of both organic matter and trace metal contamination during the <sup>15</sup>N<sub>2</sub>-enriched seawater preparation (Klawonn et al., 2015), that has been seen to enhance N<sub>2</sub> fixation in this area (Benavides et al., 2017; Moisaner et al., 2010). However, to minimize possible rate underestimations due to incomplete equilibration of the <sup>15</sup>N<sub>2</sub> gas with surrounding seawater, the final <sup>15</sup>N-enrichment of the N<sub>2</sub> pool was quantified in the incubation bottles on each profile in triplicate at 5 m and at the deep chlorophyll maximum (DCM). After incubation, 12 mL of each 4.5 L bottle were subsampled in Exetainers, fixed with HgCl<sub>2</sub> (final concentration 20 µg mL<sup>-1</sup>), and stored upside down at 4°C in the dark until analyzed onshore within 6 months after the cruise, according to Kana et al. (1994), using a Membrane Inlet Mass Spectrometer (MIMS).

For each LD station, in parallel with the last N<sub>2</sub> fixation profile (the in situ\_3 profile), we performed a replicate N<sub>2</sub> fixation profile in which bottles were incubated in on-deck incubators (see details on <https://outpace.mio.univ-amu.fr/spip.php?article135>) equipped with circulating seawater at the specified irradiances using blue screening (hereafter called ‘deck incubation method’). For these profiles, samples were collected in triplicate at 6 of the 9 depths reported above (75 %, 54 %, 36 %, 10 %, 1 % and 0.1 % surface irradiance level) in 2.3 L polycarbonate bottles amended with 2.5 mL of <sup>15</sup>N<sub>2</sub> gas (99 atom% <sup>15</sup>N, Eurisotop) for 24 h.

In both cases, incubations were stopped by gentle filtration (<0.2 bar) of the samples onto pre-combusted (450 °C, 4 h) Whatman GF/F filters (25 mm diameter, 0.7 µm nominal porosity). Filters were stored in pre-combusted glass tubes at -20 °C during the cruise, then dried at 60 °C for 24 h before analysis onshore. <sup>15</sup>N-enrichments of PN collected on filters were determined using an Elemental Analyzer coupled to an Isotope Ratio Mass Spectrometer (EA-IRMS, Integra2 Sercon Ltd).

5 The accuracy of the EA-IRMS system was systematically checked using International Atomic Energy Agency (IAEA) reference materials, AIEA-N-1 and IAEA-310A. In addition, the <sup>15</sup>N-enrichment of the ambient (unlabeled) PN was measured in one replicate at each station at the DCM and the subsurface and was used as the “initial” <sup>15</sup>N-enrichment, as termed in Montoya et al. (1996). The minimum quantifiable rate calculated using standard propagation of errors via the observed variability between replicate samples measured according to Gradoville et al. (2017) was 0.035 nmol N L<sup>-1</sup> d<sup>-1</sup>.

10 Integrated N<sub>2</sub> fixation rates over the studied layer were calculated by using the same method as for the iN-PP.

### Nitrate turbulent diffusion

NO<sub>3</sub><sup>-</sup> inputs from deep waters by turbulent mixing were estimated at the top of the nitracline. The top of the nitracline was found to fall along an isopycnal surface ( $\rho=\rho_{\text{NO}_3}$ ), the density of which was determined at each station. The NO<sub>3</sub><sup>-</sup> turbulent diffusive flux along the isopycnal surface  $\rho=\rho_{\text{NO}_3}$  was defined as:

$$15 \quad \text{Flux}_{\rho_{\text{NO}_3}}(t) = -K_z(z_{\rho_{\text{NO}_3}}, t) \times \frac{d[\text{NO}_3]}{d\rho}(\rho_{\text{NO}_3}, t) \times \frac{d\rho}{dz}(z_{\rho_{\text{NO}_3}}, t), \quad (1)$$

with  $K_z$ , the turbulent diffusion coefficient along the isopycnal  $\rho=\rho_{\text{NO}_3}$  inferred from VMP1000 measurements performed every 3-4 h during the LD station occupation, as described in Bouruet-Aubertot et al. (this issue) ;  $\frac{d[\text{NO}_3]}{d\rho}$ , the constant isopycnal slope of the nitracline, calculated for each station, and  $\frac{d\rho}{dz}$ , the vertical density gradient measured by the VMP1000 at the  $\rho_{\text{NO}_3}$  isopycnal depth  $z_{\rho_{\text{NO}_3}}$ . The time series of the NO<sub>3</sub><sup>-</sup> turbulent diffusive flux was calculated using an hourly temporal interpolation of  $K_z$  over the entire duration of each LD station. In addition, daily averages and 5-day averages were computed.

20

### Particulate matter export

Particulate matter export was quantified with three PPS5 sediment traps (1 m<sup>2</sup> surface collection, Technicap, France) deployed for 5 days at 150, 330 and 520 m at each LD station (Fig. 1). We decided to use the same configuration at the 3 LD

25 stations. The first trap was deployed at 150 m as it is always below the base of the photic layer, and therefore below the productivity layer. 520 m was used because most of the diel zooplankton vertical migrations stopped above this depth, and 330 m was chosen as an intermediary depth. Particle export was recovered in polyethylene flasks screwed on a rotary disk which allowed the flask to be automatically changed every 24-h to obtain a daily material recovery rate. The flasks were previously filled with a buffered solution (Sodium borate) of formaldehyde (final conc. 2 % and pH=8) and were stored at 4

30 °C after collection until analysis to prevent degradation of the collected material. The flask corresponding to the fifth day of

sampling on the rotary disk was not filled with formaldehyde in order to collect ‘fresh particulate matter’ for further diazotroph quantification, as described below. Thus, this last flask was not used in the particulate export computations reported in Table 1. Onshore, swimmers were handpicked from each sample, quantified and genera-identified. Exported particulate matter and swimmers were both weighed and analyzed separately on EA-IRMS (Integra2, Sercon Ltd) to quantify exported PC and PN.

### Diazotroph abundance in the traps

Triplicate aliquots of 2 to 4 mL from the flask dedicated to diazotroph quantification were filtered onto 0.2 µm Supor filters, flash frozen in liquid nitrogen and stored at -80 °C until analysis. Nucleic acids were extracted from the filters as described in Moisanter et al., (2008) with a 30 seconds reduction in the agitation step in a Fast Prep cell disruptor (Thermo, Model FP120; Qbiogene, Inc. Cedex, France) and an elution volume of 70 µl. Diazotroph abundance for *Trichodesmium* spp., UCYN-B (*Crocospaera watsonii*), UCYN-A1 (*Candidatus Atelocyanobacterium thalassa*), het-1 (*Richelia intracellularis* in symbiosis with *Rhizosolenia*), and het-2 (*Richelia intracellularis* in symbiosis with *Hemiaulus*) were quantified by qPCR analyses on the *nifH* gene using previously described oligonucleotides and assays (Foster et al. 2007; Church et al. 2005). The qPCR was conducted in a StepOnePlus system (applied Biosystems, Life Technologies, Stockholm Sweden) with the following parameters: 50 °C for 2 min, 95 °C for 10 min, and 45 cycles of 95 °C for 15 s followed by 60 °C for 1 min. Gene copy numbers were calculated from the mean cycle threshold (Ct) value of three replicates and the standard curve for the appropriate primer and probe set. For each primer and probe set, duplicate standard curves were made from 10-fold dilution series ranging from 10<sup>8</sup> to 1 gene copies per reaction. The standard curves were made from linearized plasmids of the target *nifH* or from synthesized gBlocks gene fragments (IDT technologies, Cralville, Iowa USA). Regression analyses of the results (number of cycles=Ct) of the standard curves were analyzed in Excel. Two µl of 5 KDa filtered nuclease free water was used for the no template controls (NTCs). No *nifH* copies were detected for any target in the NTC. In some samples, only 1 or 2 of the 3 replicates produced an amplification signal; these were noted as detectable but not quantifiable (< QL). A 4<sup>th</sup> replicate was used to estimate the reaction efficiency for the *Trichodesmium* and UCYN-B targets, as previously described in Short et al., (2004). Seven and two samples were below 95% in reaction efficiency for *Trichodesmium* and UCYN-B, respectively. The samples with qPCR reaction efficiency below 95% were excluded. The detection limit for the qPCR assays is 1-10 copies.

To determine directly the biovolume and C content of diazotrophs, cell sizes of *Trichodesmium* and UCYN-B were determined in samples from the photic layer of each LD station. Briefly, 2.3 L of surface (5 m) seawater were gently filtered (< 0.2 bar) onto 2 µm nominal porosity (25 mm diameter) polycarbonate filters, fixed with paraformaldehyde (final conc. 2 %) and stored at -80 °C. Cell length and width measurements were performed on 25 to 50 cells per station at x400 magnification with a Zeiss Axio Observer epifluorescence microscope. The biovolumes (BV) of *Trichodesmium* and UCYN were estimated using the equation for a cylinder and a sphere, respectively (Sun and Liu, 2003). The cellular C contents were determined by using the relation between BV and C content according to Verity et al. (1992). Given that our work was



performed in the field on wild populations, we preferred to use the biovolume estimate for C content rather than previously measured values based on published culture data. The C contents estimated here are within the range of those previously reported (Dekaezemacker and Bonnet, 2011; Dron et al., 2013; Hynes et al., 2012; Knapp et al., 2012; Luo et al., 2012). As DDAs were not easily identified on the filters, the C content was indirectly estimated. We used a C content of 1400 pg C cell<sup>-1</sup> for *Rhizosolenia* spp. (determined for the OUTPACE cruise, pers. comm. K. Leblanc) and assumed 6 cells per trichome of *Richelia intracellularis* (Foster and Zehr, 2006; Villareal, 1989), and 1 *Richelia* per diatom *Rhizosolenia* (both *Rhizosolenia* spp. and *R. intracellularis* are rarely reported as asymbiotic).

## Statistical analyses

Spearman correlation coefficients were used to examine the relationships between DCM and nitracline depths during the station occupation ( $\alpha = 0.05$ ).

A non-parametric Mann–Whitney test ( $\alpha = 0.05$ ) was used to compare N<sub>2</sub> fixation rates obtained using in situ and on-deck incubation methods.

## 3 Results

### 3.1 Hydrological background

NO<sub>3</sub><sup>-</sup> concentrations in the photic layer were below the quantification limit (0.05 µmol L<sup>-1</sup>) during the station occupation at the three LD stations. They became quantifiable below 70 m, 100 m and 120 m depth at LD A, B and C, respectively (Fig. 2) and at depths corresponding to a density anomaly of 23.59, 24.34 and 24.66 kg m<sup>-3</sup>, respectively. The latter values correspond to the top of the nitracline, the isopycnal  $\rho = \rho_{\text{NO}_3}$  (Fig. 3; Table 2). The corresponding  $\frac{d[\text{NO}_3]}{d\rho}$  were 3573, 5949 and 8888 µmol kg<sup>-1</sup> for stations LD A, LD B and LD C, respectively.

Averaged PO<sub>4</sub><sup>3-</sup> concentrations were close to or below the quantification limit (0.05 µmol L<sup>-1</sup>) from the surface to 20 m at LD A and LD B, and then increased with depth to reach 0.46 and 0.36 µmol L<sup>-1</sup> at 200 m at LD A and LD B, respectively (Fig. 2). At LD C, PO<sub>4</sub><sup>3-</sup> concentrations were always above the quantification limit and varied from 0.11 to 0.17 µmol L<sup>-1</sup> in the 0-120 m layer. Below 120 m, PO<sub>4</sub><sup>3-</sup> concentrations increased with depth to reach 0.43 µmol L<sup>-1</sup> at 200 m.

At LD A, the DCM and the  $\rho_{\text{NO}_3}$  depths were located between 60 and 100 m (Fig. 3). At LD B, the  $\rho_{\text{NO}_3}$  was located between 100 and 120 m, while the depth of the DCM increased from 25 to 70 m during the five days that the station was occupied. At LD C, the DCM and the  $\rho_{\text{NO}_3}$  were below the bottom of the photic layer (110-120 m), varying concurrently between 115 and 155 m. The DCM and  $\rho_{\text{NO}_3}$  depths were significantly correlated ( $p < 0.05$ ) at LD A and LD C and not correlated ( $p > 0.05$ ) at LD B. The depths of the photic layers, corresponding to 1 % of the surface PAR at midday, were 80-90 m at LD A, 60-70 m at LD B and 110-120 m at LD C.

### 3.2 Atmospheric deposition

Nitrate dissolution from aerosols occurred rapidly, releasing in seawater on average  $1.8 \text{ nmol m}^{-3}$  ( $26 \text{ ng m}^{-3}$ ) dissolved inorganic nitrogen. Nitrate appeared to be as nitrate aerosol since no correlation was observed between nitrate and Fe, Si, Na, Cl (K. Desboeufs, pers. Com., 2017), precluding a mixing with ash or seasalt. Extrapolated dry deposition flux (Table 2) was  
5 on average  $630 \pm 329 \text{ nmol m}^{-2} \text{ d}^{-1}$  ( $3.22 \pm 1.7 \text{ mg m}^{-2} \text{ yr}^{-1}$ ).

### 3.3 N<sub>2</sub> fixation rates

N<sub>2</sub> fixation rates measured using the in situ incubation method ranging from <QL -  $19.3 \text{ nmol N L}^{-1} \text{ d}^{-1}$  at LD A,  $0.1 - 45.0 \text{ nmol N L}^{-1} \text{ d}^{-1}$  at LD B and <QL -  $2.6 \text{ nmol N L}^{-1} \text{ d}^{-1}$  at LD C (Fig. 2). At LD A and LD B, maximum rates were measured near the surface (5 m) where they reached  $19.3$  and  $45.0 \text{ nmol N L}^{-1} \text{ d}^{-1}$ , and decreased with depth down to  $0.5 \text{ nmol N L}^{-1} \text{ d}^{-1}$  at  
10 70 and 55 m, respectively at LD A and LD B. At LD C, N<sub>2</sub> fixation rates were 2 to 20 times lower than at LD B and LD A, with a maximum of  $2.6 \text{ nmol N L}^{-1} \text{ d}^{-1}$  located around 40 m. Close to the surface (5 m), rates were below the quantification limit. At LD A and LD C, the three profiles measured on days 1, 3 and 5 at each station were similar to each other, while at LD B rates measured in the 0-40 m layer were different over the three sampling dates (Fig. 2), with rates decreasing over time.

15 N<sub>2</sub> fixation rates measured using the deck incubation method were not statistically different (Mann-Whitney paired test,  $p < 0.05$ ) from those measured using the in situ mooring line method (Fig. 2). They ranged  $0.1$  to  $21.0 \text{ nmol N L}^{-1} \text{ d}^{-1}$  at LD A,  $0.1$  to  $30.3 \text{ nmol N L}^{-1} \text{ d}^{-1}$  at LD B, and were below  $1.2 \text{ nmol N L}^{-1} \text{ d}^{-1}$  at LD C. Overall, the profiles were similar between the two methods, except the maximum at 40 m at LDC, which was not sampled with the on-deck incubation method. In addition, N<sub>2</sub> fixation rates from days 1-2-3 were not statistically different each other (Mann-Whitney paired test,  $p < 0.05$ ).

20 Integrated N<sub>2</sub> fixation rates were  $593 \pm 51$ ,  $706 \pm 302$  and  $59 \pm 16 \text{ } \mu\text{mol N m}^{-2} \text{ d}^{-1}$  at LD A, LD B and LD C, respectively using data from the in situ incubation method (Fig. 2; Table 2) and  $628 \pm 156$ ,  $942 \pm 253$  and  $56 \pm 31 \text{ } \mu\text{mol N m}^{-2} \text{ d}^{-1}$  at LD A, LD B, and LD C, respectively, using data from the deck incubation method (Fig. 2). 80 % of integrated N<sub>2</sub> fixation rate was reached at 36 m at LD A, 82 % was reached at 27 m at LD B and 78 % was reached at 60 m at LD C.

### 3.4 Vertical turbulent diffusive fluxes of nitrate

25 The averaged  $\text{NO}_3^-$  input through vertical turbulent diffusion showed strong time variability, with a typical standard deviation of the same order as the mean value (Table 3) and a strong contrast between the western station LD A and the two other stations, with mean values equal to  $24.4 \pm 24.4 \text{ } \mu\text{mol N m}^{-2} \text{ d}^{-1}$  at LD A and  $6.7 \pm 5.3$  and  $4.8 \pm 2.2 \text{ } \mu\text{mol N m}^{-2} \text{ d}^{-1}$  at LD B and LD C, respectively (Fig. 4). At LD A, a  $\text{NO}_3^-$  peak input of  $50 \text{ } \mu\text{mol N m}^{-2} \text{ d}^{-1}$  was observed on day 1 (26/02), while during days 2 and 3 (27/02 and 28/02) the daily average input was lower than the average value for the station,  
30 between  $5-10 \text{ } \mu\text{mol N m}^{-2} \text{ d}^{-1}$ , without any peak input. At the end of LD A (days 4 and 5 - 01/03 and 02/03), the strongest  $\text{NO}_3^-$  input variability was observed with instantaneous peaks reaching 46 to  $89 \text{ } \mu\text{mol N m}^{-2} \text{ d}^{-1}$ . When averaged per day,

daily input was minimum on day 2 (27/02) with  $5 \mu\text{mol N m}^{-2} \text{d}^{-1}$  and maximum on day 5 (02/03) with  $65 \mu\text{mol N m}^{-2} \text{d}^{-1}$  (red lines in Fig. 4a). At LD B, the mean daily  $\text{NO}_3^-$  input varied within a factor of  $\sim 5$ , from  $2 \mu\text{mol N m}^{-2} \text{d}^{-1}$  on day 5 (20/03) to  $11 \mu\text{mol N m}^{-2} \text{d}^{-1}$  on day 2 (17/03). The highest daily averages obtained on days 2 and 3 were explained by the occurrence of  $\text{NO}_3^-$  input peaks. At LD C,  $\text{NO}_3^-$  input heterogeneously varied between 2 and  $10 \mu\text{mol N m}^{-2} \text{d}^{-1}$  with minimum daily average on day 4 (27/03;  $3 \mu\text{mol N m}^{-2} \text{d}^{-1}$ ) and maximum daily average on day 1 (24/03;  $8 \mu\text{mol N m}^{-2} \text{d}^{-1}$ ). Similarly as for the other stations,  $\text{NO}_3^-$  input peaks were observed during the days of higher mean daily  $\text{NO}_3^-$  input. This time variability in  $\text{NO}_3^-$  input was strongly influenced by the vertical turbulent diffusion coefficient. The Kz time series showed strong variability (Table 3), with peak values occurring intermittently during periods of enhanced turbulence, thus leading to peaks in  $\text{NO}_3^-$  turbulent diffusive flux.

### 10 3.5 Particulate matter export

Mass fluxes recovered in the sediment traps at the three stations over the three sampling depths (150, 330 and 520 m) ranged from 13.6 to  $87.2 \text{ mg of Dry Weight (DW) m}^{-2} \text{d}^{-1}$  (Table 1). At LD A and LD C, fluxes decreased with depth, which was not observed at LD B. Maximum mass fluxes were measured at LD A, with  $87.2 \text{ mg DW m}^{-2} \text{d}^{-1}$  at 150 m,  $23.9 \text{ mg DW m}^{-2} \text{d}^{-1}$  at 330 m and  $22.3 \text{ mg DW m}^{-2} \text{d}^{-1}$  at 520 m. LD B presented the lowest export rate at 150 m ( $14.1 \text{ mg DW m}^{-2} \text{d}^{-1}$ ) over the three stations. At LD C,  $19.6 \text{ mg DW m}^{-2} \text{d}^{-1}$  of particulate matter was exported at 150 m and the lowest export rate was recorded at 330 m.

Particulate C (PC) and PN recovered in the sediment traps followed the same patterns as the mass fluxes (Table 1), with a maximum export rate at 150 m at LD A, a minimum export rate at LD B and LD C. However, as PN and PC were not always in the same proportion in the exported matter, it induced variations of C:N ratios at the three stations, with averaged C:N ratios of 8.2 at LD A, 9.1 at LD B and 6.2 at LD C.

The mass of swimmers (zooplankton) recovered in the traps ranged from 10.5 to  $376.1 \text{ mg DW m}^{-2} \text{d}^{-1}$  (Table 4) and accounted from 36 to 94 % of total DW. The maximum was found at 330 m at LD A, and 150 m at LD B and LD C. As for particulate matter, zooplankton C (Zoo-C) and N (Zoo-N) mass measured at each depth of each station followed the same pattern as the mass of swimmers recovered. Zoo-C ranged from 4.9 to  $129.2 \text{ mg C m}^{-2} \text{d}^{-1}$  and Zoo-N ranged from 1.1 to  $19.5 \text{ mg N m}^{-2} \text{d}^{-1}$ .

### 3.6 Direct export of diazotrophs

*Trichodesmium* abundance measured in the sediment traps at the three stations ranged from below quantification limit ( $< \text{QL}$ ) to  $2.67 \times 10^4 \text{ nifH gene copies mL}^{-1}$  of sediment material ( $< \text{QL}$  at LD C 330 m and not available at LD A 150 m), and represented less than 0.1 % of *Trichodesmium* abundance integrated over the water column at the three stations based on data of Stenegren et al., (This issue). UCYN-B abundance measured in the traps ranged for  $< \text{QL}$  to  $4.27 \times 10^3 \text{ nifH gene copies mL}^{-1}$ . It accounted from 0.1 to 10.5 % of UCYN-B abundance integrated over the water column at LD A, and  $< 0.5\%$  at LD B and LD C. DDAs abundance, restricted to het-1 (*Richelia* associated with *Rhizosolenia* diatoms), ranged from  $< \text{QL}$  to 1.99

$\times 10^4$  *nifH* gene copies  $\text{mL}^{-1}$  (< QL at LD A 150 and 520 m, LD B 150 m and LD C 330 m) and accounted up to 72.6 %, 2.9 % and 0.1 % of DDAs abundance integrated over the water column at LD A, LD B and LD C, respectively. While *het-2* (*Richelia* associated with *Hemiaulus* diatoms) were observed in the water column (Stenegren et al., This issue), they were only detected in one sediment trap sample (LD B, 330 m) and < QL in 330 m from LD A and 500 m from LD B. When converted to C, diazotrophs represented between 5.4 % and 30.6 % (Fig. 5) of the total PC measured in the traps (Table 1) at LDA, from < 0.1 to 5.0 % at LD B and < 0.1 % at LD C. *Trichodesmium* and *het-1* were the major contributors to diazotroph export at LD A and LD B (note that *Trichodesmium* data were not available for LD A (150 m)) and UCYN-B and *Het-1* were the major contributors at LD C.

## 4 Discussion

### 10 Towards a daily N-budget

The analysis of hydrographic tracers and velocity structures present during our study at the three stations reveals that horizontal variability due to advection was important at spatial scales larger than the ones sampled during each station (de Verneil et al., this issue). Thus, we consider that we sampled the same water mass during each station and only vertical exchanges controlled input and output of N in the upper water column, which allow us to perform a daily N-budget at the three stations, as summarized in Table 2.

### Contribution of $\text{N}_2$ fixation to new N input in the WTSP

The daily N-budget (Table 2) indicates that  $\text{N}_2$  fixation was the major external source of N to the WTSP, regardless of the degree of oligotrophy, and represents more than 90 % of new N to the surface ocean at every station. This contribution is higher than in previous studies performed in other oligotrophic regions impacted by  $\text{N}_2$  fixation (Table 5) such as the tropical North Atlantic (50 %, Capone et al., 2005) and Pacific (30-50 %, Dore et al., 2002; Karl et al., 2002), and higher than the average contribution at global scale (Gruber, 2008). This previously unreported high contribution of  $\text{N}_2$  fixation may have several origins.

- Extrapolated  $\text{NO}_x$  deposition from the atmosphere during OUTPACE (range:  $0.34 - 1.05 \mu\text{mol m}^{-2} \text{d}^{-1}$ ) were one order of magnitude lower than predicted with major uncertainties by global models that include wet and gas deposition for that region (Kanakidou et al., 2012). Our flux could be an underestimation as it represents only dry deposition and as gas and organic forms were not measured. At the global scale and depending on the location, organic nitrogen could represent up to 90 % of N atmospheric deposition (Kanakidou et al., 2012), and  $\text{NH}_4^+$  could account for ~40% (Dentener et al. 2006). Even if we double our estimated deposition flux, atmospheric deposition still remained low (< 1.5 %) and consequently represented a minor contribution of the new N input (Table 2). This negligible contribution of atmospheric input to the overall N budget (less than 1.5 %) therefore implies an important contribution of other terms, such as  $\text{N}_2$  fixation.

- Then,  $\text{NO}_3^-$  input by vertical turbulent diffusion appeared as the second source (1 to 8 %) of new N at the three stations. This contribution was lower than in previous studies in other oligotrophic regions (Table 5), where  $\text{NO}_3^-$  input by vertical turbulent diffusion contributes ~ 18 % of new N in the Indian South Subtropical Gyre (Fernández-Castro et al., 2015), and ~ 50 % in the Tropical North Atlantic (Capone et al., 2005). In most studies (Fernández-Castro et al., 2015; Moutin and Prieur, 2012; Painter et al., 2013), an average Kz value is used (i.e. averaged over the cruise, over a station or over depth) to determine  $\text{NO}_3^-$  input by turbulence in the photic layer. In this study we performed high frequency direct measurements of Kz and highlighted the importance of turbulent event pulses on diffusive  $\text{NO}_3^-$  input. Using a constant Kz of  $10^{-5} \text{ m}^2 \text{ s}^{-1}$  at the 3 stations decreases the  $\text{NO}_3^-$  input down to  $22.9 \mu\text{mol N m}^{-2} \text{ d}^{-1}$  at LD A and increases  $\text{NO}_3^-$  input up to 19.9 and  $25.5 \mu\text{mol N m}^{-2} \text{ d}^{-1}$  at LD B and LD C, that is 2.7 and 4.8 times higher than using a high frequency Kz for the latter two stations. The contrasted  $\text{NO}_3^-$  input observed at the three stations results from the high variability in turbulence along the west-east transects (Bouruet-Aubertot et al., this issue). Thus, using a constant Kz removes the contrasted  $\text{NO}_3^-$  input between the 3 stations (~ 4 times higher at LD A than at LDB and LD C). Consequently, using average Kz values for the turbulent diffusive flux computation can lead to significant bias. In our study,  $\text{NO}_3^-$  input was calculated at the top of the nitracline. Painter et al. (2013) have demonstrated the variability that may be introduced into the estimated  $\text{NO}_3^-$  input by the depth of the defined nitracline. With a constant Kz in the 2 cases, they estimated that  $\text{NO}_3^-$  input was 5 times lower at the top of nitracline depth than at the maximum gradient depth. In our study, the  $\text{NO}_3^-$  input would also be ~ 3-4 times higher if calculated at the maximum gradient depth rather than at the top nitracline, mainly due to the increase of the nitracline gradient up to  $48 \mu\text{mol N m}^{-4}$  (Table 3). However, in all cases, the  $\text{NO}_3^-$  input by turbulence always represented a minor contribution to the N budget.
- Finally, the high contribution of  $\text{N}_2$  fixation to new N input in the photic layer results from the intrinsically high  $\text{N}_2$  fixation rates we measured in the WTSP (especially in MA waters), that are part of the hotspot of  $\text{N}_2$  fixation reported by Bonnet et al., (2017), with rates being in the upper range of rates reported in the global  $\text{N}_2$  fixation Marine Ecosystem Data (MAREDAT) database (Luo et al., 2012). Those high  $\text{N}_2$  fixation rates are as high as westward in the Salomon Sea (Berthelot et al., 2017; Bonnet et al., 2015), extending the hotspot of  $\text{N}_2$  fixation to the whole of the WTSP (Bonnet et al., 2017).

The contribution of  $\text{N}_2$  fixation to PP was around 13-18 % in MA waters and 3 % in SPG waters. The high contribution measured in the MA region is an order of magnitude higher than that reported in previous studies performed in the Pacific Ocean (Moutin et al., 2008; Raimbault and Garcia, 2008; Shiozaki et al., 2014), the Atlantic Ocean (Fonseca-Batista et al., 2017; Rijkenberg et al., 2011) and the Mediterranean Sea (Moutin and Prieur, 2012), where it never exceeds 5%, and also slightly higher than the contribution reported from a mesocosm experiment in the New Caledonia lagoon during a UCYN bloom ( $10.8 \pm 5.0$  %; Berthelot et al., 2015). As there was low supply of  $\text{NO}_3^-$  through vertical diffusion (< 8 %) and atmospheric deposition (< 1.5 %),  $\text{N}_2$  fixation sustains nearly all new production during the austral summer in the WTSP.

## Coupling between N<sub>2</sub> fixation and export in the WTSP

Previous studies have used different methods for coupled measurements of N<sub>2</sub> fixation and export (Berthelot et al., 2015; Dore et al., 2008; Karl et al., 2012; Scharek et al., 1999a; Subramaniam et al., 2008; White et al., 2012). The Lagrangian strategy used here was designed to sample the same water mass during the experiment and therefore minimize the methodological issues associated with particulate export flux measurements using sediment traps in the open ocean (Monroy et al., 2017). The severe meteorological conditions due to the development of tropical cyclone PAM (a category 5 storm) that hit the Vanuatu Islands on March 2015, obliged us to perform the LD B station at a more easterly location than initially planned (Moutin et al., 2017). LD B was therefore sampled in a surface bloom with a DCM close to the surface (Fig. 3), in contrast to LD A and LD C sampled in a zone with a DCM near the bottom of the photic layer (Fig. 3). Thus, data from LD B, although presented together with LD A and LD C, will be discussed apart.

Stations LD A and LD C were considered as oligotrophic and ultra-oligotrophic, respectively. PP was twice higher and the DCM shallower at station LD A, compared to LD C. Furthermore, the diazotroph community composition was contrasted between the two stations, with a clear dominance of *Trichodesmium* at LD A ( $6.6 \cdot 10^4$  nifH gene copies L<sup>-1</sup> at 5 m) and lower abundance of diazotrophs, and a clear domination of UCYN-B ( $3.6 \cdot 10^3$  nifH gene copies L<sup>-1</sup> at 5 m) and het-1 ( $3.0 \cdot 10^3$  nifH copies L<sup>-1</sup> at 5 m) (Stenegren et al., this issue) at LD C. The e-ratio (e-ratio = PC export/PP) calculated at LD A (9.7 %) was higher than the e-ratio in most studied oligotrophic regions (Karl et al., 2012; Moutin and Prieur, 2012; Raimbault and Garcia, 2008), where it rarely exceeds 1 %, indicating a high efficiency of the WTSP to export C relative to PP. Moreover, the e-ratio was higher at LD A (characterized by high N<sub>2</sub> fixation rates ( $593 \mu\text{mol N m}^{-2} \text{d}^{-1}$ )) than at LD C (characterized by low N<sub>2</sub> fixation ( $59 \mu\text{mol N m}^{-2} \text{d}^{-1}$ )). This is in agreement with previous studies reporting typical e-ratios of 1 % in ultra-oligotrophic regions characterized by low N<sub>2</sub> fixation rates (like LD C), such as the eastern SPG (Moutin et al., 2008; Raimbault and Garcia, 2008) or the Mediterranean Sea (Bonnet et al., 2011), and typical e-ratios of 5 % in regions characterized by high N<sub>2</sub> fixation rates such as station ALOHA (Karl et al., 2012). Taken together, these results suggest that N<sub>2</sub> fixation would enhance particle export. This is supported by Knapp et al. (this issue) who showed that nearly all exported production was supported by N<sub>2</sub> fixation in MA waters during the OUTPACE cruise.

At station ALOHA, the e-ratio varies between 2 to 15 % and is maximum during summer export fluxes of PN, that are attributed to the direct export of DDAs (Karl et al., 2012). In the present study, we investigated the potential direct export of diazotrophs by measuring the abundance of each diazotroph group in the traps. We reveal that the export efficiency of *Trichodesmium*, i.e. the percentage of organisms present in the water column recovered in the traps (< 0.1 %), was lower than that of other diazotrophs, which is in agreement with Walsby (1992) and Chen et al. (2003), who revealed that *Trichodesmium* are rarely recovered in the sediment traps. The export efficiency of UCYN-B (2.3 % on average) and het-1 (4.0 % on average) was higher than that of *Trichodesmium*, which is consistent with Bonnet et al., (2016b) and Karl et al., (2012). In a mesocosm experiment performed in the coastal waters of New Caledonia, Bonnet et al., (2016b) revealed that UCYN-C were efficiently exported thanks to aggregation processes. In this study, the contribution of diazotrophs to PC

export was up to 30.6 % at LD A, and was mainly driven by *het-1* as the estimated *het-1* C content was higher than that of UCYN-B and *Trichodesmium*. This suggests that DDAs were efficiently exported, which is in agreement with previous studies (Karl et al., 2012; Subramaniam et al., 2008). At LD C, less than 0.1 % of the total PC measured in the traps was associated with diazotrophs, which is probably due to lower abundances of *het-1* in the traps (< DL) than at LD A and the dominance of UCYN-B (at 330 m) having low cellular C content. The contribution of diazotrophs to PC export at LD A (up to 30.6 % at 330 m) was high compared to what has been measured in a much smaller water column (15 m-high mesocosms) in New Caledonia (ca. 20 %; Bonnet et al., 2016a), and suggests that the direct export of diazotrophs should be further investigated in oligotrophic open ocean. To date, few qPCR data on *nifH* from sediment traps are available (Karl et al., 2012) to compare with our study. However, it has to be noted that we measured the highest export and e-ratio at LD A, where *Trichodesmium* dominated the diazotroph community. This suggests that most of the export was likely indirect, i.e. after the transfer of diazotroph-derived N (DDN) to the surrounding bacterial, phytoplankton and zooplankton communities, as revealed by Caffin et al., (this issue) during the same cruise.

Station LD B was studied during a surface *Trichodesmium* bloom; however observations of poor cell integrity were reported (Stenegren et al., this issue) and other evidence indicated the senescence of the bloom (de Verneil et al., this issue). Higher  $N_2$  fixation and integrated PP rates than those measured at LD A and C together, with lower PN export, resulted in an e-ratio less than 0.8 % (Table 2) at LD B. This very low export efficiency is probably related to the fact that we sampled station LD B during a collapsing *Trichodesmium* bloom (Stenegren, this issue) triggered by  $PO_4^{3-}$  starvation (de Verneil, 2017), as already reported in the WTSP (Moutin et al., 2005). The collapse of *Trichodesmium* blooms can possibly result from viral lysis (Hewson et al., 2004), mainly leading to the release of dissolved N in surrounding waters, or programmed cell death, mainly leading to rapid sinking of biomass, and may influence C export (Bar-Zeev et al., 2013). Programmed cell death (PCD) was detected at LD B (Spungin et al., this issue), indicating that this process cannot be excluded, while *Trichodesmium* were not recovered in the traps (they dominated the export of diazotrophs at 150 m, but altogether the direct export of diazotrophs never exceeded 1.1 % of total export). It is thus likely that most of the N accumulated in the phytoplankton pool (including *Trichodesmium*) was released to the dissolved pool due to grazing and viral lysis, then quickly remineralized due to high microbial activity at LD B (Van Wambeke et al., this issue), and thus would explain the low export rate measured at this station. This result is supported by the efficient transfer of DDN to the surface planktonic food web at this station, as described in Caffin et al. (This issue), and previously by Bonnet et al. (2016b) and Berthelot et al. (2016) in the WTSP. Thus, elevated Chl *a* patches such as those we sampled at LD B may be more productive areas than the ambient oligotrophic waters, but less efficient in terms of export, corresponding to the concept of “high-biomass, low-export” (HBLE) environments initially reported for the Southern Ocean (Lam and Bishop, 2007), where surface waters with high biomass were associated with low particle export at depth.

## Disequilibrium of new vs. exported production

The daily N budget computed here reveals that N input into the photic layer through atmospheric deposition, N<sub>2</sub> fixation and vertical NO<sub>3</sub><sup>-</sup> diffusion exceeded N output through organic matter export at the three studied stations. This imbalance between new and exported production is also observed in different oligotrophic regions of the ocean, such as, for instance, the SPG (Raimbault and Garcia, 2008), the Barents Sea (Olli et al., 2002; Reigstad et al., 2008; Wexels Riser et al., 2008), the North and South Atlantic gyres (Thomalla et al., 2006), and the Equatorial Pacific (Bacon et al., 1996). It should be noted that our budget was performed at the daily scale, but at the annual time scale or longer time scales, the PN export from the photic layer is supposed to balance new N input (Dore et al., 2002; Eppley and Peterson, 1979).

This imbalance between new and exported N frequently reported in the oligotrophic ocean may result from (1) a spatial decoupling between production and export, (2) a temporal decoupling between production and export and/or (3) processes other than particle export such as DON and/or zooplankton export. As we used the Lagrangian strategy described above and confirmed that we sampled the same water at all LD stations during our surveys (de Verneil et al., this issue), the first option (spatial decoupling) can be excluded. The second option (2) would mean that we performed our budget during a period corresponding to production of organic matter that was dissociated from the export that would have occurred later. Such temporal lag has already been reported in the Southern Ocean (Nodder and Waite, 2001), accompanied by biomass accumulation in the photic layer; therefore, we cannot exclude this hypothesis here. Regarding the third possible explanation, the primary process by which organic matter is exported out of the photic layer is the gravitational sinking of particles to the deep ocean (Karl et al., 1996; Knauer et al., 1990). However, two other main processes can contribute to export: physical mixing resulting in export of dissolved organic matter (Carlson et al., 1994; Carlson and Ducklow, 1995; Copin-Montégut and Avril, 1993; Toggweiler, 1989) and zooplankton diel-migrations that actively transport organic matter out of the photic layer (Longhurst et al., 1989, 1990; Longhurst and Glen Harrison, 1988; Vinogradov, 1970). In the present study, the DON export flux was limited to eddy diffusion as we performed our survey during the stratification period, and the low downward flux of DON estimated by Moutin et al., (this issue) was unable to explain the observed imbalance. However, zooplankton might play a significant role, although hard to quantify. Zooplankton living below the photic layer migrate to the surface at night, and when going down, can increase the export of dissolved and particulate organic and inorganic N through defecation, excretion or mortality (Atkinson et al., 1996; Le Borgne and Rodier, 1997; Dam et al., 1993, 1995, Longhurst et al., 1989, 1990; Longhurst and Williams, 1992; Zhang and Dam, 1998) if it occurs below a barrier to vertical mixing (i.e. nitracline or pycnocline (Longhurst et al., 1989, 1990)). Zooplankton has been reported to represent between 4.9 and 38 % of the total export flux (Table 6) in several ecosystems, and therefore should be considered in export calculations. Here, we estimated the maximum contribution of zooplankton using the higher value reported in Table 6 (38 %). It reached a maximum of 106  $\mu\text{mol N m}^{-2} \text{ d}^{-1}$  at LD A, 12  $\mu\text{mol N m}^{-2} \text{ d}^{-1}$  at LD B and 18  $\mu\text{mol N m}^{-2} \text{ d}^{-1}$  at LD C at 150 m. By applying this correction to our export values, it cannot explain the observed disequilibrium between new and exported N.



Finally, the zooplankton themselves are sampled by the traps but dead zooplankton were not distinguishable from live swimmers. In the present study, the zooplankton contribution to PN export (Table 4) was high (68 % on average), and above what has ever been measured in other oligotrophic areas such as the Mediterranean Sea (Moutin and Prieur, 2012). Here, all zooplankton recovered in traps were considered as live swimmers and were therefore discarded, which may have lead to an underestimation of the PN export and could also partly explain the observed disequilibrium between new and export production. Further studies should be undertaken to assess the contribution of living versus dead zooplankton to PN. In summary, we suggest that zooplankton plays a key role on the export in the WTSP and its contribution would increase the particulate export. Moreover, zooplankton activity can transfer N accumulated in the phytoplankton pool to the dissolved pool following grazing and related trophic processes. N from the dissolved pool is then remineralized by microbial activity and accumulates in the photic layer, thus N is not recovered in sediment traps.

### **Methodological underestimation leads to a possible higher contribution of N<sub>2</sub> fixation**

In this study we intentionally used the ‘bubble method’ to measure N<sub>2</sub> fixation rates, considering the small differences observed between this method and the method consisting in adding the <sup>15</sup>N<sub>2</sub> as dissolved in a subset of seawater previously N<sub>2</sub> degassed (Mohr et al., 2010a) in Pacific waters (Bonnet et al. 2016b, Shiozaki et al., 2015) and the high risk of sample contamination involved when manipulating sample seawater to prepare dissolved <sup>15</sup>N<sub>2</sub> (Klawonn et al., 2015). In addition to the contamination issues, preparing dissolved <sup>15</sup>N<sub>2</sub> on board represents additional time with samples sitting on the bench or rosette before incubation, which is especially critical in tropical environments. To reduce any potential underestimation, we measured the <sup>15</sup>N enrichment of the N<sub>2</sub> pool at the end of the incubation ( $7.548 \pm 0.557$  atom%, Bonnet et al., this issue), which was lower than the theoretical value of ~8.2 atom% based on gas constants calculations (Weiss, 1970). We are aware that the dissolution kinetics of <sup>15</sup>N<sub>2</sub> in the incubation bottles is progressive along the 24 h of incubation (Mohr et al., 2010a). Therefore, the <sup>15</sup>N enrichment of the N<sub>2</sub> pool measured with the MIMS likely represent maximum values, and the N<sub>2</sub> fixation rates provided in this study represent minimum values. This reinforces the conclusions of this study regarding the prominent role of N<sub>2</sub> fixation in this region. Großkopft et al. (2012) found that the discrepancy between both methods was more important when UCYN dominates the diazotroph community as compared to when *Trichodesmium* dominates. Consequently, N<sub>2</sub> fixation rates in this study are potentially more underestimated in SPG waters than in MA waters. By applying the maximum factor of underestimation found by Großkopft et al. (2012; i.e. 1.7), N<sub>2</sub> fixation in SPG waters would have been higher ( $100 \mu\text{mol N m}^{-1} \text{d}^{-1}$  instead of 59), which is still far lower than in MA waters and does not change de conclusions of this study.

### **5 Conclusion**

In this study, we successfully used a Lagrangian strategy in the WTSP to follow the same water mass during 5 days in order to perform N budgets during the stratification period (Feb. - March 2015) at 3 stations. N<sub>2</sub> fixation appeared as a substantial

biogeochemical process providing the major external source of N in the photic layer. *Trichodesmium* was the major diazotroph in the oligotrophic MA waters (LD A and LD B), while UCYN dominated the diazotroph community in the ultra-oligotrophic waters of the gyre (LD C). N<sub>2</sub> fixation contributed ~13-18 % of the estimated PP in the MA region where N<sub>2</sub> fixation rates were high, and ~3 % in the SPG water where N<sub>2</sub> fixation rates were low. As there was limited supply of NO<sub>3</sub><sup>-</sup> through vertical turbulent diffusion (< 8 %) and dry atmospheric deposition (<1 %), N<sub>2</sub> fixation accounted for nearly all new production (more than 90 % of new N). The current coupling between typical high N<sub>2</sub> fixation rates of the WTSP, with the high PN and PC export measured in this region associated with high e-ratios (up to ~10 %), suggests that N<sub>2</sub> fixation plays an important role in export during austral summer conditions in the WTSP, either directly or indirectly. The export efficiency measured here in the WTSP (LD A) is comparable to that measured in the Southern Ocean (Rembauville et al., 2015), considered as an efficient ecosystem for C export. The oligotrophic ocean represents 60 % of the global ocean surface, and therefore may play a more significant role in C export than initially considered (Baines et al., 1994; Wassmann, 1990). Even if the link between N<sub>2</sub> fixation rates and export is obvious, the possible temporal decoupling between these processes and the potential role of zooplankton need to be further investigated. Finally, as Bonnet et al. (2017) have recently shown that the WTSP is a hotspot of N<sub>2</sub> fixation, and as we have shown the importance of this process with regard to the N input and N and C export, we suggest that this region of the world ocean should be further investigated by means of oceanographic cruises and the establishment of time-series. This would give us a “big picture” of the role of N<sub>2</sub> fixation on the export in the oligotrophic ocean.

## Acknowledgements

This is a contribution of the OUTPACE (Oligotrophy from Ultra-oligoTrophy PACific Experiment) project (https://outpace.mio.univ-amu.fr/) funded by the French research national agency (ANR-14-CE01-0007-01), the LEFE-CyBER program (CNRS-INSU), the GOPS program (IRD) and the CNES (BC T23, ZBC 4500048836). The project leading to this publication has received funding from European FEDER Fund under project 1166-39417. The OUTPACE cruise (http://dx.doi.org/10.17600/15000900) was managed by MIO (OSU Institut Pytheas, AMU) from Marseilles (France). The authors thank the crew of the RV *L'Atalante* for outstanding shipboard operations. G. Rougier and M. Picheral are warmly thanked for their efficient help in CTD rosette management and data processing, as well as C. Schmechtig for the LEFE-CyBER database management. J. Louis is warmly thanked for the analyses of atmospheric DIN on board. The satellite-derived data of Sea Surface Temperature, Chl *a* concentrations and currents have been provided by CLS in the framework of the CNES funding; we warmly thank I. Pujol and G. Taburet for their support in providing these data. We acknowledge NOAA, and in particular R. Lumpkin, for providing the SVP drifters.

All data and metadata are available at the following web address: <http://www.obs-vlfr.fr/proof/php/outpace/outpace.php>. Argo DOI (<http://doi.org/10.17882/42182>).

## References

- Al-Mutairi, H. and Landry, M. R.: Active export of carbon and nitrogen at station ALOHA by diel migrant zooplankton, *Deep. Res. Part II*, 48(8–9), 2083–2103, doi:10.1016/S0967-0645(00)00174-0, 2001.
- Altabet, M. A.: Variations in nitrogen isotopic composition between sinking and suspended particles: implications for  
5 nitrogen cycling and particle transformation in the open ocean, *Deep Sea Res. Part A, Oceanogr. Res. Pap.*, 35(4), 535–554, doi:10.1016/0198-0149(88)90130-6, 1988.
- Aminot, A. and K  rouel, R.: Dosage automatique des nutriments dans les eaux marines : m  thodes en flux continu, 2007.
- Atkinson, A., Shreeve, R. S., Pakhomov, E. A., Priddle, J., Blight, S. P. and Ward, P.: Zooplankton response to a  
phytoplankton bloom near South Georgia, Antarctica, *Mar. Ecol. Prog. Ser.*, 144(1–3), 195–210, doi:10.3354/meps144195,  
10 1996.
- Bacon, M. P., Cochran, J. K., Hirschberg, D., Hammar, T. R. and Fleer, A. P.: Export flux of carbon at the equator during the  
EqPac time-series cruises estimated from <sup>234</sup>Th measurements, *Deep. Res. Part II*, 43(4–6), 1133–1153, doi:10.1016/0967-  
0645(96)00016-1, 1996.
- Baines, S. R., Pace, M. L. and Karl, D. M.: Why does the relationship between sinking flux and planktonic primary  
15 production differ between lakes and oceans?, *Limnol. Oceanogr.*, 39, 213–226, 1994.
- Bar-Zeev, E., Avishay, I., Bidle, K. D. and Berman-Frank, I.: Programmed cell death in the marine cyanobacterium  
*Trichodesmium* mediates carbon and nitrogen export, *ISME J.*, 7(12), 2340–2348, doi:10.1038/ismej.2013.121, 2013.
- Benavides, M., Berthelot, H., Duhamel, S., Raimbault, P. and Bonnet, S.: Dissolved organic matter uptake by  
*Trichodesmium* in the Southwest Pacific, *Sci. Rep.*, 7, 41315, doi:10.1038/srep41315, 2017.
- 20 Berthelot, H., Moutin, T., L’Helguen, S., Leblanc, K., H  lias, S., Grosso, O., Leblond, N., Charri  re, B. and Bonnet, S.:  
Dinitrogen fixation and dissolved organic nitrogen fueled primary production and particulate export during the VAHINE  
mesocosm experiment (New Caledonia lagoon), *Biogeosciences*, 12(13), 4099–4112, doi:10.5194/bg-12-4099-2015, 2015.
- Berthelot, H., Bonnet, S., Grosso, O., Cornet, V. and Barani, A.: Transfer of diazotroph-derived nitrogen towards non-  
diazotrophic planktonic communities: A comparative study between *Trichodesmium erythraeum* *Crocospaera watsonii* and  
25 *Cyanothece* sp., *Biogeosciences*, 13(13), 4005–4021, doi:10.5194/bg-13-4005-2016, 2016.
- Berthelot, H., Benavides, M., Moisander, P. H., Grosso, O. and Bonnet, S.: High-nitrogen fixation rates in the particulate and  
dissolved pools in the Western Tropical Pacific (Solomon and Bismarck Seas), *Geophys. Res. Lett.*, 2, 1–10,  
doi:10.1002/2017GL073856, 2017.
- Bonnet, S., Biegala, I. C., Dutrieux, P., Slemmons, L. O. and Capone, D. G.: Nitrogen fixation in the western equatorial  
30 Pacific: Rates, diazotrophic cyanobacterial size class distribution, and biogeochemical significance, *Global Biogeochem.*  
*Cycles*, 23(3), 1–13, doi:10.1029/2008GB003439, 2009.
- Bonnet, S., Grosso, O. and Moutin, T.: Planktonic dinitrogen fixation along a longitudinal gradient across the Mediterranean  
Sea during the stratified period (BOUM cruise), *Biogeosciences*, 8(8), 2257–2267, doi:10.5194/bg-8-2257-2011, 2011.

- Bonnet, S., Rodier, M., Turk-Kubo, K. A., Germineaud, C., Menkes, C., Ganachaud, A., Cravatte, S., Raimbault, P., Campbell, E., Quéroué, F., Sarthou, G., Desnues, A., Maes, C. and Eldin, G.: Contrasted geographical distribution of  $N_2$  fixation rates and nifH phylotypes in the Coral and Solomon Seas (southwestern Pacific) during austral winter conditions, *Global Biogeochem. Cycles*, 29(11), 1874–1892, doi:10.1002/2015GB005117, 2015.
- 5 Bonnet, S., Berthelot, H., Turk-Kubo, K. A., Cornet-Barthaux, V., Fawcett, S., Berman-Frank, I., Barani, A., Grégori, G., Dekaezemacker, J., Benavides, M. and Capone, D. G.: Diazotroph derived nitrogen supports diatom growth in the South West Pacific: A quantitative study using nanoSIMS, *Limnol. Oceanogr.*, 61(5), 1549–1562, doi:10.1002/lno.10300, 2016a.
- Bonnet, S., Berthelot, H., Turk-Kubo, K. A., Fawcett, S., Rahav, E., L’Helguen, S. and Berman-Frank, I.: Dynamics of  $N_2$  fixation and fate of diazotroph-derived nitrogen in a low-nutrient, low-chlorophyll ecosystem: Results from the VAHINE mesocosm experiment (New Caledonia), *Biogeosciences*, 13(9), 2653–2673, doi:10.5194/bg-13-2653-2016, 2016b.
- 10 Bonnet, S., Caffin, M., Berthelot, H. and Moutin, T.: Hot spot of  $N^2$  fixation in the western tropical South Pacific pleads for a spatial decoupling between  $N_2$  fixation and denitrification, *Proc. Natl. Acad. Sci.*, 114(14), E2800–E2801, doi:10.1073/pnas.1619514114, 2017.
- Bonnet S., Caffin, M., Berthelot, H., Grosso, O., Guieu, C. and Moutin, T.: Contribution of dissolved and particulate fractions to the Hot Spot of  $N_2$  fixation in the Western Tropical South Pacific Ocean (OUTPACE cruise), this issue.
- 15 Le Borgne, R. and Rodier, M.: Net zooplankton and the biological pump: A comparison between the oligotrophic and mesotrophic equatorial Pacific, *Deep. Res. Part II Top. Stud. Oceanogr.*, 44(9–10), 2003–2023, doi:10.1016/S0967-0645(97)00034-9, 1997.
- Böttjer, D., Dore, J. E., Karl, D. M., Letelier, R. M., Mahaffey, C., Wilson, S. T., Zehr, J. P. and Church, M. J.: Temporal variability of nitrogen fixation and particulate nitrogen export at Station ALOHA, *Limnol. Oceanogr.*, 62(1), 200–216, doi:10.1002/lno.10386, 2017.
- 20 Bourbonnais, A., Lehmann, M. F., Waniek, J. J. and Schulz-Bull, D. E.: Nitrate isotope anomalies reflect  $N_2$  fixation in the Azores Front region (subtropical NE Atlantic), *J. Geophys. Res. Ocean.*, 114(3), 1–16, doi:10.1029/2007JC004617, 2009.
- Bouruet-Aubertot, P., Cuypers, Y., Le Goff, H., Rougier, G., de Verneuil, A., Doglioli, A., Picheral, M., Yohia, C., Caffin, M., Lefèvre, D., Petrenko, A. and Moutin, T. Longitudinal contrast in small scale turbulence along 20°S in the Pacific Ocean: origin and impact on biogeochemical fluxes, this issue.
- 25 Caffin, M., Foster, R., Berthelot, H. and Bonnet, S.: Fate of  $N_2$  fixation in the Western Tropical South Pacific Ocean: Transfer of diazotroph-derived nitrogen to non-diazotrophic communities and export of diazotrophs, this issue.
- Capone, D. G., Burns, J. A., Montoya, J. P., Subramaniam, A., Mahaffey, C., Gunderson, T., Michaels, A. F. and Carpenter, E. J.: Nitrogen fixation by *Trichodesmium* spp.: An important source of new nitrogen to the tropical and subtropical North Atlantic Ocean, *Global Biogeochem. Cycles*, 19(2), 1–17, doi:10.1029/2004GB002331, 2005.
- 30 Carlson, C. A. and Ducklow, H. W.: Dissolved organic carbon in the upper ocean of the central equatorial Pacific Ocean, 1992: Daily and finescale vertical variations, *Deep. Res. Part II*, 42(2–3), 639–656, doi:10.1016/0967-0645(95)00023-J, 1995.

- Carlson, C. A., Ducklow, H. W. and Michaels, A. F.: Annual flux of dissolved organic carbon from the euphotic zone in the northwestern Sargasso Sea, *Nature*, 371(6496), 405–408, doi:10.1038/371405a0, 1994.
- Chen, Y. L. L., Chen, H. Y. and Lin, Y. H.: Distribution and downward flux of *Trichodesmium* in the South China Sea as influenced by the transport from the Kuroshio Current, *Mar. Ecol. Prog. Ser.*, 259, 47–57, doi:10.3354/meps259047, 2003.
- 5 Chen, Y.B., Dominic, B., Mellon, M. T. and Zehr, J. P.: (Circadian rhythm of nitrogenase gene expression in the diazotrophic filamentous nonheterocystous cyanobacterium *Trichodesmium* sp. strain IMS 101. *J Bacteriol* 180: 3598–3605, 1998.
- Church, M. J., Short, C. M., Jenkins, B. D., Karl, D. M. and Zehr, J. P.: Temporal Patterns of nitrogenase gene (*nifH*) expression in the oligotrophic North Pacific Ocean, *Appl. Environ. Microbiol.*, 71, 5362–5370. doi:10.1128/AEM.71.9.5362-5370.2005, 2005.
- 10 Colon-Lopez, M., Sherman, D. M. and Sherman, L. A.: Transcriptional and translational regulation of nitrogenase in light-dark- and continuous-light grown cultures of the unicellular cyanobacterium *Cyanothece* sp. strain ATCC 51142. *J. Bacteriol* 13: 4319–4327, 1997.
- Copin-Montégut, G. and Avril, B.: Vertical distribution and temporal variation of dissolved organic carbon in the North-Western Mediterranean Sea, *Deep. Res. Part I Oceanogr. Res. Pap.*, 40(10), 1963–1972, doi:10.1016/0967-0637(93)90041-Z, 1993.
- 15 de Verneil, A., Rousselet, L., Doglioli, A. M., Petrenko, A. A. and Moutin, T.: The fate of a southwest Pacific bloom: gauging the impact of submesoscale vs. mesoscale circulation on biological gradients in the subtropics, *Biogeosciences*, 14, 3471–3486, <https://doi.org/10.5194/bg-14-3471-2017>, 2017.
- 20 de Verneil A., Rousselet L., Doglioli A., Petrenko A.A., Maes, C., Bouruet-Aubertot P. and Moutin T.: OUTPACE long duration stations: physical variability, context of biogeochemical sampling, and evaluation of sampling strategy, this issue.
- d’Ovidio, F., Della Penna, A., Trull, T. W., Nencioli, F., Pujol, M. I., Rio, M. H., Park, Y. H., Cotté, C., Zhou, M. and Blain, S.: The biogeochemical structuring role of horizontal stirring: Lagrangian perspectives on iron delivery downstream of the Kerguelen Plateau, *Biogeosciences*, 12(19), 5567–5581, doi:10.5194/bg-12-5567-2015, 2015.
- 25 Dam, H. G., Miller, C. A. and Jonasdottir, S. H.: The Trophic Role of Mesozooplankton at 47-Degrees-N, 20-Degrees-W during the North-Atlantic Bloom Experiment, *Deep. Res. Part II Top. Stud. Oceanogr.*, 40(1–2), 197–212, doi:10.1016/0967-0645(93)90013-D, 1993.
- Dam, H. G., Roman, M. R. and Youngbluth, M. J.: Downward export of respiratory carbon and dissolved inorganic nitrogen by diel-migrant mesozooplankton at the JGOFS Bermuda time-series station, *Deep. Res. Part I*, 42(7), 1187–1197, doi:10.1016/0967-0637(95)00048-B, 1995.
- 30 Dekaezemacker, J. and Bonnet, S.: Sensitivity of  $N_2$  fixation to combined nitrogen forms ( $NO_3^-$  and  $NH_4^+$ ) in two strains of the marine diazotroph *Crocospaera watsonii* (Cyanobacteria), *Mar. Ecol. Prog. Ser.*, 438, 33–46, doi:10.3354/meps09297, 2011.

- Dentener, F., Drevet, J., Lamarque, J.-F., Bey, I., Eickhout, B., Fiore, A. M., Hauglustaine, D., Horowitz, L. W., Krol, M., Kulshrestha, U. C. and others: Nitrogen and sulfur deposition on regional and global scales: a multimodel evaluation, *Global Biogeochem. Cycles*, 20(4), 2006.
- Deutsch, C. A., Sarmiento, J. L., Sigman, D. M., Gruber, N. and Dunne, J. P.: Spatial coupling of nitrogen inputs and losses  
5 in the ocean, *Nature*, 445(7124), 163–167, doi:10.1038/nature05392, 2007.
- Doglioli, A. M., Nencioli, F., Petrenko, A. A., Rougier, G., Fuda, J.-L. and Grima, N.: A software package and hardware tools for in situ experiments in a Lagrangian reference frame, *J. Atmos. Ocean. Tech.*, 30, 1940–1950, 2013.
- Dore, J. E., Brum, J. R., Tupas, L. M. and Karl, D. M.: Seasonal and interannual variability in sources of nitrogen supporting export in the oligotrophic subtropical North Pacific Ocean, *Limnol. Ocean.*, 47(6), 1595–1607, 2002.
- 10 Dore, J. E., Letelier, R. M., Church, M. J., Lukas, R. and Karl, D. M.: Summer phytoplankton blooms in the oligotrophic North Pacific Subtropical Gyre: Historical perspective and recent observations, *Prog. Oceanogr.*, 76(1), 2–38, doi:10.1016/j.pocean.2007.10.002, 2008.
- Dron, A., Rabouille, S., Claquin, P., Talec, A., Raimbault, V. and Sciandra, A.: Photoperiod length paces the temporal orchestration of cell cycle and carbon-nitrogen metabolism in *Crocospaera watsonii*, *Environ. Microbiol.*, 15(12), 3292–  
15 3304, doi:10.1111/1462-2920.12163, 2013.
- Eppley, R. W. and Peterson, B. J.: Particulate organic matter flux and planktonic new production in the deep ocean, *Nature*, 282(5740), 677–680, doi:10.1038/282677a0, 1979.
- Fernández-Castro, B., Mouriño-Carballido, B., Marañón, E., Chouciño, P., Gago, J., Ramírez, T., Vidal, M., Bode, A., Blasco, D., Royer, S.-J., Estrada, M. and Simó, R.: Importance of salt fingering for new nitrogen supply in the oligotrophic  
20 ocean, *Nat. Commun.*, 6, 8002, doi:10.1038/ncomms9002, 2015.
- Fonseca-Batista, D., Dehairs, F., Riou, V., Fripiat, F., Elskens, M., Deman, F., Brion, N., Quéroque, F., Bode, M. and Auel, H.: Nitrogen fixation in the eastern Atlantic reaches similar levels in the Southern and Northern Hemisphere, *J. Geophys. Res. Ocean.*, 122(1), 587–601, 2017.
- Foster, R. A. and Zehr, J. P.: Characterization of diatom-cyanobacteria symbioses on the basis of *nifH*, *hetR* and 16S rRNA  
25 sequences, *Environ. Microbiol.*, 8(11), 1913–1925, doi:10.1111/j.1462-2920.2006.01068.x, 2006.
- Foster, R. A., Subramaniam, A., Mahaffey, C., Carpenter, E. J., Capone, D. G. and Zehr, J. P.: Influence of the Amazon River plume on distributions of free-living and symbiotic cyanobacteria in the western tropical north Atlantic Ocean, *Limnol. Oceanogr.*, 52, 517–532. doi: 10.4319/lo.2007.52.2.0517, 2007.
- García, N., Raimbault, P., Gouze, E. and Sandroni, V.: Fixation de diazote et production primaire en Méditerranée  
30 occidentale, *Comptes Rendus - Biol.*, 329(9), 742–750, doi:10.1016/j.crv.2006.06.006, 2006.
- Großkopf, T., Mohr, W., Baustian, T., Schunck, H., Gill, D., Kuypers, M. M. M., Lavik, G., Schmitz, R. a., Wallace, D. W. R. and LaRoche, J.: Doubling of marine dinitrogen-fixation rates based on direct measurements, *Nature*, 488(7411), 361–364, doi:10.1038/nature11338, 2012.

- Gruber, N.: The Marine Nitrogen Cycle: Overview and Challenges, *Nitrogen Mar. Environ.*, 1–50, doi:10.1016/B978-0-12-372522-6.00001-3, 2008.
- Gruber, N. and Sarmiento, J. L.: Global patterns of marine nitrogen fixation and denitrification, *Global Biogeochem. Cycles*, 11(2), 235–266, doi:10.1029/97GB00077, 1997.
- 5 Guieu, C., Bonnet, S., Petrenko, A., Menkes, C., Chavagnac, V., Desboeufs, C., Maes, C., and Moutin, T. : Iron from a submarine source impacts the productive layer of the Western Tropical South Pacific (WTSP), *Scientific Reports*, Submitted.
- Hannides, C. C. S., Landry, M. R., Benitez-Nelson, C. R., Styles, R. M., Montoya, J. P. and Karl, D. M.: Export stoichiometry and migrant-mediated flux of phosphorus in the North Pacific Subtropical Gyre, *Deep. Res. Part I Oceanogr. Res. Pap.*, 56(1), 73–88, doi:10.1016/j.dsr.2008.08.003, 2009.
- 10 Hewson, I., Govil, S. R., Capone, D. G., Carpenter, E. J. and Fuhrman, J. A.: Evidence of Trichodesmium viral lysis and potential significance for biogeochemical cycling in the oligotrophic ocean, *Aquat. Microb. Ecol.*, 36(1), 1–8, doi:10.3354/ame036001, 2004.
- Hynes, A. M., Webb, E. A., Doney, S. C. and Waterbury, J. B.: Comparison of cultured trichodesmium (cyanophyceae) with species characterized from the field, *J. Phycol.*, 48(1), 196–210, doi:10.1111/j.1529-8817.2011.01096.x, 2012.
- 15 JGOFS: Core measurements protocols : report of the core measurement working group, *Jt. Glob. Ocean Flux Study, JGOFS repo*, SCOR 1-40, 1988.
- Kanakidou, M., Duce, R. A., Prospero, J. M., Baker, A. R., Benitez-Nelson, C., Dentener, F. J., ... and Sarin, M. Atmospheric fluxes of organic N and P to the global ocean. *Global Biogeochemical Cycles*, 26(3), 2012.
- Kana, T. M., Darkangelo, C., Hunt, M. D., Oldham, J. B., Bennett, G. E. and Cornwell, J. C.: Membrane Inlet Mass Spectrometer for Rapid High-Precision Determination of N<sub>2</sub>, O<sub>2</sub>, and Ar in Environmental Water Samples, *Anal. Chem.*, 20 66(23), 4166–4170, doi:10.1021/ac00095a009, 1994.
- Karl, D. M., Christian, J. R., Dore, J. E., Hebel, D. V., Letelier, R. M., Tupas, L. M. and Winn, C. D.: Seasonal and interannual variability in primary production and particle flux at Station ALOHA, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 43(2–3), 539–568, doi:10.1016/0967-0645(96)00002-1, 1996.
- 25 Karl, D. M., Letelier, R. M., Tupas, L. M., Dore, J. E., Christian, J. R. and Hebel, D. V.: The role of nitrogen fixation in biogeochemical cycling in the subtropical North Pacific Ocean, *Nature*, 388(6642), 533–538, doi:10.1038/41474, 1997.
- Karl, D. M., Michaels, A. F., Bergman, B., Capone, D. G., Carpenter, E. J., Letelier, R., Lipschultz, F., Paerl, H., Sigman, D. and Stal, L.: Dinitrogen fixation in the world's oceans, *Biogeochemistry*, 57–58, 47–98, doi:10.1023/A:1015798105851, 2002.
- 30 Karl, D. M., Bates, N., Emerson, S., Harrison, P. J., Jeandel, C., Llinás, O., Liu, K. K., Marty, J.-C., Michaels, A. F., Miquel, J. C., Neuer, S., Nojiri, Y. and Wong, C. S.: Temporal studies of biogeochemical processes determined from ocean time-series observations during the JGOFS era, *Ocean Biogeochem.*, 239–265, doi:10.1007/978-3-642-55844-3\_11, 2003.

- Karl, D. M., Church, M. J., Dore, J. E., Letelier, R. M. and Mahaffey, C.: Predictable and efficient carbon sequestration in the North Pacific Ocean supported by symbiotic nitrogen fixation, *Proc. Natl. Acad. Sci.*, 109(6), 1842–1849, doi:10.1073/pnas.1120312109, 2012.
- Klawonn, I., Lavik, G., Böning, P., Marchant, H. K., Dekaezemacker, J., Mohr, W. and Ploug, H.: Simple approach for the preparation of  $^{15}\text{-}^{15}\text{N}_2$ -enriched water for nitrogen fixation assessments: Evaluation, application and recommendations, *Front. Microbiol.*, 6, doi:10.3389/fmicb.2015.00769, 2015.
- Knapp, A. N.: The sensitivity of marine  $\text{N}_2$  fixation to dissolved inorganic nitrogen, *Front. Microbiol.*, 3, 1–14, doi:10.3389/fmicb.2012.00374, 2012.
- Knapp, A. N., Sigman, D. M. and Lipschultz, F.: N isotopic composition of dissolved organic nitrogen and nitrate at the Bermuda Atlantic Time-series study site, *Global Biogeochem. Cycles*, 19(1), 1–15, doi:10.1029/2004GB002320, 2005.
- Knauer, G. A., Redalje, D. G., Harrison, W. G. and Karl, D. M.: New production at the VERTEX time-series site, *Deep Sea Res. Part A, Oceanogr. Res. Pap.*, 37(7), 1121–1134, doi:10.1016/0198-0149(90)90054-Y, 1990.
- Lam, P. J. and Bishop, J. K. B.: High biomass, low export regimes in the Southern Ocean, *Deep Sea Res. Part. II*, 54, 601–638, doi:10.1016/j.dsr2.2007.01.013, 2007.
- Longhurst, A. R. and Glen Harrison, W.: Vertical nitrogen flux from the oceanic photic zone by diel migrant zooplankton and nekton, *Deep Sea Res. Part A, Oceanogr. Res. Pap.*, 35(6), 881–889, doi:10.1016/0198-0149(88)90065-9, 1988.
- Longhurst, A. R. and Williams, R. G.: Carbon flux by seasonal vertical migrant cpoepods is a small number., *J. Plankt. Res.*, 14(1955), 1495–1509, 1992.
- Longhurst, A. R., Bedo, A. W., Harrison, W. G., Head, E. J. H., Horne, E. P., Irwin, B. and Morales, C. E.: NFLUX: a test of vertical nitrogen flux by diel migrant biota, *Deep Sea Res. Part A, Oceanogr. Res. Pap.*, 36(11), 1705–1719, doi:10.1016/0198-0149(89)90067-8, 1989.
- Longhurst, A. R., Bedo, A. W., Harrison, W. G., Head, E. J. H. and Sameoto, D. D.: Vertical flux of respiratory carbon by oceanic diel migrant biota, *Deep Sea Res. Part A, Oceanogr. Res. Pap.*, 37(4), 685–694, doi:10.1016/0198-0149(90)90098-G, 1990.
- Louis, J., Bressac, M., Pedrotti, M. L., and Guieu, C. Dissolved inorganic nitrogen and phosphorus dynamics in seawater following an artificial Saharan dust deposition event. *Frontiers in Marine Science*, 2, 27, 2015.
- Luo, Y. W., Doney, S. C., Anderson, L. A., Benavides, M., Berman-Frank, I., Bode, A., Bonnet, S., Boström, K. H., Böttjer, D., Capone, D. G., Carpenter, E. J., Chen, Y. L. L., Church, M. J., Dore, J. E., Falcón, L. I., Fernández, A., Foster, R. A., Furuya, K., Gómez, F., Gundersen, K., Hynes, A. M., Karl, D. M., Kitajima, S., Langlois, R. J., LaRoche, J., Letelier, R. M., Marañón, E., McGillicuddy, D. J., Moisander, P. H., Moore, C. M., Mouriño-Carballido, B., Mulholland, M. R., Needoba, J. A., Orcutt, K. M. M., Poulton, A., Rahav, E., Raimbault, P., Rees, A. P., Riemann, Shiozaki, T., Subramaniam, A., Tyrrell, T., Turk-Kubo, K. A., Varela, M., Villareal, T. A., Webb, E. A., White, A. E., Wu, J. and Zehr, J. P.: Database of diazotrophs in global ocean: abundance, biomass and nitrogen fixation rates, *Earth Syst. Sci. Data*, 4(1), 47–73, doi:10.5194/essd-4-47-2012, 2012.



- Mahaffey, C., Williams, R. G., Wolff, G. A., Mahowald, N. M., Anderson, W. and Woodward, M.: Biogeochemical signatures of nitrogen fixation in the eastern North Atlantic, *Geophys. Res. Lett.*, 30(6), 33–36, doi:10.1029/2002GL016542, 2003.
- Mahaffey, C., Michaels, A. F. and Capone, D. G.: The conundrum of marine N<sub>2</sub> fixation, *Am. J. Sci.*, 305(6–8 SPEC. ISS.), 546–595, doi:10.2475/ajs.305.6-8.546, 2005.
- Martínez-Pérez, C., Mohr, W., Löscher, C. R., Dekaezemacker, J., Littmann, S., Yilmaz, P., Lehnen, N., Fuchs, B. M., Lavik, G., Schmitz, R. A., LaRoche, J. and Kuypers, M. M. M.: The small unicellular diazotrophic symbiont, UCYN-A, is a key player in the marine nitrogen cycle, *Nat. Microbiol.*, 1(11), 16163, doi:10.1038/nmicrobiol.2016.163, 2016.
- Mohr, W., Großkopf, T., Wallace, D. W. R. and LaRoche, J.: Methodological underestimation of oceanic nitrogen fixation rates, *PLoS One*, 5(9), 1–7, doi:10.1371/journal.pone.0012583, 2010a.
- Mohr, W., Intermaggio, M. P. and LaRoche, J.: Diel rhythm of nitrogen and carbon metabolism in the unicellular, diazotrophic cyanobacterium *Crocospaera watsonii* WH8501. *Environmental Microbiology*, 12: 412–421, doi: 10.1111/j.1462-2920.2009.02078.x, 2010b.
- Moisander, P. H., Beinart, R. A., Voss, M., and Zehr, J. P.: Diversity and abundance of diazotrophic microorganisms in the South China Sea during intermonsoon, *ISME J.*, 2, 954–967, doi:10.1038/ismej.2008.51, 2008.
- Moisander, P. H., Beinart, R. A., Hewson, I., White, A. E., Johnson, K. S., Carlson, C. A., Montoya, J. P. and Zehr, J. P.: Unicellular Cyanobacterial Distributions Broaden the Oceanic N<sub>2</sub> Fixation Domain, *Science*, 327(5972), 1512–1514, doi:10.1126/science.1185468, 2010.
- Montoya, J. P., Voss, M., Kahler, P. and Capone, D. G.: A Simple , High-Precision , High-Sensitivity Tracer Assay for N<sub>2</sub> Fixation, *Appl. Environ. Microbiol.*, 62(3), 986–993, 1996.
- Montoya, J. P., Holl, C. M., Zehr, J. P., Hansen, A., Villareal, T. A. and Capone, D. G.: High rates of N<sub>2</sub> fixation by unicellular diazotrophs in the oligotrophic Pacific Ocean, *Nature*, 430, 1027–1031, doi:10.1038/nature02744.1., 2004.
- Monroy, P., Hernández-García, E., Rossi, V., and López, C.: Modeling the dynamical sinking of biogenic particles in oceanic flow, *Nonlin. Processes Geophys.*, 24, 293–305, <https://doi.org/10.5194/npg-24-293-2017>, 2017.
- Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., Galbraith, E. D., Geider, R. J., Guieu, C., Jaccard, S. L., Jickells, T. D., La Roche, J., Lenton, T. M., Mahowald, N. M., Marañón, E., Marinov, I., Moore, J. K., Nakatsuka, T., Oschlies, A., Saito, M. A., Thingstad, T. F., Tsuda, A. and Ulloa, O.: Processes and patterns of oceanic nutrient limitation, *Nat. Geosci.*, 6(9), 701–710, doi:10.1038/ngeo1765, 2013.
- Moutin, T. and Prieur, L.: Influence of anticyclonic eddies on the Biogeochemistry from the Oligotrophic to the Ultraoligotrophic Mediterranean (BOUM cruise), *Biogeosciences*, 9(10), 3827–3855, doi:10.5194/bg-9-3827-2012, 2012.
- Moutin, T. and Raimbault, P.: Primary production, carbon export and nutrients availability in western and eastern Mediterranean Sea in early summer 1996 (MINOS cruise), *J. Mar. Syst.*, 33–34, 273–288, doi:10.1016/S0924-7963(02)00062-3, 2002.

- Moutin, T., Van Den Broeck, N., Beker, B., Dupouy, C., Rimmelin, P. and Le Bouteiller, A.: Phosphate availability controls *Trichodesmium* spp. biomass in the SW Pacific Ocean, *Mar. Ecol. Prog. Ser.*, 297, 15–21, doi:10.3354/meps297015, 2005.
- Moutin, T., Karl, D. M., Duhamel, S., Rimmelin, P., Raimbault, P., Van Mooy, B. A. S. and Claustre, H.: Phosphate availability and the ultimate control of new nitrogen input by nitrogen fixation in the tropical Pacific Ocean, *Biogeosciences*, 5 (5), 95–109, 2008.
- Moutin, T., Doglioli, A. M., de Verneil, A., and Bonnet, S.: Preface: The Oligotrophy to the UTRa-oligotrophy PACific Experiment (OUTPACE cruise, 18 February to 3 April 2015), *Biogeosciences*, 14, 3207–3220, <https://doi.org/10.5194/bg-14-3207-2017>, 2017.
- Moutin, T., Wagener, T., Fumenia, A., Gimenez, A., Caffin, M., Lefevre, D., Leblanc, K., Bouruet- Aubertot, P., Helias-Nunige, S., Rougier, G., Grosso, O. and deVerneil, A.: Phosphate availability and the ultimate control of the biological carbon pump in the Western Tropical South Pacific Ocean (OUTPACE cruise), this issue.
- Nelson, D. M., Tréguer, P., Brzezinski, M. A., Leynaert, A. and Quéguiner, B.: Production and dissolution of biogenic silica in the ocean: Revised global estimates, comparison with regional data and relationship to biogenic sedimentation, *Global Biogeochem. Cycles*, 9(3), 359–372, doi:10.1029/95GB01070, 1995.
- Nodder, S. D. and Waite, A. M.: Is Southern Ocean organic carbon and biogenic silica export enhanced by iron-stimulated increases in biological production? Sediment trap results from SOIREE, *Deep. Res. Part II Top. Stud. Oceanogr.*, 48(11–12), 2681–2701, doi:10.1016/S0967-0645(01)00014-5, 2001.
- Olli, K., Wexels Riser, C., Wassmann, P., Ratkova, T., Arashkevich, E. and Pasternak, A.: Seasonal variation in vertical flux of biogenic matter in the marginal ice zone and the central Barents Sea, *J. Mar. Syst.*, 38(1–2), 189–204, doi:10.1016/S0924-7963(02)00177-X, 2002.
- Painter, S. C., Patey, M. D., Forryan, A., and Torres-Valdés, S.: Evaluating the balance between vertical diffusive nitrate supply and nitrogen fixation with reference to nitrate uptake in the eastern subtropical North Atlantic Ocean, *J. Geophys. Res. Ocean.*, 118, 5732–5749. doi: 10.1002/jgrc.20416, 2013.
- Petrenko, A. A., Doglioli, A. M., Nencioli, F., Kersalé, M., Hu, Z. and d'Ovidio, F.: A review of the LATEX project: mesoscale to submesoscale processes in a coastal environment, *Ocean Dyn.* 67(3-4), 513–533, 2017.
- Pujo-Pay, M. and Raimbault, P.: Improvement of the wet-oxidation procedure for simultaneous determination of particulate organic nitrogen and phosphorus collected on filters, *Mar. Ecol.-Prog. Ser.*, 105, 203–207, doi:10.3354/meps105203, 1994.
- Raimbault, P. and Garcia, N.: Evidence for efficient regenerated production and dinitrogen fixation in nitrogen-deficient waters of the South Pacific Ocean: impact on new and export production estimates, *Biogeosciences*, 5(2), 323–338, doi:10.5194/bg-5-323-2008, 2008.
- Reigstad, M., Wexels Riser, C., Wassmann, P. and Ratkova, T.: Vertical export of particulate organic carbon: Attenuation, composition and loss rates in the northern Barents Sea, *Deep. Res. Part II Top. Stud. Oceanogr.*, 55(20–21), 2308–2319, doi:10.1016/j.dsr2.2008.05.007, 2008.

- Rembauville, M., Blain, S., Armand, L., Quéguiner, B. and Salter, I.: Export fluxes in a naturally iron-fertilized area of the Southern Ocean – Part 2: Importance of diatom resting spores and faecal pellets for export, *Biogeosciences*, 12, 3171–3195, doi:10.5194/bg-12-3171-2015, 2015.
- Rijkenberg, M. J. A., Langlois, R. J., Mills, M. M., Patey, M. D., Hill, P. G., Nielsdóttir, M. C., Compton, T. J., LaRoche, J. and Achterberg E. P.: Environmental forcing of nitrogen fixation in the Eastern Tropical and Sub-Tropical North Atlantic Ocean, *PLoS ONE*, 6(12), doi:10.1371/journal.pone.0028989, 2011.
- Scharek, R., Tupas, L. M. and Karl, D. M.: Diatom fluxes to the deep sea in the oligotrophic North Pacific gyre at Station ALOHA, *Mar. Ecol. Prog. Ser.*, 182, 55–67, doi:10.3354/meps182055, 1999a.
- Scharek, R., Latasa, M., Karl, D. M. and Bidigare, R. R.: Temporal variations in diatom abundance and downward vertical flux in the oligotrophic North Pacific gyre, *Deep. Res. Part I Oceanogr. Res. Pap.*, 46(6), 1051–1075, doi:10.1016/S0967-0637(98)00102-2, 1999b.
- Shiozaki, T., Kodama, T. and Furuya, K.: Large-scale impact of the island mass effect through nitrogen fixation in the western South Pacific Ocean, *Geophys. Res. Lett.*, 41, 2907–2913, doi:10.1002/2014GL059835, 2014.
- Shiozaki, T., Nagata, T., Ijichi, M. and Furuya, K.: Nitrogen fixation and the diazotroph community in the temperate coastal region of the northwestern North Pacific, *Biogeosciences*, 12(15), 4751–4764, doi:10.5194/bg-12-4751-2015, 2015.
- Short, S. M., Jenkins, B. D. and Zehr, J. P.: The spatial and temporal distribution of two diazotrophic bacteria in the Chesapeake Bay, *Appl. Environ. Microbiol.*, 70, 2186–2192, doi:10.1128/AEM.70.4.2186-2192.2004, 2004.
- Spungin, D., Belkin, N., Foster, R., Stenegren, M., Caputo, A., Pujo-Pay, M., Leblond, N., Dupouy, C., Bonnet, S., and Berman-Frank, I.: Programmed cell death in diazotrophs and the fate of organic matter in the Western Tropical South Pacific Ocean during the OUTPACE cruise, *Biogeosciences Discuss.*, in review, 2018.
- Stenegren, M., Caputo, A., Berg, C., Bonnet, S., and Foster, R. A.: Distribution and drivers of symbiotic and free-living diazotrophic cyanobacteria in the Western Tropical South Pacific, *Biogeosciences Discuss.*, <https://doi.org/10.5194/bg-2017-63>, in review, this issue.
- Stukel, M. R., Ohman, M. D., Benitez-Nelson, C. R. and Landry, M. R.: Contributions of mesozooplankton to vertical carbon export in a coastal upwelling system, *Mar. Ecol. Prog. Ser.*, 491, 47–65, doi:10.3354/meps10453, 2013.
- Stukel, M. R., Benitez-Nelson, C. R., Decima, M., Taylor, A. G., Buchwald, C. and Landry, M. R.: The biological pump in the Costa Rica Dome: An open-ocean upwelling system with high new production and low export, *J. Plankton Res.*, 38(2), 348–365, doi:10.1093/plankt/fbv097, 2015.
- Subramaniam, A., Yager, P. L., Carpenter, E. J., Mahaffey, C., Björkman, K. M., Cooley, S., Kustka, A. B., Montoya, J. P., Sanudo-Wilhelmy, S. A., Shipe, R. and Capone, D. G.: Amazon River enhances diazotrophy and carbon sequestration in the tropical North Atlantic Ocean, *Pnas*, 105(30), 10460–10465, doi:10.1073/pnas.0710279105, 2008.
- Sun, J. and Liu, D.: Geometric models for calculating cell biovolume and surface area for phytoplankton, *J. Plankton Res.*, 25(11), 1331–1346, doi:10.1093/plankt/fbg096, 2003.

- Thomalla, S., Turnewitsch, R., Lucas, M. and Poulton, A.: Particulate organic carbon export from the North and South Atlantic gyres: The  $^{234}\text{Th}/^{238}\text{U}$  disequilibrium approach, *Deep. Res. Part II Top. Stud. Oceanogr.*, 53(14–16), 1629–1648, doi:10.1016/j.dsr2.2006.05.018, 2006.
- Toggweiler, J. R.: Is the downward dissolved organic matter (DOM) flux important in carbon transport?, *Productivity of the Ocean: Present and Past*, 65–83, 1989.
- Verity, P. G., Robertson, C. Y., Tronzo, C. R., Andrews, M. G., Nelson, J. R. and Sieracki, M. E.: Relationships between cell volume and the carbon and nitrogen content of marine photosynthetic nanoplankton, *Limnol. Oceanogr.*, 37(7), 1434–1446, doi:10.4319/lo.1992.37.7.1434, 1992.
- Villareal, T. A.: Division cycles in the nitrogen-fixing rhizosolenia (Baciliariophyceae)-richelia (nostocaceae) symbiosis, *Br. Phycol. J.*, 24(4), 357–365, doi:10.1080/00071618900650371, 1989.
- Van Wambeke F., Duhamel, S. Gimenez, A. Lefèvre, D. Pujo-Pay, M. and Moutin. T.: Dynamics of phytoplankton and heterotrophic bacterioplankton in the Western Tropical South Pacific Ocean along a gradient of diversity and activity of nitrogen fixers, this issue.
- Vinogradov, M. E.: Vertical distribution of the oceanic zooplankton, Nauka, 1970.
- Wagner, T., Guieu, C., Losno, R., Bonnet, S. and Mahowald, N. M.: Revisiting atmospheric dust export to the Southern Hemisphere ocean: Biogeochemical implications, *Global Biogeochem. Cycles*, 22(2), 1–13, doi:10.1029/2007GB002984, 2008.
- Vong, R. J., Vong, I. J., Vickers, D., and Covert, D. S.: Size-dependent aerosol deposition velocities during BEARPEX'07, *Atmos. Chem. Phys.*, 10, 5749–5758, doi:10.5194/acp-10-5749-2010, 2010
- Walsby, A. E.: The gas vesicles and buoyancy of *Trichodesmium*. In *Marine pelagic cyanobacteria: Trichodesmium and other diazotrophs*, Springer Netherlands, 141–161, 1992.
- Wassman, F.: Relationships between primary and export production in the boreal coastal zone of the North Atlantic, *Limnol. Oceanogr.*, 35, 464–471, 1990.
- Wexels Riser, C., Wassmann, P., Reigstad, M. and Seuthe, L.: Vertical flux regulation by zooplankton in the northern Barents Sea during Arctic spring, *Deep. Res. Part II Top. Stud. Oceanogr.*, 55(20–21), 2320–2329, doi:10.1016/j.dsr2.2008.05.006, 2008.
- White, A. E., Foster, R. A., Benitez-Nelson, C. R., Masqué, P., Verdeny, E., Popp, B. N., Arthur, K. E. and Prah, F. G.: Nitrogen fixation in the Gulf of California and the Eastern Tropical North Pacific, *Prog. Oceanogr.*, 109, 1–17, doi:10.1016/j.pocean.2012.09.002, 2012.
- Zhang, X. and Dam, H. G.: Downward export of carbon by diel migrant mesozooplankton in the central equatorial Pacific, 44(9–10), 2191–2202, 1998.

**Table 1: Sediment trap data at the three LD stations. Depth of collection, mean mass flux of dry weight (DW) matter, particulate C and N flux, mean C:N molar ratio. No data was collected at LD C at 520 m. No data was collected at LD C at 520 m.**

Station	depth	Mass flux	PC flux	PN flux	C:N ratio
	m	mg DW m <sup>-2</sup> d <sup>-1</sup>	mg C m <sup>-2</sup> d <sup>-1</sup>	mg N m <sup>-2</sup> d <sup>-1</sup>	106:x
LD A	150	87.2 ± 41.1	27.1 ± 12.1	3.9 ± 1.8	106:13
	330	23.9 ± 15.2	5.8 ± 4.0	0.9 ± 0.8	106:14
	520	22.3 ± 4.6	4.7 ± 1.0	0.6 ± 0.2	106:12
LD B	150	14.1 ± 6.5	3.5 ± 1.3	0.4 ± 0.2	106:10
	330	16.8 ± 9.0	3.2 ± 2.0	0.4 ± 0.2	106:11
	520	17.9 ± 10.5	3.7 ± 1.4	0.5 ± 0.2	106:12
LD C	150	19.6 ± 6.9	3.8 ± 0.8	0.7 ± 0.2	106:17
	330	13.6 ± 6.4	2.6 ± 0.9	0.5 ± 0.2	106:17
	520	-	-	-	-

**Table 2: Mean daily N-budget at the three stations LD A, LD B and LD C.**

Characteristics	Units	LD A	LD B	LD C
$\rho_{NO_3}$	kg m <sup>-3</sup>	23.59	24.34	24.66
$\frac{d[NO_3]}{d\rho}$	μmol kg <sup>-1</sup>	3573	5949	8888
Atmospheric deposition (NOx)	μmol N m <sup>-2</sup> d <sup>-1</sup>	0.90	0.51	0.52
N <sub>2</sub> fixation (in-situ)	μmol N m <sup>-2</sup> d <sup>-1</sup>	593	706	59
NO <sub>3</sub> - diffusion	μmol N m <sup>-2</sup> d <sup>-1</sup>	24	7	5
Σ N inputs	μmol N m <sup>-2</sup> d <sup>-1</sup>	618	714	65
Integrated N-PP	μmol N m <sup>-2</sup> d <sup>-1</sup>	3358	5497	1962
Export N -150 m	μmol N m <sup>-2</sup> d <sup>-1</sup>	279	31	47
Export N - 330 m	μmol N m <sup>-2</sup> d <sup>-1</sup>	64	29	36
Export N - 520 m	μmol N m <sup>-2</sup> d <sup>-1</sup>	44	37	-
e-ratio -150 m	%	9.7	0.7	2.8
e-ratio - 330 m	%	2.2	0.6	2.1
e-ratio - 520 m	%	1.5	0.8	-

**Table 3: Mean turbulent diffusion coefficient (Kz), mean nitracline gradient, mean NO<sub>3</sub><sup>-</sup> flux and associated standard deviations (n=3) over the station occupation at LD A, LD B and LD C at the top nitracline and at the maximum gradient.**

Station	Kz m <sup>2</sup> s <sup>-1</sup>	Nitracline gradient μmol N m <sup>-4</sup>	NO <sub>3</sub> <sup>-</sup> flux μmol N m <sup>-2</sup> d <sup>-1</sup>
<i>Top nitracline</i>			
LD A	1.11 x 10 <sup>-5</sup> ± 1.00 x 10 <sup>-5</sup>	23 ± 13	24.4 ± 24.4
LD B	3.59 x 10 <sup>-6</sup> ± 3.11 x 10 <sup>-6</sup>	21 ± 12	6.7 ± 5.3
LD C	2.04 x 10 <sup>-6</sup> ± 1.11 x 10 <sup>-6</sup>	27 ± 13	4.8 ± 2.2
<i>Maximum gradient</i>			
LD A	1.69 x 10 <sup>-5</sup> ± 1.15 x 10 <sup>-5</sup>	53 ± 10	79 ± 56
LD B	4.52 x 10 <sup>-6</sup> ± 3.22 x 10 <sup>-6</sup>	48 ± 6	19 ± 14
LD C	2.96 x 10 <sup>-6</sup> ± 1.84 x 10 <sup>-6</sup>	48 ± 14	21 ± 11

5

**Table 4: Zooplankton sediment traps data at the three LD stations. Depth of sampling, mean dry weight (DW) zooplankton recovered in the traps, C and N associated to zooplankton.**

Station	depth m	Zooplankton DW (Swimmers) mg m <sup>-2</sup> d <sup>-1</sup>	Zoo-C mg C m <sup>-2</sup> d <sup>-1</sup>	Zoo-N mg N m <sup>-2</sup> d <sup>-1</sup>	C:N ratio 106:x
LD A	150	82.1 ± 18.0	42.3 ± 7.6	8.2 ± 1.2	106:18
	330	376.1 ± 26.1	129.2 ± 15.5	19.5 ± 4.0	106:14
	520	14.1 ± 8.0	5.3 ± 2.2	1.3 ± 0.4	106:22
LD B	150	112.9 ± 38.1	57.1 ± 21.7	11.3 ± 4.7	106:18
	330	62.3 ± 31.6	27.3 ± 15.5	4.9 ± 2.8	106:16
	520	10.5 ± 3.0	4.9 ± 1.2	1.1 ± 0.3	106:20
LD C	150	121.3 ± 37.5	41.7 ± 14.8	10.3 ± 5.0	106:22
	330	31.0 ± 5.1	14.3 ± 5.0	2.9 ± 1.3	106:18
	520	-	-	-	

**Table 5: Contribution of N<sub>2</sub> fixation and NO<sub>3</sub><sup>-</sup> vertical diffusion to new N inputs in oligotrophic region.**

Location	Contribution to new N		Source
	N <sub>2</sub> fixation	NO <sub>3</sub> <sup>-</sup> diffusion	
Tropical North Atlantic	50 %	50 %	Capone et al., 2005
Subtropical North Atlantic	2 %	-	Mourino-Carballido et al., 2011
Subtropical South Atlantic	44 %	-	Mourino-Carballido et al., 2011
South Atlantic Gyre	21 %	24 %	Fernández-Castro et al., 2015
Indian South Subtropical Gyre	12 %	18 %	Fernández-Castro et al., 2015
Mediterranean Sea	0 – 32 %	21 – 53 %	Moutin and Prieur, 2012
			Bonnet et al., 2011
North Pacific subtropical Gyre	30 – 50 %	-	Karl et al., 2003
North Pacific subtropical Gyre	48 %	52 %	Dore et al., 2002
Western Tropical South Pacific	92 – 99 %	1 – 8 %	This study

**Table 6: Contribution of N export by active zooplankton migration to total PN export.**

Location	% of PN export	Source
Subtropical and tropical Atlantic	7.6	Longhurst et al., 1989, 1990
North Atlantic BATS	37.3	Dam et al., 1995
Equatorial Pacific	4.9	Le Borgne and Rodier, 1997
Equatorial Pacific	19.8	Le Borgne and Rodier, 1997
North Atlantic BATS	19.9	Al-Mutairi and Landry, 2001
North Pacific subtropical Gyre	38	Hannides et al., 2009
California Current Ecosystem	20	Stukel et al., 2013
Cost Rica Dome	38	Stukel et al., 2015

Figure captions

- 5 Figure 1: Position of the long duration stations sampled in this study (OUTPACE cruise): LD A in green, LD B in red and LD C in blue on a quasi-Lagrangian surface Chl *a* concentrations map. The in situ production lines were deployed in the photic layer (from 5 to 105, 80 and 180 m for LD A, LD B and LD C, respectively) and the PPS5 sediment traps were deployed at 150, 330 and 520 m.
- 10 Figure 2: Vertical profiles of net N<sub>2</sub> fixation rates (nmol N L<sup>-1</sup> d<sup>-1</sup>) estimated using in situ incubations at day 1 (in situ 1: red circles), day 3 (in situ 2 : orange circles) and day 5 (in situ 3 :purple) and using on-deck incubations (purple filled area) at stations LD A (left), LD B (middle) and LD C (right). The NO<sub>3</sub><sup>-</sup> concentrations averaged over the 5 days of station occupation are also reported (light blue squares : μmol L<sup>-1</sup>), as well as PO<sub>4</sub><sup>3-</sup> concentrations (dark blue squares: μmol L<sup>-1</sup>), and fluorescence/chlorophyll (green dots : μg L<sup>-1</sup>)
- Figure 3: Temporal evolution of PAR, DCM (green line), ρ<sub>NO3</sub> (red line), and 1 % of surface PAR (yellow dots and crosses) during the three stations' occupation period (LD A: top panel, LD B: middle panel, LD C: bottom panel).
- 15 Figure 4: Temporal evolution of upward vertical NO<sub>3</sub><sup>-</sup> flux (μmol N m<sup>-2</sup> d<sup>-1</sup>) calculated at the top of the nitracline for each station (LD A: top panel, LD B: middle panel and LD C: bottom panel) after temporal interpolation (blue). Daily mean from noon to noon in dashed orange line and occupation period mean in green line.
- Figure 5: Relative contribution of each diazotroph (*Trichodesmium* in red, UCYN-B in orange and Het-1 in yellow) to the total PC associated with diazotrophs (diazotroph-PC) in the sediment traps at 150 m (top), 330 m (middle) and 520 m (bottom), at the three stations LD A (left panel), LD B (middle panel) and LD C (right panel). Values in blue correspond to the contribution of diazotroph-PC to total PC measured in the traps. No *Trichodesmium* valid data available at LD A 150 m.








Figure 1



**Figure 2**



**Figure 3**





Figure 4



5 Figure 5