

1 The ²²⁶Ra-Ba relationship in the North Atlantic during 2 GEOTRACES-GA01

3 Emilie Le Roy¹, Virginie Sanial^{1,2,3}, Matthew A. Charette², Pieter van Beek¹, François Lacan¹,
4 Stéphanie H.M. Jacquet⁴, Paul B. Henderson², Marc Souhaut¹, Maribel I. García-Ibáñez^{5,6},
5 Catherine Jeandel¹, Fiz F. Pérez⁵, Géraldine Sarthou⁷
6

7 ¹LEGOS, Laboratoire d'Etudes en Géophysique et Océanographie Spatiales (Université de Toulouse,
8 CNRS/CNES/IRD/UPS), Observatoire Midi Pyrénées, 14 Avenue Edouard Belin, 31400 Toulouse,
9 France [emilie.le.roy@legos.obs-mip.fr]

10 ²Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods
11 Hole, MA 02543, USA

12 ³Now at Department of Marine Science, University of Southern Mississippi, Stennis Space Center, MS
13 39529, USA

14 ⁴Aix Marseille Université, CNRS/INSU, Université de Toulon, IRD, Mediterranean Institute of
15 Oceanography (MIO), UM110, 13288 Marseille, France

16 ⁵Instituto de Investigaciones Marinas (IIM, CSIC), Eduardo Cabello 6, 36208 Vigo, Spain

17 ⁶Now at Uni Research Climate, Bjerknes Centre for Climate Research, Bergen 5008, Norway

18 ⁷Laboratoire des Sciences de l'Environnement Marin (LEMAR), UMR 6539, IUEM, Technopôle Brest
19 Iroise, 29280 Plouzané, France

20 Correspondance to: Emilie Le Roy (*emilie.le.roy@legos.obs-mip.fr*)

21 **Abstract.** We report detailed sections of radium-226 (²²⁶Ra, T_{1/2} = 1602 y) activities and barium (Ba)
22 concentrations determined in the North Atlantic (Portugal-Greenland-Canada) in the framework of the
23 international GEOTRACES program (GA01 section—GEOVIDE project, May-July 2014). Dissolved
24 ²²⁶Ra and Ba are strongly correlated along the section, a pattern that may reflect their similar chemical
25 behavior. Because ²²⁶Ra and Ba have been widely used as tracers of water masses and ocean mixing, we
26 investigated more thoroughly their behavior in this crucial region for thermohaline circulation taking
27 advantage of the contrasting biogeochemical patterns existing along the GA01 section. We used an
28 Optimum Multiparameter (OMP) analysis to distinguish the relative importance of physical transport
29 (water mass mixing) and non-conservative processes (sedimentary, river, or hydrothermal inputs;
30 uptake by particles, and dissolved-particulate dynamics) on the ²²⁶Ra and Ba distributions in the North
31 Atlantic. Results show that the measured ²²⁶Ra and Ba concentrations can be explained by conservative
32 mixing for 58 and 65 % of the samples respectively, notably at intermediate depth, away from the ocean
33 interfaces. ²²⁶Ra and Ba can thus be considered as conservative tracers of water mass transport in the
34 ocean interior at the space scales considered here, namely, of the order of few thousand km. However,
35 regions where ²²⁶Ra and Ba displayed non-conservative behaviors and in some cases decoupled
36 behaviors were also identified, mostly at the ocean boundaries (seafloor, continental margins, and
37 surface waters). Elevated ²²⁶Ra and Ba concentrations found in deep waters of the West European Basin
38 suggest that lower North East Atlantic Deep Water (NEADWL) accumulates ²²⁶Ra and Ba from

1 sediment diffusion and/or particle dissolution during transport. In the upper 1500 m of the West
2 European Basin, deficiencies in ^{226}Ra and Ba are likely explained by their incorporation in planktonic
3 calcareous and siliceous shells, or in barite (BaSO_4) by substitution or adsorption mechanisms. Finally,
4 because Ba and ^{226}Ra display different source terms (mostly deep-sea sediments for ^{226}Ra and rivers for
5 Ba), strong decoupling between ^{226}Ra and Ba were observed at the land-ocean boundaries. This is
6 especially true in the shallow stations near the coasts of Greenland and Newfoundland where high
7 $^{226}\text{Ra}/\text{Ba}$ ratios at depth reflect the diffusion of ^{226}Ra from sediment and low $^{226}\text{Ra}/\text{Ba}$ ratios in the upper
8 water column reflect the input of Ba associated with meteoric waters.

9 1. Introduction

10 The primary source of radium-226 (^{226}Ra , $T_{1/2} = 1602$ y) to the ocean was found to be the diffusion
11 from deep-sea sediments following the decay of its parent isotope, ^{230}Th (Koczy, 1958; Kröll, 1953).
12 This mode of introduction led Koczy to use radium data to derive vertical eddy diffusivities and
13 velocities in the deep sea (Koczy, 1958). Since then, ^{226}Ra has been widely used to study the ocean
14 circulation and mixing at a global scale (Chung and Craig, 1980; Ku et al., 1980). In the framework of
15 the Geochemical Ocean Sections Study (GEOSECS) program, special attention was given to ^{226}Ra as its
16 solubility and half-life made it an ideal candidate as a chronometer of the global thermohaline
17 circulation. In particular, its 1602 y half-life is more adapted than the longer half-life of carbon-14 (^{14}C ,
18 $T_{1/2} = 5730$ y) that had also been used for that purpose. Therefore, the global oceanic distribution of
19 ^{226}Ra in seawater was extensively documented throughout the major ocean basins and a unique database
20 was generated during the GEOSECS program (Broecker et al., 1970, 1967; Chung et al., 1974; Ku et
21 al., 1970; Ku and Lin, 1976).

22 Barium (Ba) is an alkaline earth element like ^{226}Ra . Therefore they share a similar geochemical
23 behavior in the ocean (Chan et al., 1976; Fanning et al., 1988; Mathieu and Wolgemuth, 1973). As such,
24 Ba was proposed as a stable analog of ^{226}Ra in order to use the $^{226}\text{Ra}/\text{Ba}$ ratio as a clock in a similar
25 manner as the $^{14}\text{C}/^{12}\text{C}$ ratio. However, the recognition that ^{226}Ra and Ba participate in upper ocean
26 biological cycles (Ku and Luo, 1994) introduced additional complications for the use of the $^{226}\text{Ra}/\text{Ba}$
27 ratio as a time tracer for deep water ventilation. Both ^{226}Ra and Ba indeed increase with increasing
28 depth, thus reflecting uptake due to biological processes in surface waters, particles scavenging and
29 subsequent release at depth through the dissolution of the settling particles (Broecker et al., 1967; Ku et
30 al., 1970; Ku and Luo, 1994). ^{226}Ra and Ba are thus not only controlled by physical processes, but
31 appear to be incorporated in settling particles such as calcareous and siliceous shells, or in barite
32 (BaSO_4) that precipitates in the mesopelagic zone (Bishop, 1988; Chan et al., 1976; Dehairs et al., 1980;
33 Lea and Boyle, 1989, 1989). Hence, despite different principal sources to the ocean, rivers in the case of

1 Ba and marine sediment diffusion for ^{226}Ra , their distributions are affected by similar processes in the
2 water column. Barium displays a linear correlation with ^{226}Ra in the global ocean, resulting in a fairly
3 constant $^{226}\text{Ra}/\text{Ba}$ ratio of $2.2 \pm 0.2 \text{ dpm } \mu\text{mol}^{-1}$ (dpm, disintegrations per minute) (Chan et al., 1976;
4 Foster et al., 2004; Ku et al., 1980; Li et al., 1973; Östlund et al., 1987). Similarly, strong correlations
5 were also found between Ba-Si (silicate) and ^{226}Ra -Si although no obvious process links ^{226}Ra or Ba
6 with Si. These latter relationships appeared to be more surprising because Si is not a chemical analog of
7 Ra and Ba. It was first proposed that diatom frustules exported from the upper water column could
8 adsorb Ra and Ba, these elements being then released at depth following the dissolution of the siliceous
9 tests (Bishop, 1988; Chung, 1980; Dileep Kumar and Li, 1996). More recent studies showed that the
10 similar behaviors of Ba and Si (and alkalinity) reflect similar dissolved-particulate interactions (Jeandel
11 et al., 1996; McManus et al., 1999; Rubin et al., 2003). Indeed, Ba is not mechanistically coupled with
12 alkalinity or silicate. Rather, the observed relationships may result from the spatial coherence of
13 different carriers (barite, opal and carbonate) overprinted by hydrodynamics. The formation of biogenic
14 silica, CaCO_3 and barite in surface water and their subsequent dissolution in the deeper water column
15 may thus generate parallel oceanic distributions. While barite has been shown to be the main carrier that
16 controls the Ba water column distribution, the relationship between Ba-Ra remains unclear.

17 While the global GEOSECS program provided valuable information on the coupling between
18 biogeochemical cycles of ^{226}Ra and Ba in the ocean, several unknowns still remain. In this work, we
19 take advantage of a new worldwide program, GEOTRACES, to provide new information on the
20 distribution of ^{226}Ra and Ba and their relationship in the ocean. GEOTRACES program aims to
21 characterize the distribution of trace elements and their isotopes (TEIs) (sources, sinks, internal cycling)
22 in the ocean through a global survey consisting of ocean sections and regional process studies.

23 In the present study, we report dissolved ^{226}Ra activities and dissolved Ba concentrations in the
24 North Atlantic Ocean and Labrador Sea (GEOVIDE project, GA01 section). The North Atlantic region
25 hosts a variety of globally significant water masses with complex circulation patterns (García-Ibáñez et
26 al., this issue; Lherminier et al., 2010). This area is crucial for the thermohaline circulation, and thus for
27 global climate, through its important role in the ventilation of the deep layer of the global ocean (Seager
28 et al., 2002). As part of this process, the Meridional Overturning Circulation (MOC) includes the
29 northward transport of warm subtropical waters. These surface waters are then cooled and transformed
30 into subpolar waters, and may reach the Labrador and Irminger Seas where deep-water formation and
31 deep convection take place (Bennett et al., 1985; Pickart and Spall, 2007; Yashayaev et al., 2007). We
32 propose to study the relationship between ^{226}Ra and Ba and to test the conservative behavior of these
33 tracers in this specific region. We further document the Ra-Ba-Si relationship along the GA01 section,

1 as it was done in previous sections conducted during the GEOSECS program and more recently along
2 the GEOTRACES GA03 section.

3 **2. Materials and Methods**

4 **2.1. Study area; the GEOVIDE project**

5 The GEOTRACES GA01 section (GEOVIDE project; PIs : Géraldine Sarthou, LEMAR, France
6 and Pascale Lherminier, LOPS, France) was conducted in the North Atlantic Ocean between Lisbon,
7 Portugal, and St John's, Canada (15 May 2014-30 June 2014; Fig.1). The water samples described here
8 were collected on board the R/V *Pourquoi Pas?*. The section crossed different topographic features and
9 regions with contrasting biogeochemical patterns. It complemented the sections GA03 (U.S.-
10 GEOTRACES) and GA02 (Dutch GEOTRACES) also conducted in the Atlantic Ocean in the
11 framework of the GEOTRACES program. Seventy-eight stations were visited during the GEOVIDE
12 project.

13 **2.2. Sample collection**

14 At 15 of the 78 stations completed during the GA01 cruise, up to 22 discrete 10-L seawater samples
15 were collected through the water column from Niskin bottles. The seawater samples were passed by
16 gravity through 10 g of acrylic fibers impregnated with MnO₂ (so called "Mn fibers"), which
17 quantitatively adsorb radium isotopes (assumed to scavenge 100 % of Ra; van Beek et al., 2010; Moore
18 and Reid, 1973). High-resolution vertical profiles of ²²⁶Ra were thus built to provide a detailed ²²⁶Ra
19 section. The samples were unfiltered since particulate ²²⁶Ra activities are typically two orders of
20 magnitude lower than the dissolved ²²⁶Ra activities (van Beek et al., 2007, 2009). From the same Niskin
21 bottles, 15 mL was collected to determine the Ba concentration, so that Ba and ²²⁶Ra analyses were
22 conducted from the same initial sample, which allows us to investigate the ²²⁶Ra/Ba ratio in the
23 samples. The Ba samples were collected in pre-cleaned polypropylene bottles (rinsed three times with
24 the same seawater sample), acidified with 15 µL of HCl (10 M, Merck, Suprapur) and kept at room
25 temperature for later analysis.

26 **2.3. Analysis of dissolved ²²⁶Ra activities via ²²²Rn emanation**

27 Radium-226 was determined *via* its daughter, radon-222 (²²²Rn; T_{1/2} = 3.8 days) using a radon
28 extraction system followed by alpha scintillation counting (Key et al., 1979). The Mn Fiber samples
29 were placed into gas-tight PVC cartridges (Peterson et al., 2009) that were flushed with helium (He) for
30 5 min at 250 mL min⁻¹. The cartridges were sealed and held for approximately 2 weeks (minimum of 5

1 days) to allow for ^{222}Rn ingrowth from ^{226}Ra decay. The ^{222}Rn was then flushed out from the cartridges
2 using He and cryo-trapped in copper tubing using liquid nitrogen. The copper trap was heated to allow
3 the ^{222}Rn to be transferred to an evacuated “Lucas cells” *via* a stream of He. The “Lucas cells” are air-
4 tight chambers with inner walls coated with silver-activated zinc sulfide that emits photons when struck
5 by alpha decay particles (Key et al., 1979 ; Lucas, 1957 ; Peterson et al., 2009). The cells were held 3
6 hours to reach the secular equilibrium of all ^{222}Rn decay chain daughters. After 3 hours, the samples
7 were counted overnight on a radon counting system (Model AC/DC-DRC-MK 10-2). The counting
8 uncertainties (1SD, Standard Deviation) were within the range of 2-5 % for 10 L volume samples. All
9 samples were appropriately ingrowth and decay corrected. The combined Lucas cell and detector
10 background was ~ 7 % of the typical total measured sample activity. The method was standardized using
11 NIST (U.S. National Institute of Standards & Technology) ^{226}Ra (20 dpm) sorbed onto MnO_2 fiber and
12 analyzed in the same manner as the samples, with uncertainties (1SD) of 5 % (Charette et al., 2015;
13 Henderson et al., 2013). Vertical profiles of ^{226}Ra from the GEOTRACES GA01 (this study) and GA03
14 (Charette et al., 2015) sections that were located in close proximity off Portugal (Fig.1) were compared,
15 and showed a good agreement with increasing activities with increasing depth (Fig.S1).

16 **2.4. Analysis of dissolved Ba concentrations**

17 Barium concentrations were measured using an isotope dilution (ID) method (Freydier et al., 1995;
18 Klinkhammer and Chan, 1990) by high resolution-inductively coupled plasma-mass spectrometry (HR-
19 ICP-MS). This method was adapted to a Thermo Finnigan Element XR instrument (MIO, Marseille).
20 The Ba measurements presented here are the sum of dissolved Ba and a very small fraction of
21 particulate Ba (generally <1 % of total Ba, along GEOVIDE up to 1.3 % at the bottom of station 32 due
22 to presence of a nepheloid layer; Lemaitre et al., this issue) released from the samples as a result of the
23 acidification step. Hence, while the measurements reported herein are total Ba, they are within
24 analytical uncertainty representative of the dissolved Ba pool. The samples (0.5 mL) were spiked with
25 300 μL of a ^{135}Ba -enriched solution (93 % ^{135}Ba ; 95 nmol kg^{-1}) and diluted with 15 mL of acidified
26 (2 % HNO_3 , 14 M, Optima grade) Milli-Q grade water (Millipore). The amounts of sample, spike and
27 dilution water were assessed by weighing. The reproducibility of this method is about 1.5 % (1 RSD,
28 Relative Standard Deviation), as tested on repeated preparations of the reference solution SLRS-5
29 (NRC-CNRC river water reference material for trace metals). Average Ba values obtained for SLRS-5
30 were $13.48 \pm 0.20 \mu\text{g L}^{-1}$ (1 σ) with RSD of 1.5 %, which is in good agreement with the certified values
31 (SLRS-5 $13.4 \pm 0.6 \mu\text{g L}^{-1}$). The limit of detection calculated as three times the standard deviation of the
32 procedural blank was 0.09 nmol L^{-1} .

2.5. Multiparameter mixing model

An Optimum MultiParameter (OMP) analysis was used to distinguish the relative importance of physical transport (i.e., water mass mixing) from non-conservative processes (input from the sediments, rivers or hydrothermal vents, dissolution of particles; uptake by particles and dissolved-particulate dynamics) on the ^{226}Ra and Ba distributions in the North Atlantic. We used the OMP analysis computed for the GA01 section by Garcia Ibanez et al. (this issue) with 12 source water types (SWTs). Based on historical data reported from the North Atlantic, we defined ^{226}Ra and Ba endmember concentrations associated with each SWT (Table S1). The characteristics of SWTs (potential temperature, salinity, and geographical location) reported by Garcia Ibanez et al. (this issue) were used to determine the SWTs endmembers for ^{226}Ra and Ba. In some cases, data from the GA01 section were used for the SWT endmember (Table S1). These ^{226}Ra and Ba SWT endmembers were then used to calculate the ^{226}Ra and Ba concentrations that strictly result from mixing of the different water masses. In this way, we estimated the conservative component of ^{226}Ra and Ba, which can in turn be compared to the *in situ* concentrations to generate the non-conservative component of ^{226}Ra and Ba along the GA01 section.

The uncertainties for the SWT endmembers were considered to be ± 0.6 dpm 100 L^{-1} for ^{226}Ra and ± 1.5 nmol L^{-1} for Ba, based on typical measurement uncertainties and spatial variability. A Monte Carlo method (250 random perturbations) was used to propagate the SWT endmember uncertainties to the ^{226}Ra and Ba results of the OMP analysis. This allowed us to estimate conservative component uncertainties of ± 0.9 dpm 100 L^{-1} for ^{226}Ra and ± 4.4 nmol L^{-1} for Ba. When the measured ^{226}Ra and Ba values were different from the conservative component values (taking into account the uncertainties on these values), ^{226}Ra and Ba were considered as non-conservative. These non-conservative values can either be positive (representative of a net addition of ^{226}Ra and Ba) or negative (representative of a net removal of ^{226}Ra and Ba).

Note that the OMP analysis was not solved where non-conservative behavior of temperature and salinity is expected (that is, for waters above 100 m and for waters with salinities lower than 34.7). In these latter cases, changes in water mass properties may indeed be due to air-sea interaction or inputs of fresh waters (e.g., near Greenland shelf; Daniault et al., 2011).

3. Results

3.1. Hydrodynamic context

The OMP analysis was used to identify the different water masses (Table 1) crossing the GA01 section. The potential temperature-salinity diagram for all the GA01 stations along with the different

1 SWT endmembers used in the OMP analysis are represented in Fig.2. The salinity section is shown in
2 Fig.3. The different water masses present along the GA01 section are described below.

3 **3.1.1. Upper waters**

4 Three main water masses were found in upper waters (<1000 m) in the investigated area (Fig.3).
5 First, the Central Waters occupied the upper eastern part of the GA01 section from the Iberian Peninsula
6 to the Rockall Through (stations 1 to 26). Their distribution was associated with the circulation of the
7 North Atlantic Current (NAC). The NAC transports warm and saline waters northward, connecting the
8 subtropical and the subpolar latitudes, and is part of the upper layer of the Atlantic Meridional
9 Overturning Circulation (AMOC) in the North Atlantic subpolar gyre. The NAC flows eastward from
10 the Grand Banks of Newfoundland, splitting into four branches west of the Mid-Atlantic Ridge (MAR),
11 while incorporating local water masses (Fig.1). East of the MAR, the two northern branches of the NAC
12 flow northward into the Icelandic Basin, the Rockall Plateau and the Rockall Trough, while the two
13 southern branches flow southward into the West European Basin. The Central Waters can be identified
14 by the highest potential temperature of the entire GA01 section and are represented by two endmembers
15 called East North Atlantic Central Waters (ENACW₁₆ and ENACW₁₂). The ENACW₁₆ is warmer
16 (16 °C) than the ENACW₁₂ that can be identified with a potential temperature of 12.3 °C (Fig.2).

17 Part of the Central Waters carried by the NAC recirculates toward the Iceland Basin and the
18 Irminger Sea, leading to the formation of subpolar mode waters by mixing and cooling in the subpolar
19 gyre (Lacan and Jeandel, 2004; McCartney, 1992). Iceland Subpolar Mode Water (IcSPMW) is formed
20 in the Icelandic Basin and is located, along GA01, over the Reykjanes Ridges (stations 32 and 38)
21 (Fig.3). The IcSPMW is described by two endmembers, the SPMW₇ and the SPMW₈, which are
22 distinguished by their potential temperature of 7.0 °C and 8.0 °C, respectively (Fig.2). Once formed, the
23 IcSPMW follows the Irminger Current.

24 Finally, the Irminger Subpolar Mode Water (IrSPMW) is the result of the transformation of the
25 Central Waters and the IcSPMW, and is formed northwest of the Irminger Sea (Krauss, 1995). The
26 IrSPMW is located near Greenland (stations 53, 57 and 60) (Fig.3) (García-Ibáñez et al., 2015; Lacan
27 and Jeandel, 2004; Read, 2000).

28 **3.1.2. Intermediate waters**

29 The Subarctic Intermediate Water (SAIW) originates in the Labrador Current (Read, 2000). The
30 SAIW is associated with the advection of waters from the Labrador Sea within the NAC; it subducts
31 below the Central Waters at approximately 600 m. Low salinities (34.8 and 34.7) and potential

1 temperatures of 4.5 °C and 6 °C are representative of the two SAIWs, SAIW₄ and SAIW₆, respectively
2 (Fig.2).

3 Around the Rockall Plateau, the SAIW overlies the Mediterranean Water (MW). The MW enters
4 the North Atlantic through the Gibraltar Strait and flows northward while extending westward. The MW
5 can be identified in the West European Basin at approximately 1200 m (stations 1 and 13 in Fig.3) with
6 a salinity of 36.5 (Fig.2; García-Ibáñez et al., 2015).

7 The Labrador Sea water (LSW) is found in multiple locations and at different water depths along
8 the GA01 section (Fig.3). The LSW is formed by progressive cooling and freshening in winter due to
9 deep convection. The LSW can be characterized by its minimum salinity (34.87) and its minimum
10 potential temperature (3 °C) (Fig.2). The LSW contributes to the stratification of the interior of the
11 North Atlantic and its boundary currents and spreads at intermediate depths in three different basins
12 intersected by the GA01 section (Fig.1). The three independent pathways are: (i) northward into the
13 Irminger Sea (station 44), (ii) eastward across the MAR, through the Charlie-Gibbs fracture zone, then
14 northward into the Iceland Basin (station 32) and eastward into the West European Basin (stations 21
15 and 26), and (iii) equatorward as a major component of the North Atlantic Deep Water in the Deep
16 Western Boundary Current (DWBC), which constitutes the lower limb of the AMOC. Along these
17 paths, the LSW mixes with both the overlying and underlying water masses and becomes warmer and
18 saltier (Lazier, 1973).

19 The Polar Intermediate Water (PIW) is characterized by very low salinity (34.9) and potential
20 temperature (less than 2 °C) (Fig.2) and is defined as a separate upper core on the Greenland slope. The
21 PIW is episodically injected into the Irminger Sea and originates from either the Arctic Ocean or the
22 Greenland shelf (Falina et al., 2012; Jenkins et al., 2015; Rudels et al., 2002).

23 **3.1.3. Overflow waters and deep waters**

24 The Iceland-Scotland Overflow Water (ISOW) originates at the Iceland-Scotland sill, and entrains
25 the overlying warm saline atlantic waters (SPMW and LSW). ISOW identification features are a
26 potential temperature of 2.6 °C and a salinity of 34.98 (Fig.2; van Aken and Becker, 1996). ISOW was
27 found at stations located on the Eastern flank of the Reykjanes Ridge (stations 32 and 38) and near
28 Greenland (stations 60 and 64) at great depth (2000-3500 m) (Fig.3).

29 Overflow waters coming from the Denmark Strait mix with both the SPMW and the LSW during
30 descent into the Irminger Sea to form the Denmark Strait Overflow Water (DSOW) (Fig.1) (Read,
31 2000; Yashayaev and Dickson, 2008). DSOW is located at the northern end of the Irminger Sea
32 (station 44) and occupies the deepest part of the Greenland continental slope (stations 69 and 77)

1 (Fig.3). At bottom depth, DSOW is easily identified by a minimum potential temperature of 1.3 °C
2 (Fig.2).

3 In the Southern Ocean, the Antarctic Bottom water (AABW) is formed by deep winter convection
4 of surface waters. AABW flows to the north along the eastern side of the Atlantic and contributes to the
5 formation of the lower North East Atlantic Deep Water (NEADWL) once this water penetrates the
6 Iberian Abyssal Plain by crossing the Mid-Atlantic Ridge (Fig.1). The NEADWL is laying at the bottom
7 of the West European Basin (stations 1 to 26 in Fig.3) with a mean salinity of 34.895 and a potential
8 temperature of 1.98 °C (Fig.2). Then, the NEADWL recirculates into the Rockall Trough and meets
9 ISOW in the Iceland Basin (van Aken, 2000; McCartney, 1992; Schmitz and McCartney, 1993).

10 3.2. Distribution of ^{226}Ra and Ba along the GA01 section

11 The ^{226}Ra distribution for the GA01 section is presented in Fig.4 with Ba concentrations and
12 $^{226}\text{Ra}/\text{Ba}$ ratios. The ^{226}Ra activities and Ba concentrations in the water column range from 7 to
13 25 dpm 100 L⁻¹ and from 33.6 to 81.5 nmol L⁻¹, respectively. These data are in good agreement with
14 Atlantic data from the GEOSECS program, which range from 6.8 to 23.4 dpm 100 L⁻¹ for ^{226}Ra and
15 from 35 to 105 nmol L⁻¹ for Ba (Broecker et al., 1976).

16 For both ^{226}Ra and Ba, the vertical gradient is stronger in the eastern part of the section (West
17 European Basin) than on the western part of the section (from Reykjanes Ridge to Newfoundland). Both
18 are particularly high near the seafloor in the West European Basin (^{226}Ra : 14-25 dpm 100 L⁻¹; Ba:
19 63-82 nmol L⁻¹) and are in agreement with data previously reported for this region (Broecker et al.,
20 1976; Charette et al., 2015). At intermediate depths, Ba concentrations range from 40 to 50 nmol L⁻¹ in
21 the West European Basin (stations 1 and 21) and ^{226}Ra activities range from 9.5 to 10.9 dpm 100 L⁻¹,
22 also in good agreement with literature data (Charette et al., 2015; Schmidt and Reyss, 1996). Low ^{226}Ra
23 and Ba are found in upper waters of the West European Basin and the Iceland Basin
24 (8.1-8.9 dpm 100 L⁻¹ and 35-43 nmol L⁻¹, respectively). Intermediate ^{226}Ra activities and Ba
25 concentrations (9 dpm 100 L⁻¹ and 42 nmol L⁻¹, respectively) are observed in bottom waters in Irminger
26 and Labrador Seas. Between the Reykjanes Ridge and Newfoundland, ^{226}Ra activities range between 7-
27 10 dpm 100 L⁻¹ in surface and intermediate waters. Similar to ^{226}Ra , Ba concentrations are relatively
28 low in this area, ranging from 39-50 nmol L⁻¹. The distributions in ^{226}Ra and Ba are to a first order
29 explained by the different water masses present in the region, as discussed below.

1 4. Discussion

2 4.1. ^{226}Ra -Ba and ^{226}Ra -Ba-Si relationships

3 A linear correlation between ^{226}Ra and Ba is observed for all data collected along the GA01 section
4 (Fig.5). The slope of the ^{226}Ra -Ba linear regression obtained by this study in the North Atlantic is
5 2.5 ± 0.1 (2SD) $\text{dpm } \mu\text{mol}^{-1}$ which agrees with the slope of the ^{226}Ra -Ba linear regression of
6 $2.3 \text{ dpm } \mu\text{mol}^{-1}$ reported during the GEOSECS program for all the oceanic basins (Chan et al., 1976).
7 The intercept on the horizontal Ba axis is 4.4 nmol L^{-1} for the GA01 section, which is in agreement with
8 GEOSECS data (Chan et al., 1976; Li et al., 1973). This positive intercept may be the result of a larger
9 riverine Ba input relative to ^{226}Ra (Ku and Luo, 1994). The slope of the ^{226}Ra -Ba linear regression
10 reported from the GEOSECS program is similar from one oceanic basin to the other. The $^{226}\text{Ra}/\text{Ba}$ ratio
11 (slightly different from the slope) is also fairly constant throughout the global ocean
12 ($2.2 \pm 0.2 \text{ dpm } \mu\text{mol}^{-1}$; Östlund et al., 1987). This pattern indicates that ^{226}Ra and Ba may behave
13 similarly in the ocean. Since ^{226}Ra and Ba are incorporated in settling particles such as calcareous and
14 siliceous shells or barite (BaSO_4) by substitution or adsorption mechanisms (Bishop, 1988; Dehairs et
15 al., 1980; Lea and Boyle, 1989, 1990) and are then released at depth following the dissolution of these
16 particles, the constant $^{226}\text{Ra}/\text{Ba}$ ratio suggests that fractionation between ^{226}Ra and Ba during these
17 processes is small.

18 Investigations conducted during the GEOSECS program further concluded that ^{226}Ra and Ba were
19 tightly correlated to orthosilicic acid ($\text{Si}(\text{OH})_4$) (Chan et al., 1976; Chung, 1980; Ku et al., 1970; Ku and
20 Lin, 1976) despite the fact that ^{226}Ra , Ba, and $\text{Si}(\text{OH})_4$ exhibit different chemical behavior in the water
21 column and different source terms. A Ra-Ba-Si relationship is also observed along the GA01 section
22 (Fig.5). $\text{Si}(\text{OH})_4$ concentrations generally increase with increasing depth, with a steeper gradient in the
23 West European Basin (Introduction Paper, 2017; This issue), as it was also the case for ^{226}Ra and Ba
24 (Fig.S2). The link between ^{226}Ra , Ba and Si has been shown to reflect parallel dissolved-particulate
25 interactions between barite and biogenic silica (Bishop, 1988; Chung, 1980; Jacquet et al., 2005, 2007;
26 Jeandel et al., 1996); the main carrier of ^{226}Ra in the ocean, however, remains an open question. The
27 oceanic Ba-Si and ^{226}Ra -Si relationships may thus be the result of the interaction between ocean
28 biogeochemistry and the water mass transport.

29 In contrast to the ^{226}Ra -Ba relationship, the slope of the ^{226}Ra - $\text{Si}(\text{OH})_4$ relationship observed during
30 GEOSECS exhibited significant spatial variability from one oceanic basin to the other (Li et al., 1973).
31 First, ^{226}Ra and Si are not chemical analogues, as it is the case for ^{226}Ra and Ba. Second, the variability
32 observed in the ^{226}Ra - $\text{Si}(\text{OH})_4$ relationship may be related to the large variability of the $\text{Si}(\text{OH})_4$
33 distribution which is mostly governed by the preformed nutrient concentrations of waters feeding into

the main thermocline from surface waters of the Southern Ocean (Sarmiento et al., 2007). In the case of GA01, the ^{226}Ra - $\text{Si}(\text{OH})_4$ linear regression slope is 2.4 ± 0.9 (2SD) $10^3 \text{ dpm mol}^{-1}$. As a comparison, the ^{226}Ra - $\text{Si}(\text{OH})_4$ slope reported for the GEOTRACES-GA03 section conducted south of the GA01 section in the Atlantic Ocean was $2.1 \cdot 10^3 \text{ dpm mol}^{-1}$ (Charette et al., 2015). The slope of the ^{226}Ra - $\text{Si}(\text{OH})_4$ linear regression is $34.3 \cdot 10^3 \text{ dpm mol}^{-1}$ in the Pacific Ocean and $14.5 \cdot 10^3 \text{ dpm mol}^{-1}$ in the Antarctic Ocean. The ^{226}Ra - $\text{Si}(\text{OH})_4$ relationship has an intercept with the vertical axis of $7.3 \pm 0.1 \text{ dpm } 100 \text{ L}^{-1}$, which represents the residual ^{226}Ra resulting from the total usage of Si in surface waters (Ku et al., 1970). According to (Shannon and Cherry, 1971), the removal of ^{226}Ra in the upper waters is limited by Si. For both the ^{226}Ra -Ba and ^{226}Ra - $\text{Si}(\text{OH})_4$ relationships, several values are clearly outside of the linear regression trend (Fig.5), a pattern that indicates deviation from the relationship usually observed between ^{226}Ra and Ba (or $\text{Si}(\text{OH})_4$). Such deviations may result from non-conservative processes.

4.2. ^{226}Ra and Ba distributions and their relationship with hydrography

A striking feature of the GA01 section is that the ^{226}Ra activities and Ba concentrations are particularly high in the West European Basin below 2000 m (Fig.4), in the NEADWL. This pattern can also be observed in the GA03 section conducted south of the GA01 section (Charette et al., 2015), the two sections being separated by only ca. 500 km in that basin. The NEADWL is mainly formed from waters with a southern origin (Read, 2000). South of the Antarctic Convergence, the surface waters contain high ^{226}Ra activities from the upwelling of deep waters enriched in ^{226}Ra associated with the circumpolar current (Ku and Lin, 1976). The convection of these surface waters leads to the formation of the ^{226}Ra -rich AABW that circulates northward into the Atlantic Ocean. However, the high ^{226}Ra activities of the NEADWL cannot be solely explained by the high ^{226}Ra activities of these waters of southern origin. This will be discussed in section 4.3.1.

In contrast, the lowest ^{226}Ra activities and Ba concentrations reported on the GA01 section are associated with the Central Waters (upper waters of the West European Basin; Fig.4). Central Waters are derived from the NAC and mix with the SAIW and the SPMW. Along their path, Central Waters remain in the upper water column, and therefore are not affected by the deep sedimentary source of ^{226}Ra . West of the Iceland Basin between 200 and 400 m (stations 32 and 38), an increase in the ^{226}Ra activities and Ba concentrations is associated with the IcSPMW.

A slight increase in ^{226}Ra is observed between 1000-1600 m in the West European Basin (Fig.4; Stations 1 and 13) where a salinity maximum is identified. This pattern is associated with the MW. This is corroborated by the slightly higher Ba concentrations and lower ^{226}Ra /Ba ratios, as observed in the Western Mediterranean Sea (van Beek et al., 2009), these waters spreading westward into the Atlantic

1 Ocean. At these stations, between 30 and 79 % of the water found at 1000-1600 m is of Mediterranean
2 origin (MW), according to the OMP analysis (Garcia Ibanez et al., 2017; this issue).

3 Relatively uniform and low ^{226}Ra activities and Ba concentrations are found between the surface
4 and 2500 m in the Labrador Sea, up to 2000 m in the Iceland Basin and deeper in the Irminger Basin
5 (Fig.4). These distributions can be related to the LSW which is formed by winter convection in the
6 Labrador Sea (Fröb et al., 2016; Pickart et al., 2003; Yashayaev and Loder, 2016). When formed, the
7 LSW transports the characteristics of surface waters to the deep ocean (i.e., low ^{226}Ra activities and low
8 Ba concentrations). The LSW then spreads into the Irminger and the Iceland Basin while conserving its
9 low ^{226}Ra and Ba signatures. Relatively low ^{226}Ra activities and Ba concentrations are found at bottom
10 depths in the Irminger and Labrador Seas and may be associated with DSOW, which is also a recently
11 ventilated water mass (Lazier, 1973).

12 Finally, according to the OMP analysis, ISOW is present at several stations along the GA01
13 section (Garcia-Ibanez et al., this issue). First, on the eastern flank of the Reykjanes Ridge (station 32),
14 68 % of the water between 2700 and 3000 m is considered as ISOW. Then, in the Labrador Sea
15 (stations 69 and 77), an average of 58 % of the water between 2100 and 3000 m is identified as ISOW.
16 The slight increase in ^{226}Ra activities and Ba concentrations observed at these locations may thus be
17 related to the ISOW.

18 **4.3. Conservative versus non-conservative behavior of ^{226}Ra and Ba**

19 The use of an Optimum Multiparameter (OMP) analysis allowed us to distinguish the relative
20 importance of physical transport (i.e., mixing of water masses) from non-conservative processes on the
21 ^{226}Ra , Ba and $^{226}\text{Ra}/\text{Ba}$ ratios distributions in the North Atlantic (Fig.6). The comparison between the
22 vertical profiles of ^{226}Ra and Ba determined along the GA01 section, and those derived from OMP
23 analysis (Fig.7) clearly indicates deviations from the conservative behavior and reflects either an input
24 of ^{226}Ra or Ba (positive anomalies highlighted in red; same color code as in Fig.6) or a removal of ^{226}Ra
25 or Ba (negative anomalies highlighted in blue; same color code as in Fig.6). This comparison reveals
26 that for 58 % of the samples ^{226}Ra can be considered as conservative (activities due to mixing and
27 transport) along the GA01 section (i.e., 58 % of the ^{226}Ra anomalies are within the $[-0.9$ and
28 $0.9 \text{ dpm } 100 \text{ L}^{-1}]$ interval), whereas for 65 % of samples Ba can be considered as conservative (i.e.,
29 65 % of the Ba anomalies are within the $[-4.4$ and $4.4 \text{ nmol L}^{-1}]$ interval). A major finding of this study
30 is that ^{226}Ra and Ba are predominantly conservative at intermediate depths: mostly between 500 m to
31 2000 m, but slightly deeper in the center of deep basins such as at stations 21, 44 and 69 (Fig.6). These
32 locations correspond to the depths where the waters are far from the main sources and sinks of ^{226}Ra
33 and Ba. The non-conservative ^{226}Ra (42 % of the ^{226}Ra) is mostly distributed close to the interfaces such

1 as surface/subsurface waters and bottom waters (both in the deep West European Basin and the
2 Labrador Sea), near the main sources (seafloor or shallow sediments deposited onto the margins). The
3 non-conservative Ba is mostly distributed in the upper 1500 m and in the deep West European Basin
4 (Fig.6).

5 The $^{226}\text{Ra}/\text{Ba}$ ratios are also reported for all samples in Fig.7. The mean ratio determined along the
6 GA01 section is identical to the ratio determined during the GEOSECS program ($2.2 \pm 0.2 \text{ dpm } \mu\text{mol}^{-1}$;
7 Östlund et al., 1987). 77 % of the $^{226}\text{Ra}/\text{Ba}$ ratios determined along the GA01 section are within the
8 confidence interval [$2.0\text{--}2.4 \text{ dpm } \mu\text{mol}^{-1}$], indicating little deviation from the mean ratio, a pattern that is
9 likely related to the similar chemical behavior between ^{226}Ra and Ba.

10 4.3.1. ^{226}Ra inputs and their relationship with Ba

11 Deep waters of the West European Basin display positive ^{226}Ra and Ba anomalies (stations 1 to
12 26; Fig.7). The ^{226}Ra anomalies are initiated at shallower depths (ca. 300-2000 m) than the Ba
13 anomalies (ca. 1000—2000m) (Fig.7). As mentioned above, the NEADWl—which is the main water
14 mass of the deep West European Basin—is mainly formed from waters with a southern origin (mainly
15 AABW) that are characterized by high ^{226}Ra and Ba concentrations. However, these southern waters
16 experience a very specific history along their northward transport to the GA01 section that suggests that
17 the high ^{226}Ra activities (and Ba) of the NEADWl cannot be solely explained by the high ^{226}Ra
18 activities (and Ba) of these waters of southern origin. In order to explain the positive ^{226}Ra and Ba
19 anomalies in the deep waters of the West European Basin, we thus need to investigate the fate of ^{226}Ra
20 and Ba in the waters of southern origin that travel northward and reach section GA01. Figure 8 was
21 computed combining GEOSECS and TTO data (^{226}Ra , $\text{Si}(\text{OH})_4$, salinity and temperature) associated
22 with the AABW (Spencer, 1972) that travels northward between 60°S and 40°N in the West Atlantic
23 Basin. The same data (^{226}Ra , $\text{Si}(\text{OH})_4$, salinity and temperature) associated with the NEADWl in the
24 East Atlantic Basin and along GA01 are also reported.

25 Between 60°S and the equator, the high ^{226}Ra activities of the AABW are associated with
26 relatively low salinity, low temperature, and high $\text{Si}(\text{OH})_4$ (Fig.8). Then, while crossing the Mid-
27 Atlantic Ridge at the equator and at 11°N , the AABW undergoes several important transformations:
28 ^{226}Ra activities and $\text{Si}(\text{OH})_4$ concentrations decrease while salinity and temperature tend to increase
29 (Fig.8). The ^{226}Ra and Ba endmembers for the NEADWl were chosen at this specific location to
30 coincide both geographically and with the characteristics (salinity, temperature and $\text{Si}(\text{OH})_4$) of the
31 NEADWl endmembers used for the OMP analysis (Fig.8; Fig.S3). This endmember has been chosen far
32 from the GA01 section in the OMP analysis (García-Ibáñez et al., this issue), because between 11°N
33 and the GA01 section (Fig.8), salinity, temperature, and $\text{Si}(\text{OH})_4$ concentrations display relatively

1 constant trends indicating no major modifications. In contrast, the ^{226}Ra activities display a significant
2 spatial variability north of 11°N , and clearly increase towards the north (Fig.8). This ^{226}Ra increase is
3 thus decoupled from salinity, temperature, and $\text{Si}(\text{OH})_4$, and likely explains the positive anomalies
4 deduced from the OMP analysis in the deep West European Basin (Fig.7). The specific history of these
5 waters of southern origin (waters initially with a high ^{226}Ra activity; decrease in the ^{226}Ra activity at the
6 equator and at 11°N ; new increase of ^{226}Ra activity north of 11°N) suggest that the ^{226}Ra anomalies
7 observed in the West European Basin are explained by inputs of ^{226}Ra along the northward transport of
8 these waters.

9 The positive anomalies result from the input of ^{226}Ra (and Ba) following either i)
10 dissolution/remineralization of settling particles that incorporated ^{226}Ra and Ba in the upper water
11 column (this includes the dissolution of barite, since the waters of the Atlantic Ocean are undersaturated
12 with respect to barite; Monnin et al., 1999) and/or ii) diffusion of ^{226}Ra and Ba from deep-sea sediments
13 (Cochran and Krishnaswami, 1980) (see 4.4). Of special note are stations in the West European Basin,
14 which are especially deep (down to 5500 m). Deep sediments generally display elevated ^{230}Th activities
15 due to scavenging of ^{230}Th from the entire water column (Bacon and Anderson, 1982; Nozaki, 1984).
16 The highest dissolved ^{230}Th activities reported along the GA01 section were thus observed in the deep
17 waters of the West European Basin (Deng et al., 2017, this issue). Consequently, because ^{226}Ra is
18 produced by the decay of ^{230}Th in the sediment, the ^{226}Ra diffusive flux in this area is expected to be
19 especially high.

20 The input of ^{226}Ra in the West European Margin is accompanied by a Ba input since i) positive Ba
21 anomalies are also observed in the deep waters and ii) the $^{226}\text{Ra}/\text{Ba}$ ratios do not significantly deviate
22 from the mean GEOSECS $^{226}\text{Ra}/\text{Ba}$ ratio (Fig.7a). One exception is found at station 21 in the West
23 European Basin, which displays high $^{226}\text{Ra}/\text{Ba}$ at approximately 4000 m (up to $3.17 \text{ dpm } \mu\text{mol}^{-1}$). At
24 several stations (21, 26, 32, 38, 44, 60, 64 and 77), lower beam transmission values near the seafloor
25 indicate presence of suspended sediments likely associated with a nepheloid layer. Nepheloid layers are
26 turbid layers formed episodically by strong and intense abyssal currents that are transported along
27 isopycnals away from the site of resuspension of bottom sediments (McCave, 1986). The presence of a
28 benthic nepheloid layer is also indicated by high particulate iron concentrations at these stations
29 (Gourain et al., 2017; this issue). Such process may thus contribute to release ^{226}Ra (and potentially Ba)
30 to the deep water column, following desorption or dissolution of the particles. Similar ^{226}Ra maxima
31 have been observed in the deep waters of the Northeast Pacific suggesting that the ^{226}Ra flux is not
32 uniform over the ocean bottom even on a regional scale (Chung, 1976). Suspended particle dissolution
33 may also play a role here, notably for Ba. This will be discussed in more detailed in section 4.4.

1 Positive ^{226}Ra anomalies are also found in deep waters at several other stations located in
2 relatively deep basins (> 1200 m) along the GA01 section (e.g. stations 32, 38, 44, 60, 64, 69 and 77).
3 Most of these anomalies are associated with $^{226}\text{Ra}/\text{Ba}$ ratios higher than $2.4 \text{ dpm } \mu\text{mol}^{-1}$. The ^{226}Ra
4 positive anomalies observed at the stations mentioned above are thus best explained by the diffusion of
5 ^{226}Ra from the sediment. However, these latter stations do not exhibit a positive Ba anomaly and Ba
6 tends to be conservative. Consequently, the $^{226}\text{Ra}/\text{Ba}$ ratios in the deep waters of these stations are often
7 significantly higher than the mean GEOSECS value (stations 21, 32, 38, 60, 64; Fig.7). This pattern is
8 different to that observed in the West European Basin, a discrepancy that may be explained by the
9 different sediment composition in the two regions, by the different residence time of deep waters in
10 contact with deep-sea sediments (Chung, 1976) and/or different role played by suspended particles
11 dissolution.

12 A strong ^{226}Ra positive anomaly is observed in the deepest sample collected at station 38 above
13 the Reykjanes Ridge. It cannot be completely excluded that this positive anomaly is attributed to
14 hydrothermal vent since hydrothermal systems are known in the area (Fig.1). High particulate iron and
15 aluminum concentrations were also observed at these stations (Gourain et al., 2017 ; Menzel et al.,
16 2017 ; this issue). Enrichment in ^{226}Ra have indeed been observed in hydrothermal systems plume at
17 mid-ocean Ridges (Kadko, 1996; Kadko and Moore, 1988; Kipp et al., 2017; Rudnicki and Elderfield,
18 1992). Moreover, the ^{226}Ra enrichments are accompanied by slight Ba enrichments, which may support
19 the hydrothermal origin hypothesis, since hydrothermal venting at mid-ocean Ridge constitutes the
20 second major external source of Ba to the ocean (Edmond et al., 1979).

21 Finally, high $^{226}\text{Ra}/\text{Ba}$ ratios variations are also observed in shallow coastal waters (Fig.7c). At
22 stations 53 and 61, high $^{226}\text{Ra}/\text{Ba}$ ratios are found close to the bottom, in agreement with the input of
23 ^{226}Ra from the sediment, whereas low $^{226}\text{Ra}/\text{Ba}$ ratios are found in subsurface at stations 57, 61 and 78,
24 in association with low salinities (Fig.S2). The low $^{226}\text{Ra}/\text{Ba}$ ratios are thus explained by the input of
25 meteoritic waters in coastal areas, since such waters are known to be the predominant source of Ba to
26 the ocean (Martin and Meybeck, 1979; Wolgemuth and Broecker, 1970). At these shallow stations, the
27 different source terms between ^{226}Ra and Ba therefore leads to important variations of the $^{226}\text{Ra}/\text{Ba}$
28 ratios (Fig.7c.; stations 53, 57, 61 and 78). These observations clearly indicate that ^{226}Ra may
29 sometimes be decoupled from Ba.

30 4.3.2. ^{226}Ra removal and its relationship with Ba

31 Relatively few ^{226}Ra negative anomalies were found along the GA01 section. At the deep open-
32 ocean stations, the negative anomalies are mostly observed in the upper 1000 m (Fig.7; stations 13, 21,
33 26, 32, 38, 44 and 77), but can also be found as deep as 2000 m (i.e., stations 64 and 69). In most cases,

1 the negative ^{226}Ra anomalies are associated with significant negative Ba anomalies (stations 13,21,
2 26, 38, 44, 64 and 69). Such features are likely explained by biological mediated processes including
3 incorporation of ^{226}Ra and Ba in planktonic as calcareous and siliceous shells (Bishop, 1988), in
4 acantharian skeletons made of celestite (SrSO_4) or in barite (BaSO_4) crystals (van Beek et al., 2007;
5 Chow and Goldberg, 1960; Shannon and Cherry, 1971; Szabo, 1967; Wolgemuth and Broecker, 1970).

6 Particularly low dissolved $^{226}\text{Ra}/\text{Ba}$ ratios ($<2 \text{ dpm } \mu\text{mol}^{-1}$) are found in the upper 50 m at
7 stations 21, 32, 64, 69 and 77, a pattern that was also observed in the upper 150 m of the Sargasso Sea,
8 where van Beek et al., (2007) reported similar low dissolved $^{226}\text{Ra}/\text{Ba}$ ratios that were accompanied by
9 high $^{226}\text{Ra}/\text{Ba}$ ratios in suspended particles. This latter pattern was attributed to the incorporation of
10 ^{226}Ra and Ba in acantharian skeletons. The low dissolved $^{226}\text{Ra}/\text{Ba}$ ratios (e.g. $1.7 \text{ dpm } \mu\text{mol}^{-1}$,
11 station 77) observed in the upper 200 m along the GA01 section may thus be attributed to acantharians,
12 which have skeletons that incorporate ^{226}Ra preferentially to Ba (van Beek et al., 2007, 2009; Bernstein
13 et al., 1998). The presence of Acantharians was not studied during GEOVIDE. However, previous
14 studies reported presence of acantharians in this area, like for example in the Iceland Basin and in the
15 East Greenland Sea (Antia et al., 1993; Barnard et al., 2004; Martin et al., 2010).

16 Several phytoplankton blooms were observed along the GA01 section, as indicated by the
17 chlorophyll a concentrations (Chl-a) (Fig.S4). The largest bloom was observed in the Labrador Sea in
18 May 2014. Diatoms were the dominant species in the Irminger and Labrador Seas and on the Greenland
19 and Newfoundland margins during GA01 (up to 55 % of the total Chl-a concentration; Tonnard et al., in
20 prep). In the West European Basin, Chl-a was lower in May and June 2014 and coccolithophorids were
21 the dominant species in that area (Tonnard et al., in prep). In these two regions, diatom frustules and
22 coccolithophorids may thus contribute to the removal of ^{226}Ra and Ba (Bishop, 1988; Dehairs et al.,
23 1980), from the water column in these areas that were characterized by noticeable negative anomalies.

24 Additionally, because the Labrador Sea was sampled in June, during the decline of the bloom,
25 barite that is presumably formed following the decay of settling organic matter may also contribute to
26 the removal of Ba (and ^{226}Ra). High particulate excess Ba (Ba_{xs}) concentrations were indeed observed at
27 stations displaying significant Ba negative anomalies: Ba_{xs} concentrations reached a maximum at 400 m
28 at station 13, 21 and 26) and between 400 and 800 m near Greenland, at stations 44, 64 and 69
29 (Lemaitre et al., 2017, *Ba paper*, this issue). These Ba_{xs} profiles can be related to the relatively high
30 particulate organic carbon (POC) export flux determined at these stations (e.g. at station 69, Lemaitre et
31 al., 2017, *Export paper*; this issue). This POC flux would promote barite formation in subsurface
32 (Dehairs et al., 1980; Legeleux and Reyss, 1996) but also deeper in the water column (van Beek et al.,
33 2007). Similarly, Jullion et al., (2017)—by using a parametric OMP analyses as applied in the
34 Mediterranean Sea—also reported quantification of the non-conservative component of the Ba signal

1 and suggested that the Ba negative anomalies potentially reflected Ba subtraction during barite
2 formation occurring during POC remineralization. The winter deep convection in the Labrador Sea may
3 also potentially explain this relatively deep Ba anomalies by transporting negative Ba and ^{226}Ra
4 anomalies waters toward the deep-sea (Jullion et al., 2017). With the exception of the acantharian
5 skeletons that may impact the dissolved $^{226}\text{Ra}/\text{Ba}$ ratios in the upper 200 m, the removal of ^{226}Ra and Ba
6 that takes place deeper in the water column or that involves other processes (e.g. barite precipitation)
7 does not seem to affect significantly the dissolved $^{226}\text{Ra}/\text{Ba}$ ratios.

8 In the shallow coastal stations, lower $^{226}\text{Ra}/\text{Ba}$ ratios are observed (Fig.7). These low ratios may
9 also result from the removal of ^{226}Ra and Ba by planktonic shells and/or barite or scavenging by
10 lithogenic particles. However, because these stations are coastal stations, various processes are at play
11 in a relatively shallow water column (i.e. diffusion of ^{226}Ra from the sediments; input of Ba from
12 meteoritic water; removal of Ba and ^{226}Ra by shells and barite) thus complicating the interpretation of
13 the vertical profiles. We note that the low $^{226}\text{Ra}/\text{Ba}$ ratios observed in surface of shallow stations near
14 the coast of Greenland (stations 57 and 61) and Newfoundland (station 78) are associated with low
15 salinities (Fig.7c). This decoupling between ^{226}Ra and Ba may be explained by input of freshwater into
16 the coastal zone.

17 Finally, at several stations, a decrease in the ^{226}Ra activities is observed near the seafloor
18 (stations 13, 21, 44, 60, 64 and 77; Fig.7). Similar decreasing trends near the seafloor have been
19 reported in the Southwest Atlantic and North Pacific for ^{230}Th (Deng et al., 2014; Okubo et al., 2012), a
20 reactive element that strongly adsorbs onto suspended particles. This trend for ^{230}Th was explained by
21 nuclide scavenging at the seafloor (Deng et al., 2014; Okubo et al., 2012). Radium-226—although it is
22 much less particle-reactive than ^{230}Th —and Ba may also be scavenged by resuspended particles near
23 the seafloor and may adsorb onto the surfaces of Mn oxides (Moore and Reid, 1973). High particulate
24 trace element concentrations were also observed at stations 26, 38, and 69 and may be related to
25 nepheloid layers that impact the deep water column up to 200-300 m above the seafloor (Gourain et al.,
26 2017 ; Menzel et al., 2017 ; this issue).

27 4.4. Estimation of ^{226}Ra and Ba input fluxes into the West European Basin

28 A strong ^{226}Ra positive anomaly is observed in the NEADWI between stations 1 and 21 and below
29 3500 m. On average, it is $3.3 \text{ dpm } 100 \text{ L}^{-1}$ over this vertical section. This anomaly reflects a
30 concentration difference between the ^{226}Ra measured along GA01 and the ^{226}Ra due to water mass
31 mixing. This concentration difference is associated to the northward transport of the NEADWI,
32 estimated to be $0.9 \pm 0.3 \text{ Sv } (10^6 \text{ m}^3 \text{ s}^{-1})$ at 45°N (GA01 section) (Daniault et al., 2016; McCartney,

1992). Therefore, the positive concentration anomaly can be converted into a ^{226}Ra flux that has to be added to this water mass, $F_{\text{Input-Ra}}$, calculated as follows:

$$F_{\text{Input-Ra}} = A \times T_{\text{NEADWl}} \quad (1)$$

where A is the mean positive anomaly of ^{226}Ra (in dpm m^{-3}) determined by the OMP analysis; T_{NEADWl} is the transport associated with the NEADWl (in $\text{m}^3 \text{s}^{-1}$).

This ^{226}Ra input may be associated with a sedimentary source and/or may result from the dissolution of suspended particles. In a first place, we will study the hypothesis of the sedimentary source; the suspended particle source will be discussed in a second place.

The NEADWl at 45 °N is made of up to 92 % of the 11 °N NEADWl endmember. Therefore, the sedimentary input along the northward transport of the NEADWl is calculated across a sediment area between 11 °N and 45 °N (Fig.S3). We consider the distance of 4209 km between 11 °N and the GA01 section and the distance of 1475 km between stations 1 and 21. This leads to a horizontal area of $6.21 \times 10^6 \text{ km}^2$ (assuming a constant bathymetry), across which the sedimentary input is assumed to take place.

The ^{226}Ra flux diffusing out of bottom sediments, $F_{\text{Sed-Ra}}$ (in $\text{dpm cm}^{-2} \text{y}^{-1}$) can be calculated using Eq. (2), assuming that the anomaly is entirely explained by the sediment source:

$$F_{\text{Sed-Ra}} = \frac{F_{\text{Input-Ra}}}{S} \quad (2)$$

where S is the surface area described above (in cm^2) and $F_{\text{Input-Ra}}$ is $1.67 \times 10^8 \text{ dpm s}^{-1}$.

The calculated $F_{\text{Sed-Ra}}$ is $14.8 \pm 3.1 \times 10^{-3} \text{ dpm cm}^{-2} \text{y}^{-1}$, which is within the range of fluxes reported in the literature. For example, Cochran (1980) reported $F_{\text{Sed-Ra}}$ in the range of $1.5 \times 10^{-3} \text{ dpm cm}^{-2} \text{y}^{-1}$ for the Atlantic Ocean to $2.1 \times 10^{-1} \text{ dpm cm}^{-2} \text{y}^{-1}$ in the Pacific Ocean (Fig.8). Li et al. (1973) estimated ^{226}Ra fluxes diffusing out of the sediment in the Southern Ocean and on the Antarctic shelf of $6.2 \times 10^{-4} \text{ dpm cm}^{-2} \text{y}^{-1}$ and $1.6 \times 10^{-3} \text{ dpm cm}^{-2} \text{y}^{-1}$, respectively. The $F_{\text{Sed-Ra}}$ calculated here is thus slightly higher than the ^{226}Ra sedimentary fluxes reported in the Atlantic Ocean by Cochran (1980). Note, however, that the ^{226}Ra fluxes released from the sediments vary locally as a function of the sedimentary ^{230}Th activity, bioturbation rates, but also as a function of the sediment type and composition (Chung, 1976; Cochran, 1980). The ^{226}Ra fluxes reported in the Atlantic Ocean by Cochran (1980), which are the lowest of all basins, are only available for calcareous ooze type sediment (Cochran, 1980). The NEADWl may cross different types of sediments along its northward path in the Atlantic Ocean. This includes calcareous oozes, fine-grained calcareous sediments and clay (Dutkiewicz et al., 2015). In particular, ^{226}Ra diffusion is expected to be higher in these two latter sediment types (Cochran, 1980).

As for Ba is concerned, the mean positive anomalies deduced from the OMP analysis is 7.0 nmol L^{-1} leading to a $F_{\text{Input-Ba}}$ of 69.5 mol s^{-1} . In the same way as ^{226}Ra , a Ba sedimentary flux $F_{\text{Sed-Ba}}$

1 of $3.19 \pm 1.4 \text{ nmol cm}^{-2} \text{ y}^{-1}$ would be required to explain the Ba anomalies observed in the West
 2 European Basin. This flux is on the low end of the Ba sedimentary fluxes reported in different ocean
 3 basins, which range from 1.0 to 30 $\text{nmol cm}^{-2} \text{ y}^{-1}$ (Chan et al., 1977; Falkner et al., 1993; McManus et
 4 al., 1999; Paytan and Kastner, 1996).

5 Alternatively, the dissolution of settling particles could also contribute to the ^{226}Ra and Ba
 6 anomalies observed in the deep waters of the West European Basin. Assuming steady state, we may
 7 undertake a mass balance calculation for particulate ^{226}Ra and Ba in the same box as described above
 8 (i.e., box defined between 11 °N and the GA01 section, between stations 1 and 21 –1475 km–and
 9 between 3500 m depth and the seafloor; Fig.S5). Particles enter the box from above as settling particles,
 10 but also horizontally, carried within the water masses at 11 °N that travel northward. Particles leave the
 11 box by different processes (accumulation in the sediment or northward transport by the water masses) or
 12 alternatively may dissolve while settling in the box. In the absence of precise information about the
 13 particulate ^{226}Ra and Ba fluxes entering and exiting the box horizontally (i.e. the particulate ^{226}Ra and
 14 Ba concentrations at 11 °N and at the GA01 section), we assume that they are of equal importance and
 15 therefore that they cancel each other in the mass balance calculation.

16 The vertical particulate flux entering the box, from above, can be calculated as follows:

$$F_{Part-x} = Cp_{3500} \times Vs \times S \quad (3)$$

17
 18 where x is either ^{226}Ra or Ba, Cp_{3500} is either the particulate ^{226}Ra activities or the particulate Ba
 19 concentrations at 3500 m, Vs is the settling speed for suspended particles and S is the horizontal surface
 20 area described above ($6.21 \cdot 10^6 \text{ km}^2$).

21 We use the value of 0.007 dpm 100 L⁻¹ for the mean ^{226}Ra particulate activity at 3500 m, a value
 22 that was reported for the Atlantic Ocean, Sargasso Sea (van Beek et al., 2007) and the value of
 23 0.087 nmol L⁻¹ for the mean Ba particulate concentration at 3500 m, a value that was determined along
 24 the GA01 section (Lemaitre et al., this issue). We use the settling speeds (Vs) reported for suspended
 25 particles in the literature and that typically range from 100 to 1000 m y⁻¹ (Bacon and Anderson, 1982;
 26 Krishnaswami et al., 1976; Roy-Barman et al., 2002). The F_{Part} thus obtained range from $1.4 \cdot 10^6$ to
 27 $13.8 \cdot 10^6 \text{ dpm s}^{-1}$ for ^{226}Ra , while the F_{Part} range from 1.7 and 17.2 mol s⁻¹ for Ba. Of this total F_{Part} , some
 28 fraction may dissolve while settling, while the remainder will accumulate in the sediment. This
 29 dissolution flux is named $F_{dissolution-x}$, where x is either ^{226}Ra or Ba. We use the sediment Ba
 30 accumulation rates reported by Gingele and Dahmke (1994) in the Atlantic Ocean to calculate the
 31 particulate Ba flux that exits the box ($F_{Accumulation-Ba}$: 2.0 to 13.4 mol s⁻¹); hence, by difference the
 32 $F_{dissolution-Ba}$ is 0-15.2 mol s⁻¹ (Fig.10). This value is of the same order of magnitude of the $F_{Input-Ba}$

1 needed to explain the Ba anomalies (6.28 mol s^{-1}). Therefore, in the case of Ba, the dissolution of
 2 settling particles may entirely explain the OMPA-derived anomalies. The sediment ^{226}Ra accumulation
 3 rates can be calculated from the Ba accumulation rates estimated above, using the $^{226}\text{Ra}/\text{Ba}$ ratio
 4 determined in sinking particles collected in the Sargasso Sea near the seafloor (i.e., $1.5 \text{ dpm } \mu\text{mol}^{-1}$; van
 5 Beek et al., 2007). The sediment ^{226}Ra accumulation flux thus calculated, $F_{\text{Accumulation-Ra}}$, is $2.9 \cdot 10^6$ -
 6 $19.6 \cdot 10^6 \text{ dpm s}^{-1}$, leading to $F_{\text{dissolution-Ra}}$ of 0 - $10.9 \cdot 10^6 \text{ dpm s}^{-1}$ (Fig.10). Therefore, $F_{\text{dissolution-Ra}}$ cannot
 7 account for more than 37 % of the required ^{226}Ra flux ($F_{\text{Input-Ra}}$). This implies that even if the settling
 8 speed is high (1000 m y^{-1} ; high turnover of the particles), the particle dissolution cannot account for the
 9 entire $F_{\text{Input-Ra}}$. The remaining (minimum of 63 %) therefore has to be sustained by ^{226}Ra diffusion from
 10 the sediments.

11 While the above calculations have to be taken with caution given the numerous assumptions in the
 12 mass balance model, overall they suggest that the ^{226}Ra positive anomalies observed in the West
 13 European Basin may be explained entirely by ^{226}Ra that diffuses out of the sediments. However, it
 14 cannot be excluded that the dissolution of settling particles also contributes to the ^{226}Ra enrichment. In
 15 contrast, the Ba positive anomalies may be explained either by the diffusion of Ba from sediment or by
 16 the dissolution of settling particles or by a combination of these two processes. These conclusions are in
 17 line with the current knowledge about ^{226}Ra and Ba sources in the deep ocean (Broecker et al., 1970;
 18 Chan et al., 1976, 1977; Ku et al., 1980).

19 5. Conclusion

20 We investigated the distribution of dissolved ^{226}Ra activities and Ba concentrations in the North
 21 Atlantic Ocean along the GA01 section. To a first order, the ^{226}Ra and Ba patterns appear to be
 22 correlated to the water masses (e.g. high ^{226}Ra and Ba in NEADWL in the West European Basin; low
 23 ^{226}Ra and Ba in Central Waters; slight increase of ^{226}Ra in the MW). Using a mixing model, we show
 24 that the measured ^{226}Ra and Ba concentrations can be explained by conservative mixing for 58 % and
 25 65 % of the samples respectively, notably at intermediate depth (mostly between 1000 m and 2000 m)
 26 and slightly deeper in the middle of deep basins, away from the ocean interfaces. These locations
 27 correspond to the depths where the waters are away from the main sources of ^{226}Ra and Ba. ^{226}Ra and
 28 Ba can thus be considered as conservative tracers of water mass transport in the ocean interior at the
 29 space scales considered here, namely, of the order of few thousand km.

30 Our study also highlighted several regions where significant input or loss of ^{226}Ra and Ba takes
 31 place. In the West European Basin, the deep waters (NEADWL) accumulate both ^{226}Ra and Ba while
 32 flowing from 11°N to the GA01 section. Mass balance calculations suggest that these enrichments are
 33 predominantly explained by sediment diffusion for ^{226}Ra , with a possible contribution from suspended

1 particle dissolution, while both the sediment and suspended particle dissolution could significantly
2 contribute to the Ba enrichments. This pattern contrasts with that observed in the deep waters collected
3 elsewhere along the section that do not display Ba enrichments associated to the ^{226}Ra enrichments.
4 Bottom nepheloid layers may also contribute to the release of ^{226}Ra (and Ba) to the deep water column
5 at several stations. Interestingly, nepheloid layer processes seem to also act as potential removal of
6 ^{226}Ra (and Ba) at several other stations. Significant input of Ba—likely associated with meteoritic
7 waters—is found in the upper water column near Greenland. Finally, ^{226}Ra and Ba are removed from
8 the upper water column, likely primarily due to biological mediated processes such as incorporation of
9 ^{226}Ra and Ba into barite (BaSO_4) that are presumably formed following the decay of settling organic
10 matter and/or adsorption onto diatom frustules, a mechanism that would explain the ^{226}Ra -Ba
11 relationship reported here. Similarly, strong correlations were also found between Ba-Si and ^{226}Ra -Si
12 although no obvious process links ^{226}Ra or Ba with Si, except maybe for the adsorption of Ba and
13 (^{226}Ra) onto diatom frustules. It cannot be excluded, however, that the observed Ba-Si and ^{226}Ra -Si
14 relationships may result from the spatial coherence of different carriers overprinted by hydrodynamics.

15 Our study also provides evidence of significant decoupling between ^{226}Ra and Ba. In the upper
16 200 m, the $^{226}\text{Ra}/\text{Ba}$ ratios reach low values ($<2 \text{ dpm } \mu\text{mol}^{-1}$), a pattern that has been observed in other
17 regions and was related to acantharian skeletons that incorporate ^{226}Ra preferentially to Ba (van Beek et
18 al., 2007; Bernstein et al., 1998). Finally, deviations from the mean GEOSECS $^{226}\text{Ra}/\text{Ba}$ ratios were
19 observed in the shallow coastal waters of Greenland and Newfoundland: the predominant input of Ba
20 due to input of meteoritic waters leads to lower $^{226}\text{Ra}/\text{Ba}$ ratios whereas near the seafloor, the input of
21 sedimentary ^{226}Ra leads to higher $^{226}\text{Ra}/\text{Ba}$ ratios.

22 The absence of a stable isotope for radium led geochemists to consider Ba as a stable analog for
23 ^{226}Ra because ^{226}Ra and Ba display a similar chemical behavior, with the aim of using the $^{226}\text{Ra}/\text{Ba}$ ratio
24 as a chronometer for the thermohaline circulation. This study confirms that ^{226}Ra and Ba behave
25 similarly in the ocean interior away from external sources, both elements being predominantly
26 conservative in the studied area over distances of the order of a few thousands of km. However, this
27 study also highlights regions where ^{226}Ra and Ba deviate from a conservative behavior, an important
28 consideration when considering the balance between the large-scale oceanic circulation and biological
29 activity over long time scales. Decoupling between ^{226}Ra and Ba has been observed, in most cases at the
30 ocean boundaries as the result of dissolved ^{226}Ra and Ba external sources. In addition, suspended
31 particle dissolution may differently impact the dissolved ^{226}Ra and Ba content of intermediate and deep
32 waters (as shown for the NEADWL); such process would therefore potentially modify their $^{226}\text{Ra}/\text{Ba}$
33 ratios and would complicate the use of this ratio as a chronometer. Inclusion of the different sources and
34 sinks and particle/dissolved interactions in global ocean models should help to refine the use of the

1 $^{226}\text{Ra}/\text{Ba}$ ratio as a clock to chronometer the thermohaline circulation, as was proposed several decades
2 ago during the GEOSECS program.
3

4 Figure Caption

- 5 **Figure 1:** Station locations of the GA01 section between Portugal and Newfoundland in the North Atlantic (black and blue dots). Stations
6 investigated for ^{226}Ra and Ba are marked as blue dots. The main currents and water masses in the North Atlantic are also represented. The
7 major hydrothermal vents located near the GA01 section are indicated by black triangles. Stations investigated during the US-
8 GEOTRACES-GA03 section, also conducted in the Atlantic Ocean, are reported on the lower panel (red dots).
- 9 **Figure 2:** Potential temperature-salinity diagram—including a zoom for bottom waters—of the water samples (colored dots) from the
10 GA01 section. The properties of the source water types (based on García-Ibáñez et al., 2017; This issue) used in the Optimum
11 Multiparameter (OMP) analysis are reported with white circles. Isopycnals are also plotted (potential density referenced to 0 dbar).
- 12 **Figure 3:** Distribution of salinity (CTD data) along the GA01 section. The different water masses are also reported, following García-
13 Ibáñez et al. (2017; this issue). The station numbers are found on top of the figure. The sampling depths for ^{226}Ra are shown for each
14 vertical profile (black dots).
- 15 **Figure 4:** Distribution of (a) dissolved ^{226}Ra activities ($\text{dpm } 100 \text{ L}^{-1}$), (b) dissolved Ba concentrations (nmol L^{-1}) and $^{226}\text{Ra}/\text{Ba}$ ratio
16 ($\text{dpm } \mu\text{mol}^{-1}$) along the GA01 section. Station numbers are found on top of the panels. The sampling depths are shown for each vertical
17 profile (black dots).
- 18 **Figure 5:** Relationships between ^{226}Ra and Ba (red dots) and between ^{226}Ra and $\text{Si}(\text{OH})_4$ (blue dots) along the GA01 section in the North
19 Atlantic. The best linear fit for the two plots is also reported ($R=0.93$ for the two plots). The slopes of the relationships between ^{226}Ra -Ba
20 and between ^{226}Ra - $\text{Si}(\text{OH})_4$ are expressed in $10^{-2} \text{ dpm nmol}^{-1}$ and in $10^{-2} \text{ dpm } \mu\text{mol}^{-1}$, respectively.
- 21 **Figure 6:** Difference between the measured concentrations and those calculated by the OMP analysis, for ^{226}Ra (a), Ba (b) and (c)
22 $^{226}\text{Ra}/\text{Ba}$ ratio along the GA01 section. Positive anomalies reflect recent tracer addition, while negative ones reflect recent tracer removal.
23 Station numbers are found on top of the panels.
- 24 **Figure 7:** Vertical profiles of dissolved ^{226}Ra activities and Ba concentrations determined along the GA01 section: (a) West European
25 Basin; (b) Iceland Basin and Irminger Basin, (c) the Greenland and Newfoundland margins, and (d) Labrador Basin. As a comparison, the
26 conservative ^{226}Ra and Ba vertical profiles derived from the OMP analysis are also reported in solid grey lines. The discrepancy between
27 the two vertical profiles indicates deviation from the conservative behavior and reflects either an input of ^{226}Ra or Ba (positive anomalies
28 highlighted in red; same color code as Fig.6) or a removal of ^{226}Ra or Ba (negative anomalies highlighted in blue; same color code as
29 Fig.6). The OMP analysis has not been solved for the shallow coastal stations 53 and 78. The $^{226}\text{Ra}/\text{Ba}$ ratios are also reported, together
30 with the mean GEOSECS $^{226}\text{Ra}/\text{Ba}$ ratio ($2.2 \pm 0.2 \text{ dpm } \mu\text{mol}^{-1}$; black dashed line) together with its one standard deviation (grey shaded
31 areas). Note that the scale may be different from one station to the other and the vertical axis was cut to 1000 m. The seafloor is
32 represented by the bottom axis.
- 33 **Figure 8:** Geographical variation of ^{226}Ra activities (red dots), salinity (blue dots), temperature (yellow dots) and $\text{Si}(\text{OH})_4$ concentrations
34 (green dots) in AABW and the NEADWL between 60°S and 45°N (GA01 section) in the Atlantic Ocean based on data from the
35 GEOSECS and TTO programs. The ^{226}Ra activities, salinity, temperature and $\text{Si}(\text{OH})_4$ concentrations from GA01 are represented with
36 open circles. The values used as endmembers for the OMP analysis are also identified by the black circles. The shaded area represents the
37 region where transformation of the AABW into NEADWL takes place.
- 38 **Figure 9:** ^{226}Ra fluxes diffusing out of the sediment in relationship with bottom water ^{226}Ra activities determined in different oceanic
39 basins (P=Pacific Ocean, A=Atlantic Ocean, I=Indian Ocean and AA=Southern Ocean) by Cochran (1980). The ^{226}Ra flux calculated in
40 this study to explain the positive anomalies in the West European Basin is also reported (red dot).
- 41 **Figure 10:** ^{226}Ra and Ba flux estimations: $F_{\text{Part}-x}$ is the particulate flux entering the box and $F_{\text{Accumulation}-x}$ is the sediment accumulation flux
42 and $F_{\text{Dissolution}-x}$ is the flux of particle dissolution assuming that all the settling particles dissolve. x is either ^{226}Ra or Ba. Both maximum and
43 minimum values are shown for $F_{\text{Part}-x}$ and $F_{\text{Accumulation}-x}$. $\text{Max } F_{\text{Dissolution}-x}$ represents a maximum value since it is calculated by subtracting
44 the minimum value of $F_{\text{Accumulation}-x}$ from the maximum value of $F_{\text{Part}-x}$. $F_{\text{Dissolution}-x}$ is thus comprised between zero (if all $F_{\text{part}-x}$ accumulates
45 in the sediment) and this latter value.
- 46 **Figure S1:** Comparison of the vertical profiles of dissolved ^{226}Ra at stations 1 and 13 of the GA01 section (black and red dots,
47 respectively) and station 1 of the GA03 section (U.S.-GEOTRACES; blue dots) off Portugal.
- 48 **Figure S2:** Vertical profiles of dissolved ^{226}Ra activities and dissolved Ba concentrations with the conservative ^{226}Ra and Ba vertical
49 profiles derived from the OMP analysis, $^{226}\text{Ra}/\text{Ba}$ ratios, $\text{Si}(\text{OH})_4$ concentrations, salinity (black line) and potential temperature (red line)
50 for (a) the Iberian margin and the West European Basin, (b) the Iceland Basin and the Irminger Sea, (c) the Greenland margin, and (d) the

- 1 Labrador Sea and the Newfoundland margin. Note that the scale may be different from one station to the other and the vertical axis was cut
2 to 1000 m. The bottom is represented by the bottom axis.
- 3 **Figure S3:** Location of each endmember source water types (SWTs) used for the OMP analysis (black circles). The surface of the basin,
4 S , used to calculate the fluxes is represented by the grey hatched area.
- 5
- 6 **Figure S4:** Satellite Chlorophyll-a concentrations (MODIS Aqua from <http://oceancolor.gsfc.nasa.gov>), in mg m^{-3} during the GA01 cruise
7 in (a) May 2014 and (b) June 2014. The dashed line indicates the location of the GA01 section. Stations investigated in this work are
8 indicated by dots. White dots indicate the stations investigated during the corresponding month.
- 9 **Figure S5:** Schematic box model used to calculate the input fluxes in the West European Basin: F_{Sed-x} is the flux diffusing out of bottom
10 sediments, F_{part-x} is the vertical flux of particles entering the box from above, $F_{Accumulation-x}$ is the flux of particles accumulating in the
11 sediment and F_{H-In-x} and $F_{H-Out-x}$ represent horizontal fluxes of dissolved species or particles coming in and out of the box due to transport,
12 respectively. x is either ^{226}Ra or Ba.
- 13 **Table S1:** Characteristics and location of each endmember source water types (SWTs).
- 14 **Table S2:** ^{226}Ra activities, Ba concentrations, $^{226}\text{Ra}/\text{Ba}$ ratios, potential temperature and salinity at the different stations of the GA01
15 section.
- 16

1 Acknowledgement

2 The present research and Emilie Le Roy's fellowship are co-funded by the European Union and the
3 Région Occitanie-Pyrénées-Méditerranée (European Regional Development Fund). We are grateful to
4 the captain and crew of the N/O *Pourquoi Pas?*. The GEOVIDE project is co-funded by the French
5 national program LEFE/INSU (GEOVIDE), ANR Blanc (GEOVIDE, ANR-13-BS06-0014) and
6 RPDO (ANR-12-PDOC-0025-01), LabEX MER (ANR-10-LABX-19) and IFREMER. The
7 GEOVIDE cruise would not have been achieved without the technical skills and commitment of
8 Catherine Kermabon, Olivier Ménage, Stéphane Leizour, Michel Hamon, Philippe Le Bot, Emmanuel
9 de Saint-Léger and Fabien Pérault. We are grateful to Manon Le Goff, Emilie Grosstefan,
10 Morgane Gallinari and Paul Tréguer for Si(OH)₄ sampling and analysis. This work was also co-funded
11 by the French national program LEFE/INSU "REPAP" (PI S. Jacquet) and the U.S. National Science
12 Foundation (PI M. Charette, OCE- 1458305; OCE- 1232669). For this work M.I. García-Ibáñez and
13 F.F. Pérez were supported by the Spanish Ministry of Economy and Competitiveness through the
14 BOCATS (CTM2013-41048-P) project co-funded by the Fondo Europeo de Desarrollo Regional 2014–
15 2020 (FEDER). Several figures were constructed using Ocean Data View (Schlitzer, 2003). Therefore,
16 R. Schlitzer is warmly thanked. Satellite chlorophyll-a visualizations used in this study were produced
17 with the Giovanni online data system, developed and maintained by the NASA GES DISC.

18 References

- 19
- 20 Antia, A. N., Bauerfeind, E., Bodungen, B. v and Zeller, U.: Abundance, encystment and sedimentation
21 of acantharia during autumn 1990 in the East Greenland Sea, J. Plankton Res., 15(1), 99–114,
22 doi:10.1093/plankt/15.1.99, 1993.
- 23 Bacon, M. P. and Anderson, R. F.: Distribution of thorium isotopes between dissolved and particulate
24 forms in the deep sea, J. Geophys. Res. Oceans, 87(C3), 2045–2056, doi:10.1029/JC087iC03p02045,
25 1982.
- 26 Barnard, R. T., Batten, S., Beaugrand, G., Buckland, C., Conway, D. V. P., Edwards, M., Finlayson, J.,
27 Gregory, L. W., Halliday, N. C., John, A. W. G., Johns, D. G., Johnson, A. D., Jonas, T. D., Lindley, J.
28 A., Nyman, J., Pritchard, P., Reid, P. C., Richardson, A. J., Saxby, R. E., Sidey, J., Smith, M. A.,
29 Stevens, D. P., Taylor, C. M., Tranter, P. R. G., Walne, A. W., Wootton, M., Wotton, C. O. M. and
30 Wright, J. C.: Continuous plankton records: Plankton atlas of the North Atlantic Ocean (1958-1999). II.
31 Biogeographical charts, Mar. Ecol.-Prog. Ser., MEPS supplement(SUPPL.), 11–75, 2004.
- 32 Bennett, T., Broecker, W. S. and Hansen, J.: North Atlantic Deep Water Formation, 1985.
- 33 Bernstein, R. E., Byrne, R. H. and Schijf, J.: Acantharians: a missing link in the oceanic
34 biogeochemistry of barium, Deep Sea Res. Part Oceanogr. Res. Pap., 45(2), 491–505,
35 doi:10.1016/S0967-0637(97)00095-2, 1998.
- 36 Bishop, J. K. B.: The barite-opal-organic carbon association in oceanic particulate matter, Nature,
37 332(6162), 341–343, doi:10.1038/332341a0, 1988.
- 38 Broecker, W., Kaufman, A., Ku, T.-L., Chung, Y.-C. and Craig, H.: Radium 226 measurements from
39 the 1969 North Pacific Geosecs Station, J. Geophys. Res., 75(36), 7682–7685,
40 doi:10.1029/JC075i036p07682, 1970.

- 1 Broecker, W. S., Li, Y.-H. and Cromwell, J.: Radium-226 and Radon-222 : Concentration in Atlantic
2 and Pacific Oceans, *Science*, 158(3806), 1307–1310, 1967.
- 3 Broecker, W. S., Goddard, J. and Sarmiento, J. L.: The distribution of ^{226}Ra in the Atlantic Ocean,
4 *Earth Planet. Sci. Lett.*, 32(2), 220–235, doi:10.1016/0012-821X(76)90063-7, 1976.
- 5 Chan, L. H., Edmond, J. M., Stallard, R. F., Broecker, W. S., Chung, Y. C., Weiss, R. F. and Ku, T. L.:
6 Radium and barium at GEOSECS stations in the Atlantic and Pacific, *Earth Planet. Sci. Lett.*, 32(2),
7 258–267, 1976.
- 8 Chan, L. H., Drummond, D., Edmond, J. M. and Grant, B.: On the barium data from the Atlantic
9 GEOSECS expedition, *Deep Sea Res.*, 24(7), 613–649, doi:10.1016/0146-6291(77)90505-7, 1977.
- 10 Charette, M. A., Morris, P. J., Henderson, P. B. and Moore, W. S.: Radium isotope distributions during
11 the US GEOTRACES North Atlantic cruises, *Mar. Chem.*, 177, Part 1, 184–195,
12 doi:10.1016/j.marchem.2015.01.001, 2015.
- 13 Chow, T. J. and Goldberg, E. D.: On the marine geochemistry of barium, *Geochim. Cosmochim. Acta*,
14 20(3), 192–198, doi:10.1016/0016-7037(60)90073-9, 1960.
- 15 Chung, Y.: Radium-barium-silica correlations and a two-dimensional radium model for the world
16 ocean, *Earth Planet. Sci. Lett.*, 49(2), 309–318, doi:10.1016/0012-821X(80)90074-6, 1980.
- 17 Chung, Y., Craig, H., Ku, T. L., Goddard, J. and Broecker, W. S.: Radium-226 measurements from
18 three Geosecs intercalibration stations, *Earth Planet. Sci. Lett.*, 23(1), 116–124, doi:10.1016/0012-
19 821X(74)90038-7, 1974.
- 20 Chung, Y.-C.: A deep ^{226}Ra maximum in the Northeast Pacific, *Earth Planet. Sci. Lett.*, 32, 249–257,
21 doi:10.1016/0012-821X(76)90065-0, 1976.
- 22 Chung, Y.-C. and Craig, H.: ^{226}Ra in the Pacific Ocean, *Earth Planet. Sci. Lett.*, 49, 267–292, 1980.
- 23 Cochran, J. K.: The flux of ^{226}Ra from deep-sea sediments, *Earth Planet. Sci. Lett.*, 49(2), 381–392,
24 doi:10.1016/0012-821X(80)90080-1, 1980.
- 25 Cochran, J. K. and Krishnaswami, S.: Radium, thorium, uranium, and ^{210}Pb in deep-sea sediments and
26 sediment pore waters from the North Equatorial Pacific, *Am. J. Sci.*, 280, 849–889,
27 doi:10.2475/ajs.280.9.849, 1980.
- 28 Danialt, N., Lherminier, P. and Mercier, H.: Circulation and Transport at the Southeast Tip of
29 Greenland, *J. Phys. Oceanogr.*, 41(3), 437–457, doi:10.1175/2010JPO4428.1, 2011.
- 30 Danialt, N., Mercier, H., Lherminier, P., Sarafanov, A., Falina, A., Zunino, P., Pérez, F. F., Ríos, A. F.,
31 Ferron, B., Huck, T., Thierry, V. and Gladyshev, S.: The northern North Atlantic Ocean mean
32 circulation in the early 21st century, *Prog. Oceanogr.*, 146, 142–158, doi:10.1016/j.pocean.2016.06.007,
33 2016.
- 34 Dehairs, F., Chesselet, R. and Jedwab, J.: Discrete suspended particles of barite and the barium cycle in
35 the open ocean, *Earth Planet. Sci. Lett.*, 49(2), 528–550, doi:10.1016/0012-821X(80)90094-1, 1980.
- 36 Deng, F., Thomas, A. L., Rijkenberg, M. J. A. and Henderson, G. M.: Controls on seawater ^{231}Pa ,
37 ^{230}Th and ^{232}Th concentrations along the flow paths of deep waters in the Southwest Atlantic, *Earth*
38 *Planet. Sci. Lett.*, 390, 93–102, doi:10.1016/j.epsl.2013.12.038, 2014.

- 1 Dileep Kumar, M. and Li, Y.-H.: Spreading of water masses and regeneration of silica and ²²⁶Ra in the
2 Indian Ocean, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 43(1), 83–110, doi:10.1016/0967-
3 0645(95)00084-4, 1996.
- 4 Dutkiewicz, A., Müller, R. D., O’Callaghan, S. and Jónasson, H.: Census of seafloor sediments in the
5 world’s ocean, *Geology*, 43(9), 795–798, doi:10.1130/G36883.1, 2015.
- 6 Edmond, J. M., Measures, C., McDuff, R. E., Chan, L. H., Collier, R., Grant, B., Gordon, L. I. and
7 Corliss, J. B.: Ridge crest hydrothermal activity and the balances of the major and minor elements in the
8 ocean: The Galapagos data, *Earth Planet. Sci. Lett.*, 46(1), 1–18, doi:10.1016/0012-821X(79)90061-X,
9 1979.
- 10 Falina, A., Sarafanov, A., Mercier, H., Lherminier, P., Sokov, A. and Daniault, N.: On the Cascading of
11 Dense Shelf Waters in the Irminger Sea, *J. Phys. Oceanogr.*, 42(12), 2254–2267, doi:10.1175/JPO-D-
12 12-012.1, 2012.
- 13 Falkner, K. K., klinkhammer, G. P., Bowers, T. S., Todd, J. F., Lewis, B. L., Landing, W. M. and
14 Edmond, J. M.: The behavior of barium in anoxic marine waters, *Geochim. Cosmochim. Acta*, 57(3),
15 537–554, doi:10.1016/0016-7037(93)90366-5, 1993.
- 16 Fanning, K. A., Vargo, G. A., Bell-Torres, L. and Young, R. W.: Covariation of reactive solutes in the
17 sea, *Mar. Chem.*, 24(3-4), 215–238, 1988.
- 18 Foster, D. A., Staubwasser, M. and Henderson, G. M.: ²²⁶Ra and Ba concentrations in the Ross Sea
19 measured with multicollector ICP mass spectrometry, *Mar. Chem.*, 87(1-2), 59–71, 2004.
- 20 Freydier, R., Dupre, B. and Polve, M.: Analyses by inductively coupled plasma mass spectrometry of
21 Ba concentrations in water and rock samples. Comparison between isotope dilution and external
22 calibration with or without internal standard, *Eur. J. Mass Spectrom.*, 1(3), 283–291,
23 doi:10.1255/ejms.140, 1995.
- 24 Fröb, F., Olsen, A., Våge, K., Moore, G. W. K., Yashayaev, I., Jeansson, E. and Rajasakaren, B.:
25 Irminger Sea deep convection injects oxygen and anthropogenic carbon to the ocean interior, *Nat.*
26 *Commun.*, 7, ncomms13244, doi:10.1038/ncomms13244, 2016.
- 27 García-Ibáñez, M. I., Pardo, P. C., Carracedo, L. I., Mercier, H., Lherminier, P., Ríos, A. F. and Pérez,
28 F. F.: Structure, transports and transformations of the water masses in the Atlantic Subpolar Gyre, *Prog.*
29 *Oceanogr.*, 135, 18–36, doi:10.1016/j.pocean.2015.03.009, 2015.
- 30 Gingele, F. and Dahmke, A.: Discrete barite particles and barium as tracers of paleoproductivity in
31 south Atlantic sediments, *Paleoceanography*, 9(1), 151–168, doi:10.1029/93PA02559, 1994.
- 32 Henderson, P., Morris, P., Moore, W. and Charette, M.: Methodological advances for measuring low-
33 level radium isotopes in seawater, *J. Radioanal. Nucl. Chem.*, 296(1), 357–362, doi:10.1007/s10967-
34 012-2047-9, 2013.
- 35 Jacquet, S. H. M., Dehairs, F., Cardinal, D., Navez, J. and Delille, B.: Barium distribution across the
36 Southern Ocean frontal system in the Crozet–Kerguelen Basin, *Mar. Chem.*, 95(3–4), 149–162,
37 doi:10.1016/j.marchem.2004.09.002, 2005.
- 38 Jacquet, S. H. M., Dehairs, F., Elskens, M., Savoye, N. and Cardinal, D.: Barium cycling along WOCE
39 SR3 line in the Southern Ocean, *Mar. Chem.*, (1-2) [online] Available from:
40 <http://vliz.be/en/imis?module=ref&refid=211233&basketaction=add> (Accessed 20 January 2017), 2007.

- 1 Jeandel, C., Dupré, B., Lebaron, G., Monnin, C. and Minster, J.-F.: Longitudinal distributions of
2 dissolved barium, silica and alkalinity in the western and southern Indian Ocean, *Deep Sea Res. Part*
3 *Oceanogr. Res.*, 43, 1–31, doi:10.1016/0967-0637(95)00098-4, 1996.
- 4 Jenkins, W. J., Smethie Jr., W. M., Boyle, E. A. and Cutter, G. A.: Water mass analysis for the U.S.
5 GEOTRACES (GA03) North Atlantic sections, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 116, 6–20,
6 doi:10.1016/j.dsr2.2014.11.018, 2015.
- 7 Jullion, L., Jacquet, S. H. M. and Tanhua, T.: Untangling biogeochemical processes from the impact of
8 ocean circulation: First insight on the Mediterranean dissolved barium dynamics, *Glob. Biogeochem.*
9 *Cycles*, 2016GB005489, doi:10.1002/2016GB005489, 2017.
- 10 Kadko, D.: Radioisotopic studies of submarine hydrothermal vents, *Rev. Geophys.*, 34(3), 349–366,
11 doi:10.1029/96RG01762, 1996.
- 12 Kadko, D. and Moore, W.: Radiochemical constraints on the crustal residence time of submarine
13 hydrothermal fluids: Endeavour Ridge, *Geochim. Cosmochim. Acta*, 52(3), 659–668,
14 doi:10.1016/0016-7037(88)90328-6, 1988.
- 15 Key, R. M., Guinasso, N. L. and Schink, D. R.: Emanation of radon-222 from marine sediments, *Mar.*
16 *Chem.*, 7(3), 221–250, doi:10.1016/0304-4203(79)90041-0, 1979.
- 17 Kipp, L. E., Sanial, V., Henderson, P. B., van Beek, P., Reyss, J.-L., Hammond, D. E., Moore, W. S.
18 and Charette, M. A.: Radium isotopes as tracers of hydrothermal inputs and neutrally buoyant plume
19 dynamics in the deep ocean, *Mar. Chem.*, doi:10.1016/j.marchem.2017.06.011, 2017.
- 20 Klinkhammer, G. P. and Chan, L. H.: Determination of barium in marine waters by isotope dilution
21 inductively coupled plasma mass spectrometry, *Anal. Chim. Acta*, 232, doi:10.1016/s0003-
22 2670(00)81249-0, 1990.
- 23 Koczy, F. F.: Natural radium as a tracer in the ocean, , 18, 351–357, 1958.
- 24 Krauss, W.: Currents and mixing in the Irminger Sea and in the Iceland Basin, *J. Geophys. Res. Oceans*,
25 100(C6), 10851–10871, doi:10.1029/95JC00423, 1995.
- 26 Krishnaswami, S., Lal, D., Somayajulu, B. L. K., Weiss, R. F. and Craig, H.: Large-volume in-situ
27 filtration of deep Pacific waters: Mineralogical and radioisotope studies, *Earth Planet. Sci. Lett.*, 32(2),
28 420–429, doi:10.1016/0012-821X(76)90082-0, 1976.
- 29 Kröll, V.: Vertical Distribution of Radium in Deep-Sea Sediments, *Nature*, 171(4356), 742–742,
30 doi:10.1038/171742a0, 1953.
- 31 Ku, T.-L. and Lin, M.-C.: ²²⁶Ra distribution in the Antarctic Ocean, *Earth Planet. Sci. Lett.*, 32(2),
32 236–248, doi:10.1016/0012-821X(76)90064-9, 1976.
- 33 Ku, T.-L. and Luo, S.: New appraisal of radium 226 as a large-scale oceanic mixing tracer, *J. Geophys.*
34 *Res. Oceans*, 99(C5), 10255–10273, doi:10.1029/94JC00089, 1994.
- 35 Ku, T. L., Li, Y. H., Mathieu, G. G. and Wong, H. K.: Radium in the Indian-Antarctic Ocean south of
36 Australia, *J. Geophys. Res.*, 75(27), 5286–5292, doi:10.1029/JC075i027p05286, 1970.

- 1 Ku, T. L., Huh, C. A. and Chen, P. S.: Meridional distribution of ^{226}Ra in the eastern Pacific along
2 GEOSECS cruise tracks, *Earth Planet. Sci. Lett.*, 49, 293–308, doi:10.1016/0012-821X(80)90073-4,
3 1980.
- 4 Lacan, F. and Jeandel, C.: Denmark Strait water circulation traced by heterogeneity in neodymium
5 isotopic compositions, *Deep Sea Res. Part Oceanogr. Res. Pap.*, 51(1), 71–82, 2004.
- 6 Lazier, J. R. N.: The renewal of Labrador sea water, *Deep Sea Res. Oceanogr. Abstr.*, 20(4), 341–353,
7 doi:10.1016/0011-7471(73)90058-2, 1973.
- 8 Lea, D. and Boyle, E.: Barium content of benthic foraminifera controlled by bottom-water composition,
9 *Nature*, 338(6218), 751, doi:10.1038/338751a0, 1989.
- 10 Lea, D. W. and Boyle, E. A.: Foraminiferal reconstruction of barium distributions in water masses of
11 the glacial oceans, *Paleoceanography*, 5(5), 719–742, doi:10.1029/PA005i005p00719, 1990.
- 12 Legeleux, F. and Reyss, J.-L.: $^{228}\text{Ra}/^{226}\text{Ra}$ activity ratio in oceanic settling particles: implications
13 regarding the use of barium as a proxy for paleoproductivity reconstruction, *Deep Sea Res. Part*
14 *Oceanogr. Res. Pap.*, 43(11), 1857–1863, doi:10.1016/S0967-0637(96)00086-6, 1996.
- 15 Lherminier, P., Mercier, H., Huck, T., Gourcuff, C., Perez, F. F., Morin, P., Sarafanov, A. and Falina,
16 A.: The Atlantic Meridional Overturning Circulation and the subpolar gyre observed at the A25-OVIDE
17 section in June 2002 and 2004, *Deep Sea Res. Part Oceanogr. Res. Pap.*, 57(11), 1374–1391,
18 doi:10.1016/j.dsr.2010.07.009, 2010.
- 19 Li, Y. H., Ku, T. L., Mathieu, G. G. and Wolgemuth, K.: Barium in the Antarctic Ocean and
20 implications regarding the marine geochemistry of Ba and ^{226}Ra , *Earth Planet. Sci. Lett.*, 19(3), 352–
21 358, doi:10.1016/0012-821X(73)90085-X, 1973.
- 22 Lucas, H. F.: Improved Low-Level Alpha-Scintillation Counter for Radon, *Rev. Sci. Instrum.*, 28(9),
23 680, doi:10.1063/1.1715975, 1957.
- 24 Martin, J.-M. and Meybeck, M.: Elemental mass-balance of material carried by major world rivers,
25 *Mar. Chem.*, 7(3), 173–206, doi:10.1016/0304-4203(79)90039-2, 1979.
- 26 Martin, P., Allen, J. T., Cooper, M. J., Johns, D. G., Lampitt, R. S., Sanders, R. and Teagle, D. A. H.:
27 Sedimentation of acantharian cysts in the Iceland Basin: Strontium as a ballast for deep ocean particle
28 flux, and implications for acantharian reproductive strategies, *Limnol. Oceanogr.*, 55(2), 604–614,
29 doi:10.4319/lo.2010.55.2.0604, 2010.
- 30 Mathieu, G. G. and Wolgemuth, K.: Barium in the Antarctic Ocean and Implication regarding the marine
31 geochemistry of Ba and ^{226}Ra , *EARTH Planet. Sci. Lett.*, 19, 352–358, 1973.
- 32 McCartney, M. S.: Recirculating components to the deep boundary current of the northern North
33 Atlantic, *Prog. Oceanogr.*, 29, 283–383, 1992.
- 34 McCave, I. N.: Local and global aspects of the bottom nepheloid layers in the world ocean, *Neth. J. Sea*
35 *Res.*, 20(2), 167–181, doi:10.1016/0077-7579(86)90040-2, 1986.
- 36 McManus, J., Berelson, W. M., Hammond, D. E. and Klinkhammer, G. P.: Barium Cycling in the North
37 Pacific: Implications for the Utility of Ba as a Paleoproductivity and Paleoalkalinity Proxy,
38 *Paleoceanography*, 14(1), 53–61, doi:10.1029/1998PA900007, 1999.

- 1 Monnin, C., Jeandel, C., Cattaldo, T. and Dehairs, F.: The marine barite saturation state of the world's
2 oceans, *Mar. Chem.*, 65(3), 253–261, doi:10.1016/S0304-4203(99)00016-X, 1999.
- 3 Moore, W. S. and Reid, D. F.: Extraction of radium from natural waters using manganese-impregnated
4 acrylic fibers, *J. Geophys. Res.*, 78(36), 8880–8886, doi:10.1029/JC078i036p08880, 1973.
- 5 Nozaki, Y.: Excess ^{227}Ac in deep ocean water, *Nature*, 310(5977), 486–488, doi:10.1038/310486a0,
6 1984.
- 7 Okubo, A., Obata, H., Gamo, T. and Yamada, M.: ^{230}Th and ^{232}Th distributions in mid-latitudes of
8 the North Pacific Ocean: Effect of bottom scavenging, *Earth Planet. Sci. Lett.*, 339–340, 139–150,
9 doi:10.1016/j.epsl.2012.05.012, 2012.
- 10 Östlund, H. G., Craig, H. C., Broecker, W. S., Spencer, D. W. and GEOSECS: Shorebased
11 measurements during the GEOSECS Atlantic expedition, , doi:10.1594/PANGAEA.824123, 1987.
- 12 Paytan, A. and Kastner, M.: Benthic Ba fluxes in the central Equatorial Pacific, implications for the
13 oceanic Ba cycle, *Earth Planet. Sci. Lett.*, 142(3), 439–450, doi:10.1016/0012-821X(96)00120-3, 1996.
- 14 Peterson, R. N., Burnett, W. C., Dimova, N. and Santos, I. R.: Comparison of measurement methods for
15 radium-226 on manganese-fiber, *Limnol. Oceanogr. Methods*, 7(2), 196–205,
16 doi:10.4319/lom.2009.7.196, 2009.
- 17 Pickart, R. S. and Spall, M. A.: Impact of Labrador Sea Convection on the North Atlantic Meridional
18 Overturning Circulation, *J. Phys. Oceanogr.*, 37(9), 2207–2227, doi:10.1175/JPO3178.1, 2007.
- 19 Pickart, R. S., Straneo, F. and Moore, G. W. K.: Is Labrador sea water formed in the Irminger basin?,
20 *Deep Sea Res. Part Oceanogr. Res. Pap.*, 50(1), 23–52, 2003.
- 21 Read, J. F.: CONVEX-91: water masses and circulation of the Northeast Atlantic subpolar gyre, *Prog.*
22 *Oceanogr.*, 48(4), 461–510, doi:10.1016/S0079-6611(01)00011-8, 2000.
- 23 Roy-Barman, M., Coppola, L. and Souhaut, M.: Thorium isotopes in the western Mediterranean Sea: an
24 insight into the marine particle dynamics, *Earth Planet. Sci. Lett.*, 196(3), 161–174, doi:10.1016/S0012-
25 821X(01)00606-9, 2002.
- 26 Rubin, S. I., King, S. L., Jahnke, R. A. and Froelich, P. N.: Benthic barium and alkalinity fluxes: Is Ba
27 an oceanic paleo-alkalinity proxy for glacial atmospheric CO_2 ?, *Geophys. Res. Lett.*, 30(17), 1885,
28 doi:10.1029/2003GL017339, 2003.
- 29 Rudels, B., Fahrbach, E., Meincke, J., Budéus, G. and Eriksson, P.: The East Greenland Current and its
30 contribution to the Denmark Strait overflow, *ICES J. Mar. Sci.*, 59(6), 1133–1154,
31 doi:10.1006/jmsc.2002.1284, 2002.
- 32 Rudnicki, M. D. and Elderfield, H.: Helium, radon and manganese at the TAG and Snakepit
33 hydrothermal vent fields, 26° and 23°N , Mid-Atlantic Ridge, *Earth Planet. Sci. Lett.*, 113(3), 307–321,
34 doi:10.1016/0012-821X(92)90136-J, 1992.
- 35 Schmidt, S. and Reyss, J.-L.: Radium as internal tracer of Mediterranean Outflow Water, *J. Geophys.*
36 *Res. Oceans*, 101(C2), 3589–3596, doi:10.1029/95JC03308, 1996.
- 37 Schmitz, W. J. and McCartney, M. S.: On the North Atlantic Circulation, *Rev. Geophys.*, 31(1), 29–49,
38 doi:10.1029/92RG02583, 1993.

- 1 Seager, R., Battisti, D. S., Yin, J., Gordon, N., Naik, N., Clement, A. C. and Cane, M. A.: Is the Gulf
2 Stream responsible for Europe's mild winters?, *Q. J. R. Meteorol. Soc.*, 128(586), 2563–2586,
3 doi:10.1256/qj.01.128, 2002.
- 4 Shannon, L. V. and Cherry, R. D.: Radium-226 in marine phytoplankton, *Earth Planet. Sci. Lett.*, 11(1),
5 339–343, doi:10.1016/0012-821X(71)90189-0, 1971.
- 6 Spencer, D. W.: Geosecs II, the 1970 North Atlantic station: Hydrographic features, oxygen, and
7 nutrients, *Earth Planet. Sci. Lett.*, 16(1), 91–102, doi:10.1016/0012-821X(72)90241-5, 1972.
- 8 Szabo, B. J.: Radium content in plankton and sea water in the Bahamas, *Geochim. Cosmochim. Acta*,
9 31(8), 1321–1331, doi:10.1016/S0016-7037(67)80018-8, 1967.
- 10 van Aken, H. M.: The hydrography of the mid-latitude northeast Atlantic Ocean. I: The deep water
11 masses, *Deep-Sea Res. Part I*, 47, 757–788, doi:10.1016/S0967-0637(99)00092-8, 2000.
- 12 van Aken, H. M. and Becker, G.: Hydrography and through-flow in the north-eastern North Atlantic
13 Ocean: the NANSEN project, *Prog. Oceanogr.*, 38(4), 297–346, doi:10.1016/S0079-6611(97)00005-0,
14 1996.
- 15 van Beek, P., François, R., Conte, M., Reyss, J.-L., Souhaut, M. and Charette, M.: 228Ra/226Ra and
16 226Ra/Ba ratios to track barite formation and transport in the water column, *Geochim. Cosmochim.*
17 *Acta*, 71(1), 71–86, doi:10.1016/j.gca.2006.07.041, 2007.
- 18 van Beek, P., Sternberg, E., Reyss, J. L., Souhaut, M., Robin, E. and Jeandel, C.: 228Ra/226Ra and
19 226Ra/Ba ratios in the Western Mediterranean Sea : barite formation and transport in the water column,
20 *Geochim. Cosmochim. Acta*, 73, 4720–4737, doi:10.1016/j.gca.2009.05.063, 2009.
- 21 van Beek, P., Souhaut, M. and Reyss, J.-L.: Measuring the radium quartet (228Ra, 226Ra, 224Ra,
22 223Ra) in seawater samples using gamma spectrometry, *J. Environ. Radioact.*, 101(7), 521–529,
23 doi:10.1016/j.jenvrad.2009.12.002, 2010.
- 24 Wolgemuth, K. and Broecker, W. S.: Barium in sea water, *Earth Planet. Sci. Lett.*, 8(5), 372–378,
25 doi:10.1016/0012-821X(70)90110-X, 1970.
- 26 Yashayaev, I. and Dickson, B.: Transformation and Fate of Overflows in the Northern North Atlantic,
27 in *Arctic–Subarctic Ocean Fluxes*, edited by R. R. Dickson, J. Meincke, and P. Rhines, pp. 505–526,
28 Springer Netherlands. [online] Available from: [http://link.springer.com/chapter/10.1007/978-1-4020-](http://link.springer.com/chapter/10.1007/978-1-4020-6774-7_22)
29 [6774-7_22](http://link.springer.com/chapter/10.1007/978-1-4020-6774-7_22) (Accessed 11 April 2016), 2008.
- 30 Yashayaev, I. and Loder, J. W.: Recurrent replenishment of Labrador Sea Water and associated
31 decadal-scale variability, *J. Geophys. Res. Oceans*, 121(11), 8095–8114, doi:10.1002/2016JC012046,
32 2016.
- 33 Yashayaev, I., Bersch, M. and van Aken, H. M.: Spreading of the Labrador Sea Water to the Irminger
34 and Iceland basins, *Geophys. Res. Lett.*, 34(10), doi:10.1029/2006GL028999, 2007.
- 35
- 36
- 37

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6





Figure 7

A) West European Basin



B) Iceland and Irminger Basins






Figure 7

C) Greenland and Newfoundland Margins


Station 53




Station 61



Station 57




Station 78




D) Labrador Basin


Station 60



Station 69



Station 64



Station 77




Figure 8




Figure 9




Figure 10

