

Microbial Biobanking

Cyanobacteria-rich topsoil facilitates mine rehabilitation

Wendy Williams¹, Angela Chilton², Mel Schneemilch¹, Stephen Williams¹, Brett Neilan³, Colin Driscoll⁴

1. School of Agriculture and Food Sciences, The University of Queensland, Gatton Campus 4343 Australia
2. Australian Centre for Astrobiology and School of Biotechnology and Biomolecular Sciences, University of New South Wales, Sydney, NSW, 2052, Australia
3. School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW, 2308, Australia
4. Hunter Eco, PO Box 1047, Toronto, NSW, 2283

Correspondence to: wendy.williams@uq.edu.au

Dear Associate Editor,

Thank you for your time and effort to provide valuable comments to improve this manuscript. We trust we have addressed them all satisfactorily.

Kind regards,

Wendy Williams and co-authors

Editor: requested a revision of the introduction to reflect the following themes (refer to main comments)

- a. **Problem**
- b. **Solutions**
- c. **Properties and Ecosystem services**
- d. **Post-mine rehab recommendations**
- e. **Goals and hypothesis**

Response: The introduction has been recast to reflect the order recommended

The following issues raised throughout the ms have been addressed and detailed below:

Editor: Soil and biocrust type are not considered together in analysis...why?

The editor has enquired if there was a reason why we haven't included biocrust type with soil type in our analysis. The first point to make is that the three studies presented here (cyanobacterial community structure and function, sequencing and stockpiles) were all independent studies that were linked in the common goal but done separately. In addition, the scope of this research was focused on cyanobacteria that we have collectively called biocrusts however preliminary investigations defined these biocrusts into different successional stages (similar to Budel et al. 2009). Although these were identified across the three soil management units (SMU) aside from SMU1 they were reasonably homogenously distributed across the landscape. A characteristic of these biocrusts (SMU2 and SMU3) was that they contained a large number of lichens and

some mosses. However, due to high-level disturbance associated with the mining process it was acknowledged that cyanobacteria would be the early colonisers and lichens would be unlikely to survive. For this reason, J-A mine requested a focused study of cyanobacteria that underlined their role in terms of succession (i.e. early colonisers, soil stabilisers), nutrient cycling and their survival in topsoil stockpiles.

5 Furthermore, in relation to the sequenced biocrusts for bare, early and late successional stages, these were samples located at one site only not for each sample location. Therefore, we suggest that we could not use this data to apply to the ten sites data.

Methods

P4, L16-19 All these points should be clearly discussed (I miss some discussion about goals a) and all these questions need to be solved in the conclusions

10 Response: We have recast the aims and hypotheses to clearly reflect the goals of this research and further addressed them directly in the discussion.

P5, L14 These are not already defined, please define them

Response: We have recast this paragraph to clarify the sampling strategy, vegetation and locations within the SMU with details shown in Table S1 and S2 and Figures 1 and 2.

15 “At J-A the landscape has been characterised into three distinct soil types that were associated with vegetation communities identified as soil management units (Doudle et al., 2011; Hou and Warland, 2005). Site vegetation associations are described as follows: SMU 1 – Red Mallee: *Eucalyptus oleosa* ssp. *oleosa* = open Mallee/Myall woodland; SMU 2 – Chenopod Shrubland: *Maireana sedifolia* and *Atriplex vesicaria*; SMU 3 – Western Myall: *Acacia papyrocarpa* *Maireana sedifolia* = open Myall woodland Site 1 occurs in a transition between SMU 2 and
20 SMU 3 but was treated as most like SMU 2 (also see Table S1). Soil management units were summarised as: SMU 1 – deep calcareous yellow sands associated with dune ridges; SMU 2 – shallow calcareous sandy loams; and SMU 3 – deep calcareous sandy loam (Table S1). In the first place, sample site locations (Fig. 1) were selected based on these soil management units (SMU) and a two-year old stockpile (Table S1). Secondly, sites within these parameters (SMU) were selected for the subsequent detailed studies of cyanobacterial succession
25 and its resilience to longer-term stockpiling (Fig.2 and Section 2.4).

The biocrusts at J-A had been previously classified into three successional stages representative of the five biocrust types found growing across the landscape (Table S1, S2) (Doudle et al., 2011). Types 1–2 are light coloured, patchy, thin, and fragile cyanobacterial crusts corresponding to early stages of development; Type 3 are well established cyanobacterial crusts with establishment of some mosses and lichens corresponding to intermediate stages of development; Types 4–5 biocrusts
30 are well established with cyanolichens and/or green algal lichens and mosses corresponding with late stages of development (additional descriptions available in supplementary Table S1). As well as the three main SMU sites we also sampled a dry salt lake (Lake Ifould) to provide information on cyanobacterial species adapted to saline conditions, similar in nature to ground water used in mine operations. In this study the term biocrust covers whole crust samples that incorporated lichens

cyanobacteria and mosses in varying proportions however, the cyanobacterial component of this crust was the focus in terms of our polyphasic approach to community structure, succession and its biophysiochemical properties.”

P5, Section 2.2 As you have two factors (SMU and crust type, a two-way anova is necessary?

Response: Refer to detailed explanation above. We did not use crust type as a factor as it was not specifically identified when sampling (for this project). We used prior classifications to provide an informative backdrop for decision making in relation to the sampling strategy employed in this study.

P5, L30 please define these sites

Response: Refer to above revision and additional descriptions now included

P6, Section 2.2.2 statistical analysis for this section are still missing

Response: statistical methods added

P6, L6 half petri dish or less? Please clarify

Response: Clarified in text

P6, L11-14 was it done before other analysis (before disturbance), or was done based on a subsample, please clarify

Response: Clarified in text

P6, L15-19 was it done before other analysis (before disturbance)? or in different petri dishes

Response: Clarified in text

P6, L21 I will suggest to include that it was done on the field at this point. When it was measured? during the field sampling campaign or in a different date.

P6, L31 before or after other measurements?

Response: Section 2.2.2 has been revised to clarify the above-mentioned points

P8, L28 (last sentence) differences in what?

Response: this sentence updated to what differences

Results

P9, Section 3.1 Here you only analysed differences between SMU, what about different biocrust types, as some different biocrust types in same SMU may have larger diff than similar crust types at different SMU, it is necessary to include both factors in your analysis

Response: Please see earlier explanation regarding biocrust types vs SMU

P9, L6 significant or no?

These details are now included in text and in the table

P9, L10 (also Figure 3) see my comment on figure 3, in a similar way as I suggested on table 3, significant differences should be added in all figures and tables where you performed an analysis of differences between classes. This can be done by using superscripts

e.g. a for the group with higher values, b, c, etc

When to classes do not show significant differences, then they have same superscript

letter should be added in the same way I suggested on table 2

Response: done

Table 2 this can be solved by adding letters a, b and c for the groups with significant differences as super index

5 SMU3 a SMU2 a, as no significant differences between them are found, SMU1 b (significantly lower values than SMU 2 and 3)

Response: the letters have been added where required

Table 4 define 'Permdisp'

Response: The term Permdisp is defined in the methods where it is first mentioned.

10 We have included Permdisp definition in legend in Table 4

15

20

Microbial Biobanking

Cyanobacteria-rich topsoil facilitates mine rehabilitation

Wendy Williams¹, Angela Chilton², Mel Schneemilch¹, Stephen Williams¹, Brett Neilan³, Colin Driscoll⁴

1. School of Agriculture and Food Sciences, The University of Queensland, Gatton Campus 4343 Australia

2. Australian Centre for Astrobiology and School of Biotechnology and Biomolecular Sciences, University of New South Wales, Sydney, NSW, 2052, Australia

3. School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW, 2308, Australia

4. Hunter Eco, PO Box 1047, Toronto, NSW, 2283

Abstract

Restoration of soils post-mining requires key solutions to complex issues where the disturbance of topsoil incorporating soil microbial communities can result in a modification to ecosystem function. This research was in collaboration with Iluka Resources at Jacinth-Ambrosia (J-A) mineral sand mine located in a semi-arid chenopod shrubland in southern Australia. At J-A, assemblages of microorganisms and microflora inhabit at least half of the soil surfaces and are collectively known as biocrusts. This research encompassed a polyphasic approach to soil microbial community profiling focused on 'biobanking' viable cyanobacteria in topsoil stockpiles to facilitate rehabilitation. We found that cyanobacterial communities were compositionally diverse topsoil microbiomes. There was no significant difference in cyanobacterial community structure across soil types. As hypothesised, cyanobacteria were central to soil micro-processes, strongly supported by species richness and diversity. Cyanobacteria were a significant component of all three successional stages with 21 species identified from ten sites. Known nitrogen-fixing cyanobacteria *Symploca*, *Scytonema*, *Porphyrosiphon*, *Brasilonema*, *Nostoc* and *Gloeocapsa* comprised more than 50% of the species richness at each site and 61% of the total community richness. In the first study of its kind, we have described the response of cyanobacteria to topsoil stockpiling at various depths and ages. Cyanobacteria are moderately resilient to stockpiling at depth and over time, with average species richness greatest in the top 10 cm of the stockpiles of all ages and more viable within the first six weeks, indicating potential for biocrust re-establishment. In general, the resilience of cyanobacteria to burial in topsoil stockpiles in both the short and long term was significant, however in an arid environment recolonization and community diversity could be impeded by drought. Biocrust re-establishment during mine rehabilitation relies on the role of cyanobacteria as a means of early soil stabilisation. At J-A mine operations do not threaten the survival of any of the organisms we studied. Increased cyanobacterial biomass is likely to be a good indicator and reliable metric for the reestablishment of soil micro-processes.

1.0 Introduction

5 Mine rehabilitation is a complex process that involves many levels of understanding of difficult issues relating to ecosystem function including where the removal or burial of the bioactive soils can have knock-on effects for rehabilitation efforts such as native seedling establishment (Jasper, 2007; Tongway and Ludwig, 1996). Successful ecological restoration of arid mining sites relies on a holistic approach where microbial recolonization can serve as an indicator of the integrity of the wider ecosystem (Tongway, 1990). Microorganisms are critical components of soils that drive assorted micro-processes and impact soil ecosystem function on several levels.

Moved (insertion) [1]

10 In arid landscapes surficial crusts develop as a protective skin across the bare soil or stony interspaces between plants. These can be physical, chemical or biological (microbial) in nature. Well established biological crusts (biocrusts) are intricately interwoven and structured, high-function communities variable in composition that can include cyanobacteria, algae, lichens, mosses, liverworts, micro-fauna and bacteria. They are a significant asset to arid soil ecosystems providing a protective, nutrient-rich layer closely integrated into the soil surface (Delgado-Baquerizo et al., 2013; Maestre et al., 2012).

Deleted: Successful ecological restoration of arid mining sites relies on a holistic approach where biocrust recovery to pre-disturbance levels is integral and can serve as an indicator of the integrity of the ecosystem (Tongway, 1990).

15 In mining, the topsoil is commonly stripped and stored in stockpiles as a form of “biobanking” for redistribution at the cessation of mining. This process involves the crushing and burial of biocrusts and results in the inability of crust organisms to photosynthesise due to the lack of light. Physical disturbance can profoundly disrupt biocrust integrity, composition and physiological function where the impact is governed by site characteristics, severity, frequency and timing (Belnap and Eldridge, 2001). Biocrust burial within the natural environment as a result of environmental stress and disturbance can have severe negative impacts on biocrust integrity and function. For example, with grazing and drought significant declines were recorded in cyanobacterial richness and abundance that was linked to a reduction in soil nutrient concentrations (Rao et al., 2012; Williams and Eldridge, 2011).

25 Following disturbance, restoration and regrowth of biocrusts can take place unassisted, seasonally driven generally over many years (Belnap and Eldridge, 2001; Belnap and Gillette, 1998). Should biocrust organisms remain inactive while they are wet, cell death and decomposition commonly occurs (Kidron et al., 2012; Rao et al., 2012). Nevertheless, in dry conditions, cyanobacteria and algae are known to remain desiccated and viable for millions of years (Vishnivetskaya et al., 2003). Alternatively, assisted biocrust restoration places emphasis on the recovery of ecosystem function and necessarily
30 addresses environmental constraints. This incorporates the knowledge of ‘potential condition’ based on experience with sites of ecological similarity that have undergone disturbance and recovery (Bowker, 2007). Biocrust recovery can be altered by dust deposition, fire and climatic conditions (Weber et al., 2016). When biocrusts recover naturally soil properties change. For example, in Southern African and Spanish rangelands an incremental accumulation of soil nutrients, organic matter

and a build-up of silt and clay lead to the development of a resilient and multi-functional biocrust (Büdel et al., 2009; Maestre et al., 2012; Weber et al., 2016).

Soil disturbance results in a loss of resources and often has long lasting effects on soil stability, nutrient cycling and surface hydrology (Bowker, 2007; Tongway and Hindley, 2004). Cyanobacterial re-establishment is a key indicator of early soil surface re-stabilisation, regulation of soil moisture and the balancing of soil carbon and nitrogen (Elbert et al., 2012). Restoration of ecosystem function post-disturbance requires an appreciation of the dynamic functional status of the landscape prior to disturbance (Tongway and Ludwig, 1996), as well as an understanding of the net accumulative effects of disturbance on the components of the system. On the micro-scale, cyanobacterial species richness contributes to soil ecosystem function through micro-processes including carbon fixation through photosynthesis, atmospheric nitrogen fixation in a biological-available form, micro-nutrient breakdown and release, soil particle cohesion, regulation of moisture and soil surface structure (Delgado-Baquerizo et al., 2013; Elbert et al., 2012; Hu et al., 2002; Maestre et al., 2012 and others). Consequently, it is necessary to appreciate the micro-processes that will assist in the restoration of soil function and to monitor recovery along the way.

Research into biocrust disturbance with a focus on recovery post-mining is rare. In the Namaqualand arid lands (Namibia, South Africa) low rainfall and high winds impact the rehabilitation of degraded lands following diamond mining and grazing (Carrick and Krüger, 2007). These researchers found that cyanobacteria and non-vascular plants that form a living and protective surface crust were crucial to surface stabilisation. Jasper, (2007) also recognised the importance of soil microbial communities including cyanobacteria in post-mine rehabilitation in the Jarrah forests of south-western Australia. In the Czech Republic and Germany chrono-sequential studies of old brown coal mine sites found in younger sites that green algal biofilms and a diverse range of cyanobacteria initiated the rehabilitation of the soils (Lukešová, 2001). In serpentinite mine tailings (New South Wales, Australia), McCutcheon et al., (2016) showed filamentous cyanobacteria accelerated carbonate mineral precipitation and stabilised the tailings. They demonstrated cyanobacteria had the capacity to adsorb magnesium while acting as a nucleation site and sequestering carbon. In our current study, preliminary research identified that in undisturbed chenopod shrublands at the edge of the Nullarbor Plain (South Australia) biocrusts cover more than 45% of the soil surface, between the grass plants and post-mining rehabilitation needs to take their role into account (Doudle et al., 2011). It follows that there is a real need for focus on practical approaches that contribute to the restoration of soil function and measure relevant aspects of success through soil microbial communities and biocrust reestablishment, especially cyanobacteria (for example: Setyawan et al., 2016; Mazor et al., 1996; Fischer et al., 2014; Chiquoine et al., 2016; Doherty et al., 2015; Harris, 2003; Tongway and Hindley, 2004; Zhao et al., 2014).

Deleted: ed

Deleted: s

Deleted: investigate

Deleted: a

Deleted: In mining, the topsoil is often stripped and stored in stockpiles as a form of “biobanking” for redistribution at the cessation of mining. This process involves in the crushing and burial of biocrusts and results in the inability of crust organisms to photosynthesise due to the lack of light. To our knowledge, the effects of topsoil stockpiling on biocrust organisms such as cyanobacteria, and their recovery time, has not previously been investigated. Should biocrust organisms remain inactive while they are wet, cell death and decomposition may occur (Kidron et al., 2012; Rao et al., 2012). Nevertheless, in dry conditions, cyanobacteria and algae are known to remain desiccated and viable for millions of years (Vishnivetskaya et al., 2003).

Deleted: ¶

Microorganisms are critical components of soils that drive micro-processes and impact soil ecosystem function on several levels. Biological soil crusts (biocrusts) are topsoil assemblages of bacteria, fungi, algae, mosses, and lichen which occur where vascular plant growth is restricted, namely arid lands. Highly structured, diverse biocrusts are a significant asset to arid soil ecosystems providing a protective, nutrient rich layer closely integrated into the soil surface (Delgado-Baquerizo et al., 2013; Maestre et al., 2012). As biocrusts develop in structural complexity, the diversity of organisms is regulated by water infiltration, temperature, light and disturbance (Belnap and Eldridge, 2001; Büdel et al., 2009; Elbert et al., 2012). Early successional biocrusts are mainly composed of bacteria, algae and cyanobacteria whereas late successional biocrusts can incorporate lichens, mosses, liverworts, algae, fungi and bacteria (Büdel et al., 2009). ¶

Cyanobacteria in arid landscapes are exceptionally well-adapted to desiccation. Their polysaccharide sheaths and EPS production perform a vital role in maintaining cyanobacterial cell integrity, exchange of information and absorption of water during rehydration (Rossi et al., 2017). EPS has adhesive properties that binds non-aggregated soil particles into a protective encrusted surface that reduces the destructive impacts of wind and water (Eldridge and Leys, 2003; Rossi et al., 2017).
5 Cyanobacterial biofilms provide stabilisation of initially disturbed surfaces that pave the way for diverse microbial communities, and form bioactive crust-like layers assimilated into the soil (e.g. Büdel et al., 2009; Rossi et al., 2017; Bowker et al., 2014). As biocrusts develop in structural complexity, the diversity of organisms is regulated by water infiltration, temperature, light and disturbance (Belnap and Eldridge, 2001; Büdel et al., 2009; Elbert et al., 2012).

10 To our knowledge, the effects of topsoil stockpiling on biocrust organisms such as cyanobacteria, and their recovery time, following topsoil spreading, has not previously been investigated. In this research we focused on the cyanobacterial component of the biocrusts. This was in keeping with the mining framework in the ongoing development of informed rehabilitation plans that focuses on improved long-term outcomes. Specifically, we sought to determine whether shallow
15 biobanks of cyanobacterial-enriched topsoil would facilitate the recovery of essential soil microprocesses when re-spread following mine disturbance. The overall aims of the microbial biobanking research program were to: (a) define the cyanobacterial community structure applying a polyphasic approach with a special focus on species that drive early colonisation, nutrient cycling and soil stabilisation; and, (b) to examine the effects of stockpiling topsoil on cyanobacterial resilience to crushing and burial, and their recovery following spreading of topsoil back across mined land. We hypothesised
20 that cyanobacterial survival within a topsoil stockpile would reduce with both depth within the stockpile and elapsed time before topsoil re-spreading.

2.0 Methods

2.1 Background and site description

Jacynth-Ambrosia (J-A) heavy mineral sand mine is located on the eastern edge of the Nullarbor Plain, South Australia, across the boundary of two Regional Reserves within the Eucla Basin region. The climate is semi-arid with a mean rainfall of 185 mm, mean maximum temperature of 27.6°C and minimum of 12.1°C (further detail provided in supplementary Figure S1, www.bom.gov.au). Tertiary sediments deposited in marine and terrestrial settings and the soil distribution of the area reflects the geological history, with at least five marine transgression and regression events depositing 40-50 m of sediments (Hou and Warland, 2005). The landscape is broadly undulating with low open woodlands that have a shrub understorey with chenopod shrub lands as well as dune fields that consist of parallel dunes and inter-dune swales (Doule et al., 2011; Gillieson et al., 1996). Prior to mining disturbance, the landscape is superficially homogenous chenopod-dominated
30 vegetation but functionally patchy on the fine scale where the soil surfaces are extensively colonised by biocrusts. At J-A,

Deleted: EPS forms the bulk of the biocrust structure. EPS more than doubles the biocrusts compressive strength and increases cohesiveness by up to six times (Hu et al., 2002).

Moved up [1]: Mine rehabilitation is a complex process that involves many levels of understanding of difficult issues relating to ecosystem function where the removal or burial of the bioactive soils can have knock-on effects for rehabilitation efforts such as native seedling establishment (Jasper, 2007; Tongway and Ludwig, 1996). Successful ecological restoration of arid mining sites relies on a holistic approach where biocrust recovery to pre-disturbance levels is integral and can serve as an indicator of the integrity of the ecosystem

Deleted: ¶

Moved down [2]: Our study aimed to examine biocrusts, with a

Moved (insertion) [2]

Deleted: Our

Deleted: study aimed to

Deleted: examine biocrusts, with a focus on

Deleted: .

Deleted: in

Deleted: a

Deleted: to

Deleted: achieve

Deleted: .

Deleted: biocrust

Deleted: on

Deleted: ing

Deleted: ed lands

Deleted: We identified key aspects of biocrust and cyanobacteri(...

Deleted: goals

Deleted: biocrust

Deleted: evaluate specific roles of natural, undisturbed biocrusts(...

Deleted: in terms of key

Deleted: biogeochemical cycling

Deleted: c

Deleted: investigate

Deleted: survival

Deleted: after

Deleted: al and subsequent recovery

Deleted: .

Deleted: We hypothesised that cyanobacteria would

Deleted: be essential as ecosystem engineers in soil habitat (...)

Moved (insertion) [3]

Moved up [3]: in the topsoil stockpiles

biocrusts primarily consist of lichens, mosses, and cyanobacteria that cover around 45% of the landscape surfaces, equating to 2,000 hectares of the mining lease (Doudle et al., 2011).

At J-A the landscape has been characterised into three distinct soil types, that were associated with vegetation communities identified as soil management units (Doudle et al., 2011; Hou and Warland, 2005). Site vegetation associations are described as follows: SMU 1 – Red Mallee: *Eucalyptus oleosa* ssp. *oleosa* = open Mallee/Myall woodland; SMU 2 – Chenopod Shrubland: *Maireana sedifolia* and *Atriplex vesicaria*; SMU 3 – Western Myall: *Acacia papyrocarpa* *Maireana sedifolia* = open Myall woodland Site 1 occurs in a transition between SMU 2 and SMU 3 but was treated as most like SMU 2 (also see Table S1). Soil management units were summarised as: SMU 1 – deep calcareous yellow sands associated with dune ridges; SMU 2 – shallow calcareous sandy loams; and SMU 3 – deep calcareous sandy loam (Table S1). In the first place, sample site locations (Fig. 1) were selected based on these soil management units (SMU) and a two-year old stockpile (Table S1). Secondly, sites within these parameters (SMU) were selected for the subsequent detailed studies of cyanobacterial succession and its resilience to longer-term stockpiling (Fig.2 and Section 2.4).

The biocrusts at J-A had been previously classified into three successional stages representative of the five biocrust types found growing across the landscape (Table S1, S2) (Doudle et al., 2011). Types 1–2 are light coloured, patchy, thin, and fragile cyanobacterial crusts corresponding to early stages of development; Type 3 are well established cyanobacterial crusts with establishment of some mosses and lichens corresponding to intermediate stages of development; Types 4–5 biocrusts are well established with cyanolichens and/or green algal lichens and mosses corresponding with late stages of development (additional descriptions available in supplementary Table S1). As well as the three main SMU sites we also sampled a dry salt lake (Lake Ifould) to provide information on cyanobacterial species adapted to saline conditions, similar in nature to ground water used in mine operations. In this study the term biocrust covers whole crust samples that incorporated lichens cyanobacteria and mosses in varying proportions; however the cyanobacterial component of this crust was the focus in terms of our polyphasic approach to community structure, succession and its biophysiochemical properties.

In the process of mining and preparation of the J-A site for future rehabilitation, topsoil (0–100 mm) is stored in low stockpiles, generally less than two metres in depth where stockpiles are comprised of topsoil sourced from only one vegetation type. Later stockpiles will be returned to the surface of mined areas in the rehabilitation process, and the cyanobacterial activity therein must be quantified to enable educated planning and decision making regarding biocrust re-establishment. If low levels of biocrust organisms are detected below the top few centimetres, the addition of propagated biocrust organisms (e.g. cyanobacteria) to returned topsoil may be warranted. Activity and species richness within the stockpiled top soils may also vary with age and this may also influence the establishment of vegetation in the rehabilitation process.

Deleted: T

Deleted: (Table 1)

Deleted: called

Field Code Changed

Formatted: Font: (Default) +Body (Times New Roman)

Deleted: :

Formatted: Font: (Default) +Body (Times New Roman)

Deleted: (Doudle et al., 2011; Hou and Warland, 2005)

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman)

Deleted: 2

Deleted: locations were selected for: (a) vegetation communities, (b) soil types and underlying geology, (c) biocrust successional types, (d) areas identified for future mine operations and a dry salt lake (Lake Ifould). The dry salt lake would potentially identify cyanobacterial species adapted to saline conditions similar in nature to ground water.

2.2 Biophysical characteristics of biocrusts and cyanobacteria

2.2.1 Field Sampling

Preliminary identification of biocrust types had been determined by Doudle et al., (2011) and provided the baseline data for biocrust sampling from Sites 1–10 (Fig. 1, Table S1). The sites selected encompassed the three SMUs and five crust types (Table S1). All sites are of naturally occurring biocrusts except for Site 6 which is from a two-year-old topsoil stockpile regarded as an early (Type 1) biocrust representing recovery two years post disturbance (Fig. 2a-c). Within each site, eight 10 cm diameter samples were selected at random and removed to a depth of 1 cm using metal scraper (n=80), air dried (>40°C), and stored in Petri dishes. Each Petri dish contained approximately 80 g of crust. The samples were packed to avoid crust disruption and transported to The University of Queensland's Central Analytical Laboratory at Gatton.

2.2.2 Biocrust biophysiochemical properties

For each site, ~~about half the sample was removed from each Petri dish and fine sieved (1.70 mm). Duplicate sub-samples (~2 g), were analysed for total C and N and C:N ratio using a high temperature digestion in a vario MACRO Elemental Analyser (Elementar) (n=6 per SMU). Duplicate samples (10g) for the purpose of analysing soil pH and electrical conductivity (EC) were prepared using a 1:5 (soil to water) ratio and shaken for one hour. Following shaking, samples were left to stand for 30 min and EC measured using a Crison Conductivity Meter 525. The sample was mixed again, and pH was measured with a TPS pH meter MC-80 using an ionode IJ44C electrode. The remaining half of the Petri dish was used to determine Chlorophyll a concentration of the biocrusts. Following resurrection (by moistening) the crust was lightly homogenised and a 5 g sample used at a 1:5 ratio of (dry weight) biocrust to Dimethyl sulfoxide (DMSO) (Barnes et al., 1992) with samples placed in a warm bath (65°C) for a two-hour dark extraction, followed by centrifuging for five minutes (5000 g RCF). Chlorophyll a concentration was determined using Wellburn's, (1994) equations.~~

~~Prior to the use of the samples for analysis a pocket penetrometer (8 mm foot) was used to determine the compressive strength (kg cm⁻²) of the dry intact biocrusts samples. Overall the crust thickness was < 0.5 cm. Each sample (10 cm diameter x 2 cm depth) was placed on a solid surface and a total of twelve measurements (3 readings x 4 reps) were taken for each site. The measurement was taken at the point when the crust was broken, and the foot penetrated the crust surface.~~

~~At J-A we measured photosynthetic performance (recorded as yield, YII) of the biocrusts using a pulse-amplitude modulated (PAM) fluorometer (Pocket PAM; Gademann Instruments, Germany). The goal was to demonstrate photosynthetic yield (YII) indicative of active growth of the biocrusts, using the detection of chlorophyll fluorescence from photosystem II (PSII). The sensor was placed onto the biocrust and once started, a series of short pulses of excitation light at high intensity that is~~

Deleted: duplicate sub-samples

Deleted: were

Deleted: taken

Deleted: for each biophysiochemical analysis (n=16 per site

Deleted:). Soil was fine sieved (1.70 mm) and

Formatted: Not Highlight

Deleted: S

Deleted: samples

Deleted: were determined f

Deleted: using

Deleted: A

Deleted: P

Deleted: was measured

Deleted: aim

amplified resulting in a brief closure of PSII and the measurement of fluorescence yield based on the Genty parameter which is the quantum yield (YII) of the charge separation of PSII (Genty et al., 1989) and recorded on a scale of 0–1 for all photosynthesis. In the field this process was completed at least six times for each sample (6 x 4 reps per site).

Deleted:

5 One-way ANOVA with Tukey post-hoc tests were run to test for significant differences between SMU's and total C and N and C:N ratio (Minitab 18). Chlorophyll a and YII were independently tested with a one-way ANOVA and Tukey post-hoc test for significance between SMU.

Formatted: Font color: Auto

2.3 Cyanobacterial community structure

10

2.3.1 Microscopy of biocrust cyanobacteria

Field samples collected in section 2.2 were used to enumerate cyanobacterial richness and diversity and to classify colonies. The dried crust samples were resurrected in the glasshouse for three to five days. For each of the ten sites, twelve replicates (subsampled from the eight samples) were analysed via light microscopy. For each replicate, a minimum of two wet-mount slides incorporating six representative portions of the cyanobacterial colonies were examined (n=144 colonies per site). For the dominant land type, Chenopod shrubland (Site 8), there were an additional 10 x 6-cell multi-well plates. These were treated similarly where two slides were examined from each of the 60 multi-wells (n=120). In total > 2,184 cyanobacterial colonies were examined. Initial inspection of the biocrust and the separation of individual species were made using an Olympus SZH10 microscope at 70 x magnification. Cyanobacterial filaments or colonies were carefully extracted with forceps to recover sufficient material that included important morphological features such as their colour, encasing sheaths as well as cellular structure. Live material was examined by Nomarski differential interference contrast (DIC) microscopy with a JenaVal (Jena Zeiss) and an Olympus BX51 compound microscope (magnifications 400–1000 x). Photomicrographs were taken using an Olympus SC100 digital microscope camera, and morphological measurements of vegetative cells were made from digital images of live material taken at 400 x magnification using Olympus cellSens® digital imaging software.

Identification was performed to a species level (wherever possible) in the laboratory using the following taxonomic references: Anagnostidis and Komarek, (2005, 2005); Sant'Anna et al., (2011); Skinner and Entwisle, (2002). It was often necessary to record the closest named species as attributes varied somewhat to temperate climate and aquatic specimens described in literature. Nitrogen fixing cyanobacteria were identified based on the three recognised types: (1) heterocystous species (those with specialised N-fixing cells); (2) non-heterocystous species that fix N aerobically and; (3) non-heterocystous species that fix N anaerobically (Bergman et al., 1997; Stal, 1995). Using a graticule, abundance was ranked on a scale of 1–8 where the main taxa are ranked in decreasing order of the relative percentage area occupied in a single view

30

(Biggs and Kilroy, 2000). More than one species could be dominant, and all other taxa were ranked in relation to the dominant taxa as abundant, common, occasional and rare. To determine similarities between cyanobacterial communities, cluster analysis, SIMPROF, and non-metric multidimensional scaling (nMDS) were conducted using Primer v6 (Clarke & Gorley 2001).

5 2.3.2 16S rDNA profiling of native undisturbed biocrust microbiomes

For genomic profiling of naturally occurring successional biocrust communities, a location adjacent to Site 9 was visually determined to contain Bare, Early (Types 1-2) or Late (Types 4-5) stages of development (Table S2). Biocrust successional features were determined by morphological attributes of pigmentation, thickness and surface roughness as well as the presence/absence of lichens and mosses (Fig. 2d) (Chilton et al., 2017). Bare stage was characterised by loose soil particles with no visible biocrust structure. Samples were collected in July 2014. For each successional stage, three replicates were collected that were representative of SMUs 1–3 where a 10 cm² plot with 95% coverage of the desired biocrust stage was excised to the depth of the crust and non-aggregated soil discarded (Fig. 2e-g). Samples were processed at UNSW, Sydney.

Each biocrust replicate for Bare, Early and Late stages of development were homogenised and genomic DNA extraction performed using the FASTDNA Spin Kit for Soil (MP Bio Laboratories, USA) according to the manufacturer's instructions. Molecular libraries of the 16S rDNA V123 hypervariable region generated via PCR as per Chilton et al., (2017) and submitted to the Ramaciotti Centre for Genomics (UNSW, Australia) for a 2x300 bp sequencing run on an Illumina MiSeq instrument. Sequencing data was processed using Mothur version 1.34.0 (Schloss et al 2009) and described in detail in Chilton et al., (2017). Singleton and doubleton OTUs were removed and samples rarefied to 8598 sequences each across 3785 OTUs. The curated Greengenes database (McDonald et al 2012) was used to assign taxonomy to OTUs. Diversity values were derived using the DIVERSE function within the Primer package (Anderson et al 2008) upon standardized OTU values. ANOVA with post hoc Tukey's tests was used to test for significant differences between stages. Multivariate analyses were performed in Primer upon a Bray-Curtis dissimilarity matrix generated from square-root transformed abundance data. Samples were represented in two and three-dimensional space within a nMDS plot. Pair-wise, a posteriori comparisons of factor Stage were performed using the PERMANOVA function with 9999 Monte Carlo permutations. Homogeneity of dispersion for each stage was tested using PERMDISP.

2.4 Cyanobacterial tolerance to stockpiling

Stockpile sampling was carried out in March 2012 with samples sourced from topsoil stockpiled from areas with *Acacia papyrocarpa* (Western Myall) over-storey (Table S3). Stockpiles established at three different time points (9 months, 20 months and 29 months) as well as corresponding undisturbed sites were sampled in triplicate at six depths (0–2, 2–4, 4–6, 10, 25 and 50 cm). This resulted in 18 replicates for each soil depth for each stockpile time point. Holes were dug to >50 cm depth with a shovel and the exposed profile removed with an ethanol-wiped spatula before taking a soil sample. A second

sample at 50 cm was autoclaved to serve as a culturing control. Samples were stored in paper bags and processed at The University of Queensland. For each sample, 20 g of soil was set up in petri dishes with 9 ml of water and sealed with Parafilm®. Petri dishes were incubated for six weeks at 26°C under a 12-hour photoperiod regime and rotated weekly to prevent site specific effects. Samples were maintained as wet by the addition of sterile water within a laminar flow cabinet.

To determine cyanobacterial growth and richness, wet-mounts for each sample were examined under 16 x magnification for cyanobacterial thalli and colony size was estimated via area of coverage of the field of view. Where multiple colonies were present in a slide, the relative abundance of each was used to divide the cover between the colonies accordingly. Where no growth was observed, five soil samples were taken randomly from the sample, mounted and examined under 400 x magnification for the presence of cyanobacterial cells. ANOVAs were performed with post hoc Tukey's tests performed to determine any in relative abundance between treatments.

Deleted: differences

3.0 Results

3.1 Biophysical characteristics of biocrusts and cyanobacteria

Soil pH across the three soil management units (SMUs) ranged from pH 8.4–8.6 while the two-year old topsoil stockpile was higher at pH 8.9 (Table 1). Electrical conductivity ranged from 92–140 $\mu\text{S cm}^{-1}$. There were no significant differences in pH and EC between field samples and intact samples (data not shown). Total nitrogen was typically <0.1% across all sites and total carbon ranged between 1–2% with higher percentages generally found across SMU 2 and 3 (Table 1). The ratio between carbon and nitrogen was the greatest across SMU 3 and the two-year old topsoil stockpile, also originating from SMU 3. Percentage C and C:N ration across all SMU showed there were significant differences (p=0.052), and post hoc test showed only SMU 1 and SMU 3 means were significantly different. There were no significant differences between SMU 1 and SMU 2, or SMU 2 and SMU 3. Percent N versus SMU was significant overall (p=0.000) with all three SMU are significantly different from each other.

Moved (insertion) [5]

Moved up [5]: The ratio between carbon and nitrogen was the greatest across SMU 3 and the two-year old topsoil stockpile, also originating from SMU 3.

For the intact undisturbed biocrusts mean chlorophyll *a* concentration (10.33–13.64 $\mu\text{g g}^{-1}$ soil) differed across all sites and SMU 1 biocrusts were significantly lower ($p = 0.000$) compared to SMU 2 and 3 which were not significant (Fig. 3). The mean chlorophyll concentrations of biocrusts sourced from the two-year old topsoil stockpile ($7.49 \pm 1.01 \mu\text{g g}^{-1}$ soil) were almost half the concentration of SMU 3 ($13.53 \pm 1.74 \mu\text{g g}^{-1}$ soil), which was the origin of the topsoil. Thus Chlorophyll a concentration for the two-year old stockpiles were significantly different to SMU 1 and 2 and SMU 3 (p=0.000).

Deleted: ¶

Deleted: concentrations

Deleted: 5

Deleted: T2,

Formatted: Font: 10 pt

The photosynthetic yield measurements showed there was variability between sites (Fig. 3). The yields recorded all fell within the expected range and when data was separated into their respective SMU it provided a clearer picture of the

variability. SMU 1 and SMU 2 were not significantly different whereas SMU 3 and the two-year old stockpile were significantly different from SMU 1 and SMU 2 ($p = 0.017$).▼

Deleted: SMU 3 had the lowest mean overall while there were no differences between the others.

Deleted: ¶

▼
The penetrometer index of crust compressive strengths across the ten sites showed that Sites 4, 9 and 10 in SMU 1 were significantly different from each other as well as significantly lower than SMU 2, SMU 3 and T2, two-year old topsoil stockpile sites (Table 2). The means for SMU 1–3 were also significantly different ($p < 0.0001$), and it was confirmed by a student's t-test that the difference was between SMU 1 and SMU 2 and 3.

3.2 Cyanobacterial community structure

A total of 21 cyanobacterial species were identified across the ten sites using microscopy (Table 5; Figs. S2–S9). The majority of species richness and abundance was comprised of four filamentous genera: *Symploca* (18%), *Schizothrix* (16%), *Porphyrosiphon* (16%) and *Scytonema* (16%). Secondary cyanobacteria present variously occupied <1 to 10%, and overall made up 34% of the community. The known nitrogen-fixing cyanobacteria *Symploca*, *Scytonema*, *Porphyrosiphon*, *Brasilonema*, *Nostoc* and *Gloeocapsa* comprised more than 50% of the species richness at each site and formed 61% of the total community richness (Fig. 4). In this study cyanobacterial community structure was tested against the three different soil types to determine whether soil type was influential in determining community structure. There was no significant difference in cyanobacterial community structure across soil types, however the results do suggest some spatial structuring exists across the three SMU (Fig. 5a). The structural relationship between all samples shows greater similarities rather than dissimilarities, of which there are only a small number of samples that are significantly different (Fig. 5b). Soil type did not explain these differences. This suggests that most of the species could potentially be found anywhere across the three zones and their richness and abundance is controlled by other factors. *Symploca* occurred more frequently and was more abundant in the majority of the samples examined therefore was the most significant contributor to the community (data not shown). Individual sites generally displayed similar trends although there was some variability occurring between sites.

Of the 21 species more than half (12 species) of cyanobacteria were identified in SMU 1 where four primary genera made up 75% of the community: *Symploca*, *Schizothrix*, *Scytonema* and *Symplocastrum* (for more detail see Fig. S10). Cyanobacterial crusts from the dune regions on SMU 1 (deep calcareous yellow sands) were representative of crust types 1–3; patchy, brittle (when dry) early-successional crusts as well as formed dark crusts that were mid to late-successional and included cyanolichens (also see Doudle et al., 2011).

Cyanobacterial crusts from the chenopod shrublands and open woodlands in SMU 2 (shallow calcareous sandy loam) represented a broad range of crust types (2–5) but overall could be described as late-successional. Lichens and mosses were highly visible (also see Doudle et al., 2011). There were 21 cyanobacteria recorded: four were primary genera that made up 63% of the community including: *Schizothrix*; *Porphyrosiphon*; *Scytonema* and *Symploca* (for more detail see Fig. S11).

Cyanobacterial crusts from the open woodlands in SMU 3 (deep calcareous sandy loam, Fig. 2c) represented a broad range of crust types (2–5) but like SMU 2 could also be described as late-successional. Lichens and mosses were highly visible (see Doudle et al., 2011). There were nine cyanobacteria recorded of which four were primary genera that made up 85% of the community: *Symploca*, *Porphyrosiphon*, *Scytonema* and *Schizothrix* (for more detail see Fig S12). Cyanobacteria with the capacity to fix nitrogen contributed to 77% of the community structure.

Cyanobacterial crusts from Site 6 were from the two-year old topsoil stockpile (T2) that had originated from SMU 3 (deep calcareous sandy loam) would be described as early successional crusts with some seasonal mosses. There were eight cyanobacteria recorded of which four were primary genera that made up 84% of the community: *Symploca*; *Symplocastrum*; *Porphyrosiphon*; and *Scytonema* (also see Fig. S13). It was interesting to note that *Symplocastrum* was co-dominant with *Symploca* whereas in the other communities it ranged between 8-13%. Sub-surface species *Schizothrix* (found in top 5 mm) only contributed to 4% of the richness compared to 10-20% elsewhere. Cyanobacteria with the capacity to fix nitrogen (*Symploca*, *Porphyrosiphon*, *Scytonema* and *Brasilonema*) contributed to 61% of the community (Fig. 4).

3.3 16S rDNA profiling of native undisturbed biocrust microbiomes

Microbial community profiling using high through-put sequencing revealed cyanobacteria comprised a significant component of all three stages forming the majority of sequences in Early and Late stages (Fig. 6a). There was a diversity of morphotypes observed including simple filamentous, heterocystous and unicellular types (Fig. 6b). The most abundant genera identified were *Leptolyngbya*, *Phormidium*, *Tolypothrix*, *Nostoc*, *Brasilonema*, *Chroococcidiopsis* and *Acaryochloris*. Unclassified Nostocaceae were dominant within Bare soils while Early stages observed a relative even increase in *Phormidium*, *Brasilonema* and the unicellular genera (e.g. *Chroococcidiopsis*, *Acaryochloris*, *Xenococcaceae*). Late stage biocrusts showed a slight resurgence of Nostocaceae. There was no significant difference in the richness, evenness or diversity between the three stages (Table 3). However, there were significant differences in the composition and structure of the communities of each stage. PERMANOVA showed stage was a significant factor explaining the data variability (Pseudo- $F=4.9544$, $P(\text{perm})=0.004$, Unique perms=273). Post-hoc pair-wise comparisons showed Bare stage was significantly different to the crusted stages (Fig. 7, Table 4) while resolution between the Early and Late stages was less clear. Ordination of the samples within three dimensions showed Early and Late stages grouped separately however, no significant difference between these stages was detected via PERMANOVA (Table 3).

3.4 Cyanobacterial tolerance to stockpiling

Examination of stockpile soil samples via microscopy revealed five cyanobacterial genera: *Nostoc*, *Scytonema*, *Microcoleus*, *Porphyrosiphon* and *Leptolyngbya*. Average genera richness was highest in stockpiled samples at and above 10 cm depth for

Deleted: The Bare stage was significantly different to the crusted stages (Fig. 7, Table 4) while resolution between the Early and Late stages was less clear. Ordination of the samples within three dimensions showed Early and Late stages grouped separately however, PERMANOVA failed to identify a significant difference (Table 3). ...

all stockpile ages (Fig. 8 a, b). These differences were significant between 50 cm and 0–2 cm in all stockpiles (29-month $F_{(1,5)} = 7.00$, $p = 0.024$; 20-month $F_{(1,5)} = 8.37$, $p = 0.016$; 9-month $F_{(1,5)} = 18.00$, $p = 0.002$). Genera richness at the intermediate depths was not significantly different. Differences were significant between 25 cm and 0–2 cm in the twenty- and nine-month old stockpiles ($F_{(1,5)} = 32.73$, $p = 0.000$; $F = 16.20$, $p = 0.000$) respectively. Counterintuitively, after nine and
5 twenty months of stockpiling, average genera richness was higher in stockpiled samples when compared with undisturbed samples except for 20-month old 4–6 cm samples (Fig. 8 a, b) but this difference was only significant in the nine-month old stockpile at 10 cm depths ($F_{(1,5)} = 8.27$, $p = 0.017$). High variability between replicates accounted for the lack of significance in the rest.

10 When comparing average genera richness at different depths between stockpiles, richness was greater in material stockpiled for the least amount of time above 10 cm depth but not at or below this level (Fig. 8b). Average genera richness was variable within adjacent undisturbed areas (Fig. 8a). The variability in genera richness within replicates was high (refer to Fig. S14 a–c). The cover of genera was relatively constant between sites in the undisturbed samples, but in stockpiled samples *Nostoc cf commune* was more prevalent ($F_{(1,5)} = 5.97$, $p = 0.012$).

15 *Nostoc cf. commune*, *Nostoc* yellow, *Microcoleus* and *Leptolyngbya* were present in more stockpiled samples than undisturbed samples. Conversely, *Scytonema* and the black form of *Leptolyngbya* were more prevalent in undisturbed samples. When all samples were combined, *Nostoc cf commune* had the greatest cover in both stockpiled and undisturbed samples followed by yellow *Nostoc* in stockpiles only. The coverage by the remaining genera was similar for most other
20 genera although *Microcoleus* and *Leptolyngbya* had greater cover in stockpiled areas and *Porphyrosiphon* had slightly higher coverage in undisturbed samples. The cover of *Scytonema* was almost the same in stockpiled and undisturbed areas ($F_{(1,17)} = 0.00$, $p = 0.969$) (also see Fig. S15).

25 *Nostoc cf. commune* exhibited the best survival response in stockpiling as it had significant coverage in samples from all depths in all stockpile ages. *Nostoc* had the greatest coverage in the oldest stockpiles, followed up by the most recently created stockpiles. These differences were significant from depths of 6 cm to the surface (4–6 cm $F_{(1,5)} = 28.83$, $p = 0.000$; 2–4 cm $F_{(1,5)} = 4.89$, $p = 0.023$; 0–2 cm $F_{(1,5)} = 3.72$, $p = 0.049$). These patterns were not reflected in the total coverage by *N. cf. commune* in corresponding undisturbed areas. The *Leptolyngbya* black morphotype only occurred between 20 and 40 cm depth and were found in only one stockpile samples but were present in six samples from three adjacent areas. *Stigonema*
30 genus was found in six stockpile samples spanning all ages and in only one adjacent sample in all cases in the upper 10 cm of the soil profile.

Nostoc cf. commune, *Porphyrosiphon*, *Microcoleus* and *Scytonema* were the first genera to develop to an identifiable stage. Filaments of the *Stigonema* genus were found in low numbers and appeared to be recently formed. It was only present in

samples examined in the latter stages of the identification process. The yellow form of *Nostoc* exhibited a much slower rate of development than *Nostoc cf commune* and could only be definitively determined as a form of *Nostoc* when examined after 13 weeks of incubation. Samples from undisturbed areas sourced at depths from between 10 cm and 50 cm initially showed no visible signs of growth when examined after six weeks of incubation. Six weeks later cyanobacterial growth was evident yet in many cases had not advanced to the point where morphotypes could be distinguished.

4.0 Discussion

In terms of rehabilitation the natural capital in topsoil that has been removed in the mining process is often not recognised or poorly understood. Soils properties are crucial in the re-establishment of a raft of ecosystem services that include fertility, structure, climate regulation and biodiversity (Dominati et al., 2010). Cyanobacteria are ecosystem engineers in that they have the capacity to provide many of these crucial ecosystem services (Jones et al., 1994). This research demonstrated that the cyanobacterial communities in the J-A biocrusts were compositionally diverse topsoil microbiomes that substantially contributed to the Myall-chenopod biomes. We had hypothesised that cyanobacteria would be central to soil micro-processes and this was strongly supported by extensive species richness and diversity.

In the first study of its kind we have shown the response of cyanobacteria to topsoil stockpiling at various depths and ages. In this study we have shown cyanobacteria are moderately resilient to stockpiling at depth and over time, providing they are manipulated dry (W. Williams, unpublished data). Cyanobacteria from the top 10 cm were found to be more viable within the first six weeks and showed potential for biocrust re-establishment. We found greater cyanobacterial richness in the nine and 20-month stockpiles compared to undisturbed samples adjacent to the stockpiles. In general, the resilience of cyanobacteria to burial in topsoil stockpiles in the longer term appeared good, however in an arid environment recolonization and community diversity could be impeded by drought (Williams and Büdel, 2012)

In this study cyanobacterial community richness and abundance were not related to soil and landform type and this was further supported though the sequenced samples where no significant differences were observed. At J-A any of the cyanobacteria could conceivably occur anywhere across the landscape. Their relative abundance was most probably determined by microenvironments and microhabitats such as light (sun and shade) and chemical gradients (Stal, 2003), as well as moisture availability (Büdel et al., 2018) and soil particle size (Büdel et al., 2009).

4.1 Cyanobacterial community structure

Twenty-one cyanobacteria were recorded from 13 genera. Four species were unicellular and the remaining seventeen were filamentous. Some cyanobacteria found at J-A (*Microcoleus paludosus*, *Nostoc* sp., *Gloeocapsa*) had also been recorded at Lake Gilles (SA) about 400 km southeast of J-A (Ullmann and Büdel, 2001). Although *Microcoleus* species were recorded at

Deleted: ¶

Formatted: Font color: Blue

Deleted: ,

Deleted: with t

Deleted: likely

Moved (insertion) [4]

J-A they did not dominate the biocrust compared with many reports from the United States, Asia and elsewhere (e.g. see Belnap and Eldridge, 2001). This infers that the early colonisers such as *Microcoleus* would not play a dominant role in early stabilisation and colonisation of the soil. At J-A *Symploca* and *Scytonema* appeared to be an important colonising cyanobacterium in the biocrusts and have been recorded as playing a key role in carbon sequestration in northern Australian cyanobacterial crusts (Büdel et al., 2018).

Key cyanobacteria indicating biocrust formation and development were *Leptolyngbya*, *Phormidium*, *Tolypothrix*, *Nostoc*, *Brasilonema*, *Chroococcidiopsis* and *Acaryochloris*. These genera have consistent morphological traits with those observed via microscopy. Notably, the identification of *Brasilonema* was supported with sequencing data and had not been previously recorded in Australian soils. Simple filamentous types are often attributed with the primary crust building role, able to span inter-particle gaps within the soil via supra-cellular structures (e.g. *Microcoleus*, see Garcia-Pichel and Wojciechowski 2009). Sequencing data showed *Phormidium* was the dominant cyanobacterium for this role and it is likely that *Symploca* identified though microscopy was the principal *Phormidium* present. *Microcoleus* sp. and *Porphyrosiphon* were also identified as early colonisers however these genera are currently poorly resolved phylogenetically (Garcia-Pichel et al 2013) but share critical morphological features enabling biocrust formation and maintenance. Notwithstanding, the important crust-building cyanobacteria in this study appeared to be *Symploca* that was associated with EPS production, and principal feature of early colonising crust formation (Hu et al., 2002).

The cyanobacterial richness at J-A was determined according to their morphological features. In many cases these features (e.g. outer protective sheaths, UV protection, EPS production) provided the basis of attributes that pertained to fundamental survival strategies. Environmentally induced strategies of arid land cyanobacteria reflect their habitat, these survival traits have developed over a long evolutionary history. Many primary (common to abundant) and secondary (uncommon) cyanobacteria recorded at J-A exhibited thick gelatinous sheaths (*Porphyrosiphon*, *Schizothrix*, *Microcoleus*, *Nostoc*) or were associated with the production of EPS (*Symploca*, *Nostoc*, *Schizothrix*, *Leptolyngbya*). Filamentous cyanobacteria formed the major part of the J-A crust structure with tufts, webs or creeping masses closely intertwined (e.g. *Porphyrosiphon*, *Symploca*, *Scytonema*, *Schizothrix*, *Microcoleus*). These are often assimilated with unicellular forms (e.g. *Gloeocapsa*, *Chroococcus*, *Chroococcidiopsis*) or gelatinous colonies of *Nostoc* (see supplementary material Fig. S16 for images of growth habits).

The taxonomic status of *Brasilonema* remained uncertain and may be a variety of *Scytonema*, however, genomic data supported morphological identification and the type has also been recorded in other terrestrial habitats globally. Due to its similar morphological attributes and genomic data, in this study we called this cyanobacterium *Brasilonema* (Fiore et al., 2007; Vaccarino and Johansen, 2012). *Nostoc commune* var. *flagelliforme* had been recorded at J-A along with *Nostoc commune* across the shallow and deep sandy loams. Although *N. flagelliforme* appeared rarely, it had been previously

Moved (insertion) [7]

Moved up [7]: Notwithstanding, the important crust-building cyanobacteria in this study appeared to be *Symploca* that was associated with EPS production, and principal feature of early colonising crust formation (Hu et al., 2002).

Deleted: ¶

Deleted: ¶

Moved up [4]: Twenty-one cyanobacteria were recorded from 13 genera. Four species were unicellular and the remaining seventeen were filamentous. Some cyanobacteria found at J-A (*Microcoleus paludosus*, *Nostoc* sp., *Gloeocapsa*) had also been recorded at Lake Gilles (SA) about 400 km southeast of J-A (Ullmann and Büdel, 2001). Although *Microcoleus* species were recorded at J-A they did not dominate the biocrust compared with many reports from the United States, Asia and elsewhere (e.g. see Belnap and Eldridge, 2001). This infers that the early colonisers such as *Microcoleus* would not play a dominant role in early stabilisation and colonisation of the soil. At J-A *Symploca* and *Scytonema* appeared to be an important colonising cyanobacterium in the biocrusts and have been recorded as playing a key role in carbon sequestration in northern Australian cyanobacterial crusts (Büdel et al., 2018). ¶

documented from sites in south-western South Australia, Western Australia, Northern Territory (Skinner and Entwisle, 2002) and Victoria (W. Williams, unpublished data). A joint Spanish-Australian study has now shown that both *Nostoc commune* and *N. flagelliforme* contain the same genomic markers and cannot be separated, rather the spaghetti-like tubes that are unique ecotype likely associated with aridity (Aboal et al., 2016). This is supported by the semi-arid environment at J-A and it may be more widespread in Australia than previously recorded as it is often only clearly visible following rains.

4.2 Cyanobacterial tolerance to stockpiling

Physical disturbance of biocrusts occurs on a large scale at the J-A mine site with the removal and temporary stockpile storage of topsoil. This type of mechanical disturbance results in the burial and translocation of the biocrust. The impacts of burial within the natural environment are rarely studied. In China, artificial sand burial at shallow depths showed there were significant reductions in chlorophyll concentration, UV synthesis, total carbohydrates (EPS) and damage to photosynthetic activity (Rao et al., 2012). In a semi-arid grassland in Australia, wind-borne sand burial of cyanobacterial crusts during a severe drought resulted in a significant reduction in surface dwelling cyanobacteria and significant reductions in biological-available nitrogen (Williams and Eldridge, 2011).

Here we have shown that without further disturbance, a proportion of cyanobacteria can survive stockpiling for over two years. This is not surprising due to the recognised ability of cyanobacteria to survive in extreme environments. In previous studies, cyanobacteria have been grown from samples sourced at 18 cm depths in Japanese rice paddy soils (Fujita and Nakahara, 2006), 50 cm in the UK (Esmarch, 1914), and 70 cm depths in the USA (Moore and Karrer, 1919). Yet, the cyanobacteria from the surface depths of the stockpiles (i.e. <10 cm) appeared more resilient in the short-term.

The species sampled at J-A have a proven track record of survival under extreme conditions. *Microcoleus* and *Leptolyngbya* have survived and remained viable after up to three million years frozen in lake sediments in permafrost (Vishnivetskaya et al., 2003). Vegetative *Nostoc commune* material retains viability following several decades of storage in desiccated form (Bristol, 1919; Lipman, 1941). Reactivation of vegetative material after decades of storage was successful but several months (Lipman, 1941) to a year (Bristol, 1919) of incubation can be necessary for growth to take place. These results were reflected in the current study where growth was not observed in the undisturbed areas below 10 cm depth for several months. It may be that the longer the period of inactivity, the longer time taken for reactivation to occur (Billi and Potts, 2002; Williams et al., 2014), or less material is viable therefore it takes longer to rebuild colonies (Agrawal and Singh, 2002).

Akinetes are desiccation resistant cells produced by certain filamentous cyanobacteria that can survive for long periods. *Nostoc* and *Scytonema* produce akinetes (Kaplan-Levy et al., 2010; Tomaselli and Giovannetti, 1993) but many of the other species sampled in this study cannot, therefore alternative survival methods are in action. Heterotrophic growth is also possible for some cyanobacteria (Flores and Herrero, 2010). Cyanobacteria can survive in darkness through utilisation of alternate carbon sources in drinking water systems (Codony et al., 2003) and this may also be true for soil cyanobacteria

Table 1: Intact biocrust soil physicochemical descriptions for all sites EC = electrical conductivity in $\mu\text{S cm}^{-1}$; total percentage of nitrogen present (N%), total percentage of carbon present (C%) and carbon to nitrogen ratios (C:N) for all sites. *SMU3 is the origin of the topsoil stockpile aged two years which was not analysed with SMU1-3 as there were only two samples. Different letters indicate significant differences in columns.

Vegetation	Soil	SMU	Site	pH	EC	N%	C%	C:N
<u>Mallee</u>	Deep calcareous yellow sand	1	4.9.10	8.6	92	0.07 ^a	1.11 ^a	14.9 ^a
<u>Mvall. Mallee, Chenopod</u>	Shallow calcareous sandy loam	2	1.5.8	8.5	122	0.11 ^b	1.40 ^{a,b}	13.0
<u>Mvall</u>	Deep calcareous sandy loam	3	2,3,7	8.5	113	0.09 ^c	1.61 ^c	17.3 ^a
<u>Mvall</u>	Topsoil stockpile	<u>Origin</u> *SMU3	6	8.9	119	0.08	1.57	19.6

Table 2: Biocrust compressive strengths measured with a penetrometer, means and standard deviations (SD) for SMU 1–3 (kg cm^{-2}) tests with p-values, values in bold that are different from 0 with a significance level $\alpha=0.05$. Letters indicate significant differences

SMU	Means \pm SD	SMU 1	SMU 2	SMU 3
SMU1 ^b	2.79 \pm 1.41	0	0.001	0.005
SMU2 ^a	3.75 \pm 0.79	0.001	0	NS
SMU3 ^a	3.97 \pm 0.70	0.005	NS	0

Deleted: 3 is the origin of the topsoil stockpile aged two years

Formatted: Font: (Default) +Body (Times New Roman)

Formatted

Formatted: Font: (Default) +Body (Times New Roman)

Formatted

Formatted

Formatted

Formatted

Formatted: Font: (Default) +Body (Times New Roman)

Formatted

Formatted

Formatted: Font: (Default) +Body (Times New Roman)

Formatted

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman), Not Bold

Formatted

Formatted

Formatted

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman), Not Bold

Formatted: Font: (Default) +Body (Times New Roman)

Deleted: Vegetation

Formatted

Deleted: ¶

Deleted: , NS = not significantly different.

Deleted:

Formatted

Deleted:

Formatted

Deleted:

Formatted

Deleted: ¶

Table 3: Cyanobacterial mean (\pm Standard Error) of richness (Margalef's index), evenness (Pielou's index) and diversity (Shannon index) across successional stages. No significant difference in diversity measures was found between stages.

	Bare		Early		Late	
	Mean	SE	Mean	SE	Mean	SE
Richness (d)	167.4	5.344	146.8	5.818	142.8	15.69
Evenness (J')	0.816	0.011	0.791	0.005	0.788	0.027
Diversity (H')	5.977	0.103	5.692	0.064	5.642	0.281

5

Table 4: Permutational analysis of variance (PERMANOVA) of pair-wise comparisons of Bray-Curtis dissimilarity between biocrust stages and bare soil. P(MC) = probability values obtained using 9999 Monte Carlo permutations. A test for homogeneity of multivariate dispersions (PERMDISP) showed no significant differences in variation of spread of samples (pseudo $F=3.8068$, $P(\text{perm}) = 0.068$). Significant pair-wise differences are in bold.

Groups	t	P(perm)	Unique	
			permutations	P(MC)
Bare, Early	2.6216	0.0979	10	0.0107
Bare, Late	2.5742	0.0959	10	0.0120
Early, Late	1.2793	0.0953	10	0.1993

10

15

Deleted: ¶
¶
¶
¶
¶

Formatted: Centered
Formatted: Centered
Formatted: Centered
Formatted: Font: Bold
Formatted: Centered
Formatted: Font: Bold
Formatted: Centered

Table 5: Diversity across sites on a presence absence basis for all seasons and Lake Ifould (salt lake). Different species attributed to a genus (i.e. sp. 1,2,3) have all been separated based on their morphological features and size but could not be positively identified.

Deleted: ¶

Cyanobacterium	SMU 1	SMU 2	SMU 3	T2 Stockpile	Lake Ifould
Aphanothece					x
Brasilonema	x	x	x	x	x
Chroococciopsis		x		x	
Chroococcus sp. 1	x	x	x	x	
Chroococcus sp. 2					x
Gloeocapsa	x	x			x
Leptolyngbya	x	x			x
Microcoleus ethonoplastes					x
Microcoleus paludosus	x	x	x	x	x
Microcoleus sociatus					x
Microcoleus vaginatus	x	x			x
Nostoc commune		x	x	x	x
Nostoc flagelliforme		x			
Nostoc pruniforme		x			
Nostoc sp.	x	x			
Porphyrosiphon sp. 1	x	x	x	x	
Porphyrosiphon sp. 2		x			x
Schizothrix sp. 1		x			x
Schizothrix sp. 2	x	x	x	x	
Schizothrix sp. 3					x
Scytonema sp. 1	x	x	x	x	
Scytonema sp. 2		x			x
Scytonema sp. 3		x			
Scytonema sp. 4					x
Symploca sp. 1	x	x	x	x	x
Symploca sp. 2		x			
Symplocastrum sp. 1	x	x	x	x	x
Symplocastrum sp. 2					x
Species richness	12	21	9	10	18




Figure 1: Image of biocrust sample sites located within the vegetation associations described in Table S1 (supplied, S. Doudle). Site 11 was initially investigated but later discarded as it was a fourth replicate of the Chenopod Shrubland.




Figure 2: (a) SMU 1 (Sites 4, 9 and 10): Type 1–3 Biocrusts on deep calcareous yellow sands (dunes); (b) SMU 2 (Sites 1, 5 and 8): Primarily types 4 and 5 biocrusts on shallow calcareous sandy loam; (c) SMU 3 (Sites 2, 3 and 7): Types 1-5 biocrusts on deep calcareous sandy loam (Photographs by S. Doudle, 2011); (d) different biocrust stages (top) north of Stockpile 19, adjacent to Site 9; (e) Bare, (f) Early and (g) Late stages, also showing biocrust sample already removed from Late stage (Photographs A. Chilton).




Figure 3: Chlorophyll *a* concentration ($\mu\text{g g}^{-1}$ soil) following resurrection (2 weeks) after a desiccation (dry) period of 6 months. Soil Management Unit 1 (SMU 1) was significantly lower than SMU 2–3 ($p = 0.000$). Mean values and standard error of the mean (SEM) displayed where SMU 1 = Sites 4, 9, 10; SMU 2 = Sites 1, 5, 8; SMU 3 = Sites 2, 3, 7; T2 = two-year old Topsoil stockpile (originating from SMU 3). Photosynthetic yield (YII) of photosystem II (PSII) for SMU 1–3 and T2 (Site 6 two-year old topsoil stockpile), displaying mean values and standard error of the mean (SEM). SMU 1 and SMU 2 were significantly different from SMU 3 and T2 ($p = 0.017$), SMU 1 and SMU 2 were not significantly different from each other.

Formatted: English (Australia)


Formatted: English (Australia)

Deleted: 5

5

10

15




5

Figure 4: Cyanobacterial community structure across all sites expressed as a percentage of the total community based on mean richness and abundance scores. N-fixing cyanobacteria contributed to 61% of the community structure.

10

15



Deleted: ¶

¶

Figure 5(a): Cyanobacterial community structure based on indexed abundance and diversity across all sites displayed in an nMDS plot (Bray Curtis similarity). SMU 1-3 refer to key soil management units; SMU 2 appeared to have more range in the clusters compared to the closer groupings of SMU 1 and 3. T2 (TS2Y) is the two-year old topsoil stockpile; note these are clustered more closely. Figure 5(b): Similarities between samples within their SMU's are displayed in a Bray Curtis dendrogram. Black continuous lines show significant differences between samples ($p = 0.05$) and lighter lines indicate that most samples were not significantly different to each other.

Formatted: Justified

Deleted: ¶




Figure 6(a): Relative abundance of cyanobacteria to other bacteria within Bare soil and Early and Late stage biocrusts.

Figure 6(b): Abundance of cyanobacterial genera and groups. Green = Simple filamentous types, Blue = Heterocystic types, Purple = Unicellular. Grey = Unclassified/Other includes chloroplasts.




Figure 7: Non-metric Multidimensional Scaling of Bare soil and Early and Late Stage biocrusts within two dimensions (left) and three dimensions (right).

5


10

15


20

▼ Deleted: ¶

▼ Deleted: ¶




(a)



(b)

Figure 8: Cyanobacterial richness: (a) undisturbed were significant between 50 cm and 0–2 cm in all stockpiles ($p = 0.024$; 20-month $p = 0.016$; 9-month $p = 0.002$) and, (b) stockpiles of different ages were significant between 25 cm and 0–2 cm in the twenty- and nine-month old stockpiles ($p = 0.000$; $p = 0.000$).



Deleted:
(a)

Acknowledgements

This project was funded and supported by Iluka Resources Ltd. and the Jacinth-Ambrosia Rehabilitation team. Thanks to Sam Doudle, Iluka Rehabilitation Specialist for her recognition of the importance of biocrusts at Jacinth-Ambrosia and for initiating this research program. This was a UniQuest project that was carried out at The University of Queensland's School of Agriculture and Food Sciences, Gatton Campus and the University of New South Wales. Especially appreciated was the strong support, project advice and laboratory resources provided by Assoc. Prof Vic Galea and Katherine Raymont. Special thanks to Dr Glenn McGregor for technical advice, assistance with cyanobacterial identification and the use of specialist microscopes and facilities at the Ecosciences Precinct (Qld), as well as report editing. Special thanks to David Tongway who assisted in Landscape Function Analysis field work, and two anonymous reviewers who provided valuable insights and editing to greatly improve this manuscript.

References

- Aboal, M., Werner, O., García-Fernández, M. E., Palazón, J. A., Cristóbal, J. C. and Williams, W.: Should ecomorphs be conserved? The case of *Nostoc flagelliforme*, an endangered extremophile cyanobacteria, *J. Nat. Conserv.*, 30(Supplement C), 52–64, doi:10.1016/j.jnc.2016.01.001, 2016.
- Agrawal, S. C. and Singh, V.: Viability of dried filaments, survivability and reproduction under water stress, and survivability following heat and UV exposure in *Lyngbya martensiana*, *Oscillatoria agardhii*, *Nostoc calcicola*, *Horridium fluitans*, *Spirogyra* sp. and *Vaucheria geminata*, *Folia Microbiol. (Praha)*, 47(1), 61–67, doi:10.1007/BF02818567, 2002.
- Barnes, J. D., Balaguer, L., Manrique, E., Elvira, S. and Davison, A. W.: A reappraisal of the use of DMSO for the extraction and determination of chlorophylls a and b in lichens and higher plants, *Environ. Exp. Bot.*, 32(2), 85–100, doi:10.1016/0098-8472(92)90034-Y, 1992.
- Belnap, J. and Eldridge, D.: Disturbance and Recovery of Biological Soil Crusts, in *Biological Soil Crusts: Structure, Function, and Management*, pp. 363–383, Springer, Berlin, Heidelberg., 2001.
- Belnap, J. and Gardner, J. S.: Soil microstructure in soils of the Colorado Plateau: the role of the cyanobacterium *Microcoleus vaginatus*, *Gt. Basin Nat.*, 40–47, 1993.
- Belnap, J. and Gillette, D. A.: Vulnerability of desert biological soil crusts to wind erosion: the influences of crust development, soil texture, and disturbance, *J. Arid Environ.*, 39(2), 133–142, doi:10.1006/jare.1998.0388, 1998.
- Billi, D. and Potts, M.: Life and death of dried prokaryotes, *Res. Microbiol.*, 153(1), 7–12, doi:10.1016/S0923-2508(01)01279-7, 2002.
- Bowker, M. A.: Biological Soil Crust Rehabilitation in Theory and Practice: An Underexploited Opportunity, *Restor. Ecol.*, 15(1), 13–23, doi:10.1111/j.1526-100X.2006.00185.x, 2007.

Deleted: ¶

¶

- Bowker, M. A., Maestre, F. T., Eldridge, D., Belnap, J., Castillo-Monroy, A., Escolar, C. and Soliveres, S.: Biological soil crusts (biocrusts) as a model system in community, landscape and ecosystem ecology, *Biodivers. Conserv.*, 23(7), 1619–1637, doi:10.1007/s10531-014-0658-x, 2014.
- Bristol, B. M.: ON THE RETENTION OF VITALITY BY ALGÆ FROM OLD STORED SOILS., *New Phytol.*, 18(3-4),
5 92–107, 1919.
- Büdel, B., Darienko, T., Deuschewitz, K., Dojani, S., Friedl, T., Mohr, K. I., Salisch, M., Reisser, W. and Weber, B.: Southern African biological soil crusts are ubiquitous and highly diverse in drylands, being restricted by rainfall frequency, *Microb. Ecol.*, 57(2), 229–247, doi:10.1007/s00248-008-9449-9, 2009.
- Büdel, B., Williams, W. J. and Reichenberger, H.: Annual net primary productivity of a cyanobacteria-dominated biological
10 soil crust in the Gulf Savannah, Queensland, Australia, *Biogeosciences*, 15(2), 491, 2018.
- Carrick, P. J. and Krüger, R.: Restoring degraded landscapes in lowland Namaqualand: Lessons from the mining experience and from regional ecological dynamics, *J. Arid Environ.*, 70(4), 767–781, doi:10.1016/j.jaridenv.2006.08.006, 2007.
- Chiquoine, L. P., Abella, S. R. and Bowker, M. A.: Rapidly restoring biological soil crusts and ecosystem functions in a severely disturbed desert ecosystem, *Ecol. Appl.*, 26(4), 1260–1272, doi:10.1002/15-0973, 2016.
- 15 Codony, F., Miranda, A. M. and Mas, J.: Persistence and proliferation of some unicellular algae in drinking water systems as result of their heterotrophic metabolism, *Water Sa*, 29(1), 113–116, 2003.
- Delgado-Baquerizo, M., Morillas, L., Maestre, F. T. and Gallardo, A.: Biocrusts control the nitrogen dynamics and microbial functional diversity of semi-arid soils in response to nutrient additions, *Plant Soil*, 372(1–2), 643–654, doi:10.1007/s11104-013-1779-9, 2013.
- 20 Dodds, W. K., Gudder, D. A. and Mollenhauer, D.: The ecology of *Nostoc*, *J. Phycol.*, 31(1), 2–18, 1995.
- Doherty, K. D., Antoninka, A. J., Bowker, M. A., Ayuso, S. V. and Johnson, N. C.: A Novel Approach to Cultivate Biocrusts for Restoration and Experimentation, *Ecol. Restor.*, 33(1), 13–16, 2015.
- Dominati, E., Patterson, M. and Mackay, A.: A framework for classifying and quantifying the natural capital and ecosystem services of soils, *Ecol. Econ.*, 69(9), 1858–1868, doi:10.1016/j.ecolecon.2010.05.002, 2010.
- 25 Doudle, S., Williams, W. and Galea, V.: Improving rehabilitation outcomes using biocrusts, 8th Int. Heavy Miner. Conf. 2011, 85–97, 2011.
- Elbert, W., Weber, B., Burrows, S., Steinkamp, J., Büdel, B., Andreae, M. O. and Pöschl, U.: Contribution of cryptogamic covers to the global cycles of carbon and nitrogen, *Nat. Geosci.*, 5(7), 459–462, doi:10.1038/ngeo1486, 2012.
- Eldridge, D. J. and Leys, J. F.: Exploring some relationships between biological soil crusts, soil aggregation and wind
30 erosion, *J. Arid Environ.*, 53(4), 457–466, doi:10.1006/jare.2002.1068, 2003.
- Esmarch, F.: Investigations of the distribution of Cyanophyceae on and in various soils, 1914.
- Fiore, M. F., Sant’Anna, C. L., Azevedo, M. T. de P., Komárek, J., Kaštovský, J., Sulek, J. and Lorenzi, A. A. S.: The Cyanobacterial Genus *Brasilonema*, Gen. Nov., a Molecular and Phenotypic Evaluation1, *J. Phycol.*, 43(4), 789–798, doi:10.1111/j.1529-8817.2007.00376.x, 2007.

- Flores, E. and Herrero, A.: Compartmentalized function through cell differentiation in filamentous cyanobacteria, *Nat. Rev. Microbiol.*, 8(1), 39–50, 2010.
- Fujita, Y. and Nakahara, H.: Variations in the microalgal structure in paddy soil in Osaka, Japan: comparison between surface and subsurface soils, *Limnology*, 7(2), 83–91, doi:10.1007/s10201-006-0167-z, 2006.
- 5 Gillieson, D., Wallbrink, P. and Cochrane, A.: Vegetation change, erosion risk and land management on the Nullarbor Plain, Australia, *Environ. Geol.*, 28(3), 145–153, doi:10.1007/s002540050087, 1996.
- Harris, J. A.: Measurements of the soil microbial community for estimating the success of restoration, *Eur. J. Soil Sci.*, 54(4), 801–808, doi:10.1046/j.1351-0754.2003.0559.x, 2003.
- Hou, B. and Warland, I.: Heavy mineral sands potential of the Eucla Basin in South Australia—a world-class palaeobeach
10 placer province, *Mesa J.*, 37, 4–12, 2005.
- Hu, C., Liu, Y., Zhang, D., Huang, Z. and Paulsen, B. S.: Cementing mechanism of algal crusts from desert area, *Chin. Sci. Bull.*, 47(16), 1361–1368, doi:10.1360/02tb9301, 2002.
- Huang, T.-C., Chow, T.-J. and Hwang, I.-S.: The cyclic synthesis of the nitrogenase of *Synechococcus* RF-1 and its control at the transcription level, *FEMS Microbiol. Lett.*, 50(2–3), 127–130, 1988.
- 15 Jasper, D. A.: Beneficial Soil Microorganisms of the Jarrah Forest and Their Recovery in Bauxite Mine Restoration in Southwestern Australia, *Restor. Ecol.*, 15, S74–S84, doi:10.1111/j.1526-100X.2007.00295.x, 2007.
- Jones, C. G., Lawton, J. H. and Shachak, M.: *Organisms as Ecosystem Engineers*, in *Ecosystem Management*, pp. 130–147, Springer, New York, NY., 1994.
- Kaplan-Levy, R. N., Hadas, O., Summers, M. L., Rucker, J. and Sukenik, A.: Akinetes: dormant cells of cyanobacteria, in
20 *Dormancy and resistance in harsh environments*, pp. 5–27, Springer., 2010.
- Kidron, G. J., Barinova, S. and Vonshak, A.: The effects of heavy winter rains and rare summer rains on biological soil crusts in the Negev Desert, *CATENA*, 95(Supplement C), 6–11, doi:10.1016/j.catena.2012.02.021, 2012.
- Lipman, C. B.: The successful revival of *Nostoc commune* from a herbarium specimen eighty-seven years old, *Bull. Torrey Bot. Club*, 664–666, 1941.
- 25 Lukešová, A.: Soil Algae in Brown Coal and Lignite Post-Mining Areas in Central Europe (Czech Republic and Germany), *Restor. Ecol.*, 9(4), 341–350, doi:10.1046/j.1526-100X.2001.94002.x, 2001.
- Maestre, F. T., Castillo-Monroy, A. P., Bowker, M. A. and Ochoa-Hueso, R.: Species richness effects on ecosystem multifunctionality depend on evenness, composition and spatial pattern, *J. Ecol.*, 100(2), 317–330, doi:10.1111/j.1365-2745.2011.01918.x, 2012.
- 30 McCutcheon, J., Wilson, S. A. and Southam, G.: Microbially accelerated carbonate mineral precipitation as a strategy for in situ carbon sequestration and rehabilitation of asbestos mine sites, *Environ. Sci. Technol.*, 50(3), 1419–1427, 2016.
- Moore, G. T. and Karrer, J. L.: A subterranean algal flora, *Ann. Mo. Bot. Gard.*, 6(4), 281–307, 1919.
- Rao, B., Liu, Y., Lan, S., Wu, P., Wang, W. and Li, D.: Effects of sand burial stress on the early developments of cyanobacterial crusts in the field, *Eur. J. Soil Biol.*, 48(Supplement C), 48–55, doi:10.1016/j.ejsobi.2011.07.009, 2012.

- Reisser, W.: The hidden life of algae underground, in *Algae and Cyanobacteria in Extreme Environments*, pp. 47–58, Springer., 2007.
- Rossi, F., Mugnai, G. and Philippis, R. D.: Complex role of the polymeric matrix in biological soil crusts, *Plant Soil*, 1–16, doi:10.1007/s11104-017-3441-4, 2017.
- 5 Shaw, S. L., Chisholm, S. W. and Prinn, R. G.: Isoprene production by *Prochlorococcus*, a marine cyanobacterium, and other phytoplankton, *Mar. Chem.*, 80(4), 227–245, 2003.
- Skinner, S. and Entwisle, T.: Non-marine algae of Australia: 1. Survey of colonial gelatinous blue-green macroalgae (Cyanobacteria), *Teloepa*, 9(3), 573–599, doi:10.7751/teloepa20024003, 2002.
- Stal, L. J.: Microphytobenthos, their Extracellular Polymeric Substances, and the Morphogenesis of Intertidal Sediments, *10 Geomicrobiol. J.*, 20(5), 463–478, doi:10.1080/713851126, 2003.
- Tomaselli, L. and Giovannetti, L.: Survival of diazotrophic cyanobacteria in soil, *World J. Microbiol. Biotechnol.*, 9(1), 113–116, 1993.
- Tongway, D. and Hindley, N.: Landscape function analysis: a system for monitoring rangeland function, *Afr. J. Range Forage Sci.*, 21(2), 109–113, doi:10.2989/10220110409485841, 2004.
- 15 Tongway, D. J.: Soil and landscape processes in the restoration of rangelands., *Rangel. J.*, 12(1), 54–57, doi:10.1071/rj9900054, 1990.
- Tongway, D. J. and Ludwig, J. A.: Rehabilitation of Semiarid Landscapes in Australia. I. Restoring Productive Soil Patches, *Restor. Ecol.*, 4(4), 388–397, doi:10.1111/j.1526-100X.1996.tb00191.x, 1996.
- Ullmann, I. and Büdel, B.: Ecological determinants of species composition of biological soil crusts on a landscape scale, *20 Biol. Soil Crusts Struct. Funct. Manag.*, 203–213, 2001.
- Vaccarino, M. A. and Johansen, J. R.: *Brasilonema Angustatum* Sp. Nov. (nostocales), a New Filamentous Cyanobacterial Species from the Hawaiian Islands1, *J. Phycol.*, 48(5), 1178–1186, doi:10.1111/j.1529-8817.2012.01203.x, 2012.
- Vishnivetskaya, T. A., Spirina, E. V., Shatilovich, A. V., Erokhina, L. G., Vorobyova, E. A. and Gilichinsky, D. A.: The resistance of viable permafrost algae to simulated environmental stresses: implications for astrobiology, *Int. J. Astrobiol.*, *25* 2(3), 171–177, doi:10.1017/S1473550403001575, 2003.
- Weber, B., Bowker, M., Zhang, Y. and Belnap, J.: Natural Recovery of Biological Soil Crusts After Disturbance, in *Biological Soil Crusts: An Organizing Principle in Drylands*, pp. 479–498, Springer, Cham., 2016.
- Wellburn, A. R.: The Spectral Determination of Chlorophylls a and b, as well as Total Carotenoids, Using Various Solvents with Spectrophotometers of Different Resolution, *J. Plant Physiol.*, 144(3), 307–313, doi:10.1016/S0176-1617(11)81192-2, *30* 1994.
- Williams, W. J. and Büdel, B.: Species diversity, biomass and long-term patterns of biological soil crusts with special focus on Cyanobacteria of the *Acacia aneura* Mulga Lands of Queensland, Australia, *Algol. Stud.*, 140(1), 23–50, 2012.
- Williams, W. J. and Eldridge, D. J.: Deposition of sand over a cyanobacterial soil crust increases nitrogen bioavailability in a semi-arid woodland, *Appl. Soil Ecol.*, 49, 26–31, doi:10.1016/j.apsoil.2011.07.005, 2011.

Williams, W. J., Büdel, B., Reichenberger, H. and Rose, N.: Cyanobacteria in the Australian northern savannah detect the difference between intermittent dry season and wet season rain, *Biodivers. Conserv.*, 23(7), 1827–1844, doi:10.1007/s10531-014-0713-7, 2014.

5 Zhao, Y., Zhu, Q., Li, P., Zhao, L., Wang, L., Zheng, X. and Ma, H.: Effects of artificially cultivated biological soil crusts on soil nutrients and biological activities in the Loess Plateau, *J. Arid Land*, 6(6), 742–752, doi:10.1007/s40333-014-0032-6, 2014.