

Microbial Biobanking

Cyanobacteria-rich topsoil facilitates mine rehabilitation

Wendy Williams¹, Angela Chilton², Mel Schneemilch¹, Stephen Williams¹, Brett Neilan³, Colin Driscoll⁴

- 5
1. School of Agriculture and Food Sciences, The University of Queensland, Gatton Campus 4343 Australia
 2. Australian Centre for Astrobiology and School of Biotechnology and Biomolecular Sciences, University of New South Wales, Sydney, NSW, 2052, Australia
 3. School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW, 2308, Australia
 4. Hunter Eco, PO Box 1047, Toronto, NSW, 2283

10 *Correspondence to:* wendy.williams@uq.edu.au

Dear Associate Editor,

We appreciate your additional suggestions and editing for this manuscript and have addressed each area detailed below. We thank you for your time and input to improve this submission.

Kind regards,

15 Wendy Williams et al.

Editor:

INTRODUCTION

I suggested to work a little bit more on the introduction as it is still not fully connected.

20 Response:

We have revised the introduction and focused on your comments to connect the mining rehabilitation process and problems with the role of biocrusts.

Editor:

25 M&M

Was microscopy done based on same samples used for physicochemical analysis?
not clear in the text.

Response:

We have clarified this in the text

30

Editor:

Formatted: Left

RESULTS

Check how significant differences are presented in tables and figures

Figure 8 still not clear, what are you comparing. Moreover, all treatments should include letters.

Caption should be something like:

5 What mean age on undisturbed samples means?

Response:

We have clarified Figure 8 in the positioning of the graphs (stockpile first) and revised the caption to reflect this. (Mean) age is just the comparison to the stockpile age so the reader knows where the sample came from (i.e. adjacent to which stockpile). However we have removed the word mean as it was not relevant.

10

Editor:

Based on figure, section 3.4 still confusing for an external reader, I would like to see a clearer version with a well define story line

Response:

15 We have made further changes to this section to make it more succinct.

Editor:

DISCUSSION

20 I also suggest changing first subsection to include the link between cyanobacterial community structure, nutrient cycling and soil stabilization, as this last part is missing within the discussion

Response:

We have further revised the discussion (in full) and linked the community structure to stabilisation and nutrient cycling (refer to paragraph 1 in the discussion below):

25 “This research has demonstrated cyanobacteria to be a key component of soil microbial communities at J-A. These were compositionally diverse topsoil microbiomes that substantially contributed to the Myall-chenopod landscape. We had hypothesised that cyanobacteria would be central to soil micro-processes, and this was strongly supported by extensive species richness and diversity values. At J-A cyanobacteria contributed to soil structure and function during the early developmental stages of the biocrust. Photosynthesis drove the productivity and growth of the biocrust that initiated carbon and nitrogen cycling and resulted in increases in soil nutrient concentrations right where vascular plants might use them. The results have demonstrated how these micro-processes provide a strong foundation for the restoration of soil function.

30 Similarly, in southwestern Queensland and northern Australia cyanobacterial species richness was strongly linked to increased bioavailable nitrogen and carbon uptake (Büdel et al., 2018; Williams et al., 2018; Williams and Eldridge, 2011). Elsewhere, multiple studies have demonstrated the high value of biocrust attributes as drivers of soil micro-processes that restore soil function (e.g. (Barger et al., 2016; Belnap and Eldridge, 2001; Bowker et al., 2014; Büdel et al., 2009; Chilton et

al., 2017; Chiquoine et al., 2016; Weber et al., 2016).”

Editor:

Conclusions:

5 I also recommended to rewrote in order to ask main objectives presented:

A general conclusion based on your main objective "to determine whether shallow biobanks of cyanobacterial-enriched topsoil would facilitate the recovery of essential soil microprocesses when re-spread following mine disturbance"

and 2 specific conclusion based on specific objectives:

10

1) Define the cyanobacterial community structure with a special focus on species that drive early colonization, nutrient cycling and soil stabilization

2) to examine the effects of stockpiling topsoil on cyanobacterial resilience to crushing and burial and their recovery following spreading of topsoil back across mined land.

15

Response:

Conclusion has been fully revised to reflect comments.

20

25

30

Microbial Biobanking

Cyanobacteria-rich topsoil facilitates mine rehabilitation

Wendy Williams¹, Angela Chilton², Mel Schneemilch¹, Stephen Williams¹, Brett Neilan³, Colin Driscoll⁴

1. School of Agriculture and Food Sciences, The University of Queensland, Gatton Campus 4343 Australia
2. Australian Centre for Astrobiology and School of Biotechnology and Biomolecular Sciences, University of New South Wales, Sydney, NSW, 2052, Australia
3. School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW, 2308, Australia
4. Hunter Eco, PO Box 1047, Toronto, NSW, 2283

Correspondence to: wendy.williams@uq.edu.au

Abstract

Restoration of soils post-mining requires key solutions to complex issues where the disturbance of topsoil incorporating soil microbial communities can result in a modification to ecosystem function. This research was in collaboration with Iluka Resources at Jacinth-Ambrosia (J-A) mineral sand mine located in a semi-arid chenopod shrubland in southern Australia. At J-A, assemblages of microorganisms and microflora inhabit at least half of the soil surfaces and are collectively known as biocrusts. This research encompassed a polyphasic approach to soil microbial community profiling focused on 'biobanking' viable cyanobacteria in topsoil stockpiles to facilitate rehabilitation. We found that cyanobacterial communities were compositionally diverse topsoil microbiomes. There was no significant difference in cyanobacterial community structure across soil types. As hypothesised, cyanobacteria were central to soil micro-processes, strongly supported by species richness and diversity. Cyanobacteria were a significant component of all three successional stages with 21 species identified from ten sites. Known nitrogen-fixing cyanobacteria *Symploca*, *Scytonema*, *Porphyrosiphon*, *Brasilonema*, *Nostoc* and *Gloeocapsa* comprised more than 50% of the species richness at each site and 61% of the total community richness. In the first study of its kind, we have described the response of cyanobacteria to topsoil stockpiling at various depths and ages. Cyanobacteria are moderately resilient to stockpiling at depth and over time, with average species richness greatest in the top 10 cm of the stockpiles of all ages and more viable within the first six weeks, indicating potential for biocrust re-establishment. In general, the resilience of cyanobacteria to burial in topsoil stockpiles in both the short and long term was significant, however in an arid environment recolonization and community diversity could be impeded by drought. Biocrust re-establishment during mine rehabilitation relies on the role of cyanobacteria as a means of early soil stabilisation. At J-A mine operations do not threaten the survival of any of the organisms we studied. Increased cyanobacterial biomass is likely to be a good indicator and reliable metric for the reestablishment of soil micro-processes.

1.0 Introduction

Following the destruction of the soil profile in a post-mining landscape there is the critical need to restore ecological integrity to the system. Mine rehabilitation is a complex process that involves many levels of understanding of difficult issues relating to ecosystem function. Harsh disturbance disrupts the spatial structure of soil microbial communities. Not the least is the removal or burial of bioactive soils that will have knock-on effects for rehabilitation efforts such as nutrient cycling and plant reestablishment (Jasper, 2007; Tongway and Ludwig, 1996). Microorganisms are critical components of soils that drive assorted micro-processes and impact soil ecosystem function on several levels. Successful ecological restoration of arid mining sites relies on a holistic approach where microbial recolonization can serve as an indicator of the integrity of the wider ecosystem (Tongway, 1990).

Moved (insertion) [7]

Restoration in arid landscapes includes the reestablishment of surficial crusts that develop as a protective skin across the bare soil or stony interspaces between plants. These encrusted surfaces can be physical, chemical or biological (microbial) in nature. Well established biological crusts (biocrusts) are intricately interwoven and structured, high-functioning communities, and variable in composition (incorporating cyanobacteria, algae, lichens, mosses, liverworts, micro-fauna and bacteria). Biocrusts are a significant asset to arid soil ecosystems providing a protective, nutrient-rich layer closely integrated into the soil surface (Delgado-Baquerizo et al., 2013; Maestre et al., 2012).

Moved up [7]: Microorganisms are critical components of soils that drive assorted micro-processes and impact soil ecosystem function on several levels.

Mining severely disrupts and compromises the structure and function of biocrusts. The recognition of the absolute importance of microbial communities in the topsoil has led to the accepted practice of stripping and storing in stockpiles as a form of "biobanking" for redistribution at the cessation of mining. During this process the crushing and burial of biocrusts results in the inability of crust organisms to photosynthesise due to the lack of light. Massive soil disturbance results in a loss of structure, resources and often has long lasting effects on energy transfers, soil stability, nutrient cycling and surface hydrology (Bowker, 2007; Tongway and Hindley, 2004). Physical disturbance can profoundly disrupt biocrust integrity, composition and physiological function where the impact is governed by site characteristics, severity, frequency and timing (Belnap and Eldridge, 2001). Biocrust burial as a result of environmental stress and disturbance can result in serious impacts on biocrust microorganisms' viability and function. For example, grazing and drought caused significant declines in cyanobacterial richness and abundance which resulted in a reduction in soil nutrient concentrations (Rao et al., 2012; Williams and Eldridge, 2011).

Following disturbance, restoration and regrowth of biocrusts can take place unassisted, seasonally driven generally over many years (Belnap and Eldridge, 2001; Belnap and Gillette, 1998). Should biocrust organisms remain inactive while they are wet, cell death and decomposition commonly occurs (Kidron et al., 2012; Rao et al., 2012). Nevertheless, in dry conditions,

Moved (insertion) [8]

5 cyanobacteria and algae are known to remain desiccated and viable for millions of years (Vishnivetskaya et al., 2003). Alternatively, assisted biocrust restoration places emphasis on the recovery of ecosystem function and necessarily addresses environmental constraints. This incorporates the knowledge of 'potential condition' based on experience with sites of ecological similarity that have undergone disturbance and recovery (Bowker, 2007). Biocrust recovery can be altered by dust deposition, fire and climatic conditions (Weber et al., 2016). When biocrusts recover naturally soil properties change. For example, in Southern African and Spanish rangelands an incremental accumulation of soil nutrients, organic matter and a build-up of silt and clay lead to the development of a resilient and multi-functional biocrust (Büdel et al., 2009; Maestre et al., 2012; Weber et al., 2016).

10 Restoration of ecosystem function post-disturbance requires an appreciation of the dynamic functional status of the landscape prior to disturbance (Tongway and Ludwig, 1996), as well as an understanding of the net accumulative effects of disturbance on the components of the system. On the micro-scale, cyanobacterial species richness contributes to soil ecosystem function through micro-processes including carbon fixation through photosynthesis, atmospheric nitrogen fixation in a biological-available form, micro-nutrient breakdown and release, soil particle cohesion, regulation of moisture and soil surface structure
15 (Delgado-Baquerizo et al., 2013; Elbert et al., 2012; Hu et al., 2002; Maestre et al., 2012 and others). Consequently, it is necessary to appreciate the micro-processes that will assist in the restoration of soil function and to monitor recovery along the way.

20 Cyanobacterial re-establishment is a key indicator of early soil surface re-stabilisation, regulation of soil moisture and the balancing of soil carbon and nitrogen (Elbert et al., 2012). Cyanobacteria in arid landscapes are exceptionally well-adapted to desiccation. Their polysaccharide sheaths and EPS production perform a vital role in maintaining cyanobacterial cell integrity, exchange of information and absorption of water during rehydration (Rossi et al., 2017). EPS has adhesive properties that binds non-aggregated soil particles into a protective encrusted surface that reduces the destructive impacts of wind and water (Eldridge and Leys, 2003; Rossi et al., 2017). Cyanobacterial biofilms provide stabilisation of initially disturbed surfaces that
25 pave the way for diverse microbial communities, and form bioactive crust-like layers assimilated into the soil (e.g. Büdel et al., 2009; Rossi et al., 2017; Bowker et al., 2014). As biocrusts develop in structural complexity, the diversity of organisms is regulated by water infiltration, temperature, light and additional disturbance (Belnap and Eldridge, 2001; Büdel et al., 2009; Elbert et al., 2012).

30 Research into biocrust disturbance with a focus on recovery post-mining is rare. In the Namaqualand arid lands (Namibia, South Africa) low rainfall and high winds impact the rehabilitation of degraded lands following diamond mining and grazing (Carrick and Krüger, 2007). These researchers found that cyanobacteria and non-vascular plants that form a living and protective surface crust were crucial to surface stabilisation. Jasper, (2007) also recognised the importance of soil microbial communities including cyanobacteria in post-mine rehabilitation in the Jarrah forests of south-western Australia. In the Czech

Moved up [8]: Following disturbance, restoration and regrowth of biocrusts can take place unassisted, seasonally driven generally over many years (Belnap and Eldridge, 2001; Belnap and Gillette, 1998). Should biocrust organisms remain inactive while they are wet, cell death and decomposition commonly occurs (Kidron et al., 2012; Rao et al., 2012). Nevertheless, in dry conditions, cyanobacteria and algae are known to remain desiccated and viable for millions of years (Vishnivetskaya et al., 2003). Alternatively, assisted biocrust restoration places emphasis on the recovery of ecosystem function and necessarily addresses environmental constraints. This incorporates the knowledge of 'potential condition' based on experience with sites of ecological similarity that have undergone disturbance and recovery (Bowker, 2007). Biocrust recovery can be altered by dust deposition, fire and climatic conditions (Weber et al., 2016). When biocrusts recover naturally soil properties change. For example, in Southern African and Spanish rangelands an incremental accumulation of soil nutrients, organic matter and a build-up of silt and clay lead to the development of a resilient and multi-functional biocrust (Büdel et al., 2009; Maestre et al., 2012; Weber et al., 2016).

Republic and Germany chrono-sequential studies of old brown coal mine sites found in younger sites that green algal biofilms and a diverse range of cyanobacteria initiated the rehabilitation of the soils (Lukešová, 2001). In serpentinite mine tailings (New South Wales, Australia), McCutcheon et al., (2016) showed filamentous cyanobacteria accelerated carbonate mineral precipitation and stabilised the tailings. They demonstrated cyanobacteria had the capacity to adsorb magnesium while acting as a nucleation site and sequestering carbon. In our current study, preliminary research identified that in undisturbed chenopod shrublands at the edge of the Nullarbor Plain (South Australia) biocrusts cover more than 45% of the soil surface between the grass plants and post-mining rehabilitation needs to take their role into account (Doudle et al., 2011). It follows that there is a real need for focus on practical approaches that contribute to the restoration of soil function and measure relevant aspects of success through soil microbial communities and biocrust reestablishment, especially cyanobacteria (for example: Setyawan et al., 2016; Mazor et al., 1996; Fischer et al., 2014; Chiquoine et al., 2016; Doherty et al., 2015; Harris, 2003; Tongway and Hindley, 2004; Zhao et al., 2014).

To our knowledge, the effects of topsoil stockpiling on biocrust organisms such as cyanobacteria, and their recovery time, following topsoil spreading, has not previously been investigated. In this research we focused on the cyanobacterial component of the biocrusts. This was in keeping with the mining framework in the ongoing development of informed rehabilitation plans that focuses on improved long-term outcomes. Specifically, we sought to determine whether shallow biobanks of cyanobacterial-enriched topsoil would facilitate the recovery of essential soil microprocesses when re-spread following mine disturbance. The overall aims of the microbial biobanking research program were to: (a) define the cyanobacterial community structure applying a polyphasic approach with a special focus on species that drive early colonisation, nutrient cycling and soil stabilisation; and, (b) to examine the effects of stockpiling topsoil on cyanobacterial resilience to crushing and burial and their recovery following spreading of topsoil back across mined land. We hypothesised that cyanobacterial survival within a topsoil stockpile would reduce with both depth within the stockpile and elapsed time before topsoil re-spreading.

2.0 Methods

2.1 Background and site description

Jacinth-Ambrosia (J-A) heavy mineral sand mine is located on the eastern edge of the Nullarbor Plain, South Australia, across the boundary of two Regional Reserves within the Eucla Basin region. The climate is semi-arid with a mean rainfall of 185 mm, mean maximum temperature of 27.6°C and minimum of 12.1°C (further detail provided in supplementary Figure S1, www.bom.gov.au). Tertiary sediments deposited in marine and terrestrial settings and the soil distribution of the area reflects the geological history, with at least five marine transgression and regression events depositing 40-50 m of sediments (Hou and Warland, 2005). The landscape is broadly undulating with low open woodlands that have a shrub understorey with chenopod shrub lands as well as dune fields that consist of parallel dunes and inter-dune swales (Doudle et al., 2011; Gillieson et al., 1996). Prior to mining disturbance, the landscape is superficially homogenous chenopod-dominated vegetation but functionally

patchy on the fine scale where the soil surfaces are extensively colonised by biocrusts. At J-A, biocrusts primarily consist of lichens, mosses, and cyanobacteria that cover around 45% of the landscape surfaces, equating to 2,000 hectares of the mining lease (Doudle et al., 2011).

5 The biocrusts at J-A had been previously classified into three successional stages representative of the five biocrust types found growing across the landscape (Table S1, S2) (Doudle et al., 2011). Types 1–2 are light coloured, patchy, thin, and fragile cyanobacterial crusts corresponding to early stages of development; Type 3 are well established cyanobacterial crusts with establishment of some mosses and lichens corresponding to intermediate stages of development; Types 4–5 biocrusts are well established with cyanolichens and/or green algal lichens and mosses corresponding with late stages of development (additional
10 descriptions available in supplementary Table S1). As well as In this study the term biocrust covers whole crust samples that incorporated lichens cyanobacteria and mosses in varying proportions; however the cyanobacterial component of this crust was the focus in terms of our polyphasic approach to community structure, succession and its biophysiochemical properties.

15 At J-A the landscape has been characterised into three distinct soil types that were associated with vegetation communities identified as soil management units (Doudle et al., 2011; Hou and Warland, 2005). Site vegetation associations are described as follows: SMU 1 – deep calcareous yellow sands associated with dune ridges; Red Mallee: *Eucalyptus oleosa* ssp. *oleosa* = open Mallee/Myall woodland; SMU 2 – shallow calcareous sandy loams; Chenopod Shrubland: *Maireana sedifolia* and *Atriplex vesicaria*; SMU 3 – Western Myall: deep calcareous sandy loam; *Acacia papyrocarpa* *Maireana sedifolia* = open Myall woodland Site 1 occurs in a transition between SMU 2 and SMU 3 but was treated as most like SMU 2 (also see Table
20 S1). In the first place, sample site locations (Fig. 1) were selected based on these soil management units (SMU) and a two-year old stockpile (Table S1). Secondly, sites within these parameters (SMU) were selected for the subsequent detailed studies of cyanobacterial succession and its resilience to longer-term stockpiling (Fig.2 and Section 2.4).

25 On two occasions (July 2011 and March 2012) we sampled the biocrusts across the three main SMU sites (10 sites, see Fig. 1), in 2011 we also sampled a dry salt lake (Lake Ifould) to provide information on cyanobacterial species adapted to saline conditions, similar in nature to ground water used in mine operations.

30 In the process of mining and preparation of the J-A site for future rehabilitation, topsoil (0–100 mm) is stored in low stockpiles, generally less than two metres in depth where stockpiles are comprised of topsoil sourced from only one vegetation type. Later stockpiles will be returned to the surface of mined areas in the rehabilitation process, and the cyanobacterial activity therein must be quantified to enable educated planning and decision making regarding biocrust re-establishment. If low levels of biocrust organisms are detected below the top few centimetres, the addition of propagated biocrust organisms (e.g. cyanobacteria) to returned topsoil may be warranted. Activity and species richness within the stockpiled top soils may also vary with age and this may also influence the establishment of vegetation in the rehabilitation process.

2.2 Biophysical characteristics of biocrusts and cyanobacteria

2.2.1 Field Sampling

Preliminary identification of biocrust types had been determined by Doudle et al., (2011) and provided the baseline data for biocrust sampling from Sites 1–10 within J-A (Fig. 1). The sites selected encompassed the three SMU and five crust types (Table S1). All sites are of naturally occurring biocrusts except for Site 6 which is from a two-year-old topsoil stockpile regarded as an early (Type 1) biocrust representing recovery two years post disturbance (Fig. 2a-c). Within each site, eight 10 cm diameter samples were selected at random and removed to a depth of 1 cm using a metal scraper (n=80), air dried (>40°C), and stored in Petri dishes. Each Petri dish contained approximately 80 g of crust. The samples were packed to avoid crust disruption and transported to The University of Queensland's Central Analytical Laboratory at Gatton.

2.2.2 Biocrust biophysiochemical properties

For each site, about half the sample was removed from each Petri dish and fine sieved (1.70 mm). Duplicate sub-samples (~2 g) were analysed for total C and N and C:N ratio using a high temperature digestion in a vario MACRO Elemental Analyser (Elementar) (n=6 per SMU). Duplicate samples (10g) for the purpose of analysing soil pH and electrical conductivity (EC) were prepared using a 1:5 (soil to water) ratio and shaken for one hour. Following shaking, samples were left to stand for 30 min and EC measured using a Crison Conductivity Meter 525. The sample was mixed again, and pH was measured with a TPS pH meter MC-80 using an ionode IJ44C electrode. The remaining half of the Petri dish was used to determine Chlorophyll *a* concentration of the biocrusts. Following resurrection (by moistening) the crust was lightly homogenised and a 5 g sample used at a 1:5 ratio of (dry weight) biocrust to Dimethyl sulfoxide (DMSO) (Barnes et al., 1992) with samples placed in a warm bath (65°C) for a two-hour dark extraction, followed by centrifuging for five minutes (5000 g RCF). Chlorophyll *a* concentration was determined using Wellburn's, (1994) equations.

Prior to the use of the samples for analysis a pocket penetrometer (8 mm foot) was used to determine the compressive strength (kg cm²) of the dry intact biocrusts samples. Overall the crust thickness was < 0.5 cm. Each sample (10 cm diameter x 2 cm depth) was placed on a solid surface and a total of twelve measurements (3 readings x 4 reps) were taken for each site. The measurement was taken at the point when the crust was broken, and the foot penetrated the crust surface.

At J-A we measured photosynthetic performance (recorded as yield, YII) of the biocrusts using a pulse-amplitude modulated (PAM) fluorometer (Pocket PAM; Gademann Instruments, Germany). The goal was to demonstrate photosynthetic yield (YII) indicative of active growth of the biocrusts, using the detection of chlorophyll fluorescence from photosystem II (PSII). The sensor was placed onto the biocrust and once started, a series of short pulses of excitation light at high intensity that is amplified

resulting in a brief closure of PSII and the measurement of fluorescence yield based on the Genty parameter which is the quantum yield (YII) of the charge separation of PSII (Genty et al., 1989) and recorded on a scale of 0–1 for all photosynthesis. In the field this process was completed at least six times for each sample (6 x 4 reps per site).

- 5 One-way ANOVA with Tukey post-hoc tests were run to test for significant differences between SMU's and total C and N and C:N ratio (Minitab 18). Chlorophyll a and YII were independently tested with a one-way ANOVA and Tukey post-hoc test for significance between SMU.

2.3 Cyanobacterial community structure

2.3.1 Microscopy of biocrust cyanobacteria

- 10 ~~Prior to the destruction of the samples for analysis, a portion of~~ field samples collected (Section 2.2) ~~subsections~~ were ~~removed~~
15 ~~from six of the eight Petri dishes, placed into six-cell plates and~~ used to enumerate cyanobacterial richness and diversity and
to classify colonies. ~~These~~ dried crust samples were resurrected in the glasshouse for three to five days. For each of the ten
sites, twelve replicates (subsamped from the eight samples) were analysed via light microscopy. For each replicate, a minimum
of two wet-mount slides incorporating six representative portions of the cyanobacterial colonies were examined (n=144
colonies per site). For the dominant land type, Chenopod shrubland (Site 8), there were an additional 10 x 6-cell multi-well
plates. These were treated similarly where two slides were examined from each of the 60 multi-wells (n=120). In total > 2,184
cyanobacterial colonies were examined. Initial inspection of the biocrust and the separation of individual species were made
using an Olympus SZH10 microscope at 70 x magnification. Cyanobacterial filaments or colonies were carefully extracted
with forceps to recover sufficient material that included important morphological features such as their colour, encasing sheaths
as well as cellular structure. Live material was examined by Nomarski differential interference contrast (DIC) microscopy with
20 a Jenaval (Jena Zeiss) and an Olympus BX51 compound microscope (magnifications 400–1000 x). Photomicrographs were
taken using an Olympus SC100 digital microscope camera, and morphological measurements of vegetative cells were made
from digital images of live material taken at 400 x magnification using Olympus cellSens® digital imaging software.

- 25 Identification was performed to a species level (wherever possible) in the laboratory using the following taxonomic references:
Anagnostidis and Komarek, (2005, 2005); Sant'Anna et al., (2011); Skinner and Entwisle, (2002). It was often necessary to
record the closest named species as attributes varied somewhat to temperate climate and aquatic specimens described in
literature. Nitrogen fixing cyanobacteria were identified based on the three recognised types: (1) heterocystous species (those
with specialised N-fixing cells); (2) non-heterocystous species that fix N aerobically and; (3) non-heterocystous species that
30 fix N anaerobically (Bergman et al., 1997; Stal, 1995). Using a graticule, abundance was ranked on a scale of 1–8 where the
main taxa are ranked in decreasing order of the relative percentage area occupied in a single view (Biggs and Kilroy, 2000).
More than one species could be dominant, and all other taxa were ranked in relation to the dominant taxa as abundant, common,

occasional and rare. To determine similarities between cyanobacterial communities, cluster analysis, SIMPROF, and non-metric multidimensional scaling (nMDS) were conducted using Primer v6 (Clarke & Gorley 2001).

2.3.2 16S rDNA profiling of native undisturbed biocrust microbiomes

For genomic profiling of naturally occurring successional biocrust communities, a location adjacent to Site 9 was visually
5 determined to contain Bare, Early (Types 1-2) or Late (Types 4-5) stages of development (Table S2). Biocrust successional
features were determined by morphological attributes of pigmentation, thickness and surface roughness as well as the
presence/absence of lichens and mosses (Fig. 2d) (Chilton et al., 2017). Bare stage was characterised by loose soil particles
with no visible biocrust structure. Samples were collected in July 2014. For each successional stage, three replicates were
collected that were representative of SMUs 1–3 where a 10 cm² plot with 95% coverage of the desired biocrust stage was
10 excised to the depth of the crust and non-aggregated soil discarded (Fig. 2e-g). Samples were processed at UNSW, Sydney.

Each biocrust replicate for Bare, Early and Late stages of development were homogenised and genomic DNA extraction
performed using the FASTDNA Spin Kit for Soil (MP Bio Laboratories, USA) according to the manufacturer's instructions.
Molecular libraries of the 16S rDNA V123 hypervariable region generated via PCR as per Chilton et al., (2017) and submitted
15 to the Ramaciotti Centre for Genomics (UNSW, Australia) for a 2x300 bp sequencing run on an Illumina MiSeq instrument.
Sequencing data was processed using Mothur version 1.34.0 (Schloss et al 2009) and described in detail in Chilton et al.,
(2017). Singleton and doubleton OTUs were removed and samples rarefied to 8598 sequences each across 3785 OTUS. The
curated Greengenes database (McDonald et al 2012) was used to assign taxonomy to OTUs. Diversity values were derived
using the DIVERSE function within the Primer package (Anderson et al 2008) upon standardized OTU values. ANOVA with
20 post hoc Tukey's tests was used to test for significant differences between stages. Multivariate analyses were performed in
Primer upon a Bray-Curtis dissimilarity matrix generated from square-root transformed abundance data. Samples were
represented in two and three-dimensional space within a nMDS plot. Pair-wise, a posteriori comparisons of factor Stage were
performed using the PERMANOVA function with 9999 Monte Carlo permutations. Homogeneity of dispersion for each stage
was tested using PERMDISP.

25 2.4 Cyanobacterial tolerance to stockpiling

Stockpile sampling was carried out in March 2012 with samples sourced from topsoil stockpiled from areas with *Acacia*
papyrocarpa (Western Myall) over-storey (Table S3). Stockpiles established at three different time points (9 months, 20
months and 29 months) as well as corresponding undisturbed sites were sampled in triplicate at six depths (0–2, 2–4, 4–6, 10,
25 and 50 cm). This resulted in 18 replicates for each soil depth for each stockpile time point. Holes were dug to >50 cm depth
30 with a shovel and the exposed profile removed with an ethanol-wiped spatula before taking a soil sample. A second sample at
50 cm was autoclaved to serve as a culturing control. Samples were stored in paper bags and processed at The University of
Queensland. For each sample, 20 g of soil was set up in petri dishes with 9 ml of water and sealed with Parafilm[®]. Petri dishes

were incubated for six weeks at 26°C under a 12-hour photoperiod regime and rotated weekly to prevent site specific effects. Samples were maintained as wet by the addition of sterile water within a laminar flow cabinet. To determine cyanobacterial growth and richness, wet-mounts for each sample were examined under 16 x magnification for cyanobacterial thalli and colony size was estimated via area of coverage of the field of view. Where multiple colonies were present in a slide, the relative abundance of each was used to divide the cover between the colonies accordingly. Where no growth was observed, five soil samples were taken randomly from the sample, mounted and examined under 400 x magnification for the presence of cyanobacterial cells. Identification was carried out as per methods detailed in Section 2.2.1. ANOVAs were performed with post hoc Tukey's tests performed to determine any in differences in relative abundance between treatments.

3.0 Results

3.1 Biophysical characteristics of biocrusts and cyanobacteria

Across the three soil management units (SMU) mean soil pH ranged from 8.5–8.6 while the two-year old topsoil stockpile was higher at pH 8.9 (Table 1). Electrical conductivity ranged from 92–140 $\mu\text{S cm}^{-1}$. Total nitrogen was typically <0.1% across all sites and total carbon ranged between 1–2% with higher percentages generally found across SMU 2 and 3 (Table 1). The ratio between carbon and nitrogen was the greatest across SMU 3 and the two-year old topsoil stockpile, also originating from SMU 3. Percentage C and C:N ratio across all SMU showed there were significant differences ($p=0.052$), and post hoc test showed there were significant SMU 1 and SMU 3. There were no significant differences between SMU 1 and SMU 2, or SMU 2 and SMU 3. Percentage N versus SMU was significant overall ($p=0.000$) with all three SMU are significantly different from each other.

Mean chlorophyll *a* concentration ranged between 10.33–13.64 $\mu\text{g g}^{-1}$ soil for the intact undisturbed biocrusts differed across all sites. From the three different SMU, the lowest values were found at SMU 1 ($p = 0.000$) whereas SMU 2 and SMU 3 showed no significant differences between them (Fig. 3). There were also significant differences in chlorophyll *a* concentration of the two-year old stockpile and SMU 1, SMU 2 and SMU 3 ($p=0.000$). Mean chlorophyll concentrations of biocrusts sourced from the two-year old topsoil stockpile ($7.49 \pm 1.01 \mu\text{g g}^{-1}$ soil) were almost half those of SMU 3 ($13.53 \pm 1.74 \mu\text{g g}^{-1}$ soil), which was the origin of the topsoil stockpile.

The photosynthetic yields recorded all fell within the expected range measurements on a scale of zero to one that ranged between 0.073 (stockpile) and 0.147 (SMU 2). SMU 3 and the two-year old stockpile were significantly lower than SMU 1 and SMU 2 ($p = 0.017$) whereas SMU 1 and SMU 2 were not significantly different (Fig. 3).

Moved (insertion) [1]

Deleted: there was variability between sites SMU 1 and SMU 2 were not significantly different

Deleted: whereas

Deleted: different

Deleted: from

Moved up [1]: SMU 1 and SMU 2 were not significantly different whereas SMU 3 and the two-year old stockpile were significantly different from SMU 1 and SMU 2 ($p = 0.017$).

Deleted: ($p = 0.017$).

Deleted: The yields recorded all fell within the expected range and when data was separated into their respective SMU it provided a clearer picture of the variability. SMU 1 and SMU 2 were not significantly different whereas SMU 3 and the two-year old stockpile were significantly different from SMU 1 and SMU 2 ($p = 0.017$). ¶

Compressive strengths across Sites 4, 9 and 10 in SMU 1 were significantly different from each other as well as significantly lower than SMU 2, SMU 3 and the two-year old topsoil stockpile (Table 2). The means for SMU 1–3 were also significantly different ($p < 0.0001$), and it was confirmed by a student's t-test that the difference was between SMU 1 and SMU 2 and 3.

3.2 Cyanobacterial community structure

5 A total of 21 cyanobacterial species were identified across the ten sites using microscopy (Table 5; Figs. S2–S9). The majority of species richness and abundance was comprised of four filamentous genera: *Symploca* (18%), *Schizothrix* (16%), *Porphyrosiphon* (16%) and *Scytonema* (16%). Secondary cyanobacteria present variously occupied <1 to 10%, and overall made up 34% of the community. The known nitrogen-fixing cyanobacteria *Symploca*, *Scytonema*, *Porphyrosiphon*, *Brasilonema*, *Nostoc* and *Gloeocapsa* comprised more than 50% of the species richness at each site and formed 61% of the total community richness (Fig. 4). In this study cyanobacterial community structure was tested against the three different soil types to determine whether soil type was influential in determining community structure. There was no significant difference in cyanobacterial community structure across soil types, however the results do suggest some spatial structuring exists across the three SMU (Fig. 5a). The structural relationship between all samples shows greater similarities rather than dissimilarities, of which there are only a small number of samples that are significantly different (Fig. 5b). Soil type did not explain these differences. *Symploca* occurred more frequently and was more abundant in the majority of the samples examined therefore was the most significant contributor to the community (data not shown). Individual sites generally displayed similar trends although there was some variability occurring between sites.

Moved down [2]: This suggests that most of the species could potentially be found anywhere across the three zones and their richness and abundance is controlled by other factors.

Of the 21 species more than half (12 species) of cyanobacteria were identified in SMU 1 where four primary genera made up 75% of the community: *Symploca*, *Schizothrix*, *Scytonema* and *Symplocastrum* (for more detail see Fig. S10). Cyanobacterial crusts from the dune regions on SMU 1 (deep calcareous yellow sands) were representative of crust types 1–3; patchy, brittle (when dry) early-successional crusts as well as formed dark crusts that were mid to late-successional and included cyanolichens (also see Doudle et al., 2011).

25 Cyanobacterial crusts from the chenopod shrublands and open woodlands in SMU 2 (shallow calcareous sandy loam) represented a broad range of crust types (2–5) but overall could be described as late-successional. Lichens and mosses were highly visible (also see Doudle et al., 2011). There were 21 cyanobacteria recorded: four were primary genera that made up 63% of the community including: *Schizothrix*; *Porphyrosiphon*; *Scytonema* and *Symploca* (for more detail see Fig. S11).

30 Cyanobacterial crusts from the open woodlands in SMU 3 (deep calcareous sandy loam, Fig. 2c) represented a broad range of crust types (2–5) but like SMU 2 could also be described as late-successional. Lichens and mosses were also highly visible (see Doudle et al., 2011). There were nine cyanobacteria recorded of which four were primary genera that made up 85% of the

community: *Symploca*, *Porphyrosiphon*, *Scytonema* and *Schizothrix* (for more detail see Fig S12). Cyanobacteria with the capacity to fix nitrogen contributed to 77% of the community structure.

Cyanobacterial crusts from the two-year old topsoil stockpile that had originated from SMU 3 (deep calcareous sandy loam) would be described as early successional crusts with some seasonal mosses. There were eight cyanobacteria recorded of which four were primary genera that made up 84% of the community: *Symploca*; *Symplocastrum*; *Porphyrosiphon*; and *Scytonema* (also see Fig. S13). It was interesting to note that *Symplocastrum* was co-dominant with *Symploca* whereas in the other communities it ranged between 8-13%. Sub-surface species *Schizothrix* (found in top 5 mm) only contributed to 4% of the richness compared to 10-20% elsewhere. Cyanobacteria with the capacity to fix nitrogen (*Symploca*, *Porphyrosiphon*, *Scytonema* and *Brasilonema*) contributed to 61% of the community (Fig. 4).

3.3 16S rDNA profiling of native undisturbed biocrust microbiomes

Microbial community profiling using high through-put sequencing revealed cyanobacteria comprised a significant component of all three stages forming the majority of sequences in Early and Late stages (Fig. 6a). There was a diversity of morphotypes observed including simple filamentous, heterocystous and unicellular types (Fig. 6b). The most abundant genera identified were *Leptolyngbya*, *Phormidium*, *Tolypothrix*, *Nostoc*, *Brasilonema*, *Chroococciopsis* and *Acaryochloris*. Unclassified Nostocaceae were dominant within Bare soils while Early stages observed a relative even increase in *Phormidium*, *Brasilonema* and the unicellular genera (e.g. *Chroococciopsis*, *Acaryochloris*, *Xenococcaceae*). Late stage biocrusts showed a slight resurgence of Nostocaceae. There was no significant difference in the richness, evenness or diversity between the three stages (Table 3). However, there were significant differences in the composition and structure of the communities of each stage. PERMANOVA showed stage was a significant factor explaining the data variability (Pseudo- $F=4.9544$, $P(\text{perm})=0.004$, Unique perms=273). Post-hoc pair-wise comparisons showed Bare stage was significantly different to the crusted stages (Fig. 7, Table 4) while resolution between the Early and Late stages was less clear. Ordination of the samples within three dimensions showed Early and Late stages grouped separately (Table 3).

3.4 Cyanobacterial tolerance to stockpiling

Stockpile soil samples contained five identifiable cyanobacterial genera: *Nostoc*, *Scytonema*, *Microcoleus*, *Porphyrosiphon* and *Leptolyngbya*. Average cyanobacterial richness was highest in undisturbed and stockpiled samples at and above 10 cm depth for all stockpile ages (Fig. 8 a,b). These differences were significant between 50 cm and 0–2 cm in all stockpiles (29-month $F_{(1,5)} = 7.00$, $p = 0.024$; 20-month $F_{(1,5)} + 8.37$, $p = 0.016$; 9-month $F_{(1,5)} = 18.00$, $p = 0.002$). Genera richness at the intermediate depths was not significantly different. Richness was significantly lower between 25 cm and 0–2 cm in the twenty- and nine-month old stockpiles ($F_{(1,5)} = 32.73$, $p = 0.000$; $F = 16.20$, $p = 0.000$) respectively. Counterintuitively, after nine and twenty months of stockpiling, average genera richness was higher in stockpiled samples when compared with undisturbed samples except for 20-month old 4–6 cm samples (Fig. 8 a, b), but this difference was only significant in the nine-month old

stockpile at 10 cm depths ($F_{(1,5)} = 8.27, p = 0.017$). High variability between replicates accounted for the lack of significance in the rest.

Cyanobacterial richness between stockpiles at different depths was greater in the youngest stockpiles and at the shallowest depths (<10 cm) but not at or below this level (Fig. 8b). Average richness was variable across adjacent undisturbed areas (Fig. 8a). Comparatively the variability in genera richness across undisturbed and stockpiles within replicates was high (Fig. S14 a–c). The cover of genera was relatively constant between sites in the undisturbed samples, but in stockpiled samples *Nostoc cf commune* was more prevalent ($F_{(1,5)} = 5.97, p = 0.012$) (Fig. S15).

Nostoc cf. commune, *Nostoc sp. (yellow)*, *Microcoleus* and *Leptolyngbya* were present in more stockpiled samples than undisturbed samples. Conversely, *Scytonema* and the black form of *Leptolyngbya* were more prevalent in undisturbed samples. When all samples were combined, *Nostoc cf commune* had the greatest cover in both stockpiled and undisturbed samples followed by *Nostoc sp. (yellow)* that was only found in stockpiles. These differences were significant from depths of 6 cm to the surface (4–6 cm $F_{(1,5)} = 28.83, p = 0.000$; 2–4 cm $F_{(1,5)} = 4.89, p = 0.023$; 0–2 cm $F_{(1,5)} = 3.72, p = 0.049$). These patterns were not reflected in the total coverage by *N. cf. commune* in corresponding undisturbed areas. Coverage by the remaining genera was similar although *Microcoleus* and *Leptolyngbya* had greater cover in stockpiled areas and *Porphyrosiphon* had slightly higher coverage in undisturbed samples. The cover of *Scytonema* was almost the same in stockpiled and undisturbed areas ($F_{(1,17)} = 0.00, p = 0.969$) (Fig. S15). *Leptolyngbya* (black morphotype) only occurred between 20 and 40 cm depth and were found in only one of the stockpile samples but were present in six samples from three adjacent undisturbed areas. *Stigonema* genus was found in six stockpile samples spanning all ages, although in only one adjacent sample but in all cases in the upper 10 cm of the soil profile.

Nostoc cf. commune, *Porphyrosiphon*, *Microcoleus* and *Scytonema* were the first genera to develop to an identifiable stage. Filaments of the *Stigonema* genus were found in low numbers and appeared to be recently formed. It was only present in samples examined in the latter stages of the identification process. *Nostoc sp. (yellow)* exhibited a much slower rate of development than *Nostoc cf commune* and could only be definitively determined as a form of *Nostoc* when examined after 13 weeks of incubation. Samples from undisturbed areas sourced at depths from between 10 cm and 50 cm initially showed no visible signs of growth when examined after six weeks of incubation. Six weeks later cyanobacterial growth was evident yet in many cases had not advanced to the point where morphotypes could be distinguished.

4.0 Discussion

This research has demonstrated cyanobacteria to be a key component of soil microbial communities at J-A. These were compositionally diverse topsoil microbiomes that substantially contributed to the Myall-chenopod landscape. We had

Deleted: yellow

Moved (insertion) [3]

Deleted: only

Deleted: also see

Deleted: ¶

Nostoc cf. commune exhibited the best survival response in stockpiling as it had significant coverage in samples from all depths in all stockpile ages. *Nostoc* had the greatest coverage in the oldest stockpiles, followed up by the most recently created stockpiles.

Moved up [3]: These differences were significant from depths of 6 cm to the surface (4–6 cm $F_{(1,5)} = 28.83, p = 0.000$; 2–4 cm $F_{(1,5)} = 4.89, p = 0.023$; 0–2 cm $F_{(1,5)} = 3.72, p = 0.049$). These patterns were not reflected in the total coverage by *N. cf. commune* in corresponding undisturbed areas.

Deleted: . The

5 hypothesised that cyanobacteria would be central to soil micro-processes, and this was strongly supported by extensive species richness and diversity values. At J-A cyanobacteria contributed to soil structure and function during the early developmental stages of the biocrust. Photosynthesis drove the productivity and growth of the biocrust that initiated carbon and nitrogen cycling and resulted in increases in soil nutrient concentrations right where vascular plants might use them. The results have demonstrated how these micro-processes provide a strong foundation for the restoration of soil function. Similarly, in southwestern Queensland and northern Australia cyanobacterial species richness was strongly linked to increased bioavailable nitrogen and carbon uptake (Büdel et al., 2018; Williams et al., 2018; Williams and Eldridge, 2011). Elsewhere, multiple studies have demonstrated the high value of biocrust attributes as drivers of soil micro-processes that restore soil function (e.g. Barger et al., 2016; Belnap and Eldridge, 2001; Bowker et al., 2014; Büdel et al., 2009; Chilton et al., 2017; Chiquoine et al., 2016; Weber et al., 2016).

15 In the first study of its kind we have shown the response of cyanobacteria to topsoil stockpiling at various depths and ages. We have demonstrated cyanobacteria are moderately resilient to stockpiling at depth and over time, providing they are manipulated dry (W. Williams, unpublished data). Cyanobacteria from the top 10 cm were found to be more viable within the first six weeks and showed potential for biocrust re-establishment. Curiously, we found greater cyanobacterial richness in the nine and 20-month stockpiles compared to undisturbed samples adjacent to the stockpiles. In general, the resilience of cyanobacteria to burial in topsoil stockpiles in the longer term appeared good, however in an arid environment recolonization and community diversity could be impeded by drought (Williams and Büdel, 2012)

20 In this study cyanobacterial community richness and abundance were not related to soil and landform type and this was further supported through the sequenced samples where no significant differences were observed. At J-A any of the cyanobacteria could conceivably occur anywhere across the landscape. Their relative abundance was most probably determined by microenvironments and microhabitats such as light (sun and shade) and chemical gradients (Stal, 2003), as well as moisture availability (Büdel et al., 2018) and soil particle size (Büdel et al., 2009).

4.1 Cyanobacterial community structure

25 Environmentally induced strategies of arid land cyanobacteria reflect their habitat, these survival traits have developed over a long evolutionary history. Cyanobacterial richness at J-A was determined according to their morphological features (e.g. outer protective sheaths, UV protection, EPS production) which in many cases provided the basis of attributes that pertained to fundamental survival strategies. Filamentous cyanobacteria formed the major part of the J-A crust structure with tufts, webs or creeping masses closely intertwined (e.g. *Porphyrosiphon*, *Symploca*, *Scytonema*, *Schizothrix*, *Microcoleus*). These are often assimilated with unicellular forms (e.g. *Gloeocapsa*, *Chroococcus*, *Chroococciopsis*) or gelatinous colonies of *Nostoc* (see supplementary material Fig. S16 for images of growth habits). Simple filamentous types are often attributed with the primary crust building role, able to span inter-particle gaps within the soil via supra-cellular structures (e.g. *Microcoleus*, see

Garcia-Pichel and Wojciechowski 2009). Important crust-building cyanobacteria in this study also appeared to be *Symploca* that was associated with EPS production, a principal feature of early colonising crust formation (Hu et al., 2002). Sequencing data showed *Phormidium* was the dominant cyanobacterium for this role and it is likely that *Symploca*, identified through microscopy, was the principal *Phormidium* present. *Microcoleus* sp. and *Porphyrosiphon* were also identified as early colonisers, however these genera are currently poorly resolved phylogenetically (Garcia-Pichel et al 2013) but share critical morphological features enabling biocrust formation and maintenance. Other key cyanobacteria indicative of biocrust formation and development were *Leptolyngbya*, *Phormidium*, *Tolypothrix*, *Nostoc*, *Brasilonema*, *Chroococcidiopsis* and *Acaryochloris*. Notably, the identification of *Brasilonema*, not been previously recorded in Australian soils, was supported with sequencing data. The taxonomic status of *Brasilonema* remained uncertain and may be a variety of *Scytonema*, however, genomic data supported morphological identification and the type has also been recorded in other terrestrial habitats globally. Due to its similar morphological attributes and genomic data, in this study we called this cyanobacterium *Brasilonema* (Fiore et al., 2007; Vaccarino and Johansen, 2012). Many primary (common to abundant) and secondary (uncommon) cyanobacteria recorded at J-A exhibited thick gelatinous sheaths (*Porphyrosiphon*, *Schizothrix*, *Microcoleus*, *Nostoc*) or were associated with the production of EPS (*Symploca*, *Nostoc*, *Schizothrix*, *Leptolyngbya*).

Overall, at J-A there was a rich cyanobacterial community comprised of twenty-one species recorded from 13 genera. Four species were unicellular and the remaining seventeen were filamentous. Some cyanobacteria found at J-A (*Microcoleus paludosus*, *Nostoc* sp., *Gloeocapsa*) had also been recorded at Lake Gilles (SA) about 400 km southeast of J-A (Ullmann and Büdel, 2001). Surprisingly though, *Microcoleus* species that were recorded at J-A did not dominate the biocrust compared with many reports from the United States, Asia and elsewhere (e.g. see Belnap and Eldridge, 2001). This infers that the early colonisers such as *Microcoleus* would not necessarily play a dominant role in early stabilisation and colonisation of the soil. At J-A *Symploca* and *Scytonema* appeared to be important as early colonising cyanobacteria in the biocrusts. These species have also been recorded as playing a key role in carbon sequestration in northern Australian cyanobacterial crusts (Büdel et al., 2018).

Nostoc commune var. *flagelliforme* had been recorded at J-A along with *Nostoc commune* across the shallow and deep sandy loams. Although *N. flagelliforme* appeared rarely, it had been previously documented from sites in south-western South Australia, Western Australia, Northern Territory (Skinner and Entwisle, 2002) and Victoria (W. Williams, unpublished data). Nevertheless, it has now been documented that both *Nostoc commune* and *N. flagelliforme* are not separate species, rather the spaghetti-like tubes are unique to the ecotype and likely associated with aridity (Aboal et al., 2016). This is supported by the semi-arid environment at J-A and it may be more widespread in Australia than previously recorded as it is often only clearly visible following rains.

4.2 Cyanobacterial tolerance to stockpiling

Physical disturbance of biocrusts occurs on a large scale at the J-A mine site with the removal and temporary stockpiling of topsoil. This type of mechanical disturbance results in burial and translocation of the biocrust. The impacts of burial within the natural environment are rarely studied and have never been analysed in such stockpiles. In China, artificial sand burial at shallow depths showed there were significant reductions in chlorophyll concentration, UV synthesis, total carbohydrates (EPS) and damage to photosynthetic activity (Rao et al., 2012). In a semi-arid grassland in Australia, wind-borne sand burial of cyanobacterial crusts during a severe drought resulted in a significant reduction in surface dwelling cyanobacteria and significant reductions in biological-available nitrogen (Williams and Eldridge, 2011). In contrast, we have shown here that without further disturbance cyanobacteria can survive stockpiling for over two years, an important feature underpinning the concept of biobanking soil microbial communities. This resilience is not surprising due to the recognised ability of cyanobacteria to survive in extreme environments. In previous studies, cyanobacteria have been grown from samples sourced at 18 cm depths in Japanese rice paddy soils (Fujita and Nakahara, 2006), 50 cm in the UK (Esmarch, 1914), and 70 cm depths in the USA (Moore and Karrer, 1919). *Microcoleus* and *Leptolyngbya* have survived and remained viable after up to three million years frozen in lake sediments in permafrost (Vishnivetskaya et al., 2003). Vegetative *Nostoc commune* material retains viability following several decades of storage in desiccated form (Bristol, 1919; Lipman, 1941). Reactivation of vegetative material after decades of storage was successful but several months (Lipman, 1941) to a year (Bristol, 1919) of incubation can be necessary for growth to take place. Notwithstanding, the cyanobacteria from the surface depths of the stockpiles (i.e. <10 cm) appeared more resilient in the short-term. These results were reflected in the current study where growth was not observed in the undisturbed areas below 10 cm depth for several months. It may be that the longer the period of inactivity, the longer time taken for reactivation to occur (Billi and Potts, 2002; Williams et al., 2014), or less material is viable therefore it takes longer to rebuild colonies (Agrawal and Singh, 2002).

Akinetes are desiccation resistant cells produced by certain filamentous cyanobacteria that can survive for long periods. For example, *Nostoc* and *Scytonema* produce akinetes (Kaplan-Levy et al., 2010; Tomaselli and Giovannetti, 1993) and species like *Nostoc* have the potential to grow at low light in caves and under ice (Dodds et al., 1995), or even in darkness (Huang et al., 1988). Belnap and Gardner, (1993) reported *Microcoleus vaginatus* sheaths at depths to 10 cm and considered that, due to a lack of chlorophyll, the sheaths were remnant from a time when the surface was lower than the current day. It is possible that heterotrophic growth for which chlorophyll is unnecessary was still occurring at these depths. The fact that these organisms took much longer to grow than those sampled from upper layers would suggest that they have grown from vegetative material that has been photosynthetically inactive for long periods resulting in long lag times for growth following re-activation (Bristol, 1919; Lipman, 1941; Shaw et al., 2003). In addition, there may be the potential for photo-damage to occur as many sub-surface cyanobacteria would now be exposed in the topsoil removal and stockpiling process. This could also disrupt and slow down the recovery process once re-spread; this was observed in the laboratory following an out of season heat wave where many sub-surface species were trapped on the surface and died (W. Williams, unpublished).

In the context of rehabilitation, it is not practical to store stockpiles at very shallow depths due to the land area they would occupy. We have shown that under optimum laboratory conditions a diverse range of cyanobacteria recovered from a range of stockpile depths. The next stage of experimentation should be to monitor the recovery of cyanobacteria from stockpiled soil that has been spread under normal mine practice.

5.0 Conclusions

The Eucla Basin, in which J-A is located, is situated on the eastern edge of the Nullarbor Plain. Naturally occurring biocrusts occupy at least 45% of the landscape with a diverse community structure that contributes to landscape resilience and function. Cyanobacteria naturally occur within the first few millimetres of the surface of the biocrust where access to resources such as light and moisture are essential. Biocrusts contain cyanobacterial species such as *Phormidium* and *Leptolyngbya* that are regarded as early colonisers, and other species such as *Scytonema* and *Nostoc* that contribute bioavailable nitrogen to the nutrient cycle.

In these studies, we found that at J-A cyanobacteria were a diverse community that had a proven capacity as ecosystem engineers. Many of these cyanobacteria were early colonisers and were represented by a high proportion of filamentous and N-fixing species. Biocrust functionality is aided by a diverse cyanobacterial population and its connective properties via EPS where nutrients and resources can be exchanged. The destruction of structural integrity and fragmentation presents hurdles, however biocrust resilience, especially cyanobacteria, can over time consolidate and recolonise.

At J-A, rare earth mining processes require the removal of the topsoil, this soil is mechanically scraped off and relocated to topsoil stockpiles. These stockpiles act as a biobank in that they contain the biocrust microorganisms and are valuable eco resources. A major component of these biocrusts is cyanobacteria known for their capacity to survive extreme environments with the ability to remain in a desiccated state for long periods of time. Nevertheless, the stockpiling process destroys the macrostructure of the soil profile thus fragmenting and translocating the biocrusts throughout the stockpiles. The persistence of cyanobacteria at depth in soil stockpiles was examined and it was found that survival diminished substantially below the top few centimetres and over time. Limited access to light for photosynthesis while buried can cause cyanobacterial death. Here, our results support the early return of topsoil stockpiles to facilitate the reestablishment of biocrusts and soil microbial community function. A direct-return process planned at J-A would improve outcomes by reducing the associated impacts of topsoil storage on key microbial communities such as cyanobacteria. Further research should focus on the : (1) establishing an optimum time for topsoil storage and, (2) the relationship of N-fixing cyanobacteria to seedling recruitment and plant establishment.

Moved down [6]: Cyanobacteria are ecosystem engineers in that they have the capacity to provide many of these crucial ecosystem services (Jones et al., 1994).

Formatted: Font: (Default) +Body (Times New Roman)

Moved up [4]: The cyanobacterial richness at J-A was determined according to their morphological features. In many cases these features (e.g. outer protective sheaths, UV protection, EPS production) provided the basis of attributes that pertained to fundamental survival strategies. Environmentally induced strategies of arid land cyanobacteria reflect their habitat, these survival traits have developed over a long evolutionary history. Many primary (common to abundant) and secondary (uncommon) cyanobacteria recorded at J-A exhibited thick gelatinous sheaths (*Porphyrosiphon*, *Schizothrix*, *Microcoleus*, *Nostoc*) or were associated with the production of EPS (*Symploca*, *Nostoc*, *Schizothrix*, *Leptolyngbya*). Filamentous cyanobacteria formed the major part of the J-A crust structure with tufts, webs or creeping masses closely intertwined (e.g. *Porphyrosiphon*, *Symploca*, *Scytonema*, *Schizothrix*, *Microcoleus*). These are often assimilated with unicellular forms (e.g. *Gloeocapsa*, *Chroococcus*, *Chroococcidiopsis*) or gelatinous colonies of *Nostoc* (see supplementary material Fig. S16 for images of growth habits). ¶

Moved up [5]: Long term inactivity of vegetative material can result in long lag times for growth following re-activation (Bristol, 1919; Lipman, 1941; Shaw et al., 2003)

Moved (insertion) [6]

Moved (insertion) [4]

Moved (insertion) [5]

Deleted: In terms of rehabilitation the natural capital in topsoil that has been removed in the mining process is often not recognised or poorly understood. Soils properties are crucial in the re-establishment of a raft of ecosystem services that include fertility, structure, climate regulation and biodiversity (Dominati et al., 2010). Cyanobacteria are ecosystem engineers in that they have the capacity to provide many of these crucial ecosystem services (Jones et al., 1994). This research demonstrated that the cyanobacterial communities in the J-A biocrusts were compositionally diverse topsoil microbiomes that substantially contributed to the Myall-chenopod biomes. We had hypothesised that cyanobacteria would be central to soil micro-processes and this was strongly supported by extensive species richness and diversity. Cyanobacteria are ecosystem engineers in that they have the capacity to provide many of these crucial ecosystem services (Jones et al., 1994).¶

In the first study of its kind we have shown the response of cyanobacteria to topsoil stockpiling at various depths and ages. In this study we have shown cyanobacteria are moderately resilient to stockpiling at depth and over time, providing they are manipulated dry (W. Williams, unpublished data). Cyanobacteria from the top 10 cm were found to be more viable within the first six weeks and showed potential for biocrust re-establishment. We found greater cyanobacterial richness in the nine and 20-month stockpiles compared to undisturbed samples adjacent to the stockpiles. In general, the resilience of cyanobacteria to burial in topsoil stockpiles in the longer term appeared good, however in an arid environment (...)

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman)

Moved (insertion) [2]

Table 1: Intact biocrust soil physicochemical descriptions for all sites EC = electrical conductivity in $\mu\text{S cm}^{-1}$; total percentage of nitrogen present (N%), total percentage of carbon present (C%) and carbon to nitrogen ratios (C:N) for all sites. *SMU3 is the origin of the topsoil stockpile aged two years which was not included in the data analysis with SMU1-3 as there were only two samples. Different letters indicate significant differences in columns.

Vegetation	Soil	SMU	Site	pH	EC	N%	C%	C:N
Mallee	Deep calcareous yellow sand	1	4,9,10	8.6 ^a	92 ^a	0.07 ^c	1.11 ^b	14.9 ^{a,b}
Myall, Mallee, Chenopod	Shallow calcareous sandy loam	2	1,5,8	8.5 ^a	122 ^a	0.11 ^a	1.40 ^{a,b}	13.0 ^b
Myall	Deep calcareous sandy loam	3	2,3,7	8.5 ^a	113 ^a	0.09 ^b	1.61 ^a	17.3 ^a
Myall	Topsoil stockpile	Origin *SMU3	6	8.9	119	0.08	1.57	19.6

Table 2: Biocrust compressive strengths measured with a penetrometer, means and standard deviations (SD) for SMU 1–3 (kg cm^{-2}) tests with p-values, values in bold that are different from 0 with a significance level $\alpha=0.05$, letters indicate significant differences

SMU	Means \pm SD	SMU 1	SMU 2	SMU 3
SMU1	2.79 \pm 1.41 ^b	0	0.001	0.005
SMU2	3.75 \pm 0.79 ^a	0.001	0	NS
SMU3	3.97 \pm 0.70 ^a	0.005	NS	0

- Deleted: ^a
- Deleted: ^a
- Formatted: Superscript
- Formatted: Superscript
- Formatted: Superscript
- Formatted: Superscript
- Deleted: ^a
- Deleted: ^b
- Deleted: ^a
- Deleted: ^b
- Formatted: Superscript
- Formatted: Superscript
- Deleted: ^c
- Formatted: Superscript
- Deleted: ^c
- Formatted: Superscript
- Deleted: ^a
- Deleted: ¶
- Deleted: ^b
- Formatted: Superscript
- Deleted: ^a
- Formatted: Superscript
- Deleted: ^a
- Formatted: Superscript

Table 3: Cyanobacterial mean (\pm Standard Error) of richness (Margalef's index), evenness (Pielou's index) and diversity (Shannon index) across successional stages. No significant difference in diversity measures was found between stages.

	Bare		Early		Late	
	Mean	SE	Mean	SE	Mean	SE
Richness (d)	167.4	5.344	146.8	5.818	142.8	15.69
Evenness (J')	0.816	0.011	0.791	0.005	0.788	0.027
Diversity (H')	5.977	0.103	5.692	0.064	5.642	0.281

5

Deleted: ¶

Table 4: Permutational analysis of variance (PERMANOVA) of pair-wise comparisons of Bray-Curtis dissimilarity between biocrust stages and bare soil. P(MC) = probability values obtained using 9999 Monte Carlo permutations. A test for homogeneity of multivariate dispersions (PERMDISP) showed no significant differences in variation of spread of samples (pseudo $F=3.8068$, $P(\text{perm}) = 0.068$). Significant pair-wise differences are in bold.

Groups	t	P(perm)	Unique	
			permutations	P(MC)
Bare,				
Early	2.6216	0.0979	10	0.0107
Bare,				
Late	2.5742	0.0959	10	0.0120
Early,				
Late	1.2793	0.0953	10	0.1993

15

Table 5: Diversity across sites on a presence absence basis for all seasons and Lake Ifould (salt lake). Different species attributed to a genus (i.e. sp. 1,2,3) have all been separated based on their morphological features and size but could not be positively identified.

Cyanobacterium	SMU 1	SMU 2	SMU 3	T2 Stockpile	Lake 5 Ifould
Aphanothece					x
Brasilonema	x	x	x	x	x
Chroococciopsis		x		x	
Chroococcus sp. 1	x	x	x	x	
Chroococcus sp. 2					x
Gloeocapsa	x	x			x
Leptolyngbya	x	x			x
Microcoleus cthonoplastes					x
Microcoleus paludosus	x	x	x	x	x
Microcoleus sociatus					x
Microcoleus vaginatus	x	x			x
Nostoc commune		x	x	x	x
Nostoc flagelliforme		x			
Nostoc pruniforme		x			
Nostoc sp.	x	x			
Porphyrosiphon sp. 1	x	x	x	x	
Porphyrosiphon sp. 2		x			x
Schizothrix sp. 1		x			x
Schizothrix sp. 2	x	x	x	x	
Schizothrix sp. 3					x
Scytonema sp. 1	x	x	x	x	
Scytonema sp. 2		x			x
Scytonema sp. 3		x			
Scytonema sp. 4					x
Symploca sp. 1	x	x	x	x	x
Symploca sp. 2		x			
Symplocastrum sp. 1	x	x	x	x	x
Symplocastrum sp. 2					x
Species richness	12	21	9	10	18




Figure 1: Image of biocrust sample sites located within the vegetation associations described in Table S1 (supplied, S. Doudle). Site 11 was initially investigated but later discarded as it was a fourth replicate of the Chenopod Shrubland.




Figure 2: (a) SMU 1 (Sites 4, 9 and 10): Type 1–3 Biocrusts on deep calcareous yellow sands (dunes); (b) SMU 2 (Sites 1, 5 and 8): Primarily types 4 and 5 biocrusts on shallow calcareous sandy loam; (c) SMU 3 (Sites 2, 3 and 7): Types 1-5 biocrusts on deep calcareous sandy loam (Photographs by S. Doudle, 2011); (d) different biocrust stages (top) north of Stockpile 19, adjacent to Site 9; (e) Bare, (f) Early and (g) Late stages, also showing biocrust sample already removed from Late stage (Photographs A. Chilton).





Figure 3: Chlorophyll *a* concentration ($\mu\text{g g}^{-1}$ soil) following resurrection (2 weeks) after a desiccation (dry) period of 6 months. Soil Management Unit 1 (SMU 1) was significantly lower than SMU 2–3 ($p = 0.000$). Mean values and standard error of the mean (SEM) displayed where SMU 1 = Sites 4, 9, 10; SMU 2 = Sites 1, 5, 8; SMU 3 = Sites 2, 3, 7; T2 = two-year old Topsoil stockpile (originating from SMU 3). Photosynthetic yield (YII) of photosystem II (PSII) for SMU 1–3 and T2 (Site 6 two-year old topsoil stockpile), displaying mean values and standard error of the mean (SEM). SMU 1 and SMU 2 were significantly different from SMU 3 and T2 ($p = 0.017$), SMU 1 and SMU 2 were not significantly different from each other.

10

15



5 Figure 4: Cyanobacterial community structure across all sites expressed as a percentage of the total community based on mean richness and abundance scores. N-fixing cyanobacteria contributed to 61% of the community structure.

10

15






Figure 5(a): Cyanobacterial community structure based on indexed abundance and diversity across all sites displayed in an nMDS plot (Bray Curtis similarity). SMU 1-3 refer to key soil management units; SMU 2 appeared to have more range in the clusters compared to the closer groupings of SMU 1 and 3. T2 (TS2Y) is the two-year old topsoil stockpile; note these are clustered more closely. Figure 5(b): Similarities between samples within their SMU's are displayed in a Bray Curtis dendrogram. Black continuous lines show significant differences between samples ($p = 0.05$) and lighter lines indicate that most samples were not significantly different to each other.



Formatted Table

(a)

Deleted: .



(b)

Deleted: .

Figure 6(a): Relative abundance of cyanobacteria to other bacteria within Bare soil and Early and Late stage biocrusts. Figure 6(b): Abundance of cyanobacterial genera and groups. Green = Simple filamentous types, Blue = Heterocystic types, Purple = Unicellular. Grey = Unclassified/Other includes chloroplasts.




Figure 7: Non-metric Multidimensional Scaling of Bare soil and Early and Late Stage biocrusts within two dimensions (left) and three dimensions (right).


5

10


15

20

25




(a)




(b)

Figure 8: Cyanobacterial richness is described where: (8a) stockpiles of different ages had a significantly lower number of morphotypes at 25 cm compared to 0–2 cm in the twenty- and nine-month old stockpiles ($p = 0.000$; $p = 0.000$); (8b) adjacent to the stockpiles of three different ages (9, 20 and 29 months) the undisturbed sites had a significantly lower number of morphotypes at 50 cm compared to 0–2 cm in all stockpiles ($p = 0.024$; 20-month $p = 0.016$; 9-month $p = 0.002$).



Deleted:

Formatted Table



Deleted:

Deleted: ¶

Deleted: s: (a)

Deleted:

Deleted: were

Deleted: between

Deleted: and

Deleted: and,

Deleted: (b) stockpiles of different ages were significant between 25 cm and 0–2 cm in the twenty- and nine-month old stockpiles ($p = 0.000$; $p = 0.000$).

Acknowledgements

This project was funded and supported by Iluka Resources Ltd. and the Jacinth-Ambrosia Rehabilitation team. Thanks to Sam Doudle, Iluka Rehabilitation Specialist for her recognition of the importance of biocrusts at Jacinth-Ambrosia and for initiating this research program. This was a UniQuest project that was carried out at The University of Queensland's School of Agriculture and Food Sciences, Gatton Campus and the University of New South Wales. Especially appreciated was the strong support, project advice and laboratory resources provided by Assoc. Prof Vic Galea and Katherine Raymont. Special thanks to Dr Glenn McGregor for technical advice, assistance with cyanobacterial identification and the use of specialist microscopes and facilities at the Ecosciences Precinct (Qld), as well as report editing. Special thanks to David Tongway who assisted in Landscape Function Analysis field work, and two anonymous reviewers who provided valuable insights and editing to greatly improve this manuscript.

References

- Aboal, M., Werner, O., García-Fernández, M. E., Palazón, J. A., Cristóbal, J. C. and Williams, W.: Should ecomorphs be conserved? The case of *Nostoc flagelliforme*, an endangered extremophile cyanobacteria, *J. Nat. Conserv.*, 30(Supplement C), 52–64, doi:10.1016/j.jnc.2016.01.001, 2016.
- Agrawal, S. C. and Singh, V.: Viability of dried filaments, survivability and reproduction under water stress, and survivability following heat and UV exposure in *Lyngbya martensiana*, *Oscillatoria agardhii*, *Nostoc calcicola*, *Hormidium fluitans*, *Spirogyra* sp. and *Vaucheria geminata*, *Folia Microbiol. (Praha)*, 47(1), 61–67, doi:10.1007/BF02818567, 2002.
- Barger, N. N., Weber, B., Garcia-Pichel, F., Zaady, E. and Belnap, J.: Patterns and Controls on Nitrogen Cycling of Biological Soil Crusts, in *Biological Soil Crusts: An Organizing Principle in Drylands*, pp. 257–285, Springer, Cham., 2016.
- Barnes, J. D., Balaguer, L., Manrique, E., Elvira, S. and Davison, A. W.: A reappraisal of the use of DMSO for the extraction and determination of chlorophylls a and b in lichens and higher plants, *Environ. Exp. Bot.*, 32(2), 85–100, doi:10.1016/0098-8472(92)90034-Y, 1992.
- Belnap, J. and Eldridge, D.: Disturbance and Recovery of Biological Soil Crusts, in *Biological Soil Crusts: Structure, Function, and Management*, pp. 363–383, Springer, Berlin, Heidelberg., 2001.
- Belnap, J. and Gardner, J. S.: Soil microstructure in soils of the Colorado Plateau: the role of the cyanobacterium *Microcoleus vaginatus*, *Gt. Basin Nat.*, 40–47, 1993.
- Belnap, J. and Gillette, D. A.: Vulnerability of desert biological soil crusts to wind erosion: the influences of crust development, soil texture, and disturbance, *J. Arid Environ.*, 39(2), 133–142, doi:10.1006/jare.1998.0388, 1998.
- Billi, D. and Potts, M.: Life and death of dried prokaryotes, *Res. Microbiol.*, 153(1), 7–12, doi:10.1016/S0923-2508(01)01279-7, 2002.

- Bowker, M. A.: Biological Soil Crust Rehabilitation in Theory and Practice: An Underexploited Opportunity, *Restor. Ecol.*, 15(1), 13–23, doi:10.1111/j.1526-100X.2006.00185.x, 2007.
- Bowker, M. A., Maestre, F. T., Eldridge, D., Belnap, J., Castillo-Monroy, A., Escolar, C. and Soliveres, S.: Biological soil crusts (biocrusts) as a model system in community, landscape and ecosystem ecology, *Biodivers. Conserv.*, 23(7), 1619–1637, doi:10.1007/s10531-014-0658-x, 2014.
- 5 Bristol, B. M.: ON THE RETENTION OF VITALITY BY ALGÆ FROM OLD STORED SOILS., *New Phytol.*, 18(3-4), 92–107, 1919.
- Büdel, B., Darienko, T., Deuschewitz, K., Dojani, S., Friedl, T., Mohr, K. I., Salisch, M., Reisser, W. and Weber, B.: Southern African biological soil crusts are ubiquitous and highly diverse in drylands, being restricted by rainfall frequency, *Microb. Ecol.*, 57(2), 229–247, doi:10.1007/s00248-008-9449-9, 2009.
- 10 Büdel, B., Williams, W. J. and Reichenberger, H.: Annual net primary productivity of a cyanobacteria-dominated biological soil crust in the Gulf Savannah, Queensland, Australia, *Biogeosciences*, 15(2), 491, 2018.
- Carrick, P. J. and Krüger, R.: Restoring degraded landscapes in lowland Namaqualand: Lessons from the mining experience and from regional ecological dynamics, *J. Arid Environ.*, 70(4), 767–781, doi:10.1016/j.jaridenv.2006.08.006, 2007.
- 15 Chilton, A. M., Neilan, B. A. and Eldridge, D. J.: Biocrust morphology is linked to marked differences in microbial community composition, *Plant Soil*, 1–11, doi:10.1007/s11104-017-3442-3, 2017.
- Chiquoine, L. P., Abella, S. R. and Bowker, M. A.: Rapidly restoring biological soil crusts and ecosystem functions in a severely disturbed desert ecosystem, *Ecol. Appl.*, 26(4), 1260–1272, doi:10.1002/15-0973, 2016.
- Codony, F., Miranda, A. M. and Mas, J.: Persistence and proliferation of some unicellular algae in drinking water systems as result of their heterotrophic metabolism, *Water Sa*, 29(1), 113–116, 2003.
- 20 Delgado-Baquerizo, M., Morillas, L., Maestre, F. T. and Gallardo, A.: Biocrusts control the nitrogen dynamics and microbial functional diversity of semi-arid soils in response to nutrient additions, *Plant Soil*, 372(1–2), 643–654, doi:10.1007/s11104-013-1779-9, 2013.
- Dodds, W. K., Gudder, D. A. and Mollenhauer, D.: The ecology of Nostoc, *J. Phycol.*, 31(1), 2–18, 1995.
- 25 Doherty, K. D., Antoninka, A. J., Bowker, M. A., Ayuso, S. V. and Johnson, N. C.: A Novel Approach to Cultivate Biocrusts for Restoration and Experimentation, *Ecol. Restor.*, 33(1), 13–16, 2015.
- Dominati, E., Patterson, M. and Mackay, A.: A framework for classifying and quantifying the natural capital and ecosystem services of soils, *Ecol. Econ.*, 69(9), 1858–1868, doi:10.1016/j.ecolecon.2010.05.002, 2010.
- Doudle, S., Williams, W. and Galea, V.: Improving rehabilitation outcomes using biocrusts, *8th Int. Heavy Miner. Conf.* 2011, 85–97, 2011.
- 30 Elbert, W., Weber, B., Burrows, S., Steinkamp, J., Büdel, B., Andreae, M. O. and Pöschl, U.: Contribution of cryptogamic covers to the global cycles of carbon and nitrogen, *Nat. Geosci.*, 5(7), 459–462, doi:10.1038/ngeo1486, 2012.
- Eldridge, D. J. and Leys, J. F.: Exploring some relationships between biological soil crusts, soil aggregation and wind erosion, *J. Arid Environ.*, 53(4), 457–466, doi:10.1006/jare.2002.1068, 2003.

- Esmarch, F.: Investigations of the distribution of Cyanophyceae on and in various soils, 1914.
- Fiore, M. F., Sant'Anna, C. L., Azevedo, M. T. de P., Komárek, J., Kaštovský, J., Sulek, J. and Lorenzi, A. A. S.: The Cyanobacterial Genus *Brasilonema*, Gen. Nov., a Molecular and Phenotypic Evaluation I, J. Phycol., 43(4), 789–798, doi:10.1111/j.1529-8817.2007.00376.x, 2007.
- 5 Flores, E. and Herrero, A.: Compartmentalized function through cell differentiation in filamentous cyanobacteria, Nat. Rev. Microbiol., 8(1), 39–50, 2010.
- Fujita, Y. and Nakahara, H.: Variations in the microalgal structure in paddy soil in Osaka, Japan: comparison between surface and subsurface soils, Limnology, 7(2), 83–91, doi:10.1007/s10201-006-0167-z, 2006.
- Gillieson, D., Wallbrink, P. and Cochrane, A.: Vegetation change, erosion risk and land management on the Nullarbor Plain, Australia, Environ. Geol., 28(3), 145–153, doi:10.1007/s002540050087, 1996.
- 10 Harris, J. A.: Measurements of the soil microbial community for estimating the success of restoration, Eur. J. Soil Sci., 54(4), 801–808, doi:10.1046/j.1351-0754.2003.0559.x, 2003.
- Hou, B. and Warland, I.: Heavy mineral sands potential of the Eucla Basin in South Australia—a world-class palaeobeach placer province, Mesa J., 37, 4–12, 2005.
- 15 Hu, C., Liu, Y., Zhang, D., Huang, Z. and Paulsen, B. S.: Cementing mechanism of algal crusts from desert area, Chin. Sci. Bull., 47(16), 1361–1368, doi:10.1360/02tb9301, 2002.
- Huang, T.-C., Chow, T.-J. and Hwang, I.-S.: The cyclic synthesis of the nitrogenase of *Synechococcus* RF-1 and its control at the transcription level, FEMS Microbiol. Lett., 50(2–3), 127–130, 1988.
- Jasper, D. A.: Beneficial Soil Microorganisms of the Jarrah Forest and Their Recovery in Bauxite Mine Restoration in Southwestern Australia, Restor. Ecol., 15, S74–S84, doi:10.1111/j.1526-100X.2007.00295.x, 2007.
- 20 Jones, C. G., Lawton, J. H. and Shachak, M.: Organisms as Ecosystem Engineers, in Ecosystem Management, pp. 130–147, Springer, New York, NY., 1994.
- Kaplan-Levy, R. N., Hadas, O., Summers, M. L., Rücker, J. and Sukenik, A.: Akinetes: dormant cells of cyanobacteria, in Dormancy and resistance in harsh environments, pp. 5–27, Springer., 2010.
- 25 Kidron, G. J., Barinova, S. and Vonshak, A.: The effects of heavy winter rains and rare summer rains on biological soil crusts in the Negev Desert, CATENA, 95(Supplement C), 6–11, doi:10.1016/j.catena.2012.02.021, 2012.
- Lipman, C. B.: The successful revival of *Nostoc commune* from a herbarium specimen eighty-seven years old, Bull. Torrey Bot. Club, 664–666, 1941.
- Lukešová, A.: Soil Algae in Brown Coal and Lignite Post-Mining Areas in Central Europe (Czech Republic and Germany), Restor. Ecol., 9(4), 341–350, doi:10.1046/j.1526-100X.2001.94002.x, 2001.
- 30 Maestre, F. T., Castillo-Monroy, A. P., Bowker, M. A. and Ochoa-Hueso, R.: Species richness effects on ecosystem multifunctionality depend on evenness, composition and spatial pattern, J. Ecol., 100(2), 317–330, doi:10.1111/j.1365-2745.2011.01918.x, 2012.

- McCutcheon, J., Wilson, S. A. and Southam, G.: Microbially accelerated carbonate mineral precipitation as a strategy for in situ carbon sequestration and rehabilitation of asbestos mine sites, *Environ. Sci. Technol.*, 50(3), 1419–1427, 2016.
- Moore, G. T. and Karrer, J. L.: A subterranean algal flora, *Ann. Mo. Bot. Gard.*, 6(4), 281–307, 1919.
- Rao, B., Liu, Y., Lan, S., Wu, P., Wang, W. and Li, D.: Effects of sand burial stress on the early developments of cyanobacterial crusts in the field, *Eur. J. Soil Biol.*, 48(Supplement C), 48–55, doi:10.1016/j.ejsobi.2011.07.009, 2012.
- 5 Reisser, W.: *The hidden life of algae underground*, in *Algae and Cyanobacteria in Extreme Environments*, pp. 47–58, Springer, 2007.
- Rossi, F., Mugnai, G. and Philippis, R. D.: *Complex role of the polymeric matrix in biological soil crusts*, *Plant Soil*, 1–16, doi:10.1007/s11104-017-3441-4, 2017.
- 10 Shaw, S. L., Chisholm, S. W. and Prinn, R. G.: Isoprene production by *Prochlorococcus*, a marine cyanobacterium, and other phytoplankton, *Mar. Chem.*, 80(4), 227–245, 2003.
- Skinner, S. and Entwisle, T.: *Non-marine algae of Australia: 1. Survey of colonial gelatinous blue-green macroalgae (Cyanobacteria)*, *Telopea*, 9(3), 573–599, doi:10.7751/telopea20024003, 2002.
- Stal, L. J.: *Microphytobenthos, their Extracellular Polymeric Substances, and the Morphogenesis of Intertidal Sediments*, *Geomicrobiol. J.*, 20(5), 463–478, doi:10.1080/713851126, 2003.
- 15 Tomaselli, L. and Giovannetti, L.: *Survival of diazotrophic cyanobacteria in soil*, *World J. Microbiol. Biotechnol.*, 9(1), 113–116, 1993.
- Tongway, D. and Hindley, N.: *Landscape function analysis: a system for monitoring rangeland function*, *Afr. J. Range Forage Sci.*, 21(2), 109–113, doi:10.2989/10220110409485841, 2004.
- 20 Tongway, D. and Ludwig, J. A.: *Planning and implementing successful landscape-scale restoration*, *Restor. Ecol. New Front.*, 30, 2012.
- Tongway, D. J.: *Soil and landscape processes in the restoration of rangelands*, *Rangel. J.*, 12(1), 54–57, doi:10.1071/rj9900054, 1990.
- Tongway, D. J. and Ludwig, J. A.: *Rehabilitation of Semiarid Landscapes in Australia. I. Restoring Productive Soil Patches*, *Restor. Ecol.*, 4(4), 388–397, doi:10.1111/j.1526-100X.1996.tb00191.x, 1996.
- 25 Ullmann, I. and Büdel, B.: *Ecological determinants of species composition of biological soil crusts on a landscape scale*, *Biol. Soil Crusts Struct. Funct. Manag.*, 203–213, 2001.
- Vaccarino, M. A. and Johansen, J. R.: *Brasilonema Angustatum* Sp. Nov. (nostocales), a New Filamentous Cyanobacterial Species from the Hawaiian Islands 1, *J. Phycol.*, 48(5), 1178–1186, doi:10.1111/j.1529-8817.2012.01203.x, 2012.
- 30 Vishnivetskaya, T. A., Spirina, E. V., Shatilovich, A. V., Erokhina, L. G., Vorobyova, E. A. and Gilichinsky, D. A.: *The resistance of viable permafrost algae to simulated environmental stresses: implications for astrobiology*, *Int. J. Astrobiol.*, 2(3), 171–177, doi:10.1017/S1473550403001575, 2003.
- Weber, B., Bowker, M., Zhang, Y. and Belnap, J.: *Natural Recovery of Biological Soil Crusts After Disturbance*, in *Biological Soil Crusts: An Organizing Principle in Drylands*, pp. 479–498, Springer, Cham., 2016.

- Wellburn, A. R.: The Spectral Determination of Chlorophylls a and b, as well as Total Carotenoids, Using Various Solvents with Spectrophotometers of Different Resolution, *J. Plant Physiol.*, 144(3), 307–313, doi:10.1016/S0176-1617(11)81192-2, 1994.
- 5 Williams, W., Büdel, B. and Williams, S.: Wet season cyanobacterial N enrichment highly correlated with species richness and *Nostoc* in the northern Australian savannah, *Biogeosciences*, 15(7), 2149–2159, doi:https://doi.org/10.5194/bg-15-2149-2018, 2018.
- Williams, W. J. and Büdel, B.: Species diversity, biomass and long-term patterns of biological soil crusts with special focus on Cyanobacteria of the Acacia aneura Mulga Lands of Queensland, Australia, *Algol. Stud.*, 140(1), 23–50, 2012.
- 10 Williams, W. J. and Eldridge, D. J.: Deposition of sand over a cyanobacterial soil crust increases nitrogen bioavailability in a semi-arid woodland, *Appl. Soil Ecol.*, 49, 26–31, doi:10.1016/j.apsoil.2011.07.005, 2011.
- Williams, W. J., Büdel, B., Reichenberger, H. and Rose, N.: Cyanobacteria in the Australian northern savannah detect the difference between intermittent dry season and wet season rain, *Biodivers. Conserv.*, 23(7), 1827–1844, doi:10.1007/s10531-014-0713-7, 2014.
- 15 Zhao, Y., Zhu, Q., Li, P., Zhao, L., Wang, L., Zheng, X. and Ma, H.: Effects of artificially cultivated biological soil crusts on soil nutrients and biological activities in the Loess Plateau, *J. Arid Land*, 6(6), 742–752, doi:10.1007/s40333-014-0032-6, 2014.