



# Impact of decadal reversals of the North Ionian circulation on phytoplankton phenology

Héloïse Lavigne<sup>1</sup>, Giuseppe Civitarese<sup>2</sup>, Miroslav Gacic<sup>2</sup>, Fabrizio D'Ortenzio<sup>3</sup>

<sup>1</sup>Royal Belgian Institute of Natural Sciences, Operational Directorate Natural Environment, Gulledele 100, B-1200 Brussels, Belgium.

<sup>2</sup>Istituto Nazionale di Oceanografia e di Geofisica Sperimentale – OGS, Dip. di Oceanografia, Borgo Grotta Gigante 42/c, 34010 Sgonico (Trieste), Italy

<sup>3</sup>Laboratoire d'Océanographie de Villefranche, Université Pierre et Marie Curie et CNRS, UMR 7093, Villefranche-sur-Mer, France

*Correspondence to:* Héloïse Lavigne ([hlavigne@naturalsciences.be](mailto:hlavigne@naturalsciences.be))

**Abstract.** In the North Ionian, water circulation is characterized by a decadal alternation of cyclonic and anticyclonic regime driven by the mechanism called BiOS (Bimodal Oscillating System). The circulation regime affects the vertical dynamics and the nutrient distribution. The North Ionian is then a good study area to investigate how changes in circulation can affect phytoplankton dynamics in oligotrophic regions. From in situ observations, for each circulation regime the averaged distribution of isopycnals is provided, and a depth difference of about 80m is estimated for the nitracline between cyclonic and anticyclonic regime. Based on phytoplankton phenology metrics extracted from annual time-series of satellite ocean color data for the period 1998-2012, the cyclonic and anticyclonic regimes are compared. Results show that the average chlorophyll in March, the date of bloom initiation and the date of maximum chlorophyll were affected by circulation patterns in the North Ionian. In the center of the gyre, bloom initiation occurred in December and chlorophyll was low in March when circulation was anticyclonic, whereas during the cyclonic circulation regime, a late chlorophyll peak, likely resulting from different phytoplankton dynamics, was commonly observed in March. An additional analysis shows that the winter buoyancy losses, which govern the Mixed Layer Depth (MLD) also contribute to explain the interannual variability in bloom initiation and intensity. Two scenarios involving the relative position of the MLD and nitracline are finally developed, discussed and tested with model data to explain the different phenology patterns observed in the North Ionian.

## 1 Introduction

Phytoplankton phenology aims to analyze periodic events in phytoplankton seasonal cycles, such as blooms (Ji et al., 2010). Phytoplankton blooms are crucial events in the ocean as they contribute significantly to primary production and carbon export, (Buesseler, 1998) but also sustain the trophic food web. Thus, any changes in phytoplankton phenology may have large consequences on higher trophic levels (Edwards and Richardson, 2004; Koeller et al., 2009). Although the characterization of phytoplankton phenology in the open ocean has long been hindered by data scarcity, in the past few



years, satellite ocean color observations have offered a general view of the phenology of the surface phytoplankton biomass at global scale (Platt et al., 2010). Satellite based studies have mainly attempted to describe phytoplankton blooms (Yoder et al., 1993; Racault et al., 2012; Winder and Cloern, 2010) but have also tried to explain their spatio-temporal variability. Mixed Layer Depth (MLD) is recognized to be one of the main factors influencing phytoplankton phenology as it controls both light and nutrients availability for phytoplankton growth (Mann and Lazier, 2006). Numerous studies have attempted then to explain variability in phytoplankton phenology with MLD (Lavigne et al., 2013; Henson et al., 2006; Zhai et al., 2011) or with factors influencing it (Follows and Dutkiewicz, 2001, Ji et al., 2008). These studies have shown that, generally, a deeper winter mixing results in a delayed and stronger bloom in subtropical regions but in a weaker bloom in subpolar regions. Nevertheless, in most cases, MLD has failed to explain the whole phytoplankton variability suggesting that all the factors that impact phytoplankton phenology are still not known.

In oligotrophic environments, where nutrients are the primary factor controlling phytoplankton growth, changes in the subsurface distribution of nutrients should also affect phytoplankton phenology. Nutrient distribution is controlled by the interplay between biogeochemical and physical processes and external sources (Williams and Follows, 2003). Among the physical processes, oceanic circulation, which at global scale contributes to the transport of nutrients, can also regionally uplift or depress nutricline and modify nutrient concentrations through cyclonic and anticyclonic circulation patterns (Williams and Follows, 2003). At basin scale, the trophic status of subtropical (subpolar) gyre is determined by anticyclonic (cyclonic) circulation, which maintains the downwelling (upwelling) of the nutricline and controls the nutrient stock in subsurface waters. At smaller scales, observations and modelling experiments have demonstrated that cyclonic eddies cause the upwelling of the nutricline just above the base of the productive layer resulting in an increase of the primary production in subsurface waters (McGillicuddy et al., 1998; Falkowski et al., 1991; Salihoglu et al., 1990). These observations were confirmed by a statistical analysis carried out in the Costa Rica Dome area (Kahru et al., 2007). In fact, the authors revealed a strong coupling between the Sea Level Anomaly (SLA) and the chlorophyll anomaly with positive chlorophyll anomalies associated with negative SLA. Regarding phytoplankton phenology, little is known about the impact of the cyclonic/anticyclonic circulation on it. To our knowledge, only two recent analyses have dealt with this point. A modelling analysis demonstrated that, in long time-life cyclonic eddies of the Kuroshio current (time-life superior than one year), spring phytoplankton bloom is reinforced thanks to larger nutrient supply associated to vertical mixing (Sasai et al., 2010). In the North Western Mediterranean Sea, D'Ortenzio et al. (2014), using Bio-Argo data, suggested that large scale cyclonic circulation could be the first forcing factor of the phytoplankton bloom, and demonstrated that deep winter convection is not essential for bloom development. To further investigate the influence of upwelling (downwelling) of the nutricline on the phytoplankton seasonal cycle, we present an analysis of the phytoplankton phenology during the cyclonic and anticyclonic phases of the North Ionian Gyre (NIG, see Figure 1). Thus, our aim is to determine whether and how changes in cyclonic/anticyclonic circulation can affect interannual variability in phytoplankton phenology.

The North Ionian Sea has been selected as study area because this region of the Eastern Mediterranean Sea has the specificity to experience reversals of its surface circulation at decadal time-scale (Gacic et al., 2010). More specifically, due



- to the feedback with the Adriatic Sea, the North Ionian circulation oscillates between the cyclonic and the anticyclonic modes (Figure 1b and 1c). Variations of horizontal circulation patterns have been detected mainly from altimetric measurements, while the vertical structure of the Ionian circulation have been inferred only recently using float and oceanographic campaign data (Gacic et al., 2014). It has been shown that the Ionian behaves as a two-layer fluid and that the
- 5 decadal inversions of the NIG are generated by the spreading of the dense water formed in the Adriatic (AdDW). During the Eastern Mediterranean Transient, in the early 1990's, large amount of dense water of Aegean origin (about 3 Sv) contributed both to reinforce the anticyclonic circulation and to generate the reversal (Borzelli et al., 2009). Dense water of interannual varying density, changes the horizontal pressure gradient in the lower layer affecting the horizontal circulation over the entire water column. The mechanism at the base of the decadal reversal of the circulation regime was called Adriatic-Ionian
- 10 Bimodal Oscillating System (BiOS) (Gacic et al., 2010). From a limited number of nitrate vertical profiles, Civitarese et al. (2010) showed that NIG circulation reversals influence North Ionian biogeochemistry and especially the depth of the nitracline. However, the impact on phytoplankton remained unclear. Comparing surface chlorophyll-a measured by the CZCS (1979-1985) and SeaWiFS (1997-2000) ocean color sensors, D'Ortenzio et al. (2003) observed just a slight difference in the timing of main bloom events and, during the SeaWiFS period, a recurrent and large patch of [Chl-a]. More recently,
- 15 Mayot et al. (2016) showed that phytoplankton phenology is extremely dynamic in the North Ionian Sea with significant changes in the annual cycle of surface chlorophyll-a concentration from year to year. However, the authors did not explain this interannual variability. The present analysis attempts then to reconsider the impact of NIG circulation on phytoplankton phenology focusing on the satellite chlorophyll-a concentration (hereafter [Chl-a]) phenology and analyzing the 1998-2012 period.
- 20 In section 2, all the datasets used and the related methods are presented. Subsequently, temporal variability of the NIG circulation is analyzed and then the uplift (downlift) of isopycnals and nitracline during cyclonic (anticyclonic) regime is discussed. From the analysis of phenological metrics, differences are sought between cyclonic and anticyclonic regimes in [Chl-a] phenology. Phenological metrics are used here because they provide a synthetic method to highlight specific features of the [Chl-a] annual time-series. These metrics also demonstrated to be extremely informative over different and various
- 25 environments and regions (Follows and Dutkiewicz, 2001; Henson et al., 2009; Lavigne et al., 2013) despite the fact that they can be affected by data gaps (Cole et al., 2012) and noise (Ji et al., 2010). Focusing then on the initiation dates of [Chl-a] increase, the impact of interannual variability in the North Ionian circulation and winter buoyancy fluxes are compared and discussed. Finally, hypotheses about causes and consequences of the observed patterns in [Chl-a] phenology are proposed.



## 2 Data and Methods

### 2.1 Satellite and modelling data

- Daily [Chl-a] products, with a 4-km resolution, delivered by the Ocean Colour Climate Change Initiative (OC\_CCI) project version 1.0 (<http://www.esa-oceancolour-cci.org/>) were used. The OC\_CCI dataset, which covers the period September 1997 to July 2012, was created by band-shifting and bias-correcting MERIS and MODIS data to match SeaWiFS data and by merging the datasets with a simple average (Storm et al., 2013). Daily data were merged to obtain 8-day resolution maps. During the period from 1997 to March 2002 (only the SeaWiFS sensor available), 8-day average maps time-series contains only 80% of data but from April 2002, more than 96% of the data was available. To fill the gaps, linear interpolation was applied to each time-series. Then, time-series were then smoothed with a 3-point median filter.
- The Ssalto-Duacs Delayed-Time maps of the Absolute Dynamic Topography (ADT) (Rio et al., 2007) for the Mediterranean Sea ( $1/8^\circ$ ) computed and distributed by AVISO (<http://www.aviso.altimetry.fr/>) were downloaded for the period 1993-2014. As proposed by Gacic et al. (2014), the sea level gradient between the north and the center of the NIG was used to determine the cyclonic/anticyclonic nature of the NIG circulation. In practice, the ADT difference between rectangles S2 ( $18.75^\circ\text{E}$ - $19.25^\circ\text{E}$  and  $39.5^\circ\text{N}$ - $40^\circ\text{N}$ , see Figure 1b) and S1 ( $18.5^\circ\text{E}$ - $19^\circ\text{E}$  and  $37.5^\circ\text{N}$ - $38^\circ\text{N}$ , see Figure 1b) was computed. Then, the daily time-series was smoothed with a 30-day moving average filter (dotted line Figure 2a). As the anticyclonic (cyclonic) circulation regime is characterized by a doming (depression) of sea level, it is characterized by positive (negative) ADT difference value.
- Net air-sea heat fluxes of long-wave radiation and short-wave radiation, sensible heat flux, latent heat flux and precipitations were obtained from the European Center for Medium Range Weather Forecast (ECMWF) ERA-Interim reanalysis (Dee et al., 2011). Data was available at 6-hour intervals and downloaded at  $0.5^\circ$ -resolution. Then, daily averages were computed and the net heat flux was calculated as the sum of the four air-sea fluxes components. The surface buoyancy flux per unit area was computed according to Anderson et al. (1996), with coefficients proposed by Gacic et al. (2002) for the Mediterranean Sea. Total winter buoyancy losses were then calculated as time-integral for the period from 21 December to 20 March.
- Daily temperature and salinity profiles for the product MEDSEA\_REANALYSIS\_PHY\_006\_004 produced by the Mediterranean Forecasting System and delivered by the CMEMS service (<http://marine.copernicus.eu>) were used to derive the depth of the isopycnal  $28.9 \text{ kg m}^{-3}$  and MLD for the time range 1993-2012. MEDSEA\_REANALYSIS\_PHY\_006\_004 product results from a hydrodynamic model, supplied by the Nucleus for European Modelling of the Ocean (NEMO), with a variational data assimilation scheme for temperature and salinity vertical profiles and satellite Sea Level Anomaly along track data. The model horizontal grid resolution is  $1/16^\circ$  and the unevenly spaced vertical levels are 72. An evaluation of the product performances showed that temperature and salinity present an RMS of  $0.34^\circ$  and 0.1 psu, respectively, on average on the water column (Fратиanni et al., 2016).



## 2.2 In-situ data

71243 temperature and salinity profiles (52% from Argo floats) were collected from Coriolis web portal (<http://www.coriolis.eu.org/>) during the period 1990 from 2012, in the region comprised between 36°N-40°N and 16°E-21°E. Only the best quality (Coriolis sensu) data were retained. Then, potential density profiles were calculated and the depth of the isopycnal 28.9 kg m<sup>-3</sup> was determined.

Nitrate concentration, temperature and salinity profiles from oceanographic cruises in the area of interest (36°N-40°N, 17°E-22°E) were obtained: from POEM-BC-O91 (October 1991, 55 profiles), POEM-BC-A62 (April-May 1992, 40 profiles) and EMTEC (April-May 1999, 20 profiles). From nitrate concentration profiles, the nitracline depth was estimated as the depth of the isoline 1μM and compared to the depth of isopycnals 28.9 kg m<sup>-3</sup> (Figure 3c). The isopycnal 28.9 kg m<sup>-3</sup> was chosen here because it is located within the transition layer between the Atlantic Waters (AW, density ranging between 27.4 and 28.6 kg m<sup>-3</sup>) which are generally nutrient depleted and the Levantine Intermediate Waters (LIW, density ranging between 29.0 and 29.1 kg m<sup>-3</sup>) where nutrients accumulate (Malanotte-Rizzoli et al., 1997; Ribera d'Alcalà et al., 2003).

## 2.3 Phenological metrics

A set of phenological metrics has been computed from each annual [Chl-a] time-series. Annual time-series were defined from July of year n to June of year n+1, in order to better display the winter-to-spring transition. For each pixel, 14 annual time-series could be defined ranging from July 1998 to June 2012. Hereafter, the annual time-series are referred by year n+1 (e.g. 2000 for the annual time-series 1999-2000).

Phenological metrics were defined to describe the main characteristics of the phytoplankton annual cycle. They were selected to focus on the [Chl-a] content and on the date of the main growing period. Metrics CHL\_Year, CHL\_March and CHL\_Max refer to the yearly average [Chl-a], the March average [Chl-a] and the [Chl-a] annual maximum, respectively. CHL\_March has been defined here because previous studies demonstrated that in the Mediterranean Sea trophic situation is highly variable in March spatially and interannually ranging from oligotrophy to bloom conditions (D'Ortenzio et al., 2009; D'Ortenzio et al., 2003; Mayot et al., 2016). The bloom initiation was defined as the date of the maximum growth rate. Date of the maximum growth rate represents an objective and efficient way to determine bloom initiation date although the methodology can be affected by data noise and require then an efficient smoothing of the time-series (Brody et al., 2013). 8-day annual time-series were twice filtered with running means using a 3-point window which decreases small scale noise without strongly degrading the main pattern of the [Chl-a] annual time-series. Then, the middle date of 8-day interval corresponding to the maximum [Chl-a] increase was defined as Date\_GR\_Max. Finally, the metric indicating the date of maximum [Chl-a] is referred in the following by Date\_Max.



### 3 Results and Discussion

#### 3.1 Physical and chemical characterization of the NIG

Time-series of ADT difference between regions S1 and S2 (Figure 1b) shows decadal and seasonal variability (Figure 2). Globally, a first anticyclonic phase is evidenced from altimetry data (positive ADT difference) in 1992-1997. Anticyclonic phase is then followed by a cyclonic period (negative ADT difference) from 1998 through 2005. Then, a shorter anticyclonic period took place between 2006-2010. In 2012, circulation was again cyclonic. To determine the average circulation state during each blooming period, ADT difference values were averaged for a period ranging from October to March which largely covers the annual [Chl-a] increase (see Figure 4). Resulting values (grey circles on Figure 2a) are referred in the following as ICI, Ionian Circulation Index. Although, ICI shows an interannual variability similar to the annual averages (July to June, black diamonds on Figure 2a), it is affected by seasonal signal which has to be considered for the determination of the circulation regime during transitional phases (i.e., when ADT differences are close to zero). During these phases, the Oct.-March ADT maps (Figure 2, panels b to f) were also investigated since the impact of the small scale structures on the ADT difference criteria could be substantial. From 2005 to 2006, circulation turned from cyclonic to anticyclonic. On Figure 2b, the north of the Ionian domain is dominated by negative ADT values and, in the 34°-36°N band, a clear front, representing the Mid-Ionian Jet is observed. These patterns indicate a cyclonic circulation. In 2006 (see Figure 2c), high ADT values extend northward and the lowest ADT values are pushed along Italian and Greek coasts. This pattern represents the transition towards an anticyclonic regime, which is characterized by higher ADT value in the center of the North Ionian basin than at its periphery. From Figure 2d to Figure 2f, the transition from an anticyclonic (Figure 2d) to a cyclonic (Figure 2f) circulation regime is represented. Finally, the analysis of the ICI time-series as well as the examination of selected ADT brought us to the conclusion that from July 1998 to June 2005 and from July 2011 to June 2012, NIG circulation can be considered as cyclonic and that the period 1993-1997 and July 2007-June 2010 are clearly dominated by the anticyclonic regime. No definite circulation regime can be attributed to the years 2006 and 2011 as they correspond to transitional phases.

In the Coriolis database, 26318 potential density profiles were available to describe the anticyclonic situation (data collected between 1990 and 1996 and between 2006 and 2009) and 34753 to the cyclonic situation (data collected between 1998 and 2005 and between 2011 and 2012). Optimal interpolation applied to each dataset provides an estimation of the depth of the isopycnal 28.9 kg m<sup>-3</sup> for cyclonic and anticyclonic regimes (Figure 3 panels a and b, respectively). Resulting maps exhibit very different features. In the cyclonic pattern (Figure 3a), the depth of the isopycnal 28.9 kg m<sup>-3</sup> ranges between 40 and 120 m north of 37°N. South of 37°N, a steep gradient is observed, consistently with the position of Mid-Ionian Jet which is reinforced during the cyclonic regime of the NIG (Bessieres et al., 2013; Gacic et al., 2014). The shallowest isopycnals are located around 39°N and 18°E, offshore of the Italian coast, representing the center of the cyclone. During the anticyclonic phase, the isopycnal 28.9 kg m<sup>-3</sup> is very deep (240m) in the center of the Ionian Sea and is shallower at the border of the basin. This isopycnal depth pattern is consistent with an anticyclonic circulation.



As previously demonstrated for the global ocean (While and Haine 2010, Omand and Mahadevan, 2013), the depth of an isopycnal can be representative of the nitracline depth. This is confirmed in the North Ionian from the obtained linear relationship between the depth of the nitracline and the  $28.9 \text{ kg m}^{-3}$  isopycnal (Figure 3c,  $r^2=0.93$ ). Applying this relationship, it appears that in the center of the North Ionian, the nitracline depth ranges between 60 and 80m during the cyclonic regime whereas it reaches 150 m when NIG is anticyclonic. Considering that this nitracline depth range (i.e. 60 to 150 m) overlaps with climatological values of winter MLDs in the Ionian Sea (i.e. 80 - 130 m; D'Ortenzio et al., 2005; Houpert et al., 2015), changes in NIG circulation should have significant impact on the entrainment of nutrients to the surface layer by mixing and, consequently, on phytoplankton growth and primary production.

### 3.2 General patterns of phytoplankton phenology in the Ionian Sea

Phytoplankton phenology in the Ionian Sea is typical of very oligotrophic regions with low [Chl-a] (values range between 0.05 and  $0.5 \text{ mg m}^{-3}$  in open waters) and a seasonal cycle characterized by very low [Chl-a] values during summer and higher values during winter (see Figure 4, D'Ortenzio and Ribera d'Alcalà, 2009). Highest [Chl-a] values are observed along the coast, especially along the South Italian coast where a local upwelling can take place (D'Ortenzio et al. 2003). Offshore, a north-south gradient is observed (Figure 4 and 5a) with higher values in the northern part of the basin (above  $38^\circ\text{N}$ ). In the whole basin, lowest [Chl-a] are observed in mid-summer, then, [Chl-a] increases from October (Figure 4). Date\_GR\_Max occurs later, in December for the central Ionian Sea and in mid-January for the North Ionian and for the Italian coastal region (see Figure 5). Spatial variability in Date\_GR\_Max indicates that differences in [Chl-a] phenology should exist between northern and southern parts of the basin. This is confirmed by the analysis of the seasonal [Chl-a] pattern on the Hovmoeller diagram (Figure 4). Indeed, during some winters as for example 1999, 2000, 2003, 2005, there is a distinct local maximum in [Chl-a] generally centered on March. These March peaks are situated in the  $38^\circ\text{N}$ - $39^\circ\text{N}$  band and are never observed below  $37^\circ\text{N}$ . However, they are not observed in all situations and, for instance, no [Chl-a] peak appears in March 2001, 2007 or 2008. This special feature of the [Chl-a] seasonal cycle is related to the location of isopycnal doming area during cyclonic regime (Figure 3a), suggesting that interannual variability in [Chl-a] patterns in this region may be related to NIG circulation pattern.

### 3.3 Impact of the NIG circulation on the [Chl-a] phenology

Figure 5 displays interannual average, over period July 1998 – June 2012, of phenological metrics (left panels) as well as the anomalies for the cyclonic (1999-2005 and 2012), and the anticyclonic (2007-2010) regimes compared to interannual average. Regarding [Chl-a] content, no significant difference is observed for the annually averaged [Chl-a] content (i.e. CHL\_Year) with a relative difference ranging between -10% and +10%. During the anticyclonic regime, a small positive anomaly is generally observed along the coast while central Ionian displays negative anomalies. When only March period is considered, CHL\_March displays much stronger differences between cyclonic and anticyclonic regimes (panels e and f). Significant differences are observed in the center of the North Ionian Sea with a +15% anomaly during the cyclonic regime





and a -20% anomaly during the anticyclonic regime. In the South Ionian, no major differences are observed in March which is in agreement with the low interannual variability observed below 37°N (Figure 4). Similarly to CHL\_March, CHL\_Max shows significant differences in the central North Ionian with a positive or a negative anomaly during cyclonic and anticyclonic regimes, respectively. In South Ionian, the same feature is observed due to variability of the AW advection modulated by BiOS. Regarding metrics measuring the timing of the bloom, significant differences are again observed in the center of the North Ionian. Indeed, Date\_Max and Date\_GR\_Max are delayed by about 20 and 30 days, respectively, when NIG circulation is cyclonic. On the other hand, both are anticipated by more than 40 days when circulation is anticyclonic.

The present analysis suggests that in the North Ionian, two types of [Chl-a] phenology co-exist, one characterizing the cyclonic regime and the other one characterizing the anticyclonic regime. This is also confirmed by the analysis of annual [Chl-a] cycles in the region S3 (17.5°E-18.5°E, 38°N-39°N, Figure 6). We have chosen the position of the area S3 where to calculate spatially averaged phenological metrics so that it coincides with the region where the extremes of [Chl-a] anomalies occur (Figure 5). During the anticyclonic regime (2007-2010), all cycles show a regular increase in [Chl-a] from October to the end of January. Maximum growth rate appears between late November and late December, and maximum [Chl-a] between mid-January and mid-February (Table 1). During the cyclonic regime, as for the anticyclonic one, a first increase in [Chl-a] occurs in mid-September and extends up to January. In March, most of time-series display a narrow and strong peak in [Chl-a]. The annual maximum growth rate is clearly associated to this March peak occurring generally few days (10 days in average) before the date of [Chl-a] maximum. Although, the cyclonic mode is characterized by the March [Chl-a] peak and a higher amplitude of the annual [Chl-a], this does not significantly impact the yearly average [Chl-a].

Referring to the climatological analysis of the shapes of [Chl-a] annual cycles proposed by D'Ortenzio and Ribera d'Alcalà (2009), the anticyclonic cycles (Figure 6) are close to the so-called "NO BLOOM" shape whereas most of the cyclonic cycles could be assimilated to the "INTERMITTENT" [Chl-a] annual cycle. Lavigne et al. (2013) interpreted the "INTERMITTENT" dynamics as an intermediate state between the "NO BLOOM" dynamics and the typical dynamics of spring bloom as it can be observed in the north-western Mediterranean Sea (Marty et al., 2002). The "NO BLOOM" shape was identified in areas where winter mixing rarely reaches the nitracline depth and where light is always sufficient to support phytoplankton growth (Lavigne et al., 2013).

### 3.4 Role of the NIG circulation compared to the interannual variability in MLD (focus on the region S3)

In the North Ionian where the bimodal behavior of the circulation (i.e. cyclonic versus anticyclonic) has a direct influence on the phytoplankton phenology (see Section 3.3), interannual variability also exists (see Table 1 and Figure 6). Still, focusing on the area S3, the interannual variability of phenological metrics is compared to variability of ICI (see definition Section 3.1) and total winter buoyancy losses. The total winter buoyancy losses are considered here because they largely contribute to the formation of the mixed layer (Gill, 1982) and are thus expected to affect the bloom development (Follows and Dutkiewicz, 2001). Regarding the cyclonic regime, the main feature in [Chl-a] phenology is the absence of late [Chl-a] peak in March in 2001 and 2002 (Figure 6, Table1). Years 2001 and 2002 exhibit the lowest values for CHL\_March and





CHL\_Max and more importantly, the earliest dates for Date\_Max and Date\_GR\_Max which were anticipated to January-February and December-January, respectively (Table 1). This special pattern may be explained by the buoyancy losses which were relatively weak: -24% (2001) and -35% (2002) compared to the average for the period 1999-2012. In 2001 and 2002, shallow winter MLD probably prevented significant nutrients inputs to surface layer and consequently the spring phytoplankton bloom (see discussion below). On the other hand, in 2005 and 2012 winter buoyancy losses were particularly high (+42%, +43%, respectively) and CHL\_Max and CHL\_March show the highest values observed. More generally, we observed a positive spearman correlation between the buoyancy flux anomaly and CHL\_March (correlation = 0.79) or CHL\_Max (correlation = 0.89), suggesting that in addition to circulation regime, winter buoyancy losses affect the presence and the strength of the [Chl-a]. Regarding the anticyclonic regime and the transitional years 2006, 2011, a very low interannual variability is observed, only the year 2006 (transitional regime) displays a [Chl-a] peak in early March (Table 1) which coincides with a ICI value close to 0 (transitional regime) and a weak positive anomaly in buoyancy losses (+6%, Table 1). During the anticyclonic regime, no significant correlation was found between buoyancy losses and phenological metrics. To sum up, in the center of the North Ionian, both circulation pattern and air-sea exchanges would affect [Chl-a] cycles. Nevertheless, it seems that the impact of air-sea exchanges may be stronger when circulation is cyclonic and apparently affect the amplitude of the March peak.

Given the general oligotrophy of the Eastern Mediterranean (Ribera d'Alcalà et al., 2003), horizontal nutrient advection in the Ionian Sea should not have any significant impact on the late winter/early spring [Chl-a] peak observed during the cyclonic regime. However, the analysis of Table 1 as well as the spatial correspondence between phenological anomalies and isopycnals uplift/downlift suggest that, in the North Ionian Sea, the March bloom would be favored by cyclonic circulation which tends to uplift isopycnals and nitracline depth towards the surface and by strong negative winter buoyancy fluxes which contribute to deepen the MLD. In other words, North Ionian phytoplankton phenology is driven by the interactions between the nutrient field and MLD which is consistent with sub-tropical regions where phytoplankton growth is almost permanently limited by nutrient availability (Longhurst, 2006). From these results, we propose two scenarios to explain interannual changes in phytoplankton phenology in the North Ionian Sea. These scenarios are based on relative position of the nitracline and the winter MLD. In the first scenario (scenario A), the deepest winter MLD remains above the nitracline depth and there is no spring phytoplankton bloom whereas, in the second scenario (scenario B), winter MLD overpasses the fall nitracline depth which drives remarkable nutrient supplies to the surface and supports the development of a March bloom. The first scenario results then in a deep nitracline depth and/or a shallow MLD. Conversely, scenario B occurs if the nitracline depth is shallow and/or if the MLD is deep.

Scenario B was observed in some subtropical regions, like the BATS station in the Subtropical North Atlantic (Steinberg et al., 2001). There, vertical mixing represents one of the major processes to supply nutrients to the enlighten surface layer (Steinberg et al., 2001; Hansell and Carlson 2001). At lower latitude and generally in trades biome regions according to Longhurst (2006), trophic regimes close to scenario A are observed. For instance, at the station HOT in the North Pacific Subtropical gyre (Karl and Lucas, 1996), winter mixed layers rarely reach the deep nutrient depleted waters and a deep



chlorophyll maximum is almost permanent (Letelier et al., 2004). The rate of primary production is then explained by stochastic nutrient injections into the euphotic zone caused, for instance, by internal waves, cyclonic mesoscale eddies or wind driven Ekman pumping (Hayward, 1991; Karl 1999). In the Mediterranean Sea, Lavigne et al. (2013) showed that in the South Ionian Sea and in the Levantine Sea, the probability that MLD reaches the nitracline depth is very small. These results were corroborated by a recent analysis based on Bio-Argo floats (Pasqueron de Fommervault, 2015). However, in regions where meso-scale circulation is cyclonic (e.g. South Adriatic gyre, Rhodes gyre), there are evidences of surface nutrient enrichment caused by vertical mixing (Gacic et al., 2002; Lascaratos et al., 1993; Salihoglu et al., 1990). Thus, the Eastern Mediterranean Sea likely holds both trophic regimes; one involving subsequent surface nutrients supplies by winter vertical mixing and the other not.

- Figure 7 supports our previous hypothesis to explain interannual variability in phytoplankton phenology. Indeed, during the anticyclonic phases of the circulation regime (i.e. 1993-1997 and 2007-2010), estimated nitracline depth was relatively deep and ranged between 100 m and 150 m. Nitracline is never reached by the MLD estimated with difference density criterion of  $0.005 \text{ mg m}^{-3}$  (Brainerd and Gregg, 1995), and was rarely reached by the deepest MLD estimated with difference density criterion of  $0.03 \text{ mg m}^{-3}$  (de Boyer-Montégut et al., 2004). In addition, during the anticyclonic period, the isopycnal  $28.9 \text{ kg m}^{-3}$  never intersected the surface indicating that winter mixing is not able to penetrate this layer. During the cyclonic phases (1999-2005 and 2012), the nitracline was shallower (generally less than 100 m). Except for 2001, the shallowest MLD estimations exceeded the fall nitracline depth and reached almost the surface indicating a penetration of mixing up to the isopycnal  $28.9 \text{ kg m}^{-3}$  layer and a possible nutrient enrichment.

- A common explanation for the fall and spring bloom is the succession of a entrainment bloom (nutrient limited) in fall followed by a spring bloom (light limited, Cullen et al. 2002; Levy et al., 2005). The initiation of the spring bloom, which is initially prevented by a deficiency of light in the surface mixed layer, is generally explained by a stratification or a stabilization of the water column. Stratification can be caused by the shallowing of the mixed layer (Sverdrup, 1953), or, as recently suggested, by submesoscale eddies (Mahadevan et al., 2012). Other studies have shown that phytoplankton bloom can be induced by a decrease of the water column turbulence (Huisman et al., 1999) in response, for instance, to the inversion of the surface heat fluxes (Taylor and Ferrari, 2011). However, previous hypotheses about fall and spring phytoplankton blooms were developed for temperate and sub-polar regions and do not fit with the light and nutrient conditions of the Ionian Sea. Indeed, on one hand, in the Ionian Sea, winter solar irradiation is sufficient for supporting phytoplankton growth, according to the Sverdrup (1953) approach (Lavigne et al., 2013). On the other hand, Figure 7 shows that when the first Chl-a increase is observed in fall (October to December), MLD is shallower than the nitracline depth making the entrainment bloom hypothesis an unconvincing explanation for the fall [Chl-a] rise. A possible explanation for the early increase in [Chl-a] could be sought in the seasonal increase of Chl-a content per cell in response to a diminution of the light intensity (Winn et al., 1995; Behrenfeld et al., 2005). Indeed, from a modelling approach, Taylor et al. (1997) showed that the phytoplankton carbon to chlorophyll ratio varies by a factor of 2 in subtropical regions ( $\sim 35^\circ$ ) from early fall to December.



Spring peaks in [Chl-a], mainly observed during cyclonic circulation regime, can be considered as typical spring blooms (i. e. detrainment blooms, Cullen et al., 2002) which are light driven. Indeed, Date\_GR\_Max was sometimes observed during the MLD shallowing. However, the tight coupling between MLD deepening (deeper than nitracline) and [Chl-a] increase as well as the lack of light limitation in the Eastern Mediterranean (Lavigne et al., 2013), make us conclude that [Chl-a] increase is the result of a late winter entrainment bloom (nutrient driven). In addition, as the date of the bloom is determined from surface [Chl-a], the dilution effect during the mixed layer deepening masks the phytoplankton accumulation, causing an artificial delay in the computed date of the bloom (Behrenfeld et al., 2010).

#### 4 Summary and Conclusion

In this study we show that in the Ionian Sea two trophic regimes co-exist. In one case, the absence of spring bloom peak in March indicates the lack of nutrient supply in the surface layer. In that case, the rate of regenerated production is high all along the year and new primary production would only be sustained by sporadic nutrient injections through the nutricline. In the second case, the presence of a [Chl-a] peak in March could be explained by a significant nutrient supply by vertical mixing which sustains phytoplankton bloom and the seasonal boost in new primary production. We showed that these differences in the phytoplankton dynamics were associated with the variability of the circulation regime of the NIG. Namely, the March [Chl-a] peak was commonly observed during the cyclonic phases, whereas no peak was observed in case of anticyclonic circulation. In addition to the strength and sense of the circulation, the [Chl-a] annual cycle is affected also by the winter buoyancy losses, which determine the MLD. Thus, it appears that phytoplankton phenology is the result of the interplay between surface circulation and atmospheric conditions, affecting the nitracline depth and the MLD. Hypotheses based on the relative position of the nitracline depth and winter deepest MLD were then proposed to explain observed phenological patterns.

The dependence of phytoplankton phenology on the circulation regime in the Ionian (driven by BiOS) establishes a link between phytoplankton dynamics and the semi-closed thermohaline cell of the Eastern Mediterranean. As shown previously (Gacic et al., 2014), the Adriatic Sea, where the thermohaline cell of the Eastern Mediterranean originates, is highly sensitive to the winter air-sea heat fluxes: for example, the exceptionally severe winter of 2012 resulted in the occasional production of large amount of very high dense water in the Adriatic, whose spreading in the deep Ionian caused a partial reversal of the circulation.

In conclusion, future studies on phytoplankton dynamics in the Ionian Sea should not take into account just the local circulation regime, but winter climatic conditions over the Adriatic Sea as well.



## Aknowledgements

The authors thank the ESA Ocean Colour CCI Team for providing OC-CCI chlorophyll data; NASA for providing SeaWiFS, MODIS and MERIS chlorophyll data; ECMWF for providing meteorological data and AVISO with support from CNES for distributing altimeter products produced by Ssalto/Duacs. The Coriolis Centre for in situ Oceanographic Data and the  
5 National Oceanographic Data Center of the OGS (NODC-OGS) are acknowledged for the collection of the CTD and nitrate profiles that were made available to this study. This study has been conducted using E.U. Copernicus Marine Service Information. We are also grateful to Dhyan Aranha for proof-reading this manuscript. We acknowledge the European Commission "Cofunded by the European Union under FP7- People - Co-funding of Regional, National and International Programmes, GA n. 600407" and of the RITMARE Flagship Project, for the support to H. L. during her stay at OGS. This  
10 work is also the contribution to the French "Equipementd'Avenir" NAOS project (ANR J11R107-F) and to the PERSEUS EU Project. M.G. was partially supported by Croatian Science Foundation under theproject SCOOL (IP-2014-09-5747).

## References

- Anderson, S. P., Weller, R. A., and Lukas, R. B.: Surface buoyancy forcing and the mixed layer of the Western Pacific warm pool: observations and 1d model results, *Journal of Climate*, 9(12), 3056–3085, 1996.
- 15 Behrenfeld, M. J.: Abandoning sverdrup's critical depth hypothesis on phytoplankton blooms, *Ecology*, 91(4), 977–989, doi: 10.1890/09-1207.1, 2010.
- Behrenfeld, M. J., Boss, E., Siegel, D.A., and Shea, D. M.: Carbon-based ocean productivity and phytoplankton physiology from space, *Global Biogeochemical Cycles*, 19(1), GB1006, doi: 10.1029/2004GB002299, 2005.
- Bessi eres, L., Rio, M., Dufau, C., Boone, C., and Pujol, M.: Ocean state indicators from myocean altimeter products, *Ocean*  
20 *Science*, 9(3), 545, 2013.
- Borzelli, G. L. E., Gacic, M., Cardin, V., and Civitarese G.: Eastern Mediterranean transient and reversal of the Ionian sea circulation, *Geophysical Research Letters*, 36(15), 2009.
- Brainerd, K. E. and Gregg, M. C.: Surface mixed and mixing layer depths, *Deep Sea Research Part I: Oceanographic Research Papers*, 42(9), 1521–1543, doi: 16/0967-0637(95)00068-H, 1995.
- 25 Brody, S. R., Lozier, M. S., and Dunne, J. P.: A comparison of methods to determine phytoplankton bloom initiation, *Journal of Geophysical Research: Oceans*, 2013.
- Buesseler, K. O.: The decoupling of production and particulate export in the surface ocean, *Global Biogeochemical Cycles*, 12(2), 297–310, 1998.
- Civitarese, G., Gacic, M., Lipizer, M., and Eusebi Borzelli, G. L.: On the impact of the Bimodal Oscillating System (BiOS)  
30 on the biogeochemistry and biology of the Adriatic and Ionian seas (Eastern Mediterranean), *Biogeosciences*, 7, 3987–3997, 2010.



- Cole, H., Henson, S., Martin, A., and Yool, A.: Mind the gap: The impact of missing data on the calculation of phytoplankton phenology metrics, *Journal of Geophysical Research: Oceans* (1978–2012), 117(C8), 2012.
- Cullen, J. J., Franks, P., Karl, D., and Longhurst, A.: Physical influences on marine ecosystem dynamics, in *The sea, ideas and observations on progress in the study of the seas*, vol. 12, A.R. Robinson, J. McCarthy, and B.J. Rothschild eds., pp. 297–336, John Wiley & Sons Inc, 2002.
- de Boyer Montegut, C., Madec, G., Fischer, A., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: an examination of profile data and a profile-based climatology, *Journal of Geophysical Research-Oceans*, 109(C12), doi: 10.1029/2004JC002378, 2004.
- Dee, D., et al.: The era-interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137(656), 553–597, 2011.
- D’Ortenzio, F., Ragni, M., Marullo, S., and Ribera d’Alcalá, M.: Did biological activity in the Ionian sea change after the Eastern Mediterranean Transient? Results from the analysis of remote sensing observations, *Journal of Geophysical Research: Oceans* (1978–2012), 108(C9), 2003.
- D’Ortenzio, F., Iudicone, D., Montegut, C., Testor, P., Antoine, D., Marullo, S., Santoleri, R., and Madec, G.: Seasonal variability of the mixed layer depth in the Mediterranean sea as derived from in situ profiles, *Geophysical Research Letters*, 32(12), doi: 10.1029/2005GL022463, 2005.
- D’Ortenzio, F. and Ribera d’Alcalá, M.: On the trophic regimes of the Mediterranean sea: a satellite analysis, *Biogeosciences*, 6(2), 139–148, doi: 10.5194/bg-6-139-2009, 2009.
- D’Ortenzio, F., et al.: Observing mixed layer depth, nitrate and chlorophyll concentrations in the Northwestern Mediterranean: A combined satellite and no3 profiling floats experiment, *Geophysical Research Letters*, 2014.
- Edwards, M. and Richardson A. J.: Impact of climate change on marine pelagic phenology and trophic mismatch, *Nature*, 430(7002), 881–884, doi: 10.1038/nature02808, 2004.
- Falkowski, P. G., Ziemann, D., Kolber, Z., and Bienfang, P. K.: Role of eddy pumping in enhancing primary production in the ocean, *Nature*, 352, 55–58, 1991.
- Follows, M. and Dutkiewicz, S.: Meteorological modulation of the North Atlantic spring bloom, *Deep Sea Research Part II: Topical Studies in Oceanography*, 49(1–3), 321–344, doi: 10.1016/S0967-0645(01)00105-9, 2001.
- Fратиanni, C., Clementi, E., and Simoncelli, S.: Quality information document for Mediterranean physics reanalysis product: *Medsea\_reanalysis\_phy\_006\_004*, Tech. Rep. CMEMS-Med-QUID-006-004-V2, MED-MFC, 2016.
- Gacic, M., et al.: Extreme winter 2012 in the Adriatic: an example of climatic effect on the BiOSrhythm, *Ocean Science*, 10(3), 513–522, 2014.
- Gacic, M., Borzelli, G., Civitarese, G., Cardin, V., and Yari, S.: Can internal processes sustain reversals of the ocean upper circulation? The Ionian sea example, *Geophysical Research Letters*, 37(9), 2010.



- Gacic, M., Civitarese, G., Miserocchi, S., Cardin, V., Crise, A., and Mauri, E.: The open-ocean convection in the Southern Adriatic: a controlling mechanism of the spring phytoplankton bloom, *Continental Shelf Research*, 22(14), 1897–1908, doi: 10.1016/S0278-4343(02)00050-X, 2002.
- Gill, A. E.: *Atmosphere-ocean dynamics*, vol. 30, Academic press, 1982.
- 5 Hansell, D. A. and Carlson, C. A.: Biogeochemistry of total organic carbon and nitrogen in the Sargasso Sea: control by convective overturn, *Deep Sea Research Part II: Topical Studies in Oceanography*, 48(8), 1649–1667, 2001.
- Hayward, T. L.: Primary production in the North Pacific central gyre: a controversy with important implications, *Trends in Ecology & Evolution*, 6(9), 281 – 284, doi: [http://dx.doi.org/10.1016/0169-5347\(91\)90005-I](http://dx.doi.org/10.1016/0169-5347(91)90005-I), 1991.
- Henson, S. A., Robinson, I., Allen, J. T., and Waniek, J. J.: Effect of meteorological conditions on interannual variability in
- 10 timing and magnitude of the spring bloom in the Irminger basin, North Atlantic, *Deep Sea Research Part I: Oceanographic Research Papers*, 53(10), 1601–1615, doi: 10.1016/j.dsr.2006.07.009, 2006.
- Henson, S. A., Dunne, J. P., and Sarmiento, J. L.: Decadal variability in North Atlantic phytoplankton blooms, *Journal of Geophysical Research*, 114, C04,013, 2009.
- Houpert, L., Testor, P., de Madron, X. D., Somot, S., D’Ortenzio, F., Estournel, C., and Lavigne, H.: Seasonal cycle of the
- 15 mixed layer, the seasonal thermocline and the upper-ocean heat storage rate in the Mediterranean Sea derived from observations, *Progress in Oceanography*, 132, 333–352, 2015.
- Huisman, J., Oostveen, P. V., and Weissing, F. J.: Critical depth and critical turbulence: two different mechanisms for the development of phytoplankton blooms, *Limnology and Oceanography*, 44(7), 1781–1787, 1999.
- Ji, R., Davis, C. S., Chen, C., Townsend, D. W., Mountain, D. G., and Beardsley, R. C.: Modeling the influence of low-
- 20 salinity water inflow on winter-spring phytoplankton dynamics in the Nova Scotian shelf–gulf of Maine region, *Journal of Plankton Research*, 30(12), 1399–1416, 2008.
- Ji, R., Edwards, M., Mackas, D. L., Runge, J. A., and Thomas, A. C: Marine plankton phenology and life history in a changing climate: current research and future directions, *Journal of Plankton Research*, 32(10), 1355–1368, doi: 10.1093/plankt/fbq062, 2010.
- 25 Kahru, M., Fiedler, P., Gille, S., Manzano, M. and Mitchell, B.: Sea level anomalies control phytoplankton biomass in the Costa Rica dome area, *Geophysical Research Letters*, 34(22), doi: 10.1029/2007GL031631, 2007.
- Karl, M. D.: A sea of change: biogeochemical variability in the North Pacific subtropical gyre, *Ecosystems*, 2(3), 181–214, doi: 10.1007/s100219900068, 1999.
- Karl, D. M. and Lukas, R.: The Hawaii Ocean Time-series (HOT) program: background, rationale and field implementation,
- 30 *Deep Sea Research Part II: Topical Studies in Oceanography*, 43(2-3), 129–156, doi: 10.1016/0967-0645(96)00005-7, 1996.
- Koeller, P., et al.: Basin-scale coherence in phenology of shrimps and phytoplankton in the North Atlantic ocean, *Science*, 324(5928), 791–793, doi: 10.1126/science.1170987, 2009.
- Lascaratos, A., Williams, R. G., and Tragou, E.: A mixed-layer study of the formation of Levantine Intermediate Water, *Journal of Geophysical Research*, 98(C8), PP. 14,739–14,749, doi: 10.1029/93JC00912, 1993.



- Lavigne, H., D'Ortenzio, F., Migon, C., Claustre, H., Testor, P., d'Alcalà, M. R., Lavezza, R., Houpert, L. and Prieur, L.: Enhancing the comprehension of mixed layer depth control on the Mediterranean phytoplankton phenology, *Journal of Geophysical Research: Oceans*, 118, 3416–3430, doi: 10.1002/jgrc.20251, 2013.
- Letelier, R. M., Karl, D. M., Abbott, M. R., and Bidigare, R. R.: Light driven seasonal patterns of chlorophyll and nitrate in the lower euphotic zone of the North Pacific subtropical gyre, *Limnology and Oceanography*, 49(2), 508–519, 2004.
- Lévy, M., Lehahn, Y., Andre, J.-M., Memery, L., Loisel, H., and Heifetz, E.: Production regimes in the northeast atlantic: a study based on Sea-viewing Wide Field-of-view Sensor (SeaWiFS) chlorophyll and ocean general circulation model mixed layer depth, *Journal of Geophysical Research: Oceans*, 110(C7), 2156–2202, doi: 10.1029/2004JC002771, c07S10, 2005.
- Longhurst, A. R.: *Ecological Geography of the Sea*, 2 ed., Academic Press, 2006.
- 10 Mahadevan, A., D'Asaro, E., Lee, C., and Perry, M. J.: Eddy-driven stratification initiates north Atlantic spring phytoplankton blooms, *Science*, 337(6090), 54–58, 2012.
- Malanotte-Rizzoli, P., et al.: A synthesis of the Ionian Sea hydrography, circulation and water mass pathways during POEM-Phase-I, *Progress In Oceanography*, 39(3), 153–204, doi: 10.1016/S0079-6611(97)00013-X, 1997.
- Mann, K., and Lazier, J.: *Dynamics of marine ecosystems: biological-physical interactions in the oceans*, Wiley-Blackwell, 15 2006.
- Marty, J.-C., Chiavérini, J., Pizay, M.-D., and Avril, B.: Seasonal and interannual dynamics of nutrients and phytoplankton pigments in the Western Mediterranean Sea at the DYFAMED time-series station (1991-1999), *Deep Sea Research Part II: Topical Studies in Oceanography*, 49(11), 1965–1985, doi: 10.1016/S0967-0645(02)00022-X, 2002.
- Mayot, N., D'Ortenzio, F., Ribera d'Alcala, M., Lavigne, H., and Claustre, H.: Interannual variability of the Mediterranean trophic regimes from ocean color satellites, *Biogeosciences*, 13, 1901–1917, doi:10.5194/bg-13-1901-2016, 2016.
- 20 McGillicuddy, D., Robinson, A., Siegel, D., Jannasch, H., Johnson, R., Dickey, T., McNeil, J., Michaels, A., and Knap, A.: Influence of mesoscale eddies on new production in the Sargasso Sea, *Nature*, 394(6690), 263–266, 1998.
- Omand, M. M. and Mahadevan, A.: Large-scale alignment of oceanic nitrate and density, *Journal of Geophysical Research: Oceans*, 118(10), 5322–5332, 2013.
- 25 Pasqueron de Fommervault, O., D'Ortenzio, F., Mangin, A., Serra, R., Migon, C., Claustre, H., Lavigne, H., Ribera d'Alcalà, M., Prieur, L., Taillandier, V., Schmechtig, C., Poteau, A., Leymarie, E., Besson F., and Obolensky, G.: Seasonal variability of nutrient concentrations in the Mediterranean Sea: contribution of Bio-Argo floats. *Journal of Geophysical Research*, 120, doi:10.1002/2015JC011103, 2015.
- Platt, T., Sathyendranath, S., White III, G. N., Fuentes-Yaco, C., Zhai, L., Devred, E., and Tang, C.: Diagnostic properties of phytoplankton time series from remote sensing, *Estuaries and Coasts*, 33(2), 428–439, 2010.
- 30 Racault, M.-F., Le Quéré, C., Buitenhuis, E., Sathyendranath, S., and Platt, T.: Phytoplankton phenology in the global ocean, *Ecological Indicators*, 14(1), 152–163, doi: 10.1016/j.ecolind.2011.07.010, 2012.
- Ribera d'Alcalà, M., Civitarese, G., Conversano, F., and Lavezza, R.: Nutrient ratios and fluxes hint at overlooked processes in the Mediterranean sea, *Journal of Geophysical Research*, 108, 16 PP., doi: 10.1029/2002JC001650, 2003.





- Rio, M.-H., Poulain, P.-M., Pascal, A., Mauri, E., Larnicol, G., and Santoleri, R.: A Mean Dynamic Topography of the Mediterranean Sea computed from altimetric data, in-situ measurements and a general circulation model, *J. Marine Syst.*, 65, 484–508, 2007.
- Salihoglu, I., Saydam, C., Basturk, Ö., Yilmaz, K., Gocmen, D., Hatipoglu, E., and Yilmaz, A.: Transport and distribution  
5 of nutrients and chlorophyll-a by mesoscale eddies in the northeastern Mediterranean, *Marine chemistry*, 29, 375–390, 1990.
- Sasai, Y., Richards, K. J.: A. Ishida, and H. Sasaki, Effects of cyclonic mesoscale eddies on the marine ecosystem in the Kuroshio extension region using an eddy-resolving coupled physical-biological model, *Ocean Dynamics*, 60(3), 693–704, 2010.
- Steinberg, D., Carlson, C., Bates, N., Johnson, R., Michaels, A., and Knap, A.: Overview of the US JGOFS Bermuda  
10 Atlantic Time-series Study (BATS): a decade-scale look at ocean biology and biogeochemistry, *Deep-Sea Research Part II-Topical Studies in Oceanography*, 48(8-9), 1405–1447, 2001.
- Storm, T., Boettcher, M., Grant, M., Zuhlke, M., Fomferra, N., Jackson, T., and Sathyendranath, S.: Ocean colour climate change initiative (oc\_cci) - phase one - product user guide, Tech. rep., Plymouth Marine Laboratory, 2013.
- Sverdrup, H. U.: On conditions for the vernal blooming of phytoplankton, *ICES Journal of Marine Science*, 18, 287–295,  
15 1953.
- Taylor, A., Geider, R. J., and Gilbert, F.: Seasonal and latitudinal dependencies of phytoplankton carbon-to-chlorophyll a ratios: results of a modelling study, *Marine Ecology-Progress Series*, 152, 51–66, 1997.
- Taylor, J. and Ferrari, R.: Shutdown of turbulent convection as a new criterion for the onset of spring phytoplankton blooms, *Limnol. Oceanogr.*, 56(6), 2293–2307, 2011.
- 20 While, J. and Haines, K.: A comparison of the variability of biological nutrients against depth and potential density, *Biogeosciences*, 7(4), 1263–1269, 2010.
- Williams, R. G. and Follows, M. J.: Physical transport of nutrients and the maintenance of biological production, in *Ocean Biogeochemistry*, pp. 19–51, Springer, 2003.
- Winn, C. D., Campbell, L., Christian, J. R., Letelier, R. M., Hebel, D. V., Dore, J. E., Fujieki, L., and Karl, D. M.: Seasonal  
25 variability in the phytoplankton community of the north pacific subtropical gyre, *Global Biogeochemical Cycles*, 9(4), 605–620, doi: 10.1029/95GB02149, 1995.
- Winder, M. and Cloern, J. E.: The annual cycles of phytoplankton biomass, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1555), 3215–3226, doi: 10.1098/rstb.2010.0125, 2010.
- Yoder, J. A. and Kennelly, M. A.: Seasonal and ENSO variability in global ocean phytoplankton chlorophyll derived from 4  
30 years of SeaWiFS measurements, *Global Biogeochemical Cycles*, 17(4), doi: 10.1029/2002GB001942, 2003.
- Yoder, J. A., McClain, C. R., Feldman, G. C., and Esaias, W. E.: Annual cycles of phytoplankton chlorophyll concentrations in the global ocean: a satellite view, *Global Biogeochemical Cycles*, 7(1), PP. 181–193, doi: 199310.1029/93GB02358, 1993.



Zhai, L., Platt, T., Tang, C., Sathyendranath, S., and Walls, R. H.: Phytoplankton phenology on the scotian shelf, ICES Journal of Marine Science: Journal du Conseil, 68(4), 781–791, 2011.



5 **Table 1: Average Ionian Circulation Index (ICI, ADT difference values between regions S1 and S2 averaged over October-March), winter buoyancy fluxes (WBF) and phenological metrics in the region S3. ICI is expressed in cm and WBF are expressed in percent relative difference compared to the 1999-2012 interannual average. CHL\_Year (yearly average [Chl-a]), CHL\_March (March average [Chl-a]) and CHL\_Max (annual maximum [Chl-a]) are expressed in  $\text{mg m}^{-3}$ . Numbers in brackets are standard deviations. For Date\_Max (date of annual maximum [Chl-a]), Date\_GR\_Max (date of maximum growth rate in [Chl-a]) standard deviations are expressed in days.**

year	regime	ICI	WBF	CHL_Year	CHL_March	CHL_Max	Date_Max	Date_GR_Max
1999	Cyclonic	-3.522	-7.7 (2.9)	0.19 (0.011)	0.36 (0.07)	0.45 (0.12)	16-March (37)	04-March (39)
2000	Cyclonic	-4.186	26.2 (3.2)	0.18 (0.008)	0.30 (0.04)	0.46 (0.16)	02-April (29)	12-March (38)
2001	Cyclonic	-1.826	-23.8 (0.8)	0.15 (0.008)	0.17 (0.01)	0.29 (0.03)	19-Jan. (34)	30-Dec. (28)
2002	Cyclonic	-2.905	-34.9 (0.6)	0.17 (0.011)	0.26 (0.03)	0.33 (0.05)	28-Feb. (14)	31-Jan. (41)
2003	Cyclonic	-3.966	5.3 (0.6)	0.17 (0.008)	0.26 (0.03)	0.39 (0.06)	02-April (2)	20-March (15)
2004	Cyclonic	-2.395	15.2 (2.2)	0.17 (0.008)	0.34(0.04)	0.46 (0.10)	24-March (6)	20-March (6)
2005	Cyclonic	-1.281	42.2 (3.1)	0.18 (0.013)	0.44 (0.06)	0.90 (0.19)	24-March (2)	20-March (3)
2006	Transition	+2.49	6.2 (1.8)	0.17 (0.014)	0.28 (0.03)	0.34 (0.06)	08-March (22)	31-Jan. (31)
2007	Anticyclonic	+7.5	-17.3 (2.0)	0.14 (0.019)	0.20 (0.03)	0.24 (0.04)	20-Feb. (26)	20-Nov. (39)
2008	Anticyclonic	+5.53	-23.9 (2.5)	0.16 (0.024)	0.20 (0.08)	0.32 (0.09)	19-Jan. (28)	30-Dec. (47)
2009	Anticyclonic	+2.73	12.4 (2.9)	0.16 (0.019)	0.20 (0.02)	0.36 (0.08)	04-Feb. (11)	06-Dec.(18)
2010	Anticyclonic	+0.75	-18.1 (3.2)	0.19 (0.025)	0.25 (0.03)	0.36 (0.08)	11-Jan. (11)	14-Dec. (26)
2011	Transition	+0.28	-26.3 (2.9)	0.18 (0.015)	0.22 (0.02)	0.33 (0.07)	18-Dec. (17)	12-Nov. (22)
2012	Cyclonic	-3.795	44.6 (3.8)	0.16 (0.010)	0.35 (0.07)	0.46 (0.13)	24-March (6)	03-Dec. (2)




Figure 1: Study area (panel a) and schematic representation of the North Ionian upper-layer circulation during anticyclonic (panel b) and cyclonic (panel c) regimes. AW, LSW and LIW refer as Atlantic Water, Levantine Surface Water and Levantine Intermediate Water, respectively. See text for a definition of regions S1, S2 (Section 2.1) and S3 (Section 3.2). The underlying maps represent the 1995 (b) and 1999 (c) average absolute dynamic topography, respectively.

5




Figure 2: Panel a: Dotted line is the ADT difference between regions S1 and S2 ( $S1 - S2$ , see Figure 1c). Annual periods used for the phenology analysis, which range from July of year  $n$  to June of year  $n+1$  are featured by white and grey bands. For each of them, black diamonds and grey circles represent the averaged ADT difference during periods October to March (also named ICI) and July to June, respectively. The zero level differentiates cyclonic (negative values) and anticyclonic (positive values) regimes. Panels b to f represent the average ADT pattern for different periods. Squares represent the regions S1 (south) and S2 (north)





Figure 3: Climatological maps of the estimation of the depth of the isopycnal  $28.9 \text{ kg m}^{-3}$  for cyclonic (panel a) and anticyclonic (panel b) regimes. Data were spatially interpolated with kriging. Relationship between the depth of the nitracline and the depth of isopycnal  $28.9 \text{ kg m}^{-3}$  in the North Ionian Sea (panel c) from in situ data collected during three oceanographic cruises (see colors).





**Figure 4:** Hovmöller diagram of the satellite [Chl-a] along the 18.5°E meridian. [Chl-a] is expressed in  $\text{mg m}^{-3}$ . Solid lines indicate 1-year periods (January to December) and dotted lines 3-month periods.








Figure 5: left panels: interannual average values (1998–2012) of phenological metrics. CHL\_Year, CHL\_March and CHL\_Max are expressed in  $\text{mg m}^{-3}$ . Middle and right panels show the anomaly of metrics compared to the interannual average for the cyclonic (middle panels) and anticyclonic (right panels) periods. They are expressed in percent difference for CHL\_Year, CHL\_March and CHL\_Max, and in days for Date\_Max and Date\_GR\_Max. Solid black lines refer to areas with significant difference between cyclonic and anticyclonic patterns ( $p\text{-value} < 0.05$ ).




Figure 6:  $[\text{Chl-a}] (\text{mg m}^{-3})$  annual cycles for the cyclonic and anticyclonic regimes averaged in the region S3 (see Figure 1).




Figure 7: Time-series, for the region S3, of the mean nitracline depth (solid black line), depth of the isopycnal  $28.9 \text{ kg m}^{-3}$  (solid grey line) and MLD computed with a density criteria of  $0.005 \text{ kg m}^{-3}$  (solid red line) and  $0.03 \text{ kg m}^{-3}$  (dotted red line) from a surface reference of 10m. Nitracline depth was derived from the depth of the isopycnal  $28.9 \text{ kg m}^{-3}$  with the linear relationship established on Figure 3c. Circles represent the mean Date\_GR\_Max dates for the region S3.

5