

"Interactive comment on "Simulating sustained yield harvesting adaptive to future climate change" by Rasoul Yousefpour et al."

Anonymous Referee #1

Received and published: 15 February 2018

Report #1

Submitted on 12 Jun 2018 Anonymous Referee #1	
Anonymous during peer-review: Yes No Anonymous in acknowledgements of published article: Yes No	
Recommendation to the editor	
1) Scientific significance Does the manuscript represent a substantial contribution to scientific progress within the scope of this journal (substantial new concepts, ideas, methods, or data)?	Excellent Good Fair Poor
2) Scientific quality Are the scientific approach and applied methods valid? Are the results discussed in an appropriate and balanced way (consideration of related work, including appropriate references)?	Excellent Good Fair Poor
3) Presentation quality Are the scientific results and conclusions presented in a clear, concise, and well structured way (number and quality of figures/tables, appropriate use of English language)?	Excellent Good Fair Poor
For final publication, the manuscript should be accepted as is accepted subject to technical corrections accepted subject to minor revisions reconsidered after major revisions I am willing to review the revised paper. I am not willing to review the revised paper. rejected	
Suggestions for revision or reasons for rejection (will be published if the paper is accepted for final publication)	
I am grateful to the authors for considering my comments by extending the discussion and making more clear the limitations of their modelling. However, still it is very misleading to impose a concept about "sustained yield" (SY) by estimating wood harvest rates, which are perceived as providing a potential, but which have nothing to do with the actually possible harvest. An information about a harvest when harvesting is hardly possible (e.g. because of inaccessibility) or even impossible provides actually not a potential. This should better not be converted into carbon in harvested wood pools. A harvest potential is usually something	

which may potentially be used in future, but which is currently not used for some reasons. A potential SY is thus possibly not what the authors actually reveal with their simulations, but rather a kind of a change rate in biomass stocks. It would be much better to avoid SY. If harvesting is to be implemented in these models, and this would certainly be a good idea, it makes only sense if the harvesting regime would at least show some realism.

R1: We thank the reviewer for his/her additional comments, which made it clear to us that the terminology and framing of our paper can lead to confusion in particular when considering the interdisciplinary context of our study. We thank the editor for a chance to modify the manuscript accordingly. The following major changes were implemented:

- 1- We outline in the introduction in detail the different concepts dealt with in our study and how they relate to each other and we have clarified our terminology throughout the manuscript. Specifically we now distinguish between a “growth potential” and a “harvest potential”, where the second accounts for the reviewer’s and editor’s concern that much area may not be suited for management (see next bullet point), while the first is now clarified as a purely ecological potential. Accordingly, we rename the applied harvest concept to “growth-based harvest (GB)” to clearly state the relation between forest growth potential under changing climate and CO₂ concentration and the amount of wood harvest as regrown annually. We discuss the similarity to the sustained yield concept (targeting the annual increment), but refrain from mixing this management practice with our approach throughout the rest of the manuscript.*
- 2- We provide new figures masking out inaccessible forest area by overlaying a map of managed forest area to truly account for real potentials of forest harvest management for mitigating CO₂ (MF). This map indicates forest areas subject to conservation, infrastructural limits, or not being influenced by human activities so far due to other reasons by Kraxner et al., 2017. The resulting numbers of harvest and mitigation potentials are lower than our estimates based on growth potential. The impact of environmental changes on harvest potentials, however, turns out to remain important. The key message of our paper, that simulating wood harvest needs to account for these changes, remains valid qualitatively, but becomes stronger as it can now better be linked to considerations of actual future harvest regimes, as suggested by the review.*
- 3- We extended substantially the discussion of the trustworthiness of our model with respect to processes like CO₂-fertilization as suggested by the editor. In particular, we now included the findings of a recent study that used a similar model version like ours but included the nitrogen cycle explicitly (it found a low sensitivity of the land carbon cycle to nitrogen limitation). We also point the reader to new evidence that CO₂-fertilization of our model may be in line with observations. We further added new simulation results of forcing GB and MF by present-day climate to show the effects of transient climate and CO₂ concentrations on forest growth and harvest potential. This makes it easier for the reader to identify the effect of climate changes in all components, from growth potential, to harvest potential, to relevance for mitigation.*

These changes have substantially improved the quality of our submitted (see differences below). We hope this response satisfies the high standards of Biogeosciences.

1 **Title Page**

2

3 **Title:** Simulating growth-based harvest adaptive to future climate change

Gelöscht: sustained yield harvesting

4 **List of authors:** Rasoul Yousefpour^{1,2*}, Julia E.M.S. Nabel¹, Julia Pongratz^{1,3}

5 ¹ Land in the Earth System, Max Planck Institute for Meteorology, Bundesstr. 53, 20143
6 Hamburg, Germany

7 ² Chair of Forestry Economics and Forest Planning, University of Freiburg, 79106 Freiburg,
8 Germany

9 ³ Now at Department of Geography, Ludwig-Maximilians-Universität, 80333 München,
10 Germany

11

12 ***Corresponding author:** Rasoul Yousefpour (Rasoul.yousefpour@ife.uni-freiburg.de, Tel:
13 +49-761-2033688)

14

15

Gelöscht:

Gelöscht:

17 **Abstract**

18

19 Forests are the main source of biomass production from solar energy and take up globally
20 around 2.4 ± 0.4 PgC per year. Future changes in climate may affect forest growth and
21 productivity. Currently, state-of-the-art Earth system models use prescribed wood harvest rates
22 in future climate projections. These rates are defined by integrated assessment models (IAMs)
23 only accounting for regional wood demand and largely ignoring the supply side from forests.

24 Therefore, we assess how global growth and harvest potentials of forests change when they are
25 allowed to respond to changes in environmental conditions. For this, we simulate wood harvest
26 rates oriented towards the actual rate of forest growth. Applying this growth-based harvest rule
27 (GB) in "JSBACH", the land component of the Max-Planck-Institute's Earth System Model,
28 forced by several future climate scenarios, we realized a growth potential twice to four times
29 ($3-9 \text{ PgCy}^{-1}$) the harvest rates prescribed by IAMs ($1-3 \text{ PgCy}^{-1}$). Limiting GB to managed forest
30 area (MF), we simulated a harvest potential of $3-7 \text{ PgCy}^{-1}$, two to three times higher than IAMs.

31 This highlights the need to account for the dependence of forest growth on climate. To account
32 for long term effects of wood harvest as integrated in IAMs, we added a life cycle analysis
33 showing that the higher supply with MF as an adaptive forest harvesting rule may improve the
34 net mitigation effects of forest harvest during the 21st century by sequestering carbon in
35 anthropogenic wood products.

36

37 **Keywords:** Adaptation to climate change, Mitigation, Mortality, Carbon forestry, sustainable
38 forest management, Global forest model

39

Gelöscht: apply the concept

Gelöscht: "sustained yield" (SY)

Gelöscht: represent a climate-adaptive forest management allowing a

Gelöscht: rate

Gelöscht: SY

Gelöscht: wood harvest amount

Gelöscht: SY

Gelöscht: (max. 379 PgC).

Gelöscht:

Gelöscht:

49 1. Introduction

50 Forest ecosystems play a major role in taking up global CO₂ emissions and affect global climate
51 conditions through a range of complex biophysical and biogeochemical processes. Forests are
52 the main source of biomass production from solar energy through photosynthesis and are
53 estimated to take up globally around $2.4 \pm 0.4 \text{ PgCy}^{-1}$ (Pan et al., 2011). A large part of this
54 uptake can be attributed to direct and indirect human interference: Direct human impact by
55 forest management creates young forests sequestering carbon during regrowth (Houghton et al.,
56 2012), and provides material for fossil-fuel substitution (Nabuurs et al., 2007). However, forest
57 utilization and interaction of management with large-scale natural disturbances, such as forest
58 fires, may emit tonnes of CO₂ immediately to the atmosphere and act as a source of CO₂
59 emissions (Bonan, 2008). Indirect human impact alters environmental conditions, in particular
60 climate and atmospheric CO₂ concentrations, which historically has caused a carbon uptake by
61 the terrestrial vegetation (Le Quéré et al., 2018). Any change in environmental conditions
62 affects forest growth, risks of hazards, and productivity and, consequently, the amount of wood
63 that can be harvested (Temperli et al., 2012; Sohngen and Tian, 2016).

64 The effects of changes in environmental conditions on the state of the biosphere are represented
65 in state-of-the-art Earth system models (ESMs). However, the description of forest management
66 in these models is largely independent of environmental changes: So far, ESMs employ
67 prescribed wood harvest amounts. These are derived from national statistics for the historical
68 period and from global integrated assessment models (IAMs) for future scenarios. IAMs
69 determine the wood harvest rates based on the supply of woody materials from vegetation and
70 demands of regional industries and population (van Vuuren et al., 2011). However, changes in
71 the supply via forest growth and changed structural conditions especially under climate change
72 and increasing CO₂ concentrations are ignored. The main drivers of these models are economic,
73 i.e. market price, and population growth scenarios and forest harvest decisions are only reactive

Gelöscht: .

Gelöscht: Quere

Gelöscht: 2015

Gelöscht: .

Gelöscht:

Gelöscht:

78 to the assumed socioeconomic scenarios and do not take forest ecosystem dynamics and growth
79 into account.

80 In this study we investigate the relevance of changes in environmental conditions for the growth
81 potential of forests and subsequently their harvest potentials. Moreover, we explore the
82 ecological potential of world forest resources for wood production and the implications for
83 carbon mitigation. To assess growth and harvest potentials, we investigate forest growth under
84 various future climate scenarios. We allow forests to be harvested and to regrow in response to
85 the respective changes in environmental conditions, in all scenarios such that the growth
86 increment is removed each year, i.e. the biomass stocks are neither reduced nor increased. We
87 call this “growth-based” harvesting (GB). Removing the annual increment mirrors the forest
88 management concept of “sustained yield”. Managing for sustained yield is a strong
89 sustainability policy applied in sustainable forest management, which aims to maintain forest
90 stocks as natural capital and controls wood extraction (Luckert and Williamson, 2005).

91 According to the sustained yield concept, the maximum wood harvest rate to utilize forest
92 resources equals the actual rate of forest growth. Exceeding regrowth rates would result in the
93 exploitation of forest ecosystems and would decrease forest yield and productivity. On the other
94 hand, minimalistic usage, i.e. falling below regrowth rates, would not be an optimal allocation

95 of forest resources from the perspective of production. However, the traditional concept of
96 sustained yield management, as defined above, does not account for changes in the growth rates
97 (Luckert und Williamson, 2005), although forest growth rates are highly dependent on the
98 environmental conditions (Collins et al., 2018). It has been noted before that any decision about
99 forest management should take into account the effects of changes in climate and CO₂
100 concentrations on forest growth (Yousefpour et al., 2012; Hickler et al., 2015; Sohngen and
101 Tian, 2016; Sohngen et al., 2016) and consequently on the harvest rate (Temperli et al., 2012;
102 Jönsson et al., 2015). Here we demonstrate how altered growth potentials translate into higher
103 harvest potentials under an adaptive growth-based harvest. We idealize the concept for this

Gelösch: allowing wood harvest decisions to respond to

Gelösch: For this we design a thought experiment applying the concept of “sustained yield” (SY) to illustrate the potential consequences of representing adaptive forest management in ESMs and compare the results with wood harvest prescribed from IAMs. SY

Gelösch: SY

Gelösch: Forest

Gelösch: , especially climate and CO₂ concentrations, which are projected to be substantially altered for future climate scenarios as compared to the conditions observed in the past

Gelösch: . 2013). Therefore,

Gelösch: Sohngen et al., 2016;

Gelösch: Jönsson et al., 2015;

Gelösch: The traditional

Gelösch: of SY, as defined above, does not account for changes

Gelösch:

Gelösch:

122 study in that GB is applied world-wide irrespective of the accessibility of the forest for forest
123 management activities, but allowing for dependence of wood harvest on altered climate
124 conditions and CO₂ concentrations. To link these results to actual harvest potentials, we overlay
125 information on the accessibility of forest areas represented as managed forest area (Kraxner et
126 al., 2017).

127 Keeping in mind the above-mentioned problem of not accounting for changing environmental
128 conditions in global forest utilization modelling, the goal of this study is to establish a modeling
129 framework that allows harvesting rates to respond interactively to environmental changes. We
130 further assess the maximum potential of global forest resources for wood production and the
131 long term CO₂ mitigation effects of wood harvest, which are implicit or explicit drivers of forest
132 utilization in IAMs. We compare the outcome of the growth-based harvest with the outcome
133 when applying prescribed wood harvest amounts from three different Representative
134 Concentration Pathways (RCPs) realized by IAMs and commonly used by ESMs as an external
135 forcing (Hurt et al., 2011). Since harvested material is used in the IAMs to estimate the amount
136 of bioenergy wood, which in turn is needed in the IAMs to analyze energy and carbon mitigation
137 policies, we perform a first-order assessment of the CO₂ consequences of altering the harvest
138 rates in response to climate. Similarly and to determine the mitigation potential by wood
139 products we allocate the harvested material to products of different lifetimes according to FAO
140 country-specific statistics (FAOSTAT, 2016). The change in atmospheric carbon content
141 resulting from the release of CO₂ by the decay of these products is quantified accounting for
142 compensating fluxes by the ocean and the terrestrial vegetation (Maier-Reimer und
143 Hasselmann, 1987). The net mitigation effect of wood harvest is then defined as the difference
144 between the total amount of harvested material and the change in atmospheric carbon content.

Gelöscht: the regrowth rates (Luckert und Williamson, 2005). Therefore,

Gelöscht: concept is idealized

Gelöscht: this study,

Gelöscht: However, we remain consistent with the aim of applying sustained yield

Gelöscht: safeguard

Gelöscht: current level

Gelöscht: wood stocks in the forest. We emphasize that our idealized SY approach realizes global forest resources potentials for wood production assuming full accessibility of all

Gelöscht: .

Gelöscht: goals of this study are (1) to establish a coupled, interactive harvest-climate module based on the concept of sustained yield to account for the reaction of the forest production system to climate change and increasing CO₂ concentration in the atmosphere, and (2) to

Gelöscht: as applied

Gelöscht: by an active and dynamic management strategy.

Gelöscht: sustained yield modelling approach

Gelöscht: are

Gelöscht:

Gelöscht:

167 2. Materials and methods

168 2.1. Dynamic global vegetation model JSBACH

169 We implemented the **GB** harvesting rule in JSBACH, the land component of the MPI-ESM
170 (Reick et al., 2013). In the applied version of JSBACH vegetation is represented by 12 plant
171 functional types (PFTs) including six woody PFTs. Each PFT is globally endowed with
172 properties in relation to integrated processes in JSBACH and PFT-specific phenology, albedo,
173 morphology, and photosynthetic parameters (Pongratz et al., 2009). Organic carbon is stored in
174 three vegetation pools: living tissue as “green”, woody material as “wood”, sugar and starches
175 as “reserve pool”, and two soil pools with a fast (about 1 year) and a slow (about 100 years)
176 turnover time (Raddatz et al., 2007). Wood harvest activities do not change the area or
177 characteristics of different PFTs, but affect the carbon pools of woody PFTs (forests and shrubs)
178 by removing carbon from the wood pool, resembling trees’ stem and branches removal via
179 harvesting (Reick et al., 2013). Harvest thus affects the vegetation carbon stocks, but the model
180 does not represent a feedback of the harvest activity on the forest productivity.

181 We applied JSBACH in ‘offline’ mode, i.e. not coupled to the atmosphere, but driven by the
182 CMIP5 output of the MPI-ESM (Giorgetta et al., 2013) from experiments with CO₂ forcing
183 according to three different RCP scenarios (RCP2.6, RCP4.5 and RCP8.5) **for the year 2006-**
184 **2100**. We used a T63/1.9° horizontal resolution and conducted our simulations with
185 disturbances due to fire and wind. The simulations were conducted without dynamic vegetation
186 and without land-use transitions to prevent changes in the areas occupied by the different PFTs
187 and to be thus able to isolate the effects of forest management activities. Further details on the
188 model version and the simulation setup are given in the supplementary material (S1).

189 2.2. RCPs wood harvest

190 The current standard module for anthropogenic land cover and land-use change in JSBACH is
191 based on the harmonized land-use protocol (Reick et al., 2013), which provides land-use

Gelöscht: sustained yield

Gelöscht: With this model we simulated the effect of forest management activities, i.e. wood harvest according to different harvesting rules, on the future (2006-2100) carbon balance of terrestrial ecosystems. ¶

Gelöscht: .

Gelöscht:

Gelöscht:

198 scenarios for the period 1500-2100 (Hurt et al., 2011). As part of this protocol, a set of globally
199 gridded harvest maps from the IAM implementations of the RCPs is provided (Hurt et al.,
200 2011). In JSBACH simulations, the harvest prescribed in these maps is fulfilled taking above-
201 ground carbon of all vegetation pools and all PFTs proportionally to the different pool sizes. In
202 this study, however, we concentrate on the carbon harvested from the wood pool of the woody
203 PFTs, which by far contributes most of the harvested volume.

Gelöscht: paper

204 2.3. Growth-based (GB) harvesting rule to estimate growth potentials

Gelöscht: Sustained yield (SY)

205 As an alternative for the prescribed harvest maps, we implemented the GB harvesting rule,
206 which allows for adaptive wood harvesting reacting to changes in wood increments, and
207 accordingly dependent on climate and CO₂ conditions. We define the GB rule as the allowance
208 to harvest specific volumes of wood to the extent of the average increment (i.e. the average
209 annual growth). Applying GB, we aim to stabilize the wood carbon pool in the woody PFTs at
210 the level of a selected reference period. In the current paper we selected the maximum level for
211 the present period (1996-2005) simulated with JSBACH (see S1). Using a reference level
212 determined from the last ten years of the historical simulation allows us to keep the standing
213 wood on the present level and to account for the dependence of forest growing stocks (carbon
214 pools) to disturbances, silvicultural interventions and varying environmental conditions. Under
215 the GB harvesting rule, the wood harvest is only allowed to reduce the wood carbon pool down
216 to the reference level. Aside from environmentally driven decreases, the wood carbon pool thus
217 nearly remains constant over the whole simulation time.

Gelöscht: SY

Gelöscht: SY

Gelöscht: the SY rule

Gelöscht: ¶

Gelöscht: SY

218 219 2.4. Simulation runs with JSBACH

220 We conducted six simulations (Table 1) from 2006-2100, all starting from the same initial state
221 (see S1). The simulations differ in the applied harvest rule and in their climate and CO₂ forcing.
222 While the different RCP harvest maps were applied in simulations with the corresponding MPI-

Gelöscht:

Gelöscht:

230 ESM RCP forcing, each MPI-ESM RCP forcing was additionally run applying the GB
 231 harvesting rule.

Gelöscht: SY harvest

232 **Table 1**

233 2.5. Growth-based harvesting restricted to managed forests (MF)

234 To infer from the growth potentials simulated under GB how much biomass could potentially
 235 be harvested (harvest potentials), we conduct a post-processing step overlaying a map that
 236 masks out forest areas subject to conservation, infrastructural limits, or not being influenced by
 237 human activities so far due to other reasons (Kraxner et al., 2017). Applying nearest neighbor
 238 interpolation on the 1 km² spatially explicit map of primary forest intensity (0%-100%; Fig. 6
 239 in Kraxner et al., 2017) we derived a T63 map of primary forest area. This static map was used
 240 to filter the growth-based harvest determined in the GB simulations for 2006 to 2100, to only
 241 account for managed forests (MF) in the mitigation assessment.

242 2.6. Analysis of wood harvest impacts on forest disturbances and natural mortality

243 To analyze the mechanisms driving differences in GB and RCP wood harvest amounts we can
 244 formulate changes in above-ground wood carbon stocks over time (dCw/dt) as carbon gains
 245 from net primary production allocated to the wood pools (NPP_w) minus losses due to natural
 246 disturbances and anthropogenic management (i.e., wood harvest, h):

Gelöscht: SY

247
$$\frac{dCw}{dt} = NPP_w - \frac{Cw}{\tau} - h \quad (1)$$

248 In this conceptual formulation, the loss due to natural disturbances depends on the size of the
 249 carbon stock and a time constant (τ). As net primary production in our model does not depend
 250 on harvest, GB growth potentials (p_{GB}) and RCP harvest can be related as

Gelöscht: RCP

Gelöscht: SY

251
$$p_{GB} = h_{RCP} + \left(\frac{Cw_{RCP}}{\tau_{RCP}} - \frac{Cw_{GB}}{\tau_{GB}} \right) + \left(\frac{dCw_{RCP}}{dt} - \frac{dCw_{GB}}{dt} \right) \quad (2)$$

Gelöscht:
$$h_{SY} = h_{RCP} + \left(\frac{Cw_{RCP}}{\tau_{RCP}} - \frac{Cw_{SY}}{\tau_{SY}} \right) + \left(\frac{dCw_{RCP}}{dt} - \frac{dCw_{SY}}{dt} \right) \quad (2)$$

Gelöscht:

Gelöscht:

258 The amount of **growth potential under GB** can thus be split into several terms: The first term is
259 the reference harvest rate of the RCPs. The second term accounts for the difference in loss due
260 to natural disturbances in the RCP and the **GB** simulation. In JSBACH this can further be split
261 into differences in losses due to background mortality, such as self-thinning of forests, due to
262 fire, and due to windbreak. JSBACH explicitly integrates two modules for the simulation of
263 fire and wind disturbances depending on climate and carbon pools. The third term accounts for
264 the changes in the above-ground wood pool realized over time in the simulations. As shown
265 below, the RCP harvest results in an increase of above-ground woody biomass over the 21st
266 century for all three scenarios. For **GB**, on the other hand, dC_{WGB}/dt should theoretically be
267 close to zero over time as **GB** aims to sustain the above-ground carbon pools of woody PFTs;
268 however, reductions in NPP due to less favorable climatic conditions or increased disturbances
269 can entail negative dC_{WGB}/dt . To summarize, **GB** includes the RCP wood harvest and,
270 moreover, makes use of additionally accumulated carbon and eventually reduced mortalities to
271 adapt harvest decisions to the novel climate and forest growing conditions.

Gelöscht: sustained yields

Gelöscht: SY

Gelöscht: SY

Gelöscht: $dC_{W_{SY}}$

Gelöscht: SY

Gelöscht: $dC_{W_{SY}}$

Gelöscht: SY

272 2.7.Accounting for the mitigation potential of forest management in the Earth system
273 We account for long term effects of wood harvest, as in IAMs, by approaching a life cycle
274 analysis. Many wood products have lifetimes of decades to centuries. Here, we assess the effect
275 on atmospheric carbon content when harvested carbon is transferred, at least to a part, to longer-
276 lived product pools, instead of entering the atmosphere immediately. We compare this
277 “mitigation effect” achievable by the wood products harvested under the **GB** concept **after the**
278 **map of managed forest area is overlaid (MF)** to those achievable according to the three RCP
279 harvest maps. To this end, we distinguish three anthropogenic wood product pools -- bioenergy,
280 paper, and construction -- with 1, 10, and 100 year life times, respectively, as are typically
281 assumed in global modeling studies (Houghton et al., 1983; McGuire et al., 2001).

Gelöscht: SY

Gelöscht:

Gelöscht:

290 To allocate the wood biomass harvested in our JSBACH simulations to different product pools,
291 we made use of FAO country-specific statistics reporting wood production in fourteen different
292 categories (FAOSTAT, 2016). For our analysis, we assume that the production technology and
293 allocated percentage of each country's total wood production to these fourteen categories
294 remains constant at 2005 levels over the 21st century and used these percentages to allocate
295 wood biomass harvest from JSBACH (remapped to countries - see a calculus example in
296 supplementary material S2). The fourteen categories are then assigned to the three distinguished
297 anthropogenic wood product pools. We assume that the harvested material entering one of these
298 three product pools in a year decays at a rate of 1/lifetime, i.e. that all material used for
299 bioenergy is respired to the atmosphere within the same year it is harvested, while the material
300 entering the paper and construction pool is emitted at a constant rate over the following 10 or
301 100 years, respectively. The emissions at a given year for paper and construction pools are
302 therefore composed of a fraction of that year's harvest, but also of the legacy of material
303 harvested earlier, yielding annual emissions E from all three product pools as follows:

$$304 \quad E(t) = f_b h(t) + \sum_{s=t-9}^t \frac{1}{10} f_p h(s) + \sum_{s=t-99}^t \frac{1}{100} f_c h(s) \quad (1)$$

305 Here, f for bioenergy (b), paper (p), and construction wood (c) are the fractions with which the
306 harvested biomass is assigned to the product pools (see S2). We call E "emissions from product
307 decay" in the following.

308 To account for the fact that the emissions from product decay leave the atmosphere over time
309 to be taken up by the terrestrial biosphere and the ocean, we apply the response function (Eq.
310 2) by Maier-Reimer and Hasselmann (1987). Convolution of this response function with the
311 time series of annual emissions from product decay until year t results in the change in
312 atmospheric carbon content in that year, C(t) (Eq. 3).

313

$$314 \quad G(t) = A_0 + \sum_{p=1}^4 A_p \exp^{-t/\tau_p} \quad (2)$$

Gelöscht:
Gelöscht:

315 $C(t) = \int_0^t G(t-s) \cdot E(s) ds$ (3)

316

317 Emissions are present in the atmosphere as they occur and, therefore, $G(0) = 1$ and $A_0 = 1 -$
318 $\sum_p A_p$. The constants A_p and the time constants τ_p are fitted for $p > 0$ using one of the best fits
319 found by Maier-Reimer und Hasselmann (1987): the sum of four exponential terms with time
320 constants τ_1, τ_2, τ_3 and τ_4 of approximately 1.9, 17.3, 73.6, and 362.9 years, and constants $a_1,$
321 a_2, a_3 and a_4 of 0.098, 0.249, 0.321, and 0.201. Accordingly, Equation 2 is an exponential
322 function that accounts for the uptake of CO_2 by ocean and land over time and Eq. (3) integrates
323 the accumulated amount of total CO_2 concentrations in the atmosphere at each time step
324 regarding past and present emissions. The mitigation effect of wood products is then determined
325 as the difference between the harvested material and the change in atmospheric carbon content.

326 3. Results

327 3.1. Comparison of **GB** and RCP **harvesting**

328 Above-ground woody biomass is simulated to increase by the end of the 21st century for the
329 RCP wood harvest (Figure 1a), despite an increase of the amounts of wood harvest (Figure 1b).
330 This implies that the changes in environmental conditions lead to a larger accumulation of
331 woody biomass than is removed by the increased harvest. **Depending on the RCP, the simulated**
332 **increase in above-ground woody biomass may reach 133% (425 PgC in 2100) of the initial**
333 **level in 2005 (320 PgC) for RCP8.5 and substantially higher levels of 128% and 117% for**
334 **RCP4.5 and RCP2.6, respectively (Figure 1a).** The temporal pattern of this increase, with strong
335 increase only in the first half of the century for RCP2.6 or throughout the century for RCP8.5,
336 reflects the projected evolution of changes in CO_2 and climate (Collins et al., 2018).
337 For the **GB** rule, woody biomass remains more or less constant over time (Figure 1a), as the
338 average annual increment is removed by harvest by definition of the **GB** rule (see Methods).

Gelöscht: sustained-yield

Gelöscht: harvest rates

Gelöscht: 2013).

Gelöscht: SY

Gelöscht: SY

Gelöscht:

Gelöscht:

344 Consequently, the growth potential of global forest resources under GB is simulated to be as
 345 high as 9 PgCy⁻¹ at the end of the century subject to the realization of RCP8.5 climatic
 346 conditions, or about 4 to 6 PgCy⁻¹ for the other two scenarios (Figure 1b). About two thirds of
 347 the growth potential lie in managed forest areas and are thus potentially harvestable (Figure 1b,
 348 MF-harvest curves). The MF harvest potentials are thus twice to three times (3-7 PgCy⁻¹) as
 349 high as those of prescribed wood harvest simulated by IAMs for the RCPs. Note that, as
 350 described in the methods, managed forest areas refer to the present-day state and may expand
 351 in the future, which would further increase the harvest potential. These figures are harvestable
 352 wood biomass amount and differ from commercially useable timber including bioenergy, paper,
 353 and construction woody biomass (see 2.7 and 3.3).

354 We map the geographical distribution of RCP harvest as well as growth and harvest potential
 355 under the GB harvesting rule applied to all global forest (GB) and managed forest areas (MF)
 356 to recognize regional hotspots (Figure 2). Central Latin America including the accessible parts
 357 of the Amazon forests, large parts of North America, the accessible parts of central Africa,
 358 eastern Asia and Europe including Russia can be recognized under all climate scenarios as
 359 hotspots for allocation of simulated harvest activities. The large harvest potentials of the supply-
 360 based harvest in the tropics contrast with the patterns of the demand-based RCP harvest; in
 361 particular in RCP2.6 and RCP4.5 much of the global harvest is provided from eastern North
 362 America, central Europe and East Asia. A reasonable proportion of GB harvest amount in the
 363 tropics is masked out in MF as inaccessible forest area; nevertheless the tropics contribute a
 364 large harvest potential from wood supply side in both GB and MF.

365 **Figure 1**

366 **Figure 2**

Gelöscht: wood harvest

Gelöscht: SY

Gelöscht: SY wood

Gelöscht: amounts

Gelöscht: four

Gelöscht: Depending on RCPs, the simulated

Gelöscht: in above-ground woody biomass may reach 133% (425 PgC in 2100) of the initial level in 2005 (320 PgC) for RCP8.5 and substantially higher levels of 128% and 117% for RCP4.5 and RCP2.6, respectively (Figure 1b). All the

Gelöscht: 6

Gelöscht: allocation

Gelöscht: potentials, applying both SY

Gelöscht: RCPs wood

Gelöscht: .

Gelöscht: the

Gelöscht: Figures

Gelöscht: in

Gelöscht: maps

Gelöscht: SY

Gelöscht: SY

Gelöscht:

Gelöscht:

388 3.2 Separation of the processes underlying the growth potentials under future climate
389 scenarios

390 The harvest potential under the GB harvesting rule in JSBACH exceeds RCPs wood harvest
391 defined by IAMs not only because of taking into account changes in growth rates caused by
392 changed environmental conditions, but also due to avoided mortality and disturbances (see
393 methods section). Figure 3 shows the separation of the growth potential underlying the GB
394 harvest into changes in standing wood as compared to RCP harvest, avoided background
395 mortality, natural fire, and wind disturbances, and the amount prescribed originally by RCPs.

396 The largest contribution to the growth potential under the GB harvesting rules exceeding the
397 RCP harvest is the lower background mortality, which is directly related to lower accumulation
398 of woody biomass (see Figure 1a). This lower accumulation also leads to the decreased carbon
399 losses from fire and wind disturbances. Depending on the climate scenario (RCPs) the simulated
400 reduction of mortality and disturbances add up to 2-5 PgCy⁻¹ at the end of the century. Under
401 the RCP harvest, woody biomass is simulated to mostly increase beyond what is required by
402 the increasing harvest rates (see Figure 1). Harvesting this “surplus”, i.e. the increase of
403 standing biomass over time by applying RCP harvest rates and harvesting less biomass than the
404 annual increment provides, also contributes to the larger growth potentials under the GB
405 harvesting rule. The temporal evolution is different from that of avoided mortality and
406 disturbances, reflecting the projected changes in CO₂ and climate. Greater fluctuation of the
407 growth potential compared to the RCPs’ annual wood harvest amounts is because of the direct
408 dependency of the forest’s productivity on climate fluctuations.

409 **Figure 3**

410 3.3. Mitigation potential of GB versus RCP wood harvest
411 We show the mitigation potential of forest resources in the 21st century under growth-based
412 harvesting of global forest (GB) and managed forest (MF) areas versus the RCP wood harvest

Gelöscht: Wood

Gelöscht: impacts on forest disturbances and regrowth
The realized SY harvest

Formatiert: Schriftfarbe: Akzent 1

Gelöscht: climatic

Gelöscht: realized SY

Gelöscht: higher harvesting

Gelöscht: SY

Gelöscht: and thus to sustained yield and increased
harvest potentials of over 100%, over 200%, and over
250% for RCP2.6, and RCP4.5 and RCP8.5, respectively.

Gelöscht: lower

Gelöscht: growth

Gelöscht: SY harvest

Gelöscht: yearly wood harvest from sustained yield
comparing

Gelöscht: to climate and accordingly fluctuation of forest
growth and

Gelöscht: SY

Gelöscht: applying SY harvest rule

Gelöscht:

Gelöscht:

432 prescribed from IAMs in Figure 4. Due to the larger harvested amounts, the mitigation potential
433 is higher for GB and MF compared to RCP harvest and the magnitude depends on the
434 underlying climate scenario. The advantage of growth-based harvesting lies in storing a larger
435 amount of carbon in wood products whilst keeping above-ground woody carbon pools constant.
436 These aspects are largely ignored by IAMs. Table 2 below shows the net mitigation potentials
437 of world forest resources (GB and MF against RCP harvest) by wood harvest at the middle and
438 end of the 21st century (2050 and 2100). The highest mitigation effect is achieved in the GB8.5
439 scenario with 140.6 PgC and 379.1 PgC up to 2050 and 2100, respectively. These figures
440 account for 278% and 287% more global carbon storage than in the RCP8.5 scenario with
441 prescribed RCP wood harvest with 50.6 and 132.1 PgC mitigation up to 2050 and 2100,
442 respectively. Only considering current managed forests, the mitigation effect realized for MF8.5
443 still reaches a maximum mitigation potential of 109.3 and 295.8 PgC up to 2050 and 2100,
444 respectively.

445 **Table 2**

446 **Figure 4**

447 4. Discussion

448 RCPs define wood harvest in each region according to scenarios realized by IAMs about social
449 and economic developments in the 21st century, but independent of ecological capacities of
450 forest ecosystems. Although the growth-based harvesting rule realizes potentially a larger wood
451 harvest amount than the RCPs, it remains as per definition a sustained-yield forest harvesting
452 approach and guarantees sustainability of the current ecological conditions at each region with
453 respect to standing biomass. However, as a consequence, regions with low standing biomass,
454 for example due to extensive historical harvest, will maintain these low biomass levels. Below
455 we discuss the effectivity of GB in adapting to new environmental conditions and the mitigation
456 potential and highlight the missing issues in our simulation analysis, especially about the

Gelöscht: 4a-f

Gelöscht: SY

Gelöscht: (RCPs).

Gelöscht: SY

Gelöscht: SY

Gelöscht: SY8

Gelöscht:

Gelöscht: 287

Gelöscht: 286

Gelöscht: SY

Gelöscht: SY

Gelöscht:

Gelöscht:

468 provisioning of multiple goods and services (e.g. biodiversity, forest health), and the future
469 research themes about integration of diversified management strategies in ESMs.

470 Accounting for the climate state in simulating future forest harvest is crucial (Temperli et al.,

471 2012; Sohngen and Tian, 2016). Accordingly, the novelty of the applied **GB** in this study is the

472 dynamic nature of this management approach based on the ecology of forest ecosystems and

473 climatic and atmospheric conditions. According to Schelhaas et al. (2010), an accelerated level

474 of wood harvest to reduce the vulnerability of European old forests to wind and fire disturbances

475 is needed to stop the current built-up of growing stock. Applying **GB** in this study realized an

476 increased wood harvest rate for European forests (see Figure 2) showing first signs of carbon

477 sink saturation and high vulnerability to natural disturbances (Nabuurs and Maseret, 2013).

478 Global studies of this nature are largely missing due to the lack of data and forest ecosystems

479 complexity on global scale. Our **idealized** simulations suggest that **GB** does not only effectively

480 safeguard sustainability of the current forest biomass on the global scale, but also positively

481 affects the resistance of forest resources against natural disturbances and efficiently utilizes

482 forest growth and productivity potentials, (see Figure 3). Our estimates are, of course, sensitive

483 to the choice of reference level: In this study, we applied the maximum current (1996-2005)

484 above-ground wood biomass as the reference level. Any changes in this reference may affect

485 the realized harvest **potentials** and should be carefully defined regarding ecological potentials

486 and economic implications.

487 In our simulations, future environmental changes are mostly beneficial for accumulation of

488 forest biomass, apparent from increasing standing biomass in the RCP harvest scenarios or the

489 increase of **GB and MF** harvest rates over the 21st century. This is in line with other studies

490 projecting above-ground forest carbon storage to increase in the future (e.g. Tian et al., 2016).

491 These effects of environmental changes on forest growth are largely missing in the IAMs

492 providing the wood harvest scenarios to **dynamic global vegetation models (DGVMs)** and

Gelöscht: SY

Gelöscht: SY

Gelöscht: idealised

Gelöscht: SY

Gelöscht: .

Gelöscht: amounts

Gelöscht: SY

Gelöscht:

Gelöscht:

500 ESMs. The RCP wood harvest rates are based on the demand for wood and bioenergy as the
501 main driver of decisions by IAMs on forest harvest. For example, RCP8.5 applies the forest
502 sector model DIMA (Riahi et al., 2011), which is a spatial model for simulating forestry
503 processes to meet specific regional demand on wood and bioenergy. RCP4.5 bases wood
504 harvest rates solely on the price of carbon affected by emissions and mitigation potentials of
505 forestry and agricultural activities (Hurt et al., 2011). Finally, RCP2.6 relies on the forecasted
506 demand on timber and fuelwood from forest resources and applies a series of forest
507 management rules (plantation, clear cutting, selective logging) to meet this demand as the only
508 driver of wood harvest rate in the IMAGE model (Stehfest et al., 2014). IAMs do not account
509 for the fact that the demand side may also be influenced by the availability of the resource and,
510 accordingly, the increased biomass stocks projected for the future would likely lead to larger
511 wood harvest rates than IAMs simulate by assuming present-day growth conditions. The extent
512 to which accounting for environmental changes may influence estimates of harvestable material
513 (e.g. apparent from comparisons of GB and MF harvest potentials under RCP2.6 as compared
514 to RCP8.5, see Figure 1) highlights the need to include these effects in models, such as IAMs,
515 that estimate future wood harvest. Our study is limited to considering biomass growth, albeit in
516 interaction with soil conditions also responding to the altered climate. In reality, harvest
517 decisions would consider further variables that depend on environmental conditions, such as
518 the maximum soil expectation value, which are not explicitly simulated neither in our model
519 nor in IAMs.

520 Note that the estimates of GB wood harvest as provided by our model are not meant as plausible
521 estimates of actual future harvest, which as described before depends not just on resource
522 availability and accessibility of areas, but is demand driven by other economic and political
523 considerations. Limiting GB to available managed forest area, MF realized less harvest
524 potential than GB, however, still a larger amount than RCP and with a higher mitigation
525 potential (see Table 2). Also, actual future harvest will interact with other land-use decisions

Gelöscht: SY

Gelöscht: rates

Gelöscht: SY

Gelöscht: , including considerations e.g. of conservation and accessibility of forests.

Gelöscht:

Gelöscht:

531 such as changes in forest cover due to agricultural expansion, but also afforestation. We have
532 further not accounted for the effects of wood harvest on biodiversity, forest health, and other
533 ecosystem services. Chaudhary et al. (2015) state that the effect of forest management on the
534 species richness, for example, highly depends on the management regime applied. They refer
535 to literature reporting a positive effect of logging activities on species richness as a result of
536 establishing early successional colonizers. Additionally, applying selective logging approaches
537 (e.g. future crop trees of targeted species) for forest management may enhance forest recovery
538 and reduce unintended changes in species composition (Luciana de Avila et al., 2017). Instead
539 of actual forest harvest that considers all these aspects in its decision-making, our study
540 provides an estimate of the ecological potentials for wood harvest. However, the change in
541 resource potentials with climate change forms the ecological basis for realistic decision-making.

542 There is uncertainty in simulating ecosystem response to environmental changes. Regional
543 forest inventories show an increase in biomass due to historical environmental changes
544 (excluding effects of land-use change) (Pan et al., 2011). The largest sinks are found by these
545 studies to be in the tropical regions, coinciding with our simulated regions of largest potentials
546 for additional wood harvest. Also the other regions showing larger potential for wood harvest
547 under **GB and MF** than RCP, such as North America, Europe, Russia and East Asia, currently
548 exhibit carbon uptake due to historical environmental changes. This gives some confidence in
549 the robustness of our results, in particular since most models project the carbon sink in
550 vegetation to continue for the future; however, its magnitude is uncertain (Sitch et al., 2008). A
551 large source of uncertainty is the strength of the CO₂-fertilization effect (Kauwe et al., 2013;
552 Hickler et al., 2015), which reflects in a large spread across models in estimates of global total
553 (vegetation plus soil) terrestrial carbon stocks (Arora et al., 2013) and of vegetation productivity
554 (Zaehle et al., 2014). To better assess these effects, we additionally simulated the future **GB**
555 **and MF** harvest potentials under present-day climate and CO₂ conditions (see simulations of
556 **GBpd and MFpd** in the supplementary material (S1)). These simulations led to a wood harvest

Gelöscht: SY

Gelöscht: for SY harvest

Gelöscht: ; Kauwe et al., 2013)

Gelöscht: SY

Gelöscht: amount

Gelöscht: This simulation

Gelöscht:

Gelöscht:

563 potential larger than that with RCPs harvest rates and rather constant harvest over time (~3.2
 564 and 2.7 PgC annually for GBpd and MFpd, respectively, see S1-Figure 1). The harvest amount
 565 of GBpd is equal to RCP8.5 harvest amount at the end of the century in our simulations.
 566 Differences between GBpd and MFpd and the simulations forced by the different RCPs as well
 567 as differences among the latter illustrate the effects of changes in climate and CO₂ concentration
 568 on forest growth and resulting harvest potentials. The differences in wood harvest amounts
 569 between the harvest simulations based on GB and MF and those with prescribed RCP wood
 570 harvest rates in the first simulation year show differences of applying the supply-based harvest
 571 rule (GB and MF) versus the demand-based RCPs under current environmental conditions. The
 572 geographic allocation of growth and harvest potentials for GBpd and MFpd (see S1-Figure 2)
 573 resembles those under RCPs, however, with higher global values. That the GBpd and MFpd
 574 harvest potential are higher than the RCP harvest implies that the larger potentials, as compared
 575 to RCP harvest are partly attributable to the harvest simulated by IAMs not using the full
 576 sustained, ecological potential (e.g. due to real-world demand). However, the harvest potentials
 577 under RCP climate all grow substantially larger than the harvest potential under present-day
 578 climate. This depicts the isolated effect of environmental changes, particularly CO₂ fertilization,
 579 on the simulated potential harvest.

581 A further uncertainty in the model we used is that our model did not explicitly account for a
 582 nitrogen cycle. Nitrogen may become a limiting factor for the additional uptake of carbon in
 583 vegetation, although future climate change might also lead to higher nutrient availability due to
 584 faster decomposition rates (Friedlingstein and Prentice, 2010). Further, nitrogen deposition may
 585 reduce nitrogen limitation (Churkina et al., 2010), and it is not predictable if artificial
 586 fertilization of managed forests may find wide-spread application in the future. Overall,
 587 therefore, quantifications of effects of future climate change on global carbon stocks derived
 588 from individual models have to be treated with care. Our model includes present-day nitrogen

Gelöscht: biomass

Gelöscht: S1-Figure1

Gelöscht: SY under present-day climate

Gelöscht: SY

Gelöscht: SY

Gelöscht: RCPs

Gelöscht: SY

Gelöscht: geographical

Gelöscht: SY under present-day climate

Gelöscht: SY

Gelöscht: lower

Gelöscht: harvest amount.

Gelöscht: SY

Gelöscht: under present-day climate is

Gelöscht: we simulated under SY

Gelöscht: constraints such as conservation and accessibility issues discussed above).

Gelöscht: SY

Gelöscht: rates

Gelöscht: SY

Gelöscht: harvesting

Gelöscht: , which

Gelöscht: However, the location of the largest potentials of SY

Gelöscht:

Gelöscht:

613 limitation implicitly by choice of photosynthetic parameters and includes structural limits
614 prohibiting development of wood densities beyond observational values. Tests with a similar
615 model version as ours but representing an explicit nitrogen cycle suggest a rather small
616 sensitivity of the land carbon cycle to nitrogen limitation under CO₂ increases and climate
617 changes in the range of the RCP scenarios investigated here (Goll et al., 2017). The increase in
618 gross primary production (GPP) over the industrial era of our model (or similar versions) lie at
619 the high end, but within the range of a wide range of other models (Anav et al., 2013); recent
620 evidence from long-term atmospheric carbonyl sulfide (COS) records shows that models with
621 high GPP growth are most consistent with observations (Campbell et al., 2017). The location
622 of the largest potentials of GB and partly MF harvest simulated in our study being in the tropical
623 forests is consistent with the large carbon sinks derived from inventories for past environmental
624 change (Pan et al., 2011).

625 GB harvest was simulated to mitigate 255-380 PgC, depending on the realized RCP, through
626 wood product usage for the period 2006-2100, from global forest resources. Moreover, it
627 accounted for sustaining the above-ground wood carbon pool at the reference level of 1996-
628 2005. A comprehensive mitigation study, however, should take into account the total carbon
629 balance of forest ecosystems including soil plus litter carbon. Growth enhanced by
630 environmental changes, as simulated to lead to accumulation of woody biomass in the RCP
631 harvest simulations (Fig. 1a), may lead to larger input to the soil (if not removed by wood
632 harvest). However, soil carbon pools respond differently to environmental changes than forest
633 biomass. In particular, soil carbon models generally assume enhanced soil respiration under
634 higher temperatures (Friedlingstein et al., 2006), which may substantially offset the additional
635 carbon uptake by the vegetation (Ciais et al., 2013). As these processes act the same in our
636 simulations of GB, MF and RCP harvesting rules (as they share the same climate scenarios),
637 effects of environmental changes on soil carbon will likely not substantially affect our
638 comparison of GB, MF and RCP harvest in relative terms, but may alter the net carbon balance

Gelöscht: SY

Gelöscht: 2005

Gelöscht: .

Gelöscht: SY

Gelöscht: SY

Gelöscht: harvest

Gelöscht: SY

Gelöscht:

Gelöscht:

646 in each of them. Further, the usage of wood products implies removal of carbon off-field. This
647 can lead to depletion of soil plus litter carbon stocks. Observational data generally found small
648 decreases of soil carbon, but substantial reduction of deadwood material (Erb et al., 2017). Such
649 effects must be expected to be stronger for **GB and MF** harvest with its larger harvested **biomass**
650 than for RCP harvest, reducing on-site carbon stocks, but consequently also soil respiration.
651 Estimating a mitigation potential based on the net carbon balance of vegetation, soil plus litter
652 and product pools therefore would depend on the actual size of soil and vegetation carbon pools
653 and the lifetimes of products relative to the lifetimes of the on-site carbon, which are further
654 subject to a changing climate. There is not a unique life time for anthropogenic wood products
655 pools in the literature. Lifetime of construction wood, for example, spanning from 67 years in
656 Härtl et al. (2017), up to 160-200 years in van Kooten et al. (2007) are applied in recent studies.
657 Regarding global variation of carbon turnover rate, Carvalhais et al. (2013) find mean turnover
658 times of 15 and 255 years for carbon residing in vegetation and soil near to Equator and higher
659 Latitude over 75°, respectively. Regarding the uncertainty about life time of anthropogenic
660 wood pools, we stay consistent with the applied figures in FAO statistics (FAOSTAT, 2016)
661 and other land carbon budget studies (Houghton et al., 1983; McGuire et al., 2001).

662 Despite carbon fluxes being the focus of **land-use** change as mitigation tool (e.g., UNFCCC,
663 2012), forest management may enhance or mitigate climate change by a range of other
664 mechanisms such as a change in surface albedo (e.g., Rautiainen et al., 2011; Otto et al., 2014)
665 or turbulent heat fluxes (e.g., Miller et al., 2011). Such biogeophysical effects needed to be
666 accounted for in a complete assessment of the mitigation potentials, as has been done for global
667 land cover change (Pongratz et al., 2011) or for forest management on local (Bright et al., 2011)
668 or regional scale (Naudts et al., 2016). These biogeophysical effects are particularly important
669 for the local climate (Winckler et al., 2017). In our study, we restrict estimates of mitigation
670 potential to carbon fluxes only and thus focus on the perspective of mitigating global

Gelöscht: SY

Gelöscht: amounts

Gelöscht:

Gelöscht:

Gelöscht:

674 greenhouse gas concentrations. This further allows for a direct comparison of the wood harvest
675 scenarios provided as part of the RCPs.

676 Different from economic models, ESMs do not consider costs associated with early mitigation
677 measures and thereby implicitly assume a zero social discount rate, meaning that there is no
678 preference for immediate mitigation. However, the discount rate plays a major role to find
679 economically the most efficient mitigation action (Stern, 2007). van Kooten et al. (1999)
680 analyzed the sensitivity of investments for carbon sequestration to discount rate in western
681 Canada and found that applying zero discount may not provide enough incentive for increasing
682 carbon storage. However, most forest carbon cost studies are inconsistent in using terms,
683 geographic scope, assumptions, program definitions, and methods (Richards and Stokes, 2004)
684 and may not truly assess carbon sequestration potentials of forest ecosystems. Therefore, if
685 there were a social preference for prompt climate change mitigation, carbon sinks later in the
686 century should be discounted. Regarding the discussion on discount rate, Johnston and van
687 Kooten (2015) argue that applying sufficiently high discount rates in substituting biomass for
688 fossil fuels never leads to carbon neutrality.

689 5. Conclusions

690 We recommend that future research on integration of management strategies in DGVMs and
691 ESMs should regard ecological sustainability as well as socio-economic challenges. In reality
692 and today, forest management is more of a gamble than a scientific debate (Bellassen and
693 Luyssaert, 2014) and there is no consensus in applying a certain forest harvest rule (e.g. **GB**)
694 among forest owners, decision-makers and local users. The rationale to manage forest resources
695 sustainably and efficiently is generally recognized and implemented (Luckert and Williamson,
696 2005; Elbakidze et al., 2013). However, the process of forest management decision-making is
697 based on the past experiences with a business-as-usual strategy (BAU). Adaptation to future
698 environmental change and minimizing the risks associated with climate change impacts is

Gelöscht: SY

Gelöscht:

Gelöscht:

700 recently fully integrated in forest research (Lindner et al., 2014), however, remains in
701 experimental level in implementation (Yousefpour and Hanewinkel, 2015). Mitigation, in turn,
702 is of public interest and there are some attempts internationally to account for mitigation effects
703 of forest management in carbon policy. International programs such as the Kyoto protocol
704 encourage forest managers to store carbon in the forest stocks on the ground applying financial
705 instruments such as tax reduction and direct purchase of carbon offsets. Therefore, inclusion of
706 financial aspects in global forest management modelling and decision-making may help to put
707 scientific results into practice (Hanewinkel et al., 2013). This suggestion is in line with van
708 Vuuren et al. (2011) about the necessity of strengthening the cooperation between integrated
709 assessment models (IAMs) and Earth system modelling communities to improve the
710 understanding of interactions and joint development of environmental and human systems. Our
711 study is the first implementation to account for the climate-dependence of forest growth on
712 global scale for harvest potentials. It suggests the importance of considering this dependence:
713 the **growth-based harvest** approach (GB) as applied in this study may realize wood harvest
714 potentials twice to four times as high as those of prescribed wood harvest simulated by IAMs
715 for the RCPs and **would closely** triple the net mitigation effects of wood products. **By limiting**
716 GB to managed forests (MF), we simulated a lower harvest potential than GB, still two to three
717 times more than in the IAMs, which could double the net mitigation effect of wood harvest
718 potential in the 21st century. To move from estimates of potentials to actual harvest rates,
719 climate-dependent forest growth needs to be integrated with socio-economic factors to fully
720 incorporate economic aspects of forestry practices within a dynamic forest growth and yield
721 modelling system.

Gelöscht: of SY

Gelöscht: sustained-yield

Gelöscht: rates

Gelöscht: may

Gelöscht:

Gelöscht:

726 Code availability

727 Scripts used in the analysis and other supplementary information that may be useful in
728 reproducing the authors' work are archived by the Max Planck Institute for Meteorology and
729 can be obtained by contacting publications@mpimet.mpg.de.

730 Data availability

731 Primary data are archived by the Max Planck Institute for Meteorology and can be obtained by
732 contacting publications@mpimet.mpg.de.

733 Sample availability

734 None

735 Appendices

736 None

737 Supplement link (will be included by Copernicus)

738 Supplementary includes two main files: a word document *S1* on the “details on JSBACH, ~~the~~
739 model version, ~~the simulation setup, and the additional simulation with present day forcing~~”
740 and a zipped excel file *S2* as an example of how “mitigation potentials of woody products in
741 their life time” ~~are calculated.~~

Gelöscht: its

Gelöscht: and

Gelöscht: ”.

742 Team list

743 See authors list

744 Author contribution

745 R.Y. and J.P. initiated the study. R.Y. performed the model simulations. J.N. implemented the
746 code changes. All authors contributed to analyzing the simulations and writing the manuscript.

747 Competing interests

748 The authors declare that they have no conflict of interest.

Gelöscht:

Gelöscht:

752 Acknowledgments

753 This work was supported by the German Research Foundation's Emmy Noether Program (PO
754 1751/1-1). Computational resources were made available by the German Climate Computing
755 Center (DKRZ) through support from the German Federal Ministry of Education and Research
756 (BMBF).

757 References

758 [Anav, A., Friedlingstein, P., Kidston, M., Bopp, L., Ciais, P., Cox, P., Jones, C., Jung, M.,
759 Myneni, R., Zhu, Z.: Evaluating the Land and Ocean Components of the Global Carbon Cycle
760 in the CMIP5 Earth System Models. *Journal of Climate* 26, 6801–6843, 2013](#)

761 [Arora, V. K., Boer, G. J., Friedlingstein, P., Eby, M., Jones, C. D., Christian, J. R., Bonan, G.,
762 Bopp, L., Brovkin, V., Cadule, P., Hajima, T., Ilyinam, T., Lindsay, K., Tjiputra, J. F., and Wu,
763 T.: Carbon-concentration and carbon-climate feedbacks in CMIP5 Earth system models,
764 *Journal of climate*, 26, 5289–5314, 2013.](#)

765 [Bellassen, V. and Luysaert, S.: Carbon sequestration: Managing forests in uncertain times,
766 *Nature News*, 2014.](#)

767 [Bonan, G. B.: Forests and climate change: forcings, feedbacks, and the climate benefits of
768 forests, *Science \(New York, N.Y.\)*, 320, 1444–1449, doi:10.1126/science.1155121, 2008.](#)

769 [Bright, R. M., Stromman, A. H., and Peters, G. P.: Radiative Forcing Impacts of Boreal Forest
770 Biofuels: A Scenario Study for Norway in Light of Albedo, *ENVIRONMENTAL SCIENCE
771 & TECHNOLOGY*, 45, 7570–7580, doi:10.1021/es201746b, 2011.](#)

772 [Campbell, J. E., Berry, J. A., Seibt, , Smith, Montzka, S. A., Launois, T., Belviso, S., Bopp, L.,
773 Laine, M.: Large historical growth in global terrestrial gross primary production, *Nature* volume
774 544, 84-87, 2017](#)

775 [Churkina, G., Zaehle, S., Hughes, J., Viovy, N., Chen, Y., Jung, M., Heumann, B. W.,
776 Ramankutty, N., Heimann, M., and Jones, C.: Interactions between nitrogen deposition, land](#)

Formatiert: Englisch (USA)

Gelöscht:

Gelöscht:

777 cover conversion, and climate change determine the contemporary carbon balance of Europe,
778 Biogeosciences, 7, 2749–2764, doi:10.5194/bg-7-2749-2010, 2010.

779 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, A.,
780 Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao S., and Thornton, P.:
781 Carbon and Other Biogeochemical Cycles, in: Climate Change 2013: The Physical Science
782 Basis. Contribution of Working Group I to the Fifth Assessment Report of the
783 Intergovernmental Panel on Climate Change, 465–570, 2013.

784 Elbakidze, M., Andersson, K., Angelstam, P., Armstrong, G. W., Axelsson, R., Doyon, F.,
785 Hermansson, M., Jacobsson, J., and Pautov, Y.: Sustained yield forestry in Sweden and Russia:
786 how does it correspond to sustainable forest management policy?, *Ambio*, 42, 160–173,
787 doi:10.1007/s13280-012-0370-6, 2013.

788 Erb, K.-H., Luysaert, S., Meyfroidt, P., Pongratz, J., Don, A., Kloster, S., Kuemmerle, T.,
789 Fetzel, T., Fuchs, R., Herold, M., Haberl, H., Jones, C. D., Marin-Spiotta, E., McCallum, I.,
790 Robertson, E., Seufert, V., Fritz, S., Valade, A., Wiltshire, A., and Dolman, A. J.: Land
791 management: data availability and process understanding for global change studies, *GLOBAL*
792 *CHANGE BIOLOGY*, 23, 512–533, doi:10.1111/gcb.13443, 2017.

793 FAOSTAT: FAOSTAT–Forestry: <http://faostat.fao.org/site/626/default.aspx#ancor>, 2016.

794 Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Bloh, W. von, Brovkin, V., Cadule, P., Doney,
795 S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W.,
796 Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.-
797 G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate-carbon
798 cycle feedback analysis: Results from the (CMIP)-M-4 model intercomparison, *Journal of*
799 *climate*, 19, 3337–3353, 2006.

800 Friedlingstein, P. and Prentice, I. C.: Carbon-climate feedbacks: a review of model and
801 observation based estimates, *CURRENT OPINION IN ENVIRONMENTAL*
802 *SUSTAINABILITY*, 2, 251–257, doi:10.1016/j.cosust.2010.06.002, 2010.

803 Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V.,
804 Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Ilyina, T.,
805 Kinne, S., Kornblueh, L., Matei, D., Mauritsen, T., Mikolajewicz, U., Mueller, W., Notz, D.,
806 Pithan, F., Raddatz, T., Rast, S., Redler, R., Roeckner, E., Schmidt, H., Schnur, R.,
807 Segschneider, J., Six, K. D., Stockhause, M., Timmreck, C., Wegner, J., Widmann, H., Wieners,
808 K.-H., Claussen, M., Marotzke, J., and Stevens, B.: Climate and carbon cycle changes from
809 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase
810 5, *J. Adv. Model. Earth Syst.*, 5, 572–597, doi:10.1002/jame.20038, 2013.

811 Goll, S. D., Winkler, A. J., Raddatz, Th., Dong, N., Prentice I. C., Ciais, Ph., Brovkin, V.:
812 Carbon–nitrogen interactions in idealized simulations with JSBACH (version 3.10), *Geosci.*
813 *Model Dev.*, 10, 2009-2030, 2017

814 Hanewinkel, M., Cullmann, D. A., Schelhaas, M.-J., Nabuurs, G.-J., and Zimmermann, N. E.:
815 Climate change may cause severe loss in the economic value of European forest land, *NATURE*
816 *CLIMATE CHANGE*, 3, 203–207, doi:10.1038/NCLIMATE1687, 2013.

817 Hickler, T., Rammig, A., and Werner, C.: Modelling CO2 Impacts on Forest Productivity,
818 *Current Forestry Reports* 1: 69-80: [http://link.springer.com/article/10.1007/s40725-015-0014-](http://link.springer.com/article/10.1007/s40725-015-0014-8)
819 [8](http://link.springer.com/article/10.1007/s40725-015-0014-8), 2015.

820 Houghton, R. A., House, J. I., Pongratz, J., Werf, G. R. van der, DeFries, R. S., Hansen, M. C.,
821 Le Quéré, C., and Ramankutty, N.: Carbon emissions from land use and land-cover change,
822 *Biogeosciences*, 9, 5125–5142, 2012.

823 Hurtt, G. C., Chini, L. P., Frohking, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P.,
824 Hibbard, K., Houghton, R. A., Janetos, A., Jones, C. D., Kindermann, G., Kinoshita, T., Klein
825 Goldewijk, K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P.,
826 van Vuuren, D. P., and Wang, Y. P.: Harmonization of land-use scenarios for the period 1500–
827 2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting
828 secondary lands, *Climatic Change*, 109, 117–161, doi:10.1007/s10584-011-0153-2, 2011.

829 Jönsson, A. M., Lagergren, F., and Smith, B.: Forest management facing climate change - an
830 ecosystem model analysis of adaptation strategies, *Mitig Adapt Strateg Glob Change*, 20, 201–
831 220, doi:10.1007/s11027-013-9487-6, 2015.

832 Kauwe, M. G. d., Medlyn, B. E., Zaehle, S., Walker, A. P., Dietze, M. C., Hickler, T., Jain, A.
833 K., Luo, Y. Q., Parton, W. J., Prentice, I. C., Smith, B., Thornton, P. E., Wang ShuSen, Wang
834 YingPing, Warlind, D., Weng, E. S., Crous, K. Y., Ellsworth, D. S., Hanson, P. J., Kim
835 HyunSeok, Warren, J. M., Oren, R., and Norby, R. J.: Forest water use and water use efficiency
836 at elevated CO₂: a model-data intercomparison at two contrasting temperate forest
837 FACE sites, *GLOBAL CHANGE BIOLOGY*, 19, 1759–1779, 2013.

838 Kraxner, F., Schepaschenko, D., Fussler, S., Lunnanc, A., Kindermann, G., Aokie, K.,
839 Dürauer, M., Shvidenko, A., See, L.: Mapping certified forests for sustainable management
840 - A global tool for information improvement through participatory and collaborative mapping.
841 *Forest Policy and Economics* 83:10–18, 2017

842 Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Pongratz, J., Manning, A. C.,
843 Korsbakken, J. I., Peters, G. P., Canadell, J. G., Jackson, R. B., Boden, T. A., Tans, P. P.,
844 Andrews, O. D., Arora, V. K., Bakker, D. C. E., Barbero, L., Becker, M., Betts, R. A., Bopp,
845 L., Chevallier, F., Chini, L. P., Ciais, P., Cosca, C. E., Cross, J., Currie, K., Gasser, T., Harris,
846 I., Hauck, J., Haverd, V., Houghton, R. A., Hunt, C. W., Hurtt, G., Ilyina, T., Jain, A. K., Kato,
847 E., Kautz, M., Keeling, R. F., Goldewijk, K. K., Kortzinger, A., Landschutzer, P., Lefevre, N.,
848 Lenton, A., Lienert, S., Lima, I., Lombardozzi, D., Metz, N., Millero, F., Monteiro, P. M. S.,
849 Munro, D. R., Nabel, J., Nakaoka, S., Nojiri, Y., Padin, X. A., Peregon, A., Pfeil, B., Pierrot,
850 D., Poulter, B., Rehder, G., Reimer, J., Rodenbeck, C., Schwinger, J., Seferian, R., Skjelvan, I.,
851 Stocker, B. D., Tian, H. Q., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der
852 Werf, G. R., van Heuven, S., Viovy, N., Vuichard, N., Walker, A. P., Watson, A. J., Wiltshire,
853 A. J., Zaehle, S., Zhu, D.: Global Carbon Budget 2017, *Earth System Science Data*. 10:405–
854 448, 2018.

- Gelöscht:** Quere
- Gelöscht:** Moriarty, R.,
- Gelöscht:** Peters, G. P., Ciais, P.,
- Gelöscht:** Jones, S. D.,
- Gelöscht:** Tans, P., Arneeth, A.,
- Gelöscht:** Bozec, Y., Canadell, J. G., Chini, L. P.,
- Gelöscht:** E.,
- Gelöscht:** Hoppema, M
- Gelöscht:** House, J. I
- Gelöscht:** Johannessen, T.,
- Gelöscht:** F., Kitidis, V
- Gelöscht:** Koven, C., Landa, C. S., Landschuetzer
- Gelöscht:** .
- Gelöscht:** Marland, G., Mathis, J. T.,
- Gelöscht:** Olsen,
- Gelöscht:** Ono, T., Peng, S., Peters, W
- Gelöscht:** Raupach, M. R., Regnier, P., Roedenbeck, C., Saito, S., Salisbury, J. E., Schuster, U
- Gelöscht:** Segsneider, J., Steinhoff, T
- Gelöscht:** Sutton, A. J., Takahashi, T
- Gelöscht:** Wang, Y.-
- Gelöscht:** Wanninkhof, R
- Gelöscht:** ., and Zeng, N
- Gelöscht:** carbon budget 2014, EARTH SYSTEM SCIENCE DATA, 7, 47–85, doi:10.5194/essd-7-47-2015, 2015
- Gelöscht:**
- Gelöscht:**

881 Lindner, M., Fitzgerald, J. B., Zimmermann, N. E., Reyer, C., Delzon, S., van der Maaten, E.,
882 Schelhaas, M.-J., Lasch, P., Eggers, J., van der Maaten-Theunissen, M., Suckow, F., Psomas,
883 A., Poulter, B., and Hanewinkel, M.: Climate change and European forests: What do we know,
884 what are the uncertainties, and what are the implications for forest management?, *JOURNAL*
885 *OF ENVIRONMENTAL MANAGEMENT*, 146, 69–83, doi:10.1016/j.jenvman.2014.07.030,
886 2014.

887 Luckert, M. K. and Williamson, T.: Should sustained yield be part of sustainable forest
888 management?, *Can. J. For. Res.*, 35, 356–364, doi:10.1139/x04-172, 2005.

889 Maier-Reimer, E. and Hasselmann, K.: Transport and storage of CO₂ in the ocean - an inorganic
890 ocean-circulation carbon cycle model, *Climate Dynamics*, 1987.

891 Miller, S. D., Goulden, M. L., Hutyra, L. R., Keller, M., Saleska, S. R., Wofsy, S. C., Silva
892 Figueira, A. M., da Rocha, H. R., and Camargo, P. B. de: Reduced impact logging minimally
893 alters tropical rainforest carbon and energy exchange, *PROCEEDINGS OF THE NATIONAL*
894 *ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA*, 108, 19431–19435,
895 doi:10.1073/pnas.1105068108, 2011.

896 Nabuurs, G.-J., Lindner, M., Verkerk, P. J., Gunia, K., Deda, P., Michalak, R., and Grassi, G.:
897 First signs of carbon sink saturation in European forest biomass, *Nature Climate change*, 3,
898 792–796, doi:10.1038/nclimate1853, 2013.

899 Nabuurs, G.-J., Maseraet, O.: Forestry. In *Climate Change 2007: Mitigation. Contribution of*
900 *Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate*
901 *Change* [B. Metz, O.R. Davidson, P.R. Bosch, R. Dave, L.A. Meyer (eds)], 2013.

902 Naudts, K., Chen YiYing, McGrath, M. J., Ryder, J., Valade, A., Otto, J., and Luysaert, S.:
903 Europe's forest management did not mitigate climate warming, *Science (Washington)*, 351,
904 597–600, 2016.

905 Otto, J., Berveiller, D., Breon, F.-M., Delpierre, N., Geppert, G., Granier, A., Jans, W., Knohl,
906 A., Kuusk, A., Longdoz, B., Moors, E., Mund, M., Pinty, B., Schelhaas, M.-J., and Luysaert,

907 S.: Forest summer albedo is sensitive to species and thinning: how should we account for this
908 in Earth system models?, *Biogeosciences*, 11, 2411–2427, doi:10.5194/bg-11-2411-2014,
909 2014.

910 Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L.,
911 Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W., McGuire,
912 A. D., Piao, S., Rautiainen, A., Sitch, S., and Hayes, D.: A Large and Persistent Carbon Sink in
913 the World's Forests, *SCIENCE*, 333, 988–993, doi:10.1126/science.1201609, 2011.

914 Pongratz, J., Reick, C. H., Raddatz, T., Caldeira, K., and Claussen, M.: Past land use decisions
915 have increased mitigation potential of reforestation, *GEOPHYSICAL RESEARCH LETTERS*,
916 38, doi:10.1029/2011GL047848, 2011.

917 Pongratz, J., Reick, C. H., Raddatz, T., and Claussen, M.: Effects of anthropogenic land cover
918 change on the carbon cycle of the last millennium, *Global Biogeochem. Cycles*, 23, n/a-n/a,
919 doi:10.1029/2009GB003488, 2009.

920 Raddatz, T. J., Reick, C. H., Knorr, W., Kattge, J., Roeckner, E., Schnur, R., Schnitzler, K.-G.,
921 Wetzel, P., and Jungclaus, J.: Will the tropical land biosphere dominate the climate-carbon
922 cycle feedback during the twenty-first century?, *Climate Dynamics*, 29, 565–574,
923 doi:10.1007/s00382-007-0247-8, 2007.

924 Rautiainen, M., Stenberg, P., Mottus, M., and Manninen, T.: Radiative transfer simulations link
925 boreal forest structure and shortwave albedo, *BOREAL ENVIRONMENT RESEARCH*, 16,
926 91–100, 2011.

927 Reick, C. H., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and
928 anthropogenic land cover change in MPI-ESM, *J. Adv. Model. Earth Syst.*, 5, 459–482,
929 doi:10.1002/jame.20022, 2013.

930 Riahi, K., Rao, S., Krey, V., Cho, C., Chirkov, V., Fischer, G., Kindermann, G., Nakicenovic,
931 N., and Rafaj, P.: RCP 8.5—A scenario of comparatively high greenhouse gas emissions,
932 *Climatic Change*, 109, 33–57, doi:10.1007/s10584-011-0149-y, 2011.

933 Richards, K. R. and Stokes, C.: A review of forest carbon sequestration cost studies: a dozen
934 years of research, *Climatic Change*, 63, 1–48, 2004.

935 Schelhaas, M.-J., Hengeveld, G., Moriondo, M., Reinds, G. J., Kundzewicz, Z. W., ter Maat,
936 H., and Bindi, M.: Assessing risk and adaptation options to fires and windstorms in European
937 forestry, *Mitig Adapt Strateg Glob Change*, 15, 681–701, doi:10.1007/s11027-010-9243-0,
938 2010.

939 Sitch, S., Huntingford, C., Gedney, N., Levy, P. E., Lomas, M., Piao, S. L., Betts, R., Ciais, P.,
940 Cox, P., Friedlingstein, P., Jones, C. D., Prentice, I. C., and Woodward, F. I.: Evaluation of the
941 terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five
942 Dynamic Global Vegetation Models (DGVMs), *GLOBAL CHANGE BIOLOGY*, 14, 2015–
943 2039, doi:10.1111/j.1365-2486.2008.01626.x, 2008.

944 Sohngen, B. and Tian, X.: Global climate change impacts on forests and markets, *Forest Policy
945 and Economics*, 72, 18–26, doi:10.1016/j.forpol.2016.06.011, 2016.

946 Sohngen, B., Tian, X., and Kim, J.: Global Climate Change Impacts on Forests and Markets,
947 *Forest Policy and Economics*, 18–26, 2016.

948 Stehfest, E., van Vuuren, D., Kram, T., and Bouwman, L.: Integrated Assessment of Global
949 Environmental Change with IMAGE 3.0, PBL Netherlands Environmental Assessment
950 Agency, The Hague, 2014, 2014.

951 Stern, N.: *The Economics of Climate Change: The Stern Review*, Cambridge University Press,
952 2007.

953 Temperli, C., Bugmann, H., and Elkin, C.: Adaptive management for competing forest goods
954 and services under climate change, *Ecological Applications*, 2065–2077, 2012.

955 Tian, X., Sohngen, B., Kim, J. B., Ohrel, S., and Cole, J.: Global climate change impacts on
956 forests and markets, *ENVIRONMENTAL RESEARCH LETTERS*, 11, doi:10.1088/1748-
957 9326/11/3/035011, 2016.

958 UNFCCC: Report of the Conference of the Parties serving as the meeting of the Parties to the
959 Kyoto Protocol on its seventh session, held in Durban from 28 November to 11 December 2011:
960 Addendum decisions adopted by the Conference of the Parties serving as the meeting of the
961 Parties to the Kyoto Protocol, United Nations, Geneva, Nairobi, New York, Vienna, 49 pp.,
962 2012.

963 van Kooten, G. C., Kremer-Nozic, E., Stennes, B., and van Gorkom, R.: Economics of fossil
964 fuel substitution and wood product sinks when trees are planted to sequester carbon on
965 agricultural lands in western Canada, *Can. J. For. Res.*, 29, 1669–1678, doi:10.1139/cjfr-29-11-
966 1669, 1999.

967 van Vuuren, D. P., Stehfest, E., den Elzen, Michel G. J., Kram, T., van Vliet, J., Deetman, S.,
968 Isaac, M., Klein Goldewijk, K., Hof, A., Mendoza Beltran, A., Oostenrijk, R., and van Ruijven,
969 B.: RCP2.6: Exploring the possibility to keep global mean temperature increase below 2°C,
970 *Climatic Change*, 109, 95–116, doi:10.1007/s10584-011-0152-3, 2011.

971 Winckler, J., Reick, C. & Pongratz, J.: Robust identification of local biogeophysical effects of
972 land-cover change in a global climate model. *Journal of Climate*, 30, 1159-
973 1176, doi:10.1175/JCLI-D-16-0067.1, 2017

974 Yousefpour, R. and Hanewinkel, M.: Forestry professionals' perceptions of climate change,
975 impacts and adaptation strategies for forests in south-west Germany, *Climatic Change*, 130,
976 273–286, doi:10.1007/s10584-015-1330-5, 2015.

977 Yousefpour, R., Jacobsen, J. B., Thorsen, B. J., Meilby, H., Hanewinkel, M., and Oehler, K.: A
978 review of decision-making approaches to handle uncertainty and risk in adaptive forest
979 management under climate change (vol 69, pg 1, 2012), *ANNALS OF FOREST SCIENCE*,
980 69, 531, doi:10.1007/s13595-012-0192-5, 2012.

981 Zaehle, S., Medlyn, B. E., Kauwe, M. G. de, Walker, A. P., Dietze, M. C., Hickler, T., Luo, Y.,
982 Wang, Y.-P., El-Masri, B., Thornton, P., Jain, A., Wang, S., Warlind, D., Weng, E., Parton, W.,
983 Iversen, C. M., Gallet-Budynek, A., McCarthy, H., Finzi, A. C., Hanson, P. J., Prentice, I. C.,

Gelöscht:

Gelöscht:

984 Oren, R., and Norby, R. J.: Evaluation of 11 terrestrial carbon-nitrogen cycle models against
985 observations from two temperate Free-Air CO₂ Enrichment studies, *NEW PHYTOLOGIST*,
986 202, 803–822, doi:10.1111/nph.12697, 2014.

Gelöscht:

Gelöscht:

987 Figure's Captions

988 **Figure 1** Development of **global standing** wood carbon pools forced by three different RCP
989 scenarios and subject to the harvesting rules of the representative concentration pathways
990 (RCP2.6, RCP4.5 and RCP8.5) **or subject to growth-based harvesting (GB2.6, GB4.5, and**
991 **GB8.5) (1a). Development of RCP wood harvest rates, of the growth potential of forests**
992 **under GB, and of the harvest potential under GB limited to global managed forest area**
993 **(MF2.6, MF4.5, and MF8.5) (1b). All lines are smoothed over 10 years.**

994 **Figure 2** **Spatial distribution of the harvest realized in JSBACH when harvest rates are**
995 **prescribed from the representative concentration pathways (left panels), of the harvest**
996 **potential applying the growth-based harvesting rule to available managed forest area**
997 **(right panels) and of the underlying growth potential (middle panels). All values are**
998 **summed over the entire simulated period (2006-2100).**

999 **Figure 3** Composition of **growth-based harvest (GB)** forced by different climate change
1000 scenarios (RCP2.6, RCP4.5, and RCP8.5 in figures a, b, and c, respectively). dCw/dt refers
1001 to the difference in changes in above-ground woody biomass between representative
1002 concentration pathways' and **GB** harvest (where changes in biomass in **GB** are by
1003 construction of the harvest rule close to 0), **BG**mort refers to the difference in woody
1004 carbon losses between RCP and **GB** harvest due to background mortality, **Fire** to that due
1005 to fire disturbance, and **Wind** to that due to wind disturbance. **GB** and RCP harvest are
1006 as in Figure 1b.

1007 **Figure 4** **Net mitigation potentials from the growth potential under the growth-based**
1008 **harvesting rule (GB) (a, b, c), representative concentration pathways' (RCP) harvest (d,**
1009 **e, f), and GB harvest limited to managed forest area (MF) (g, h, i). Left axes show the**
1010 **annual carbon fluxes due to harvested material and product decay changing atmospheric**

Gelöscht: (1a) and realized wood harvest (1b)

Gelöscht: and sustained yield (SY2.6, SY4.5, and SY8.5).
Lines

Gelöscht: Allocation

Gelöscht: wood

Gelöscht: applying sustained yield (three left figures) and

Gelöscht: three right figures) rules

Gelöscht: different

Gelöscht: regions

Gelöscht: sustained-yield (SY)

Formatiert: Schriftart: Fett

Gelöscht: SY

Formatiert: Schriftart: Fett

Gelöscht: SY

Formatiert: Schriftart: Fett

Gelöscht: SY

Formatiert: Schriftart: Fett

Gelöscht: SY

Formatiert: Schriftart: Fett

Formatiert: Beschriftung, Block

Gelöscht: Mitigation

Gelöscht: of simulated wood harvest

Gelöscht: sustained yields (SY

Gelöscht:) and

Formatiert: Schriftart für komplexe Schriftzeichen:
Nicht Fett

Gelöscht:

Gelöscht:

1029 CO₂ concentration, and the mitigation potential of wood products as the difference of
1030 both. Right axes **accumulate** the annual figures over time.

- Gelöscht:** accumulates
- Gelöscht:** ¶
- Formatiert:** Schriftart für komplexe Schriftzeichen:
Nicht Fett

- Gelöscht:**
- Gelöscht:**

1033 **Table 1: JSBACH simulations conducted in this study with the applied harvesting rule**
 1034 **and climate and CO₂ forcing.**

Name	Harvest rule	MPI-ESM forcing
GB2.6	GB	RCP2.6
GB4.5	GB	RCP4.5
GB8.5	GB	RCP8.5
RCP2.6	RCP2.6 map	RCP2.6
RCP4.5	RCP4.5 map	RCP4.5
RCP8.5	RCP8.5 map	RCP8.5

- Gelöscht: SY2
- Gelöscht: SY
- Gelöscht: SY4
- Gelöscht: SY
- Gelöscht: SY8
- Gelöscht: SY

1035

1036

- Gelöscht:
- Gelöscht:

1043 **Table 2 Net mitigation potentials of GB, MF and RCP harvest at the middle and end of**
 1044 **the 21st century**

Gelöscht: Mitigation

Gelöscht: SY

Applied harvest rule	Harvested wood (PgC)		Mitigation effect (PgC)	
	2050	2100	2050	2100
RCP2.6	58.1	137.6	38.3	85.1
RCP4.5	62.9	147.2	40.7	90.2
RCP8.5	76.5	211.8	50.6	132.1
GB2.6	192.7	421.3	124.5	255.0
GB4.5	210.0	513.9	136.4	314.7
GB8.5	215.0	609.4	140.6	379.1
MF2.6	148.3	324.3	96.6	199.5
MF4.5	161.6	395.1	105.6	244.9
MF8.5	166.4	472.9	109.3	295.8

Gelöscht: SY2

Gelöscht: SY4

Gelöscht: SY8


1045

1046


Gelöscht:

Gelöscht:

1052 **Figure 1**



1053
1054




Gelöscht:

Gelöscht:

Gelöscht:


1056 **Figure 2**

1057



1058

1059




Gelöscht:

Gelöscht:

Gelöscht:


1061 **Figure 3**

1062



1063



1064



Gelöscht:

Gelöscht:

Gelöscht:



Gelöscht:

Gelöscht:

Gelöscht:

S1: Supplementary material 1

A1. JSBACH simulations

Simulations were conducted with revision 7277 of cosmos-landveg-fom, a svn branch of revision 7215 of cosmos-landveg, the former JSBACH development branch of the department "The Land in the Earth System" of the Max Planck Institute for Meteorology.

Simulations were executed on the IBM Power 6 machine BLIZZARD at the German Climate Computing Center (DKRZ).

Growth-based (GB) forest harvesting is implemented in this model version as described in the methods section of the main text. A modification over earlier JSBACH versions is that wood harvest applies just to harvesting from all woody PFTs and specifically from the above-ground wood carbon pool. To isolate the effects of different wood harvest rules, we do not apply land-cover change and dynamic biogeographic vegetation shifts for our future scenarios. We take into account changes in wood carbon pool, natural mortality and forest disturbances to determine the net annual increment of the above-ground wood carbon pool as the maximum amount to be harvested from forest areas.

Gelöscht: Sustained-yield (SY)

Gelöscht: use

A1.1 Initial state in 2006

All simulations described in the paper started in 2006 from the same initial conditions. These conditions base on a spin-up of the terrestrial system state using the MPI-ESM climate from the historical (1850-2005) CMIP5 experiment (Giorgetta et al., 2013) and land-use change and wood harvest data from Hurtt et al. (2011).

The initial state was derived carrying out three consecutive simulations. (I) An initial simulation with JSBACH to spin-up photosynthesis, phenology, hydrology and running climatic means required by the disturbance module of JSBACH. This simulation was forced by cycling the first

30 years (1850-1879) of the historical CMIP5 experiment for four times. Wood harvest was fixed to the level of the initial year 1850 and no land-use change was applied. (II) A simulation with the stand-alone carbon cycle module of JSBACH to equilibrate the carbon pools with respect to the driving climate. This simulation was forced by NPP, LAI and climatic means, resulting from the preceding JSBACH simulation. (III) A second JSBACH simulation resuming the first JSBACH simulation, but starting from the equilibrated carbon pools. In this second simulation the full transient (1850-2005) climate from the historical CMIP5 experiment was used and land-use change and wood harvest were prescribed according to the data from Hurtt et al. (2011).

A1.2 Reference level for **GB**

An important decision for our study is the definition of the reference level of the wood carbon pool to be kept constant in the future applying **GB**. As one of the main goals of our study is to estimate potentials for wood harvesting under future climate scenarios, consistent with the historic past, we refer to the current level of wood carbon pools. The reference level for the **GB** simulations was therefore derived from the maximum simulated wood carbon per grid-cell and PFT in the period from 1996 to 2005 under the historical JSBACH simulation (see A1.1 simulation III). Because the historical JSBACH simulation was subject to land-use and land-cover changes maximum wood carbon densities were used instead of wood carbon stocks.

A1.3 Simulation of **GB** and **MF** under present-day climate

These simulations (GBpd and MFpd) keep the current level of wood carbon pools constant as described above in A1.2, **and apply the GB harvest rule to global and managed forest area.** However, **they are** not forced by a transient but a cycled detrended present-day climate of the

Gelöscht: SY

Gelöscht: SY

Gelöscht: SY

Gelöscht: SY

Gelöscht: This simulation (SYpd) keeps

Gelöscht: .

Gelöscht: it is

period 1991-2020 with a constant CO₂ concentration (381 ppm) as the average value of the period (1991-2020). S1-Figure 1 shows the development of wood carbon pool and harvested amount resulting in the simulation GBpd compared to the 6 simulations described in the main text. GBpd realizes a higher wood harvest (+3.2 PgC) than RCPs (~1.2 PgC) at the beginning of the simulations and equals the RCP8.5 wood harvest at the end of the century. GBpd diverges from the GB2.6, GB4.5, and GB8.5 wood harvest amounts largely towards the end of the century and remains below these figures. The geographical allocation of realized wood harvest amount as shown below in GBpd in S1-Figure 2 resembles largely the other GBs (see Figure 2 in the manuscript), however, the amount of harvested wood is lower. Values for the simulated wood harvest from MFpd is lower than GBpd because of limiting forest harvest to managed forest area (excluding primary forest area). S1-Figure 3 shows the net mitigation of MF forced by present day climate. Logically, the annual harvest amount stay more or less constant (~3.2 PgC) in the 21st century. This is exactly resembling the concept of sustained yield if no changes in forest growth is expected. As a result, MFpd would result in a lower net mitigation potential (~150 PgC) than GB and MF (see Figure 4 in text for details), applying the same life cycle analysis described in section 2.6.

Gelöscht: SYpd

Gelöscht: SYpd

Gelöscht: SYpd

Gelöscht: wood harvest amount by SY2

Gelöscht: SY4

Gelöscht: SY8

Gelöscht: SYpd

Gelöscht: SYs

Formatiert: Englisch (USA)

Formatiert: Links, Zeilenabstand: Mehrere 1.08 ze

S1-Figure 1 Development of **global standing wood carbon pools** forced by three different RCP scenarios, and a present-day (pd) forcing, subject to the harvesting rules of the representative concentration pathways (RCP2.6, RCP4.5 and RCP8.5) or subject to growth-based harvesting (GB2.6, GB4.5, GB8.5, and GBpd) (1a). Development of RCP wood harvest rates, of the growth potential of forests under GB and of the harvest potential under GB limited to global managed forest area (MF2.6, MF4.5, MF8.5, and MFpd) (1b). All lines are smoothed over 10 years.



Gelöscht: (1a) and realized wood harvest (1b)

Gelöscht: ,

Gelöscht: climate, and

Gelöscht: and sustained yield (SY2.6, SY4.5, SY8.5, and SYpd). Lines

Formatiert: Schriftart: Nicht Fett






Gelöscht: --- RCP2.6 --- RCP4.5 --- RCP8.5

Gelöscht: --- GB2.6 --- GB4.5 --- GB8.5 --- GBpd --- MF2.6 --- MF4.5 --- MF8.5 --- MFpd

S1-Figure 2 Allocation of wood harvest applying **growth-based harvesting rule to the global forest area (GB)** and **limited to managed forest area (MF)** under **present-day forcing** summed over the entire simulated period (2006-2100).

Gelöscht: sustained yield

Gelöscht: climate to different forest regions



Gelöscht:


References¶

Giorgetta MA et al. (2013) Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5. *J. Adv. Model. Earth Syst.* 5:572–597.¶

Hurt, GC et al (2011) Harmonization of land-use scenarios for the period 1500–2100. 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands. *Climatic Change* 109:117–161.

Formatiert: Dänisch

S1-Figure 3 Net mitigation potentials of simulated wood harvest from growth-based harvest rule applied to managed forest area under cycled present day forcing (MFpd). Left axis shows the annual carbon fluxes due to harvested material and product decay changing atmospheric CO₂ concentration, and the mitigation potential of wood products as the difference of both. Right axis accumulates the annual figures over time.



References

- Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Ilyina, T., Kinne, S., Kornblueh, L., Matei, D., Mauritsen, T., Mikolajewicz, U., Mueller, W., Notz, D., Pithan, F., Raddatz, T., Rast, S., Redler, R., Roeckner, E., Schmidt, H., Schnur, R., Segschneider, J., Six, K. D., Stockhause, M., Timmreck, C., Wegner, J., Widmann, H., Wieners, K.-H., Claussen, M., Marotzke, J., and Stevens, B.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, *J. Adv. Model. Earth Syst.*, 5, 572–597, doi:10.1002/jame.20038, 2013.
- Hurtt, G. C., Chini, L. P., Frohling, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P., Hibbard, K., Houghton, R. A., Janetos, A., Jones, C. D., Kindermann, G., Kinoshita, T., Klein Goldewijk, K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D. P., and Wang, Y. P.: Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands, *Climatic Change*, 109, 117–161, doi:10.1007/s10584-011-0153-2, 2011.
- Kraxner, F., Schepaschenko, D., Fussler, S., Lunnanc, A., Kindermann, G., Aokie, K., Dürauer, M., Shvidenko, A., See, L.: Mapping certified forests for sustainable management - A global tool for information improvement through participatory and collaborative mapping. *Forest Policy and Economics* 83:10–18, 2017