


1 Supplement


2



3
4
5


Figure S1. Location of the nine study sites (Globcover). All sites are located in old growth tropical forests.

6



7

8 **Figure S2** : Description of steps taken from the original Lidar canopy height model to the final LCA product. This
9 example is a 400 m by 400 m area in BCI, with a Height threshold of 30 m and the minimum number of contiguous
10 pixels set to 100.



11 **Figure S3.** Lidar derived images: shaded relief of topography and canopy height model (CHM), at 1 m resolution.
 12 Mean canopy height (MCH), standard deviation of canopy height and elevation range are reported.
 13
 14

15 **S.1 Estimating aboveground biomass from forest inventories.**

16
17 For trees with no height measurement, a site specific DBH height model was used to infer tree
18 height in each site, as described in previous studies (Feldpauch et al., 2012; Meyer et al., 2013,).
19 Wood density was extracted from the global wood density database for tropical trees (Chave et
20 al., 2009; Zanne et al., 2009) for each tree, based on its level of botanical identification (species,
21 genus, family). Trees with no botanical identification were assigned the average wood density of
22 the plot. Average wood density of each site was calculated as the unweighted average of all trees
23 within a site, or was taken from a previous study in the case of Cotriguaçu (Fearnside, 1997).
24 Average wood density of large trees was calculated as the unweighted average of trees with
25 $DBH \geq 50$ cm in each site. Tree level AGB was aggregated at the plot level using a commonly
26 used allometric regression model for moist tropical forests (Eq. (S1), Chave et al., 2005), except
27 for La Selva and Chocó, for which a model for wet tropical forest (Eq. (S2), Chave et al., 2005)
28 and a local allometric model (Eq. (S3), Duque et al., 2017) were used, respectively.

$$29 \quad AGB_{moist} = 0.0509 \times \rho DBH^2 H \quad (S1)$$

$$30 \quad AGB_{wet} = 0.0776 \times (\rho DBH^2 H)^{0.940} \quad (S2)$$

$$31 \quad AGB_{Chocó} = 0.089 \times (\rho DBH^2 H)^{0.951} \quad (S3)$$

32 where trunk diameter (DBH in cm) is measured during the inventory, tree height (H , in m) is
33 either measured in the field or estimated from a local DBH – H model, and specific gravity or
34 wood density (ρ in $g\ cm^{-3}$) is known for each tree. AGB (in kg of dry biomass) of individual
35 trees estimated using the former equations was used to calculate plot level AGB density ($Mg\ ha^{-1}$)
36 by summing over the biomass of all stems within each plot. Ground estimated AGB density is
37 henceforth referred to as AGB_{inv} . Estimating AGB with these allometric models has been

38 reported to have a standard error of 12.5 % when using height as a parameter, against 19.5 %
39 when height is not available (Chave et al., 2005).

40

41 **Table S1.** Coefficients, R^2 and RMSE for the local models used to estimate AGB from MCH in the different sites,
 42 based on available ground data.

Site	a	b	R^2	RMSE	plot size
Antimary	0.1687	2.2544	0.62	45.3	0.25
BCI	2.3257	1.4336	0.68	28.14	1
Chocó	10.7328	1.0509	0.66	33.34	0.25
Cotriguacu	3.0249	1.3895	-	-	-
La Selva	2.1241	1.3889	0.84	11.45	1
Manaus	3.0249	1.3895	0.59	43.73	0.25
Nouragues	5.7963	1.242	0.46	59.19	1
Paracou	5.707	1.239	0.58	40.49	1
Tapajos	3.0249	1.3895	0.59	43.73	0.25

43

44

45

46 **Table S2.** Matrices of canopy height thresholds (20 m to 50 m, presented with a 5 m increment for readability
 47 purposes) and crown area thresholds (50 to 200 pixels or m²), represented respectively by the minimum height
 48 considered for segmentation and by the minimum number of contiguous 1m pixels in a segment. Values are given in
 49 percentage of coverage of the 1km² subsets.

Site	Number of pixels	20 m threshold	25 m threshold	30 m threshold	35 m threshold	40 m threshold	45 m threshold	50 m threshold
Antimary	50	75.30	48.32	22.19	8.23	2.12	0.46	0.06
	100	75.03	47.40	20.92	7.64	1.92	0.43	0.05
	150	74.84	46.55	19.82	7.03	1.75	0.41	0.04
	200	74.81	45.89	18.93	6.33	1.60	0.34	0.04
BCI	50	70.06	51.53	29.19	11.66	2.51	0.25	0.05
	100	69.79	50.92	28.21	11.08	2.30	0.21	0.05
	150	69.48	50.38	27.44	10.52	2.06	0.17	0.05
	200	69.35	49.96	26.67	10.06	1.95	0.16	0.05
Chocó	50	70.48	48.64	23.22	7.72	1.88	0.24	0.02
	100	70.20	47.58	21.31	6.45	1.53	0.18	0.00
	150	69.97	46.81	19.83	5.52	1.17	0.09	0.00
	200	69.84	46.05	18.19	4.83	0.94	0.06	0.00
Cotriguaçu	50	80.07	53.85	25.93	8.12	1.65	0.24	0.02
	100	79.91	52.87	24.70	7.33	1.43	0.21	0.02
	150	79.85	52.08	23.23	6.48	1.26	0.18	0.02
	200	79.84	51.71	21.92	5.70	1.12	0.18	0.02
La Selva	50	74.32	50.22	24.00	7.88	1.95	0.27	0.06
	100	74.15	49.30	22.71	7.17	1.70	0.25	0.05
	150	74.05	48.69	21.57	6.55	1.47	0.21	0.05
	200	74.01	48.03	20.57	6.01	1.22	0.11	0.03
Manaus	50	86.07	66.42	33.69	9.02	1.63	0.41	0.08
	100	86.04	65.99	32.72	7.91	1.48	0.38	0.07
	150	86.03	65.69	31.20	6.99	1.17	0.34	0.03
	200	86.00	65.51	29.87	6.08	1.05	0.32	0.03
Nouragues	50	85.14	69.77	48.77	28.82	13.34	4.05	0.79
	100	85.05	69.37	47.86	27.73	12.45	3.56	0.60
	150	84.93	69.16	47.23	26.67	11.41	3.02	0.44
	200	84.89	68.87	46.62	25.73	10.30	2.51	0.29
Paracou	50	84.56	63.40	31.82	9.56	1.83	0.17	0.00
	100	84.50	62.90	30.18	8.47	1.46	0.10	0.00
	150	84.46	62.43	28.64	7.36	1.10	0.05	0.00
	200	84.42	62.18	27.61	6.30	0.82	0.23	0.00
Tapajós	50	79.61	66.61	51.61	33.46	15.15	4.06	0.60
	100	79.46	66.24	51.37	32.89	14.42	3.56	0.48
	150	79.35	65.98	45.89	31.80	13.51	3.19	0.35
	200	79.34	65.69	50.18	30.92	12.64	2.67	0.26


50
51

52 **Table S3.** Coefficients, R^2 , RMSE and bias for the models used to estimate AGB from MCH without and with wood
 53 density (WD) as a weighting factor

Model	Equation	a	b	R^2	RMSE	Bias	R^2 cross-val	RMSE cross-val	Bias cross-val
m_MCH	$AGB = aMCH^b$	0.91	1.77	0.68	54.76	-0.3	0.67	55.38	-0.54
m_MCH_wd	$AGB = a(MCH \times WD)^b$	9.29	1.23	0.79	44.23	0.02	0.79	44.78	0.01

54
 55
 56

57



58

59 **Figure S4.** Distribution of R^2 between DBH thresholds and AGB_{Lidar} in 1 ha subareas (a) and AGB_{inv} in 1 ha inventory
60 plots (b). DBH was calculated from tree height using Chave's E factor model (Chave et al., 2014)

61

62

63 **S.2 LCA and ground data**

64 We looked at the relation between LCA and other stand level metrics: AGB_{inv} of large trees,
65 stand basal area, Lorey's height (H_L), and the number of large trees. Lorey's height is defined by

66 $H_L = \frac{\sum_{i=1}^N BA_i H_i}{\sum_{i=1}^N BA_i}$ where H_i is the height of stem i , and BA_i is the basal area of stem i . Weighting

67 tree height with basal area increases the relative importance of large trees in a stand (Lefsky,
68 2010).

69 There was a significant correlation between LCA and AGB_{inv} in all calibration sites ($R^2_{max} = 0.77$
70 in La Selva), and even more so between LCA and AGB_{inv} of large trees ($R^2_{max} = 0.83$ in La
71 Selva) (Table S4). LCA correlated with Lorey's height, especially at La Selva but less so at
72 Nouragues ($R^2_{LaSelva} = 0.86$, $R^2_{Nouragues} = 0.21$). We also found that LCA and total basal area are
73 correlated. The relationship between LCA and the number of large trees is significant in all sites,
74 but it is lower than other structural metrics in Paracou ($R^2 = 0.32$). Among study sites, La Selva
75 and BCI show higher correlations between LCA and ground derived metrics. The strong
76 correlation of LCA with plot level AGB_{inv} as well as with AGB_{inv} of large trees suggested that a
77 model based on LCA can be used as an estimator of AGB.

78

79 **Table S4** : Coefficients of correlation (R^2) between LCA and ground data derived metrics : AGB_{inv} , AGB_{inv} of
 80 trees ≥ 50 cm DBH, basal area, Lorey's height and number of large trees in BCI, La Selva, Nouragues and Paracou.

	BCI	La Selva	Nouragues	Paracou
AGB_{inv}	0.64	0.77	0.48	0.64
AGB_{50}	0.68	0.83	0.47	0.69
BA	0.63	0.53	0.41	0.43
LH	0.57	0.86	0.21	0.63
N_{70}	0.64	0.76	0.46	0.32

81

82

83 **S.3 Contribution of Large Trees to AGB_{inv}**


84 First, we explored the role of wood density as a key variable in allometric models to scale the
85 tree volume or size to AGB_{inv} (Brown et al., 1989; Chave et al., 2005; Chave et al., 2014,
86 Ngomanda et al., 2014). We examined the variations of wood density as a function of DBH
87 classes in all sites and for all trees greater than 10 cm arranged in 10 cm DBH bins.

88 We then assessed the proportion of large trees by examining the DBH frequency distribution of
89 trees at different sites using the field inventory data, and compared it to the distribution of
90 AGB_{inv} for the same DBH ranges, using 1 cm bins to create DBH classes.

91 **Results**

92 DBH and wood density: At BCI, mean wood density decreased significantly with increasing
93 DBH ($p < 0.001$, $R^2 = 0.92$), which is consistent with the results of a previous study (Chave et al.,
94 2004). On the contrary, at Chocó, mean wood density increased significantly with increasing
95 DBH ($p < 0.001$, $R^2 = 0.8$). No significant trend was found at the other study sites (Fig. S5). These
96 results show that stand average wood density and wood density of large trees are not
97 significantly different in four sites and can therefore be used interchangeably in a model using
98 these data. However, using one or the other will have an impact on the results of a model
99 calibrated using data from BCI or Chocó. Mean wood density of trees ≥ 50 cm is 0.49 in BCI (vs.
100 stand average $WD = 0.54 \text{ g cm}^{-3}$) and 0.66 in Chocó (vs. stand average $WD = 0.60 \text{ g cm}^{-3}$).

101



102

103 **Figure S5.** Wood density and DBH classes in BCI, Chocó, La Selva, Manaus, Nouragues and Tapajós. The decrease
 104 in mean wood density is significant in BCI ($p < 0.001$, $R^2 = 0.92$). The increase in mean wood density is significant in
 105 Chocó ($p < 0.001$, $R^2 = 0.8$) but there is no significant wood density change across DBH classes in the other sites
 106 ($p > 0.1$). Points with no standard deviation bars mean that only one tree was present in the class.

107

108 We found that using mean wood density of trees ≥ 50 cm DBH in our LCA model gives slightly

109 better results than using mean wood density (see Eq. (3)), with R^2 of 0.78, RMSE of 45.34 Mg

110 ha^{-1} and bias of -0.36 Mg ha^{-1} .

111

112 DBH and AGB : The relationship between AGB_{inv} and the DBH frequency distribution was

113 similar in BCI, La Selva, Nouragues and Tapajós (Fig. S6). In these sites, more than 50 % of the

114 total AGB_{inv} was contributed by around 5 % (between 3 % and 6 %) of the stems, corresponding


115 to a threshold DBH of 58 cm, 50 cm, 55 cm and 61 cm respectively for the four study sites.

116 Similar results were found in a recent study focusing on the contribution on large trees to AGB

117 in Central Africa (Bastin et al., 2015). Also, 80 % of the AGB stock was in trees with $\text{DBH} \geq 30$

118 cm DBH at BCI, 26 cm at La Selva, 32 cm at Nouragues and 29 cm at Tapajós, which echoes

119 results published on BCI (Chave et al., 2001). These represent only 20 % of the total number of
 120 trees in each site (only 15 % in Tapajós). Chocó and Manaus had far less large trees, 50 % of
 121 their total AGB_{inv} being explained by respectively 10 % and 7 % of the total number of trees,
 122 corresponding to a DBH greater than only 28 cm and 40 cm. In these two sites, 80 % of AGB_{inv}
 123 is found in trees above 17 cm and 23 cm, respectively, representing 37 % and 31 % of their total
 124 number of trees. These findings corroborate with previous studies from other sites and suggests
 125 that large trees can explain AGB variations and could potentially be used to estimate biomass
 126 without having to measure all trees in a plot (Bastin et al., 2015; Slik et al., 2013). Note that
 127 some ground data from Manaus and Nouragues were included in Silk et al.'s dataset).



128
 129 **Figure S6.** Cumulation of of AGB_{inv} and of number of trees (total number of trees ≥ 10 cm DBH) as a function of
 130 DBH in BCI, Chocó, La Selva, Manaus, Nouragues and Tapajós plots.
 131

132
 133 **References (Supplement)**

135 Chave, J., Riéra, B., and Dubois, M. A.: Estimation of biomass in a neotropical forest of French
 136 Guiana: spatial and temporal variability. *J. Trop. Ecol.*, 17(01), 79-96, 2001.
 137

138 Chave, J., Andalo, C., Brown, S., Cairns, M., Chambers, J., Eamus, D., Folster, H., Fromard, F.,
139 Higuchi, N., Kira, T., Lescure, J. P., Nelson, B., Ogawa, H., Puig, H., Rié ra, B., and Ya-
140 T.: Tree allometry and improved estimation of carbon stocks and balance in tropical forests,
141 *Oecologia*, 145, 87–99, 2005.

142

143 Chave, J., Coomes D. A., Jansen S, Lewis, S. L., Swenson, N. G. and Zanne, A. E.: Towards a
144 worldwide wood economics spectrum. *Ecol. Lett.* 12(4), 351-366.
145 <http://dx.doi.org/10.1111/j.1461-0248.2009.01285.x>, 2009.

146

147 Duque, A., Saldarriaga, J., Meyer, V., and Saatchi, S.: Structure and allometry in tropical forests
148 of Chocó, Colombia. *Forest Ecol. Manag.*, 405, 309-318, 2017.

149

150 Feldpausch, T. R., Lloyd, J., Lewis, S. L., Brienen, R. J., Gloor, M., Monteagudo Mendoza, A.,
151 Lopez-Gonzalez, G., Banin, L., Abu Salim, K., Affum-Baffoe, K. and Alexiades, M.: Tree height
152 integrated into pantropical forest biomass estimates. *Biogeosciences*, pp.3381-3403, 2012.

153

154 Lefsky, M. A.: A global forest canopy height map from the Moderate Resolution Imaging
155 Spectroradiometer and the Geoscience Laser Altimeter System, *Geophys. Res. Lett.*, 37, L15401,
156 doi:10.1029/2010GL043622, 2010.

157

158 Ngomanda, A., Obiang, N. L. E., Lebamba, J., Mavouroulou, Q. M., Gomat, H., Mankou, G. S.
159 and Picard, N.: Site-specific versus pantropical allometric equations: Which option to estimate the
160 biomass of a moist central African forest?. *Forest Ecol. Manag.*, 312, 1-9., 2014.

161

162 Zanne, A. E., Lopez-Gonzalez, G., Coomes, D. A., Ilic, J., Jansen, S., Lewis, S. L., Miller, R. B.,
163 Swenson, N. G., Wiemann, M. C., and Chave, J.: Global wood density database. *Dryad*. Identifier:
164 <http://hdl.handle.net/10255/dryad.235>, 2009.

165

166

167