

# ***Interactive comment on “Distribution and rates of nitrogen fixation in the western tropical South Pacific Ocean constrained by nitrogen isotope budgets” by Angela N. Knapp et al.***

**Angela N. Knapp et al.**

anknapp@fsu.edu

Received and published: 15 April 2018

Response to Reviewers We thank the reviewers for their constructive feedback. Based on their suggestions, the following changes have been made:

We have included a figure illustrating the location of the field work (Fig. 1a), as well as included additional data to provide more context for the results. The additional data include a section plot showing the  $\text{NO}_3^- + \text{NO}_2^-$   $\delta^{15}\text{N}$  at additional stations across the transect (Fig. 1b), as well as the fluorescence trace on the figure of the  $\delta^{15}\text{N}$  budgets to indicate the range in depth and magnitude of productivity within the euphotic zone (Fig. 1c).

Printer-friendly version

Discussion paper



Additional changes in response to specific comments are described below.

Review #2 The manuscript by Knapp and colleagues estimates the input of N via N<sub>2</sub> fixation using a biogeochemical approach in the western south Pacific Ocean. The manuscript is well-written, and explains the conclusions well, addressing the relevant references. However, this reviewer considers that a few minor changes need to be addressed before publication.

Specific comments. I do agree with Reviewer 1 in two points. First, though it is clear that this manuscript is related to many other publications coming from OUTPACE probably showing a map of station, having the map here will be useful for readers beginning from this work, instead of having to look for the geographical context on their own. Second, it will be useful to extend a bit the context of the sediment traps, maybe adding an additional line in Fig 1 showing either the mixed layer depth or the The units of Average PN<sub>sink</sub> flux in Table 1 and in the Results section are not the same. The text is in  $\mu\text{mol N}$ , and the table in  $\text{mmol N}$ . I recommend units in agreement in both parts.

As suggested by both reviewers, we have included a map in Fig. 1, as well as additional data, please see the above description. We also thank the Reviewer for their careful attention to detail – the units for the mass flux in Table 1 have been changed so that they are consistent with the units in the text, as well as with the units for N<sub>2</sub> fixation, which improves the readability of the manuscript.

Some parts of section 3.2 could be moved to the Methods section, while others seem to fit better in the Discussion, as a first subsection 4.1. My suggestion is that P7 L25 to P8 L14 and P9 L4 to P11 L7 move to the Discussion, while P8 L15 to L23 up to “of the source NO<sub>3</sub>” move to the Methods. This way the Results subsection is reduced to the description of the results themselves (P8 L23 to P9 L3). P3 L26.

We appreciate the Reviewer’s suggestion, and have moved the last two paragraphs of the original Results section to the beginning of the Discussion section as suggested. However, we felt that the paragraph describing NO<sub>3</sub>-+NO<sub>2</sub>-  $\delta^{15}\text{N}$  gradients with depth

(in the original manuscript P8 L15 to L23), that the Reviewer suggested be moved to Methods, was better suited to the Results section because it felt out of sequence to move a description of the  $\text{NO}_3\text{+NO}_2$ -  $\delta^{15}\text{N}$  gradients with depth to methods before those values were described. Additionally, since we were not interpreting these gradients, just describing the measurements, we felt this text best fit in the Results section. Similarly, we felt that the first paragraph of the Results section was better suited there than in the Discussion, since it describes what the quantitatively relevant terms are in the regional  $\delta^{15}\text{N}$  budgets.

Just curiosity, but is there a reason for using the term Oxygen Deficient Zones instead of the most widely used Oxygen Minimum Zones (OMZs)?

Oxygen deficient zones (ODZs) has become the preferred term to identify water columns “where the water column oxygen concentration is so low (low nanomolar range) that oxygen respiration is precluded and denitrification and other low-oxygen (suboxic) metabolisms predominate” (Devol, 2015, Annual Reviews of Marine Science)”, and is used to differentiate from other water columns which have higher oxygen concentrations, but all of which have a minimum in oxygen concentrations at some depth that typically coincides with the depth of peak rates of oxic respiration (i.e., remineralization).

P5-P6. The description of the geochemical tools could be moved to the Methods section. And it could be more intuitive to begin the name of the variables by  $_{15}\text{N}$ -xx. It is a bit confusing reading  $\text{NO}_3\text{+NO}_2$   $_{15}\text{N}$ , for instance.

The text has been moved as suggested.

P7 L16. What do the authors mean with “thermocline  $\text{NO}_3\text{+NO}_2$ ”? Do they refer to subsurface  $\text{NO}_3\text{+NO}_2$  as in section 3.2, or  $\text{NO}_x$  produced in the thermocline?

Here we refer to  $\text{NO}_3\text{+NO}_2$ - in the thermocline, i.e., between roughly 100 and 800 m, which is plotted in Fig. 1c. This depth range records the isotopic signature of  $\text{NO}_3$ -

[Printer-friendly version](#)[Discussion paper](#)

reduction processes that occurred in the ODZs of the eastern tropical South Pacific. However, in section 3.2 when we use the term “subsurface”, it is to refer to the upper portion of the thermocline, i.e., below the euphotic zone, and specifically the depth range over which the majority of the  $\text{NO}_3^- + \text{NO}_2^-$  that fuels phytoplankton growth in the euphotic zone is sourced.

---

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2017-564>, 2018.

**BGD**

Interactive  
comment

Printer-friendly version

Discussion paper

