

Interplay of community dynamics, temperature, and productivity on the hydrogen isotope signatures of lipid biomarkers

S. Nemiah Ladd^{1,2}, Nathalie Dubois^{2,3}, Carsten J. Schubert^{1,4}

¹Department of Surface Waters – Research and Management, Eawag, Swiss Federal Institute of Aquatic Science and Technology, 6047 Kastanienbaum, Switzerland

²Department of Earth Sciences, ETH Zürich, 8092 Zürich, Switzerland

³Department of Surface Waters – Research and Management, Eawag, Swiss Federal Institute of Aquatic Science and Technology, 8600 Dübendorf, Switzerland

⁴Institute of Biogeochemistry and Pollutant Dynamics, ETH Zürich, 8092 Zürich, Switzerland

10 *Correspondence to:* S. Nemiah Ladd (Nemiah.Ladd@eawag.ch)

Abstract. The hydrogen isotopic composition ($\delta^2\text{H}$) of lipid biomarkers has diverse applications in the fields of paleoclimatology, biogeochemistry, and microbial community dynamics. Large changes in hydrogen isotope fractionation have been observed among microbes with differing core metabolisms, while environmental factors including temperature and nutrient availability can affect isotope fractionation by photoautotrophs. Much effort has gone into studying these effects under laboratory conditions with single species cultures. Moving beyond controlled environments and quantifying the natural extent of these changes in freshwater lacustrine settings and identifying their causes is essential for robust application of $\delta^2\text{H}$ values of common short-chain fatty acids as a proxy of net community metabolism, and of phytoplankton specific biomarkers as a paleohydrologic proxy.

This work targets the effect of community dynamics, temperature, and productivity on $^2\text{H}/^1\text{H}$ fractionation in lipid biomarkers through a comparative time series in two central Swiss lakes: eutrophic Lake Greifen and oligotrophic Lake Lucerne. Particulate organic matter was collected from surface waters at six time points throughout the spring and summer of 2015, and $\delta^2\text{H}$ values of short chain fatty acids, as well as chlorophyll-derived phytol and the diatom biomarker brassicasterol, were measured. We paired these measurements with in situ incubations conducted with $\text{NaH}^{13}\text{CO}_3$, which were used to calculate the production rates of individual lipids in lake surface water. As algal productivity increased from April to June, net discrimination against ^2H in Lake Greifen increased by as much as 148 ‰ for individual fatty acids. During the same time period in Lake Lucerne, net discrimination against ^2H increased by as much as 58 ‰ for individual fatty acids. A large portion of this signal is likely due to a greater proportion of heterotrophically derived fatty acids in the winter and early spring, which are displaced by more ^2H -depleted fatty acids as phytoplankton productivity increases. Smaller increases in ^2H discrimination for phytol and brassicasterol suggest that a portion of the signal is due to changes in net photoautotrophic ^2H fractionation, which may be caused by increasing temperatures, a shift from maintenance to high

growth, or changes in the community assemblage. The fractionation factors for brassicasterol were significantly different between the two lakes, suggesting that its hydrogen isotope composition may be more sensitive to nutrient regime than is the case for fatty acids or phytol.

1 Introduction

5 Compound specific hydrogen isotope measurements of lipid biomarkers are an emerging tool with diverse applications to microbial community dynamics (Osburn et al., 2011; Heinzlmann et al., 2016), organic matter cycling (Jones et al., 2008; Li et al., 2009), and paleoclimatology (Sachse et al., 2012 and sources therein). The hydrogen isotopic composition of source water exerts a first order control on lipid hydrogen isotopes (expressed as $\delta^2\text{H} = ({}^2\text{H}/{}^1\text{H}_{\text{Sample}})/({}^2\text{H}/{}^1\text{H}_{\text{VSMOW}}) - 1$) (Sessions et al., 1999; Sauer et al., 2001; Sachs, 2014). However, a number of variables can influence the offset between the $\delta^2\text{H}$ values
10 between lipids and source water, which is typically expressed by the fractionation factor $\alpha_{\text{Lipid-Water}} = ({}^2\text{H}/{}^1\text{H}_{\text{Lipid}})/({}^2\text{H}/{}^1\text{H}_{\text{Water}})$.

For short-chain ($C < 20$) fatty acids, which can be synthesized by a diverse range of organisms, including photoautotrophs, chemoautotrophs, and heterotrophs, core metabolism typically exerts a large control on $\alpha_{\text{Lipid-Water}}$, with variability in $\delta^2\text{H}$ values exceeding 500 ‰ for organisms grown on the same source water (Zhang et al., 2009a; Osburn et al., 2011; Heinzlmann et al. 2015a; Osburn et al., 2016). These metabolic differences have led to the suggestion that $\delta^2\text{H}$ values of
15 short-chain fatty acids can be used as an indicator of net community metabolism (Zhang et al., 2009a; Osburn et al., 2011; Heinzlmann et al., 2016; Osburn et al., 2016). This application has previously been assessed in coastal marine settings (Heinzlmann et al., 2016) and hot springs (Osburn et al., 2011), but not in lakes.

20 The $\delta^2\text{H}$ values of lipids produced exclusively by photoautotrophs, such as alkenones and certain sterols, have received particular attention as a proxy for past water isotopes (Sessions et al., 1999; Sauer et al., 2001; Huang et al., 2004; Sachse et al. 2012; Sachs, 2014), which is useful for paleoclimatologists seeking to reconstruct changes in temperature, moisture source, and the balance of precipitation to evaporation, all of which influence the $\delta^2\text{H}$ values of water (Craig and Gordon, 1965; Gat, 1996; Henderson and Schuman, 2009; Steinmann et al., 2013). The hydrogen isotopic composition of lipids
25 produced by cyanobacteria and eukaryotic algae is well correlated with those of source water in laboratory and field settings (Sauer et al., 2001; Huang et al., 2004; Englebrecht and Sachs, 2005; Zhang and Sachs, 2007; Sachse et al., 2012), and is stable under near surface temperatures and pressures for carbon-bound hydrogen (Sessions et al., 2004; Schimmelmann et al., 2006). Hydrogen isotopes of biomarkers from eukaryotic algae have been successfully applied to infer changes in past climate using sediment cores from diverse lakes (Huang et al., 2002; Sachs et al., 2009; Smittenberg et al., 2011; Atwood

and Sachs, 2014; Zhang et al., 2014; Nelson and Sachs, 2016; Richey and Sachs, 2016; Randlett et al., 2017) and marine settings (Pahnke et al., 2007; van der Meer et al., 2007; van der Meer et al., 2008; Leduc et al., 2013; Vasiliev et al., 2013; Kasper et al., 2014; Vasiliev et al., 2017).

5 However, among photoautotrophs, there is increasing evidence that $\alpha_{\text{Lipid-Water}}$ is not constant, and can change with variables such as salinity, species, light availability, growth rate, and temperature (summarized in Table 1) (Sachs, 2014 and sources therein; Chivall et al., 2014; M'boule et al., 2014; Nelson and Sachs, 2014; Heinzelmann et al., 2015b; Sachs and Kawka, 2015; van der Meer et al., 2015; Wolhowe et al., 2015; Maloney et al., 2016; Sachs et al., 2016; Sachs et al., 2017). While the array of secondary isotope effects may appear daunting, these relationships can provide useful information about past
10 environmental changes in their own right, and developing a thorough understanding of them is important for robust interpretations of $\delta^2\text{H}_{\text{Lipid}}$ values from phytoplankton.

Most previous investigations into variability in $\alpha_{\text{Lipid-Water}}$ in algal lipid biosynthesis have been done with controlled cultures of eukaryotes in laboratory settings. While similar relationships between salinity and $\alpha_{\text{Lipid-Water}}$ have been observed for
15 eukaryotic algal and cyanobacterial lipids in both laboratory (Schouten et al., 2006; Chivall et al., 2014; M'boule et al., 2014; Heinzelmann et al., 2015b; Maloney et al., 2016; Sachs et al., 2016) and field calibrations (Sachse and Sachs, 2008; Sachs and Schwab, 2011; Nelson and Sachs, 2014), the temperature and growth rate effects observed in cultures have yet to be assessed in lacustrine settings where photoautotrophic $\delta^2\text{H}_{\text{Lipid}}$ values are likely to be applied to reconstruct past hydroclimate. In contrast to cultures, lake water contains a diverse and dynamic community of phytoplankton, most of whom
20 contribute lipids to the sediment that cannot be attributed to one particular species. The culturing data that exist are limited to a few species, many of which are only found in marine environments.

In order to evaluate the significance of temperature and growth rate effects on the hydrogen isotopic composition of algal lipids produced in lakes, we collected monthly samples of particulate organic matter in two lakes in central Switzerland
25 throughout the spring and summer of 2015. Both lakes experience similar changes in surface water temperature during this time period, but one of them (Lake Greifen) is characterized by high nutrient availability and increasing algal productivity and biomass throughout the spring and early summer. The other lake (Lake Lucerne) is oligotrophic and had relatively low, constant rates of algal productivity throughout the study period. We paired measurements of hydrogen isotope fractionation with in situ incubations designed to determine lipid production rates, allowing us to distinguish between the effects of
30 productivity and temperature on hydrogen isotope fractionation.

In addition to measuring $\delta^2\text{H}$ values of brassicasterol (24-methyl cholest-5,22-dien-3 β -ol) and phytol, lipids that are produced exclusively by photoautotrophs, we also analysed short-chain fatty acids ($n\text{C}_{14:0}$, $n\text{C}_{16:0}$, $n\text{C}_{16:1}$, $n\text{C}_{18:1}$), which, although they are typically the most abundant lipids in algal and cyanobacterial cells, are also synthesized by heterotrophic and chemoautotrophic microbes. The time series of fatty acid $\delta^2\text{H}$ values from an oligotrophic and eutrophic lake presented here is the first opportunity to assess how changes in net community metabolism might be recorded by these compounds in lakes.

2 Methods

2.1 Site description

Lake Greifen (Greifensee) is a small perialpine lake, located in the eastern fringes of the Zurich metropolitan area at 47° 21' N and 8° 40' E (Fig. 1). The lake has a surface area of 24 km², and a maximum depth of 32 m. The lake is fed by three small brooks, and has one main outlet, the Glatt Canal. Lake Greifen experienced severe eutrophication in the mid-20th century (Hollander et al., 1992; Keller et al., 2008). Strict government regulations on nutrient inputs were imposed in the 1970s, and the water quality in the lake has since improved, but its deep water remains anoxic and nutrient levels in the upper water column are still elevated. Winter overturn in the lake brings additional nutrients to the surface water, resulting in large phytoplankton blooms in the spring and summer as temperature and light availability increase (McKenzie, 1982). All samples from Lake Greifen were collected from the northern part of the lake, near a permanent platform maintained by Eawag (at 47° 21.99' N, 8° 39.89' E).

Lake Lucerne (Vierwaldstättersee) is a large perialpine lake, located in central Switzerland at 47° 0' N and 8° 30' E (Fig. 1). The lake, which has a total surface area of 116 km², is formed of seven distinct basins, of which the deepest is 214 m. The lake is fed by four alpine rivers: the Reuss, Muota, Engelberger Aa, and Sarner Aa, and its primary outflow is Reuss river from the northwest tip of the lake. Although Lake Lucerne experienced a mild eutrophication event in the 1970s, it is oligotrophic today (Bürgi et al., 1999; Bührer and Ambühl, 2001; Thevenon et al., 2012). All samples from Lake Lucerne were collected from the center of Kreuztricher basin (near 8° 21' N, 47° 0' E), with a water depth of 96 m.

2.2 Sample collection

Particulate material in each lake was collected at approximately monthly intervals throughout the spring and summer of 2015 (mid-April through early September). Surface water (~0.5 m water depth) was filtered onto a pre-combusted 142 mm diameter GF/F filter (0.7 μm pore size) using a WTS-LV Large Volume Pump (McLane, Massachusetts, USA). Pumping began at 7 L/min and continued until the flow rate decreased to 4 L/min or until 25 minutes had passed. All filters were

collected at midday on sunny or mostly sunny days. Filters were wrapped in combusted aluminum foil and stored in a cool box on ice until transport to the laboratory, where they were stored at -20 °C until analysis.

5 Water samples were collected from surface water before and after pumping began. Samples were collected in 4 mL screw cap vials, sealed with electrical tape, and stored at room temperature prior to analysis. Depth profiles of temperature, conductivity, pH, turbidity, and dissolved oxygen were collected for the upper 20 m of the water column each sampling day at the beginning and end of filtration using a multiparameter CTD probe (75M, Sea & Sun Marine Tech, Trappenkamp, Germany).

10 On the morning of each sampling day, 4 x 12.5 L of surface water was collected in acid-rinsed, autoclaved, transparent carboys for in situ incubations. In two of the four carboys, 1 mL of concentrated $\text{NaH}^{13}\text{CO}_3$ solution was added. The other two carboys were not isotopically labelled. Carboys were mixed and attached to a fixed, floating line so that they stayed in the upper 50 cm of lake water throughout the day. After 6 hours, they were retrieved and the contents were filtered onto a pre-combusted 142 mm diameter GF/F filter using a peristaltic pump. Water samples for DIC analyses were collected in 12
15 mL exetainers prior to isotopic labelling, after labelling but before incubation, and after incubation. These samples were sterile filtered through a 0.2 μm syringe filter and stored in the dark at 4 °C prior to analysis.

2.3 Water isotope measurements

Surface water isotope samples were filtered through a 25 mm syringe filter with a 0.45 μm polyethersulfone membrane to
20 remove particulate matter. Water $\delta^2\text{H}$ and $\delta^{18}\text{O}$ values were measured by Cavity Ring Down Spectroscopy (CRDS) on a L-2120i Water Isotope Analyzer (Picarro, Santa Clara, CA, USA) at ETH-Zurich. Each sample was injected seven times in sequence, and the first four values were discarded to avoid any memory effects from the previous sample. Three water standards with known $\delta^2\text{H}$ values of ranging from -161 ‰ to 7 ‰ and $\delta^{18}\text{O}$ values ranging from -22.5 ‰ to 0.9 ‰ were injected at the beginning and end of each sequence, as well as after every 10 samples. These standards were used to correct
25 measured values to the VSMOW scale and to account for any instrumental drift over the course of the sequence. Average standard deviations were 0.4 ‰ for hydrogen isotopes and 0.06 ‰ for oxygen isotopes.

2.4 DIC concentrations and $\delta^{13}\text{C}$ measurements

DIC concentrations were measured on a TOC-L_{CSH/CHN} Total Organic Carbon Analyzer (Shimadzu, Kyoto, Japan). Solutions
30 with DIC concentrations ranging from 5 mg/L to 100 mg/L were injected at the beginning of the sequence to form a calibration curve, and one standard of 50 mg/L was run after every five samples. Samples were analyzed in triplicate.

3.7 mL exetainers were prepared for $\delta^{13}\text{C}$ measurements of DIC by adding 100 μL of concentrated H_3PO_4 and filling the headspace with He. 1 mL of sample water was added with a syringe through the septa of the exetainer. Samples were allowed to equilibrate overnight before analysis. Carbon isotope values were measured on an Isotope Ratio Mass Spectrometer (IRMS) (Isoprime, Stockport, United Kingdom). A standard of known isotopic composition was analyzed after every 6 samples. All samples were measured in duplicate.

2.5 Lipid extraction and purification

An internal standard containing $n\text{C}_{19}$ -alkanol, $n\text{C}_{19}$ -alkanoic acid, and 5α -cholestane was quantitatively added to freeze-dried filters, which were extracted in 30 mL of 9:1 Dichloromethane (DCM)/Methanol (MeOH) in a SOLVpro Microwave Reaction System (Anton Paar, Graz, Austria) at 70 $^\circ\text{C}$ for 5 minutes (Randlett et al., 2017), centrifuged, and the supernatant containing the total lipid extract (TLE) was poured off and evaporated under a gentle stream of N_2 . The TLE was saponified with 3 mL of 1 N KOH in MeOH and 2 mL of solvent-extracted nanopure H_2O for 3 hours at 80 $^\circ\text{C}$ (Randlett et al., 2017), after which the neutral fraction was extracted with hexane. Subsequently, the aqueous phase was acidified to $\text{pH} = 2$, and the protonated fatty acids were extracted with hexane.

Neutral fractions were further purified using silica gel column chromatography, following a scheme modified from Randlett et al. (2017). The sample was dissolved in hexane and loaded onto a 500 mg / 6 mL Isolute Si gel column (Biotage, Uppsala, Sweden). *N*-alkanes were eluted in 4 mL of hexane, aldehydes and ketones in 4 mL of 1:1 Hexane/DCM, alcohols in 4 mL of 19:1 DCM/MeOH, and remaining polar compounds in 4 mL of MeOH. The alcohol fraction was acetylated with 25 μL of acetic anhydride and 25 μL of pyridine for 30 minutes at 70 $^\circ\text{C}$. The $\delta^2\text{H}$ and $\delta^{13}\text{C}$ values of the added acetyl group were determined by analyzing acetylated and unacetylated $n\text{C}_{10}$ -alkanol.

Further purification was necessary in order to obtain base line separation of brassicasterol for $\delta^2\text{H}$ measurements. This was achieved by loading the acetylated alcohol fraction onto 500 mg of Si gel impregnated with AgNO_3 (10% by weight, Sigma Aldrich) in a 6 mL glass cartridge. The first fraction, containing *n*-alkanols and phytol, was eluted with 20 mL of 4:1 hexane/DCM, the second fraction, containing stanols and singly unsaturated sterols (such as cholesterol) with 20 mL of 1:1 hexane/DCM, the third fraction, containing most doubly unsaturated sterols including brassicasterol, with 16 mL of DCM, and the remaining compounds with 4 mL of ethyl acetate.

30

Fatty acid fractions were methylated with 1 mL of BF₃ in MeOH (14% by volume, Sigma Aldrich) for 2 hours at 100 °C. After methylation, 2 mL of nanopure H₂O was added to the sample and the fatty acid methyl esters (FAMES) were extracted with hexane. The δ²H and δ¹³C values of the added methyl group were determined by methylating phthalic acid of known isotopic composition (Arndt Schimmelmann, Indiana University).

5

FAMES and brassicasterol were quantified by gas chromatography – flame ionization detection (GC-FID) (Shimadzu, Kyoto, Japan). Samples were injected by an AOC-20i autosampler (Shimadzu) through a split/splitless injector operated in splitless mode at 280 °C. The GC column was an InertCap 5MS/NP (0.25 mm x 30 m x 0.25 μm) (GL Sciences, Japan) and it was heated from 70 °C to 130 °C at 20 °C/min, then to 320 °C at 4 °C/min, and held at 320 °C for 20 minutes. FAMES were identified by comparing their retention times to an external standard (Fatty Acid Methyl Ester mix from Sulpelco, Ref 47885-U). Brassicasterol and phtylol were identified by comparing their retention times to those obtained by analyzing a subset of samples by gas chromatography – mass spectrometry (GC-MS) under identical conditions. In order to determine how much of the compound was in the original sample, peak areas were normalized to those of the internal standard. Peak areas were quantified relative to an external calibration curve in order to determine suitable injection volumes for isotopic analysis.

15

2.6 Lipid δ²H and δ¹³C measurements

The stable isotope values of individual FAMES and brassicasterol were measured by gas chromatography – isotope ratio mass spectrometry (GC-IRMS). A GC-1310 gas chromatograph (Thermo Scientific, Bremen, Germany) equipped with an InertCap 5MS/NP (0.25 mm x 30 m x 0.25 μm) (GL Sciences, Japan) was interfaced to a Delta Advantage IRMS (Thermo Scientific) with a ConFlow IV interface (Thermo Scientific). Samples were injected with a TriPlusRSH autosampler to a PTV inlet operated in splitless mode at 280 °C. The oven was heated from 80 °C to 215 °C at 15 °C/min, then to 320 °C at 5 °C/min, and then was held at 320 °C for 10 minutes. Hydrogen isotope samples were pyrolyzed at 1420 °C after they eluted from the GC column. Carbon isotope samples were combusted at 1020 °C after elution.

25

Raw isotope values were converted to the VSMOW (hydrogen) and VPDB (carbon) scales using Thermo Isodat 3.0 software and pulses of a reference gas that was measured at the beginning and end of each analysis. Sample δ²H and δ¹³C values were further corrected using the slope and intercept of measured and known values of isotopic standards (*n*C_{17, 19, 21, 23, 25, 28, and 34}-alkanes, Arndt Schimmelmann, Indiana University), which were run at the beginning and end of each sequence, as well as after every 6 to 8 sample injections. Offsets between measured and known values for these standards were used to correct for any drift over the course of the sequence or any isotope effects associated with peak area or retention time. The standard

30

deviation for these standards averaged 4 ‰ and the average offset from their known values was 2 ‰ for hydrogen isotopes. For carbon isotopes, the average standard deviation of isotopic standards was 0.4 ‰ and the average offset from known values was 0.1 ‰ over the period of analysis.

- 5 An additional standard of nC_{29} -alkane was measured three times per sequence, corrected in the same way as the samples, and used for quality control. The standard deviation of these measurements was 4 ‰ for hydrogen and 0.5 ‰ for carbon over the period of analysis. The H_3^+ factor was measured at the beginning of each sequence and averaged 3.6 ± 0.3 during the analysis period. Samples were corrected for hydrogen and carbon added during derivatization using isotopic mass balance, and reported errors represent propagated errors from replicate measurements and the uncertainties associated with the added
10 hydrogen.

2.7 Calculated lipid production rates

Lipid production rates were calculated using Eq. (1) (modified from Popp et al. 2006):

$$15 \quad \text{Production rate} = (\delta^{13}C_l - \delta^{13}C_n) / (\delta^{13}C_{DIC} - \delta^{13}C_n) * (C_l/t) \quad (1)$$

- where $\delta^{13}C_l$ is the $\delta^{13}C$ value of the target compounds from labelled incubations, $\delta^{13}C_n$ is that from unlabelled incubations, $\delta^{13}C_{DIC}$ is the $\delta^{13}C$ value of DIC, C_l is the concentration of the lipid at the end of the incubation, and t is the duration of the incubation. Residence times – assuming a steady state, the amount of time needed to replace all molecules of a given lipid –
20 were calculated by dividing C_l by the production rate, which reduces to Eq. (2):

$$\text{Residence time} = t * (\delta^{13}C_{DIC} - \delta^{13}C_n) / (\delta^{13}C_l - \delta^{13}C_n) \quad (2)$$

2.8 Statistics

- 25 PRISM software (Graphpad Software Inc., La Jolla, CA, USA) was used to carry out all statistical analyses. Ordinary least squares regression was used to determine relationships among fractionation factors, temperature, and lipid production rates. Regression lines are only shown where the slope of the regression was significantly different from 0 at the $p < 0.05$ level. The results of all linear regression analyses are presented in Table 2. Differences between the slopes of various regressions were assessed using a two-tailed test of the null hypothesis that both slopes are equal. Differences in the mean values of
30 replicate measurements were determined using an unpaired, two-tailed t-test, and were considered significantly different for $p < 0.05$.

3. Results

3.1 Lipid concentrations and production rates

Lipid concentrations increased significantly in Lake Greifen from April to July, and then declined slightly from July to September, except for phytol and $nC_{16:1}$ fatty acid, which had increasing concentrations into the late summer (Fig. 2a). $nC_{16:0}$ fatty acid had the highest concentrations, while those of $nC_{16:1}$ fatty acid were usually the lowest, except in April-May, when $nC_{14:0}$ was the least abundant fatty acid. Brassicasterol concentrations were 1-2 orders of magnitude smaller than those of fatty acids and phytol concentrations were intermediate (Fig. 2a). Lipid concentrations in Lake Lucerne were generally an order of magnitude lower than in Lake Greifen (Fig. 2b). Fatty acid concentrations increased significantly from April to May in Lake Lucerne, and were then relatively stable throughout the rest of the time series (Figure 2b). Again, $nC_{16:0}$ fatty acid had the highest concentrations and $nC_{16:1}$ was the least abundant fatty acid. Phytol concentrations were typically an order of magnitude lower than those of fatty acids in Lake Lucerne, and increased slightly over the course of the time series. Brassicasterol concentrations were an order of magnitude lower still, and reached a maximum in June (Fig. 2b).

In both lakes, fatty acid production rates were highest for $nC_{16:0}$ (palmitic acid), followed by $nC_{18:x}$ (unsaturated C_{18} fatty acids, primarily $nC_{18:1n9c}$, or oleic acid), $nC_{14:0}$ (myristic acid), and $nC_{16:1}$ (palmitoleic acid) (Fig. 2c and 2d). Phytol and brassicasterol production rates were 2 - 3 orders of magnitude lower in both lakes than those of fatty acids (Fig. 2c and 2d). Lipid production rates were up to three times higher in Lake Greifen than in Lake Lucerne (Fig. 2c and 2d). Lipid production rates generally increased from May to July and then remained high in Lake Greifen, while in Lake Lucerne they were relatively constant throughout the study period (Fig. 2c and 2d).

Residence times – or the amount of time necessary to replace all molecules of a given compound assuming steady state – of individual lipids were calculated according to Eq. (2) (Sect. 2.7) and were typically shortest for $nC_{14:0}$, $nC_{16:0}$, and $nC_{18:x}$ fatty acids, with values as low as 9 ± 1 hr in Lake Greifen in May, and as low as 12 ± 3 hr in Lake Lucerne in August (Table 3). Of the fatty acids, $nC_{16:1}$ had the longest residence times, reaching 60 ± 13 hr in Lake Greifen in May and 62 ± 27 hr in Lake Lucerne in June (Table 3). Brassicasterol residence times were the longest of any lipid in the first part of the time series, but were exceeded by phytol for the last two sampling dates (Table 3).

3.2 Lipid δ^2H and $\alpha_{\text{lipid-water}}$ values

In both lakes lipid δ^2H values typically decreased over the spring and summer (Fig. 2e and 2f). This effect was most pronounced for fatty acids in Lake Greifen. For example, $nC_{16:0}$ fatty acid δ^2H values declined by 133 ‰ (from -172 ‰ to -

305 ‰) from April to August in Lake Greifen, while they only declined by 53 ‰ (from -249 ‰ to -302 ‰) over the same time period in Lake Lucerne (Fig. 2e and 2f). During the same time period water $\delta^2\text{H}$ values increased slightly in Lake Greifen (from -73 ‰ to -65 ‰) and were relatively constant in Lake Lucerne (fluctuating between -82 ‰ and -86 ‰) (S2). Changes in the fractionation factor between fatty acids and surface water ($\alpha_{\text{lipid-water}}$) were therefore linked closely to changes in fatty acid $\delta^2\text{H}$ values. In Lake Greifen, $\alpha_{\text{lipid-water}}$ for $n\text{C}_{16:0}$ -fatty acid decreased from 0.891 to 0.743 from April to August (Fig. 2g), while in Lake Lucerne, it decreased from 0.821 to 0.763 (Fig. 2h). Similar patterns were observed for $\alpha_{\text{lipid-water}}$ for $n\text{C}_{14:0}$, $n\text{C}_{16:1}$ and $n\text{C}_{18:X}$ fatty acids (Fig. 2g and 2h). Values for $\alpha_{\text{lipid-water}}$ were less variable for phytol and brassicasterol than for fatty acids, although they also declined from April to May. Brassicasterol was always ^2H -depleted relative to fatty acids in Lake Greifen, and was depleted in ^2H relative to all fatty acids except $n\text{C}_{14:0}$ in Lake Lucerne (Fig. 2g and 2h). Phytol $\delta^2\text{H}$ values were the most ^2H depleted of any lipid measured in either lake (Fig. 2g and 2h).

Overall, fatty acid $\alpha_{\text{lipid-water}}$ values were negatively correlated with lake surface temperature in both lakes ($R^2 = 0.32$, $p = 0.004$ in Lake Greifen; $R^2 = 0.24$, $p = 0.01$ in Lake Lucerne) (Fig. 3a; Table 2). The slope of the relationship was significantly steeper ($p = 0.03$) in Lake Greifen than in Lake Lucerne ($m = -0.006 \pm 0.002$ in Lake Greifen and -0.003 ± 0.001 in Lake Lucerne). Significant correlations were observed between lake surface temperature and $\alpha_{\text{lipid-water}}$ values for most fatty acids, but not for $n\text{C}_{16:1}$ in Lake Greifen and $n\text{C}_{14:0}$ in Lake Lucerne (Fig. 3; Table 2). Significant relationships between lake surface temperature and $\alpha_{\text{lipid-water}}$ values were not observed in either lake for brassicasterol or phytol (Fig. 3; Table 2).

Fatty acid production rates were not correlated with $\alpha_{\text{lipid-water}}$ values in either lake (Table 2). Among individual fatty acids, only $n\text{C}_{16:1}$ fatty acids from Lake Greifen had a significant negative correlation between $\alpha_{\text{lipid-water}}$ values and production rate ($R^2 = 0.84$; $p = 0.03$) (Table 2). Brassicasterol and phytol production rates were not correlated with $\alpha_{\text{lipid-water}}$ in either lake (Table 2), although brassicasterol $\alpha_{\text{lipid-water}}$ values from Lake Lucerne cluster as a significantly higher group than in Lake Greifen ($p = 0.0004$).

4. Discussion

In both lakes, the most striking feature of the lipid $\delta^2\text{H}$ values in the particulate organic matter (POM) is the significant decrease that occurs for most lipids during the spring (April – June) (Fig. 2e and 2f). As the lake water $\delta^2\text{H}$ values increased slightly (Greifen) or remained constant (Lucerne) throughout the summer, this trend indicates a decrease in $\alpha_{\text{lipid-water}}$ (Fig. 2g and 2h). There are a number of factors that could contribute to this decline in $\alpha_{\text{lipid-water}}$, but they generally group into two categories: changes in lipid source or changes in environmental variables, such as temperature, light, and productivity.

4.1 Potential changes in lipid source

Hydrogen isotope fractionation for short-chain fatty acids varies significantly among organisms with different metabolisms (Zhang et al., 2009a; Osburn et al., 2011; Heinzemann et al., 2015a). In general, fatty acids from heterotrophs grown on tricarboxylic acid (TCA)-cycle precursors are most enriched in ^2H , followed by heterotrophs grown on sugars, then photoautotrophs, and finally chemoautotrophs (Zhang et al., 2009a; Osburn et al., 2011; Heinzemann et al., 2015a). This variability can be greater than 500 ‰, and has led to the suggestion that the $\delta^2\text{H}$ values of ubiquitous compounds such as palmitic acid can be used as a proxy of net community metabolism. For example, at a coastal site in the North Sea, fatty acid chain-weighted average $\delta^2\text{H}$ values declined by more than 40 ‰ during the spring phytoplankton bloom, which was attributed to increased contributions from photoautotrophs (Heinzemann et al., 2016).

In Lakes Lucerne and Greifen, large decreases in fatty acid $\delta^2\text{H}$ values from April to May coincide with increases in fatty acid concentrations of 1-2 orders of magnitude (Fig. 2). It is therefore possible that the ^2H -enriched April samples represent a wintertime background of mixed heterotrophic and autotrophic derived compounds. As phytoplankton productivity ramped up with warmer temperatures, water column stratification, and longer daylight hours in the spring, newly produced fatty acids from photoautotrophs could have overwhelmed the heterotrophic signature, causing the net fatty acid $\delta^2\text{H}$ values to decrease. The increase in phytoplankton cell density (Fig. 4) from April onward in both lakes is supportive of increased contributions of fatty acids from phytoplankton as the study period progressed.

For Lake Greifen, a simple isotopic mass balance indicates that 31 % of the total $n\text{C}_{16:0}$ fatty acid would need to come from heterotrophs with $\alpha_{\text{lipid-water}}$ values of 1.200 (the maximum observed by Zhang et al., 2009a) in mid-April if the remaining $n\text{C}_{16:0}$ fatty acid was derived from phytoplankton with $\alpha_{\text{lipid-water}}$ values of 0.750 (assuming mid-summer $\alpha_{\text{lipid-water}}$ values represent the phytoplankton end member). Similar calculations suggest that 16% of $n\text{C}_{16:0}$ fatty acid would need to come from heterotrophic bacteria in mid-April in Lake Lucerne in order to account for the 50 ‰ decrease in $n\text{C}_{16:0}$ fatty acid $\delta^2\text{H}$ values over the course of the summer. These calculations assume that all heterotrophic bacteria use the highest $\alpha_{\text{lipid-water}}$ value ever observed for heterotrophs, and that they primarily use the TCA-cycle, rather than glycolysis. If, as is likely, there is diversity in $\alpha_{\text{lipid-water}}$ values for short-chain fatty acids produced by the heterotrophic bacteria, and at least some of the heterotrophs are relying primarily on glycolysis, the portion of these fatty acids from heterotrophic sources in April would need to be even higher than the values calculated above. This would necessitate a proportionally larger winter heterotrophic contribution of fatty acids than was observed in the coastal North Sea (Heinzemann et al., 2016), and it seems likely that other variables may contribute to the springtime decline in fatty acid $\delta^2\text{H}$ values.

Contributions from heterotrophs are also an improbable explanation for ^2H -enriched brassicasterol and phytol in April. Brassicasterol is a sterol that is commonly used as a biomarker for diatoms, although it has also been detected in some non-diatom eukaryotic phytoplankton (Volkman et al., 1998; Volkman, 2003; Rampen et al., 2010; Taipale et al., 2016) and occasionally in plant oils (Zarrouk et al., 2009). Since brassicasterol is not produced by bacterial sources, it seems improbable that the 25 ‰ (Greifen) and 19 ‰ (Lucerne) decreases in its $\delta^2\text{H}$ values from April to May could be due to proportionately greater heterotrophic contributions during the winter, as suggested for fatty acids. The 46 ‰ April – May decrease in $\delta^2\text{H}$ values of Lake Greifen phytol, the side-chain of chlorophyll molecules, is also unlikely to be caused by heterotrophic contributions in the early spring. Although phytol is produced by some photoheterotrophs, these typically have similar $\alpha_{\text{lipid-water}}$ values to photoautotrophs (Zhang et al., 2009a).

However, seasonal changes in the phytoplankton community composition alone could be a significant source of variability in $\alpha_{\text{lipid-water}}$ over the course of the study period. Hydrogen isotope fractionation for $n\text{C}_{16:0}$ fatty acid has been demonstrated to vary by over 100 ‰ among five different species of freshwater green algae grown in laboratory batch cultures (Zhang and Sachs, 2007). Such variations may be due to different enzymes involved in lipid synthesis among different species, or to the colony-forming behaviour of the two species with higher $\alpha_{\text{lipid-water}}$ values. Smaller species-dependent variations in $\alpha_{\text{lipid-water}}$ have been observed in cultures of haptophytes (~30 ‰ offset between alkenones in *G. oceanica* and in *E. huxleyi*; Schouten et al., 2006). Since there is limited data from culturing experiments, it is not possible to say how widespread such interspecies variability is. It is possible that most phytoplankton display similar magnitudes of hydrogen isotope fractionation during lipid synthesis under similar conditions. However, it is equally possible that lipid synthesis varies among species in ways that are not yet understood.

Given this uncertainty, and the significant changes in abundance of different phytoplankton taxa in Lake Greifen over the course of 2015 (Fig. 4a), contributions of short-chain fatty acids from different species of algae with different magnitudes of hydrogen isotope fractionation could account for some or all of the seasonal variability in $\alpha_{\text{lipid-water}}$. A comparable data set of algal species counts does not exist for Lake Lucerne from 2015, but bi-monthly data has been compiled from 2014 (Fig. 4b). Some changes in relative distributions of taxa are similar between the two lakes; for example, both Lake Greifen in 2015 and Lake Lucerne in 2014 experienced a peak in golden algae (*Chrysophyceae*) in June, and elevated abundance of cyanobacteria in late summer and late autumn (Fig. 4). Other trends differ starkly between the two lakes. Green algae (*Chlorophyceae*) are largely absent from Lake Lucerne, while they make up a significant portion of the algal community in Lake Greifen. Notably, the relative abundance of green algae steadily increased from May to September in Lake Greifen

(Fig. 4), during which time $\alpha_{\text{lipid-water}}$ values declined at a greater rate for most compounds than they did in Lake Lucerne (Fig. 2). If green algae tend to have lower $\alpha_{\text{lipid-water}}$ values than other algal taxa, their greater abundance in Lake Greifen throughout the summer could account for the greater decline in $\alpha_{\text{lipid-water}}$ over the course of the time series than in Lake Lucerne.

5

4.2 Relationships between seasonal environmental gradients and $\alpha_{\text{lipid-water}}$ values

Even for lipids produced in controlled cultures of eukaryotic algae, several factors have been shown to influence hydrogen isotope fractionation, including salinity, light availability, temperature, and growth rate (Summarized in Table 1) (Sachs, 2014 and sources therein; Chivall et al., 2014; M'boule et al., 2014; Heinzlmann et al., 2015b; Sachs and Kawka, 2015; van der Meer et al., 2015; Wolhowe et al., 2015; Maloney et al., 2016; Sachs et al., 2016; Sachs et al., 2017). Of these, salinity can be excluded as a source of variability freshwater lakes such as Greifen and Lucerne. The effect of light availability on $\alpha_{\text{lipid-water}}$ has only been detected at low light levels (below 250 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) (van der Meer et al., 2015; Wolhowe et al., 2015; Sachs et al., 2017). Although photosynthetically available radiation (PAR) was not measured as part of the present study, all samples were collected from lake surface water at a mid-latitude northern hemisphere site during boreal spring and summer, and it is unlikely that PAR was less than 250 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ at any sampling date (Pinker and Laszlo, 1992), meaning that the effect of light intensity is unlikely to be a source of the observed seasonal variability in $\alpha_{\text{lipid-water}}$ in lake surface water.

Lake Surface Temperature (LST) varied from 11 to 27 °C in Lake Greifen and from 11 to 25 °C in Lake Lucerne over the study period (Fig. 2i and 2j), and therefore may have contributed to the seasonal changes in lipid $\delta^2\text{H}$ values. In laboratory cultures of eukaryotic algae, $\alpha_{\text{lipid-water}}$ values for acetogenic lipids have been shown to decrease by 0.002 – 0.004 per °C, resulting in more depleted $\delta^2\text{H}$ values (Schouten et al., 2006; Zhang et al., 2009b; Wolhowe et al., 2009; Wolhowe et al., 2015). Increased hydrogen isotope fractionation at higher temperatures has been attributed to (i) changes in the relative activity of different enzymes involved in lipid synthesis at different temperatures, (ii) changes in the relative amount of NADPH from the pentose phosphate cycle as temperature changes, and (iii) the potential for hydrogen tunnelling at higher temperatures as substrate-enzyme complex vibrations increase (Sachs, 2014 and references therein). The relationship between temperature and hydrogen isotope fractionation in cultures is similar to that observed for fatty acids in Lake Lucerne, where $\alpha_{\text{lipid-water}}$ decreases by 0.003 ± 0.001 per °C (Fig. 3; Table 2). The relationship between $\alpha_{\text{lipid-water}}$ and temperature for fatty acids in Lake Greifen (-0.006 ± 0.002 per °C) (Fig. 3; Table 2) is much steeper than that observed in culturing studies.

30

If the influence of temperature on hydrogen isotope fractionation is consistent among laboratory cultures and lakes, warmer temperatures can account for the entire seasonal change in $\alpha_{\text{lipid-water}}$ for fatty acids in Lake Lucerne. However, increasing temperatures would only be able to explain part of the decrease in fatty acid $\delta^2\text{H}$ values in Lake Greifen over the course of the spring and summer. At most, temperature could account for half of the decrease in $\alpha_{\text{lipid-water}}$ in Lake Greifen, assuming a
5 consistent relationship to that observed in cultures.

Brassicasterol and phytol $\alpha_{\text{lipid-water}}$ values do not have strong relationships with lake surface temperature. There is no correlation between lake surface temperature and $\alpha_{\text{lipid-water}}$ values for phytol in either lake, nor for brassicasterol in Lake Lucerne. In Lake Greifen temperature and $\alpha_{\text{Brassicasterol-water}}$ are negatively correlated, although the relationship is not quite
10 significant (Fig. 3f; Table 2). The slope of the relationship between $\alpha_{\text{Brassicasterol-water}}$ and lake surface temperature in Lake Greifen is significantly shallower than that observed for fatty acids (-0.002 ± 0.001 for brassicasterol vs. -0.008 ± 0.002 for $n\text{C}_{16:0}$ fatty acid). Given these relatively weak relationships, it seems unlikely that temperature influences hydrogen isotope fractionation of either brassicasterol or phytol. The negative correlations between temperature and $\alpha_{\text{lipid-water}}$ values for fatty acids may therefore be an artefact of the probable increase in photoautotrophically-derived compounds as it became warmer
15 throughout the spring (Section 4.1). On the other hand, the relationship between temperature and $\alpha_{\text{lipid-water}}$ values in cultures has only been observed for *n*-alkanoic acids and alkenones, both of which are acetogenic lipids. It is thus possible that temperature is partially responsible for the decreases in $\alpha_{\text{lipid-water}}$ values as temperature increased in Lakes Lucerne and Greifen, but that the responsible mechanism is specific to lipids produced acetogenically, and does not affect isoprenoids, such as sterols and phytol.

20
Phytoplankton productivity and biomass also increased with temperature in the spring and summer in Lakes Greifen and Lucerne, with a more marked effect in nutrient-rich Greifen (Fig. 2a-d, Fig 4). This trend could also partially explain the increase in ^2H fractionation and decrease in lipid $\delta^2\text{H}$ values that co-occurred with rising temperatures. Increased nutrient availability and higher growth rates have been shown to result in lower $\alpha_{\text{lipid-water}}$ values during lipid synthesis for eukaryotic
25 algae in laboratory settings, with lipids more ^2H depleted relative to source water as growth rate increases (Schouten et al., 2006; Zhang et al., 2009b; Sachs and Kawka, 2015; Wolhowe et al., 2015). This relationship is most likely caused by increased contributions of hydrogen from relatively enriched NADPH from the oxidative pentose phosphate cycle under low-growth, nutrient stressed conditions, at the expense of relatively depleted hydrogen from photosystem I (Schmidt et al., 2003; Sachs and Kawka, 2015). If April samples include a higher proportion of lipids from organisms in a low-growth
30 maintenance phase, they should therefore be relatively enriched in ^2H . As light availability and water column stratification

became more amenable to photosynthesis later in the spring, relatively ^2H -depleted lipids produced with NADPH from photosystem I would be expected to become more abundant, bringing the net $\delta^2\text{H}$ and $\alpha_{\text{lipid-water}}$ values down.

5 Bottle incubations to determine lipid production rates were unfortunately not conducted for the first sampling in April, when the most enriched lipid $\delta^2\text{H}$ values were measured. For the remaining five sampling dates, there were not significant correlations between lipid production rate and $\alpha_{\text{lipid-water}}$, with the exception of $n\text{C}_{16:1}$ in Lake Greifen (Table 2). Lipid concentrations and production rates are not a direct proxy for growth rate, as a higher percentage of algal biomass is typically allocated to lipids under low nutrient and slow growth conditions (Rossler, 1990; Williams and Laurens, 2010). However, higher lipid production rates for the whole community (rather than on a per cell basis) will co-occur with higher growth rates. The large increase in fatty acid concentrations from April to May in both lakes, as well as smaller increases in brassicasterol concentrations, may indicate that the greatest community-wide change in growth rate occurred between those two months, and contributes in part to the decrease in fatty acid, phytol, and brassicasterol $\delta^2\text{H}$ values from April to May.

4.3 Comparison of mean $\alpha_{\text{lipid-water}}$ in lakes with different trophic statuses

15 For phytol, $n\text{C}_{14:0-}$, $n\text{C}_{16:0-}$, and $n\text{C}_{18:x-}$ fatty acids, there was no significant difference in $\alpha_{\text{lipid-water}}$ between the oligotrophic and eutrophic lake. However, significant differences in $\alpha_{\text{lipid-water}}$ do exist for brassicasterol (0.045 ± 0.008 ; $p = 0.0004$) and $n\text{C}_{16:1}$ -fatty acid (0.058 ± 0.022 ; $p = 0.02$) (Fig. 2). For brassicasterol, $\alpha_{\text{lipid-water}}$ is lower in Lake Greifen (0.712 ± 0.006) than in the less productive Lake Lucerne (0.757 ± 0.005). This result would be consistent with decreased hydrogen isotope fractionation (higher α values) for sterols in more nutrient-limited systems, as predicted by culturing experiments (Zhang et al. 2009b; Sachs & Kawka, 2015). The fact that this strong difference in fractionation between the two lakes is observed only for brassicasterol may be because it is the most source specific of the biomarkers that were analyzed. Phytol and short-chain fatty acids are produced by all photoautotrophs and may be dominated by phytoplankton who are optimized to grow under the nutrient regimes of each system, while relatively more of the brassicasterol may come from taxa who are nutrient-stressed and relying more on the pentose phosphate pathway than photosystem I.

25

Alternatively, the difference in $\alpha_{\text{Brassicasterol-water}}$ values between the two lakes could be due to variable contributions of brassicasterol from different phytoplankton sources. Even though brassicasterol is produced by fewer organisms than short-chain fatty acids and phytol, it still has multiple sources (Volkman et al., 1998; Volkman, 2003; Rampen et al., 2010; Taipale et al., 2016). Species-specific differences in hydrogen isotope fractionation have not been observed for sterols, but have been reported for fatty acids and alkenones (Schouten et al., 2006; Zhang & Sachs, 2007), making this an unconstrained possibility that could be responsible for the difference in $\alpha_{\text{Brassicasterol-water}}$ between the oligotrophic and eutrophic lake.

30

Different sources could also account for the difference in $\alpha_{\text{lipid-water}}$ for $n\text{C}_{16:1}$ -fatty acid, which displays higher $\alpha_{\text{lipid-water}}$ values in the more productive lake, and therefore cannot be explained by the nutrient effect observed in cultures.

5. Conclusions

- 5 We measured $\delta^2\text{H}$ values of short-chain fatty acids, phytol, and the diatom biomarker brassicasterol in surface water particulate organic matter in two lakes in central Switzerland with different trophic states at six time points throughout the spring and summer of 2015. Measurements were paired with in situ incubations with ^{13}C -enriched DIC that allowed us to calculate lipid production rates.
- 10 In April in both lakes, lipid concentrations were at their lowest and lipid $\delta^2\text{H}$ values were at their highest. In the case of short-chain fatty acids, which are produced by both photoautotrophic and heterotrophic microbes, the relatively high fractionation factors observed in the spring are consistent with a greater proportion of these compounds being derived from heterotrophs (Zhang et al., 2009a; Osburn et al., 2011; Heinzemann et al., 2015a). As phytoplankton productivity increased throughout the springtime, net $\alpha_{\text{lipid-water}}$ values declined to the range more commonly associated with photoautotrophs. The
- 15 observed decline in $\alpha_{\text{lipid-water}}$ for fatty acids in oligotrophic Lake Lucerne was similar to that observed during the spring bloom in the North Sea (Heinzemann et al., 2016), but was nearly three times as large in eutrophic Lake Greifen.

Changing contributions from heterotrophs cannot explain all of the decline in $\alpha_{\text{lipid-water}}$ from April to May, since this was also observed to a lesser extent in phytol and brassicasterol, compounds produced exclusively by photoautotrophs. Several factors

20 could be responsible for changes in photoautotrophic $\alpha_{\text{lipid-water}}$ throughout the spring, including temperature, growth rate, and species assemblage. Fractionation factors were inversely correlated with temperature for most fatty acids in each lake, and the slope of this relationship in Lake Lucerne (Fig. 3) was consistent with laboratory cultures, which suggest that $\alpha_{\text{lipid-water}}$ decreases with temperature by 0.002 – 0.004 per °C for acetogenic lipids (Zhang et al., 2009b; Wolhowe et al., 2009). Slower growth rates in the early spring could also result in higher $\alpha_{\text{lipid-water}}$ values at this time, as low growth rates correlate

25 with higher $\alpha_{\text{lipid-water}}$ values in cultures (Schouten et al., 2006; Zhang et al., 2009b; Sachs & Kawka, 2015). Finally, changes in phytoplankton species assemblage could have contributed to changes in $\alpha_{\text{lipid-water}}$ over time, as hydrogen isotope fractionation has been observed to vary among eukaryotic algal species grown in culture (Schouten et al., 2006; Zhang & Sachs, 2007; Heinzemann et al., 2015a)

While average fractionation factors for most lipids were consistent between the two lakes, average $\alpha_{\text{lipid-water}}$ values for brassicasterol were 0.045 ± 0.008 lower in Lake Greifen relative to Lake Lucerne, suggesting that sterol hydrogen isotopes may be more sensitive to nutrient availability than those of fatty acids and phytol.

5 Author contribution

S. N. Ladd designed the study with input from N. Dubois and C. J. Schubert. S. N. Ladd and N. Dubois collected the samples. S. N. Ladd processed and measured the samples. S. N. Ladd, N. Dubois, and C. J. Schubert contributed to data interpretation. S. N. Ladd prepared the manuscript with contributions from N. Dubois and C. J. Schubert. The authors declare that they have no conflict of interest.

10

Acknowledgements

This research was funded by a National Science Foundation Earth Sciences Postdoctoral Fellowship (Award #1452254) to NL and Eawag internal funds. Alois Zwyssig and Alfred Lück assisted with sample collection. Serge Robert and Julian Stauffer assisted with sample preparation and laboratory analyses. Daniel Montluçon at ETH-Zürich measured the water isotopes. Algal counts were conducted by Esther Keller as part of Eawag's Department of Aquatic Ecology's long-term monitoring program. We had productive conversations with Ashley Maloney, Daniel Nelson, Julian Sachs, Blake Matthews, and Romana Limberger that improved the study design and interpretation of results. Magdalena Osburn and Rienk Smittenberg provided helpful reviews that significantly strengthened the manuscript. We are grateful for all of their contributions.

20 References

- Atwood, A. R. and Sachs, J. P.: Separating ITCZ-and ENSO-related rainfall changes in the Galápagos over the last 3 kyr using D/H ratios of multiple lipid biomarkers, *Earth Planet. Sci. Lett.*, 404, 408-419, doi:10.1016/j.epsl.2014.07.038, 2014.
- Bührer, H. and Ambühl, H.: Lake Lucerne, Switzerland, a long term study of 1961-1992, *Aquat. Sci.*, 63, 432-456, doi:10.1007/s00027-001-8043-8, 2001.
- 25 Bürgi, H. R., Heller, C., Gaebel, S., Mookerji, N., and Ward, J. V.: Strength of coupling between phyto-and zooplankton in Lake Lucerne (Switzerland) during phosphorus abatement subsequent to a weak eutrophication, *J. Plankton Res.*, 21, 485-507, doi:10.1093/plankt/21.3.485, 1999.

- Chivall D., M'Boule D., Sinke-Schoen D., Damsté J. S. S., Schouten S., and van der Meer M. T. J.: The effects of growth phase and salinity on the hydrogen isotopic composition of alkenones produced by coastal haptophyte algae, *Geochim. Cosmochim. Acta* 140, 381–390, doi:10.1016/j.gca.2014.05.043, 2014.
- Craig H. and Gordon L.: Deuterium and oxygen 18 variations in the ocean and the marine atmosphere, in *Proceedings of a Conference on Stable Isotopes in Oceanographic Studies and Paleotemperatures* (ed E. Tongioli), CNR-Laboratorio di Geologia Nucleare, Pisa, Italy, 9-130, 1965.
- Englebrecht, A. C. and Sachs, J. P.: Determination of sediment provenance at drift sites using hydrogen isotopes and unsaturation ratios in alkenones. *Geochim. Cosmochim. Acta*, 69, 4253-4265, doi:10.1016/j.gca.2005.04.011, 2005.
- Gat, J. R., Oxygen and hydrogen isotopes in the hydrologic cycle. *Annu. Rev. Earth Planet. Sci.* 24, 225-262, doi:10.1146/annurev.earth.24.1.225, 1996.
- Heinzelmann, S. M., Villanueva, L., Sinke-Schoen, D., Damsté, J. S. S., Schouten, S., and van der Meer, M. T. J.: Impact of metabolism and growth phase on the hydrogen isotopic composition of microbial fatty acids, *Front. Microbio.*, 6, 408, doi:10.3389/fmicb.2015.00408, 2015a.
- Heinzelmann, S. M., Chivall, D., M'Boule, D., Sinke-Schoen, D., Villanueva, L., Damsté, J. S. S., Schouten, S., and Van der Meer, M. T. J.: Comparison of the effect of salinity on the D/H ratio of fatty acids of heterotrophic and photoautotrophic microorganisms, *FEMS microbiology letters*, 362, fmv065, doi:10.1093/femsle/fmv065, 2015b.
- Heinzelmann, S. M., Bale, N. J., Villanueva, L., Sinke-Schoen, D., Philippart, C. J. M., Damsté, J. S. S., Smede, J., Schouten, S., and van der Meer, M. T. J.: Seasonal changes in the D/H ratio of fatty acids of pelagic microorganisms in the coastal North Sea, *Biogeosci.*, 13, 5527-5539, doi:10.5194/bg-13-5527-2016, 2016.
- Henderson, A. K. and Shuman, B. N.: Hydrogen and oxygen isotopic compositions of lake water in the western United States, *Geol. Soc. Am. Bull.*, 121, 1179-1189, doi:10.1130/B26441.1, 2009.
- Hollander, D. J., McKenzie, J. A., and Ten Haven, H. L.: A 200 year sedimentary record of progressive eutrophication in Lake Greifen (Switzerland): implications for the origin of organic-carbon-rich sediments, *Geology*, 20, 825-828, doi:10.1130/0091-7613(1992)020<0825:AYSROP>2.3.CO;2, 1992.
- Huang, Y., Shuman, B., Wang, Y., and Webb III, T.: Hydrogen isotope ratios of palmitic acid in lacustrine sediments record late Quaternary climate variations. *Geology* 30, 1103-1106, doi:10.1130/0091-7613(2002)030<1103:HIROPA>2.0.CO;2, 2002.
- Huang, Y., Shuman, B., Wang, Y. and Webb, T.: Hydrogen isotope ratios of individual lipids in lake sediments as novel tracers of climatic and environmental change: a surface sediment test, *J. Paleolimnol.*, 31, 363-375, doi:10.1023/B:JOPL.0000021855.80535.13, 2004.

- Jones, A. A., Sessions, A. L., Campbell, B. J., Li, C., and Valentine, D. L.: D/H ratios of fatty acids from marine particulate organic matter in the California Borderland Basins, *Org. Geochem.*, 39, 485-500, doi:10.1016/j.orggeochem.2007.11.001, 2008.
- Kasper S., van der Meer M. T. J., Metz A., Zahn R., Damsté J. S. S., and Schouten S.: Salinity changes in the Agulhas leakage area recorded by stable hydrogen isotopes of C37 alkenones during Termination I and II. *Clim. Past.* 10, 251–260, doi:10.5194/cp-10-251-2014, 2014.
- Keller, B., Wolinska, J., Manca, M., and Spaak, P.: Spatial, environmental and anthropogenic effects on the taxon composition of hybridizing *Daphnia*, *Philos. T. Roy. Soc. B.*, 363, 2943-2952, doi:10.1098/rstb.2008.0044, 2008.
- Leduc, G., Sachs, J. P., Kawka, O. E., and Schneider, R. R.: Holocene changes in eastern equatorial Atlantic salinity as estimated by water isotopologues. *Earth Planet. Sci. Lett.*, 362, 151-162, doi:10.1016/j.epsl.2012.12.003, 2013.
- Li, C., Sessions, A. L., Kinnaman, F. S., and Valentine, D. L.: Hydrogen-isotopic variability in lipids from Santa Barbara Basin sediments, *Geochim. Cosmochim. Acta*, 73, 4803-4823, doi:10.1016/j.gca.2009.05.056, 2009.
- Maloney, A. E., Shinneman, A. L., Hemeon, K., and Sachs, J. P.: Exploring lipid $2\text{H}/1\text{H}$ fractionation mechanisms in response to salinity with continuous cultures of the diatom *Thalassiosira pseudonana*, *Org. Geochem.*, 101, 154-165. doi:10.1016/j.orggeochem.2016.08.015, 2016.
- M'boule, D., Chivall, D., Sinke-Schoen, D., Sinninghe Damsté, J. S. S., Schouten, S., and van der Meer M. T. J.: Salinity dependent hydrogen isotope fractionation in alkenones produced by coastal and open ocean haptophyte algae, *Geochim. Cosmochim. Acta* 130, 126–135, doi:10.1016/j.gca.2014.01.029, 2014.
- McKenzie, J. A.: Carbon-13 cycle in Lake Greifen: a model for restricted ocean basins, in: *Nature and origin of Cretaceous carbon-rich facies* (eds. Schlanger, S. O. and Cita, M.), Academic Press, London, UK, 197-208, 1982.
- Nelson, D. B. and Sachs, J. P.: The influence of salinity on D/H fractionation in dinosterol and brassicasterol from globally distributed saline and hypersaline lakes, *Geochim. Cosmochim. Acta*, 133, 325–339, doi:10.1016/j.gca.2014.03.007, 2014.
- Nelson, D. B. and Sachs, J. P.: Galápagos hydroclimate of the Common Era from paired microalgal and mangrove biomarker $^2\text{H}/^1\text{H}$ values, *P. Natl. Acad. Sci. USA*, 113, 3476-3481, doi:10.1073/pnas.1516271113, 2016.
- Osburn, M. R., Sessions, A. L., Pepe-Ranney, C., and Spear, J. R.: Hydrogen-isotopic variability in fatty acids from Yellowstone National Park hot spring microbial communities, *Geochim. Cosmochim. Acta*, 75, 4830-4845, doi:10.1016/j.gca.2011.05.038, 2011.
- Osburn, M. R., Dawson, K. S., Fogel, M. L., and Sessions, A. L.: Fractionation of hydrogen isotopes by sulphate- and nitrate-reducing bacteria, *Front. Microbio.*, 7, 1166m doi:10.2289/fmicb.2016.01166, 2016.
- Pahnke, K., Sachs, J. P., Keigwin, L., Timmermann, A., and Xie, S.: Eastern tropical Pacific hydrologic changes during the past 27,000 years from D/H ratios in alkenones, *Paleoceanography* 22, PA4214, doi:10.1029/2007PA001468, 2007.

- Pinker, R. T. and Laszlo, I.: Modeling surface solar irradiance for satellite applications on a global scale, *J. Appl. Meteorol.*, 31, 194-211, doi:10.1175/1520-0450(1992)031<0194:MSSIFS>2.0.CO;2, 1992.
- Popp, B. N., Prahl, F.G., Wallsgrove R. J., and Tanimoto, J. K.: Seasonal patterns of alkenone production in the subtropical oligotrophic North Pacific, *Paleoceanography*, 21, PA1004, doi:10.1029/2005PA001165, 2006.
- 5 Rampen, S. W., Abbas, B. A., Schouten, S., and Damsté J. S. S.: A comprehensive study of sterols in marine diatoms (*Bacillariophyta*): implications for their use as tracers for diatom productivity. *Limnol. Oceanogr.* 55, 91–105, doi:10.4319/lo.2010.55.1.0091, 2010.
- Randlett, M. E., Bechtel, A., van der Meer, M. T., Peterse, F., Litt, T., Pickarski, N., Kwiecien, O., Stockhecke, M., Wehrli, B., and Schubert, C. J.: Biomarkers in Lake Van sediments reveal dry conditions in eastern Anatolia during 110.000–10.000
 10 years BP, *Geochem. Geophys. Geosy.*, 18, doi:10.1002/2016GC006621, 2017
- Richey, J. N. and Sachs, J. P.: Precipitation changes in the western tropical Pacific over the past millennium, *Geology*, 44, 671-674, doi:10.1130/G37822.1, 2016.
- Roessler, P. G.: Environmental control of glycerolipid metabolism in microalgae: commercial implications and future research directions, *J. Phycol.*, 26, 393-399, doi:10.1111/j.0022-3646.1990.00393.x, 1990.
- 15 Sachs, J. P.: Hydrogen isotope signatures in the lipids of phytoplankton, in: *Treatise on Geochemistry*, 2nd Edition, eds: Holland, H. D. and Turekian, K. K., Elsevier, Oxford, UK, Vol. 12, 79-94, doi:10.1016/B978-0-08-095975-7.01007-X, 2014.
- Sachs, J. P. and Kawka, O. E.: The influence of growth rate on $2\text{H}/1\text{H}$ fractionation in continuous cultures of the coccolithophorid *Emiliana huxleyi* and the diatom *Thalassiosira pseudonana*, *PloS one*, 10, e0141643,
 20 doi:10.1371/journal.pone.0141643, 2015.
- Sachs, J. P. and Schwab, V. F.: Hydrogen isotopes in dinosterol from the Chesapeake Bay estuary, *Geochim. Cosmochim. Acta*, 75, 444–459, doi:10.1016/j.gca.2010.10.013, 2011.
- Sachs, J. P., Sachse, D., Smittenberg, R. H., Zhang, Z., Battisti, D. S., and Golubic S.: Southward movement of the Pacific intertropical convergence zone AD 1400–1850, *Nat. Geosci.*, 2, 519–525, doi:10.1038/ngeo554, 2009.
- 25 Sachs, J. P., Maloney, A. E., Gregersen, J., and Paschall, C. Effect of salinity on $2\text{H}/1\text{H}$ fractionation in lipids from continuous cultures of the coccolithophorid *Emiliana huxleyi*, *Geochim. Cosmochim. Acta*, 189, 96-109, doi:10.1016/j.gca.2016.05/041, 2016.
- Sachs, J. P., Maloney, A. E., and Gregersen, J.: Effect of light on $2\text{H}/1\text{H}$ fractionation in lipids from continuous cultures of the diatom *Thalassiosira pseudonana*, *Geochim., Cosmochim., Acta*, 209, 204-215, doi:10.1016/j.gca.2017.04.008, 2017.
- 30 Sachse, D. and Sachs, J. P.: Inverse relationship between D/H fractionation in cyanobacterial lipids and salinity in Christmas Island saline ponds, *Geochim. Cosmochim. Acta*, 72, 793–806, doi:10.1016/j.gca.2007.11.022, 2008.

- Sachse, D., Billault, I., Bowen, G. J., Chikaraishi, Y., Dawson, T. E., Feakins, S. J., Freeman, K. H., Magill, C. R., McInerney, F. A., van der Meer, M. T. J., Polissar, P., Robins, R. J., Sachs, J. P., Schmidt, H. L., Sessions, A. L., White, J. W. C., West, J. B., and Kahmen, A.: Molecular paleohydrology: interpreting the hydrogen-isotopic composition of lipid biomarkers from photosynthesizing organisms. *Annu. Rev. Earth Planet. Sci.* 40, 212-249, doi:10.1146/anurev-earth-042711-1055353, 2012.
- Sauer, P. E., Eglinton, T. I., Hayes, J. M., Schimmelmann, A., and Sessions, A. L.: Compound-specific D/H ratios of lipid biomarkers from sediments as a proxy for environmental and climatic conditions, *Geochim. Cosmochim. Acta*, 65, 213–222, doi:10.1016/S0016/7037(00)00520-2, 2001.
- Schimmelmann, A., Sessions, A. L., and Mastalerz, M.: Hydrogen isotopic (D/H) composition of organic matter during diagenesis and thermal maturation. *Annu. Rev. Earth Planet. Sci.*, 34, 501-533, doi:10.1146/annurev.earth.34.031405.125011, 2006.
- Schmidt, H. L., Werner, R. A., and Eisenreich, W.: Systematics of ^2H patterns in natural compounds and its importance for the elucidation of biosynthetic pathways, *Phytochem. Rev.*, 2, 61-85, doi:10.1023/B:PHYT.0000004185.92648.ae, 2003.
- Schouten, S., Ossebaar, J., Schreiber, K., Kienhuis, M. V. M., Langer, G., Benthien, A., and Bijma, J.: The effect of temperature, salinity and growth rate on the stable hydrogen isotopic composition of long chain alkenones produced by *Emiliania huxleyi* and *Gephyrocapsa oceanica*, *Biogeosci.* 3, 113–119, doi:10.5194/bg-3-113-2006, 2006.
- Sessions, A. L., Sylva, S. P., Summons, R. E., and Hayes, J. M.: Isotopic exchange of carbon-bound hydrogen over geologic timescales, *Geochim. Cosmochim. Acta*, 68, 1545-1559, doi:10.1016/j.gca.2003.06.004, 2004.
- Smittenberg, R. H., Saenger, C., Dawson, M. N., and Sachs, J. P.: Compound-specific D/H ratios of the marine lakes of Palau as proxies for West Pacific Warm Pool hydrologic variability, *Quatern. Sci. Rev.* 30, 921–933, doi:10.1016/j.quascirev.2011.01.012, 2011.
- Steinman, B. A., Abbott, M. B., Nelson, D. B., Stansell, N. D., Finney, B. P., Bain, D. J., and Rosenmeier, M.F.: Isotopic and hydrologic responses of small, closed lakes to climate variability: Comparison of measured and modeled lake level and sediment core oxygen isotope records, *Geochim. Cosmochim. Acta*, 105, 455-471, doi:10.1016/j.gca.2012.11.027, 2013.
- Taipale, S. J., Hiltunen, M., Vuorio, K., and Petomaa, E.: Suitability of phytosterols alongside fatty acids as chemotaxonomic biomarkers for phytoplankton, *Front. Plant Sci.*, 7:212, doi: 10.3389/fpls.2016.00212, 2016.
- Thevenon, F., Adatte, T., Poté, J., and Spangenberg, J. E.: Recent human-induced trophic change in the large and deep perialpine Lake Lucerne (Switzerland) compared to historical geochemical variations, *Palaeogeogr., Palaeoclimatol., 363*, 37-47, doi:10.1016/j.palaeo.2012.08.010, 2012.

- van der Meer, M., Baas, M., Rijpstra, W., Marino, G., Rohling, E., Damsté, J. S. S., and Schouten S.: Hydrogen isotopic compositions of long-chain alkenones record freshwater flooding of the Eastern Mediterranean at the onset of sapropel deposition, *Earth Planet. Sci. Lett.*, 262, 594–600, doi:10.1016/j.epsl.2007.08.014, 2007.
- van der Meer, M., Sangiorgi, F., Baas, M., Brinkhuis, H., Damsté J. S. S., and Schouten, S.: Molecular isotopic and dinoflagellate evidence for Late Holocene freshening of the Black Sea, *Earth Planet. Sci. Lett.*, 267, 426-434, doi:10.1016/j.epsl.2007.12.001, 2008.
- van der Meer, M. T., Benthien, A., French, K. L., Epping, E., Zondervan, I., Reichart, G. J., Bijma, J., Damsté J. S. S., and Schouten, S.: Large effect of irradiance on hydrogen isotope fractionation of alkenones in *Emiliania huxleyi*, *Geochim. Cosmochim. Acta*, 160, 16-24, doi:10.1016/j.gca.2015.03.024, 2015.
- 10 Vasiliev, I., Reichart, G. J., and Krijgsman, W.: Impact of the Messinian Salinity Crisis on Black Sea hydrology—Insights from hydrogen isotopes analysis on biomarkers, *Earth Planet. Sci. Lett.*, 362, 272–282, doi:10.1016/j.epsl.2012.11.038, 2013.
- Vasiliev, I., Mezger, E. M., Lugli, S., Reichart, G. J., Manzi, V., and Roveri, M.: How dry was the Mediterranean during the Messinian salinity crisis?, *Palaeogeogr. Palaeoclimatol.*, 471, 120-133, doi:10.1016/j.palaeo.2017.01.032, 2017.
- 15 Volkman, J. K.: Sterols in microorganisms, *Appl. Microbiol. Biotechnol.*, 60, 495–506, doi:10.1007/s00253-002-1172-8, 2003.
- Volkman, J. K., Barrett, S. M., Blackburn, S. I., Mansour, M. P., Sikes, E. L., and Gelin F.: Microalgal biomarkers: a review of recent research developments. *Org. Geochem.*, 29, 1163–1179, doi:10.1016/S0146-6380(98)00062-X, 1998.
- Williams, P. J. L. B., and Laurens, L. M.: Microalgae as biodiesel & biomass feedstocks: review & analysis of the biochemistry, energetics & economics, *Energy & Environmental Science*, 3, 554-590, doi:10.1039/B924978H, 2010.
- 20 Wolhowe, M. D., Prahl, F. G., Probert, I., and Maldonado M.: Growth phase dependent hydrogen isotopic fractionation in alkenone-producing haptophytes, *Biogeosci.*, 8, 1681–1694, doi:10.5194/bg-6-1681-2009, 2009.
- Wolhowe, M. D., Prahl, F. G., Langer, G., Oviedo, A. M., and Ziveri, P.: Alkenone δD as an ecological indicator: A culture and field study of physiologically-controlled chemical and hydrogen-isotopic variation in C 37 alkenones, *Geochim. Cosmochim. Acta*, 162, 166-182, doi:10.1016/j.gca.2015.04.034, 2015.
- Zarrouk, W., Carrasco-Pancorbo, A., Zarrouk, M., Segura-Carretero, A., and Fernandez-Gutierrez, A.: Multi-component analysis (sterols, tocopherols and triterpenic dialcohols) of the unsaponifiable fraction of vegetable oils by liquid chromatography– atmospheric pressure chemical ionization–ion trap mass spectrometry, *Talanta* 80, 924–934, doi:10.1016/j.talanta.2009.08.022, 2009.
- 30 Zhang, Z. and Sachs, J. P.: Hydrogen isotope fractionation in freshwater algae: 1. Variations among lipids and species, *Org. Geochem*, 38, 582–608, doi:10.1016/j.orggeochem.2006.12.004, 2007.

Zhang, X., Gillespie, A., and Sessions, A.: Large D/H variations in bacterial lipids reflect central metabolic pathways, P. Natl. Acad. Sci. USA, 106, 12580-12586, doi:10.1073/pnas.0903030106, 2009a.

Zhang, Z., Sachs, J. P., and Marchetti, A.: Hydrogen isotope fractionation in freshwater and marine algae: II. Temperature and nitrogen limited growth rate effects, Org. Geochem, 40, 428-439, doi:10.1016/j.orggeochem.2008.11.002, 2009b.

5 Zhang, Z., Leduc, G., and Sachs, J. P.: El Niño evolution during the Holocene revealed by a biomarker rain gauge in the Galápagos Island, Earth Planet. Sci. Lett., 404, 420-434, doi:10.1016/j.epsl.2014.07.013, 2014.

10

15

20

25

30

Table 1 Summary of expected changes in $\alpha_{\text{Lipid-Water}}$ in response to different environmental variables, based on laboratory cultures and field studies in marine settings

Variable	Sign of correlation with $\alpha_{\text{Lipid-Water}}$	Magnitude	References
Temperature	Negative	2 – 4 ‰ per °C	Zhang et al., 2009b; Wolhowe et al., 2009
Growth Rate	Negative	~30 ‰ per division day ⁻¹	Schouten et al., 2006; Zhang et al., 2009b; Sachs and Kawka, 2015; Wolhowe et al., 2015
Nutrient Availability	Negative	~40 ‰ difference between nutrient limited and nutrient replete cultures	Zhang et al., 2009b; Wolhowe et al., 2015
Light Availability	Positive	Below ~250 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, ~0.2‰ per $\mu\text{mol photons m}^{-2} \text{s}^{-1}$	van der Meer et al., 2015; Wolhowe et al., 2015; Sachs et al., 2017
Salinity	Positive	0.5 – 3 ‰ per practical salinity unit (PSU)	Schouten et al. 2006; Sachse and Sachs, 2008; Sachs and Schwab, 2011; Chivall et al., 2014; M'boule et al., 2014; Nelson and Sachs, 2014; Heinzlmann et al., 2015b; Maloney et al., 2016; Sachs et al., 2016
Species Assemblage	Variable	Differences up to 160 ‰ observed for <i>n</i> C16:0 fatty acid among species growing under identical conditions	Schouten et al., 2006; Zhang & Sachs, 2007

5

10

Table 2 Summary of linear regression statistics; bolded relationships are significant at the $p < 0.05$ level.

	LAKE GREIFEN					LAKE LUCERNE				
α Lipid-Water vs. Temperature										
Lipid	Slope	y-intercept	R ²	p	n	Slope	y-intercept	R ²	p	n
All Fatty acids	-0.006 ± 0.002	0.95 ± 0.04	0.32	0.004	24	-0.003 ± 0.001	0.85 ± 0.02	0.24	0.015	24
<i>n</i> C14:0 Fatty acid	-0.004 ± 0.001	0.86 ± 0.03	0.75	0.03	6	-0.0012 ± 0.0009	0.80 ± 0.02	0.28	0.28	6
<i>n</i> C16:0 Fatty acid	-0.008 ± 0.002	0.98 ± 0.03	0.87	0.006	6	-0.003 ± 0.001	0.84 ± 0.02	0.66	0.049	6
<i>n</i> C16:1 Fatty acid	-0.005 ± 0.003	0.98 ± 0.07	0.35	0.22	6	-0.004 ± 0.001	0.89 ± 0.02	0.74	0.028	6
<i>n</i> C18:x Fatty acid	-0.006 ± 0.0003	0.97 ± 0.007	0.99	<0.0001	6	-0.004 ± 0.001	0.89 ± 0.02	0.70	0.037	6
Phytol	-0.001 ± 0.001	0.66 ± 0.03	0.07	0.61	6	0.002 ± 0.001	0.62 ± 0.03	0.47	0.20	5
Brassicasterol	-0.002 ± 0.001	0.75 ± 0.02	0.63	0.11	5	0 ± 0.001	0.76 ± 0.02	3*10 ⁻⁵	0.99	6
α Lipid-Water vs. Lipid Production Rate										
Lipid	Slope	y-intercept	R ²	p	n	Slope	y-intercept	R ²	p	n
All Fatty acids	-0.01 ± 0.01	0.83 ± 0.02	0.10	0.18	20	-0.04 ± 0.03	0.81 ± 0.01	0.09	0.19	20
<i>n</i> C14:0 Fatty acid	0.00 ± 0.02	0.76 ± 0.02	0.0002	0.98	5	-0.2 ± 0.2	0.82 ± 0.05	0.25	0.39	5
<i>n</i> C16:0 Fatty acid	0.01 ± 0.02	0.78 ± 0.05	0.02	0.83	5	-0.01 ± 0.04	0.83 ± 0.03	0.46	0.21	5
<i>n</i> C16:1 Fatty acid	-0.17 ± 0.04	0.89 ± 0.01	0.84	0.03	5	0.6 ± 0.6	0.76 ± 0.06	0.22	0.42	5
<i>n</i> C18:x Fatty acid	-0.01 ± 0.02	0.85 ± 0.04	0.08	0.64	5	-0.09 ± 0.05	0.86 ± 0.02	0.57	0.14	5
Phytol	0 ± 1	0.63 ± 0.03	0.03	0.77	5	0 ± 7	0.66 ± 0.02	0.001	0.95	5
Brassicasterol	2 ± 2	0.70 ± 0.01	0.34	0.42	4	5 ± 43	0.76 ± 0.03	4*10 ⁻³	0.92	5

5

10

Table 3: Mean residence times in hours of lipids in lake surface water, calculated according to equation 2

Date	nC14:0	nC16:0	nC16:1	nC18:x	Brassicasterol	Phytol
<i>Lake Greifen</i>						
May 11, 2015	30 ± 9	19 ± 4	60 ± 13	19 ± 3	59 ± 26	42 ± 10
June 5, 2015	9 ± 2	9 ± 1	21 ± 2	9 ± 1	58 ± 34	32 ± 8
July 2, 2015	19 ± 1	20 ± 1	49 ± 5	20 ± 3	165 ± 51	40 ± 6
Aug. 11, 2015	15 ± 3	17 ± 2	41 ± 5	16 ± 2	81 ± 34	123 ± 23
Sept. 8, 2015	10 ± 3	12 ± 2	31 ± 8	12 ± 2	76 ± 37	127 ± 26
<i>Lake Lucerne</i>						
May 13, 2015	16 ± 6	16 ± 5	27 ± 8	17 ± 2	124 ± 22	99 ± 19
June 3, 2015	20 ± 12	18 ± 6	62 ± 27	16 ± 4	208 ± 56	106 ± 17
July 7, 2015	19 ± 8	19 ± 5	34 ± 18	17 ± 5	114 ± 39	94 ± 31
July 31, 2015	22 ± 9	19 ± 4	38 ± 8	18 ± 4	164 ± 59	399 ± 36
Aug. 31, 2015	20 ± 4	22 ± 3	41 ± 6	12 ± 3	258 ± 61	248 ± 42

5

10

15

20

Figure Captions

Figure 1: Map of Switzerland with locations of Lake Greifen and Lake Lucerne indicated. Base map from d-maps (http://www.d-maps.com/carte.php?num_car=2648&lang=en).

5

Figure 2: Time series of lipid concentrations in $\mu\text{g/L}$ (panels a and b), lipid production rates in $\mu\text{g L}^{-1} \text{hr}^{-1}$ (panels c and d), lipid $\delta^2\text{H}$ values in ‰ relative to VSMOW (panels e and f), $\alpha_{\text{lipid-water}}$ values (panels g and h), and lake surface temperature (panels i and j) for Lake Greifen (left column) and Lake Lucerne (right column) during the spring and summer of 2015. Panels a-d are plotted on an exponential scale to accommodate the large range in lipid concentrations and production rates. Error bars represent 1 standard deviation of replicate measurements, and are propagated to include uncertainties from multiple sources in calculated production rates and $\alpha_{\text{lipid-water}}$ values. In cases where error bars are not visible, they are smaller than the marker size.

Figure 3: Relationships between $\alpha_{\text{lipid-water}}$ values and lake surface temperatures in Lakes Greifen (panel a) and Lucerne (panel b) throughout the spring and summer of 2015. Error bars are propagated 1σ uncertainty from replicate measurements of surface water and lipid $\delta^2\text{H}$ values. In cases where error bars are not visible, they are smaller than the marker size. Shading represents 95% confidence intervals of the linear regression. Statistics associated with each curve are summarized in Table 2, and plots of individual compounds are available in the supplemental file.


20

Figure 4: Cell counts (individuals per liter) for all algae and for most common taxa of algae in (a) Lake Greifen throughout 2015 and (b) Lake Lucerne in 2014. The scale of the y-axis differs between the two panels. Data from long-term monitoring program run by the department of Aquatic Ecology at Eawag.

25

30

Figure 1:



5

10

Figure 2:







Figure 3:



5

10

Figure 4:

