


## **Supplementary Info**

We analyzed how CO<sub>2</sub> emissions vary spatially along radial transects across a sample of volcanoes from around the world. We used published data from a variety of volcanoes, either in the form of individual point measurements or spatially simulated flux maps. Because magnitude of CO<sub>2</sub> emissions varies significantly between volcanoes, we took the log<sub>10</sub> of the flux or concentration values and then transformed these values into a relative scale from 0 to 1, where 1 was the maximum flux or concentration at a specific volcano. For some groups of transects that had similar magnitudes of CO<sub>2</sub> emissions they were all put on the same relative scale. All CO<sub>2</sub> data was then linearly detrended. The distance along the radial transect was also put into a relative scale from 0 to 1, where 0 was the beginning of the transect and 1 was the end of the transect. The volcanoes and publications we took data from are as follows: Arenal (Williams-Jones, 1997), Teide (Hernández et al., 1998), Furnas (Viveiros et al., 2010), Vesuvius (Frondini et al., 2004), Vulcano (Schwandner et al., 2004), Cerro Negro (Salazar et al., 2001), Etna (Allard et al., 1991), Irazú (our collected data), and Mammoth Mountain (Werner et al., 2014). The data plotted in the supplemental Figure 1 show that all volcanoes studied show significant variations in volcanic CO<sub>2</sub> flux across their flanks, likely due to varying permeability (substrate) and fracture-controlled advective transport of CO<sub>2</sub> feeding these emissions from depth. These variations allow to study volcanically enhanced levels of CO<sub>2</sub> emissions as proxies for future atmospheric conditions via two modes: a local, substrate- and altitude-independent mode on the order of 50-200 meters, and a broad-scale enhancement mode, covering much of the volcanic flanks (distances on the order of 5-20 km) at lower levels of enhancements.



**Fig. S1:** CO<sub>2</sub> flux varies significantly along radial transects from 11 volcanoes around the world.

Species	Latitude	Longitude	Max CO <sub>2</sub> flux (g m <sup>-2</sup> day <sup>-1</sup> )	Mean CO <sub>2</sub> flux (g m <sup>-2</sup> day <sup>-1</sup> )	CO <sub>2</sub> flux $1\sigma$	$\delta^{13}\text{C}$ (‰)	CCI $1\sigma$	Chlorophyll concentration (umol m <sup>-2</sup> )	Chlorophyll concentration $1\sigma$	Fv/Fm	Fv/Fm $1\sigma$	$g_s$	$g_s \ 1\sigma$	
A. acuminata	10° 0'14.42"N	83°46'7.31"W	7.1	6.1	0.7	-26.1	28.7	8.6	452.4	135.5	0.793	0.017	83.5	7.8
A. acuminata	10° 0'17.87"N	83°46'9.20"W	18.5	11.1	6.6	-26.58	32.3	2.7	485.6	41.1	0.812	0.013	153.5	9.2
A. acuminata	10° 0'18.52"N	83°46'7.11"W	6.5	4.1	2.5	-24.48	25.7	8.0	423.5	132.7	0.793	0.011		
A. acuminata	10° 0'17.69"N	83°46'8.72"W	18.5	11.1	6.6	-26.22	33.0	4.7	491.8	70.0	0.806	0.013	241.0	79.2
A. acuminata	9°59'52.79"N	83°46'13.90"W	8.8	8.0	0.8	-25.76	20.8	5.6	371.9	100.7	0.801	0.010	131.5	17.7
A. acuminata	9°59'52.45"N	83°46'14.53"W	13.0	10.6	2.3	-26.24	40.7	6.7	556.1	91.5	0.824	0.005	331.5	51.6
B. nitida	9°58'60.00"N	83°50'9.15"W	15.7	9.6	4.2		56.1	29.6	669.4	352.8	0.838	0.018		
B. nitida	9°59'16.85"N	83°50'26.77"W	18.1	16.8	1.8		11.9	0.1	259.6	2.5	0.805	0.009		
B. nitida	10° 0'41.38"N	83°50'4.43"W	43.4	37.5	8.4		44.0	1.9	582.0	24.8	0.832	0.004		
B. nitida	10° 0'14.38"N	83°46'6.65"W	18.7	13.2	5.1	-26.76	83.1	1.4	834.6	14.5	0.799	0.009		
B. nitida	10° 0'18.54"N	83°46'7.15"W	6.5	4.1	2.5	-28.12	35.2	12.3	510.9	178.5	0.823	0.009		
B. nitida	10° 0'15.89"N	83°46'10.65"W	31.5	25.0	5.9	-26.22	80.3	1.9	819.0	19.4	0.795	0.012	160.0	0.0
B. nitida	9°59'56.46"N	83°46'31.92"W	5.7	4.6	1.1	-27.5	86.4	15.9	852.7	156.6	0.840	0.012	171.3	48.9
B. nitida	9°59'57.83"N	83°46'33.99"W	19.7	15.7	9.9	-26.72	99.5	15.3	921.9	141.5	0.840	0.012	200.5	14.8
B. nitida	9°59'55.11"N	83°46'36.36"W	21.5	15.6	7.9	-27.46	65.7	15.1	731.8	167.8	0.830	0.009	199.5	38.9
O. xalapensis	9°58'60.00"N	83°50'9.15"W	15.7	9.6	4.2		25.7	2.1	424.0	34.2	0.804	0.007		
O. xalapensis	10° 0'11.84"N	83°49'39.52"W	9.3	9.3	4.7		31.9	0.9	482.0	14.1	0.888	0.038		
O. xalapensis	10° 0'20.32"N	83°46'7.94"W	8.8	6.2	3.7	-26.31	22.7	2.0	393.2	35.3	0.783	0.016	190.0	0.0
O. xalapensis	9°59'56.41"N	83°46'12.18"W	12.2	5.7	4.5	-26.58	29.3	12.2	458.5	191.2	0.784	0.024	232.5	99.7
O. xalapensis	9°59'51.80"N	83°46'14.70"W	24.2	10.6	11.8	-25.51	39.4	5.2	546.1	72.2	0.806	0.010	321.0	45.3
O. xalapensis	9°59'52.21"N	83°46'12.76"W	10.0	7.0	3.0	-26.03	59.6	15.9	692.3	185.1	0.807	0.001	267.0	97.6
O. xalapensis	9°59'50.79"N	83°46'12.73"W	5.5	4.0	1.7	-27.16	66.5	13.9	736.8	153.5	0.804	0.012	306.0	59.4
O. xalapensis	9°59'56.47"N	83°46'31.92"W	5.7	4.6	1.1	-26.46	67.0	14.5	740.1	160.4	0.801	0.013	330.5	20.5

Table S1: Data used to generate all plots.

Species	Latitude	Longitude	Altitude (m)	Pressure (mbars)	Humidity (%)	PAR ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ )	Air T (°C)	Leaf T (°C)	Wind (m/s)	Aspect (degrees)	Slope (degrees)	DBH (cm)
<i>A. acuminata</i>	10° 0'14.42"N	83°46'7.31"W	2638	734.4	64	198	15.8	16	0	270	5	24.9
<i>A. acuminata</i>	10° 0'17.87"N	83°46'9.20"W	2640		70.55	150	12.5	13			0	14.3
<i>A. acuminata</i>	10° 0'18.52"N	83°46'7.11"W	2636	734.6	77.1	110	13.7	14.1	0	215	60	23
<i>A. acuminata</i>	10° 0'17.69"N	83°46'8.72"W	2633	734.8	84.5	173	15.5	15.4	0	290	5	45.3
<i>A. acuminata</i>	9°59'52.79"N	83°46'13.90"W	2432		64.9	200	15	15.4		135	25	58
<i>A. acuminata</i>	9°59'52.45"N	83°46'14.53"W	2434		65.2	325	15.4	15.5		110	20	90.3
<i>B. nitida</i>	9°58'60.00"N	83°50'9.15"W	3016	700	86.5	363	10.8	11.3	0	140	45	22.7
<i>B. nitida</i>	9°59'16.85"N	83°50'26.77"W	2968			94	12	11.2		20	50	50.2
<i>B. nitida</i>	10° 0'41.38"N	83°50'4.43"W	2322	763.7	84.4	86	12.9	13.4	0.5	120	55	43.5
<i>B. nitida</i>	10° 0'14.38"N	83°46'6.65"W	2619	735.9	67.6	800	12.6	13.4	1.7	190	35	15.3
<i>B. nitida</i>	10° 0'18.54"N	83°46'7.15"W	2615	736.4	89.5	172	13.4	12.9	0.4	215	60	11.5
<i>B. nitida</i>	10° 0'15.89"N	83°46'10.65"W	2625		85.7	334	13.6	13.8	1	200	15	26
<i>B. nitida</i>	9°59'56.46"N	83°46'31.92"W	2515	745.8	49.3	128	13.7	14	0	250	25	190
<i>B. nitida</i>	9°59'57.83"N	83°46'33.99"W	2511	746.2	67.7	157	13.9	13.9	0.2	180	5	180
<i>B. nitida</i>	9°59'55.11"N	83°46'36.36"W	2514	745.9	78.2	98	12.8	12.9	0	125	10	150
<i>O. xalapensis</i>	9°58'60.00"N	83°50'9.15"W	3016	700	86.5	43	10.4	10	0	45	58	15
<i>O. xalapensis</i>	10° 0'11.84"N	83°49'39.52"W	2101	785.4	100	37	15.4	15.2	0	80	30	11
<i>O. xalapensis</i>	10° 0'20.32"N	83°46'7.94"W	2619	736.1	77.9	170	13.5		0	150	55	15.4
<i>O. xalapensis</i>	9°59'56.41"N	83°46'12.18"W	2437	753	64.7	29	12	11.8	0	190	15	20.2
<i>O. xalapensis</i>	9°59'51.80"N	83°46'14.70"W	2438		65.5	325	14.7	14.7		190	55	20.3
<i>O. xalapensis</i>	9°59'52.21"N	83°46'12.76"W	2439		65.7	388	15.5	14.9		310	40	25
<i>O. xalapensis</i>	9°59'50.79"N	83°46'12.73"W	2438	752.9	65.8	338	17.7	17.7	0	285	60	22.5
<i>O. xalapensis</i>	9°59'56.47"N	83°46'31.92"W	2515	745.8	49.3	553	13.9	17.4	0	250	25	27.2

Table S2: Supplementary info for data presented in Table S1.