



Bipolar carbon and hydrogen isotope constraints of the Holocene methane budget

Jonas Beck^{1,2}, Michael Bock^{1,2}, Jochen Schmitt^{1,2}, Barbara Seth^{1,2}, Thomas Blunier³, Hubertus Fischer^{1,2}

¹Climate and Environmental Physics, Physics Institute, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland

²Oeschger Centre for Climate Change Research, University of Bern, 3012 Bern, Switzerland

³Centre for Ice and Climate, Niels Bohr Institute, University of Copenhagen, Juliane Maries Vej 30, 2100 Copenhagen, Denmark

Correspondence to: Jonas Beck (beck@climate.unibe.ch)

Abstract. Atmospheric methane concentration shows a well-known decrease over the first half of the Holocene following the northern hemisphere summer insolation before it started to increase again to preindustrial values. There is a debate about what caused this change in the methane concentration trend, in particular, whether an early anthropogenic influence or natural emissions led to the reversal of the atmospheric CH₄ concentration. Here, we present new methane concentration and stable hydrogen and carbon isotope data measured on ice core samples from both Greenland and Antarctica over the Holocene. With the help of a two-box model and the full suite of CH₄ parameters, the new data allow us to quantify the total methane emissions in the northern and southern hemispheres separately as well as their isotopic signatures, while interpretation of isotopic records of only one hemisphere may lead to erroneous conclusions. For the first half of the Holocene our results indicate a decrease in northern and southern hemisphere CH₄ emissions by more than 30 Tg CH₄/yr in total accompanied by a drop in the northern carbon isotopic source signature of about -3‰. This cannot be explained by a change in the source mix alone, but requires shifts in the isotopic signature of the sources themselves caused by changes in the precursor material for the methane production. In the second half of the Holocene global CH₄ emissions increased by about 30 Tg CH₄/yr, while preindustrial isotopic emission signatures remained more or less constant. However, our results show that the increase of methane emissions starting in the mid-Holocene took place in the southern hemisphere, while northern hemisphere emissions started to increase only about 2000 years ago. Accordingly, natural emissions in the southern tropics appear to be the main cause of the CH₄ increase starting 5000 years ago in contradiction to an early anthropogenic influence on the global methane budget by East Asian land use changes.

1 Introduction

Atmospheric methane (CH₄) is a potent greenhouse gas and its concentrations are strongly coupled to the Earth's climate system. Due to the human influence on the Earth system, the CH₄ concentration ([CH₄]) in the atmosphere has increased by a factor of 2.5 (relative to the preindustrial level) over the last centuries and contributes significantly to the human-induced radiative forcing (Dlugokencky et al., 2005; Etheridge et al., 1998). Today both anthropogenic CH₄ sources (rice agriculture, livestock, fossil fuel production, anthropogenic biomass burning and landfills) and natural CH₄ sources (natural wetlands, wildfires, geologic emissions, wild animals (including termites) and marine CH₄ hydrates) contribute to the global CH₄ emissions (Kirschke et al., 2013). The main mechanism removing CH₄ from the atmosphere is the chemical reaction of CH₄ with OH radicals in the troposphere. Together with the other sink processes (such as stratospheric loss, reaction with Cl radicals in the marine boundary layer and soil uptake) the OH sink determines the recent atmospheric life time of CH₄, which is between 8.4 and 10.6 years (Naik et al., 2013).

Despite the fact that the total CH₄ emissions today are well known, there is still considerable debate about the individual contributions of (in particular small) CH₄ source and their variability. Moreover, a long-standing debate exists whether long-



term Holocene CH₄ trends are naturally caused by wetland emissions (Singarayer et al., 2011) or whether an anthropogenic influence started already 4-5000 years ago by early human land-use changes (Ruddiman, 2003) (see below). Accordingly, to better assess the human influence on the global methane cycle, robust estimates of the natural (pre-anthropogenic) CH₄ emissions and their climate-coupled variations are required. Only polar ice cores can provide this information as they represent a direct, albeit low-pass filtered archive of trapped air and, therefore, offer the possibility to investigate the atmospheric composition of the past. With such ice core measurements the millennial to centennial evolution of the atmospheric [CH₄] has been determined back to 800 thousand years before present (ka BP), where present refers to the year 1950, from the Antarctic EPICA Dome C (EDC) ice core showing glacial/interglacial variations in atmospheric [CH₄] by a factor of about 2 (Loulergue et al., 2008). More important for the assessment of the human influence on CH₄ levels are their variations during the Holocene, i.e. our current interglacial that started about 10,000 years ago. In this time period [CH₄] decreased by about 100 parts per billion (ppb) until about 5000 yr ago (Figure 1) after an early maximum during the preboreal period (about 11.5-10 ka BP) (Blunier et al., 1995; Brook et al., 1996; Flückiger et al., 2002; Schilt et al., 2010). CH₄ values started to slowly increase again in the mid-Holocene until the fast anthropogenic increase resulting from the industrialisation starting around 200 years ago (MacFarling Meure et al., 2006). High resolution CH₄ measurements on the West Antarctic Ice Sheet (WAIS) and Greenland GISP2 (Greenland Ice Sheet Project 2) ice cores (Mitchell et al., 2013) show also centennial CH₄ variations of 20-30 ppb over the last 2800 years superimposed on the long-term increase.

With the trend reversal in the mid-Holocene leading to a “bowl shape” (Sowers, 2010) of [CH₄], the Holocene differs from many previous interglacials, where the [CH₄] steadily dropped to glacial levels concurrently with the northern hemisphere summer insolation. A controversial hypothesis postulates that CH₄ emissions caused by early human land use were responsible for the CH₄ turnaround about 5 ka BP (Ruddiman, 2003; Ruddiman and Thomson, 2001). According to these authors, early farming activities, mainly rice agriculture in the eastern part of China, led to significant releases of CH₄ long before the industrial era (Ruddiman et al., 2008) and, thus, the authors propose that the “bowl” was shaped by human activities. Another explanation supported by a model-based study explains the late Holocene CH₄ rise by calling upon increased emissions from southern tropical wetlands due to an insolation-driven strengthening of the monsoon in the western Amazon (Singarayer et al., 2011). Accordingly, this model is able to produce a global [CH₄] rise in the Holocene driven by changes in the orbital forcing only. Note that due to the different orbital configuration during the last interglacial (MIS5.5), this model is also able to reproduce the steadily declining [CH₄] during MIS5.5, when strongly declining boreal CH₄ emissions overcompensate increasing southern tropical wetland emissions.

An important fact is that the two scenarios explain the Holocene CH₄ rise with additional emissions taking place in different geographic regions. By measuring the [CH₄] on ice cores from both polar regions (Greenland and Antarctica), the inter polar difference (IPD) can be calculated. This quantity can be used to draw conclusions about the hemispheric distribution of the CH₄ emissions (Brook et al., 2000; Chappellaz et al., 1997; Mitchell et al., 2013; Yang et al., 2017). Another tool to investigate the processes involved in the CH₄ cycle are measurements of the stable isotope signature in ice core CH₄. The CH₄ released by the various sources are associated with typical hydrogen and carbon isotopic signatures $\delta\text{D-CH}_4$ (D representing Deuterium or ²H) (Quay et al., 1999; Walter et al., 2008; Whiticar, 1993; Whiticar and Schaefer, 2007) and $\delta^{13}\text{C-CH}_4$ (Etioppe et al., 2008; Quay et al., 1999; Walter et al., 2008; Whiticar, 1993; Whiticar and Schaefer, 2007). In addition, the sink processes lead to a strong fractionation (Feilberg et al., 2005; Levine et al., 2011b; Snover and Quay, 2000; Whiticar and Schaefer, 2007), which leave atmospheric CH₄ being strongly enriched in heavy isotopes relative to the isotopic emission signatures. Changes of the relative contributions in sources lead to an alteration of the atmospheric CH₄ isotopic composition. CH₄ isotope measurements on ice cores, therefore, allow us to deduce the role the different processes related to the CH₄ cycle played in the past. By measuring both $\delta\text{D-CH}_4$ and $\delta^{13}\text{C-CH}_4$ from the Greenland GISP2 ice core over the Holocene, Sowers (2010) was able to



provide first evidence of possible changes in the CH₄ precursor material due to a shift in the C3/C4 plant ratio, but could not unambiguously answer the question about the Holocene CH₄ anomaly.

In this study we combined for the first time comprehensive information on the CH₄ IPD and measurements of the carbon and hydrogen isotopic signature of CH₄ by measuring all three parameters – the [CH₄], δD-CH₄ and δ¹³C-CH₄ – on ice samples from both polar regions with a high analytical precision and improved resolution. Furthermore, we use a two-box model approach, which allows us to deconvolve the atmospheric signal, to calculate the total emissions and their isotopic signatures in each hemisphere.

As mentioned above, changes in the sinks leave an imprint on the isotopic composition of CH₄ in the atmosphere as well. However, the CH₄ sinks are thought to scale proportionally with the source (not changing the CH₄ life time) and only play a minor role for the observed atmospheric [CH₄] variation since the last glacial maximum (Levine et al., 2011a; Martinerie et al., 1995). Therefore, the discussion of this study centres around CH₄ source processes, hence emissions of different CH₄ source categories.

2 Method

2.1 Measurements

The stable isotope data (δD-CH₄ and δ¹³C-CH₄) presented in this study were measured at the University of Bern on samples from the NGRIP (North GREENland Ice core Project) ice core from Greenland. Antarctic samples from the EDML (EPICA Dronning Maud Land) and the TALDICE (TALos Dome Ice CorE) ice cores were analysed for δD-CH₄ and δ¹³C-CH₄, respectively (Bock et al., 2017). Note that for both parameters the samples from Greenland and Antarctica have been measured during the same time interval with the same measurement system by the same operators and in randomised order. This procedure is crucial to avoid any systematic error (e.g. due to a long-term drift of the system) and ensures unbiased results in our source deconvolution.

The two isotopic parameters were measured at the University of Bern on two independent measurement systems designed for this specific purpose as described in detail in previous publications (Bock et al., 2010; Schmitt et al., 2014). They follow the same principal procedure: the air is (i) extracted from the ice sample by purging the melt water, (ii) water vapour, CO₂ and the major air compounds (N₂, O₂ and Ar) are removed from the gas sample, (iii) CH₄ is separated from other trace gases using gas chromatography (GC), (iv) CH₄ is pyrolysed to H₂ or combusted to CO₂ in the δD and δ¹³C systems, respectively, and (v) the isotopic composition of H₂ and CO₂ is measured using an isotope-ratio mass spectrometer (IRMS). For both systems effects of interfering masses, specifically Krypton, in the mass spectrometer are avoided by technical adaptations (Schmitt et al., 2013). In the case of δD-CH₄, the focussed H₂ peak is separated from other components using a second GC separation (post-pyrolysis trapping) (Bock et al., 2014). In the δ¹³C-CH₄ system, the CH₄-derived CO₂ is captured and delayed using a fused silica trap (at –196°C) after the combustion oven and thereby separated from Krypton (Schmitt et al., 2014). Note that many δD-CH₄ and δ¹³C-CH₄ values based on such a gas chromatography - mass spectrometry method reported in the literature are not corrected for this effect, leading to significant offsets especially for low CH₄ concentrations.

The δD-CH₄ values are determined relative to our primary air standard ‘Air Controlé’, a recent clean air ([CH₄] = 1971 ± 7 ppb), which has been cross-referenced to –93.6 ± 2.2 ‰ with respect to (wrt) Vienna Standard Mean Ocean Water (VSMOW) using ‘Alert 2002/11’, an air collected at Alert Station, Canada (Bock et al., 2014). The latter was referenced by Poss (2003) to be –82.15 ± 0.3 ‰ wrt VSMOW, which anchors our δD values to the isotope scale of the University of Heidelberg (Bock



et al., 2010, 2014). Note that so far no internationally uniform $\delta\text{D-CH}_4$ standardisation scale has been established leading to systematic interlaboratory offsets as described in Umezawa et al. (2017). The measurement precision (1σ) of 2.3 ‰ was derived by calculating the pooled standard deviation of multiple NGRIP replicate (vertically neighbouring samples) measurements. The reference gas for the $\delta^{13}\text{C-CH}_4$ measurements is ‘Boulder’, an ambient air diluted with CH_4 -free air ($[\text{CH}_4]$ = 1508.18 ± 0.17 ppb) calibrated by the National Oceanic and Atmospheric Administration (NOAA) to a $\delta^{13}\text{C-CH}_4$ value of -47.34 ± 0.02 ‰ with respect to Vienna Pee Dee Belemnite (VPDB) (Schmitt et al., 2014; Umezawa et al., 2017). The measurement precision (1σ) derived from replicate measurements is better than 0.15 ‰ for ice core samples.

2.2 Monte Carlo spline approximation of millennial CH_4 changes

To compare values from different ice core datasets (with different and variable temporal resolutions and measurement precisions) interpolation of the data to a common age scale is needed. Additionally, the records have to be low-pass filtered to the same (millennial) time scales that can be resolved in the isotope records. Therefore, in this study smoothing splines following Enting (1987) are used to calculate a continuous low-pass filtered record from the discretely measured values. The stiffness of the spline curve is characterised by its cutoff period. A Python (<http://www.python.org/>) routine was developed to calculate splines in a Monte Carlo manner, where the data points are varied within a normal distribution within their standard deviation defined by the individual uncertainty of the measured values. This procedure allows for the calculation of the average of the splines representing the best guess mean evolution of the data on the time scale resolved by the spline. Furthermore, using the standard deviation of all splines, the spread of the individual splines can be translated to an uncertainty of the mean spline. This spline calculation is used for various purposes in this study. The choice of the cutoff period has a large influence on the outcome of the spline and is, therefore, always documented wherever it is selected manually. For some applications (e.g. in the routines described in the sections 2.3.2 and 2.3.3), however, we apply an objective rule to define an appropriate cutoff period for a data set: wherever the cutoff period is not specified explicitly, it is set to eight times the median of the data resolution.

If the sampling rate in some parts of a dataset is much higher than in others, this part has a larger impact on the spline than other coarser resolved sections. For short periods of large signal change (e.g. the 8.2 ka event for CH_4) such an oversampling relative to other parts of the dataset may lead to significant artefacts in the calculated spline. Wherever this might be an issue, an additional step was introduced into the spline calculation routine, where the data were down-sampled in a randomised way within the Monte Carlo loop (i.e. for each individual spline) to end up with a sub set of a prescribed number of data points, where the chance of data points to get picked are inversely proportional to the local data density. The down-sampled data are still unequally distributed, but the local average data density is similar over the whole time period studied.

2.3 Data correction and compilation

For further analysis of our new data, we complemented our $[\text{CH}_4]$ records with data from other studies from the GRIP (Greenland Ice core Project), GISP2, TALDICE, EDML, WAIS (West Antarctic Ice Sheet) and EDC ice cores shown in Table 1. However, combining different datasets introduces various challenges. In the following paragraphs we describe how these difficulties have been addressed.

2.3.1 Time synchronisation

All ice cores have individually determined gas age scales, which allow us to assign an age to each sample. Since we compare values from different ice cores (and from both hemispheres) small deviations in the age calculations have significant consequences for the interpretation (e.g. artefacts in the deconvolution (see section 2.4)). In an attempt to synchronise the



different datasets, we linearly stretched and compressed the individual age scales between manually-chosen tie points in three steps:

1. Wherever possible, we tied the data sets of each of the cores to the same reference age scale. For CH₄ datasets that reach far enough to the present, the fast CH₄ increase of the last centuries was used to synchronise the time scales. As a reference we used [CH₄] data from Law Dome firn air measurements (MacFarling Meure et al., 2006) and an atmospheric [CH₄] history derived from NEEM firn air (Buizert et al., 2012) representing the CH₄ evolution in Antarctica and in Greenland, respectively. Both records cover the entire anthropogenic CH₄ increase and overlap with recent atmospheric measurements (CSIRO, 2017; Dlugokencky et al., 2017). For the early Holocene the layer-counted NGRIP δ¹⁸O-H₂O record was used as an absolute time marker. Where the CH₄ data show strong signals such as at the 8.2 ka event and the Younger Dryas-Holocene transition these were synchronised to the corresponding δ¹⁸O signal, assuming that there is no significant lag between the temperature changes recorded in δ¹⁸O and the response of CH₄ source to this change (Baumgartner et al., 2014).

2. With the use of splines of different cutoff periods and their time derivatives we were looking for typical features and signals in the [CH₄] evolution of the different data sets within the same hemisphere to define additional tie points and synchronise the data sets for each hemisphere. However, within both hemispheres the temporal alignments of all the distinguishable CH₄ features were already within the limitation of the data resolution. Therefore, no additional tie points could be defined at this stage.

3. In the third synchronisation step we aligned the data of both hemispheres after applying the offset corrections (see section 2.3.2). Additional time corrections (for the southern hemisphere data) could be defined by looking at the fine structure of the [CH₄] evolution (e.g. the mid-Holocene [CH₄] minimum (around 5400 a BP) or the local [CH₄] maximum before the early Holocene decrease (around 9850 a BP)). Over the late Holocene period Mitchell et al. (2013) provided already synchronised data for the GISP2 and the WAIS ice core based on decadal to centennial variability; we retained their time scale over this period.

Table 2 lists the original age scales of the different ice cores. The tie points (original age and age correction) used to achieve an optimal temporal synchronisation between the CH₄ datasets are shown in Table 3. Temporal shifts to data with ages younger than the first tie point and older than the last tie point were chosen to be identical to those of the first and last tie points, respectively. Please note that for the ice cores, where the primary age scales provide an age uncertainty, all but three tie point corrections (GRIP: 273 a; TALDICE: 127 a and 6042 a) lie within the 1σ-range of the uncertainty given for the individual age scales.

2.3.2 Offset correction

Another issue that can complicate compiling multiple datasets are concentration offsets. The CH₄ datasets we use to complement our new data were measured with different methods, in different labs, and at different times over the last decades. If we use the information about the relative evolution of the methane concentration documented by other measurement series, one has to be aware that the absolute level might show an offset to our data due to calibration and extraction issues. Therefore, we correct all datasets for such potential offsets. To do so, we calculated residuals defined as the difference between the spline values of the CH₄ data from other labs evaluated at the age of our data and the CH₄ mixing ratios of our data. A least-squares optimisation routine was used to find the offsets for the spline values of the different data sets. The offset corrections applied to the data are shown in Table 1. The GISP2 and WAIS datasets by Mitchell et al. (2013) are treated as a single dataset, as



they were measured in the same lab and during the same measurement campaign, and their difference (Greenland-Antarctica) represents the true IPD. In Figure 1 the synchronised and offset-corrected CH₄ time series as used in this study are shown.

2.3.3 Outlier detection

The GISP2 datasets from Brook et al. (1996) and Brook (2009) show an increased incidence of anomalously high [CH₄] values (relative to other data sets) in the mid-Holocene. As this ice is from the brittle ice zone (the zone where air bubbles and clathrates coexist in situ, and which is prone to bad core quality due to damage during the pressure and temperature relaxation process after the core was brought to the surface) this is probably due to infiltration of modern air into the ice samples in (micro) cracks opened in the ice after core retrieval (Gow et al., 1997; Neff, 2014). To avoid an artificial increase of the splined signal, which would result in a significant blow-up of the IPD, we identify and remove such outliers within the depth interval 650-1400 m of the GISP2 ice core. To do so we used an iterative elimination approach. In each step, first the two GISP2 datasets were corrected for offsets relative to each other (Brook et al. (1996) was shifted to fit with Brook (2009) by 36 ppb) and combined to one data set. Thereafter, for each GISP2 data point the difference between its value and the spline of all other GISP2 data in the brittle zone was calculated and divided by the 1 σ uncertainty of the Monte Carlo spline at its age. If the largest (positive) relative residuum exceeded the threshold value of 4, the corresponding data point was removed. This procedure was repeated until no outliers could be found any more. Table 4 shows the data points that were identified and eliminated through this method (also shown in Figure 1). In total 15 out of 95 data points (within this depth range) were removed, reflecting the bad core quality in this depth interval. Note that we make the assumption here that the [CH₄] evolution in the atmosphere is relatively smooth. As we focus only on millennial changes in the data here, this assumption does not affect our conclusions.

2.3.4 Isotope data corrections

All stable isotope data have been corrected for gravitational settling. This process takes place in the firm column and leads to an enrichment of the heavier isotopes at the depth where the air bubbles are finally closed off and the gas is trapped in the ice (Schwander, 1996). Using $\delta^{136}\text{Xe}$ data, which come as a side product during the $\delta^{13}\text{C}$ -CH₄ measurement of the Bern $\delta^{13}\text{C}$ system, and additional published $\delta^{15}\text{N}$ -N₂ data (Capron et al., 2013; Eggleston et al., 2016; Landais et al., 2006), we experimentally derived this gravitational enrichment and directly correct the CH₄ stable isotope data, as this gravitational enrichment is only dependent on the mass difference between the two isotopes considered. The correction for the TALDICE $\delta^{13}\text{C}$ values increases from the present to the early Holocene from 0.28 ‰ to 0.39 ‰. The NGRIP $\delta^{136}\text{Xe}$ data do not show a systematic signal. Thus, a constant value of 0.32 ‰ is subtracted from the NGRIP isotope values. Only very few $\delta^{15}\text{N}$ -N₂ data points in the early Holocene were available to correct the EDML δD -CH₄ record. Since in the case of δD -CH₄ the measurement error and the observed signal are much larger than the gravitational enrichment, it is adequate to assume that this correction is also constant over time without any consequences for the interpretation of the data. For the EDML δD -CH₄ record the correction of 0.46 ‰ was applied.

Due to the relatively constant climatic conditions and no rapid [CH₄] changes occurring over the Holocene, thermal diffusion (Severinghaus et al., 1998) and diffusive fractionation (Buizert et al., 2013) – other processes potentially leading to an artefact in the stable isotope signature – are not considered in this study.

2.4 Deconvolution model

Our deconvolution model is a tool to disentangle the different processes involved in the atmospheric methane cycle. More precisely, it allows us to obtain quantitative information about the emission processes that led to the local tropospheric



composition at the ice core sites in the past. In other similar studies different approaches of varying complexity and with different boundary conditions have been presented (Baumgartner et al., 2012; Fischer et al., 2008; Mitchell et al., 2013). For this study we use a two-box model where the boxes represent the northern and southern tropospheric hemispheres.

- 5 The following six deconvolution equations allow us to use the six measured parameters (the tropospheric $[\text{CH}_4]$ (c_x) and the two stable isotope ratios (iR_x) for each hemisphere to calculate the emissions (E_x) and their isotope ratios (${}^iR_{E_x}$) for both boxes analytically (with x and y representing north and south and i the mass numbers 2 or 13 for deuterium and ${}^{13}\text{C}$):

$$E_x = m^* \cdot \left(\frac{r_x \cdot c_x}{\tau_x} + \frac{c_x}{2 \cdot \theta} - \frac{c_y}{2 \cdot \theta} \right) \quad (1)$$

$${}^iR_{E_x} = \frac{m^*}{E_x} \cdot \left(\frac{r_x \cdot c_x}{\tau_x} \cdot {}^iR_x \cdot {}^i\alpha_x + \frac{c_x}{2 \cdot \theta} \cdot {}^iR_x - \frac{c_y}{2 \cdot \theta} \cdot {}^iR_y \right) \quad (2)$$

- The first term in equations Eq. (1) and Eq. (2) represents the loss due to the CH_4 sinks, and the other two terms represent the hemispheric mixing (out of and into the box, with θ being the mixing time between the hemispheres as explained in more detail below). The parameter $m^* = \frac{m_0}{c_0}$ translates atmospheric concentrations (ppb) to total atmospheric inventories (in Tg) using the mean atmospheric concentration value $c_0 = 1650 \text{ ppb}$ of the year 1987 and the corresponding global CH_4 burden $m_0 = 4800 \text{ Tg}$ (Steele et al., 1992). The transport between the boxes is quantified by the exchange time $\theta = 1.56 \text{ a}$, which is determined through a calibration using recent sulphur hexafluoride (SF_6) data (described in section 2.4.1).

- 15 Note that the sizes of the two well-mixed boxes are not equal; the sizes are defined by the global mean annual latitude of the inter-tropical convergence zone (ITCZ) $\varphi_{ITCZ} = 5^\circ\text{N}$ (Marshall et al., 2014), which acts as a barrier to the hemispheric exchange of air masses. The inequality of the box sizes is reflected in Eq. (1) and Eq. (2) by

$$r_x = \frac{V_x}{V_{atm}} = \frac{1}{2} (1 \mp \sin(\varphi_{ITCZ})), \quad (3)$$

- the ratio of the box volume relative to the total atmospheric volume. The global mean life time of CH_4 quantifies the total methane sink and is set to $\tau = 8.4 \text{ a}$ (as in the 4-box model used in Fischer et al. (2008) and in Bock et al. (2017)). According to the relative importance given in Kirschke et al. (2013) (average of bottom-up estimates of all three time intervals) the sink is proportionally divided into the four different sink processes (s_{OH} , s_{strat} , s_{soil} , s_{Cl}) shown in Table 5. While s_{OH} and s_{strat} are thought to be equally distributed north and south of the ITCZ and, therefore, are set equal in the two boxes, s_{soil} and s_{Cl} are partitioned according to the area ratios of land and open water, respectively, derived from an 0.5° -by- 0.5° resolution land-cover database (Channan et al., 2014). The life time of CH_4 in each box

$$\tau_x = \tau \cdot \frac{\sum s_{tot}}{\sum s_x} \cdot r_x, \quad (4)$$

- 25 is determined by the north-south distribution of the sinks and the relative box sizes.

- The different sink processes lead to strong fractionations in both isotopes (Feilberg et al., 2005; Houweling et al., 2000; Quay et al., 1999; Snover and Quay, 2000 and references therein) and yield the hemispheric isotope fractionation factors ${}^i\alpha_x = 1 + {}^i\varepsilon_x$, with the fractionations ${}^i\varepsilon_x$ shown in Table 5. As in our model with only two tropospheric boxes s_{strat} is applied to the total tropospheric CH_4 inventory instead of the isotopically more depleted stratospheric CH_4 , the literature values for the stratospheric sink fractionation have to be adjusted to ${}^2\varepsilon_{strat} = -170 \text{ ‰}$ and ${}^{13}\varepsilon_{strat} = -13.1 \text{ ‰}$. These values have been empirically derived to match the tropospheric isotopic signature of the forward 4-box model described by Fischer et al. (2008) which contains two stratospheric boxes. First, using the identical CH_4 emission setup in both models, the north-south air mass exchange of the 2-box model has been increased to equal the tropospheric $[\text{CH}_4]$ values with those of the 4-box



model (IPD). In a second step, the fractionations of the stratospheric sink ${}^i\varepsilon_{strat}$ in the 2-box model have been adjusted to match the tropospheric $\delta\text{D-CH}_4$ and $\delta^{13}\text{C-CH}_4$ values of the two models.

As the individual data points from ice core $[\text{CH}_4]$ and isotope measurements are subject to uncertainty in both time and values even after above homogenisation steps, we only use the millennial variations in CH_4 , $\delta\text{D-CH}_4$ and $\delta^{13}\text{C-CH}_4$ in our deconvolution model. To this end we calculated low-pass filtered versions of the measured data using the Monte Carlo spline approximations described in section 2.2, which allow us to quantify the mean and the uncertainty of the millennial variations. The uncertainty of the splines of the tropospheric values are transmitted to the emission values using Gaussian error propagation. Due to the mentioned limitations we refrain from analysing shorter-scale variability.

10

2.4.1 SF₆ calibration of the two box model

The same approach used to set up the CH_4 deconvolution described in the previous section (Eq. (1) and (2)) allows us to formulate the general equations for the change of the concentration of any trace gas species in the two hemisphere boxes:

$$\frac{dc_x}{dt} \cdot r_x \cdot m^* = E_x - S_x - \frac{m^*}{2 \cdot \theta} (c_x - c_y) \quad (1)$$

with $\frac{dc_x}{dt}$ representing the change rate of the hemispheric concentration, m^* the translation factor linking a concentration to an atmospheric inventory, E_x and S_x the emissions and the sinks in Box x , and r_x the ratio of the box volumes depending on the mean annual latitude of the ITCZ φ_{ITCZ} as described above, and x and y standing for north and south (or vice versa).

SF₆ is a strong greenhouse gas with a life time of more than 1000 years and is solely of anthropogenic origin mainly emitted in the northern hemisphere ($r_{E_n} = 95\%$, $r_{E_s} = 5\%$) (Geller et al., 1997, p.6; Kovács et al., 2017; Levin et al., 2010). NOAA/ESRL (2017) measurements taken at Alert Station (Canada), Summit (Greenland), Palmer Station (Antarctica) and South Pole (Antarctica) record the SF₆ evolution in both polar regions over the last decades. Since the target in this study is the long-time trend and not seasonal variations, the NOAA/ESRL data are smoothed using a smoothing spline with a cutoff period of 2 years. For every year in the recorded period (1996-2008 CE) the annual SF₆ increase ($\frac{dc_x}{dt}$) and the mean concentration (c_x and c_y) are calculated. The records of the global emissions (E_{tot}), the global atmospheric mean SF₆ concentration (c_{tot}) and the total atmospheric inventory (m_{tot}) published by Levin et al. (2010) are used to calculate the hemispheric emissions $E_x = r_{E_x} \cdot E_{tot}$ and the translation factor $m^* = \text{mean} \left(\frac{m_{tot}}{c_{tot}} \right) = 25.7 \frac{\text{Gg SF}_6}{\text{ppt}}$. With the approximation of a negligible sink term (since life time exceeds the time scale of interest by 2-3 orders of magnitude) Eq. (1) can be solved for the hemispheric exchange time:

20

25

$$\theta = \frac{m^*(c_x - c_y)}{2 \left(\frac{dc_x}{dt} \cdot r_x \cdot m^* - r_{E_x} \cdot E_{tot} \right)} \quad (2)$$

Using Eq. (2) the exchange time can be calculated for each year and hemisphere. With $\varphi_{ITCZ} = 5.0^\circ\text{N}$ we get the average value of $\theta = 1.56 \pm 0.17 \text{ a}$.

30

In reality the air is not perfectly mixed within each hemisphere, which means that our ice core measurements do not perfectly represent the hemispheric average. However, by using the high latitude SF₆ data for the model calibration, we inherently correct for the fact that a signal subject to the inter-hemispheric exchange has to be transported to the poles before it is captured in the ice. Accordingly, θ has to be interpreted as an inter-polar rather than an interhemispheric exchange time.

35



2.4.2 Deconvolution of artificial time series

The interpretation of the output of our inversion model is challenging and not always intuitive. To get a better understanding of the model output, some simplified experiments using artificial time series have been performed and are shown in this study for didactic reasons. In a first approach all six parameters are held constant over time. In the panels on the left in Figure 2 the tropospheric $[\text{CH}_4]$ (a), and the isotopic signatures $\delta\text{D}-\text{CH}_4$ (b) and $\delta^{13}\text{C}-\text{CH}_4$ (c) are shown for the northern box in red and for the southern box in blue. The constant values (solid lines) represent the levels of the splines of our measurements at 2 ka BP and serve as input to the deconvolution model. The panels on the right show the deconvolution results, i.e., the integrated emissions (d) and their mean isotopic signatures (e, f) that are emitted into the two hemisphere boxes according to our model. As the input parameters are constant, the emission signal calculated by the deconvolution to achieve the prescribed tropospheric values must be constant in all parameters as well. However, although the northern hemisphere concentrations are only 6 % larger than the southern ones, a large difference in the emission strengths ($E_n \approx 2.2 \cdot E_s$) is needed to maintain the IPD in the $[\text{CH}_4]$ compensating the air mass exchange between the atmospheres. Note also that due to the sink fractionation the isotopic signature of atmospheric CH_4 is strongly enriched in the heavy isotopes compared to the emissions (see Table 5).

For the second experiment (dashed lines in Figure 2), the IPD of the tropospheric isotope signatures ($\text{IPD}_{\delta\text{D}}$ and $\text{IPD}_{\delta^{13}\text{C}}$) have been changed. We kept the southern values constant, while we linearly changed the northern values so that the minimal and the maximal $\text{IPD}_{\delta\text{D}}$ and $\text{IPD}_{\delta^{13}\text{C}}$ values match the splined isotope data, respectively (dashed lines in (b) and (c)). This leads to changes in the calculated isotope signatures of the emissions, whereas the emission strengths are unaffected. As shown (dashed lines) in panels (e) and (f) the isotopic signature for both the northern and the southern hemisphere emissions need to be altered to yield the given atmospheric values (of constant $[\text{CH}_4]$ and isotopic signatures changing only in one hemisphere). To achieve the declining isotope signal in the northern troposphere, the isotope signatures of emissions from the northern hemisphere have to become isotopically more depleted in heavy isotopes (further on conveniently referred to as isotopically “lighter”, equivalent to lower values δ -values) over time. At the same time the emissions in the southern hemisphere compensate for the isotopically lighter CH_4 mixed down from the northern box and, therefore, become more enriched in the heavy isotope (isotopically “heavier”, equivalent to higher δ -values) over time. Since the southern source is much weaker than the northern source (more northern-sourced CH_4 is mixed from north to south than southern-sourced CH_4 is mixed from south to north), the change in the isotopic signatures of the emissions turns out to be even larger in the southern hemisphere.

The dotted line in panel (a) in Figure 2 shows a third scenario, where all tropospheric parameters are kept constant except for the $[\text{CH}_4]$ in the northern hemisphere. Similar to the former experiment, the $[\text{CH}_4]$ in the northern box is varied between values representing the minimal and the maximal measured IPD_{CH_4} . The dotted curves in the right panels (d-f) in Figure 2 indicate that this change has an influence on all six emission parameters of our model. The emission strength in the northern box increases to produce the rising $[\text{CH}_4]$. Declining emission strength in the southern hemisphere compensates for the increasing CH_4 north-to-south flux due to the rising IPD_{CH_4} . The CH_4 that enters a box by atmospheric mixing is already fractionated by the sinks and isotopically heavier than the newly emitted CH_4 . If the relative amount of CH_4 input by mixing is decreased for the northern (increased for the southern) box, the emission signatures become less (more) depleted in heavy isotopes to maintain the tropospheric CH_4 isotopic signatures.

Since the variations for both experiments were done based on a reasonable range (observed changes in IPDs), the span for the calculated values of the emission parameters also provides a range for the changes that are needed to alter the measured tropospheric IPDs in the Holocene. Note, however, that the variation scenarios (e.g. with a fully decoupling of atmospheric concentration and the isotopic composition of CH_4) do not represent changes we expect to observe in nature and are presented for didactic reasons only.



3 Results

All tropospheric data (this study complemented with published [CH₄] data from Blunier et al., (1995); Brook et al., (1996); Brook, (2009); Chappellaz et al., (1993, 1997); Flückiger et al., (2002); Mitchell et al., (2013); Schilt et al., (2010), see section 2.3, Figure 1 and Table 1) are shown in Figure 3 in the left-hand panels. The smoothing splines, which are all calculated with the same cutoff period of 3000 years, are shown including their 1 σ uncertainty bands. In panel (a) the composite of the time-synchronised and offset-corrected CH₄ data is shown over the Holocene. The diamond-shaped symbols represent the NGRIP and TALDICE data of this study measured during the $\delta^{13}\text{C-CH}_4$ measurement campaign. Overall, our data compilation after the above mentioned corrections supports previous trends: after high concentrations in the Preboreal (ca. 11.5-10 ka BP) the data show a decline of the atmospheric [CH₄] over the first half of the Holocene followed by a reversal of this trend and an accelerating increase from the mid- to the late Holocene. The preindustrial level is comparable to the [CH₄] during the Preboreal warm period. The splines emphasise this long-term shape nicely but, deliberately, the splines do not reproduce the fine structure in the data (see for example Mitchell et al. (2013)). The difference between the [CH₄] values in the northern and southern hemispheres (the IPD_{CH₄}) is also clearly visible in the data. The concentration measured on ice cores from Greenland (red) is on average 45 ppb higher than the values derived from Antarctic ice cores (blue), which agrees with the result of Chappellaz et al. (1997).

In panel (b) the measured $\delta\text{D-CH}_4$ values show a pronounced difference of -16.3‰ between northern and southern values, which is rather stable over the whole period of investigation. Limited by the coarser resolution of the EDML $\delta\text{D-CH}_4$ data, the smoothed curves and the error bands illustrate the probable evolution of the true tropospheric signals only on millennial timescales and longer. Compared to the large [CH₄] changes, the $\delta\text{D-CH}_4$ signal over the Holocene is relatively small and similar in the two datasets. The values become isotopically lighter in the first half of the Holocene and heavier again in the second half. Note that our NGRIP $\delta\text{D-CH}_4$ data differ strongly from the dataset from the GISP2 ice core published by Sowers (2010) (see Figure 4). First, there is a difference of the mean level between the two datasets of 10 to 15 ‰, which can be attributed to an inter-laboratory difference (Bock et al., 2014; Umezawa et al., 2017). Second, the GISP2 data suggested a shift of about 20 ‰ towards heavier values from 5 to 1 ka BP, which is not confirmed by our data. This disagreement, however, may be largely attributed to the much larger scattering of the GISP2 dataset, which does not allow to quantify an unambiguous trend in $\delta\text{D-CH}_4$.

In panel (c) the $\delta^{13}\text{C-CH}_4$ measured on ice from the NGRIP and TALDICE ice cores are shown. Both datasets show a gradual trend towards lighter values over the first half of the Holocene and almost constant values over the second half. The point of inflection is well-defined at about 5 ka BP. At about the same time the $\delta^{13}\text{C-CH}_4$ signals measured in the two hemispheres diverge from a difference of -0.35‰ to -0.63‰ , which is rather constant before and after the point of inflection. Our NGRIP $\delta^{13}\text{C-CH}_4$ record is well in line with the GISP2 data by Sowers (2010) (see Figure 4), which have been subsequently corrected for Krypton interference during the measurement (Schmitt et al., 2013).

Our deconvolution model calculates the emissions that led to the measured tropospheric signals shown in the panels on the left. In panel (d) the strength of the integrated emissions (in Tg CH₄ per year) in each of the two troposphere boxes are shown. Additionally, the total CH₄ global emission (sum of north and south) are shown in grey. Note that the uncertainty (1 σ) in emission fluxes is very small and much smaller than the long-term changes in the emission fluxes over the Holocene indicated by the spline average. This uncertainty reflects only the error in emissions fluxes for changes on millennial time scales that are resolved in our spline. As indicated in high-resolution data (for example Mitchell et al., 2013) the centennial variability of atmospheric CH₄ is high and the uncertainties of reconstructed emission fluxes on such shorter time scales would be significantly higher as well.



The following observations can be made from the deconvolution results: The CH₄ emissions into the northern box are more than two times higher than the emissions into the southern box. After a decrease of roughly 30 Tg CH₄/a in the early Holocene the northern emissions remain relatively constant between 8 and 2 ka BP before increasing by 15 Tg CH₄/a. In the southern hemisphere the emissions show a slight increase in the first 2,000 years of the record followed by a 22 Tg CH₄/a decrease until 4-5 ka BP after which emissions increase by 25 Tg CH₄/a in the second half of the Holocene.

The weighted averaged δD signature of all the integrated CH₄ emissions is shown in panel (e) for each hemisphere. With values of about -295 ‰ the northern emission δD signature is on average 30 ‰ lighter than the southern hemisphere emissions. The δD-E_x signal shows long-term variations around the Holocene mean that are in antiphase between the northern and southern hemisphere, however these variations in each hemisphere are not significantly different from their respective Holocene mean within their 2σ uncertainty. The global mean δD-E (grey line) remains essentially constant. Based on these two observations, we attribute the δD-E_x fluctuations in each hemisphere to the measurement uncertainty of the individual data points of the low-resolution δD-CH₄ EDML record which affect the spline reconstruction.

Our experiments with artificial time series (see section 2.4.2) show that a change in the isotopic signature in one hemisphere has a large influence of different sign on the emission isotope signal in the other hemisphere (Figure 2). Hence, the absence of a long-term trend in δD-E_x over the Holocene with the significant changes in the atmospheric [CH₄] is remarkable and a strong constraint on the CH₄ budget. However, the large uncertainty in δD-E_x does not allow us to make robust conclusions about millennial variations in the hydrogen isotopic signature of CH₄ emissions.

The δ¹³C-CH₄ values of the emissions shown in panel (f) indicate a significant shift of the northern emission to heavier values over the Holocene, which is especially pronounced in the time interval before 4 ka BP. In the south δ¹³C-E_s shows also a slight negative trend over the first half of the Holocene, which however, is still well within the 1σ deconvolution uncertainty. A stronger positive trend, however, occurs during the second half of the Holocene, which is still within the 2σ uncertainty of the reconstruction. The global mean of emissions reflects the coherent trends in the measured atmospheric values, with a strong negative trend of -1.8‰ over the first 5000 years of the Holocene.

3.1 Sensitivity studies

The size-ratio of the two boxes of the deconvolution model is defined by the annual mean latitude of the ITCZ φ_{ITCZ} . In fact, there is evidence from different proxies of a southward migration of the ITCZ during the Holocene (Haug et al., 2001; McGee et al., 2014; Zhao and Harrison, 2012). Since it is not possible to deduce robust numbers for the position and the movement of the global mean ITCZ from local studies, φ_{ITCZ} has been kept constant over the time at a reasonable Holocene value in our deconvolution (described in section 2.4). However, sensitivity runs have been performed to quantify the impact of this assumption on the conclusions drawn from our deconvolution results.

As described in section 2.4.1, SF₆ calibration has been used to determine the inter-hemispheric mixing time θ associated to different values of φ_{ITCZ} . For the first sensitivity study these two parameters have always been fitted in parallel to ensure that the different model configurations all fit the SF₆ constraint in an optimal way. In Figure 5 the splines of the Holocene data (left panels) and the corresponding emission values derived by the deconvolution calculations (right panels) are shown. In the sensitivity runs, the deconvolution equations have been solved using other (φ_{ITCZ} , ε)-couples with φ_{ITCZ} moving from 0.0°N to 10.0°N. In all six emission parameters, these variations only lead to small changes relative to the total signal variability. For the isotopic signatures of the emissions, all curves lay well within the 1σ error of the 5.0°N results.



A change of the mean annual position of the ITCZ has an influence on the box size distributions. This has a direct impact on the north-south partitioning of the individual CH₄ sinks and thus on all six deconvolution parameters. Therefore, another sensitivity run was carried out, where only φ_{ITCZ} was varied and θ kept constant at the best guess value. As shown in Figure 6, the sensitivity of the deconvolution to a change of φ_{ITCZ} is very low if θ is kept constant. Note that the southward migration of the ITCZ over the Holocene is thought to be smaller than 1° (McGee et al., 2014). The results of this second set of experiments where only one parameter was changed (i.e. the constraints of the SF₆ calibration were not fulfilled) also showed that on average variations in φ_{ITCZ} and θ contribute by a similar amount to the deviation from the 5°N solutions (see Figure 6).

10 4 Discussion

The main goal of this study is to profit from the improved CH₄ multi-parameter dataset using our inversion model in order to evaluate the different scenarios and to constrain the global CH₄ budget over the Holocene. To this end, we consider the early, mid- and late Holocene separately.

4.1 Early Holocene (11-8 ka BP)

15 The significant drop in the northern emission strength together with the slight shift towards isotopically lighter carbon emissions in the northern hemisphere from 11 to 8 ka BP is challenging to account for. One potential solution to account for the observed concentration shift would be a general weakening of the boreal CH₄ sources as a result of the decreasing northern summer insolation (Berger and Loutre, 1991). However, this should lead to a trend in both isotope emission signatures towards heavier values in the northern hemisphere, as in general the high-latitude CH₄ emissions from wetlands are isotopically light
20 in both isotopes (Walter et al., 2008; Whiticar and Schaefer, 2007). Accordingly, this conflicts with the stable isotope results from our deconvolution. For $\delta D-E_n$ the deconvolution might be affected by one single measurement point (EDML $\delta D-CH_4$) at 8.19 yr BP forcing the emission signature curves of the two hemispheres apart and indicating that $\delta D-E_n$ becomes lighter. Nevertheless, even if this single data point is removed, the deconvolution signal would be constant over this period but would not show a trend toward heavier values as expected from a decline in boreal emissions.

25

Based on ¹⁴C-dating Walter et al. (2007) estimated the CH₄ emissions from thermokarst lakes to decrease from 26 to 5 Tg CH₄/a in the period from 11 to 8 ka BP in line with the decreasing trend in northern hemisphere emissions in our deconvolution. To match the result of the isotope deconvolution, another isotopically heavy northern source (e.g. biomass burning) has to be reduced simultaneously to compensate for the lack of isotopically light thermokarst CH₄ emissions. However, charcoal records
30 from sediments do not indicate a decline of wild fires in the northern hemisphere but rather the opposite during the Holocene (Daniau et al., 2012).

Part of the answer of this puzzle may lie in the combined temporal evolution of the strengths and isotopic signature of the boreal wetlands. Due to land ice and permafrost retreat in the early Holocene, minerotrophic fens, characterised by relatively
35 high CH₄ emissions which are not as strongly depleted in ¹³C, turn into ombrotrophic bogs over time with more ¹³C-depleted but also much weaker CH₄ emissions (Ding et al., 2005; Hornibrook, 2013; Yu et al., 2013). This reduction of CH₄ emissions in the course of conversion of fens to bogs requires a very depleted source signature of the bogs to quantitatively explain our deconvolution results and may therefore not suffice to explain the $\delta^{13}C-E_n$ shift towards lighter values in the early Holocene. Furthermore, an enrichment trend in $\delta D-E_n$ (of about 5‰) would be expected if the δD -depleted high-latitude emissions
40 became relatively less important, while – if at all – we see a 20‰ depletion in $\delta D-E_n$ from 10 to 8 ka BP. Also the isotopic



signature of the water used in methanogenesis has a direct imprint on the hydrogen isotopic signature of the emitted CH₄ (Whiticar and Schaefer, 2007). Speleothem δ¹⁸O records confirm an isotopic shift towards lighter values in the meteoric water in the summer monsoon regions of the northern hemisphere from 11 to 8 ka BP (Wang et al., 2014) and an inverse relationship in the southern hemisphere, in line with our deconvolution. However, the speleothem records also suggest opposite trends over
5 the rest of the Holocene, which are not seen in our deconvolution results. In addition, isotopically light melt water from the retreating northern hemisphere ice sheets (for example in proglacial lakes) may contribute to a depletion in the hydrogen isotopic signature of boreal CH₄ source in the early Holocene.

4.2 Mid-Holocene (8-5.5 ka BP)

Our deconvolution results show that the decrease in atmospheric CH₄ concentrations over this interval are mainly caused by
10 declining southern hemisphere emissions. This decrease in the southern hemisphere CH₄ emission from 8 to 5.5 ka BP - during the time of the strongest increase of austral summer insolation (Berger and Loutre, 1991) - appears counterintuitive but is in line with the model results by Singarayer et al. (2011) explaining it with changes in tropical rainfall location. This is also the period when the ITCZ is thought to have started moving southward as a response to the change in the orbital forcing (Haug et al., 2001; McGee et al., 2014; Zhao and Harrison, 2012), which has an influence on the hemispheric CH₄ source distributions.
15 CH₄ emissions from regions that were attributed to the southern hemisphere box in the early Holocene may now contribute CH₄ to the northern hemisphere due to the shift of the ITCZ. Note that such a change not only has an influence on the spatial emission distribution, but also on other parameters, which are relevant for the deconvolution model, where the box sizes are kept constant over time. Our sensitivity studies (see section 3.1) show, however, that a movement of the ITCZ in the model has only a small effect on the deconvolution results.

20

As discussed above the δD signal of the CH₄ emissions is mostly constant in both hemispheres. This is expected if one hemisphere box gains emissions from tropical wetlands, characterised by stable isotope signatures close to the integrated source mix, at the cost of the other box. The δ¹³C shift of the emissions could be related to a shift in the ratio of C3 to C4 plants as suggested by Sowers (2010), while a further evolution of northern fens to bogs is incompatible with the relatively constant
25 emissions in the northern hemisphere. Under Holocene climate conditions, with high availability of humidity and CO₂, C3 plants successfully compete against the relatively ¹³C-depleted C4 plants and thereby alter the carbon isotopic signature of the precursor material of methanogenesis (Farquhar et al., 1989). This process was also suggested by Möller et al. (2013) to explain the close CO₂ correlation of the δ¹³C-CH₄ signal and the decoupling from the [CH₄] in the glacial period.

4.3 Late Holocene (5.5-1 ka BP)

30 The early anthropogenic influence hypothesis (Ruddiman and Thomson, 2001) calls upon increasing emissions from rice paddies mainly in Eastern Asia. In our box model such emissions would be located in the northern hemisphere box and, therefore, lead to an increase in northern emissions after 5 ka BP. However, the deconvolution shows no significant increase in E_n before 2 ka BP. In contrast at 5.5 ka BP, at the time when all six presented tropospheric records imply a change in the global CH₄ cycle, the deconvolution indicates an amplification of southern hemisphere CH₄ emissions. This clearly supports
35 the idea of a natural Holocene wetland CH₄ emission increase as suggested by Singarayer et al. (2011). In contrast, a southern anthropogenic source can be excluded since the southern hemisphere was barely populated in the mid-Holocene (Kaplan et al., 2011). The observed shift in δD-E_s and especially δ¹³C-E_s towards heavier values in this interval cannot be explained simply by increased tropical wetland emissions in the Amazon but requires also an enhancement of a southern hemisphere CH₄ source strongly enriched in ¹³C. Increasing emissions from tropical and subtropical wild fires in the southern hemisphere,
40 which are also documented in charcoal records (Daniau et al., 2012), could have become increasingly important and would readily explain the joint information derived from deconvolution results on southern hemisphere CH₄ emissions and its stable



isotopic signatures. Other isotopically heavy sources such as geological emissions and marine CH₄ hydrates are unlikely to dramatically gain importance only in one hemisphere and can, therefore, be ruled out as important players driving the observed CH₄ changes.

5 Conclusions

- 5 The presented records of [CH₄], δD-CH₄ and δ¹³C-CH₄ measured on samples from polar ice cores from both hemispheres provide valuable insights into the Holocene CH₄ cycle. A significant IPD for δ¹³C and δD, as already documented for [CH₄], exists over the entire period of investigation. While the [CH₄] data confirm the well-known Holocene evolution, the two stable isotope records provide additional insights. This shows the value of our multi isotope approach needed not to draw the wrong conclusions. A two-box model approach has been used to deconvolve the measured signals. The calculated hemispheric CH₄ emission signatures, characterised by different trends in the two hemispheres for δ¹³C and no statistically significant variation in δD, show a decoupling of the emission signatures of the two isotopes. Therefore, we attribute the observed changes of the atmospheric CH₄ isotopes to shifts in the isotopic source signatures of individual CH₄ sources rather than to changes in the global CH₄ source mix.
- 10
- 15 In the early Holocene (11-8 ka BP) we associate the decline of northern hemisphere emissions and the δ¹³C trend towards lower values with the natural evolution of high latitude wetlands from fens (high CH₄ emissions) to bogs (lower but strongly ¹³C depleted CH₄ emissions). An alternative explanation calling upon a significant decrease of CH₄ emissions from thermokarst lakes requires compensation by other processes to fulfil the δ¹³C constraint, e.g. a significant decline of wild fire activity in the northern hemisphere without such evidence. In the mid-Holocene (8-5.5 ka BP) the deconvolution shows a decline of the southern hemisphere CH₄ emissions and an ongoing shift of the δ¹³C emission signatures towards lighter values, which might be related to the southward migration of the ITCZ and the change of the C3-to-C4 plant ratio altering the carbon isotopic composition of the precursor material for CH₄ production. During the second half of the Holocene (5.5-1 ka BP), which shows a major increase in CH₄ emissions accompanied by a shift towards higher δ¹³C signatures in the southern hemisphere, our results favour the 'Holocene anomaly' to be caused by natural CH₄ emissions (both wetlands and wildfires) in the southern tropics rather than by early rice agriculture in east Asia.
- 20
- 25

The CH₄ stable isotope data provide a powerful constraint on the Holocene CH₄ system (e.g. to benchmark CH₄ emission models), however, the relatively large uncertainty ranges in the calculated emission parameters still limit our ability to draw more robust conclusions. The deconvolution would probably benefit from an increased temporal resolution in the southern hemisphere δD-CH₄ record. However, since the concentration data have large influence on all six deconvolution parameters, we see the largest potential to further constrain the Holocene CH₄ emissions by actually improving the [CH₄] records. Better IPD_{CH₄} data measured on the same system during the same measurement campaign in high temporal resolution, which ensures an accurate synchronisation (such as the data by Mitchell et al. (2013) for the last 2,800 years), would bring us a large step forward in further constraining the Holocene CH₄ cycle. This is especially true for the Greenland ice cores within the brittle ice zone (e.g. GISP2 records for depths between 650 and 1400 m).

30

35

Data availability. Data presented in this study are available at (URL Pangaea page)

Acknowledgments. The research leading to these results has received funding from the European Research Council (ERC) under the European Union's Seventh Framework Programme FP7/2007-2013 ERC Grant 226172 [ERC Advanced Grant Modern Approaches to Temperature Reconstructions in Polar Ice Cores (MATRICs)] and the Swiss National Science Foundation. This work is a contribution to the North-GRIP ice core project, which is directed and organised by the Department of Geophysics at the Niels Bohr Institute for Astronomy, Physics and Geophysics, University of Copenhagen. It

40



is supported by funding agencies in Denmark (SNF), Belgium (FNRS-CFB), France (IFRTP and NSU/CNRS), Germany (AWI), Iceland (RannIs), Japan (MEXT), Sweden (SPRS), Switzerland (SNF), and the United States (NSF). Furthermore, this work is a contribution to the European Project for Ice Coring in Antarctica (EPICA), a joint European Science Foundation/European Commission scientific program funded by the European Union and national contributions from

5 Belgium, Denmark, France, Germany, Italy, The Netherlands, Norway, Sweden, Switzerland, and the United Kingdom. The main logistic support was provided by Institut Polaire Français Paul-Emile Victor (IPEV) and PNRA (at Dome C) and AWI (at Dronning Maud Land). The Talos Dome Ice Core Project (TALDICE), a joint European program led by Italy, is funded by national contributions from Italy, France, Germany, Switzerland, and the United Kingdom. The main logistical support was provided by Programma Nazionale di Ricerche in Antartide (PNRA) at Talos Dome.

10 This work is EPICA publication no. xx and TALDICE publication no. yy.

6 References

- Baumgartner, M., Schilt, A., Eicher, O., Schmitt, J., Schwander, J., Spahni, R., Fischer, H. and Stocker, T. F.: High-resolution inter-polar difference of atmospheric methane around the Last Glacial Maximum, *Biogeosciences*, 9 [online] Available from: <http://www.diva-portal.org/smash/record.jsf?pid=diva2:810843> (Accessed 17 March 2017), 2012.
- 15 Baumgartner, M., Kindler, P., Eicher, O., Floch, G., Schilt, A., Schwander, J., Spahni, R., Capron, E. F. N., Chappellaz, J., Leuenberger, M., Fischer, H. and Stocker, T. F.: NGRIP CH₄ concentration from 120 to 10 kyr before present and its relation to a $\delta^{15}\text{N}$ temperature reconstruction from the same ice core, *Clim. Past*, 10(2), 903–920, doi:10.5194/cp-10-903-2014, 2014.
- Bazin, L., Landais, A., Lemieux-Dudon, B., Toyé Mahamadou Kele, H., Veres, D., Parrenin, F., Martinerie, P., Ritz, C., Capron, E., Lipenkov, V., Loutre, M.-F., Raynaud, D., Vinther, B., Svensson, A., Rasmussen, S. O., Severi, M., Blunier, T.,
- 20 Leuenberger, M., Fischer, H., Masson-Delmotte, V., Chappellaz, J. and Wolff, E.: An optimized multi-proxy, multi-site Antarctic ice and gas orbital chronology (AICC2012): 120–800 ka, *Clim. Past*, 9(4), 1715–1731, doi:10.5194/cp-9-1715-2013, 2013.
- Berger, A. and Loutre, M.-F.: Insolation values for the climate of the last 10 million years, *Quat. Sci. Rev.*, 10(4), 297–317, 1991.
- 25 Blunier, T., Chappellaz, J., Schwander, J., Stauffer, B. and Raynaud, D.: Variations in atmospheric methane concentration during the Holocene epoch, *Nature*, 374(6517), 46–49, doi:10.1038/374046a0, 1995.
- Bock, M., Schmitt, J., Behrens, M., Möller, L., Schneider, R., Sapart, C. and Fischer, H.: A gas chromatography/pyrolysis/isotope ratio mass spectrometry system for high-precision δD measurements of atmospheric methane extracted from ice cores, *Rapid Commun. Mass Spectrom.*, 24(5), 621–633, doi:10.1002/rcm.4429, 2010.
- 30 Bock, M., Schmitt, J., Beck, J., Schneider, R. and Fischer, H.: Improving accuracy and precision of ice core $\delta\text{D}(\text{CH}_4)$ analyses using methane pre-pyrolysis and hydrogen post-pyrolysis trapping and subsequent chromatographic separation, *Atmospheric Meas. Tech.*, 7(7), 1999–2012, doi:10.5194/amt-7-1999-2014, 2014.
- Bock, M., Schmitt, J., Beck, J., Seth, B., Chappellaz, J. and Fischer, H.: Glacial/interglacial wetland, biomass burning, and geologic methane emissions constrained by dual stable isotopic CH₄ ice core records, *Proc. Natl. Acad. Sci.*, 114(29), E5778–E5786, doi:10.1073/pnas.1613883114, 2017.
- 35 Brook, E. J.: Methane measurements from the GISP2 and Siple Dome ice cores, [online] Available from: <http://nsidc.org/data/nsidc-0440>, 2009.
- Brook, E. J., Sowers, T. and Orchardo, J.: Rapid variations in atmospheric methane concentration during the past 110,000 years, *Science*, 273(5278), 1087–1091, doi:10.1126/science.273.5278.1087, 1996.
- 40 Brook, E. J., Harder, S., Severinghaus, J., Steig, E. J. and Sucher, C. M.: On the origin and timing of rapid changes in atmospheric methane during the last glacial period, *Glob. Biogeochem. Cycles*, 14(2), 559–572, doi:10.1029/1999GB001182, 2000.
- Buizert, C., Martinerie, P., Petrenko, V. V., Severinghaus, J. P., Trudinger, C. M., Witrant, E., Rosen, J. L., Orsi, A. J., Rubino, M. and Etheridge, D. M.: Gas transport in firn: multiple-tracer characterisation and model intercomparison for NEEEM, Northern Greenland, *Atmospheric Chem. Phys.*, 12(9), 4259–4277, doi:10.5194/acp-12-4259-2012, 2012.
- 45 Buizert, C., Sowers, T. and Blunier, T.: Assessment of diffusive isotopic fractionation in polar firn, and application to ice core trace gas records, *Earth Planet. Sci. Lett.*, 361, 110–119, doi:10.1016/j.epsl.2012.11.039, 2013.



- Buizert, C., Cuffey, K. M., Severinghaus, J. P., Baggenstos, D., Fudge, T. J., Steig, E. J., Markle, B. R., Winstrup, M., Rhodes, R. H., Brook, E. J., Sowers, T. A., Clow, G. D., Cheng, H., Edwards, R. L., Sigl, M., McConnell, J. R. and Taylor, K. C.: The WAIS Divide deep ice core WD2014 chronology - Part 1: Methane synchronization (68-31 ka BP) and the gas age-ice age difference, *Clim. Past*, 11, 153–173, doi:10.5194/cp-11-153-2015, 2015.
- 5 Capron, E., Landais, A., Buiron, D., Cauquoin, A., Chappellaz, J., Debret, M., Jouzel, J., Leuenberger, M., Martinerie, P., Masson-Delmotte, V., Mulvaney, R., Parrenin, F. and Prié, F.: Glacial–interglacial dynamics of Antarctic firn columns: comparison between simulations and ice core air- $\delta^{15}\text{N}$ measurements, *Clim Past*, 9(3), 983–999, doi:10.5194/cp-9-983-2013, 2013.
- Channan, S., Collins, K. and Emanuel, W. R.: Global mosaics of the standard MODIS land cover type data, *Univ. Md. Pac. Northwest Natl. Lab. Coll. Park Md. USA*, 30, 2014.
- Chappellaz, J., Blunier, T., Raynaud, D., Barnola, J. M., Schwander, J. and Stauffer, B.: Synchronous changes in atmospheric CH_4 and Greenland climate between 40 and 8 kyr BP, *Nature*, 366(6454), 443, doi:10.1038/366443a0, 1993.
- Chappellaz, J., Blunier, T., Kints, S., Dällenbach, A., Barnola, J.-M., Schwander, J., Raynaud, D. and Stauffer, B.: Changes in the atmospheric CH_4 gradient between Greenland and Antarctica during the Holocene, *J. Geophys. Res. Atmospheres*, 15 102(D13), 15987–15997, doi:10.1029/97JD01017, 1997.
- CSIRO: Latest Cape Grim greenhouse gas data, [online] Available from: <https://www.csiro.au/en/Research/OandA/Areas/Assessing-our-climate/Latest-greenhouse-gas-data> (Accessed 27 November 2017), 2017.
- Daniau, A.-L., Bartlein, P. J., Harrison, S. P., Prentice, I. C., Brewer, S., Friedlingstein, P., Harrison-Prentice, T. I., Inoue, J., Izumi, K., Marlon, J. R., Mooney, S., Power, M. J., Stevenson, J., Tinner, W. and Andrić, M.: Predictability of biomass burning in response to climate changes, *Glob. Biogeochem. Cycles*, 26(4), doi:doi:10.1029/2011GB004249, 2012.
- Ding, W., Cai, Z. and Tsuruta, H.: Plant species effects on methane emissions from freshwater marshes, *Atmos. Environ.*, 39(18), 3199–3207, doi:10.1016/j.atmosenv.2005.02.022, 2005.
- 15 Dlugokencky, E. J., Myers, R. C., Lang, P. M., Masarie, K. A., Crotwell, A. M., Thoning, K. W., Hall, B. D., Elkins, J. W. and Steele, L. P.: Conversion of NOAA atmospheric dry air CH_4 mole fractions to a gravimetrically prepared standard scale, *J. Geophys. Res. Atmospheres*, 110(D18), D18306, doi:10.1029/2005JD006035, 2005.
- Dlugokencky, E. J., Lang, P. M., Crotwell, A. M., Mund, J. W., Crotwell, M. J. and Thoning, K. W.: Atmospheric methane dry air mole fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1983–2016, [online] Available from: ftp://aftp.cmdl.noaa.gov/data/trace_gases/ch4/flask/surface/README_surface_flask_ch4.html (Accessed 20 30 October 2017), 2017.
- Eggleston, S., Schmitt, J., Bereiter, B., Schneider, R. and Fischer, H.: Evolution of the stable carbon isotope composition of atmospheric CO_2 over the last glacial cycle, *Paleoceanography*, 31(3), 434–452, doi:10.1002/2015PA002874, 2016.
- Enting, I. G.: On the use of smoothing splines to filter CO_2 data, *J. Geophys. Res. Atmospheres*, 92(D9), 10977–10984, 1987.
- Etheridge, D. M., Steele, L. P., Francey, R. J. and Langenfelds, R. L.: Atmospheric methane between 1000 A.D. and present: Evidence of anthropogenic emissions and climatic variability, *J. Geophys. Res. Atmospheres*, 103(D13), 15979–15993, doi:10.1029/98JD00923, 1998.
- 35 Etiopé, G., Milkov, A. V. and Derbyshire, E.: Did geologic emissions of methane play any role in Quaternary climate change?, *Glob. Planet. Change*, 61(1), 79–88, doi:10.1016/j.gloplacha.2007.08.008, 2008.
- Farquhar, G. D., Ehleringer, J. R. and Hubick, K. T.: Carbon isotope discrimination and photosynthesis, *Annu. Rev. Plant Biol.*, 40(1), 503–537, 1989.
- 40 Feilberg, K. L., Griffith, D. W., Johnson, M. S. and Nielsen, C. J.: The ^{13}C and D kinetic isotope effects in the reaction of CH_4 with Cl , *Int. J. Chem. Kinet.*, 37(2), 110–118, doi:10.1002/kin.20058, 2005.
- Fischer, H., Behrens, M., Bock, M., Richter, U., Schmitt, J., Loulergue, L., Chappellaz, J., Spahni, R., Blunier, T., Leuenberger, M. and Stocker, T. F.: Changing boreal methane sources and constant biomass burning during the last termination, *Nature*, 452(7189), 864–867, doi:10.1038/nature06825, 2008.
- 45



- Flückiger, J., Monnin, E., Stauffer, B., Schwander, J., Stocker, T. F., Chappellaz, J., Raynaud, D. and Barnola, J.-M.: High-resolution Holocene N₂O ice core record and its relationship with CH₄ and CO₂, *Glob. Biogeochem. Cycles*, 16(1), 10–1–10–8, doi:10.1029/2001GB001417, 2002.
- 5 Geller, L. S., Elkins, J. W., Lobert, J. M., Clarke, A. D., Hurst, D. F., Butler, J. H. and Myers, R. C.: Tropospheric SF₆: Observed latitudinal distribution and trends, derived emissions and interhemispheric exchange time, *Geophys. Res. Lett.*, 24(6), 675–678, doi:10.1029/97GL00523, 1997.
- Gow, A. J., Meese, D. A., Alley, R. B., Fitzpatrick, J. J., Anandakrishnan, S., Woods, G. A. and Elder, B. C.: Physical and structural properties of the Greenland Ice Sheet Project 2 ice core: A review, *J. Geophys. Res.-Oceans*, 102(C12), 26559–26575, doi:10.1029/97JC00165, 1997.
- 10 Haug, G. H., Hughen, K. A., Sigman, D. M., Peterson, L. C. and Röhl, U.: Southward migration of the intertropical convergence zone through the Holocene, *Science*, 293(5533), 1304–1308, doi:10.1126/science.1059725, 2001.
- Hornibrook, E. R. C.: The stable carbon isotope composition of methane produced and emitted from northern peatlands, in *Carbon Cycling in Northern Peatlands*, edited by A. J. Baird, L. R. Belyea, X. Comas, A. S. Reeve, and L. D. Slater, pp. 187–203, American Geophysical Union, Washington D.C., 2013.
- 15 Houweling, S., Dentener, F. and Lelieveld, J.: Simulation of preindustrial atmospheric methane to constrain the global source strength of natural wetlands, *J. Geophys. Res. Atmospheres*, 105(D13), 17243–17255, doi:10.1029/2000JD900193, 2000.
- Kaplan, J. O., Krumhardt, K. M., Ellis, E. C., Ruddiman, W. F., Lemmen, C. and Goldewijk, K. K.: Holocene carbon emissions as a result of anthropogenic land cover change, *The Holocene*, 21(5), 775–791, doi:10.1177/0959683610386983, 2011.
- Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Fraser, P. J., Krumpel, P. B., Lamarque, J. F., Langenfelds, R. L., Le Quéré, C., Naik, V., O’Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R., Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., Van Der Werf, G. R., Voulgarakis, A., Van Weele, M., Weiss, R. F., Williams, J. E., and Zeng, G.: Three decades of global methane sources and sinks, *Nat. Geosci.*, 25(10), 813–823, doi:10.1038/ngeo1955, 2013.
- 20 Kovács, T., Feng, W., Totterdill, A., Plane, J. M. C., Dhomse, S., Gómez-Martín, J. C., Stiller, G. P., Haenel, F. J., Smith, C., Forster, P. M., García, R. R., Marsh, D. R. and Chipperfield, M. P.: Determination of the atmospheric lifetime and global warming potential of sulfur hexafluoride using a three-dimensional model, *Atmos. Chem. Phys.*, 17(2), 883–898, doi:10.5194/acp-17-883-2017, 2017.
- 30 Landais, A., Barnola, J. M., Kawamura, K., Caillon, N., Delmotte, M., Van Ommen, T., Dreyfus, G., Jouzel, J., Masson-Delmotte, V., Minster, B., Freitag, J., Leuenberger, M., Schwander, J., Huber, C., Etheridge, D. and Morgan, V.: Firn-air δ¹⁵N in modern polar sites and glacial-interglacial ice: a model-data mismatch during glacial periods in Antarctica?, *Quat. Sci. Rev.*, 25(1–2), 49–62, doi:10.1016/j.quascirev.2005.06.007, 2006.
- Levin, I., Naegler, T., Heinz, R., Osusko, D., Cuevas, E., Engel, A., Ilmberger, J., Langenfelds, R. L., Neining, B., Rohden, C. v., Steele, L. P., Weller, R., Worthy, D. E. and Zimov, S. A.: The global SF₆ source inferred from long-term high precision atmospheric measurements and its comparison with emission inventories, *Atmos Chem Phys*, 10(6), 2655–2662, doi:10.5194/acp-10-2655-2010, 2010.
- 35 Levine, J. G., Wolff, E. W., Jones, A. E., Sime, L. C., Valdes, P. J., Archibald, A. T., Carver, G. D., Warwick, N. J. and Pyle, J. A.: Reconciling the changes in atmospheric methane sources and sinks between the Last Glacial Maximum and the pre-industrial era, *Geophys. Res. Lett.*, 38(L23804), 1–6, doi:10.1029/2011GL049545, 2011a.
- Levine, J. G., Wolff, E. W., Jones, A. E. and Sime, L. C.: The role of atomic chlorine in glacial-interglacial changes in the carbon-13 content of atmospheric methane, *Geophys. Res. Lett.*, 38(L04801), 1–6, doi:10.1029/2010GL046122, 2011b.
- Loulergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola, J.-M., Raynaud, D., Stocker, T. F. and Chappellaz, J.: Orbital and millennial-scale features of atmospheric CH₄ over the past 800,000 years, *Nature*, 453(7193), 383–386, doi:10.1038/nature06950, 2008.
- 45 MacFarling Meure, C., Etheridge, D., Trudinger, C., Steele, P., Langenfelds, R., Van Ommen, T., Smith, A. and Elkins, J.: Law Dome CO₂, CH₄ and N₂O ice core records extended to 2000 years BP, *Geophys. Res. Lett.*, 33(14), doi:10.1029/2006GL026152, 2006.



- Marshall, J., Donohoe, A., Ferreira, D. and McGee, D.: The ocean's role in setting the mean position of the Inter-Tropical Convergence Zone, *Clim. Dyn.*, 42(7–8), 1967–1979, doi:10.1007/s00382-013-1767-z, 2014.
- Martinerie, P., Brasseur, G. P. and Granier, C.: The chemical composition of ancient atmospheres: A model study constrained by ice core data, *J. Geophys. Res.*, 100(D7), 14,291–14,304, doi:10.1029/95JD00826, 1995.
- 5 McGee, D., Donohoe, A., Marshall, J. and Ferreira, D.: Changes in ITCZ location and cross-equatorial heat transport at the Last Glacial Maximum, Heinrich Stadial 1, and the mid-Holocene, *Earth Planet. Sci. Lett.*, 390, 69–79, doi:10.1016/j.epsl.2013.12.043, 2014.
- Meese, D. A., Gow, A. J., Grootes, P., Stuiver, M., Mayewski, P. A., Zielinski, G. A., Ram, M., Taylor, K. C. and Waddington, E. D.: The accumulation record from the GISP2 Core as an indicator of climate change throughout the Holocene, *Science*, 266(5191), 1680–1682, doi:10.1126/science.266.5191.1680, 1994.
- 10 Mitchell, L., Brook, E., Lee, J. E., Buizert, C. and Sowers, T.: Constraints on the late Holocene anthropogenic contribution to the atmospheric methane budget, *Science*, 342(6161), 964–966, doi:10.1126/science.1238920, 2013.
- Möller, L., Sowers, T., Bock, M., Spahni, R., Behrens, M., Schmitt, J., Miller, H. and Fischer, H.: Independent variations of CH₄ emissions and isotopic composition over the past 160,000 years, *Nat. Geosci.*, 6(10), 885–890, doi:10.1038/NGEO1922, 2013.
- 15 Naik, V., Voulgarakis, A., Fiore, A. M., Horowitz, L. W., Lamarque, J.-F., Lin, M., Prather, M. J., Young, P. J., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R., Eyring, V., Faluvegi, G., Folberth, G. A., Josse, B., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Van Noije, T. P., Plummer, D. A., Righi, M., Rumbold, S. T., Skeie, R., Shindell, D. T., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., and Zeng, G.: Preindustrial to presentday changes in tropospheric hydroxyl radical and methane lifetime from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmospheric Chemistry and Physics*, 13, 5277–5298, doi:10.5194/acp-13-5277-2013, 2013.
- 20 Neff, P. D.: A review of the brittle ice zone in polar ice cores, *Ann. Glaciol.*, 55(68), 72–82, doi:10.3189/2014AoG68A023, 2014.
- NOAA/ESRL: Combined Sulfur hexafluoride data from the NOAA/ESRL Global Monitoring Division, [online] Available from: <https://www.esrl.noaa.gov/gmd/hats/combined/SF6.html> (Accessed 10 October 2017), 2017.
- 25 Poss, C.: Untersuchung der Variabilität des atmosphärischen Methanhaushalts hochpolarer Breiten anhand eines regionalen Trajektorienmodells und der Messung stabiler Isotope, Universität Heidelberg. [online] Available from: <http://www.ub.uni-heidelberg.de/archiv/3966> (Accessed 15 September 2017), 2003.
- Quay, P., Stutsman, J., Wilbur, D., Snover, A., Dlugokencky, E. and Brown, T.: The isotopic composition of atmospheric methane, *Glob. Biogeochem. Cycles*, 13(2), 445–461, doi:10.1029/1998GB900006, 1999.
- 30 Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Andersen, M. L., Johnsen, S. J., Larsen, L. B., Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E. and Ruth, U.: A new Greenland ice core chronology for the last glacial termination, *J. Geophys. Res.-Atmospheres* 111 D06102, (10.1029/2005JD006079), doi:10.1029/2005JD006079, 2006.
- 35 Rasmussen, S. O., Bigler, M., Blockley, S. P., Blunier, T., Buchardt, S. L., Clausen, H. B., Cvijanovic, I., Dahl-Jensen, D., Johnsen, S. J., Fischer, H., Gkinis, V., Guillevic, M., Hoek, W. Z., Lowe, J. J., Pedro, J. B., Popp, T., Seierstad, I. K., Steffensen, J. P., Svensson, A. M., Vallenga, P., Vinther, B. M., Walker, M. J. C., Wheatley, J. J. and Winstrup, M.: A stratigraphic framework for abrupt climatic changes during the Last Glacial period based on three synchronized Greenland ice-core records: refining and extending the INTIMATE event stratigraphy, *Quat. Sci. Rev.*, 106(Supplement C), 14–28, doi:10.1016/j.quascirev.2014.09.007, 2014.
- 40 Ruddiman, W. F.: The anthropogenic greenhouse era began thousands of years ago, *Clim. Change*, 61(3), 261–293, 2003.
- Ruddiman, W. F. and Thomson, J. S.: The case for human causes of increased atmospheric CH₄ over the last 5000 years, *Quat. Sci. Rev.*, 20(18), 1769–1777, doi:10.1016/S0277-3791(01)00067-1, 2001.
- Ruddiman, W. F., Guo, Z., Zhou, X., Wu, H. and Yu, Y.: Early rice farming and anomalous methane trends, *Quat. Sci. Rev.*, 27(13), 1291–1295, doi:10.1016/j.quascirev.2008.03.007, 2008.
- 45 Schilt, A., Baumgartner, M., Schwander, J., Buiron, D., Capron, E., Chappellaz, J., Louergue, L., Schüpbach, S., Spahni, R. and Fischer, H.: Atmospheric nitrous oxide during the last 140,000 years, *Earth Planet. Sci. Lett.*, 300(1), 33–43, doi:10.1016/j.epsl.2010.09.027, 2010.



- Schmitt, J., Seth, B., Bock, M., van der Veen, C., Möller, L., Sapart, C. J., Prokopiou, M., Sowers, T., Röckmann, T. and Fischer, H.: On the interference of Kr during carbon isotope analysis of methane using continuous-flow combustion–isotope ratio mass spectrometry, *Atmospheric Meas. Tech.*, 6, 1425–1445, doi:10.5194/amt-6-1425-2013, 2013.
- Schmitt, J., Seth, B., Bock, M. and Fischer, H.: Online technique for isotope and mixing ratios of CH₄, N₂O, Xe and mixing ratios of organic trace gases on a single ice core sample, *Atmos Meas Tech*, 7(8), 2645–2665, doi:10.5194/amt-7-2645-2014, 2014.
- Schwander, J.: Gas Diffusion in Firn, in *Chemical Exchange Between the Atmosphere and Polar Snow*, edited by E. W. Wolff and R. C. Bales, pp. 527–540, Springer Berlin Heidelberg, Berlin, Heidelberg., 1996.
- Seierstad, I. K., Abbott, P. M., Bigler, M., Blunier, T., Bourne, A. J., Brook, E., Buchardt, S. L., Buizert, C., Clausen, H. B., Cook, E., Dahl-Jensen, D., Davies, S. M., Guillevic, M., Johnsen, S. J., Pedersen, D. S., Popp, T. J., Rasmussen, S. O., Severinghaus, J. P., Svensson, A. and Vinther, B. M.: Consistently dated records from the Greenland GRIP, GISP2 and NGRIP ice cores for the past 104 ka reveal regional millennial-scale $\delta^{18}\text{O}$ gradients with possible Heinrich event imprint, *Quat. Sci. Rev.*, 106(Supplement C), 29–46, doi:10.1016/j.quascirev.2014.10.032, 2014.
- Severinghaus, J. P., Sowers, T., Brook, E. J., Alley, R. B. and Bender, M. L.: Timing of abrupt climate change at the end of the Younger Dryas interval from thermally fractionated gases in polar ice, *Nature*, 391(6663), 141–146, doi:10.1038/34346, 1998.
- Singarayer, J. S., Valdes, P. J., Friedlingstein, P., Nelson, S. and Beerling, D. J.: Late Holocene methane rise caused by orbitally controlled increase in tropical sources, *Nature*, 470(7332), 82–85, doi:10.1038/nature09739, 2011.
- Snover, A. K. and Quay, P. D.: Hydrogen and carbon kinetic isotope effects during soil uptake of atmospheric methane, *Glob. Biogeochem. Cycles*, 14(1), 25–39, doi:10.1029/1999GB900089, 2000.
- Sowers, T.: Atmospheric methane isotope records covering the Holocene period, *Quat. Sci. Rev.*, 29(1), 213–221, doi:10.1016/j.quascirev.2009.05.023, 2010.
- Steele, L. P., Dlugokencky, E. J., Lang, P. M., Tans, P. P., Martin, R. C. and Masarie, K. A.: Slowing down of the global accumulation of atmospheric methane during the 1980s, *Nature*, 358(6384), 313–316, doi:10.1038/358313a0, 1992.
- Umezawa, T., Brenninkmeijer, C. A., Röckmann, T., van der Veen, C., Tyler, S. C., Fujita, R., Morimoto, S., Aoki, S., Sowers, T. and Schmitt, J.: Intercomparisons of $\delta^{13}\text{C}$ and δ^{D} measurements of atmospheric CH₄ for combined use of datasets from different laboratories, *Atmospheric Meas. Tech. Discuss.*, 2017.
- Veres, D., Bazin, L., Landais, A., Kele, H. T. M., Lemieux-Dudon, B., Parrenin, F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., Chappellaz, J., Rasmussen, S., Severi, M., Svensson, A., Vinther, B. and Wolff, E. W.: The Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years, *Clim. Past*, 9(4), 1733–1748, doi:10.5194/cp-9-1733-2013, 2013.
- Vinther, B. M., Clausen, H. B., Johnsen, S. J., Rasmussen, S. O., Andersen, K. K., Buchardt, S. L., Dahl-Jensen, D., Seierstad, I. K., Siggaard-Andersen, M.-L., Steffensen, J. P., Svensson, A., Olsen, J. and Heinemeier, J.: A synchronized dating of three Greenland ice cores throughout the Holocene, *J. Geophys. Res. Atmospheres*, 111(D13), doi:10.1029/2005JD006921, 2006.
- Walter, K. M., Edwards, M. E., Grosse, G., Zimov, S. A. and Chapin, F. S.: Thermokarst lakes as a source of atmospheric CH₄ during the last deglaciation, *science*, 318(5850), 633–636, doi:10.1126/science.1142924, 2007.
- Walter, K. M., Chanton, J. P., Chapin, F. S., Schuur, E. A. G. and Zimov, S. A.: Methane production and bubble emissions from arctic lakes: Isotopic implications for source pathways and ages, *J. Geophys. Res. Biogeosciences*, 113(G3), doi:10.1029/2007JG000569, 2008.
- Wang, P. X., Wang, B., Cheng, H., Fasullo, J., Guo, Z. T., Kiefer, T. and Liu, Z. Y.: The global monsoon across timescales: coherent variability of regional monsoons, *Clim. Past*, 10(6), 2007–2052, doi:10.5194/cp-10-2007-2014, 2014.
- Whitcar, M. and Schaefer, H.: Constraining past global tropospheric methane budgets with carbon and hydrogen isotope ratios in ice, *Philos. Trans. R. Soc. Lond. Math. Phys. Eng. Sci.*, 365(1856), 1793–1828, doi:10.1098/rsta.2007.2048, 2007.
- Whitcar, M. J.: Stable isotopes and global budgets, in *Atmospheric methane: sources, sinks, and role in global change*, pp. 138–167, Springer, Berlin., 1993.
- Yang, J.-W., Ahn, J., Brook, E. J. and Ryu, Y.: Atmospheric methane control mechanisms during the early Holocene, *Clim. Past*, 13(9), 1227, doi:10.5194/cp-13-1227-2017, 2017.



Yu, Z., Loisel, J., Turetsky, M. R., Cai, S., Zhao, Y., Frohling, S., MacDonald, G. M. and Bubier, J. L.: Evidence for elevated emissions from high-latitude wetlands contributing to high atmospheric CH₄ concentration in the early Holocene, *Glob. Biogeochem. Cycles*, 27(1), 131–140, doi:10.1002/gbc.20025, 2013.

Zhao, Y. and Harrison, S. P.: Mid-Holocene monsoons: a multi-model analysis of the inter-hemispheric differences in the responses to orbital forcing and ocean feedbacks, *Clim. Dyn.*, 39(6), 1457–1487, 2012.




Figure 1: CH₄ concentration data compilation. Discrete [CH₄] data over the Holocene used for the CH₄ data compilation on synchronised gas ages scales and corrected for offsets between datasets (references are listed in Table 1). The δ¹⁸O-H₂O data from the layer-counted NGRIP ice core have been used as an absolute time marker for fast climatic changes (e.g. the Younger Dryas-Holocene transition and the 8.2 ka event) assuming an essentially synchronous change in rapid temperature variations and atmospheric CH₄. The markers at the bottom of the plot indicate the tie points of the synchronised age scales (see Table 3).




Figure 2: Artificial time series and deconvolution results. Artificial tropospheric $[\text{CH}_4]$ (a) and stable isotope time series (b), (c) used as input for the deconvolution model. The calculated mean hemispheric CH_4 emission strengths (d) and their isotopic signatures (e), (f) show the dependence on the different input parameters. Three experiments are performed: all input values constant (solid lines), all constant but the northern tropospheric CH_4 isotope values varied (dashed lines) and all constant but $[\text{CH}_4]$ varied (dotted lines). To be consistent with the wording in the text, time is running from right to left. Note the inverted y-axes in the panels (b), (c), (e) and (f).




Figure 3: Data and deconvolution. The panels on the left (a-c) show the concentration and the two stable isotope signatures of CH₄ measured in ice cores representing the tropospheric values over the Holocene. The concentration data are complemented with data from other studies (see Table 1). All three parameters were measured on ice cores from Greenland (red) and from Antarctica (blue and cyan) representing the northern and the southern troposphere, respectively. A spline function with a cutoff period of 3000 years has been used to calculate the smoothed evolution of the tropospheric signals represented by the lines and the 1σ error bands. A deconvolution model is used to calculate the hemispheric CH₄ emissions shown on the panels on the right. In panel (d) the emission strengths are shown for both hemisphere boxes together with the total CH₄ emissions. Panel (e) and (f) show the mean isotopic signatures of the emissions for both stable isotopes of CH₄. All the error bands represent 1σ uncertainties. Note the inverted y-axes in the panels (b), (c), (e) and (f).

10




Figure 4: Comparison of stable CH₄ isotope data. This study's stable CH₄ isotope values of the NGRIP ice cores are shown together with the data published by Sowers (2010). The δD-CH₄ data sets (panel (a)) differ in the absolute level and the variance, whereas the δ¹³C-CH₄ data (panel (b)) are in good agreement. Note that all data are corrected for gravitational fractionation in the firm column. All data of this study are free of Krypton interference, whereas the GISP2 δ¹³C-CH₄ record has been subsequently corrected for it (Schmitt et al., 2013). Note the inverted y-axes.




Figure 5: Sensitivity study “SF₆ calibration”. In the panels (a-c) the data splines with the 1σ uncertainty bands of the measured tropospheric values are shown. In panels (d-f) the calculated strengths and the isotopic signature of the emissions are shown for the three different φ_{ITCZ} and θ setups to investigate the sensitivity of the deconvolution to the SF₆ calibration. Note the inverted y-axes in the panels (b), (c), (e) and (f).

5




Figure 6: Sensitivity study “ITCZ position”. In the panels (a-c) the data splines with the 1σ uncertainty error bands of the measured tropospheric values are shown. In panels (d-f) the calculated strengths and the isotopic signature of the emissions are shown for the three different φ_{ITCZ} and the best guess value for θ to investigate the sensitivity of the deconvolution to a shift of the ITCZ. Note the inverted y-axes in the panels (b), (c), (e) and (f).

5



Table 1: Datasets used for CH₄ data compilation. The CH₄ offset correction is added to the data to correct for inter-laboratory and time period-related offsets (see section 2.3.2). Additional datasets used for the time matching (e.g. covering the anthropogenic CH₄ increase or the Younger Dryas-Holocene transition) are listed. *GRIP data compilation done by J. Flückiger in 2001 containing data from various publications, e.g. Blunier et al. (1995), Chappellaz et al. (1993, 1997).

Data used for CH ₄ compilation			
Data set name	Data type	Reference	CH ₄ shift (ppb)
NGRIP_Be	NGRIP ice core	This study	0
GRIP_FI01	GRIP ice core	GRIP data Flückiger, 2001*	17.69
GISP2_Br96	GISP2 ice core	(Brook et al., 1996)	34.13
GISP2_Br09	GISP2 ice core	(Brook, 2009)	-1.81
GISP2_Mi13	GISP2 ice core	(Mitchell et al., 2013)	-6.15
TALDICE_Be17	TALDICE ice core	This study	0
TALDICE_Sc10	TALDICE ice core	(Schilt et al., 2010)	-2.09
EDML_Sc10	EDML ice core	(Schilt et al., 2010)	-4.63
WAIS_Mi13a	WAIS (WDC05A) ice core	(Mitchell et al., 2013)	-6.15
WAIS_Mi13b	WAIS (WDC06A) ice core	(Mitchell et al., 2013)	-6.15
EDC_FI02	EDC ice core	(Flückiger et al., 2002)	4.23
Additional data used for time matching			
Data set name	Data type	Reference	
NGRIP_d18O	NGRIP ice core $\delta^{18}\text{O}(\text{H}_2\text{O})$	(Rasmussen et al., 2006; Vinther et al., 2006)	
NGRIP_Ba12	NGRIP ice core	(Baumgartner et al., 2014)	
NEEM_Bu12	NEEM firn model	(Buizert et al., 2012)	
NOAA_Alert	Alert flasks	(Dlugokencky et al., 2017)	
LawDome_Ma06	Law Dome firn and ice core	(MacFarling Meure et al., 2006)	
CapeGrim_csiro	Cape Grim flasks	CSIRO (27.11.2017)	

5



Table 2: Ice cores and age scales. Short names of the ice cores and initial gas age scales used for the CH₄ compilation.

Ice core	Initial age scale	References
NGRIP	AICC2012 ice core chronology	(Bazin et al., 2013; Veres et al., 2013)
GRIP	GICC05	(Rasmussen et al., 2014; Seierstad et al., 2014)
GISP2	Mitchell, Meese/Sowers	(Meese et al., 1994; Mitchell et al., 2013)
TALDICE	AICC2012 ice core chronology	(Bazin et al., 2013; Veres et al., 2013)
EDML	AICC2012 ice core chronology	(Bazin et al., 2013; Veres et al., 2013)
EDC	AICC2012 ice core chronology	(Bazin et al., 2013; Veres et al., 2013)
WAIS	Mitchell, Buizert WD2004	(Buizert et al., 2015; Mitchell et al., 2013)



Table 3: Time synchronisation. Tie points of the reference time scale for the individual ice cores (left column, in aBP) and time corrections (right column, in a) applied with respect to the core dependent time scale listed in Table 2. The correction values are linearly interpolated between the tie points and held constant before the earliest and after the latest tie points.

NGRIP		GRIP		GISP2		TALDICE		EDML		EDC		WAIS	
294	0	273	-40	2752	-5	127	-134	2747	0	349	-225	2752	-5
11800	-150	8157	30	8172	-15	2835	-88	3754	39	2959	-212	3753	40
		11606	34	11771	-165	3825	-32	4538	63	3961	-168	4532	69
						4596	5	5279	130	4741	-140	5269	140
						5325	84	6008	215	5478	-69	5993	230
						6042	181	6361	129	6204	19	6344	146
						6389	101	8107	36	6555	-65	8079	64
						8106	37	8164	22	8292	-149	8228	28
						8252	4	8254	2	8353	-164	9869	-13
						9876	-20	9875	-19	8438	-182	10391	110
						10393	108	10391	110	9397	-179	11591	49
						11580	60	11576	64	9989	-133		
										10462	39		
										11547	93		



Table 4: GISP2 outliers. Data points from the two GISP2 data sets by Brook et al. (2009) and Brook (1996) that were removed due to elevated CH₄ values probably due to contamination related to the ice sample quality in the brittle zone. Please note that the age and the concentration values are corrected for the age shift and the CH₄ correction, respectively.

Removed data			
depth (m)	age (aBP)	CH ₄ (ppm)	dataset
655.1	2731.9	698.8	GISP2_Br09
736.1	3199.1	680.8	GISP2_Br96
819.7	3694.8	678.9	GISP2_Br09
932.7	4411.4	661.8	GISP2_Br09
943.1	4474.0	643.8	GISP2_Br09
992.3	4804.6	677.4	GISP2_Br96
1019.0	4989.5	642.0	GISP2_Br09
1059.0	5272.2	631.8	GISP2_Br09
1177.3	6186.3	639.6	GISP2_Br09
1194.2	6314.4	653.8	GISP2_Br96
1202.9	6382.8	652.1	GISP2_Br96
1253.9	6810.3	687.8	GISP2_Br09
1274.8	6978.6	677.8	GISP2_Br96
1389.7	7993.8	675.8	GISP2_Br96
1392.0	8011.9	682.2	GISP2_Br96



Table 5: Values used in our deconvolution model. Relative strengths, isotopic fractionations ($^2\epsilon$ and $^{13}\epsilon$) and north-south distribution of the individual sink processes (ratio n/s) and the parameters for each hemisphere used in our CH_4 deconvolution model.

sink processes				
	rel. strength	$^2\epsilon$ (‰)	$^{13}\epsilon$ (‰)	ratio n/s
S_{OH}	0.820	-231	-5.4	246.0/246.0
S_{strat}	0.092	-170	-13.1	27.5/27.5
S_{soil}	0.047	-80	-22.0	19.6/8.4
S_{Cl}	0.042	-315	-55.0	9.6/15.4
numbers used for deconvolution equations				
	τ_x (a)	$^2\epsilon_x$ (‰)	$^{13}\epsilon_x$ (‰)	r_x
north box	7.60	-218.3	-7.52	0.456
south box	9.22	-225.5	-7.72	0.544