

Referee #4 comments and responses:

In general, this article will be able to contribute to understand impact on marine biota by Fukushima derived radio activities. The data presented in this article about Ag-110m in neon flying squid is interesting.

Response: The authors thank the anonymous reviewer for their useful comments. We largely agree with the points raised and revised the manuscript accordingly.

Specific points:

1, It is however, a serious question about measurement of Ag-110m activity concentration in seawater which is a base of estimation of CR. The authors report that Ag-110g activity concentrations were below detection limit, but no description how they measure the Ag-110m activity concentration in seawater. In some conditions, Ag-110m can not be extracted well from sea water because Ag-110m exist in organic form. Therefore the authors should state how they measure the Ag-110m activity concentration and need to show reliability of their measurements.

Response: One paragraph describing the analytical method for seawater samples was added to the manuscript and the reliability of the method was demonstrated.

2, Another bog problem in this article is handling manner of the maximum number of significant digit throughout main text and Tables 1 and 2. In general based on the results, the maximum number of significant digit should be two or three, not four or more in case of this article. It is also needed to show the number as consistent number of significant digit throughout the article.

Response: The maximum number of significant digit was changed into 2 in Table 1 and Table 2 and throughout the article main text.

**Artificial radionuclides in neon flying squid from the northwestern Pacific in 2011
following the Fukushima accident**

Abstract:

In order to better understand the impact of the Fukushima Daiichi Nuclear Power Plant (FDNPP) accident on a commercial marine species, neon flying squid (*Ommastrephes bartramii*), samples obtained from the northwestern Pacific in November 2011 were analyzed for a range of artificial and natural radionuclides (Cs-134, Cs-137, Ag-110m, U-238, Ra-226 and K-40). Short-lived radionuclides Cs-134 and Ag-110m released from the Fukushima Nuclear Power Plant (NPP) accident were found in the samples, with an extremely high water-to-organism concentration ratio for Ag-110m ($> 2.9 \times 10^4$). While accident-derived radionuclides were present, their associated dose rates for the squid were far lower than the relevant benchmark of $10 \mu\text{Gy h}^{-1}$. For human consumers ingesting these squid, the dose contribution from natural radionuclides, including Po-210, was far greater ($>99.9\%$) than that of Fukushima-accident radionuclides ($<0.1\%$). The whole-body to tissue and whole-body to gut concentration ratios were calculated and reported, providing a simple method to estimate the whole-body concentration in environmental monitoring programs, and filling a data gap for concentration ratios in cephalopods. Our results help fill data gaps on uptake of NPP radionuclides in the commercially important Cephalopoda class and add to scarce data on open-ocean nekton in the northwestern Pacific shortly after the Fukushima accident.

Key words:

Fukushima NPP Accident; squid; concentration ratios; radiological dose; Silver-110m.

1. Introduction

The Fukushima Daiichi Nuclear Power Plant (NPP) accident, which was caused by the combined effect of the March 2011 earthquake and subsequent tsunami, resulted in increased levels of artificial radioactivity in the marine environment to the east of Japan (IAEA, 2015). The radioactive releases, dominated by radiocesium, were transported eastwards in surface water across the mid-latitude North Pacific at a speed of $3\text{-}7 \text{ km day}^{-1}$ ($3.5\text{-}8.0 \text{ cm s}^{-1}$) and dispersed widely in the North Pacific within a few years (Aoyama et al., 2013; Smith et al., 2015), which raised concerns about the

potential impact on marine biota and human consumers of seafood products.

A large amount of research has been conducted to determine the level of artificial radionuclides in biota samples and to assess the relevant radiological impact on both human and marine species. However, most studies have focused on the concentration of radiocesium in fish (Johansen, 2015; Wada, 2016), and only a few publications have reported on radionuclides in other marine species (Buesseler et al., 2012; Yu et al., 2015). Few data are available for open-ocean locations as compared to coastal areas, especially from 2011. Filling these data gaps will improve and expand understanding of the dynamics of cesium in the early months following the accident.

Ommastrephes bartramii (neon flying squid) is a migratory squid species that is commercially important, consumed by humans, and is common in both the Pacific Ocean and circumglobal temperate and tropical waters. It feeds near the surface on small fish and is thus a potential accumulator of radiocesium via dietary and water pathways. Moreover, cephalopods have a strong capability to accumulate silver in their bodies (Miramand and Bentley, 1992; Bustamante et al., 2004), potentially indicating uptake of the short-lived (0.70 year half-life) Ag-110m released from the Fukushima Daiichi NPP accident. Similarly, the presence of Cs-134 (2.1 year half-life) in samples would also indicate a pathway from Fukushima Daiichi NPP releases. Therefore, specimens captured at locations in the North Pacific may serve as bio-indicators of the presence, strength, and movement of the radioactive signal from the Fukushima Daiichi accident.

This study assessed samples of *O. bartramii* obtained from the northwestern Pacific in November 2011 for a range of artificial and natural radionuclides (Cs-134, Cs-137, Ag-110m, U-238, Ra-226 and K-40). The radiological dose rates and relevant risk levels were determined for the squid, as well as potential dose rates for human consumers of squid. Consistent with international efforts to compile transfer data, concentration ratios (whole-body to water and whole-body to tissue) were calculated and reported, including those for different age classes of squid.

2. Materials and methods

2.1. Sample collection and analytical procedure

Thirteen composite samples of *O. bartramii* samples with a total weight of 126.2 kg were obtained

by bait fishing in open water in the northwestern Pacific. Six sampling locations were selected within the area of 34°-39°N to 145°-149°E to investigate eastward deposition and oceanic migration pathways of radionuclide releases from the Fukushima Daiichi NPP (Fig 1Fig-1). To ensure sample mass was sufficient to reach minimum detectable activity (MDA) levels for key radionuclides, composite samples of multiple specimens from the same sampling site were used. For those sites with enough sample mass, the specimens were divided into different composite categories according to body weight. Specimens with a body mass less than 1 kg were categorized as “small”, those between 1 kg and 2 kg were categorized as “medium” and those heavier than 2 kg were categorized as “large.” The samples were frozen at -18°C on board for transport to the laboratory for subsequent analysis.

带格式的: 字体: Times New Roman

Squid samples were dissected after thawing into muscle and gut tissues (other soft tissues, including the digestive tract, gills, heart, gonads and associated glands), dried at 50°C, and then ashed at 450°C. The fresh weight and ash weight of the composite samples were recorded. The ash was sealed in cylindrical 75-mm diameter containers and was then subjected to a planar HPGe (High Purity Germanium) spectrometry for detection of gamma-emitting radionuclides.

At each station, 60 L surface seawater samples, were collected with a submersible pump, stored in polyethylene barrels with acidification to pH = 2, and taken back to land-based laboratory for analysis. With carriers of CsCl, AgNO₃, CoCl₂·6H₂O and FeNH₄(SO₄)₂·12H₂O added into the samples, Ag-110m in the seawater was collected with AgCl precipitation, Cs-134 and Cs-137 were collected with AMP (Ammonium Molybdophosphate) precipitation, and ⁵⁸Co, ⁶⁰Co, ⁵⁴Mn and ⁶⁵Zn were collected with Fe(OH)₃ precipitation, sequentially. The precipitate was collected with suction filtration, ashed at 450 °C, weighed, boxed, and then subjected to HPGe spectrometry. Method validation was carried out with standard solutions of Cs-137, Ag-110m, Co-60, Mn-54 and Zn-65.

- 带格式的: 字体: 五号
- 带格式的: 字体: 五号, 下标
- 带格式的: 字体: 五号
- 带格式的: 字体: 五号, 下标
- 带格式的: 字体: 五号
- 带格式的: 字体: 五号
- 带格式的: 字体: 五号, 下标
- 带格式的: 字体: 五号
- 带格式的: 字体: 五号, 下标
- 带格式的: 字体: 五号
- 带格式的: 字体: 五号, 下标
- 带格式的: 字体: 五号
- 带格式的: 字体: 五号, 下标
- 带格式的: 字体: 五号

Gamma rays from artificial radionuclides (Cs-134, Cs-137, Ag-110m, Co-58, Co-60, Mn-54 and Zn-65) and natural radionuclides (K-40, Ra-226 and U-238) were analyzed using a planar HPGe detector (Model BE6530 with Multi Channel Analyzer Lynx system; Canberra, U.S.A.). Detection efficiencies for the geometry used were 2.75%, 2.72%, 2.56%, 2.15%, 1.51%, 2.10%, 1.65%, 1.46%, 4.30% and 8.52%, for Cs-134, Cs-137, Ag-110m, Co-58, Co-60, Mn-54, Zn-65, K-40, Ra-226 and

U-238, respectively. The counting time for each sample was 24 hr. Genie 2000 software was used to analyze the respective peaks in the energy spectrum. The concentrations were corrected for decay to the initial date of the nuclear accident on 12 March 2011, when the first hydrogen explosion occurred in Unit 1 of the FDNPP (Wakeford, 2011).




Fig 1 Map of sampling sites

2.2. Dose assessment for squid

The ERICA Assessment Tool (version 1.2) (Brown et al., 2008) was used with Tier 2 assessment to evaluate the radiological risk to squid from the study areas in 2011. The ERICA Tool has the capability to specify organism sizes, and in this study, average mass (1.3 kg) and dimensions (ellipsoid equivalent of 0.3 m, 0.1 m, and 0.085 m for length, width, and height, respectively) from the specimens were used to calculate dose rates. The dimensions of the average *O. bartramii* are very similar to the standard ERICA “pelagic fish” and therefore, the dose rates, as calculated by ERICA, are also similar. The measured activity concentrations in the whole-body

of ^{137}Cs , ^{134}Cs , $^{110\text{m}}\text{Ag}$, ^{226}Ra and ^{238}U in the samples were used as dose calculation input. The maximum tissue activity concentrations were used for a more conservative result. As *O. bartramii* are migratory, their radionuclide tissue levels represent an integrated accumulation from recently traversed areas in the open ocean. Since the exact migratory routes are unknown, the external dose rates to the squid were calculated using the average of water radioactivity levels in the study capture region (average of samples across all sampling locations). In this instance, using the average is reasonable because the external dose rates for artificial radionuclides were much lower than the internal dose rates. As a result, variable water activity concentrations had little influence on overall dosages. For internal dose rates for squid, the dose conversion coefficients (DCCs) were calculated within the ERICA tool (supplemental). The occupancy factors were 100% in water, and weighting factors of internal low beta, internal beta/gamma and internal alpha were set as 3, 1 and 10, respectively.

2.3. Dose from ingesting squid

Committed effective doses (Sv) for human consumers of squid were estimated using standard exposure-to-dose conversion factors (DCFs) for ingestion from the ICRP Compendium of dose coefficients based on ICRP Publication 60 (ICRP, 1999). Key DCFs are 1.30×10^{-8} and 1.90×10^{-8} Sv Bq $^{-1}$ for Cs-137 and Cs-134, respectively (DCFs provided in the supplemental material). The factors are multiplied by intake (e.g., kg yr $^{-1}$) to obtain committed effective doses for the consumer. In this study, the annual intake rate of seafood by an adult consumer is assumed to be 20 kg yr $^{-1}$ (consistent with world per capita fish and related seafood consumption (FAO, 2016)). As a conservative assumption, the entire 20 kg yr $^{-1}$ for a hypothetical consumer is assumed to be sourced from the squid of the study area east of the Fukushima Daiichi NPP (in practice, only a small percentage of a seafood diet would be sourced from this region). As most dose to human consumers of seafood typically comes from the natural radionuclide Po-210 (~89% (Johansen et al., 2015)), the seafood ingestion dose rates here were compared with and without Po-210 to provide a context of the relative influence of Fukushima NPP accident radionuclides. For this comparison, a generic Po-210 seafood value of 15 Bq kg $^{-1}$ was used based on Hosseini et al. (2010) and consistent with the conservative (lognormal 95th percentile) based on the limited squid data in (Carvalho, 2011; Heyraud et al., 1994; Waska et al., 2008).

2.4. Whole-body concentration ratios

The water-to-organism whole-body concentration ratio ($CR_{WB:Water}$) used here is defined as:

$$CR_{WB:Water} = \frac{\text{Whole-Body Activity Concentration (fresh mass) (Bq/kg-wet)}}{\text{Water Activity Concentration (Bq/L)}} \quad (1)$$

The whole-body activity of a radionuclide was estimated using a mass balance approach (Yankovich et al., 2010) to reconstruct the amount of radionuclide in the whole-body of the squid. The whole-body to tissue concentration ratio ($CR_{WB:Tissue}$) was estimated as:

$$CR_{WB:Tissue} = \frac{\sum[\text{Tissue}_t \text{ Activity Concentration (fresh mass) Tissue}_t \text{ fresh mass fraction}]}{\text{Tissue}_t \text{ Activity Concentration (fresh mass)}} \quad (2)$$

3. Results and discussion

3.1. Description of *O. bartramii* specimens

In total, 98 specimens were obtained from 6 stations. The mass of the specimens ranged from 118 g to 2551 g, with an average of 1347 g. Sixty percent of the specimens weighed 701 g to 1700 g. The trunk length of the specimens ranged from 115 mm to 440 mm, on average 333 mm. Seventy-five percent of the specimens had a length greater than 290 mm (adult size), suggesting that the majority of the specimens hatched in the winter of 2010 or spring in 2011 and had been living for 8 to 11 months (Wang and Chen, 2005). Combining the estimated age of the squid, and assuming residence in the general region east of Fukushima Prefecture, it can be inferred that most specimens had been accumulating radionuclides since the Fukushima Daiichi NPP accident. —However, a minor proportion (the small size category) may have hatched after the accident and had shorter exposure times.

3.2. Activity concentrations and CRs in squid

The activity levels of radionuclides in Table 1 indicate that all *O. bartramii* size classes had accumulated radionuclides from Fukushima Daiichi NPP releases as indicated by Cs-134 and Ag-110m. The squid specimens had a strong capability to concentrate Ag in their bodies. The maximum activity of Ag-110m in the whole body of *O. bartramii* reached 9 Bq/kg, as compared to that in water, which was below the MDA of 0.22 Bq/m³, indicating a maximum concentration factor higher than 4×10^4 for all size classes. The mean CRs for Ag-110m were calculated as $> (2.953.0 \pm 0.9 \times)$

$\times 10^4 \pm 9.84 \times 10^3$ (Table 2), using the MDA as the activity of seawater in Equation (1).

Although this estimate contains large uncertainties because of using MDA of Ag-110m as the water concentration, these Ag data provide new insights for international researchers. Additionally, they fill a gap, as the relevant international database (Wildlife Transfer Parameter Database; www.wildlifetransferdatabase.org) and IAEA Technical Reports Series No.422 have entries for Ag uptake in the mollusk category (3.6×10^4 and 6×10^4 , respectively), but none specifically for squid/cephalopods.

The mean CR_{WB} values for Cs-134 and Cs-137 in *O. bartramii* were $6.33 (\pm 2.80 \text{ S.D.})$ and $5.57-6 (\pm 2.59-6 \text{ S.D.})$, respectively. These values are similar to previously published mean concentration factors for Cs ranging from 2 to 14 in cephalopods (Bustamante P, 2006; IAEA, 1978; Ishii et al., 1978; Suzuki et al., 1978; IAEA, 2004). The slightly lower CR_{WB} in this study is well within the range of expected variation, which can be very high for water-to-organism CR values (e.g. reported CRs for Cs-137 in marine fish range over nearly an order of magnitude) (Beresford, 2010). The activity concentration of ^{137}Cs in the research area reached a maximum of $\sim 600 \text{ Bq m}^{-3}$ in June 2011 and soon decreased to below 100 Bq m^{-3} (Aoyama et al., 2016). Considering the temporal change of radiocesium in seawater and its relatively short biological half-life (~ 70 days) in marine organisms, in this study, the CR calculation used mean Cs-134 and Cs-137 seawater activity concentrations (35.1 and 36.2 Bq m^{-3} , respectively) from this study's November 2011 sampling, which were similar to the $\sim 50 \text{ Bq m}^{-3}$ reported for the same open ocean area July-December timeframe (Kaeriyama, 2017).

The results also showed that both Cs-134 and Cs-137 were concentrated mainly in the muscle of the squid. Cesium behaves similarly to potassium in biota and tends to be distributed to the muscle tissue. These results for the open ocean, real-world conditions are consistent with previous laboratory results of 80+% accumulation in the muscle and head of cuttlefish after only 8 hours of exposure to water (Bustamante et al., 2004; Bustamante et al., 2006). In contrast, for Ag, the open ocean squid had 95% Ag in the gut versus muscle. This result was also consistent with the laboratory cuttlefish which had 98% Ag in the gut following a single spiked feeding and 29 d depuration (Bustamante et al., 2004). From the same study, within the gut, accumulation of Ag is dominant in

the digestive gland.

The smallest squid samples had the highest concentration factors for Cs-134, Cs-137, Ag-110m and U-238 (Fig 3). Despite their inferred shorter exposure times (shorter lifespan), the higher accumulation occurred in the smaller size class compared to the larger size class. These results are consistent with observed Cs depuration rates in juvenile cephalopods (*Sepia officinalis*) being ~four times slower than that of adults, but both relatively fast (adult cuttlefish have a biological half-life of 16 days for Cs and 9 days for Ag (Bustamante P, 2006)). This previous study suggests the radiocesium accumulation and depuration in *O. bartramii* is relatively rapid and that our results primarily reflect recent (~ several months) exposure rather than longer-term accumulation.

The levels of activity for ^{58}Co , ^{60}Co , ^{54}Mn and ^{65}Zn in the samples were all below the MDA (0.22 mBq/g-ash).




Fig 2 Activity concentrations of Cs-134 (left) and Ag-110m (right) in squid tissues.

Table 1 Radionuclide levels in composite samples (Bq/kg-fresh mass)

Size	Tissues	Cs-137		Cs-134		Ag-110m		K-40		Ra-226		U-238	
		Range	Average	Range	Average	Range	Average	Range	Average	Range	Average	Range	Average
All (n=13)	M	0.10-0.46	0.27±0.12	0.06-0.39	0.22±0.10	0.06-1.293	0.36±0.33	56.29-94.805	76.05±9.40	nd-0.07	0.03±0.03	0.16-1.778	0.59±0.44
	G	nd-0.33	0.05±0.10	nd-0.34	0.06±0.11	8.10-56.27	36.857±12.02	9.72-72.37	53.03±15.776	nd-0.89	0.28±0.28	nd-26.897	5.40±7.60
	WB	0.08-0.38	0.23±0.10	0.05-0.31	0.20±0.09	1.70-9.04	6.495±1.972_0	53.674-88.09	72.13±8.455	nd-0.17	0.07±0.05	0.27-5.32	1.354±1.40
Large (n=5)	M	0.13-0.46	0.26±0.13	0.09-0.39	0.22±0.11	0.06-0.36	0.24±0.13	67.628-94.805	80.551±11.49	nd-0.07	0.04±0.03	0.33-0.94	0.63±0.26
	G	ND	ND	ND	ND	32.25-40.501	36.14±3.192	49.5750-58.889	53.00±3.667	nd-0.68	0.24±0.27	nd-7.899	2.64±3.22
	WB	0.11-0.38	0.21±0.11	0.08-0.31	0.18±0.09	5.53-7.78	6.54±0.83	64.555-88.09	75.766±9.8610	nd-0.15	0.07±0.06	0.54-2.109	0.97±0.63
Medium (n=6)	M	0.10-0.41	0.27±0.14	0.06-0.34	0.22±0.12	0.06-0.46	0.25±0.19	56.29-78.789	72.30±8.12	nd-0.05	0.02±0.02	0.16-0.86	0.40±0.28
	G	nd-0.13	0.03±0.05	nd-0.21	0.04±0.08	8.10-45.856	31.40±13.05	9.72-67.738	47.808±21.18	nd-0.53	0.18±0.19	nd-10.44	3.051±3.84
	WB	0.08-0.35	0.23±0.12	0.05-0.29	0.19±0.11	1.70-8.091	5.64±2.34	53.674-73.944	68.16±7.56	nd-0.11	0.04±0.04	0.27-2.45	0.86±0.80
Small (n=2)	M	0.21-0.34	0.27±0.09	0.24-0.27	0.25±0.02	0.65-1.293	0.971_0±0.455	73.26-78.889	76.07±3.974_0	nd-0.05	0.02±0.03	0.41-1.778	1.091±0.971_0
	G	0.20-0.33	0.27±0.09	0.19-0.34	0.27±0.11	53.64-56.27	54.955±1.862	65.30-72.37	68.839±4.995_0	0.44-0.89	0.67±0.32	12.00-26.897	19.45±10.531
	WB	0.21-0.34	0.27±0.09	0.25-0.26	0.26±0.00	8.90-9.04	8.979_0±0.10	73.13-76.767	74.955±2.576	0.07-0.17	0.12±0.07	2.24-5.32	3.768±2.192

* Tissues: M – muscle, G – gut, WB – whole-body. **ND: level was below the minimum detectable activity.

203 **3.3. Whole-body to muscle and whole-body to gut concentration ratios**

204 Most of non-human biota radiation dose-assessing models focus on estimation of dose rates
205 using the *whole-body* activity concentrations of radionuclides (Brown et al., 2008; DOE, 2004).

206 However, muscle tissue (vs. whole-body) is measured in most monitoring programs, which
207 typically focus on seafood tissues consumed by humans. Therefore, there exists a need for
208 whole-body to tissue concentration ratios that allow for estimation of whole-body
209 concentrations from commonly measured tissue data (Yankovich et al., 2010).

210 The whole-body to muscle and whole-body to gut concentration ratios for radionuclides in
211 squid samples are listed in Table 2. For many radionuclides, the tissue-specific concentrations
212 for the small squids tend to be higher than those for large squids. The uncertainty of the whole-
213 body to gut CRs for Cs-137 and Cs-134 are relatively high because of the comparatively low
214 level and large activity range of radiocesium in gut samples. The CRs presented here are
215 calculated for the non-equilibrium conditions following the accident. This issue is somewhat
216 compensated for by using the average activity concentrations that have accumulated over time,
217 albeit over the relatively short lifespans of the squid. Equilibrium conditions are generally not
218 achieved in natural systems, and our results, all CRs should be considered in context. Further
219 research is necessary to obtain a better estimation the biokinetics of uptake in squid and of the
220 whole-body to gut CRs for Cs-137 and Cs-134.

221

Table 2 Concentration ratios for radionuclides in 2011 following the accident (see text).

CR*	Size	Cs-137	Cs-134	Ag-110m	K-40	Ra-226	U-238
WB-M	All	0.93±0.28	0.94±0.30	41.872±39.49	1.04±0.283	2.758±1.60	2.364±1.364
	Large	0.82±0.01	0.82±0.01	38.899±30.21	0.94±0.01	2.42±1.697	1.64±0.798
	Medium	0.85±0.03	0.86±0.06	47.908±50.29	0.96±0.01	2.00±1.53	2.354±1.20
	Small	1.00±0.00	1.01±0.071	10.30±4.675	0.99±0.02	3.586±0.5	4.22±1.73
WB-G	All	2.596±2.50	2.293±2.44	0.18±0.02	1.30±0.182	0.33±0.29	0.24±0.03
	Large	NA**	NA**	0.18±0.01	1.43±0.182	0.50±0.43	0.28±0.01
	Medium	4.152±4.152	3.54±3.54	0.18±0.02	1.253±0.12	0.24±0.09	0.24±0.01
	Small	1.03±0.01	1.04±0.40	0.16±0.00	1.091±0.12	0.17±0.02	0.19±0.01
WB-W***	All	6.33±2.80	5.576±2.596	$\frac{>(2.953.0 \times 10^4 \pm 0.9.8.94)}{\times 10^3 10^4}$	6.172±0.71	14.665±11.922	37.978±39.39
	Large	5.90±2.91	5.182±2.60	$\frac{>(2.973.0 \times 10^4 \pm 0.4) 3.76}{\times 10^{3d}}$	6.42±0.84	16.35±12.46	27.34±17.898
	Medium	6.30±3.172	5.33±3.04	$\frac{>(2.62.55 \times 10^4 \pm 1.051) \times}{10^4}$	5.989±0.697	9.566±9.40	24.11±22.43
	Small	7.52±2.52	7.293±0.061	$\frac{>(4.081 \pm 0.0) \times 10^4 + 4.50 \times}{10^2}$	6.354±0.22	25.766±15.07	$\frac{(1.1 \pm 0.6) \times}{10^2 106.09 \pm 61.85}$

222 * CR: WB-M represents whole-body to muscle concentration ratios, WB-G represents whole-body to gut concentration ratios, and WB-W represents whole-body to water concentration ratios.

223 ** NA: Data not available because radioactivity of specific radionuclides in at least one tissue was below the MDA.

224 *** Values for Cs-134 and Cs-137 were calculated using mean Cs-134 and Cs-137 seawater activity concentrations of 35.1 and 36.2 Bq m⁻³, and the values for Ag-110m were calculated using the225 MDA of Ag-110m in seawater (0.22 Bqm⁻³).

226 **3.4. Dose assessment results**

227 **3.4.1. Dose rates for squid**

228 The internal radiological dose rates in squid from artificial radionuclides (^{110m}Ag , ^{134}Cs and ^{137}Cs)
229 were collectively much higher than the external dose rates (Fig. 4). This is consistent with the
230 observed accumulation of radionuclides inside the squid body as compared with that in the
231 surrounding seawater. The internal dose rates from FDNPP-associated artificial radionuclides were
232 lower, by two orders of magnitude, than those from the natural radionuclides measured in this study.
233 Only approximately 1.4% of the total dose rate is estimated to have come from the Fukushima
234 Daiichi NPP releases. The total dose rate for squid is $0.15 \mu\text{Gy}\cdot\text{h}^{-1}$ from radionuclides measured in
235 this study, and increases to approximately $0.61 \mu\text{Gy}\cdot\text{h}^{-1}$ when adding Po-210, a natural radionuclide
236 significant dose contributor in marine organisms (assumes 0.001 Bq/L in seawater and a generic
237 marine value of 15 Bq/kg - whole-body fresh mass which is consistent with a general value in
238 Hosseini et al (2010) and with the lognormal 95th percentile of limited squid Po-210 data (Carvalho,
239 2011; Heyraud et al., 1994; Waska et al., 2008)). When median squid data are used (3 Bq/kg WB,
240 FM), the total dose rate is $0.25 \mu\text{Gy}\cdot\text{h}^{-1}$. Regardless of using the median or 95th percentile, these
241 dose rates are much lower than the most conservative screening benchmark dose rate of $10 \mu\text{Gy}\cdot\text{h}^{-1}$
242 (Garnier-Laplace, 2008). The dose calculations used the measured activity concentrations in the
243 squid (not CRs), and the calculated dose rates represent a point in time (November 2011) with
244 likely higher doses prior to, and lower doses following, the sampling date. However, the relatively
245 low values indicate that a more detailed (e.g., pulse-dynamic uptake) dose calculation is not
246 necessary in this case. Overall, results indicate that the radioactive releases from the Fukushima
247 accident would not have a significant adverse effect on *O. bartramii* individuals or populations
248 living in the study area.

249




Fig 3 Dose rates ($\mu\text{Gy}\cdot\text{h}^{-1}$) from measured radionuclides in squid samples

250
251
252

253 **3.4.2. Dose rates for human consumers of seafood**

254 From the radionuclides measured in edible squid tissue (muscle), a committed effective ingestion
 255 dose of 0.010 mSv (median; minimum = 0.007 mSv, maximum = 0.014 mSv) would have occurred
 256 in a hypothetical human consumer of 20 kg yr⁻¹ of squid from the study area (based on squid
 257 captured in November 2011). The doses calculated here are hypothetical and are intended to be
 258 conservative overestimates given the unrealistic assumption that all of the consumer's yearly
 259 seafood came from the study area. If consumption of Po-210 (from a natural background) is also
 260 included, the total dose increases to 0.30 mSv, with almost all derived from Po-210 using a
 261 conservative generic value as described above (Table 3). Of this dose (including Po-210), less than
 262 0.1% is estimated to have been sourced from the Fukushima Daiichi NPP. This is consistent with
 263 previous findings that natural radionuclides provided far greater dose rates to potential consumers
 264 of Pacific tuna (Fisher et al., 2013), and even for seafood sourced within a few kilometers of the
 265 Fukushima Daiichi NPP in 2013 (Johansen et al., 2015). The dose contribution from the Fukushima
 266 Daiichi NPP releases for squid consumption of this study are far below the 1 mSv per year
 267 recommended constraint for prolonged exposure by the public from nuclear facility releases (ICRP,
 268 1999).

269

270 **Table 3** Ingestion dose estimates for human consumers of the squid in this study (Sv y⁻¹ based on
 271 20 kg consumption of study squid).

	minimum	median	maximum	% this study*
K-40	6.98×10^{-6}	9.43×10^{-6}	1.18×10^{-5}	3.12%
Ag-110m	3.36×10^{-9}	2.02×10^{-8}	7.22×10^{-8}	0.01%
Cs-134	2.28×10^{-8}	8.36×10^{-8}	1.48×10^{-7}	0.03%
Cs-137	2.60×10^{-8}	7.02×10^{-8}	1.20×10^{-7}	0.02%
Ra-226		1.68×10^{-7}	3.92×10^{-7}	0.06%
U-238	1.44×10^{-7}	5.31×10^{-7}	1.59×10^{-6}	0.18%
Po-210**	1.44×10^{-5}	2.92×10^{-4}	1.08×10^{-3}	96.59%

272 * Based on median activity concentration values this study (Table 1 data, average of all sizes).

273 ** Po-210 from generic published data (Carvalho, 2011; Hosseini et al., 2010).

274

275 4. Conclusions

276 Elevated levels of Cs-134 and Ag-110m from the Fukushima NPP accident were found in the squid
 277 (*O. bartramii*) samples collected from NW Pacific in November 2011. The whole-body to water
 278 CRs for Ag-110m in squid were found to be as high as 4×10^4 L/kg in the smallest samples, with a
 279 mean value of 2.95×10^4 L/kg in all the samples, indicating that squid was a good bioindicator for
 280 Ag-110m from the Fukushima NPP accident. The radiological dose contribution from the
 281 Fukushima Daiichi NPP releases for squid living in the study area in 2011, and for human consumers
 282 of these squid, were both far below the recommended dose limits. By comparison, natural
 283 radionuclides, particularly Po-210, provided greater dose rates by several orders of magnitude. This
 284 study filled a gap in international transfer data by providing concentration ratios for several key
 285 NPP-associated radionuclides in the whole-body and tissues of an open ocean cephalopod.

286

287 Acknowledgements

288 This study was partially supported by the Scientific Research Foundation of the Third Institute of
 289 Oceanography, SOA (2015010), the Northwestern Pacific Marine Environmental Monitoring
 290 Project, the Coordinated Research Project (CRP K41017) and Regional Cooperative Agreement
 291 Project (IAEA/RCA RAS7028) of the International Atomic Energy Agency (IAEA), the National
 292 Key Scientific Instrument and Equipment Development Project (2016YFF0103905), the
 293 International Organizations and Conferences Project of the State Oceanic Administration of China,
 294 and the Public Science and Technology Research Project of the Ocean (201505005-1).

295

296 References

297

298 Aoyama, M., Tsumune, D., and Hamajima, Y.: Distribution of ¹³⁷Cs and ¹³⁴Cs in the North Pacific
299 Ocean: impacts of the TEPCO Fukushima-Daiichi NPP accident, *Journal of Radioanalytical and*
300 *Nuclear Chemistry*, 296, 535-539, 2013.

301 Aoyama, M., Hamajima, Y., Hult, M., Uematsu, M., Oka, E., Tsumune, D., and Kumamoto, Y.: ¹³⁴Cs
302 and ¹³⁷Cs in the North Pacific Ocean derived from the March 2011 TEPCO Fukushima Dai-
303 ichi Nuclear Power Plant accident, Japan. Part one: surface pathway and vertical distributions,
304 *Journal of oceanography*, 72, 53-65, 2016.

305 Beresford, N. A.: The transfer of radionuclides to wildlife, *Radiat Environ Biophys*, 49, 505-508,
306 2010.

307 Brown, J. E., Alfonso, B., Avila, R., Beresford, N. A., Copplestone, D., Prohl, G., and Ulanovsky, A.:
308 The ERICA tool, *Journal of Environmental Radioactivity*, 99, 1371-1383, 2008.

309 Buesseler, K. O., Jayne, S. R., Fisher, N. S., Rypina, I. I., Baumann, H., Baumann, Z., Breier, C. F.,
310 Douglass, E. M., George, J., and Macdonald, A. M.: Fukushima-derived radionuclides in the
311 ocean and biota off Japan, *Proceedings of the National Academy of Sciences*, 109, 5984-5988,
312 2012.

313 Bustamante, P., Teyssié, J. L., Danis, B., Fowler, S. W., Miramand, P., Cotret, O., and Warnau, M.:
314 Uptake, transfer and distribution of silver and cobalt in tissues of the common cuttlefish *Sepia*
315 *officinalis* at different stages of its life cycle, *Marine Ecology Progress Series*, 269, 185-195,
316 2004.

317 Bustamante, P., Teyssié, J. L., Fowler, S. W., and Warnau, M.: Assessment of the exposure pathway
318 in the uptake and distribution of americium and cesium in cuttlefish (*Sepia officinalis*) at
319 different stages of its life cycle, *Journal of Experimental Marine Biology and Ecology*, 331(2),
320 198-207, 2006.

321 Bustamante P, T. J. L., Fowler S W, et al. : Assessment of the exposure pathway in the uptake and
322 distribution of americium and cesium in cuttlefish (*Sepia officinalis*) at different stages of its
323 life cycle, *Journal of Experimental Marine Biology and Ecology*, 331(2), 198-207, 2006.

324 Carvalho, F. P.: Polonium (²¹⁰Po) and lead (²¹⁰Pb) in marine organisms and their transfer in
325 marine food chains, *Journal of Environmental Radioactivity*, 102, 462-472, 2011.

326 DOE, U. S.: RESRAD BIOTA: a tool for implementing a graded approach to biota dose evaluation,
327 Washington, DC, 2004.

328 FAO: The State of World Fisheries and Aquaculture 2016 (SOFIA) Contributing to food security
329 and nutrition for all, Rome, 200, 2016.

330 Fisher, N. S., Beaugelin-Seiller, K., Hinton, T. G., Baumann, Z., Madigan, D. J., and Garnier-Laplace,
331 J.: Evaluation of radiation doses and associated risk from the Fukushima nuclear accident to
332 marine biota and human consumers of seafood, Proceedings of the National Academy of
333 Sciences, 110, 10670-10675, 10.1073/pnas.1221834110, 2013.

334 Heyraud, M., Cherry, R.D., Oschadleus, H.D., Augustyn, C.J., Cherry, M.I., and Sealy, J.C.. Polonium-
335 210 and Lead-210 in edible molluscs from near the Cape of Good Hope: Sources of variability
336 in polonium-210 concentrations. *J. Environ. Radioact.* 24, 253-272, 1994.

337 Hosseini, A., Beresford, N. A., Brown, J. E., Jones, D. G., Phaneuf, M., Thorrying, H., and Yankovich,
338 T.: Background dose-rates to reference animals and plants arising from exposure to naturally
339 occurring radionuclides in aquatic environments, *Journal of radiological protection*, 30, 235,
340 2010.

341 IAEA: The Radiological Basis of the IAEA Revised Definition and Recommendations Concerning
342 High-level Radioactive Waste Unsuitable for Dumping at Sea. In: IAEA-TECDOC-211, IAEA,
343 Vienna, 1978.

344 IAEA: Sediment distribution coefficients and concentration factors for biota in the marine
345 environment, IAEA, Vienna, 2004.

346 ICRP: Protection of the Public in Situations of Prolonged Radiation Exposure, Publication 82,
347 International Commission on Radiological Protection, Ottawa, ON, 1999.

348 Ishii, T., Suzuki, H., and Koyanagi, T.: Determination of trace elements in marine organisms, 1:
349 Factors for variation of concentration of trace element, *Bulletin of the Japanese Society of*
350 *Scientific Fisheries*, 1978.

351 Johansen, M. P., Ruedig, E., Tagami, K., Uchida, S., Higley, K., and Beresford, N. A.: Radiological
352 dose rates to marine fish from the Fukushima Daiichi accident: the first three years across the
353 North Pacific, *Environmental science & technology*, 49, 1277-1285, 2015.

354 Kaeriyama, H.: Oceanic dispersion of Fukushima derived radioactive cesium: a review, *Fisheries*

355 Oceanography, 26, 99-113, 2017.

356 Miramand, P., and Bentley, D.: Concentration and distribution of heavy metals in tissues of two
357 cephalopods, *Eledone cirrhosa* and *Sepia officinalis*, from the French coast of the English
358 Channel, *Marine Biology*, 114, 407-414, 1992.

359 Smith, J. N., Brown, R. M., Williams, W. J., Robert, M., Nelson, R., and Moran, S. B.: Arrival of the
360 Fukushima radioactivity plume in North American continental waters, *Proceedings of the*
361 *National Academy of Sciences*, 112, 1310-1315, 2015.

362 Suzuki, Y., Nakahara, M., and Nakamura, R.: Accumulation of cesium-137 by useful mollusca,
363 *Nippon Suisan Gakkaishi*, 44, 325-329, 1978.

364 Wada, T., Fujita, T., Nemoto, Y., Shimamura, S., Mizuno, T., Sohtome, T., Kamiyama, K., Narita, K.,
365 Watanabe, M., Hatta, N., Ogata, Y., Morita, T., and Igarashi, S.. Effects of the nuclear disaster
366 on marine products in Fukushima: An update after five years. *Journal of environmental*
367 *radioactivity*, 164, 312-324, 2016.

368 Wakeford, R.: And now, Fukushima, *Journal of radiological protection*, 31, 167, 2011.

369 Wang, R., and Chen, X.: *The world ocean economic squid resources and Fisheries*, Beijing, 2005.

370 Waska, H., Kim, S., Kim, G., Kang, M.R., Kim, G.B.. Distribution patterns of chalcogens (S, Se,
371 Te, and ²¹⁰Po) in various tissues of a squid, *Todarodes pacificus*. *Sci. Total Environ.* 392,
372 218-224, 2008.

373 Yankovich, T. L., Beresford, N. A., Wood, M. D., Aono, T., Andersson, P., Barnett, C. L., Bennett, P.,
374 Brown, J. E., Fesenko, S., and Fesenko, J.: Whole-body to tissue concentration ratios for use in
375 biota dose assessments for animals, *Radiation and environmental biophysics*, 49, 549-565,
376 2010.

377 Yu, W., He, J., Lin, W., Li, Y., Men, W., Wang, F., and Huang, J.: Distribution and risk assessment of
378 radionuclides released by Fukushima nuclear accident at the northwest Pacific, *Journal of*
379 *Environmental Radioactivity*, 142, 54-61, 2015.

380