

5 Dear Editor,

Thanks for evaluating positively our manuscript. We also thank Reviewer 1, Reviewer 2 and David Sebag for their stimulating and constructive comments on our manuscript.

We have now revised the manuscript according to the reviewers suggestions, we describe and explain below the different changes in the revised version of the manuscript.

10 Our responses to reviewers are indicated in blue while the **locations of changes** in the revised manuscript are indicated **in red**.

We also did some minor modifications in the processing of raw Rock-Eval 6 data (**p.5 1.30–31 in the revised manuscript**). All calculations (RE6 parameters, correlations, principal component analysis, random forests regression model) have been performed again with these modifications, marginally modifying the results that
15 have all been updated **in the text** and in **Fig. 3, Fig. 4, Fig. 5 and Table 2** of the revised manuscript.

We are looking to hearing from you.

Yours sincerely,

Lauric Cécillon

20

Reviewer 1's comments

Major point 1: use of the concept of residence time in the manuscript

(1) Comment from Referee 1

5 One important source of confusion in this manuscript is the use of the concept residence time. Notice that in soils one must distinguish between the concepts of age and transit (residence) time (Bruun et al., 2004; Manzoni et al., 2009; Derrien and Amelung, 2011). What the authors are trying to estimate here is an indication of the age of the SOM, not the residence time. These should be more clearly treated in the introduction and the discussion. Currently, the use of these terms is ambiguous.

(2) Response to Referee 1's comment

10 We agree with Reviewer 1 that it is important to distinguish between the concepts of age and residence time of organic carbon in soils. However, our methodology does not focus on either the estimation of the age nor on the estimation of the residence time of soil organic carbon (SOC). Our methodology aims at estimating the size of the centennially persistent SOC (CPsoc) pool, a SOC pool whose mean age and mean residence time are both assumed to be high (e.g. several centuries) but for which precise definitions are not necessary. The only required property for defining the CPsoc pool is not its mean age or its mean residence time, but the non-significant change in its size in periods inferior to the century. Specifically, we defined the CPsoc pool (gC.kg⁻¹ soil) at each site as the constant term of an exponential plus constant model fitted on the temporal evolution of SOC under bare fallow treatment. The only assumption that we made was that, given our data set (temporal evolution of SOC under 5 to 8 decades of bare fallow treatment or associated non-bare fallow treatment), the size of the CPsoc pool remained constant (i.e. did not change significantly, see also our response to major point 2 below).

(3) Proposed changes in manuscript

25 Overall, we thus did not make any specific estimation of the age or the residence time of the CPsoc pool, but we propose to treat this point more clearly in a revised version of our manuscript (introduction and discussion sections) to avoid any confusion regarding the concepts of age and residence time of organic carbon in soils. (p.4 1.31-33 in the revised manuscript)

Major point 2: centennially persistent soil organic carbon is not inert

(1) Comment from Referee 1

30 The other important issue in this manuscript is also conceptual. The model presented in equ. 1 and used to compute the centennially persistent pool is, in my opinion, inappropriate. It assumes that an amount of 'inert' carbon c is sitting there doing nothing and it will never decompose. This is highly unlikely, because there's

always some small probability that carbon that is stabilized either by mineral association or protection in aggregates, would get consumed by microorganisms and respired as CO₂.

(2) Response to Referee 1's comment

5 We fully agree with Reviewer 1 that organic carbon is not biogeochemically inert in soils vis-à-vis microbial decomposition, and that the CPsoc pool is also (though very slowly) gaining and losing carbon. However, regarding the CPsoc pool, its high residence time (e.g. several centuries; not precisely defined here, see above) masks changes in its size at the time scale of our sample set (5 to 8 decades). We thus pragmatically and operationally considered the CPsoc pool to be mathematically inert (the constant term in an exponential plus constant model) given our data set, though we agree that the CPsoc pool is a biogeochemically stable but not inert SOC pool. We argue that since all models are simplifications of reality, assuming an inert SOC pool is acceptable for a SOC pool with very low decomposition rate, as implemented in many widely used SOC models such as RothC. We have considered performing a double exponential (without constant) as suggested by Reviewer 1, but we think that a constant is mathematically the most appropriate way to model the CPsoc pool with our data (maximum 80 years of decomposition). Furthermore, we think that the argument that a double exponential model does not add parameters is not true: $y(t)=y_0*a*exp(-k_1*t) + y_0*(1-a)*exp(-k_2*t)$ has 3 parameters – but $y(t)= a + y_0*(1-a)*exp(-k_1*t)$ has only 2. Parsimony (Ockham's razor) and equifinality of more complex model lead us to propose keeping a single exponential plus constant model to estimate the size of the CPsoc pool (gC.kg-1 soil) in the bare fallow treatments of each study site.

(3) Proposed changes in manuscript

20 We propose to revise the manuscript to clarify that the CPsoc pool is not biogeochemically inert in our view, but that it is mathematically more sound to mathematically simulate it as a constant on such a dataset. (p.4 1.31–33 and p.5 1.1–2 in the revised manuscript)

Reviewer 2's comments

25 Major point 1: Validation of the multivariate regression model to predict the size of the centennially persistent SOC (CPsoc) pool in “new” soils

(1) Comment from Referee 2

30 Machine learning is used to find the best regression model predicting the proportion of CPsoc. A high R² (0.91) in the calibration dataset is impressive and shows that the thermal stability (RE6) can be linked to biogeochemical stability, which has been shown before. Now, the interesting thing is the validation: the authors report the same R² for the validation set and show that they scatter as nicely along the 1:1 line as the calibration dataset does: Of course, this is the case because the dataset was randomly split, although the samples were not independent but originated from only 4 experiments. So the question is, will the results be similarly good, if for example 3 sites are used to train the model and 1 site is used for validation? This would

give a much more honest picture on the validity of the approach. I guess that the prediction would not be as good: According to the Barre et al 2016 paper, at least the three presented thermal stability parameters HI, OI and T50CO2_ox which played an intermediate to important role also in the present study, varied considerably across sites. Also Figure 3A indicates that the ‘thermal signature’ of the samples is really site dependent. So for me the question is: Can this product really hold what the authors are promising, e.g. in the last sentence of the abstract: ‘This model can thus be used to predict...’? This has to be clarified and if not the case be discussed with much more caveats. Uncertainties are already huge and they would probably inflate if new samples shall be predicted.

(2) Response to Referee 2’s comment

We randomly split our sample set into a calibration and a validation set, therefore including soil samples from the four study sites in both the calibration and the validation set. We thus agree with Reviewer 2 that the validation of the multivariate regression model is not based on truly independent soil samples, even though samples from the validation set were not used in the calibration set. As discussed in the manuscript, the good fit obtained for the current validation set (Figure 4 in the manuscript) indicates that the multivariate regression model can be used to predict the size of the CPsoc pool with a known uncertainty in soils with pedoclimates similar to those found in the four study sites (Versailles, Grignon, Rothamsted and Ultuna; Supplementary material S1).

However, Reviewer 2 asks an important question: “will the results be similarly good, if for example 3 sites are used to train the model and 1 site is used for validation?”

Since each site used in this study has a specific pedoclimate (even the two sites with a similar climate, Versailles and Grignon, have different soil mineralogy, with carbonate soils in Grignon and soils developed in loess in Versailles), we have to slightly rephrase Reviewer 2’s question regarding the specificity of our data set. In fact, Reviewer 2 asks if the multivariate regression model is able to predict the size of the CPsoc pool in soils with a different pedoclimate (i.e. a pedoclimate not included in the calibration set). Or more generally, can the model predict the CPsoc proportion outside of the studied sites?

We agree with Reviewer 2 that testing the multivariate regression model sensitivity to pedoclimate (i.e. by validating it on a site with a new pedoclimate) would provide useful information to the readers regarding its applicability on soils from different pedoclimates.

We argue that a necessary prerequisite for applying the multivariate regression model on “new” soil samples from a different pedoclimate is the thermal similarity between the “new” soil samples and samples of the calibration set (i.e. similar range of values for the 30 Rock-Eval parameters that were used as predictors in the multivariate regression model).

All soils from Grignon (carbonate site), and some samples from Versailles and Rothamsted show some important specificities regarding their thermal characteristics, while all soils from Ultuna had thermal characteristics similar to some samples from Versailles and Rothamsted (see the PCA plot in Figure 3 in the manuscript).

(3) Proposed changes in manuscript

We therefore propose to add a second validation scheme to test the sensitivity of the multivariate regression model to a different pedoclimate. Ultuna can be used as a truly independent validation site with similar thermal characteristics but a different pedoclimate than the calibration set.

5 The results of the predictions for Ultuna samples, using samples from Versailles, Grignon and Rothamsted for calibration of the multivariate regression model are shown in Figure 1 (see below). As expected, the R² strongly decreases, yet the error of prediction of the model does not increase strongly (RMSEP = 0.09 vs. 0.07 in the previous validation scheme).

10 Overall, these new results illustrate the sensitivity of the multivariate regression model to a very different pedoclimate (different climate and soil mineralogy), yet they clearly show the potential of the model based on Rock-Eval analysis for predicting the proportion of CPsoc in “new” soil samples. We thus propose to include and discuss them thoroughly in a revised version of the manuscript.

(Abstract Section p.2 1.1–15, Material and Methods Section p.8 1.23–29, Results Section p.11 1.8–11 and 1. 16–20, Discussion Section p.13 1.26–32 and new Figure 5 in the revised manuscript)

15

Specific comments:

(1) Comment from Referee 2

P5, line 8: Why such a huge intercept in the regression (0.4)?

(2) Response to Referee 2’s comment

20 The relatively high value of the intercept may be linked to the fact that when estimating the total organic carbon content, a small amount of organic carbon is not taken into account by the commercial software of Rock-Eval 6 (organic carbon being volatilized as CO or CO₂ at high temperatures, that may be inorganic carbon in carbonated soils). Underestimation of SOC concentration by RE6 has already been reported (e.g. Saenger et al., 2013). As soils from Grignon contain carbonates, we chose the same metric of SOC_RE6 for all samples, even if
25 they are slightly biased towards lower values.

(1) Comment from Referee 2

P6, line 23: Fixed standard deviation for SOC concentration data? Why, and what is it exactly derived from?

(2) Response to Referee 2’s comment

30 We used a fixed value for the standard deviation for SOC concentration data obtained from elemental analyzer (SOC_EA; 0.75 gC.kg⁻¹ soil). As stated in the manuscript, this value is a conservative estimate of the standard

deviation of SOC_EA data estimated by Barré et al. (2010) for the same soils. In the latter paper, the authors stated that “standard deviation [was] estimated from 15 replicate determinations of C in soil samples taken from the same plot in Grignon in 1959 (Barré et al., 2010). The measured standard deviation was 0.3 gC.kg⁻¹. As the C contents at the different LTBF sites were determined on composite samples from the same plot at each site, it was considered that the a priori error on measurements should be less than 0.5 gC.kg⁻¹ for every site”. They finally applied a standard deviation of 0.5 gC.kg⁻¹ for every site except Versailles, where the final standard deviation for SOC concentration data was 0.75 gC.kg⁻¹ (Barré et al., 2010).

(1) Comment from Referee 2

P12, line 28: I thought it was 30 RE6 parameters, here it says it was 25?

10 (2) Response to Referee 2’s comment

We have indeed used a total of 30 RE6 parameters in the multivariate regression model, but only 25 of them are RE6 temperature parameters (i.e. unit: °C). In this section of the manuscript (P12, line 28), we only discuss the outcomes of the 25 temperature parameters. We discuss the outcomes of the remaining 5 RE6 parameters (TLHC-index, I-index, R-index, HI and OIRE6) later in the manuscript (P13, lines 12-25).

15 (1) Comment from Referee 2

Discussion 1: Is very technical and focused on the specific RE6 method and related papers. Authors miss the chance to broaden the perspective and discuss this approach to estimate CPsoc in comparison to other approaches or to establish a clear link to kinetic models.

(2) Response to Referee 2’s comment

20 Only one out of three sections of the discussion (section 4.2) is devoted to the discussion of the specific RE6 method and related papers. We agree that this section 4.2 is rather technical but we think that critically discussing some technical limitations of our approach is necessary. Section 4.1 of the discussion already broadens the scope of our study and discusses different methodologies that have been used to produce estimates of the CPsoc concentration under various pedoclimatic conditions (i.e. in other long term agronomic experiments or using other analytical techniques and/or models such as radiocarbon (¹⁴C) data and steady-state SOC turnover model).

We agree with Reviewer 2 that discussing our approach to estimate CPsoc in comparison to other approaches may be useful to the reader.

(3) Proposed changes in manuscript

30 We propose to add some references to alternative techniques used for initializing the size of SOC kinetic pools in models of SOC dynamics (e.g. Falloon et al., 1998; Zimmermann et al., 2007) in section 4.3 of the discussion.

(Discussion Section p.15 l.25–28 in the revised manuscript)

(1) Comment from Referee 2

Discussion 2: Is very positive about the overall results (and sample set), and although uncertainty was a clearly stated focus of the study it is not really taken up here: Yes, the sample set is truly unique, but this is also the problem: How uncertain will the CPsoc estimation in soils be, which do not have bare fallow treatments?

5 (2) Response to Referee 2's comment and (3) Proposed changes in manuscript

Following our response to the major point raised by Reviewer 2 (see above), we propose to revise the manuscript to discuss in section 4.3 the new results regarding the multivariate regression model sensitivity to pedoclimate (i.e. prediction on “new” soils from a different pedoclimate).

10 **References:**

- Barré, P., Eglin, T., Christensen, B.T., Ciais, P., Houot, S., Kätterer, T., van Oort, F., Peylin, P., Poulton, P.R., Romanenkov, V., and Chenu, C.: Quantifying and isolating stable soil organic carbon using long-term bare fallow experiments. *Biogeosciences*, 7, 3839–3850, 2010.
- Falloon, P., Smith, P., Coleman, K., and Marshall, S.: Estimating the size of the inert organic matter pool from total soil organic carbon content for use in the Rothamsted carbon model. *Soil Biol. Biochem.*, 30, 1207–1211, 1998.
- Saenger, A., Cécillon, L., Sebag, D., and Brun, J.J.: Soil organic carbon quantity, chemistry and thermal stability in a mountainous landscape: A Rock-Eval pyrolysis survey. *Org. Geochem.*, 54, 101–114, 2013.
- Zimmermann, M., Leifeld, J., Schmidt, M.W.I., Smith, P., Fuhrer, J.: Measured soil organic matter fractions can be related to pools in the RothC model. *Eur. J. Soil Sci.*, 58, 658–667, 2007.

A model based on Rock-Eval thermal analysis to quantify the size of the centennially persistent organic carbon pool in temperate soils

Lauric Cécillon^{1,2}, François Baudin³, Claire Chenu⁴, Sabine Houot⁴, Romain Jolivet², Thomas Kätterer⁵, Suzanne Lutfalla², Andy Macdonald⁶, Folkert van Oort⁴, Alain F. Plante⁷, Florence Savignac³, Laure [N. Soucémarianadin](#)², Pierre Barré²

¹Université Grenoble Alpes, Irstea, UR [EMGRLESSEM](#), F-38402 St-Martin-d'Hères, France

²Laboratoire de Géologie de l'ENS, PSL Research University, CNRS UMR 8538, F-75005, Paris, France

³Sorbonne Université-UPMC-Univ Paris 06, Institut des Sciences de la Terre de Paris, F-75005 Paris, France

⁴AgroParisTech - INRA, UMR 1402 ECOSYS, F-78850 Thiverval-Grignon, France

⁵Department of Ecology, Swedish University of Agricultural Sciences, 750 07 Uppsala, Sweden

⁶Department of Sustainable Agriculture Sciences, Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ, UK

⁷Earth and Environmental Science, University of Pennsylvania, Philadelphia, USA

Correspondence to: Lauric Cécillon (lauric.cecillon@irstea.fr)

Abstract. Changes in global soil carbon stocks have considerable potential to influence the course of future climate change. However, a portion of soil organic carbon (SOC) has a very long residence time (> 100 years) and may not contribute significantly to terrestrial greenhouse gas emissions during the next century. The size of this persistent SOC reservoir is presumed to be large. Consequently, it is a key parameter required for the initialization of SOC dynamics in ecosystem and Earth system models, but there is considerable uncertainty in the methods used to quantify it. Thermal analysis methods provide cost-effective information on SOC thermal stability that has been shown to be qualitatively related to SOC biogeochemical stability. The objective of this work was to build the first quantitative ~~thermal analysis based~~ model of the size of the centennially persistent SOC pool based on thermal analysis. We used a unique set of 118 archived soil samples from four agronomic experiments in Northwestern Europe with long-term bare fallow and non-bare fallow treatments (*e.g.* manure amendment, cropland and grassland), as a sample set for which estimating the size of the centennially persistent SOC pool is relatively straightforward. At each experimental site, we estimated the average concentration of centennially persistent SOC and its uncertainty by applying a Bayesian curve fitting method on the observed declining SOC concentration over the duration of the long-term bare fallow treatment. Overall, the estimated concentrations of centennially persistent SOC ranged from 5 to 11 gC.kg⁻¹ soil (lowest and highest boundaries of four 95% confidence intervals). Then, by dividing site-specific concentrations of persistent SOC by the total SOC concentration ~~of 118 archived soil samples from long-term bare fallow and non-bare fallow treatments~~, we could estimate the proportion of centennially persistent SOC in the 118 archived soil samples and the associated uncertainty. The proportion of centennially persistent SOC ranged from 0.14 (standard deviation of 0.01) to 1 (standard deviation of 0.15). Samples were subjected to thermal analysis by Rock-Eval 6 that generated a series of 30 parameters reflecting their SOC thermal stability and bulk chemistry. We trained a non-parametric machine learning algorithm (random forests multivariate regression model) to predict the proportion of centennially persistent SOC in new soils using Rock-Eval 6 thermal parameters as predictors. We evaluated the model predictive performance with two different strategies. We first used a calibration set (n = 88) and a validation set (n = 30) with soils from all sites. Second, to test the sensitivity of the model to pedoclimate, we built a calibration set with soil samples from three out of the four sites (n = 84). The sample set was split into a calibration set (n = 88) and a validation set (n = 30). We trained a non-parametric machine learning algorithm (random forests multivariate regression model) that ~~The multivariate regression model~~ accurately predicted the size-proportion of ~~the~~ centennially persistent SOC ~~pool using Rock-Eval 6 thermal parameters as predictors~~ in the calibration set (pseudo R² = 0.91, RMSEC = 0.06) and the validation set composed of soils from all sites (R² = 0.9492, RMSEP = 0.07, n = 30). The uncertainty of the model predictions ~~obtained using the multivariate regression model~~ was quantified by a Monte Carlo approach that produced conservative 95% prediction intervals across the ~~30 samples of the~~ validation set. The predictive performance of the model decreased when

~~predicting the proportion of centennially persistent SOC in soils from one fully independent site with a different pedoclimate, yet the mean error of prediction only slightly increased ($R^2 = 0.53$, RMSEP = 0.10, $n = 34$). This model based on Rock-Eval 6 thermal analysis can thus be used to predict the proportion of centennially persistent SOC with known uncertainty in new soil samples from different pedoclimates, at least for sites that have similar Rock-Eval 6 thermal characteristics to those included in the calibration set. This model based on Rock-Eval 6 thermal analysis can thus be used to predict the proportion of centennially persistent SOC with known uncertainty in new soil samples from similar pedoclimates.~~ Our study ~~reinforces the evidence that there is a link between~~~~strengthens the evidence for a link between~~ the thermal and biogeochemical stability of soil organic matter, and demonstrates that Rock-Eval 6 thermal analysis can be used to quantify the size of the centennially persistent organic carbon pool in temperate soils.

10 1 Introduction

Soils exert a key regulation of the atmospheric greenhouse gas concentrations on a decadal timescale through the net carbon source and sink status of their organic carbon reservoir (Amundson, 2001; Eglin et al., 2010). However, a portion of the soil organic carbon (SOC) reservoir may not contribute significantly to the net exchange of CO_2 and CH_4 between atmosphere and land during the next century because its residence time exceeds 100 years and its rate of carbon inputs is low (Trumbore, 1997; He et al., 2016). The size of ~~theis~~ centennially persistent SOC pool is presumed to be large (*i.e.* between one and two thirds of total SOC) and dependent on geochemical parameters such as soil texture and mineralogy (Buyanovsky and Wagner, 1998a; Trumbore, 2009; Mills et al., 2014; Mathieu et al., 2015). However, the amount of centennially persistent organic carbon in soils is highly uncertain as it cannot be estimated accurately by current analytical methods (Post and Kwon, 2000; von Lützow et al., 2007; Bruun et al., 2008). Physico-chemical procedures attempting to isolate SOC with high residence time from bulk SOC have proven unsatisfactory because of indications that such fractions are a mixture of ancient and recent SOC (von Lützow et al., 2007; Trumbore, 2009; Lutfalla et al., 2014). Even the well-established radiocarbon (^{14}C) analytical technique cannot precisely determine the size of the centennially persistent SOC pool (Schrumppf and Kaiser, 2015; Menichetti et al., 2016). The importance of better information on the size of the centennially persistent SOC pool has been emphasized recently (International Soil Carbon Initiative, 2011; ~~Bailey et al., in press~~; Bispo et al., 2017; ~~Bailey et al., 2018; Harden et al., 2018in press~~), stressing the need for operational and standardized metrics or proxies to accurately quantify SOC persistent at the centennial timescale. The general lack of information on the size and turnover rate of measurable SOC pools hampers the initialization of SOC pools in dynamic models, questioning their predictions of the evolution of the global SOC reservoir (Falloon and Smith, 2000; Luo et al., 2014; Feng et al., 2016; He et al., 2016; Sanderman et al., 2016). Luo et al. (2016) and He et al. (2016) recently claimed that optimizing parameter~~s~~ estimation with global data sets on SOC pools and fluxes was the highest priority to reduce biases among Earth system models.

~~During-In~~ the past decade, thermal stability of organic carbon has been proposed as a good surrogate for its biogeochemical stability in litter and soils (*e.g.* Rovira et al., 2008; Plante et al., 2009; Gregorich et al., 2015). Several studies using thermal analysis techniques such as thermogravimetry and differential scanning calorimetry ~~during-with~~ ramped combustion have shown that the fast-cycling SOC pool determined as the amount CO_2 respired in laboratory incubation experiments was thermally labile (Plante et al., 2011; Leifeld and von Lützow, 2014; Campo and Merino, 2016). Recently, studies using thermal analysis under oxidative or inert (pyrolysis) reaction atmosphere coupled with evolved gas analysis have shown a high and positive correlation between the thermally stable SOC and persistent SOC determined using ^{14}C measurements (Plante et al., 2013), and between thermally stable SOC and mineral-associated SOC isolated by a classical SOC physical fractionation scheme (Saenger et al., 2015). Using long-term bare fallow (LTBF) soils kept free of vegetation for several decades (*i.e.* with negligible carbon inputs), Barré et al. (2016) recently showed that persistent SOC was low in energy and

thermally stable. While there appears to be strong qualitative links between thermal and biogeochemical stability of SOC, there is to date no established quantitative link between the size of the persistent SOC pool and SOC thermal characteristics.

The objective of this work was to design a reliable, routine method based on a thermal analysis technique (Rock-Eval 6; RE6) to quantify centennially persistent SOC in a range of temperate soil types. First, we compiled a set of reference soil samples from four long-term agronomic experiments in Northwestern Europe with long-term bare fallow treatments. The SOC concentration of LTBF treatments can be used to estimate the size of the persistent SOC pool of a particular site, as proposed by Rühlmann (1999) and Barré et al. (2010). Here, we refined estimates of the persistent SOC concentration previously published by Barré et al. (2010) for the four sites used in this study. We then used these values to estimate the proportion of centennially persistent SOC in 118 archived soil samples (time series) from LTBF and non-LTBF treatments of these four sites. The last step consisted in analyzing these reference samples using RE6 thermal analysis, and building a multivariate regression model to relate RE6 information on SOC thermal stability and bulk chemistry to the estimated proportion of centennially persistent SOC. In this work, we aimed at delivering a model based on thermal analysis-based model with reliable prediction intervals around the predicted values of the size of the centennially persistent SOC pool. We thus had a particular focus focused on the uncertainty of the estimated proportion of centennially persistent SOC and its propagation in the multivariate regression model.

2 Materials and methods

2.1 Reference soil sample set with estimated size of the centennially persistent SOC pool

The reference soil sample set was built using samples from four long-term agronomic experimental sites in Northwestern Europe (Versailles, France, Grignon, France, Rothamsted, United Kingdom, Ultuna, Sweden, Supplementary material S1). Each of the four sites includes a LTBF treatment, with bare fallow durations ranging from 48 years at Grignon to 79 years at Versailles. For all experimental sites, we also included non-LTBF treatments that have increased or maintained their total SOC concentrations over time, or sustained smaller losses than the LTBF treatment. The selected non-LTBF treatments included manure amendments (Versailles), straw or composted straw amendments (Grignon), continuous grassland (Rothamsted), and continuous cropland (Ultuna). Soil samples from each site and treatment have been regularly collected and archived since the initiation of the experiments. A total of 118 topsoil samples (0–20 to 0–25 cm depth, Supplementary material S1) ~~samples~~ were selected from the archives of LTBF and non-LTBF treatments to build the reference sample set. Samples were selected from two or three field replicate plots with a decadal frequency in each decade from the initiation of the experiments up to 2007 (Grignon), 2008 (Versailles, Rothamsted) or 2009 (Ultuna) to obtain a sample set with the widest possible range of proportions of centennially persistent SOC. The non-LTBF treatments and multiple sites also added to the diversity of land-use, climate and parent material. For each sample, total SOC concentration was measured by dry combustion with an elemental analyzer (SOC_{EA} , gC.kg^{-1} soil) after decarbonation when necessary according to NF ISO 10694 (1995).

Based on the decline in total SOC concentration over the duration of the LTBF treatment, Barré et al. (2010) estimated the concentration of centennially persistent SOC at each site using a Bayesian curve fitting method applied to each LTBF field replicate plot. Here, we refined those site-specific estimates by (i) applying a similar Bayesian curve fitting method on combined SOC concentration data from all LTBF field replicate plots of each site (four field replicate plots for Ultuna and Rothamsted, six field replicate plots for Versailles and Grignon), and (ii) using new SOC concentration data up to 2014 for Rothamsted and 2015 for Ultuna, increasing their LTBF duration to 55 years for Rothamsted and 59 years for Ultuna.

For each site, we assumed that the temporal evolution of LTBF SOC concentration, $\gamma(t)$, followed an exponential decay function:

$$\gamma(t) = ae^{-bt} + c, \quad (1)$$

5 where $\gamma(t)$ (unit = gC.kg^{-1} soil) is the LTBF SOC concentration at time t , t (unit = year) is the time under bare fallow, and a , b and c are fitting parameters. Parameter a (unit = gC.kg^{-1} soil) corresponds to the amplitude of the decay and b (unit = y^{-1}) is the characteristic decay rate. The parameter c (unit = gC.kg^{-1} soil) represents a theoretically inert portion of SOC. We considered this parameter as a site-specific metric of the centennially persistent SOC concentration. In our view, the centennially persistent SOC pool is not biogeochemically inert, it has a mean age and a mean residence time that are both assumed to be high (e.g. several centuries) though not precisely defined in this study. As a result, its small decline cannot be observed at the timescale of this study and we thus considered that it is mathematically more sound to simulate the centennially persistent SOC pool concentration at each experimental site as a constant. We used a Bayesian inference method to compute site-specific estimates of centennially persistent SOC concentration and associated uncertainties (detailed in the Section 2.3.1).

10
15 The proportion of centennially persistent SOC (CP_{SOC}) in each soil sample was then calculated as the ratio of the site-specific CP_{SOC} concentration to the total SOC concentration of the sample:

$$\text{CP}_{\text{SOC}} \text{ proportion [sample]} = \frac{\text{CP}_{\text{SOC}} \text{ concentration [site]}}{\text{SOC concentration [sample]}}, \quad (2)$$

20 where the unit of CP_{SOC} concentration [site] and SOC concentration [sample] is gC.kg^{-1} soil. The CP_{SOC} proportions of five samples that were slightly above 1 were set to 1. In these calculations, we assumed that at each site, the concentration of CP_{SOC} was the same in the LTBF and non-LTBF treatments and was constant with time. The details related to the estimation of the uncertainty on the CP_{SOC} proportion of each sample are reported in the Section 2.3.2.

2.2 Thermal analysis of soil samples by Rock-Eval 6

25 The 118 soil samples from the reference set were analyzed with a RE6 Turbo device (Vinci Technologies) using the basic set-up for the analysis of soil organic matter (Behar et al., 2001; Disnar et al., 2003). The RE6 technique provided measurements from the sequential pyrolysis and oxidation of *ca.* 40 mg of finely ground ($< 250 \mu\text{m}$) soil per sample (Fig. 1). Volatile hydrocarbon effluents from pyrolysis were detected and quantified with flame ionization detection (FID), while the evolution of CO and CO₂ gases ~~were~~ was quantified by infrared detection during both pyrolysis and oxidation stages. Pyrolysis was carried out from 200 °C to 650 °C in a N₂ atmosphere with a heating rate of 30 °C minute⁻¹, while the oxidation was carried out from 300 °C to 850 °C in a laboratory air atmosphere (with O₂) with a heating rate of 20 °C minute⁻¹. The RE6 technique generated five thermograms per sample (Fig. 1, *i.e.* volatile hydrocarbon (HC) effluents during pyrolysis, CO₂ during pyrolysis, CO₂ during oxidation, CO during pyrolysis, and CO during oxidation). On average, the organic carbon yield of the RE6 analysis was greater than 96.5% of SOC_{EA} for the soils of the reference sample set ($\text{SOC}_{\text{RE6}} = 0.966 \times \text{SOC}_{\text{EA}} + 0.403$, $R^2 = 0.97$, $n = 118$).

30
35 For each RE6 thermogram, we determined the temperatures corresponding to each incremental proportion of the amount of gases evolved during the pyrolysis and oxidation stages. Upper temperatures of 850 °C (CO oxidation thermogram), 650 °C

(HC pyrolysis thermogram), 611 °C (CO₂ oxidation thermogram) and 560 °C (CO and CO₂ pyrolysis thermograms) were chosen for signal integration (Fig. 1), thereby excluding any interference of soil carbonates (Behar et al., 2001). Thermal decomposition of carbonates was indeed observed beyond 560 °C (CO and CO₂ pyrolysis thermograms) and 611 °C (CO₂ oxidation thermogram) for the site of Grignon (data not shown). For each RE6 thermogram, signal integration was performed on the offset-corrected thermogram, using sample-specific offset values estimated by the RE6 Turbo device. For the three pyrolysis thermograms, signal integration started after an isotherm step of 200 s at 200 °C. Finally, we retained 5 of these temperature parameters per thermogram: T₁₀, T₃₀, T₅₀, T₇₀, T₉₀ which respectively represent the temperatures corresponding to the evolution of 10, 30, 50, 70, and 90% of the amount of evolved gases for each sample and for each of the five different thermograms (HC, CO₂ pyrolysis, CO₂ oxidation, CO pyrolysis, CO oxidation).

- 10 For the HC pyrolysis thermogram we also determined three parameters reflecting a proportion of thermally resistant or labile hydrocarbons: a parameter representing the proportion of hydrocarbons evolved between 200 °C and 450 °C (thermo-labile hydrocarbons, TLHC-index, modified from Saenger et al., 2015), the I-index representing the preservation of thermally labile immature hydrocarbons (after Sebag et al., 2016), and the R-index representing the proportion of hydrocarbons thermally stable at 400°C (after Sebag et al., 2016). Those three RE6 parameters were calculated as follows:

15

$$\text{TLHC-index} = \frac{\text{Area of HC pyrolysis thermogram [200 °C–450 °C]}}{\text{Total area of HC pyrolysis thermogram}}, \quad (3)$$

$$\text{I-index} = \log_{10} \left(\frac{\text{proportion of HC pyrolysis thermogram [200 °C–400 °C]}}{\text{proportion of HC pyrolysis thermogram [400 °C–460 °C]}} \right), \quad (4)$$

20
$$\text{R-index} = \frac{\text{Area of HC pyrolysis thermogram [400 °C–650 °C]}}{\text{Total area of HC pyrolysis thermogram}}, \quad (5)$$

- Using the HC pyrolysis thermogram, we determined a parameter reflecting SOC bulk chemistry, the hydrogen index (HI, mgHC.g⁻¹C), that corresponds to the quantity of pyrolyzed hydrocarbons relative to SOC_{RE6}. Using the CO and CO₂ pyrolysis thermograms, we determined another parameter reflecting SOC bulk chemistry, the oxygen index (OI_{RE6}, mgO₂.g⁻¹C) corresponding to the oxygen yield as CO and CO₂ during thermal pyrolysis of soil organic matter divided by the total SOC (SOC_{RE6}) of the sample. The HI correlates with the elemental H:C atomic ratio of SOC and the OI_{RE6} correlates with the elemental O:C atomic ratio of SOC (Espitalié et al., 1977).
- 25

Overall, we thus calculated for each soil sample a series of 30 RE6 parameters reflecting SOC thermal stability and bulk chemistry to be used in subsequent statistical and modelling analyses.

- 30 Signal integration of the RE6 thermograms and calculation of the RE6 temperature parameters were performed with R v.3.4.3 (R Core Team, 2017) and the hyperSpec (Beleites and Sergio, 2014), pracma (Borchers, 2015) and stringr (Wickham, 2015) packages.

2.3 Statistical analysis

2.3.1 Bayesian inference of site-specific CP_{SOC} concentrations and uncertainties

At each site, the CP_{SOC} concentration was estimated as the model parameter c of the exponential decay function described in Eq. (1). To estimate the value of this parameter and assess its uncertainty, we sampled the posterior Probability Density Function (PDF) of the model parameters in Eq. (1), which is given by Bayes' theorem as a function of the prior PDF (*i.e.* what we know before collecting data) and the likelihood (*i.e.* how likely is it to predict the data given a set of parameters). The posterior PDF is the combination of our prior knowledge and of the information carried by the data, including measurement uncertainties. For a model vector \mathbf{m} (containing the parameters a , b and c) and a data vector \mathbf{d} of all the measurements of SOC concentrations, the posterior PDF, $P(\mathbf{d} | \mathbf{m})$, is $P(\mathbf{d} | \mathbf{m}) \propto P(\mathbf{m})P(\mathbf{m} | \mathbf{d})$, with $P(\mathbf{m})$ the prior PDF on the model parameters and $P(\mathbf{m} | \mathbf{d})$ the likelihood.

We chose uniform PDFs for the model parameters, a , b , and c to be as uninformative as possible. We use the Gaussian form of the likelihood function, such as $P(\mathbf{m} | \mathbf{d}) \propto e^{-\frac{1}{2}(\mathbf{d} - \gamma(\mathbf{t}))^T \mathbf{C}_d^{-1}(\mathbf{d} - \gamma(\mathbf{t}))}$, where \mathbf{t} is the vector of all observation times and \mathbf{C}_d is the data covariance matrix describing the uncertainties on the SOC measurements. We consider a conservative standard deviation for SOC concentration data ($0.75 \text{ gC.kg}^{-1} \text{ soil}$) estimated by Barré et al. (2010) for the same soils. We use a Metropolis algorithm to draw 3×10^4 samples from the posterior PDF with a burning phase of 3.7×10^5 steps. We can then derive the mean and standard deviation for the parameter c from the posterior PDF.

2.3.2 Estimating the uncertainty of CP_{SOC} proportion in each sample

Based on our assessment of the uncertainties on SOC concentration data and site-specific CP_{SOC} concentrations (see above), we propagated these errors to estimate the uncertainty on the CP_{SOC} proportion in each soil sample. This was estimated by calculating the standard deviation (sd) of the CP_{SOC} proportion for each sample as follows:

sd (CP_{SOC} proportion [sample]) =

$$CP_{soc} \text{ proportion [sample]} \times \sqrt{\left(\frac{\text{sd}(CP_{soc} \text{ concentration [site]})}{CP_{soc} \text{ concentration [site]}}\right)^2 + \left(\frac{\text{sd}(SOC \text{ concentration [sample]})}{SOC \text{ concentration [sample]}}\right)^2},$$

(6)

~~where abbreviation sd stands for standard deviation.~~

2.3.3 Statistical relationships between RE6 parameters and CP_{SOC} proportion

The reference sample set was randomly split into a calibration set ($n = 88$ samples) and a validation set ($n = 30$ samples). The correlations between the 30 RE6 parameters and the CP_{SOC} proportion were assessed with a non-parametric Spearman's rank correlation test on the calibration set ($n = 88$). A principal component analysis (PCA) of the 30 centered and scaled RE6 parameters was performed for the calibration set to (i) summarize the variance of SOC thermal stability and bulk chemistry on a single factorial map, and (ii) illustrate the correlations among RE6 parameters. Correlations between the CP_{SOC} proportion in calibration soils and their principal component scores were determined using Spearman's rank correlation tests, and its relationships with the 30 RE6 parameters were further illustrated by projecting the CP_{SOC} proportion variable in the

PCA correlation plot. The RE6 data of the soils from the validation set were projected on the same PCA factorial map to check that the validation set was representative of the calibration set.

2.3.4 Random forests regression model to predict CP_{SOC} proportion from RE6 parameters

5 A multivariate regression model was built to relate CP_{SOC} proportion in the reference samples from the calibration set (response vector or dependent variable \mathbf{y} , $n = 88$) to their SOC thermal stability and bulk chemistry, summarized by a matrix of predictor variables (\mathbf{X}) made of the 30 centered and scaled RE6 parameters. The non-parametric and non-linear machine learning technique of random forests (RF, Breiman, 2001; Strobl et al., 2009) was used to build this model. The random forests regression model was based on a forest of 1000 diverse regression trees made of splits and nodes. A random forests learning algorithm combines bootstrap resampling and random variable selection. Each of the 1000 regression trees was thus
10 grown on a bootstrapped subset of the calibration set (*i.e.* containing about two thirds of “in-bag” calibration samples) by randomly sampling 10 out of the 30 RE6 parameters as candidates at each split of the tree, and using a minimum size of terminal tree nodes of five soil samples. The random forests regression model was then used to predict the proportion of CP_{SOC} in the validation set ($n = 30$), a prediction corresponding to the mean of the predicted values across the 1000 regression trees.

15 The performance of the random forests regression model for predicting CP_{SOC} proportion was assessed by statistics comparing the RF-predicted *vs.* reference (estimated) values of the sample set. The performance statistics were calculated on: (i) the “out-of-bag” soil samples of the calibration set and (ii) the soil samples of the validation set. Out-of-bag samples are observations from the calibration set not included in the learning sample set for a specific tree that can be used as a “built-in” test set for calculating its prediction accuracy (Strobl et al., 2009). The performance statistics included the
20 coefficient of determination (pseudo-R² for the calibration set or R² for the validation set) and the root-mean-square error of calibration or prediction (RMSEC for the calibration set or RMSEP for the validation set). The ratio of performance to deviation (RPD) and the bias of the random forests regression model were additionally calculated for the validation set. The relative importance (*i.e.* ranking) of each of the 30 RE6 parameters for the prediction of the proportion of CP_{SOC} in the RF regression model was computed as the unscaled permutation accuracy (Strobl et al., 2009).

25 Additionally, the sensitivity of the RF regression model to pedoclimate was assessed by examining its predictive performance for a calibration set based on soils from three sites (Versailles, Grignon, Rothamsted, $n = 84$) and a fully independent validation set based on soils from a different pedoclimate (Ultuna, $n = 34$) but with similar RE6 thermal characteristics to those of the calibration set (see Section 3.2). Soils from Ultuna have indeed a higher clay content (from 11% to 20% more clays), and experience a lower mean annual temperature (from 4 °C to 5.2 °C lower temperature), and a
30 lower mean annual precipitation (from 116 mm to 179 mm lower precipitation) than the soils of the three other sites (Supplementary material S1).

2.3.5 Error propagation in the random forests regression model

Since our objective was to deliver a model based on thermal analysis ~~based model~~ with reliable prediction intervals around the predicted values of the CP_{SOC} proportion, we estimated the prediction uncertainty of the random forests model for new
35 soil samples. We used a methodology recently published by Coulston et al. (2016) to approximate prediction uncertainty for random forests regression models, and adapted it to explicitly take into account the uncertainty on reference values of CP_{SOC} proportion (Eq. (6)) that were used to build the model (Supplementary material S2).

Briefly, we sampled with replacement (*i.e.* bootstrapped) the calibration set (\mathbf{y}, \mathbf{X}) 2000 times to obtain 2000 bootstrap samples ($\mathbf{y}^{*b}, \mathbf{X}^{*b}$) that were used to parametrize 2000 random forest models (\mathbf{RF}^{*b}). To incorporate the uncertainty on reference values of CP_{SOC} proportion, each of the 2000 bootstrapped vectors (\mathbf{y}^{*b}) contained values of CP_{SOC} proportion that were randomly sampled from normal distributions with means and standard deviations of the CP_{SOC} proportion of the corresponding soil samples from the calibration set (Eq. (6)). For each bootstrap sample of the calibration set, resampling discarded approximately 37% of the data ($\mathbf{y}^{*b}, \mathbf{X}^{*b}$) that were used for prediction. We obtained an error assessment dataset made of 2000 vectors of observed (reference) values \mathbf{y}^{*b} , predicted values $\hat{\mathbf{y}}^{*b}$ (mean of the predictions across 1000 regression trees for each observation), and $\text{var}(\hat{\mathbf{y}})^{*b}$ (variance of the predictions across 1000 regression trees for each observation). For each observation of the 2000 bootstrap samples, we calculated a metric τ allowing to scale between $\text{var}(\hat{\mathbf{y}})$ that can be calculated for any soil sample by the random forests regression model, and the squared prediction error $(\mathbf{y} - \hat{\mathbf{y}})^2$ that is only available for the reference sample set. The metric τ was calculated as follows (Coulston et al., 2016):

$$\tau = \sqrt{\frac{(\mathbf{y} - \hat{\mathbf{y}})^2}{\text{var}(\hat{\mathbf{y}})}}, \quad (7)$$

A Monte Carlo approach was used to estimate $\hat{\tau}$, the 95th percentile of all calculated τ values for all out-of-bag observations of the 2000 bootstraps (Supplementary material S2). This $\hat{\tau}$ value was such that 95% of the predictions of the CP_{SOC} proportion lie within $\hat{\tau} \times \text{sd}(\hat{\mathbf{y}})$ of the true value of CP_{SOC} proportion (*i.e.* 95% prediction intervals). As $\text{sd}(\hat{\mathbf{y}})$, the standard deviation of the predictions of the CP_{SOC} proportion across 1000 regression trees, can be calculated by the random forests regression model for any soil sample, this approach allows the calculation of 95% prediction intervals on any new soil sample for which only X data (30 RE6 parameters) are available. We calculated the 95% prediction intervals ($\hat{\mathbf{y}} \pm \hat{\tau} \times \text{sd}(\hat{\mathbf{y}})$) for the validation set ($n = 30$) to examine whether those intervals included the true (estimated) values of CP_{SOC} proportion. More details on the procedure to approximate prediction uncertainty for random forests regression models are provided in Coulston et al. (2016). We finally checked how the error on CP_{SOC} proportion propagated into the random forests regression model by (i) comparing the value of $\hat{\tau}$ with or without incorporating the uncertainty on reference values of CP_{SOC} proportion in the algorithm, and (ii) by comparing the sizes of the 95% prediction intervals calculated for the validation soil samples with their respective 95% confidence intervals (determined by multiplying their standard deviation calculated in Eq. (6) by 1.96).

The Bayesian inference method was performed with Python 2.7 and the PyMC library (Patil et al., 2010). All other statistical analyses were performed with R v.3.4.3 (R Core Team, 2017) and the factoextra package for running PCA (Kassambara, 2015), the randomForest package for running the random forests regression models (Liaw and Wiener, 2002) and the boot package for bootstrapping (Davison and Hinkley, 1997; Canty and Ripley, 2015).

30 3 Results

3.1 CP_{SOC} concentration at each site and CP_{SOC} proportion in reference soil samples

The Bayesian inference of the parameter c of the exponential decay function (Eq. (1)) yielded site-specific estimates of the CP_{SOC} concentration with 95% confidence intervals (Eq. (1), Table 1, Fig. 2). Estimated CP_{SOC} concentrations ranged from 6.22 gC.kg^{-1} soil at Versailles to 10.46 gC.kg^{-1} soil at Rothamsted. The uncertainty on CP_{SOC} concentration was lower at Rothamsted (standard deviation of 0.27 gC.kg^{-1} soil) and Versailles (standard deviation of 0.31 gC.kg^{-1} soil) than at Ultuna (standard deviation of 0.88 gC.kg^{-1} soil) and Grignon (standard deviation of 1.00 gC.kg^{-1} soil).

Overall, the wide range in total SOC concentrations within and across sites (from 5 to 46 gC.kg⁻¹ soil, Table 1) combined with an assumed constant CP_{SOC} concentration within each site, resulted in a reference sample set with a wide spectrum of CP_{SOC} proportions ranging from 0.14 to 1 (Eq. (2), Table 1). The uncertainty (standard deviation) on the values of CP_{SOC} proportion ranged from 0.01 to 0.15 for the reference sample set (Eq. (6), Supplementary material S3). High uncertainties were found for high values of CP_{SOC} proportion (*i.e.* samples with longer time periods under bare fallow treatment), with a modulation by the site-specific CP_{SOC} concentration uncertainty (Grignon > Ultuna > Versailles > Rothamsted, Table 1), as expected from Eq. (6) (Supplementary material S3).

The random splitting of the reference sample set generated calibration and validation sample sets with similar mean values, range of values and standard deviations for both total SOC concentration and CP_{SOC} proportion (Table 1).

10 3.2 Relationships between RE6 parameters and CP_{SOC} proportion

The 30 RE6 parameters showed contrasted correlations with the CP_{SOC} proportion in the calibration set (Table 2). Most RE6 temperature parameters showed positive correlations with the CP_{SOC} proportion, with Spearman's ρ above 0.8 for four of them (the RE6 temperature parameter corresponding to 50% of CO₂ gas evolution during the pyrolysis stage, T_{50_CO2_PYR}, and the RE6 temperature parameters corresponding to 30%, 50% and 70% of CO₂ gas evolution during the oxidation stage, T_{30_CO2_OX}, T_{50_CO2_OX}, T_{70_CO2_OX}, Table 2).

Conversely, five RE6 temperature parameters showed significant negative correlations with the CP_{SOC} proportion (T_{10_HC_PYR}, T_{10_CO_PYR}, T_{30_CO_PYR}, T_{50_CO_PYR}, T_{70_CO_PYR}, Table 2). ~~Out of t~~The three RE6 parameters reflecting a proportion of thermally resistant or labile hydrocarbons (~~only the TLHC-index, showed a weakly significant negative Spearman's ρ with the CP_{SOC} proportion, the I-index, and the R-index~~) ~~showed~~ no correlations with the CP_{SOC} proportion (Table 2). The two RE6 parameters reflecting SOC bulk chemistry showed highly significant correlations with the CP_{SOC} proportion (Table 2), the HI being negatively correlated and the OI_{RE6} being positively correlated.

The PCA of the centered and scaled RE6 parameters illustrates the correlations among those 30 variables in the calibration set (Fig. 3). A continuum of CP_{SOC} proportion values was observed in the reference samples along the first two principal components (Fig. 3A), and projecting the CP_{SOC} proportion in the PCA correlation circle further highlighted the relationships between this variable and the 30 RE6 parameters (Fig. 3B). The CP_{SOC} proportion variable had a strongly negative projected loading score on PC1 (Fig. 3B), as well as negative projected loadings on PC2 (Fig. 3B) and PC3 (data not shown). The scores of the calibration soils on the first three principal components were indeed significantly and negatively correlated with the CP_{SOC} proportion ($\rho = -0.671$, p -value < 0.001 for PC1, $\rho = -0.4936$, p -value < 0.001 for PC2, $\rho = -0.25$, p -value < 0.05 for PC3), such that a large part (82%) of the variance of the 30 RE6 parameters was linked to the CP_{SOC} proportion in the calibration set.

The random splitting of the reference sample set generated calibration and validation sample sets with similar RE6 thermal characteristics as illustrated by their similar distribution on the factorial map of the first two principal components of the PCA (Fig. 3A). Soils from the site of Grignon (with carbonates) showed specific RE6 thermal characteristics (Fig. 3A). Some soils from the sites of Rothamsted and Versailles with high CP_{SOC} proportions also showed specific RE6 thermal signatures (Fig. 3A). Conversely, all soils from the site of Ultuna showed similarities regarding their RE6 thermal characteristics with certain soil samples from other sites (Versailles and Rothamsted, Fig. 3A).

3.3 Performance of the regression model using RE6 parameters to predict CP_{SOC} proportion

The random forests regression model performed very well in predicting the CP_{SOC} proportion in the reference sample set using the 30 RE6 parameters as predictors (Fig. 4). Both performance statistics on the calibration set (pseudo-R² = 0.9489, RMSEC = 0.0607, n = 88) and on the validation set (R² = 0.9492, RMSEP = 0.07, n = 30) demonstrated the good predictive power of the regression model based on RE6 thermal analysis. The predictive performance of the random forests model based on RE6 thermal analysis (RE6-RF) was altered when soil samples from a pedoclimate different (site of Ultuna, n = 34) to the calibration set (Versailles, Rothamsted, Grignon), but with similar RE6 thermal characteristics (see Section 3.2), were used for validation (Fig. 5). The coefficient of determination of the model decreased (R² = 0.53), yet its mean error of prediction did not increase strongly (RMSEP = 0.10, Fig. 5).

Propagating the estimated uncertainties on the values of CP_{SOC} proportion increased the size of the prediction intervals of RE6-RF regression model. Indeed, the value of \hat{t} increased from 1.72–83 to 2.40–12 when the uncertainty on CP_{SOC} proportion was integrated in the algorithm described at Section 2.3.5. The horizontal and vertical error bars on Fig. 4 illustrate the global error propagation on the CP_{SOC} proportion estimates in the RE6-RF regression model for the validation soil sample set. The values of the total width of the 95% confidence interval (reference estimations of CP_{SOC} proportion, horizontal error bars in Fig. 4) were 0.03 (minimum total width), 0.58 (maximum total width) and 0.24 (mean total width) for the soil samples of the validation set (n = 30). For the 95% prediction intervals (RE6-RF predictions of CP_{SOC} proportion, vertical error bars in Fig. 4), the uncertainties increased to 0.44–11 (minimum total width), 0.67–66 (maximum total width) and 0.35–37 (mean total width). The thirty 95% prediction intervals for RE6-RF predictions of CP_{SOC} proportion in the validation set all included their respective reference estimation of CP_{SOC} proportion (Fig. 4).

Out of the 30 RE6 parameters tested by the random forests model as possible predictor variables of the CP_{SOC} proportion in the calibration set, the RE6 temperature parameters corresponding to 50% and 70% of CO₂ gas evolution during the pyrolysis stage (T_{50_CO2_PYR}, T_{70_CO2_PYR}) and to 30% of CO₂ gas evolution during the oxidation stage (T_{30_CO2_OX}) showed the highest importance scores (based on permutation accuracy importance calculations, Table 2). The twelve-eight most important RE6 parameters for predicting the CP_{SOC} proportion were temperature parameters calculated on the five different RE6 thermograms (Table 2). The two RE6 parameters reflecting SOC bulk chemistry (OI_{RE6} and HI) were of medium importance to predict the CP_{SOC} proportion, while the RE6 parameters reflecting a proportion of thermally resistant or labile hydrocarbons (I-index, R-index and TLHC-index) were of weak importance (Table 2).

4 Discussion

4.1 A unique soil sample set with accurate and contrasted values of CP_{SOC}

Adding new SOC concentration data for Rothamsted (up to 2014) and Ultuna (up to 2015), and combining SOC concentration data from all LTBF field replicate plots of each site decreased the uncertainty on the site-specific estimates of the CP_{SOC} concentration (Fig. 2), compared with the previous estimations published by Barré et al. (2010). Indeed, the total width of the 95% confidence interval around the estimation of the site-specific CP_{SOC} concentration slightly decreased from 1.4 to 1.2 gC.kg⁻¹ soil at Versailles and from 4.96 to 3.92 gC.kg⁻¹ soil at Grignon, and strongly decreased from 7.24 to 3.46 gC.kg⁻¹ soil at Ultuna and from 5.98 to 1.06 gC.kg⁻¹ soil at Rothamsted (Table 1, Fig. 2, Barré et al., 2010). The mean estimated values of the CP_{SOC} concentration were marginally changed at Versailles (6.22 vs. 6.12 gC.kg⁻¹ soil in Barré et al., 2010) and Grignon (7.12 vs. 6.80 gC.kg⁻¹ soil in Barré et al., 2010), but strongly modified (increased) at Ultuna (6.95 vs. 3.90 gC.kg⁻¹ soil in Barré et al., 2010) and Rothamsted (10.46 vs. 2.72 gC.kg⁻¹ soil in Barré et al., 2010, Table 1).

Our results obtained under four contrasted pedoclimates of Northwestern Europe indicate a minimum value of 5 gC.kg⁻¹ soil (lowest boundary of the four 95% confidence intervals, Table 1) and a maximum value of 11 gC.kg⁻¹ soil (highest boundary of the four 95% confidence intervals, Table 1) for CP_{SOC} concentration in topsoils (0–20 to 0–25 cm depth). These estimates are close, yet below the CP_{SOC} concentration value of 12 gC.kg⁻¹ soil estimated by Buyanovsky and Wagner (1998b) for the topsoil (0–20 cm depth) of the Sanborn long-term (100 years) agronomic experiment (Columbia, Missouri, USA). Our estimates of topsoil CP_{SOC} concentration are also well below the value of 16 gC.kg⁻¹ soil estimated by Franko and Merbach (2017) in the topsoil (0–30 cm depth) of the long-term (28 years) bare fallow experiment of Bad Lauchstädt (Central Germany). The soil type in Bad Lauchstädt (Haplic Chernozem) and its high concentration of slow-cycling black carbon (estimated at 2.5 gC.kg⁻¹ soil, Brodowski et al., 2007) may explain this difference, as well as the relatively short time period under bare fallow (higher uncertainty on the inferred CP_{SOC} concentration).

Among the wide range of CP_{SOC} proportions (0.14 to 1) of our reference sample set, high values of CP_{SOC} proportions (> 0.6) were obtained only for soils which had been under bare fallow for a long period of time: after several years or decades with negligible C inputs and sustained SOC decomposition (Table 1). Similarly, the low values of CP_{SOC} proportions (< 0.25) of our reference sample set were obtained for soils without vegetation but receiving high amounts of manure amendments at Versailles (Table 1). It could be argued that CP_{SOC} proportion values obtained for bare soils with or without organic matter amendments may not be representative of CP_{SOC} proportions of soils under conventional management practices. However, it is interesting to notice that soils of the reference sample set with vegetation and experiencing classical management practices (grassland at Rothamsted, cropland at Ultuna) also showed a wide range of CP_{SOC} proportions, from 0.25 to 0.56 (Table 1). Moreover, other studies have shown the high variability of CP_{SOC} proportion in soils. For instance, Falloon et al. (1998) listed a series of published values of CP_{SOC} proportions ranging from 0.13 to 0.59. More recently, Mills et al. (2014) published a large dataset of CP_{SOC} proportions in uncultivated topsoils (*ca.* 15 cm depth). They estimated CP_{SOC} proportions using a global dataset of topsoil radiocarbon (¹⁴C) data and a steady-state SOC turnover model with a fixed mean residence time of 1000 years for persistent SOC. Their estimates of CP_{SOC} proportions varied greatly from 0.03 to 0.98 (mean = 0.48, standard deviation = 0.22, n = 232, soils with inconsistent negative modeled SOC pools values were removed), with significantly higher CP_{SOC} proportions in non-forest than in forest uncultivated ecosystems (Mills et al., 2014).

Overall, those combined results illustrate the wide range of CP_{SOC} concentrations and proportions in topsoils that may depend upon pedoclimate, land-use and management practices. ~~Additionally, these results show the value of LTBF experiments to investigate the long-term dynamics of SOM.~~

4.2 A quantitative link between the long-term biogeochemical stability of SOC and its thermal stability and bulk chemistry

This work ~~reinforces the evidence that there is a link between~~ ~~strengthens the existence of a link between~~ SOC persistence in ecosystems and its thermal stability, providing evidence of the first quantitative link between thermal and *in-situ* long-term (> 100 years) biogeochemical SOC stability (Fig. 4). The regression model yields accurate RE6-RF predictions of CP_{SOC} proportions with 95% prediction intervals that fully propagate the uncertainties originating from the calibration set that was used to build the model. Predictions on the validation set illustrate that the error propagation scheme provides highly conservative 95% prediction intervals of the CP_{SOC} proportion in new samples, all intervals including their respective reference estimate of CP_{SOC} proportion (Fig. 4). Despite rather large prediction intervals, the RE6-RF regression model clearly discriminates soils with small CP_{SOC} proportions from samples with large CP_{SOC} proportions (Fig. 4). This model based on RE6 thermal analysis can thus be used to predict the size of the CP_{SOC} pool with known uncertainty in new soil samples from similar pedoclimates and with thermal characteristics (*i.e.* RE6 predictor variables) similar to those of the reference sample set.

Our results also highlight the sensitivity of the RE6-RF regression model to pedoclimate. Decreased predictive performance of the model (as assessed by the coefficient of determination) was indeed observed when predicting the CP_{SOC} proportion in new soils with similar RE6 thermal characteristics but from a different pedoclimate (Fig. 5). However, the mean error of prediction of the model only slightly increased when predicting the CP_{SOC} proportion in soils from the fully independent site of Ultuna (Fig. 5). Overall, those results illustrate the potential of the model based on RE6 thermal analysis to predict the proportion of CP_{SOC} in new soil samples from different pedoclimates, at least for sites that have similar RE6 thermal characteristics to those of the calibration set.

Our results also illustrate the complex relationships between thermal analysis-based parameters of SOC stability and the CP_{SOC} proportion. The hypothesis behind the use of SOC thermal stability as a proxy of its biogeochemical stability implies positive correlations between the size of the CP_{SOC} pool and temperature parameters derived from thermal analysis such as the 25 RE6 temperature parameters calculated in this study. Significant positive correlations with the CP_{SOC} proportions were indeed found for the majority (15-14 out of 25) of RE6 temperature parameters, with very high and positive Spearman's ρ values for some of them (Table 2). This was notably the case of the RE6 temperature parameter corresponding to 50% of CO₂ gas evolution during the oxidation stage, T_{50-CO₂-OX} that had been previously shown to systematically increase with bare fallow duration on the same soils by Barré et al. (2016). This study extends the results of Barré et al. (2016) towards a quantitative link between RE6 temperature parameters and SOC persistence (direct correlations and predictions of the size of the CP_{SOC} pool rather than time under bare fallow treatment). It also extends those results to non-bare fallow soils: bare soils receiving organic amendments (at Grignon and Versailles), cropland soils (Ultuna) and grassland soils (Rothamsted). Conversely, ten-11 RE6 temperature parameters showed no significant correlation or significant negative correlations with the CP_{SOC} proportion. Weak or negative correlations occurred principally for temperature parameters calculated on thermograms of the pyrolysis stage of the RE6 analysis: for all-parameters of the HC and CO thermograms (except T_{90-HC-PYR}) and lowest temperature parameters of the HC and CO₂ thermograms (Table 2). Negative correlations contradict the above-mentioned hypothesis, with the evolution of a similar proportion of the total amount of gases (HC pyrolysis effluents or CO) occurring at lower temperatures for samples with high CP_{SOC} proportions than for soils with low CP_{SOC} proportions. A possible explanation for this unexpected observation could be that the pyrolysis of SOC in samples with high proportion of CP_{SOC} may undergo an enhanced pyrolysis catalyst effect by soil minerals (Auber, 2009), which are relatively more abundant in those samples generally characterized by low total SOC concentrations.

Despite the fact that three RE6 parameters used here, i.e. the TLHC-index, the I-index, and the R-index, had originally been proposed as useful qualitative metrics of soil carbon dynamics, reflecting a proportion of thermally resistant or labile hydrocarbons (Disnar et al., 2003; Sebag et al., 2006; Saenger et al., 2013, 2015; Sebag et al., 2016), those parameters were weakly correlated (TLHC index) or not correlated (I index, R index) to the CP_{SOC} proportion. Furthermore, they also had a weak importance in the random forests model predictions of the CP_{SOC} proportion (Table 2). The poor link between those three RE6 parameters and the CP_{SOC} proportion may be explained by the high residence time of CP_{SOC} (> 100 years). Indeed, so far those parameters have been related to the proportion of SOC present in the particulate organic matter fraction (size > 50 μ m, density < 1-1.6), a SOC pool characterized by a residence time in soils generally below 20 years (Saenger et al., 2015; Soucémarianadin et al., 2018).

The two RE6 parameters reflecting SOC bulk chemistry showed highly significant correlations with the CP_{SOC} proportion. This confirms, and extends to vegetated soils, the observed decreasing trend for HI and increasing trend for OI_{RE6} (except at Versailles where soils have high pyrogenic C content which has high proportions of pyrogenic carbon) with bare fallow duration observed by Barré et al. (2016) on the bare fallow treatments of the same experimental sites. Soils with high proportions of CP_{SOC} are thus characterized by an oxidized and H-depleted organic matter.

4.3 Perspectives to improve and foster RE6 thermal analysis based predictions of the size of the CP_{SOC} pool

Future developments of this work must extend the Rock-Eval 6 thermal analysis regression model to a wider range of pedoclimates and to other biomes. As sites with LTBF treatments are not widespread, complementing the reference sample set may be achieved by using soils that have different soil forming factors (*e.g.* climate, parent material) and (i) which are sampled from long-term (> 50 to 100 years) experiments with contrasted SOC inputs, enabling the estimation of their CP_{SOC} concentration (Buyanovsky and Wagner, 1998a; 1998b), or (ii) for which the mean SOC age is known from radiocarbon data, enabling the estimation of the size of their persistent SOC pool (Trumbore, 2009; Mills et al., 2014).

Another development of this work will involve elucidating the fundamental mechanisms linking the biogeochemical stability of SOC with its thermal stability (*see e.g.* Leifeld and von Lützow, 2014). This was beyond the scope of this work, yet it ~~remains-constitutes~~ an exciting field of research that should be addressed in the future, as highlighted by the unexpected observations discussed in Section 4.2 and by other recent works that found no relationships between the thermal oxidation of SOC between 200 °C and 400 °C and the size of SOC pools with shorter residence times in soils (below or above *ca.* 18 years, Schiedung et al., 2017).

Overall, this work demonstrates the value of Rock-Eval 6 as a routine method for quantifying the size of the centennially persistent SOC pool with known uncertainty in temperate soils. The relatively low cost of the Rock-Eval 6 technique and the robustness of the thermal analysis regression model makes it possible to apply it to soil monitoring networks across a continuum of scales, as a reliable proxy of SOC persistence. This may be part of the framework proposed by O'Rourke et al. (2015) to better understand SOC processes at the biosphere to biome scales, and should be added to the soil carbon cycling proxies recently listed by Bailey et al. (*in press*2018). Mapping persistent SOC at large scales may allow the identification of regional hotspots of centennially persistent SOC that may contribute little to climate change by 2100. It may also provide information on the sustainability of additional SOC storage from soil carbon sequestration strategies such as those promoted by the international 4 per 1000 initiative in agriculture and forestry (<https://www.4p1000.org/>; Dignac et al., 2017; Minasny et al., 2017; Soussana et al., *in press*). A global map of centennially persistent SOC based on this empirical RE6 thermal analysis model could also be useful for improving the parameterization of ~~models simulating~~ SOC dynamics in Earth system models (Falloon and Smith, 2000; Luo et al., 2014; He et al., 2016). Indeed, this model based on RE6 thermal analysis appears as a robust and operational alternative to existing techniques used to initialize the size of the CP_{SOC} pool in models of SOC dynamics (such as the methods of Falloon et al. (1998) or Zimmermann et al. (2007) that estimate the size of the inert SOC pool in the RothC model). The integration of large-scale information on the size of SOC kinetic pools may ~~indeed~~ provide an adequate complement to the global data sets on SOC fluxes that are currently under development and restructuring (Hashimoto et al., 2015; Luo et al., 2016; Harden et al., *in press*2018).

Author contribution

LC and PB designed the study. PB and CC collected the archived soils. TK, SH, FvO and AM provided the archived soils and associated metadata. FB and FS performed the RE6 thermal analyses. RJ wrote the Python code. LC wrote the R codes and performed all statistical analysis. All authors contributed to the interpretation of the results. LC prepared the manuscript with contributions from all co-authors.

Acknowledgements

This work was funded by ADEME and EC2CO (CARACAS project). Pierre Barré, Lauric Cécillon, Laure Soucémariadin and Suzanne Lutfalla thank the financial support of the Mairie de Paris (Emergences Programme). We thank Thomas Eglin (ADEME), [Nicolas Bouton and Jean Espitalié \(Vinci Technologies\)](#), [David Sebag \(Normandie Univ., University of Lausanne\)](#) and two anonymous reviewers for ~~their~~ valuable suggestions on this work. We thank INRA and AgroParisTech for access to and maintenance of the Versailles 42 plots and the Grignon 36 plots long-term experiments and their sample archives. We thank Rothamsted Research and the Lawes Agricultural Trust for access to archived samples and the BBSRC for support under the Institute National Capabilities program grant (BBS/E/C/000J0300). We gratefully acknowledge the Faculty of Natural Resources and Agricultural Sciences [of the Swedish University of Agricultural Sciences \(SLU\)](#) for providing funds for maintenance of the Ultuna long-term field experiment and for its sample archive.

Supplementary material

(see Supplementary material manuscript)


References

- Amundson, R.: The carbon budget of soils. *Annu. Rev. Earth Planet. Sci.*, 29, 535–562, 2001.
- 15 Auber, M.: Effet catalytique de certains inorganiques sur la sélectivité des réactions de pyrolyse rapide de biomasses et de leurs constituants. PhD Thesis, Institut National Polytechnique de Lorraine, 296 p, 2009.
- Bailey, V.L., Bond-Lamberty, B., DeAngelis, K., Grandy, A.S., Hawkes, C.V., Heckman, K., Lajtha, K., Phillips, R.P., Sulman, B.N., Todd-Brown, K.E.O., and Wallenstein, M.D.: Soil carbon cycling proxies: Understanding their critical role in predicting climate change feedbacks. *Glob. Change Biol.*, [24, 895–905, 2018doi:10.1111/gcb.13926, in press.](#)
- 20 Barré, P., Plante, A.F., Cécillon, L., Lutfalla, S., Baudin, F., Bernard, S., Christensen, B.T., Eglin, T., Fernandez, J.M., Houot, S., Kätterer, T., Le Guillou, C., Macdonald, A., van Oort, F., and Chenu, C.: The energetic and chemical signatures of persistent soil organic matter. *Biogeochemistry* 130, 1–12, 2016.
- Barré, P., Eglin, T., Christensen, B.T., Ciais, P., Houot, S., Kätterer, T., van Oort, F., Peylin, P., Poulton, P.R., Romanenkov, V., and Chenu, C.: Quantifying and isolating stable soil organic carbon using long-term bare fallow experiments. *Biogeosciences*, 7, 3839–3850, 2010.
- 25 Behar, F., Beaumont, V., and De B. Penteadó, H.L.: Rock-Eval 6 technology: performances and developments. *Oil Gas Sci. Technol.*, 56, 111–134, 2001.
- Beleites, C. and Sergo, V.: hyperSpec: a package to handle hyperspectral data sets in R. R package version 0.98-20140523, 2014.
- 30 Bispo, A., Andersen, L., Angers, D., Bernoux, M., Brossard, M., Cécillon, L., Comans, R., Harmsen, J., Jonassen, K., Lamé, F., Lhuillery, C., Maly, S., Martin, E., Mcelnea, A., Sakai, H., Watabe, Y., and Eglin, T.: Accounting for carbon stocks in soils and measuring GHGs emission fluxes from soils: do we have the necessary standards? *Front. Environ. Sci.* 5, 41, 2017.
- Borchers, H.W.: pracma: Practical Numerical Math Functions. R package version 1.8.8, 2015.
- Breiman, L.: Random forests. *Mach. Learn.*, 45, 5–32, 2001.
- 35 Brodowski, S., Amelung, W., Haumaier, L., and Zech, W.: Black carbon contribution to stable humus in German arable soils. *Geoderma*, 139, 220–228, 2007.
- Bruun, S., Thomsen, I.K., Christensen, B.T., and Jensen, L.S.: In search of stable soil organic carbon fractions: a comparison of methods applied to soils labelled with ¹⁴C for 40 days or 40 years. *Eur. J. Soil Sci.*, 59, 247–256, 2008.

- Buyanovsky, G.A. and Wagner, G.H.: Carbon cycling in cultivated land and its global significance. *Glob. Change Biol.*, 4, 131–141, 1998a.
- Buyanovsky, G.A. and Wagner, G.H.: Changing role of cultivated land in the global carbon cycle. *Biol. Fertil. Soils*, 27, 242–245, 1998b.
- 5 Campo, J. and Merino, A.: Variations in soil carbon sequestration and their determinants along a precipitation gradient in seasonally dry tropical forest ecosystems. *Glob. Change Biol.*, 22, 1942–1956, 2016.
- Canty, A. and Ripley, B.: *boot: Bootstrap R (S-Plus) Functions*. R package version 1.3-17, 2015.
- Coulston, J.W., Blinn, C.E., Thomas, V.A., and Wynne, R.H.: Approximating prediction uncertainty for random forest regression models. *Photogramm. Eng. Remote Sens.*, 82, 189–197, 2016.
- 10 Davison, A.C. and Hinkley, D.V.: *Bootstrap Methods and Their Applications*. Cambridge University Press, Cambridge. ISBN 0-521-57391-2, 1997.
- Dignac, M.F., Derrien, D., Barré, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T., Freschet, G.T., Garnier, P., Guenet, B., Hedde, M., Klumpp, K., Lashermes, G., Maron, P.A., Nunan, N., Roumet, C., and Basile-Doelsch, I.: Increasing soil carbon storage: mechanisms, effects of agricultural practices and proxies. A review. *Agron. Sustain. Dev.*, 37, 14, 2017.
- 15 Disnar, J.R., Guillet, B., Keravis, D., Di-Giovanni, C., and Sebag, D.: Soil organic matter (SOM) characterization by Rock-Eval pyrolysis: scope and limitations. *Org. Geochem.*, 34, 327–343, 2003.
- Eglin, T., Ciais, P., Piao, S.L., Barré, P., Bellassen, V., Cadule, P., Chenu, C., Gasser, T., Koven, C., Reichstein, M., and Smith, P.: Historical and future perspectives of global soil carbon response to climate and land-use changes. *Tellus Ser. B-Chem. Phys. Meteorol.*, 62, 700–718, 2010.
- 20 Espitalié, J., Madec, M., Tissot, B., Mennig, J.J., and Leplat, P.: Source rock characterization method for petroleum exploration. 9th annual offshore technology conference, May 2-5 1977, Houston, USA, OTC 2935, DOI: 10.4043/2935-MS, 1977.
- Falloon, P. and Smith, P.: Modelling refractory soil organic matter. *Biol. Fertil. Soils*, 30, 388–398, 2000.
- Falloon, P., Smith, P., Coleman, K., and Marshall, S.: Estimating the size of the inert organic matter pool from total soil organic carbon content for use in the Rothamsted carbon model. *Soil Biol. Biochem.*, 30, 1207–1211, 1998.
- 25 Feng, W., Shi, Z., Jiang, J., Xia, J., Liang, J., Zhou, J., and Luo, Y.: Methodological uncertainty in estimating carbon turnover times of soil fractions. *Soil Biol. Biochem.*, 100, 118–124, 2016.
- Franko, U. and Merbach, I.: Modelling soil organic matter dynamics on a bare fallow Chernozem soil in Central Germany. *Geoderma*, 303, 93–98, 2017.
- 30 Gregorich, E.G., Gillespie, A.W., Beare, M.H., Curtin, D., Sanei, H., and Yanni, S.F.: Evaluating biodegradability of soil organic matter by its thermal stability and chemical composition. *Soil Biol. Biochem.*, 91, 182–191, 2015.
- Harden, J.W., Hugelius, G., Ahlström, A., Blankinship, J.C., Bond-Lamberty, B., Lawrence, C.R., Loisel, J., Malhotra, A., Jackson, R.B., Ogle, S., Phillips, C., Ryals, R., Todd-Brown, K., Vargas, R., Vergara, S.E., Cotrufo, M.F., Keiluweit, M., Heckman, K.A., Crow, S.E., Silver, W.L., DeLonge, M., and Nave, L.E.: Networking our science to characterize the state, vulnerabilities, and management opportunities of soil organic matter. *Glob. Change Biol.*, 24, [e705–e718](https://doi.org/10.1111/geb.13896), 2018doi: 10.1111/geb.13896, in press.
- 35 Hashimoto, S., Carvalhais, N., Ito, A., Migliavacca, M., Nishina, K., and Reichstein, M.: Global spatiotemporal distribution of soil respiration modeled using a global database. *Biogeosciences*, 12, 4121–4132, 2015.
- He, Y., Trumbore, S.E., Torn, M.S., Harden, J.W., Vaughn, L.J.S., Allison, S.D., and Randerson, J.T.: Radiocarbon constraints imply reduced carbon uptake by soils during the 21st century. *Science*, 353, 1419–1424, 2016.
- 40 Kassambara, A.: *factoextra: extract and visualize the results of multivariate data analyses*. R package version 1.0.3, 2015.
- Leifeld, J. and von Lütow, M.: Chemical and microbial activation energies of soil organic matter decomposition. *Biol. Fertil. Soils*, 50, 147–153, 2014.
- Liaw, A. and Wiener, M.: Classification and Regression by randomForest. *R News*, 2, 18–22, 2002.

- Luo, Y., Ahlström, A., Allison, S.D., Batjes, N.H., Brovkin, V., Carvalhais, N., Chappell, A., Ciais, P., Davidson, E.A., Finzi, A., Georgiou, K., Guenet, B., Hararuk, O., Harden, J.W., He, Y., Hopkins, F., Jiang, L., Koven, C., Jackson, R.B., Jones, C.D., Lara, M.J., Liang, J., McGuire, A.D., Parton, W., Peng, C., Randerson, J.T., Salazar, A., Sierra, C.A., Smith, M.J., Tian, H., Todd-Brown, K.E.O., Torn, M., van Groenigen, K.J., Wang, Y.P., West, T.O., Wei, Y., Wieder, W.R., Xia, J., Xu, X., Xu, X., and Zhou, T.: Toward more realistic projections of soil carbon dynamics by Earth system models. *Glob. Biogeochem. Cycle*, 30, doi:10.1002/2015GB005239, 2016.
- Luo, Z.K., Wang, E.L., Fillery, I.R.P., Macdonald, L.M., Huth, N., and Baldock, J.: Modelling soil carbon and nitrogen dynamics using measurable and conceptual soil organic matter pools in APSIM. *Agric. Ecosyst. Environ.*, 186, 94–104, 2014.
- 10 Lutfalla, S., Chenu, C., and Barré, P.: Are chemical oxidation methods relevant to isolate a soil pool of centennial carbon? *Biogeochemistry*, 118, 135–139, 2014.
- Mathieu, J.A., Hatté, C., Balesdent, J., and Parent, E.: Deep soil carbon dynamics are driven more by soil type than by climate: a worldwide meta-analysis of radiocarbon profiles. *Glob. Change Biol.*, 21, 4278–4292, 2015.
- Menichetti, L., Kätterer, T., and Leifeld, J.: Parametrization consequences of constraining soil organic matter models by total carbon and radiocarbon using long-term field data. *Biogeosciences*, 13, 3003–3019, 2016.
- 15 Mills, R.T.E., Tipping, E., Bryant, C.L., and Emmett, B.A.: Long-term organic carbon turnover rates in natural and semi-natural topsoils. *Biogeochemistry*, 118, 257–272, 2014.
- Minasny, B., Malone, B.P., McBratney, A.B., Angers, D.A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.S., Cheng, K., Das, B.S., Field, D.J., Gimona, A., Hedley, C.B., Hong, S.Y., Mandal, B., Marchant, B.P., Martin, M., McConkey, B.G., Mulder, V.L., O'Rourke, S., Richer-de-Forges, A.C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C.C., Vågen, T.G., van Wesemael, B., and Winowiecki, L.: Soil carbon 4 per mille. *Geoderma*, 292, 59–86, 2017.
- NF ISO 10694: Determination of organic and total carbon after dry combustion (elementary analysis). AFNOR, 1995.
- O'Rourke, S.M., Angers, D.A., Holden, N.M., and McBratney, A.B.: Soil organic carbon across scales. *Glob. Change Biol.*, 21, 3561–3574, 2015.
- 25 Patil, A., Huard, D., and Fonnesbeck, C.J.: PyMC: Bayesian stochastic modelling in Python. *J. Stat. Softw.*, 35, 1–81, 2010.
- Plante, A.F., Beupré, S.R., Roberts, M.L., and Baisden, T.: Distribution of radiocarbon ages in soil organic matter by thermal fractionation. *Radiocarbon*, 55, 1077–1083, 2013.
- Plante, A.F., Fernández, J.M., Haddix, M.L., Steinweg, J.M., and Conant, R.T.: Biological, chemical and thermal indices of soil organic matter stability in four grassland soils. *Soil Biol. Biochem.*, 43, 1051–1058, 2011.
- 30 Plante, A.F., Fernández, J.M., and Leifeld, J.: Application of thermal analysis techniques in soil science. *Geoderma*, 153, 1–10, 2009.
- Post, W.M. and Kwon, K.C.: Soil carbon sequestration and land-use change: processes and potential. *Glob. Change Biol.*, 6, 317–327, 2000.
- 35 R Core Team: R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing, 2017.
- Rovira, P., Kurz-Besson, C., Coîteaux, M.M., and Ramón Vallejo, V.: Changes in litter properties during decomposition: A study by differential thermogravimetry and scanning calorimetry. *Soil Biol. Biochem.*, 40, 172–185, 2008.
- Rühlmann, J.: A new approach to estimating the pool of stable organic matter in soil using data from long-term field experiments. *Plant Soil*, 213, 149–160, 1999.
- 40 Saenger, A., Cécillon, L., Poulénard, J., Bureau, F., De Danieli, S., Gonzalez, J.M., and Brun, J.J.: Surveying the carbon pools of mountain soils: A comparison of physical fractionation and Rock-Eval pyrolysis. *Geoderma*, 241–242, 279–288, 2015.

- Saenger, A., Cécillon, L., Sebag, D., and Brun, J.J.: Soil organic carbon quantity, chemistry and thermal stability in a mountainous landscape: A Rock-Eval pyrolysis survey. *Org. Geochem.*, 54, 101–114, 2013.
- Sanderman, J., Baisden, W.T., and Fallon, S.: Redefining the inert organic carbon pool. *Soil Biol. Biochem.*, 92, 149–152, 2016.
- 5 Schiedung, M., Don, A., Wordell-Dietrich, P., Alcántara, V., Kuner, P., and Guggenberger, G.: Thermal oxidation does not fractionate soil organic carbon with differing biological stabilities. *J. Plant Nutr. Soil Sci.*, 180, 18–26, 2017.
- Schrumpf, M. and Kaiser, K.: Large differences in estimates of soil organic carbon turnover in density fractions by using single and repeated radiocarbon inventories. *Geoderma*, 239–240, 168–178, 2015.
- Sebag, D., Verrecchia, E.P., Cécillon, L., Adatte, T., Albrecht, R., Aubert, M., Bureau, F., Cailleau, G., Copard, Y., Decaens, T., Disnar, J.R., Hetényi, M., Nyilas, T., and Trombino, L.: Dynamics of soil organic matter based on new Rock-Eval indices. *Geoderma*, 284, 185–203, 2016.
- 10 Sebag, D., Disnar, J.R., Guillet, B., Di Giovanni, C., Verrecchia, E.P., and Durand, A.: Monitoring organic matter dynamics in soil profiles by “Rock-Eval pyrolysis”: bulk characterization and quantification of degradation. *Eur. J. Soil Sci.*, 57, 344–355, 2006.
- 15 Soil Carbon Initiative: Outcomes Document Soil Carbon Summit. Dow Sustainability Fund and ALCOA, 11p, 2011.
- Soucémarianadin, L., Cécillon, L., Chenu, C., Baudin, F., Nicolas, M., Girardin, C., and Barré, P.: Is Rock-Eval 6 thermal analysis a good indicator of soil organic carbon lability? – A method comparison study in forest soils. *Soil Biol. Biochem.*, 117, 108–116, 2018.
- Soussana, J.F., Luffalla, S., Ehrhardt, F., Rosenstock, T., Lamanna, C., Havlík, P., Richards, M., Wollenberg, E.L., Chotte, J.L., Torquebiau, E., Ciais, P., Smith, P., and Lal, R.: Matching policy and science: Rationale for the ‘4 per 1000 - soils for food security and climate’ initiative. *Soil Tillage Res.*, doi: 10.1016/j.still.2017.12.002, in press.
- 20 Strobl, C., Malley, J., and Tutz, G.: An introduction to recursive partitioning: rationale, application and characteristics of classification and regression trees, bagging and random forests. *Psychol. Methods*, 14, 323–348, 2009.
- Trumbore, S.E.: Radiocarbon and soil carbon dynamics. *Annu. Rev. Earth Planet. Sci.*, 37, 47–66, 2009.
- 25 Trumbore, S.E.: Potential responses of soil organic carbon to global environmental change. *Proc. Natl. Acad. Sci. U. S. A.*, 94, 8284–8291, 1997.
- von Lütow, M., Kögel-Knabner, I., Ekschmitt, K., Flessa, H., Guggenberger, G., Matzner, E., and Marschner, B.: SOM fractionation methods: relevance to functional pools and to stabilization mechanisms. *Soil Biol. Biochem.*, 39, 2183–2207, 2007.
- 30 Wickham, H: stringr: Simple, Consistent Wrappers for Common String Operations. R package version 1.0.0. <http://CRAN.R-project.org/package=stringr>, 2015.
- [Zimmermann, M., Leifeld, J., Schmidt, M.W.I., Smith, P., Fuhrer, J.: Measured soil organic matter fractions can be related to pools in the RothC model. *European Journal of Soil Science*. 58, 658–667, 2007.](#)



5 Figure 1: Detail of the sequential pyrolysis and oxidation stages of Rock-Eval 6 (RE6) thermal analysis, and of the five thermograms used to derive the 30 RE6 parameters reflecting SOC thermal stability and bulk chemistry. The grey area under each RE6 thermogram represents the portion of the signal unaffected by soil carbonates that was used to calculate RE6 temperature parameters (modified after Behar et al., 2001; Saenger et al., 2013).







Figure 2: Evolution of SOC concentration (gC.kg⁻¹ soil) with time for the bare fallow plots of each experimental site, and representation of the 3×10^4 fitted exponential decay functions (Bayesian curve fitting method) from which a site-specific CP_{SOC} concentration (model parameter c) and its 95% confidence interval were determined (histogram in the upper right side of each scatter plot). At each site, the 95% confidence interval around the CP_{SOC} concentration was determined as $c \pm 1.96 \text{ sd}(c)$, where c is the model parameter c in Eq. (1) and $\text{sd}(c)$ is its standard deviation ~~calculated in Eq. (6)~~.



5 Figure 3: Principal component analysis (PCA) of the 30 RE6 parameters of the calibration soil sample set (n = 88). A: the scores of the calibration samples on the first two principal components are represented in the factorial map, as well as the projected principal component scores of the validation samples (n = 30). A color scale is used to represent the CP_{SOC} proportion (determined using Eq. (2)) in all samples. B: PCA loadings of the 30 RE6 parameters and projection of the CP_{SOC} proportion variable in the PCA correlation circle. Note that despite some of the 30 RE6 parameters are significantly correlated, all parameters are included in the analysis.



5 Figure 4: Performance of the random forests regression model based on Rock-Eval 6 thermal analysis (RE6-RF) for predicting the CP_{SOC} proportion. The performance statistics on the calibration set ($n = 88$) and on the validation set ($n = 30$) of the RE6-RF multivariate regression model are shown. Horizontal bars represent the estimated uncertainty (95% confidence intervals) on the reference CP_{SOC} proportion values of the validation set, calculated as: CP_{SOC} proportion [sample] $\pm 1.96 \times sd(CP_{SOC}$ proportion [sample]). Vertical bars represent the propagated errors (95% confidence intervals) on the RE6-RF predicted CP_{SOC} proportion values of the validation sample set, calculated as $\hat{y} \pm \hat{\tau} \times sd(\hat{y})$ (see Section 2.3.5), with a $\hat{\tau}$ value of 2.10-12 (Fig. S1 Supplementary material S2). Abbreviations: *RMSEC*, root-mean-square error of calibration; *RMSEP*, root-mean-square error of prediction; *RPD*, ratio of performance to deviation; *sd*, standard deviation.



5 **Figure 5:** Performance of the random forests regression model based on Rock-Eval 6 thermal analysis (RE6-RF) for predicting the CP_{soC} proportion, for a calibration set based on soils from three sites (Versailles, Rothamsted, Grignon; n = 84) and a validation set based on soils from a different pedoclimate but with similar RE6 thermal characteristics than those of the calibration set (Ultuna; n = 34).

Table 1: Measured total SOC concentrations, estimated site-specific CP_{SOC} concentrations, and resulting CP_{SOC} proportions in four long-term agronomic experimental sites used to generate calibration and validation soil sample sets. Abbreviations: LTBF, long-term bare fallow; min, minimum; max, maximum; sd, standard deviation; CI, confidence interval.

Site	Treatments (number of samples)	SOC concentration (gC.kg ⁻¹ soil) mean (min, max, sd)	CP_{SOC} concentration (gC.kg ⁻¹ soil) mean (95 % CI)	CP_{SOC} proportion mean (min, max, sd)
Versailles	Manure (n = 20)	27.9 (17.1, 45.5, 8.2)	6.22 (5.62–6.82)	0.24 (0.14, 0.36, 0.07)
	LTBF (n = 20)	10.5 (5.4, 19.7, 4.4)		0.67 (0.32, 1.00, 0.24)
Rothamsted	Grassland (n = 8)	36.8 (31.8, 42.6, 4.8)	10.46 (9.93–10.99)	0.29 (0.25, 0.33, 0.04)
	LTBF (n = 12)	17.7 (9.7, 30.5, 7.5)		0.68 (0.34, 1.00, 0.25)
Ultuna	Cropland (n = 23)	15.8 (12.4, 20.3, 2.2)	6.95 (5.22–8.68)	0.45 (0.34, 0.56, 0.06)
	LTBF (n = 11)	12.0 (9.1, 16.3, 2.4)		0.60 (0.43, 0.76, 0.12)
Grignon	Straw or composted straw (n = 12)	12.9 (11.7, 14.2, 0.8)	7.12 (5.16–9.08)	0.55 (0.50, 0.60, 0.03)
	LTBF (n = 12)	11.8 (8.4, 14.7, 1.9)		0.62 (0.48, 0.85, 0.11)
Calibration set (n = 88)		18.0 (5.5, 45.5, 9.5)		0.50 (0.14, 1.00, 0.21)
Validation set (n = 30)		16.1 (5.4, 38.8, 7.9)		0.53 (0.16, 1.00, 0.23)
All samples (n = 118)		17.5 (5.4, 45.5, 9.1)		0.51 (0.14, 1.00, 0.21)

5

Table 2: Spearman's rank correlation coefficient test between the 30 RE6 parameters and the CP_{SOC} proportion, and variable importance (ranking) of the 30 RE6 parameters to predict CP_{SOC} proportion in the random forests model based on Rock-Eval 6 thermal analysis (RE6-RF, calibration soil sample set, n = 88). Symbols for *p*-values: * *p* < 0.001; ** *p* < 0.01; * *p* < 0.05; NS *p* > 0.05 = not significant.**

RE6 parameter	Spearman's ρ with CP_{SOC} proportion	<i>p</i>-value	Variable importance to predict CP_{SOC} proportion in the RE6-RF regression model (rank)
T ₁₀ _HC_PYR	-0.36	***	20
T ₃₀ _HC_PYR	-0.12	NS	30
T ₅₀ _HC_PYR	0.00	NS	29
T ₇₀ _HC_PYR	0.09	NS	21
T ₉₀ _HC_PYR	0.55	***	18
T ₁₀ _CO_PYR	-0.21	*	7
T ₃₀ _CO_PYR	-0.39	***	12
T ₅₀ _CO_PYR	-0.33	**	14
T ₇₀ _CO_PYR	-0.23	*	23
T ₉₀ _CO_PYR	-0.09	NS	22
T ₁₀ _CO ₂ _PYR	0.07	NS	13
T ₃₀ _CO ₂ _PYR	0.71	***	4
T ₅₀ _CO ₂ _PYR	0.80	***	1
T ₇₀ _CO ₂ _PYR	0.78	***	2
T ₉₀ _CO ₂ _PYR	0.66	***	17
T ₁₀ _CO_OX	0.50	***	5
T ₃₀ _CO_OX	0.71	***	16
T ₅₀ _CO_OX	0.64	***	28
T ₇₀ _CO_OX	0.42	***	27
T ₉₀ _CO_OX	0.14	NS	25
T ₁₀ _CO ₂ _OX	0.72	***	10
T ₃₀ _CO ₂ _OX	0.83	***	3
T ₅₀ _CO ₂ _OX	0.82	***	6
T ₇₀ _CO ₂ _OX	0.80	***	8
T ₉₀ _CO ₂ _OX	0.54	***	19
I-index	0.04	NS	26
R-index	-0.01	NS	24
TLHC-index	-0.02	NS	15
HI	-0.78	***	9
OI_{RE6}	0.42	***	11

Supplementary material to:

A model based on Rock-Eval thermal analysis to quantify the size of the centennially persistent organic carbon pool in temperate soils

5 Lauric Cécillon^{1,2}, François Baudin³, Claire Chenu⁴, Sabine Houot⁴, Romain Jolivet², Thomas Kätterer⁵, Suzanne Lutfalla², Andy Macdonald⁶, Folkert van Oort⁴, Alain F. Plante⁷, Florence Savignac³, Laure N. Soucémariadin², Pierre Barré²

¹Université Grenoble Alpes, Irstea, UR **LESSEM**, F-38402 St-Martin-d'Hères, France

²Laboratoire de Géologie de l'ENS, PSL Research University, CNRS UMR 8538, F-75005, Paris, France

10 ³Sorbonne Université-UPMC-Univ Paris 06, Institut des Sciences de la Terre de Paris, F-75005 Paris, France

⁴AgroParisTech - INRA, UMR 1402 ECOSYS, F-78850 Thiverval-Grignon, France

⁵Department of Ecology, Swedish University of Agricultural Sciences, 750 07 Uppsala, Sweden


⁶Department of Sustainable Agriculture Sciences, Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ, UK

⁷Earth and Environmental Science, University of Pennsylvania, Philadelphia, USA


15 *Correspondence to:* Lauric Cécillon (lauric.cecillon@irstea.fr)

Supplementary material S1: Basic characteristics of the four long-term agronomic experimental sites used for the reference soil sample set. Abbreviation: *LTFB*, long-term bare fallow.

Site	Latitude, longitude	Mean annual temperature (°C)	Mean annual precipitation (mm)	Land use before experiment	Sampling depth (cm)	Soil texture (%, clay/silt/sand)
Versailles, France	48°48N, 2°08E	10.7	628	grassland	25	17/57/26
Rothamsted, United Kingdom	51°82N, 0°35E	9.5	712	grassland	23	25/62/13
Ultuna, Sweden	59°49N, 17°38E	5.5	533	arable	20	36/41/23
Grignon, France	48°51N, 1°55E	10.7	649	grassland	25	16/54/30



Supplementary material S2: Error propagation scheme in the random forests regression model (adapted from Coulston *et al.* (2016) and modified).



Supplementary material S3: Standard deviation of the CP_{SOC} proportion as a function of the CP_{SOC} proportion in the reference soil sample set (n = 118).

Supplementary reference

Coulston, J.W., Blinn, C.E., Thomas, V.A., and Wynne, R.H.: Approximating prediction uncertainty for random forest regression models. *Photogramm. Eng. Remote Sens.*, 82, 189–197, 2016.