



## Distribution of Fe isotopes in particles and colloids in the salinity gradient along the Lena River plume, Laptev Sea

Sarah Conrad<sup>1,\*</sup>, Johan Ingrí<sup>1</sup>, Johan Gelting<sup>1</sup>, Fredrik Nordblad<sup>1</sup>, Emma Engström<sup>1,2</sup>, Ilia Rodushkin<sup>1,2</sup>,  
5 Per S. Andersson<sup>3</sup>, Don Porcelli<sup>4</sup>, Örjan Gustafsson<sup>5</sup>, Igor Semiletov<sup>6,7,8</sup>, and Björn Öhlander<sup>1</sup>

<sup>1</sup>Department of Chemical Engineering and Geosciences, Luleå University of Technology, Luleå, Sweden

<sup>2</sup>ALS Laboratory Group, ALS Scandinavia AB, Aurorum 10, Luleå, Sweden

<sup>3</sup>Department of Geosciences, Swedish Museum of Natural History, Stockholm, Sweden

<sup>4</sup>Department of Earth Sciences, Oxford University, Oxford, United Kingdom

10 <sup>5</sup>Department of Environmental Science and Analytical Chemistry, Stockholm University, Stockholm, Sweden

<sup>6</sup>International Arctic Research Center (IARC), University of Alaska, Fairbanks, AK, USA

<sup>7</sup>Pacific Oceanological Institute (POI), Far Eastern Branch of the Russian Academy of Sciences (FEBRAS), Vladivostok, Russia

<sup>8</sup>Tomsk National Research Politecnical University, Tomsk, Russia

15 *Correspondence to:* Sarah Conrad (sarah.conrad@ltu.se)

**Abstract.** Riverine Fe input is the primary Fe source to the ocean. This study is focused on the distribution of Fe along the Lena River freshwater plume in the Laptev Sea using samples from a 600 km long transect in front of the Lena River mouth. Separation of the particulate ( $>0.22\mu\text{m}$ ), colloidal ( $0.22\mu\text{m} - 1\text{kDa}$ ), and truly dissolved ( $<1\text{kDa}$ ) fractions of Fe was carried out. The total Fe concentrations ranged from 0.15 to 57  $\mu\text{M}$  with Fe dominantly as particulate Fe. The loss of  $> 99\%$  of particulate Fe and about 90% of the colloidal Fe was observed across the shelf, while the truly dissolved phase was almost constant across the Laptev Sea. Thus, the truly dissolved Fe could be an important source of bioavailable Fe for plankton in the central Arctic Ocean, together with the colloidal Fe. Fe-isotope analysis showed that the particulate phase and the sediment below the Lena River freshwater plume had negative  $\delta^{56}\text{Fe}$  values (relative to IRMM-14). The colloidal Fe phase showed negative  $\delta^{56}\text{Fe}$  values close to the river mouth (about  $-0.20\text{\textperthousand}$ ) and positive  $\delta^{56}\text{Fe}$  values in the outermost stations (about 25  $+0.10\text{\textperthousand}$ ).

We suggest that the shelf zone acts as a sink for Fe particles and colloids with negative  $\delta^{56}\text{Fe}$  values, representing chemically reactive ferrihydrites. While the positive  $\delta^{56}\text{Fe}$  values of the colloidal phase within the outer Lena River freshwater plume, might represent Fe-oxyhydroxides, which remain in the water column, and will be the predominant  $\delta^{56}\text{Fe}$  composition in the Arctic Ocean.

30



## 1 Introduction

The cycling of Fe is a key component for understanding water quality and biogeochemical processes. Iron is the 4<sup>th</sup> most abundant element in the continental crust (Wedepohl, 1995). The concentration in seawater is low compared to riverine input (Martin and Gordon, 1991). The riverine input of Fe is the most important contribution to the oceanic Fe budget (Raiswell and 5 Canfield, 2012). Iron behaves non-conservatively during the mixing of freshwater and seawater and is removed to sediments (Boyle et al., 1977; Eckert and Sholkovitz, 1976; Gustafsson et al., 2000; Sholkovitz, 1978, 1976) since Fe-rich particles and colloids flocculate and settle in this mixing zone (Sholkovitz, 1978).

It has been recognised that dissolved Fe is related to dissolved organic carbon (DOC) in freshwater (Perdue et al., 1976) and so to investigate the pathways for organic carbon (OC) in the Arctic, knowledge about Fe cycling and the coupling between 10 the boreal-arctic watershed and the Arctic basin is crucial. Different forms and sizes of Fe and OC can be separated using a variety of filtration techniques. These include both membrane filtration (0.22 – 0.7 µm) and ultrafiltration (1kDa, 10kDa, or 30kDa) and size fractions are thus often operationally defined as particulate matter (larger than 0.22µm or 0.7µm), colloidal (smaller than particles, but do not pass an ultrafilter) and truly dissolved phases (passing through an ultrafilter). Due to the technical complexity with ultrafiltration, including the extensive filtration time, there are few ultrafiltration Fe data available 15 (Guo and Santschi, 1996; Ingri et al., 2000; Pokrovsky et al. 2012). Truly dissolved Fe data are scarce and deliver insights into this part of the Fe pool.

Previous studies showed that there is a relationship between Fe and OC in the dissolved fraction and found two main forms of Fe compounds: Fe-OC and Fe oxyhydroxides (Escoube et al., 2015; Hirst et al., 2017; Ilina et al., 2013; Ingri et al., 2006, 2000; Kritzberg et al., 2014; Pokrovsky et al., 2010, 2006; Pokrovsky and Schott, 2002; Raiswell and Canfield, 2012; Stolpe 20 et al., 2013). It has also been shown that humic substances (HS) are associated with newly formed Fe oxyhydroxides in freshwater (Pédrot et al., 2011; Tipping, 1981). The behaviour of these Fe and OC particles and colloids during estuarine mixing depend on their chemical reactivity, which is defined by their size and speciation (Poulton and Raiswell, 2005; Tagliabue et al., 2017). Hirst et al. (2017) found that about 70% of the total suspended Fe in the Lena River to be in the form 25 of reactive ferrihydrite. These ferrihydrites are independent particles within a network of amorphous particulate OC (POC) and are attached to the surfaces of primary organic matter and clay particles (Hirst et al., 2017).

Carbon-iron cycling is complex, and stable Fe isotope data show that the isotopic compositions might be used to investigate chemical pathways for Fe and OC during weathering and estuarine mixing in the boreal-arctic region (Dos Santos Pinheiro et al., 2014; Escoube et al., 2015, 2009; Ilina et al., 2013; Ingri et al., 2006; Mulholland et al., 2015; Poitrasson, 2006; Poitrasson et al., 2014). The  $^{56}\text{Fe}/^{54}\text{Fe}$  and  $^{57}\text{Fe}/^{54}\text{Fe}$  ratios are defined relative to the international reference material IRMM-14 and are 30 expressed as deviations from the standard in parts per thousand, or  $\delta$  notation (in per mill ‰), as

$$\delta^{56}\text{Fe} = \left[ \frac{(^{56}\text{Fe}/^{54}\text{Fe})_{\text{sample}}}{(^{56}\text{Fe}/^{54}\text{Fe})_{\text{IRMM-14}}} - 1 \right] * 10^3, \quad (1)$$



$$\delta^{57}\text{Fe} = \left[ \frac{(\text{Fe}^{57}/\text{Fe}^{54})_{\text{sample}}}{(\text{Fe}^{57}/\text{Fe}^{54})_{\text{IRMM-14}}} - 1 \right] * 10^3, \quad (2)$$

Using this definition, the continental crust has a  $\delta^{56}\text{Fe}$  value of  $0.07 \pm 0.02\text{\textperthousand}$ . In low-temperature environments the  $\delta^{56}\text{Fe}$  can vary by about  $5\text{\textperthousand}$  (Anbar, 2004; Beard et al., 2003; Dauphas and Rouxel, 2006; Fantle and DePaolo, 2004; Rouxel A. Bekker, 5 K.J. Edwards, 2005). The variations in  $\delta^{56}\text{Fe}$  can be used to trace different Fe phases in rivers (Dos Santos Pinheiro et al., 2014; Ilina et al., 2013; Ingri et al., 2006; Poitrasson et al., 2014) and to map the origin of Fe (Conway and John, 2014). Isotope fractionation processes result in a  $\delta^{56}\text{Fe}$  value that can be higher or lower compared to continental crust. The Fe isotopic composition is impacted by redox reactions (Wiederhold et al., 2006), complexation with organic ligands, and inorganic speciation of Fe, as well as the immobilization of Fe by precipitation and adsorption (Beard et al., 2003, 1999; Beard and 10 Johnson, 2004; Brantley et al., 2001; Bullen et al., 2001; Icopini et al., 2004; Poitrasson and Freydier, 2005; Skulan et al., 2002; Welch et al., 2003). These processes can yield either negative or positive  $\delta^{56}\text{Fe}$  values, depending on the initial Fe isotopic composition and the fractionation factor. Recent studies showed that subarctic and temperate rivers, with high Fe and 15 OC concentrations, have low  $\delta^{56}\text{Fe}$  values in the particulate phase, while the dissolved phase has high  $\delta^{56}\text{Fe}$  (Escoube et al., 2015, 2009; Ilina et al., 2013; Ingri et al., 2006; Rouxel et al., 2008; Severmann et al., 2006). Also, high  $\delta^{56}\text{Fe}$  values have 20 been reported in the Low Molecular Weight (LMW) fraction ( $< 10\text{kDa}$ ), while colloids and particles showed high  $\delta^{56}\text{Fe}$  values (Ilina et al., 2013). Furthermore, seasonal variations in the Fe isotopic composition and Fe speciation have been reported (Allard et al., 2004; Escoube et al., 2015; Ingri et al., 2006).

This study presents Fe concentrations and Fe isotope compositions in the particulate, colloidal, and truly dissolved phase along the Lena River freshwater plume in the Laptev Sea. The Lena River – Laptev Sea transect is stratified, with a freshwater layer 25 that is on top of more saline and dense, deep waters and plays an important role in the transport of Fe and the distribution of Fe isotopes in the Arctic Ocean. The main objectives were to study the distribution of Fe in the Lena River – Laptev Sea transect and the variations in the partitioning of Fe between the different size fractions, as well as to identify the impact of processes such as mixing, transformation, and removal by settling on the export of Fe to the deeper ocean.

## 2 Sampling Site and Analytical Methods

### 2.1 Study Area

The Lena River is 4,387 km long and has the 8<sup>th</sup> largest discharge in the world. It is the 2<sup>nd</sup> largest river draining into the Arctic Ocean and flows into the Laptev Sea (Fig. 1). The Lena watershed covers an area of  $2.46 \times 10^6 \text{ km}^2$  (Rachold et al., 1996) and is bound by the Verkhoyansk Mountain Ridge in the northeast and the central Siberian uplands in the west. Larch forests cover 72% of the watershed area and shrublands about 12% (Wagner, 1997; Walter and Breckle, 2002). Permafrost underlays 78–30 93% of the watershed (Zhang et al., 1999) and extends to depths of up to 1,500 m (Anisimov and Reneva, 2009). The annual discharge to the Arctic Ocean is  $581 \text{ km}^3$  (Yang et al., 2002). During spring flood, late May to June, 31–45% of the annual



runoff occurs (Amon et al., 2012). The Lena River delivers 5.6–5.8 Tg of DOC into the Arctic Ocean annually (Holmes et al., 2012; Raymond et al., 2007), along with about 0.4 Tg of particulate OC (Semiletov et al., 2011). More than 50% of the total OC (TOC) is delivered during a two-month period in summer, with 6.6 Tg year<sup>-1</sup> in June (Le Fouest et al., 2013) and 3.5 Tg year<sup>-1</sup> in July (Kutscher et al., 2017). The run-off from the Lena River accounts for more of 70% of the overall river inflow to  
5 the Laptev Sea (Antonov, 1967). The freshwater plume in the Laptev Sea is a mixing zone of about 600 km length and 50 km wide (Fig. 2). A low salinity freshwater plume overlies denser highly saline Arctic seawater (Alling et al., 2010). The Lena River plume can be divided into an inner and an outer plume based on a sharp increase in salinity, with salinities up to 5 in the inner plume and up to 15 in the outer plume (Alling et al., 2010). Both parts of the plume are separated by a strong halocline  
10 at about 10 m depth from the underlying dense Arctic sea water that has salinities up to 35 (Alling et al., 2010; Chester, 2003; Martin et al., 1993).

## 2.2 Sampling and Processing

The samples were collected in August 2008 during the International Siberian Shelf Study (ISSS-08) from the RV Yacob Smirnitskyi. The ISSS-08 was part of the International Polar Year (IPY) and the Arctic GEOTRACES programs. The sampling transect is 600 km long, stretching from off the Lena River mouth across the Laptev Sea, and samples from ten stations were  
15 collected (Fig. 1, Fig. 2 and Table 1). Additionally, surface sediment (upper 2 cm) samples were taken from the Kara, Laptev and East Siberian Seas (Fig. 1). Samples from this region collected during this cruise have also been studied for DOC (Alling et al., 2010; Bröder et al., 2016; Karlsson et al., 2016; Salvadó et al., 2017), dissolved inorganic carbon (Alling et al., 2012), POC (Karlsson et al. 2016; Sánchez-García et al., 2011), nutrients and alkalinity (Anderson et al., 2009; Pipko et al., 2017), and stable O isotopes (Rosén et al., 2015).

20

25

30



**Table 1: Sampling stations in the Laptev Sea of the ISSS-08 research cruise. Temperature and salinity data for the shelf sediment sample locations are obtained from the overlying bottom waters.**

Station	Lat.	Long.	Date	Water depth	Salinity	Temp.	
	dec.	dec.		m	°C		
Lena River freshwater plume	<b>YS-128</b>	76.987	130.356	17/09/08	51	20.66	-1.43
	<b>YS-4*</b>	75.987	129.984	23/08/08	44	11.45	-1.54
	<b>YS-5</b>	75.266	130.017	24/08/08	42	7.03	-1.56
	<b>YS-6</b>	74.724	130.016	24/08/08	32	5.23	-1.61
	<b>YS-7</b>	74.132	130.000	24/08/08	16	6.32	-1.26
	<b>YS-8</b>	73.566	130.008	24/08/08	13	5.34	-0.78
	<b>YS-9</b>	73.366	129.997	25/08/08	23	8.47	-1.13
	<b>YS-10</b>	73.184	129.996	25/08/08	20	3.78	-0.89
	<b>YS-11</b>	73.019	129.989	25/08/08	11	2.67	-0.32
	<b>YS-14*</b>	71.630	130.050	27/08/08	8	1.28	11.14
Shelf sediment locations	<b>YS-2</b>	73.405	72.995	19/08/08	30	7.85	-1.09
	<b>YS-3</b>	73.492	79.885	19/08/08	35	5.54	-1.06
	<b>YS-13</b>	71.968	131.701	26/08/08	21	3.79	-1.03
	<b>YS-26</b>	72.460	150.596	31/08/08	16	19.18	-0.72
	<b>YS-28</b>	72.651	154.185	01/09/08	28	19.73	-0.86
	<b>YS-30</b>	71.358	152.153	01/09/08	9	19.31	1.19
	<b>YS-39</b>	71.217	169.347	04/09/08	44	27.56	-1.64

\* also sampled for sediment

All water samples, besides YS-14, were collected between 2.5 m and 5.0 m depth using a peristaltic pump and acid-cleaned, 5 silicon tubing. The tubing was attached to a flagpole, which was mounted to the bow of the ship. To avoid contamination from the ship, the flagpole was extended about 10 m in front of the ship. The samples were pumped into a 25 L container, which was rinsed with MQ water between each station. Station YS-14 was sampled at 4.0 m depth using a 60 L Go-Flo® water sampler. All equipment in contact with the samples were cleaned with 5% HNO<sub>3</sub>, rinsed with MQ water, and dried in a HEPA-filtered clean air hood. Membrane filtration was carried out within 12 hours of sampling. All water samples were stored in 10 acid-cleaned polyethylene (PE) bottles and acidified, and all nitrocellulose filters (0.22 µm, Millipore®) were stored at -18°C until further analysis (Ödman et al., 1999). Samples for particulate organic carbon were filtered with 0.7 µm GF/F glassfiber filters (Whatman®). The filters were pre-combusted for four hours at 450°C to limit the C blank.



Sediment samples were taken with a GEMAX gravity corer and a Van Veen grab sampler as described earlier (Vonk et al., 2012).

During cross-flow ultrafiltration the sample water ( $< 0.22\mu\text{m}$ ) flows across a membrane surface at a constant pressure. This process prevents clogging, since while particles smaller than the membrane cut-off can pass, larger suspended particles remain 5 circulating in the sample water. The sample water progressively decreases in volume as the permeate crosses the filter, and the larger colloids and particles remain in the retentate, which therefore are progressively concentrated. The cross-flow ratio ( $\text{CFR} = Q_R/Q_P$ , where  $Q_R$  and  $Q_P$  are the flow rates of the retentate and permeate, respectively) (Forsberg et al., 2006; Ingri et al., 2000; Larsson et al., 2002) was kept between 60 and 100 to achieve an overall concentration factor larger than 10 (Conc. Fact. =  $(V_P + V_R)/V_R$ , where  $V_P$  and  $V_R$  are the final volumes of the permeate and retentate, respectively). For the concentration 10 factors and cross-flow ratios, see Table 2. In this study, the water used for ultrafiltration was pre-filtered through a membrane ( $< 0.22\mu\text{m}$ ) prior to introduction into the MilliPore® Prep/Scale ultrafiltration system, which had a cutoff of 1 kDa. Thus, the permeate is  $< 1$  kDa, while the retentate includes colloids between  $< 0.22\mu\text{m}$  and 1 kDa.

Table 2: Cross-flow ultrafiltration details

Sample	Retentate ( $V_R$ ) litre	Permeate ( $V_P$ ) litre	Conc. Factor	Retentate	Permeate	Cross-flow
				$Q_R$ (ml/min)	$Q_P$ (ml/min)	ratio (CFR)
YS-128	0.97	16.4	18	>3,000	30-50	60-100
YS-4	1.14	11.8	11	3,000	30	100
YS-11	1.10	10.5	11	>3,000	30-50	60-100
YS-14	0.61	12.2	21	>3,000	30-50	60-100

### 15 2.3 Analytical Methods

Iron concentrations and isotopic compositions were measured at ALS Scandinavia AB. High purity Suprapure® acids were used throughout sample treatment and analysis. Organic carbon analyses were carried out at Stockholm University (for analytical details, see Alling et al., 2010; Sánchez-García et al., 2011).

For element analysis, the water samples were diluted (2-200 fold) with 10 %  $\text{HNO}_3$ . The degree of dilution was dependent on 20 the salinity of the sample. At least two dilutions of each sample were carried out; one high dilution for determination of major elements and one low dilution for minor and trace elements. For Fe analysis, the samples were diluted by a factor of 50. In order to analyze the particles on the filters, the filters were treated with a 1,000:1 mixture of  $\text{HNO}_3/\text{HF}$  overnight, followed by closed-vessel microwave-assisted digestion. Prior to analysis, the digests were further diluted in 10%  $\text{HNO}_3$ .

Multi-elemental analysis of the water and filter samples was performed on an Inductively Coupled Plasma Sector Field Mass 25 Spectrometer (ICP-SFMS, ELEMENT2 Thermo Scientific) at ALS Scandinavia AB. The measurement procedure combines



internal standardization and external calibration. For internal standardization, indium was added to all the solutions (Rodushkin et al., 2005; Rodushkin and Ruth, 1997).

The blanks of digested filters (0.22 $\mu$ m) for Fe were 2.79  $\mu$ g/L, which is about 0.25% of the average Fe concentration in the samples for the Lena River sampling transect. Replicated measurements of sample concentrations showed a precision of  $\pm 3\%$  5 (n=4).

For the Fe isotope ratio measurements, water samples and digested filters were evaporated to dryness, and the residue was re-dissolved in 1 mL 9M HCl. Iron was separated from the matrix by ion exchange, with a recovery rate above 95%. The Fe isotope compositions in separated fractions from filters and water samples were measured using a Multi Collector Inductively Coupled Plasma Mass Spectrometer (MC-ICP-MS, NEPTUNE PLUS<sup>®</sup>, Thermo Scientific) equipped with micro-concentric 10 nebulizer and tandem cyclonic/Scott double pass spray chamber. Instrumental mass biases were corrected by sample-standard bracketing using IRMM-14 CRM, while an internal standard (Ni) was added to all samples and used to correct for instrumental drift. Each sample was measured twice with the sample-standard bracketing method. Detailed information on the correction procedures can be found in Baxter et al. (2006). During the Fe isotope analysis,  $\delta^{56}\text{Fe}$  and  $\delta^{57}\text{Fe}$  were measured. In the three 15 isotopes plot of  $\delta^{56}\text{Fe}$  and  $\delta^{57}\text{Fe}$  all samples plot on a single mass fractionation line (Fig. S1). We only discuss the  $\delta^{56}\text{Fe}$  in this study, although all Fe isotope data are reported in Table 3 including  $2\sigma$  (n=4;).

20

25

30



Table 3: Fe isotope data for the sediment and the particulate and colloidal phase.

Particulate >0.22μm				
Station	$\delta^{56/54}\text{Fe}$	2σ	$\delta^{57/54}\text{Fe}$	2σ
	‰	‰	‰	‰
YS-128	-0.289	0.050	-0.487	0.024
YS-4	-0.406	0.126	-0.735	0.114
YS-6	-0.360	0.014	-0.644	0.082
YS-8	-0.130	0.008	-0.266	0.136
YS-11	-0.067	0.040	-0.106	0.008
YS-14	-0.048	0.106	-0.097	0.114

Colloidal 1kDa-0.22μm				
Station	$\delta^{56/54}\text{Fe}$	2σ	$\delta^{57/54}\text{Fe}$	2σ
	‰	‰	‰	‰
YS-128	0.112	0.069	0.233	0.050
YS-4	0.102	0.079	0.277	0.038
YS-11	-0.227	0.089	-0.312	0.298
YS-14	-0.171	0.015	-0.267	0.030

Surface Sediment				
Station	$\delta^{56/54}\text{Fe}$	2σ	$\delta^{57/54}\text{Fe}$	2σ
	‰	‰	‰	‰
YS-13	-0.233	0.070	-0.324	0.006
YS-4	-0.220	0.040	-0.355	0.028
YS-26	-0.209	0.002	-0.298	0.116
YS-14	-0.250	0.110	-0.404	0.100
YS-2	-0.351	0.150	-0.439	0.102
YS-3	-0.230	0.024	-0.396	0.106
YS-11	-0.083	0.022	-0.209	0.094
YS-28	-0.131	0.074	-0.220	0.118
YS-30	-0.102	0.028	-0.185	0.088
YS-39	-0.241	0.086	-0.403	0.124



### 3 Results

All concentration results are reported in the supplement (Tab. S2 and Tab. S3).

#### 3.1 Organic Carbon distributions in the Lena River plume

The DOC concentrations show a small variation of between 300 and 440  $\mu\text{M}$  in the surface waters of the inner and outer plume

5 (Fig. 3). The average DOC concentration of 410  $\mu\text{M}$  in the surface water of the Lena River freshwater plume has been reported by Alling et al. (2010) and is similar to previous studies (Cauwet and Sidorov, (1996): 300–600  $\mu\text{M}$ ). It has been shown that DOC is behaving conservatively during mixing between Lena River water and Arctic Ocean water along the sampling profile (Alling et al., 2010; Opsahl et al., 1999; Pugach et al., 2018). The POC concentrations decrease from high values (89  $\mu\text{M}$ ) close to the coast to low values (8  $\mu\text{M}$ ) in the outer plume (Fig 3; S2). In the inner plume (YS-14 to YS-10) the POC 10 concentrations are high, between 89  $\mu\text{M}$  and 36  $\mu\text{M}$ , whereas in the outer plume the POC concentrations were almost constant, with an average value of about 12  $\mu\text{M}$ . The overall average POC concentration of about 28  $\mu\text{M}$  has been earlier reported by Sánchez-García et al. (2011).

#### 3.2 Iron concentrations in the Lena River freshwater plume

Three size fractions were analyzed for Fe: particulate Fe (pFe,  $> 0.22 \mu\text{m}$ ), colloidal Fe (cFe, 1kDa-0.22  $\mu\text{m}$ ), and truly 15 dissolved Fe (dFe,  $< 1 \text{ kDa}$ ). The total Fe (tFe) concentration was calculated as the sum of pFe, cFe, and dFe (S3).

The pFe concentration decreased from 56 to 0.1  $\mu\text{M}$  along the Lena River freshwater plume (Fig. 4). Between the inner and the outer plumes (i.e. between YS-10 and YS-9), the pFe concentration dropped to 0.9  $\mu\text{M}$ , a loss of 98% of pFe. The cFe concentration decreased from 0.6 to 0.1  $\mu\text{M}$  along the freshwater plume, a loss of about 85% cFe (Fig 4). The concentration of dFe was low, at around 8 nM, and relatively constant along the plume (Fig. 4). In total, a loss of >99% tFe was observed 20 between the first station (YS-14) and the last station (YS-128).

We observed non-conservative behaviour of pFe during mixing between Lena River water and Arctic Ocean water, while cFe showed generally conservative behaviour, with an almost linear correlation with salinity (Fig. 5). The pFe concentrations below 1  $\mu\text{M}$  also showed an almost linear correlation at salinities above 5 in the outer plume. In the inner plume, at salinities below 5, the pFe showed non-conservative behaviour.

#### 25 3.3 Iron isotopes in the Lena River freshwater plume

The Fe isotope compositions in the particulate and the colloidal phases, as well as in the surface sediments, are reported in Fig. 6. The  $\delta^{56}\text{Fe}$  values in the particles varied between  $-0.05 \pm 0.11\text{\textperthousand}$  (YS-14) in the inner plume and  $-0.41 \pm 0.12\text{\textperthousand}$  (YS-4) in the outer plume (Fig. 6), with the  $\delta^{56}\text{Fe}$  values in the outer plume all lower compared to the inner plume. The cFe show negative  $\delta^{56}\text{Fe}$  values (average  $-0.20 \pm 0.06\text{\textperthousand}$ ) in the inner plume and positive  $\delta^{56}\text{Fe}$  values (average  $0.11 \pm 0.08\text{\textperthousand}$ ) in the outer plume. 30 The surface sediments from the Laptev Sea all had negative  $\delta^{56}\text{Fe}$  values ( $-0.23 \pm 0.08\text{\textperthousand}$  and  $-0.25 \pm 0.12\text{\textperthousand}$ ). Surface



sediments obtained from 10 samples in other parts of the ESAS showed only small variations (Fig 1 and Fig. 6; Tab. 3; Tab. S4).

#### 4 Discussion

In the Laptev Sea close to the river mouth about 18% of the total OC was present as POC and this was apparently rapidly lost  
5 during mixing (Fig. 5) In the outer plume only about 2% of the total OC was present as POC. It has been suggested that POC  
in the Lena River freshwater plume is transported in different forms, including large particles, which can sink, and almost  
neutrally buoyant flocculates of humic substances (Gustafsson and Gschwend, 1997; Gustafsson et al., 2000; Sánchez-García  
et al., 2011). The POC, which is associated with larger particles ( $> 0.7\text{ }\mu\text{m}$ ), will settle close to land, whereas the humic  
substance flocculates will travel further out (Vonk et al., 2010).

##### 10 4.1 Iron behaviour in the Lena River freshwater plume

The pFe concentrations found in the Laptev Sea close to the shore are higher than the average pFe concentration in the Lena  
River, but similar to the highest river values up to  $32\text{ }\mu\text{M}$  (Hirst et al., 2017). The cFe and dFe in the Lena River (Hirst et al.,  
2017) showed higher average concentrations (cFe:  $1.5\text{ }\mu\text{M}$ ; dFe:  $54\text{ nM}$ ) that are similar to concentrations found in the Lena  
River – Laptev Sea transect. Within the Arctic Ocean, dissolved Fe (cFe + dFe) concentrations vary between 0.2 and  $63\text{ nM}$   
15 and the concentrations depend on distance to the shore and depths of sampling, with generally higher values in surface waters  
as well as close to the bottom sediment, which might be related to resuspension, sinking of brines, or resuspension from the  
sediment of Fe (Klunder et al., 2012; Thuróczy et al., 2011). The cFe concentrations are higher close to the coast and decreasing  
in the outer plume to values that are similar to cFe concentrations reported from further out in the Arctic Ocean. Estuarine  
processes, including flocculation and sedimentation, are the primary cause for the sharp decrease of particulate and dissolved  
20 Fe concentrations along the transect from the river towards the open Arctic Ocean. The distribution of Fe between the different  
phases show that pFe is the dominant Fe-phase in the inner plume system (with a pFe/cFe ratio of about 90). However, most  
of the pFe is lost in the inner plume close to the shore and the ratio pFe/cFe decreases towards a ratio of about 1 in the outer  
plume.

We observed non-conservative mixing of pFe at salinities lower than 5 and conservative mixing at salinities higher than 5.  
25 Recent studies showed that the majority of particulate Fe ( $70\pm15\%$ ) coming from the Lena River is in the form of chemically  
reactive ferrihydrite (Hirst et al., 2017). Organic C hinders the coagulation of the particles during riverine transport, but in the  
estuarine mixing zone the negatively charged particles will react with seawater cations and form larger aggregates (Boyle et  
al., 1977). This results in flocculation and sinking to sediments in the Lena River – Laptev Sea transect and can thus explain  
the observed non-conservative behaviour (Martin et al., 1993). This process is a common feature for Fe that is observed in  
30 other estuaries and is responsible for at least 80% loss of “dissolved” riverine Fe (Boyle et al., 1977; Figuères et al., 1978;  
Guieu et al., 1996; Windom, H. L., Beck, K., Smith, 1971). The large amount of pFe (98%) lost in the inner Lena River



freshwater plume is likely due removal of chemically reactive ferrihydrite, which is the main form of particulate Fe in the Lena River. Furthermore, it has been shown that about 20% of OC in the Eurasian Arctic Shelf is bound to reactive Fe phases (Salvadó et al., 2015). It has also been shown that part of the ferrihydrite might be transported via surface attachment to POC in a network of organic fibrils (Hirst et al., 2017). The attachment of POC to the ferrihydrite possibly reduces the density of 5 Fe-oxyhydroxides (Passow, 2004), allowing both POC and pFe to be transported into the Arctic Ocean, where they are present at about 2% of their initial concentration in rivers. Concentrations of pFe and cFe in the form of Fe oxyhydroxides show a linear correlation with salinity at salinities  $>5$ , suggesting that these particles and colloids are less affected by changes in ionic strength. The truly dissolved Fe ( $<1\text{kDa}$ ) concentrations along the freshwater plume are almost constant around 8nM. These 10 observations are in accordance with previous studies in the Laptev Sea where dissolved Fe concentrations of  $>10$  nM has been reported (Klunder et al., 2012). It has also been reported that about 74 to 83% of the dissolved Fe is present in the truly dissolved phase in the Arctic Ocean (Thuróczy et al., 2011). Slagter et al. (2017) report dissolved Fe concentration of 2.6 nM in the Transpolar Drift, which is transporting surface water from Siberian great rivers, e.g. Lena River, across the Arctic Ocean into the Atlantic. Available evidence indicates that the Ob River similarly contributes Fe into the open Arctic Ocean. Along 15 the Ob River, the dFe shows relatively constant dFe concentrations of 36–44 nM in the 10kDa fraction (Dai and Martin, 1995), which are somewhat higher than reported here for the Lena possibly due to a larger ultrafiltration cutoff size. The conservative behaviour of dFe concentrations along a salinity gradient has been examined in estuarine mixing experiments, and it has been shown that freshwater Fe oxyhydroxide colloids aggregate into much larger particles in contact with seawater, whereas the truly dissolved phase was virtually unaffected (Gustafsson et al., 2000; Stolpe and Hassellöv, 2007). The observation that the truly dissolved phase is less affected by the increase in salinity suggests that this phase can be transported through estuaries 20 and further out into the open ocean (Laglera and Van Den Berg, 2009).

River water is the most important source of Fe for the central Arctic Ocean (Klunder et al., 2012) and estuarine processes significantly modify the amount and distribution of Fe between different fractions, and therefore also the bioavailability of the river-derived Fe. The loss of Fe-OC aggregates close to the shoreline might also cause a great loss of phosphorous and thus contribute to the suggested “rusty carbon sink” (Lalonde et al., 2012; Salvadó et al., 2015).

#### 25 **4.2 Iron isotopes in the Lena River freshwater plume**

The measured  $\delta^{56}\text{Fe}$  compositions in the Lena River plume are broadly similar to those reported in previous studies in other arctic/subarctic regions. In these areas, within the fully oxidized water column, the pFe phase show negative  $\delta^{56}\text{Fe}$  values, while the dissolved phase generally shows higher values (Escoube et al., 2015, 2009; Ingri et al., 2006; Staubwasser et al., 2013; Zhang et al., 2015). Hirst et al. (in prep.) show a seasonal dependence of the  $\delta^{56}\text{Fe}$  composition of the dissolved fraction 30 (colloidal and truly dissolved) in the Lena River, with summer flow  $\delta^{56}\text{Fe}$  values higher than those of the continental crust, and spring flood  $\delta^{56}\text{Fe}$  values of the dissolved phase lower than those of the continental crust. Therefore, the Lena River can be a source of colloidal Fe with both negative and positive isotope compositions. Along the freshwater plume the colloidal Fe phase has two different Fe isotope compositions, positive and negative  $\delta^{56}\text{Fe}$  values, therefore it might also represent water



masses from different seasons. This would suggest that the water masses in the inner plume represent spring flood discharge, whereas the water masses in the outer plume represent summer flow discharge. In contrast, Alling et al. (2010), claim that the age of the entire freshwater plume is approximately two months. All measured DOC samples (400–420  $\mu\text{M}$ ) from their study plot on a mixing line of Lena River water measured in August and Arctic Deep water. If the water would represent spring

5 flood discharge, which has much higher DOC concentrations (1170  $\mu\text{M}$ ), their samples would plot on a different mixing line (Alling et al., 2010).

Sundman et al., (2014) measured the speciation of Fe in stream water samples with X-ray absorption spectroscopy and found iron-organic complexes with mixed speciation states of Fe as Fe(II, III)-OC and Fe(III)oxyhydroxides associated to OC. The variations in the distributions of Fe between the different species in the iron-organic complexes are controlled by pH and OC concentrations (Neubauer et al., 2013; Sundman et al., 2013). The Fe speciations of these complexes regulate the Fe isotopic

10 composition. When Fe(II) is oxidized to Fe(III), the heavy  $^{56}\text{Fe}$  is enriched in the Fe(III) phase whereas Fe(II) becomes depleted in the  $^{56}\text{Fe}$  isotope (Bullen et al., 2001; Homoky et al., 2012; Rouxel et al., 2008; Severmann et al., 2006; Welch et al., 2003; Wu et al., 2011). Laboratory experiments of the oxidative precipitation of Fe(II) to Fe(III), which can occur in natural streams, show an overall fractionation factor of 0.9. Hence, Fe(III)oxyhydroxides should show an enrichment of  $^{56}\text{Fe}$  in oxidized river water, while Fe(II, III)-OC complexes should show a depletion of  $^{56}\text{Fe}$ . The existence of two different Fe colloid pools,

15 composed of organic-rich and Fe rich particles, was shown by Pokrovsky and Schott (2002) in small boreal rivers. Fe-isotope data from this study show the existence of two colloidal Fe phases with different  $\delta^{56}\text{Fe}$  within the Lena River – Laptev Sea transect. The Fe isotope variation along the plume and the composition of the surface sediment suggest that the chemically reactive ferrihydrite represent colloids and particles, with a negative  $\delta^{56}\text{Fe}$  value, sedimenting close to the shoreline. The Fe-oxyhydroxides that remain in the water column could then be responsible for the positive  $\delta^{56}\text{Fe}$  values in the colloidal phase

20 in the outer plume. Therefore, in this case the Lena River is an important source of positive  $\delta^{56}\text{Fe}$  values to the Arctic Ocean, along with small OC-rich arctic and subarctic rivers (Ilina et al., 2013; Pokrovsky et al., 2014).

The surface sediments in the shelf areas along the Laptev Sea have  $\delta^{56}\text{Fe}$  values of -0.2‰ (Figure 6). This value results from the removal of particulate and colloidal Fe(II, III)oxyhydroxides from the water column and burial in the sediment. As seen in earlier studies, flocculation during estuarine mixing did not fractionate the Fe isotopic composition of the colloids and particles

25 (Bergquist and Boyle, 2006; Escoube et al., 2009; Fantle and DePaolo, 2004; Poitrasson et al., 2014).

## 5. Conclusions

Close to the coast and within the inner part of the river plume, the concentration of pFe dominates the total Fe budgets. In the outer part of the plume, the pFe and cFe concentrations are almost equal, as more than 99% of the total Fe is lost. The loss of

30 pFe, most likely in the form of chemically reactive ferrihydrite, results from increasing ionic strength, due to increasing salinities, which promotes flocculation. The coagulation and removal appear at the beginning of the mixing zone at low salinities (0–5). Colloidal Fe concentrations are almost constant along the inner plume and the decrease along the outer plume



due to conservative mixing. The truly dissolved Fe shows little variation along the Lena River freshwater plume. Therefore, the river-derived truly dissolved fraction could be an important source of bioavailable Fe, along with colloidal Fe, which may affect the primary production in the central Arctic Ocean.

The Fe isotope compositions in the Lena River freshwater plume provide clear indications of which forms of Fe reach the deep ocean basin. There are significant differences between the particulate and colloidal phases. The negative  $\delta^{56}\text{Fe}$  values, found in the colloidal and particulate phases, are lost during estuarine mixing and buried in the sediment. These negative  $\delta^{56}\text{Fe}$  values seem to represent chemically reactive ferrihydrite. Within the colloidal phase, we measured positive  $\delta^{56}\text{Fe}$  values further out in the plume, which likely represent Fe-oxyhydroxide, which remain buoyant in the water column, transported along the Lena River freshwater plume into the Arctic Ocean.

10 Climate warming is increasing discharge and accompanying OC and Fe from land to the ocean. Increasing the amount of colloidal and truly dissolved Fe, which is passing the estuarine mixing zone will lead to a higher Fe flux towards the Arctic Ocean.

### Competing interests

The authors declare that they have no conflict of interest.

### 15 Acknowledgment

The ISSS-08 program was supported by the Knut and Alice Wallenberg Foundation, the Far Eastern Branch of the Russian Academy of Sciences, the Swedish Research Council (621-2004-4039 and 211-621-2007), the U.S. National Oceanic and Atmospheric Administration, the Russian Foundation for Basic Research, the Swedish Polar Research Secretariat, and the Stockholm University Bert Bolin Centre for Climate Research. Ö.G. also acknowledge a Distinguished Professor Grant from the Swedish Research Council (VR Contract No. 2017-05687), an Advanced Grant from the European Research Council (ERC-AdG CC-TOP project #695331). I.S. acknowledges the Russian Government (grant No. 14.Z50.31.0012). The ISSS-08 program is part of the IPY (International Polar Year) and the GEOTRACES program.

### References

Allard, T., Menguy, N., Salomon, J., Calligaro, T., Weber, T., Calas, G., Benedetti, M.F., 2004. Revealing forms of iron in river-borne material from major tropical rivers of the Amazon Basin (Brazil). *Geochim. Cosmochim. Acta* 68, 3079–3094. <https://doi.org/10.1016/J.GCA.2004.01.014>

Alling, V., Porcelli, D., Mörth, C.M., Anderson, L.G., Sanchez-Garcia, L., Gustafsson, Ö., Andersson, P.S., Humborg, C., 2012. Degradation of terrestrial organic carbon, primary production and out-gassing of CO<sub>2</sub> in the Laptev and East Siberian



Seas as inferred from  $\delta^{13}\text{C}$  values of DIC. *Geochim. Cosmochim. Acta* 95, 143–159.  
<https://doi.org/10.1016/j.gca.2012.07.028>

Alling, V., Sanchez-Garcia, L., Porcelli, D., Pugach, S., Vonk, J.E., Van Dongen, B., Mörth, C.M., Anderson, L.G., Sokolov, A., Andersson, P., Humborg, C., Semiletov, I., Gustafsson, Ö., 2010. Nonconservative behavior of dissolved organic carbon  
5 across the Laptev and East Siberian seas. *Global Biogeochem. Cycles* 24, 1–15. <https://doi.org/10.1029/2010GB003834>

Amon, R.M.W., Rinehart, A.J., Duan, S., Loucheouarn, P., Prokushkin, A., Guggenberger, G., Bauch, D., Stedmon, C., Raymond, P.A., Holmes, R.M., McClelland, J.W., Peterson, B.J., Walker, S.A., Zhulidov, A. V., 2012. Dissolved organic matter sources in large Arctic rivers. *Geochim. Cosmochim. Acta* 94, 217–237. <https://doi.org/10.1016/j.gca.2012.07.015>

Anbar, A.D., 2004. Iron stable isotopes: Beyond biosignatures. *Earth Planet. Sci. Lett.* 217, 223–236.  
10 [https://doi.org/10.1016/S0012-821X\(03\)00572-7](https://doi.org/10.1016/S0012-821X(03)00572-7)

Anderson, L.G., Jutterström, S., Hjalmarsson, S., Wählström, I., Semiletov, I.P., 2009. Out-gassing of CO<sub>2</sub> from Siberian Shelf seas by terrestrial organic matter decomposition. *Geophys. Res. Lett.* 36, L20601.  
<https://doi.org/10.1029/2009GL040046>

Anisimov, O., Reneva, S., 2009. Permafrost and Changing Climate: The Russian Perspective. [http://dx.doi.org/10.1579/0044-1547\(2006\)35\[169:PACCTR\]2.0.CO;2](http://dx.doi.org/10.1579/0044-1547(2006)35[169:PACCTR]2.0.CO;2)

15 Antonov, V.S., 1967. The Mouth Area of the Lena, in: *The Hydrographic Review*. Gidrometeoizdat, Leningrad.

Baxter, D.C., Rodushkin, I., Engström, E., Malinovsky, D., 2006. Revised exponential model for mass bias correction using an internal standard for isotope abundance ratio measurements by multi-collector inductively coupled plasma mass spectrometry. *J. Anal. At. Spectrom.* 21, 427. <https://doi.org/10.1039/b517457k>

20 Beard, B.L., Johnson, C.M., 2004. Ancient Earth and Other Planetary Bodies. *Rev. Mineral.* 55, 319–357.  
<https://doi.org/10.2138/gsrmg.55.1.319>

Beard, B.L., Johnson, C.M., Cox, L., Sun, H., Nealson, K.H., Aguilar, C., 1999. Iron isotope biosignatures. *Science* (80-. ). 285, 1889–1891. <https://doi.org/10.1126/science.285.5435.1889>

Beard, B.L., Johnson, C.M., Von Damm, K.L., Poulsen, R.L., 2003. Iron isotope constraints on Fe cycling and mass balance  
25 in oxygenated Earth oceans. *Geology* 31, 629–632. [https://doi.org/10.1130/0091-7613\(2003\)031<0629:IICOFC>2.0.CO;2](https://doi.org/10.1130/0091-7613(2003)031<0629:IICOFC>2.0.CO;2)

Bergquist, B.A., Boyle, E.A., 2006. Iron isotopes in the Amazon River system: Weathering and transport signatures. *Earth Planet. Sci. Lett.* 248, 39–53. <https://doi.org/10.1016/j.epsl.2006.05.004>

Boyle, E.A., Edmond, J.M., Sholkovitz, E.R., 1977. The mechanism of iron removal in estuaries. *Geochim. Cosmochim. Acta* 41, 1313–1324. [https://doi.org/10.1016/0016-7037\(77\)90075-8](https://doi.org/10.1016/0016-7037(77)90075-8)

30 Brantley, S.L., Liermann, L., Bullen, T.D., 2001. Fractionation of Fe isotopes by soil microbes and organic acids. *Geology* 29, 535. [https://doi.org/10.1130/0091-7613\(2001\)029<0535:FOFIBS>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<0535:FOFIBS>2.0.CO;2)

Bröder, L., Tesi, T., Salvadó, J.A., Semiletov, I.P., Dudarev, O.V., Gustafsson, Ö. 2016. Fate of terrigenous organic matter  
across the Laptev Sea from the mouth of the Lena River to the deep sea of the Arctic interior. *Biogeosci.* 13, 17, 5003–5019.  
<https://doi.org/10.5194/bg-13-5003-2016>



Bullen, T.D., White, A.F., Childs, C.W., Vivit, D. V., Schultz, M.S., 2001. Demonstration of a significant iron isotope fractionation in nature. *Geology* 29, 699–702. [https://doi.org/10.1130/0091-7613\(2001\)029<0699:DOSAII>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<0699:DOSAII>2.0.CO;2)

Cauwet, G., Sidorov, I., 1996. The biogeochemistry of Lena River: organic carbon and nutrients distribution. *Mar. Chem.* 53, 211–227. [https://doi.org/10.1016/0304-4203\(95\)00090-9](https://doi.org/10.1016/0304-4203(95)00090-9)

5 Chester, R. (Roy), 2003. *Marine geochemistry*, 2. [rev.] ed. ed. Blackwell Pub, Malden.

Conway, T.M., John, S.G., 2014. Quantification of dissolved iron sources to the North Atlantic Ocean. *Nature* 511, 212–215. <https://doi.org/10.1038/nature13482>

Dai, M.-H., Martin, J.-M., 1995. First data on trace metal level and behaviour in two major Arctic river-estuarine systems (Ob and Yenisey) and in the adjacent Kara Sea, Russia. *Earth Planet. Sci. Lett.* 131, 127–141. [https://doi.org/10.1016/0012-821X\(95\)00021-4](https://doi.org/10.1016/0012-821X(95)00021-4)

10 Dauphas, N., Rouxel, O., 2006. Mass spectrometry and natural variations of iron isotopes. *Mass Spectrom. Rev.* 25, 515–520. <https://doi.org/10.1002/mas.20078>

Dos Santos Pinheiro, G.M., Poitrasson, F., Sondag, F., Cochonneau, G., Vieira, L.C., 2014. Contrasting iron isotopic compositions in river suspended particulate matter: The Negro and the Amazon annual river cycles. *Earth Planet. Sci. Lett.* 15 394, 168–178. <https://doi.org/10.1016/j.epsl.2014.03.006>

Eckert, J.M., Sholkovitz, E.R., 1976. The flocculation of iron, aluminum and humates from river water by electrolytes. *Geochim. Cosmochim. Acta* 40, 847–848.

Escoube, R., Rouxel, O.J., Pokrovsky, O.S., Schroth, A., Max Holmes, R., Donard, O.F.X., 2015. Iron isotope systematics in Arctic rivers. *Comptes Rendus - Geosci.* 347, 377–385. <https://doi.org/10.1016/j.crte.2015.04.005>

20 Escoube, R., Rouxel, O.J., Sholkovitz, E., Donard, O.F.X., 2009. Iron isotope systematics in estuaries: The case of North River, Massachusetts (USA). *Geochim. Cosmochim. Acta* 73, 4045–4059. <https://doi.org/10.1016/j.gca.2009.04.026>

Fantle, M.S., DePaolo, D.J., 2004. Iron isotopic fractionation during continental weathering. *Earth Planet. Sci. Lett.* 228, 547–562. <https://doi.org/10.1016/j.epsl.2004.10.013>

Figuères, G., Martin, J.M., Meybeck, M., 1978. Iron behaviour in the Zaire Estuary. *Netherlands J. Sea Res.* 12, 329–337.

25 Forsberg, J., Dahlqvist, R., Gelting-Nyström, J., Ingri, J., 2006. Trace metal speciation in brackish water using diffusive gradients in thin films and ultrafiltration: comparison of techniques. *Environ. Sci. Technol.* 40, 3901–5. <https://doi.org/10.1021/es0600781>

Guieu, C., Huang, W.W., Martin, J.M., Yong, Y.Y., 1996. Outflow of trace metals into the Laptev Sea by the Lena River. *Mar. Chem.* 53, 255–267. [https://doi.org/10.1016/0304-4203\(95\)00093-3](https://doi.org/10.1016/0304-4203(95)00093-3)

30 Guo, L., Santschi, P., 1996. A critical evaluation of cross-flow ultrafiltration technique for sampling colloidal organic carbon in seawater. *Mar. Chem.* 55, 113–127. [https://doi.org/10.1016/S0304-4203\(96\)00051-5](https://doi.org/10.1016/S0304-4203(96)00051-5)

Gustafsson, C., Gschwend, P.M., 1997. Aquatic colloids: Concepts, definitions, and current challenges. *Limnol. Oceanogr.* 42, 519–528. <https://doi.org/10.4319/lo.1997.42.3.0519>



Gustafsson, Ö., Widerlund, A., Andersson, P.S., Ingri, J., Roos, P., Ledin, A., 2000. Colloid dynamics and transport of major elements through a boreal river - Brackish bay mixing zone. *Mar. Chem.* 71, 1–21. [https://doi.org/10.1016/S0304-4203\(00\)00035-9](https://doi.org/10.1016/S0304-4203(00)00035-9)

Hirst, C., Andersson, P., Murphy, M.J., Schmitt, M., Kutscher, L., Maximov, T., Mörth, C.-M., Porcelli, D., n.d. Seasonal variations in the stable Fe isotope composition of particles and colloids in the Lena River basin, NE Russia.

Hirst, C., Andersson, P.S., Shaw, S., Burke, I.T., Kutscher, L., Murphy, M.J., Maximov, T., Pokrovsky, O.S., Mörth, C.M., Porcelli, D., 2017. Characterisation of Fe-bearing particles and colloids in the Lena River basin, NE Russia. *Geochim. Cosmochim. Acta* 213, 553–573. <https://doi.org/10.1016/j.gca.2017.07.012>

Holmes, R.M., McClelland, J.W., Peterson, B.J., Tank, S.E., Bulygina, E., Eglinton, T.I., Gordeev, V. V., Gurtovaya, T.Y., 10 Raymond, P.A., Repeta, D.J., Staples, R., Striegl, R.G., Zhulidov, A. V., Zimov, S.A., 2012. Seasonal and Annual Fluxes of Nutrients and Organic Matter from Large Rivers to the Arctic Ocean and Surrounding Seas. *Estuaries and Coasts* 35, 369–382. <https://doi.org/10.1007/s12237-011-9386-6>

Homoky, W.B., Severmann, S., McManus, J., Berelson, W.M., Riedel, T.E., Statham, P.J., Mills, R.A., 2012. Dissolved oxygen and suspended particles regulate the benthic flux of iron from continental margins. *Mar. Chem.* 134–135, 59–70.

15 <https://doi.org/10.1016/j.marchem.2012.03.003>

Icopini, G.A., Anbar, A.D., Ruebush, S.S., Tien, M., Brantley, S.L., 2004. Iron isotope fractionation during microbial reduction of iron: The importance of adsorption. *Geology* 32, 205–208. <https://doi.org/10.1130/G20184.1>

Ilina, S.M., Poitrasson, F., Lapitskiy, S.A., Alekhin, Y. V., 2013. Extreme iron isotope fractionation between different size colloids of boreal organic-rich waters. *Geochim. Cosmochim. Acta* 101, 96–111. <https://doi.org/10.1016/j.gca.2012.10.023>

20 Ingri, J., Widerlund, A., Land, M., Gustafsson, Ö., Andersson, P., Öhlander, B.,

Ingri, J., Malinovsky, D., Rodushkin, I., Baxter, D.C., Widerlund, A., Andersson, P., Gustafsson, Ö., Forsling, W., Öhlander, B., 2006. Iron isotope fractionation in river colloidal matter. *Earth Planet. Sci. Lett.* 245, 792–798. <https://doi.org/10.1016/j.epsl.2006.03.031>

Ingri, J., Widerlund, A., Land, M., Gustafsson, Ö., Andersson, P., Öhlander, B., 2000. Temporal variations in the fractionation 25 of the rare earth elements in a Boreal river; the role of colloidal particles. *Chem. Geol.* 166, 23–45. [https://doi.org/10.1016/S0009-2541\(99\)00178-3](https://doi.org/10.1016/S0009-2541(99)00178-3)

Karlsson, E., Gelting, J., Tesi, T., van Dongen, B., Andersson, A., Semiletov, I., Charkin, A., Dudarev, O., Gustafsson, Ö., 2016. Different sources and degradation state of dissolved, particulate, and sedimentary organic matter along the Eurasian Arctic coastal margin. *Global Biogeochem. Cycles*, 30, 898–919, doi:10.1002/2015GB005307

30 Klunder, M.B., Bauch, D., Laan, P., De Baar, H.J.W., Van Heuven, S., Ober, S., 2012. Dissolved iron in the Arctic shelf seas and surface waters of the central Arctic Ocean: Impact of Arctic river water and ice-melt. *J. Geophys. Res. Ocean.* 117, 1–18. <https://doi.org/10.1029/2011JC007133>

Kritzberg, E.S., Villanueva, A.B., Jung, M., Reader, H.E., 2014. Importance of boreal rivers in providing iron to marine waters. *PLoS One* 9. <https://doi.org/10.1371/journal.pone.0107500>



Kutscher, L., Mörth, C.M., Porcelli, D., Hirst, C., Maximov, T.C., Petrov, R.E., Andersson, P.S., 2017. Spatial variation in concentration and sources of organic carbon in the Lena River, Siberia. *J. Geophys. Res. Biogeosciences* 122, 1999–2016. <https://doi.org/10.1002/2017JG003858>

Laglera, L.M., Van Den Berg, C.M.G., 2009. Evidence for geochemical control of iron by humic substances in seawater. *Limnol. Oceanogr.* 54, 610–619. <https://doi.org/10.4319/lo.2009.54.2.0610>

Lalonde, K., Mucci, A., Ouellet, A., Gélinas, Y., 2012. Preservation of organic matter in sediments promoted by iron. *Nature* 483, 198–200. <https://doi.org/10.1038/nature10855>

Larsson, J., Ingri, J., Gustafsson, Ö., 2002. Evaluation and optimization of two complementary cross-flow ultrafiltration systems toward isolation of coastal surface water colloids. *Environ. Sci. Technol.*, Vol. 36, No. 10, 2236–2241.

10 <https://doi.org/10.1021/ES010325V>

Le Fouest, V., Babin, M., Tremblay, J.-É., 2013. The fate of riverine nutrients on Arctic shelves. *Biogeosciences* 10, 3661–3677. <https://doi.org/10.5194/bg-10-3661-2013>

Martin, J.M., Guan, D.M., Elbazpoulichet, F., Thomas, A.J., Gordeev, V. V., 1993. Preliminary Assessment of the Distributions of Some Trace-Elements (as, Cd, Cu, Fe, Ni, Pb and Zn) in a Pristine Aquatic Environment - the Lena River Estuary (Russia). *Mar. Chem.* 43, 185–199. [https://doi.org/10.1016/0304-4203\(93\)90224-C](https://doi.org/10.1016/0304-4203(93)90224-C)

Martin J.H., Gordon M.R., F.E.S., 1991. Iron Limitation? *Limnol. Ocean.* 36, 1793–1802. <https://doi.org/10.4319/lo.1991.36.8.1793>

Mulholland, D.S., Poitrasson, F., Boaventura, G.R., Allard, T., Vieira, L.C., Santos, R.V., Mancini, L., Seyler, P., 2015. Insights into iron sources and pathways in the Amazon River provided by isotopic and spectroscopic studies. *Geochim. Cosmochim. Acta* 150, 142–159. <https://doi.org/10.1016/j.gca.2014.12.004>

Neubauer, E., Köhler, S.J., Von Der Kammer, F., Laudon, H., Hofmann, T., 2013. Effect of pH and stream order on iron and arsenic speciation in boreal catchments. *Environ. Sci. Technol.* 47, 7120–7128. <https://doi.org/10.1021/es401193j>

Ödman, F., Ruth, T., Pontér, C., 1999. Validation of a field filtration technique for characterization of suspended particulate matter from freshwater. Part I. Major elements. *Appl. Geochemistry* 14, 301–317. [https://doi.org/10.1016/S0883-2927\(98\)00050-X](https://doi.org/10.1016/S0883-2927(98)00050-X)

Opsahl, S., Benner, R., Amon, R.M.W., 1999. Major flux of terrigenous dissolved organic matter through the Arctic Ocean. *Limnol. Oceanogr.* 44, 2017–2023. <https://doi.org/10.4319/lo.1999.44.8.2017>

Passow, U., 2004. Switching perspectives: Do mineral fluxes determine particulate organic carbon fluxes or vice versa? *Geochemistry, Geophys. Geosystems* 5, n/a-n/a. <https://doi.org/10.1029/2003GC000670>

30 Pédrot, M., Boudec, A. Le, Davranche, M., Dia, A., Henin, O., 2011. How does organic matter constrain the nature, size and availability of Fe nanoparticles for biological reduction? *J. Colloid Interface Sci.* 359, 75–85. <https://doi.org/10.1016/j.jcis.2011.03.067>

Perdue, E.M., Beck, K.C., Helmut Reuter, J., 1976. Organic complexes of iron and aluminium in natural waters. *Nature* 260, 418–420. <https://doi.org/10.1038/260418a0>



Pipko, I.I., S.P. Pugach, I.P. Semiletov, L.G. Anderson, N.E. Shakhova, Ö. Gustafsson, I.A. Repina, E. A. Spivak, A. N. Charkin, A. N. Salyuk, K. P. Shcherbakova, E. V. Panova, and Dudarev O. V.: The dynamics of the carbon dioxide system in the outer shelf and slope of the Eurasian Arctic Ocean //Ocean Sci., 13, 997–1016, 2017, <https://doi.org/10.5194/os-13-997-2017>

5 Pugach, S.P., Irina I. Pipko, Natalia E. Shakhova, Evgeny A. Shirshin, Irina V. Perminova, Örjan Gustafsson, Valery G. Bondur, and Semiletov I. P. DOM and its optical characteristics in the Laptev and East Siberian seas: Spatial distribution and inter-annual variability (2003–2011) // Ocean Sci., 14, 87–103, 2018, <https://doi.org/10.5194/os-14-87-2018>

Poitrasson, F., 2006. On the iron isotope homogeneity level of the continental crust. Chem. Geol. 235, 195–200. <https://doi.org/10.1016/j.chemgeo.2006.06.010>

10 Poitrasson, F., Cruz Vieira, L., Seyler, P., Márcia dos Santos Pinheiro, G., Santos Mulholland, D., Bonnet, M.P., Martinez, J.M., Alcantara Lima, B., Resende Boaventura, G., Chmeleff, J.Ô., Dantas, E.L., Guyot, J.L., Mancini, L., Martins Pimentel, M., Ventura Santos, R., Sondag, F., Vauchel, P., 2014. Iron isotope composition of the bulk waters and sediments from the Amazon River Basin. Chem. Geol. 377, 1–11. <https://doi.org/10.1016/j.chemgeo.2014.03.019>

Poitrasson, F., Freydier, R., 2005. Heavy iron isotope composition of granites determined by high resolution MC-ICP-MS. Chem. Geol. 222, 132–147. <https://doi.org/10.1016/j.chemgeo.2005.07.005>

Pokrovsky, O.S., Schott, J., 2002. Iron colloids/organic matter associated transport of major and trace elements in small boreal rivers and their estuaries (NW Russia). Chem. Geol. 190, 141–179. [https://doi.org/10.1016/S0009-2541\(02\)00115-8](https://doi.org/10.1016/S0009-2541(02)00115-8)

Pokrovsky, O.S., Schott, J., Dupré, B., 2006. Trace element fractionation and transport in boreal rivers and soil porewaters of permafrost-dominated basaltic terrain in Central Siberia. Geochim. Cosmochim. Acta 70, 3239–3260. <https://doi.org/10.1016/j.gca.2006.04.008>

15 Pokrovsky, O.S., Shirokova, L.S., Viers, J., Gordeev, V. V., Shevchenko, V.P., Chupakov, A. V., Vorobieva, T.Y., Candaupad, F., Causserand, C., Lanzanova, A., Zouiten, C., 2014. Fate of colloids during estuarine mixing in the Arctic. Ocean Sci. 10, 107–125. <https://doi.org/10.5194/os-10-107-2014>

Pokrovsky, O.S., Viers, J., Shirokova, L.S., Shevchenko, V.P., Filipov, A.S., Dupré, B., 2010. Dissolved, suspended, and 20 colloidal fluxes of organic carbon, major and trace elements in the Severnaya Dvina River and its tributary. Chem. Geol. 273, 136–149. <https://doi.org/10.1016/j.chemgeo.2010.02.018>

Pokrovsky, O.S., Shirokova, L.S., Zabelina, S.A., Vorobieva, T.Ya., Moreva, O.Yu., Klimov, S.I., Chupakov, A.V., Shorina, N.V., Kokryatskaya, N.M., Audry, S., Viers, J., Zoutien, C., Freydier, R., 2012. Size fractionation of trace elements in a seasonally stratified boreal lake: control of organic matter and iron colloids. Aquat. Geochem. 18, 115–139. <https://doi.org/10.1007/s10498-011-9154-z>

Poulton, S.W., Raiswell, R., 2005. Chemical and physical characteristics of iron oxides in riverine and glacial meltwater sediments. Chem. Geol. 218, 203–221. <https://doi.org/10.1016/j.chemgeo.2005.01.007>

Rachold, V., Alabyan, A., Hubberken, H.-W., Korotaev, V.N., Zaitsev, A.A., 1996. Sediment transport to the Laptev Sea—hydrology and geochemistry of the Lena River. Polar Res. 15, 183–196. <https://doi.org/10.3402/polar.v15i2.6646>



Raiswell, R., Canfield, D.E., 2012. The Iron Biogeochemical Cycle Past and Present. *Geochemical Perspect.* 1, 1–220. <https://doi.org/10.7185/geochempersp.1.1>

Raymond, P.A., McClelland, J.W., Holmes, R.M., Zhulidov, A. V., Mull, K., Peterson, B.J., Striegl, R.G., Aiken, G.R., Gurtovaya, T.Y., 2007. Flux and age of dissolved organic carbon exported to the Arctic Ocean: A carbon isotopic study of the five largest arctic rivers. *Global Biogeochem. Cycles* 21, 1–9. <https://doi.org/10.1029/2007GB002934>

Rodushkin, I., Nordlund, P., Engström, E., Baxter, D.C., 2005. Improved multi-elemental analyses by inductively coupled plasma-sector field mass spectrometry through methane addition to the plasma. *J. Anal. At. Spectrom.* 20, 1250–1255. <https://doi.org/10.1039/b507886e>

Rodushkin, I., Ruth, T., 1997. Determination of Trace Metals in Estuarine and Sea-water Reference Materials by High Resolution Inductively Coupled Plasma Mass Spectrometry. *J. Anal. At. Spectrom.* 12, 1181–1185. <https://doi.org/10.1039/a702486j>

Rosén, P.-O., Andersson, P.S., Alling, V., Mörth, C.-M., Björk, G., Semiletov, I., Porcelli, D., 2015. Ice export from the Laptev and East Siberian Sea derived from  $\delta^{18}\text{O}$  values. *J. Geophys. Res. Ocean.* 120, 5997–6007. <https://doi.org/10.1002/2015JC010866>

Rouxel, O., Sholkovitz, E., Charette, M., Edwards, K.J., 2008. Iron isotope fractionation in subterranean estuaries. *Geochim. Cosmochim. Acta* 72, 3413–3430. <https://doi.org/10.1016/j.gca.2008.05.001>

Rouxel, A., Bekker, K.J., Edwards, O.J., 2005. Iron isotope constraints on the Archaen and Paleoproterozoic Ocean redox state. *Science* (80-). 307, 1088–1091.

Salvadó, J.A., Tesi, T., Andersson, A., Ingrí, J., Dudarev, O. V., Semiletov, I.P., Gustafsson, Ö., 2015. Organic carbon remobilized from thawing permafrost is resequestered by reactive iron on the Eurasian Arctic Shelf. *Geophys. Res. Lett.* 42, 8122–8130. <https://doi.org/10.1002/2015GL066058>

Salvadó, J.A., Bröder, L., Andersson, A., Semiletov, I.P., Gustafsson, Ö., 2017. Release of black carbon from thawing permafrost estimated by sequestration fluxes in the East Siberian Arctic Shelf recipient. *Global Biogeochemical Cycles*, 31, 1501–1515. <https://doi.org/10.1002/2017GB005693>

Sánchez-García, L., Alling, V., Pugach, S., Vonk, J., Van Dongen, B., Humborg, C., Dudarev, O., Semiletov, I., Gustafsson, Ö., 2011. Inventories and behavior of particulate organic carbon in the Laptev and East Siberian seas. *Global Biogeochem. Cycles* 25, 1–13. <https://doi.org/10.1029/2010GB003862>

Semiletov, I.P., Pipko, I.I., Shakhova, N.E., Dudarev, O. V., Pugach, S.P., Charkin, A.N., McRoy, C.P., Kosmach, D., Gustafsson, Ö., 2011. Carbon transport by the Lena River from its headwaters to the Arctic Ocean, with emphasis on fluvial input of terrestrial particulate organic carbon vs. carbon transport by coastal erosion. *Biogeosciences* 8, 2407–2426. <https://doi.org/10.5194/bg-8-2407-2011>

Severmann, S., Johnson, C.M., Beard, B.L., McManus, J., 2006. The effect of early diagenesis on the Fe isotope compositions of porewaters and authigenic minerals in continental margin sediments. *Geochim. Cosmochim. Acta* 70, 2006–2022. <https://doi.org/10.1016/J.GCA.2006.01.007>



Sholkovitz, E.R., 1978. The flocculation of dissolved Fe, Mn, Al, Cu, Ni, Co and Cd during estuarine mixing. *Earth Planet. Sci. Lett.* 41, 77–86. [https://doi.org/10.1016/0012-821X\(78\)90043-2](https://doi.org/10.1016/0012-821X(78)90043-2)

Sholkovitz, E.R., 1976. Floculation of dissolved organic and inorganic matter during the mixing of river water and seawater. *Geochim. Cosmochim. Acta* 40, 831–845.

5 Skulan, J.L., Beard, B.L., Johnson, C.M., 2002. Kinetic and equilibrium Fe isotope fractionation between aqueous Fe(III) and hematite. *Geochim. Cosmochim. Acta* 66, 2995–3015. [https://doi.org/10.1016/S0016-7037\(02\)00902-X](https://doi.org/10.1016/S0016-7037(02)00902-X)

Slagter, H.A., Reader, H.E., Rijkenberg, M.J.A., Rutgers van der Loeff, M., De Baar, H.J.W.W., Gerringa, L.J.A., Salvadó, J.A., Tesi, T., Andersson, A., Ingri, J., Dudarev, O. V., Semiletov, I.P., Gustafsson, Ö., Kutscher, L., Mört, C.M., Porcelli, D., Hirst, C., Maximov, T.C., Petrov, R.E., Andersson, P.S., Klunder, M.B., Bauch, D., Laan, P., De Baar, H.J.W.W., Van Heuven, S., Ober, S., Forsberg, J., Dahlqvist, R., Gelting-Nyström, J., Ingri, J., Pokrovsky, O.S., Shirokova, L.S., Viers, J., Gordeev, V. V., Shevchenko, V.P., Chupakov, A. V., Vorobieva, T.Y., Caudadap, F., Causserand, C., Lanzanova, A., Zouiten, C., 2017. Trace metal speciation in brackish water using diffusive gradients in thin films and ultrafiltration: comparison of techniques. *J. Geophys. Res. Biogeosciences* 40, 8122–8130. <https://doi.org/10.1016/j.marchem.2017.10.005>

10 Staubwasser, M., Schoenberg, R., Von Blanckenburg, F., Krüger, S., Pohl, C., 2013. Isotope fractionation between dissolved and suspended particulate Fe in the oxic and anoxic water column of the Baltic Sea. *Biogeosciences* 10, 233–245. <https://doi.org/10.5194/bg-10-233-2013>

Stolpe, B., Guo, L., Shiller, A.M., 2013. Binding and transport of rare earth elements by organic and iron-rich nanocolloids in alaskan rivers, as revealed by field-flow fractionation and ICP-MS. *Geochim. Cosmochim. Acta* 106, 446–462. <https://doi.org/10.1016/j.gca.2012.12.033>

20 Stolpe, B., Hassellöv, M., 2007. Changes in size distribution of fresh water nanoscale colloidal matter and associated elements on mixing with seawater. *Geochim. Cosmochim. Acta* 71, 3292–3301. <https://doi.org/10.1016/j.gca.2007.04.025>

Sundman, A., Karlsson, T., Laudon, H., Persson, P., 2014. XAS study of iron speciation in soils and waters from a boreal catchment. *Chem. Geol.* 364, 93–102. <https://doi.org/10.1016/j.chemgeo.2013.11.023>

Sundman, A., Karlsson, T., Persson, P., 2013. An experimental protocol for structural characterization of Fe in dilute natural 25 waters. *Environ. Sci. Technol.* 47, 8557–8564. <https://doi.org/10.1021/es304630a>

Tagliabue, A., Bowie, A.R., Boyd, P.W., Buck, K.N., Johnson, K.S., Saito, M.A., 2017. The integral role of iron in ocean biogeochemistry. *Nature* 543, 51–59. <https://doi.org/10.1038/nature21058>

Thuróczy, C.-E., Gerringa, L.J.A., Klunder, M., Laan, P., Le Guittou, M., de Baar, H.J.W., 2011. Distinct trends in the speciation of iron between the shallow shelf seas and the deep basins of the Arctic Ocean. *J. Geophys. Res.* 116, C10009.

30 <https://doi.org/10.1029/2010JC006835>

Tipping, E., 1981. The adsorption of aquatic humic substances by iron oxides. *Geochim. Cosmochim. Acta* 45, 191–199. [https://doi.org/10.1016/0016-7037\(81\)90162-9](https://doi.org/10.1016/0016-7037(81)90162-9)

Vonk, J.E., van Dongen, B.E., Gustafsson, Ö., 2010. Selective preservation of old organic carbon fluvially released from sub-Arctic soils. *Geophys. Res. Lett.* 37, n/a-n/a. <https://doi.org/10.1029/2010GL042909>



Vonk, J.E., Sánchez-García, L., van Dongen, B.E., Alling, V., Kosmach, D., Charkin, A., Semiletov, I.P., Dudarev, O.V., Shakhova, N., Roos, P., Eglinton, T.I., Andersson, A., Gustafsson, Ö., 2012. Activation of old carbon by erosion of coastal and subsea permafrost in Arctic Siberia. *Nature* 489, 137-140. <https://doi.org/10.1038/nature11392>

Wagner, V., 1997. Analysis of a Russian landscape map and landscape classification for use in computer-aided forestry research. IIASA Interim Report IR-97-54. 56.

Walter, H., Breckle, S.-W., 2002. *Walter's Vegetation of the earth : the ecological systems of the geo-biosphere*. Springer.

Wedepohl, K.H., 1995. INGERSON LECTURE The composition of the continental crust. *Geochim. Cosmochim. Acta* 59, 1217-1232. [https://doi.org/10.1016/0016-7037\(95\)00038-2](https://doi.org/10.1016/0016-7037(95)00038-2)

Welch, S.A., Beard, B.L., Johnson, C.M., Braterman, P.S., 2003. Kinetic and equilibrium Fe isotope fractionation between aqueous Fe(II) and Fe(III). *Geochim. Cosmochim. Acta* 67, 4231-4250. [https://doi.org/10.1016/S0016-7037\(03\)00266-7](https://doi.org/10.1016/S0016-7037(03)00266-7)

Wiederhold, J.G., Kraemer, S.M., Teutsch, N., Borer, P.M., Halliday, A.N., Kretzschmar, R., 2006. Iron isotope fractionation during proton-promoted, ligand-controlled, and reductive dissolution of goethite. *Environ. Sci. Technol.* 40, 3787-3793. <https://doi.org/10.1021/es052228y>

Windom, H. L., Beck, K., Smith, R., 1971. Transport of trace metals to the Atlantic Ocean by three southeastern rivers. *Southeast Geol.* 12, 169-181.

Wu, L., Beard, B.L., Roden, E.E., Johnson, C.M., 2011. Stable Iron Isotope Fractionation Between Aqueous Fe(II) and Hydrous Ferric Oxide. *Environ. Sci. Technol.* 45, 1847-1852. <https://doi.org/10.1021/es103171x>

Yang, D., Kane, D.L., Hinzman, L.D., Zhang, X., Zhang, T., Ye, H., 2002. Siberian Lena River hydrologic regime and recent change. *J. Geophys. Res. Atmos.* 107, ACL 14-1-ACL 14-10. <https://doi.org/10.1029/2002JD002542>

Zhang, F., Zhu, X., Yan, B., Kendall, B., Peng, X., Li, J., Algeo, T.J., Romaniello, S., 2015. Oxygenation of a Cryogenian ocean (Nanhua Basin, South China) revealed by pyrite Fe isotope compositions. *Earth Planet. Sci. Lett.* 429, 11-19. <https://doi.org/10.1016/J.EPSL.2015.07.021>

Zhang, T., Barry, R.G., Knowles, K., Heginbottom, J.A., Brown, J., 1999. Statistics and characteristics of permafrost and ground-ice distribution in the Northern Hemisphere<sup>1</sup>. *Polar Geogr.* 23, 132-154. <https://doi.org/10.1080/10889379909377670>

25

30






Figure 1: Sampling stations in the Arctic Ocean. Black dots mark the stations in the detailed ESAS map. Along the Lena River-Laptev Sea transect membrane filtration and/or ultrafiltration was carried out. The sampling stations of this study follow the Lena River freshwater plume. The green numbers display  $\delta^{56}\text{Fe}$  values, measured in the uppermost sediment.




**Figure 2: The salinity gradient along the Lena River transect. The freshwater builds an almost 10m thick surface layer in the Laptev Sea, and the plume itself extends over an area of about 50 times 600km. The plume is divided into an inner and outer plume between station YS8 and YS11 by a sharp increase of salinity.**

5



**Figure 3: Dissolved (< 0.70 µm) and particulate (> 0.70 µm) organic carbon concentrations along the Lena River freshwater plume in the Laptev Sea. Close to the Lena River mouth POC constitutes about 20% of the TOC input, while at the outermost station it is only 2% of the TOC.**



5 **Figure 4: Particulate, colloidal and truly dissolved Fe concentrations along the Lena River freshwater plume. Concentrations of pFe and cFe decreased along the salinity gradient, while the concentrations of truly dissolved Fe is almost constant. Note the logarithmic scale and the sharp decrease of pFe between the inner and the outer plume.**




Figure 5: The colloidal and particulate Fe concentrations plotted versus salinity. Note the y-axis break due to the high range of pFe in the inner plume. The linear correlation between pFe and salinity is based on the data points below 1  $\mu\text{M}$  pFe. In the low salinity environment, the pFe is much higher compared to the cFe, whereas at salinities above 5 the differences are smaller.




Figure 6: Iron isotope values along the Lena River freshwater plume and the uppermost sediment of the ESAS. The error bars represent  $\pm 2 \sigma$ , in some cases the symbol is larger than the error. The  $\delta^{56}\text{Fe}$  values of pFe are negative at all stations, values close to zero close to the coast and more negative towards the open sea. The  $\delta^{56}\text{Fe}$  values of the cFe are negative in the inner plume and positive in the outer plume. The  $\delta^{56}\text{Fe}$  of the sediment samples were around -0.2 ‰, displaying the overall composition of the entire ESAS area.