

1 **Predominance of methanogens over methanotrophs in rewetted** 2 **fens characterized by high methane emissions**

3 Xi Wen^{1,14*}, Viktoria Unger^{2*}, Gerald Jurasinski², Franziska Koebsch², Fabian Horn¹, Gregor
4 Rehder³, Torsten Sachs⁴, Dominik Zak^{5,6}, Gunnar Lischeid^{7,8}, Klaus-Holger Knorr⁹, Michael E.
5 Böttcher¹⁰, Matthias Winkel^{1,11}, Paul L. E. Bodelier¹², and Susanne Liebner^{1,13}.

6 ¹Section 5.3 Geomicrobiology, GFZ German Research Centre for Geosciences, Helmholtz Centre
7 Potsdam, Telegrafenberg, Potsdam, 14473, Germany

8 ²Landscape Ecology and Site Evaluation, Faculty for Agricultural and Environmental Sciences,
9 Rostock University, Rostock, 18059, Germany

10 ³Department of Marine Chemistry, Leibniz Institute for Baltic Sea Research, Warnemünde, 18119,
11 Germany

12 ⁴Section 1.4 Remote Sensing, GFZ German Research Centre for Geosciences, Helmholtz Centre
13 Potsdam, Telegrafenberg, Potsdam, 14473, Germany

14 ⁵Department of Bioscience, Aarhus University, Silkeborg, 8600, Denmark

15 ⁶Department of Chemical Analytics and Biogeochemistry, Leibniz Institute of Freshwater Ecology
16 and Inland Fisheries, Berlin, 12587, Germany

17 ⁷Institute of Landscape Hydrology, Leibniz Center for Agricultural Landscape Research,
18 Münchberg, 15374, Germany

19 ⁸Institute of Earth and Environmental Science, University of Potsdam, Potsdam, 14476, Germany

20 ⁹Institute of Landscape Ecology, University of Münster, Münster, 48149, Germany

21 ¹⁰Geochemistry and Stable Isotope Biogeochemistry, Leibniz Institute for Baltic Sea Research,
22 Warnemünde, 18119, Germany

23 ¹¹Water and Environmental Research Center, Institute of Northern Engineering, University of
24 Alaska Fairbanks, 306 Tanana Loop, 99775, Fairbanks, AK, USA

25 ¹²Department of Microbial Ecology, Netherlands Institute of Ecology (NIOO-KNAW),
26 droevendaalsesteeg 10, Wageningen, 6708PB, the Netherlands

27 ¹³University of Potsdam, Institute of Biochemistry and Biology, Potsdam, 14469, Germany

28 ¹⁴College of Electrical Engineering, Northwest Minzu University, Lanzhou, 730070, China

29 *Correspondence to:* Viktoria Unger (viktoria.unger@uni-rostock.de), Franziska Koebsch
30 (franziska.koebsch@uni-rostock.de)

31 *Shared first authorship – the two first authors contributed equally to preparation of this work

32 **Abstract.** The rewetting of drained peatlands alters peat geochemistry and often leads to sustained
33 elevated methane emission. Although this methane is produced entirely by microbial activity, the

34 distribution and abundance of methane-cycling microbes in rewetted peatlands, especially in fens,
35 is rarely described. In this study, we compare the community composition and abundance of
36 methane-cycling microbes in relation to peat porewater geochemistry in two rewetted fens in
37 northeastern Germany, a coastal brackish fen and a freshwater riparian fen, with known high
38 methane fluxes. We utilized 16S rRNA high-throughput sequencing and quantitative polymerase
39 chain reaction on 16S rRNA, *mcrA*, and *pmoA* genes to determine microbial community
40 composition and the abundance of total bacteria, methanogens, and methanotrophs. Electrical
41 conductivity was more than three times higher in the coastal fen than in the riparian fen, averaging
42 5.3 and 1.5 mS cm⁻¹, respectively. Porewater concentrations of terminal electron acceptors varied
43 within and among the fens. This was also reflected in similarly high intra- and inter-site variations
44 of microbial community composition. Despite these differences in environmental conditions and
45 electron acceptor availability, we found a low abundance of methanotrophs and a high abundance
46 of methanogens, represented in particular by *Methanosaetaceae*, in both fens. This suggests that
47 rapid re/establishment of methanogens and slow re/establishment of methanotrophs contributes to
48 prolonged increased methane emissions following rewetting.

49 **1 Introduction**

50 Rewetting is a technique commonly employed to restore ecological and biogeochemical
51 functioning of drained fens. However, while rewetting may reduce carbon dioxide (CO₂) emissions
52 (Wilson et al. 2016), it often increases methane (CH₄) emissions in peatlands that remain inundated
53 following rewetting. On a 100-year time scale, CH₄ has a global warming potential 28 times
54 stronger than CO₂ (Myhre et al. 2013), and the factors that contribute to the magnitude and duration
55 of increased emissions are still uncertain (Joosten et al. 2015, Abdalla et al. 2016). Thus,
56 elucidating the dynamics of post-rewetting CH₄ exchange is of strong interest for both modelling
57 studies and peatland management projects (Abdalla et al. 2016). Although a recent increase in
58 rewetting projects in Germany and other European countries has prompted a number of studies of
59 methane cycling in rewetted peatlands (e.g., Jerman et al. 2009, Hahn-Schöfl et al. 2011, Urbanová
60 et al. 2013, Hahn et al. 2015, Vanselow-Algan et al. 2015, Zak et al. 2015, Emsens et al. 2016,
61 Putkinen et al. 2018), the post-rewetting distribution and abundance of methane-cycling microbes

62 in rewetted fens has seldom been examined (but see Juottonen et al. 2012, Urbanová et al. 2013,
63 Putkinen et al. 2018).

64 Peat CH₄ production and release is governed by a complex array of interrelated factors including
65 climate, water level, plant community, nutrient status, site geochemistry, and the activity of
66 microbes (i.e., bacteria and archaea) that use organic carbon as energy source (Segers 1998,
67 Abdalla et al. 2016). To date, the vast majority of studies in rewetted fens have focused on
68 quantifying CH₄ emission rates in association with environmental variables such as water level,
69 plant community, and aspects of site geochemistry (Abdalla et al. 2016). Site geochemistry indeed
70 plays an important role for methanogenic communities, as methanogenesis is suppressed in
71 presence of thermodynamically more favorable terminal electron acceptors (TEAs, Blodau 2011).
72 Due to a smaller pool of more favorable electron acceptors and high availability of organic carbon
73 substrates, organic-rich soils such as peat rapidly establish methanogenic conditions post-
74 rewetting (Segers 1998, Keller and Bridgham 2007, Knorr and Blodau 2009). Despite their
75 decisive role as producers (i.e., methanogens) and consumers (i.e., methanotrophs) of CH₄ (Conrad
76 1996), only a few studies have combined a characterization of the CH₄-cycling microbial
77 community, site geochemistry, and observed trends in CH₄ production. Existing studies have been
78 conducted in oligotrophic and mesotrophic boreal fens (e.g., Juottonen et al. 2005, Yrjälä et al.
79 2011, Juottonen et al. 2012), alpine fens (e.g., Liebner et al. 2012, Urbanová et al. 2013, Cheema
80 et al. 2015, Franchini et al. 2015), subarctic fens (Liebner et al. 2015), and incubation experiments
81 (e.g., Jerman et al. 2009, Knorr and Blodau 2009, Urbanová et al. 2011, Emsens et al. 2016).
82 Several studies on CH₄-cycling microbial communities have been conducted in minerotrophic
83 temperate fens (e.g., Cadillo-Quiroz et al. 2008, Liu et al. 2011, Sun et al. 2012, Zhou et al. 2017),
84 but these sites were not subject to drainage or rewetting. Direct comparisons of *in situ* abundances
85 of methanogens and methanotrophs in drained versus rewetted fens are scarce (Juottonen et al.
86 2012, Putkinen et al. 2018), and the studied sites, so far, are nutrient-poor fens with acidic
87 conditions.

88 While studies of nutrient-poor and mesotrophic boreal fens have documented post-rewetting CH₄
89 emissions comparable to or lower than at pristine sites (Komulainen et al. 1998, Tuittila et al. 2000,
90 Juottonen et al 2012), studies of temperate nutrient-rich fens have reported post-flooding CH₄
91 emissions dramatically exceeding emissions in pristine fens (e.g., Augustin and Chojnicki 2008,
92 Hahn et al. 2015). These high emissions typically occur together with a significant dieback in
93 vegetation, a mobilization of nutrients and electron acceptors in the upper peat layer, and increased
94 availability of dissolved organic matter (Zak and Gelbrecht 2007, Hahn-Schöfl et al. 2011, Hahn
95 et al. 2015, Jurasinski et al. 2016). High CH₄ fluxes may continue for decades following rewetting,
96 even in bogs (Vanselow-Algan et al. 2015). Hence, there is an urgent need to characterize CH₄-
97 cycling microbial communities and geochemical conditions in rewetted minerotrophic fens. In this
98 study, we therefore examined microbial community composition and abundance in relation to
99 post-flooding geochemical conditions in two rewetted fens in northeastern Germany. In both fens,
100 CH₄ emissions increased dramatically after rewetting, to over 200 g C m⁻² a⁻¹ (Augustin and
101 Chojnicki 2008, Hahn-Schöfl et al. 2011, Hahn et al. 2015, Jurasinski et al. 2016). Average annual
102 CH₄ emissions have decreased in both fens since the initial peak (Franz et al. 2016, Jurasinski et
103 al. 2016). Nevertheless, fluxes remained higher than under pre-flooding conditions (*ibid.*), and
104 higher than in pristine fens (Urbanová et al. 2013, Minke et al 2016). In the Hütelmoor in 2012,
105 average CH₄ emissions during the growing season were 40 g m⁻² (Koebsch et al. 2015). In
106 Zarnekow, average CH₄ emissions were 40 g m⁻² for the year 2013 (Franz et al. 2016). In
107 comparison, a recent review paper (Abdalla et al. 2016) estimated an average flux of 12 ± 21 g C
108 m⁻² a⁻¹ for pristine peatlands.

109 We expected patterns in microbial community composition would reflect the geochemical
110 conditions of the two sites and hypothesized a high abundance of methanogens relative to
111 methanotrophs in both fens. We also expected acetoclastic methanogens, which typically thrive in
112 nutrient-rich fens (Kelly et al. 1992, Galand 2005), to dominate the methanogenic community in
113 both fens.

114 2 Methods

115 2.1 Study sites

116 The nature reserve “Heiligensee and Hütelmoor” (‘Hütelmoor’ in the following, approx. 540 ha,
117 54°12'36.66" N, 12°10'34.28" E), is a coastal, mainly minerotrophic fen complex in Mecklenburg-
118 Vorpommern (NE Germany) that is separated from the Baltic Sea by a narrow (~100 m and less)
119 dune dike (Fig. 1a and b). The climate is temperate in the transition zone between maritime and
120 continental, with an average annual temperature of 9.1 °C and an average annual precipitation of
121 645 mm (data derived from grid product of the German Weather Service, reference climate period:
122 1981–2010). Episodic flooding from storm events delivers sediment and brackish water to the site
123 (Weisner and Schernewski 2013). The vegetation is a mixture of salt-tolerant macrophytes, with
124 dominant to semi-dominant stands of *Phragmites australis*, *Bolboschoenus maritimus*, *Carex*
125 *acutiformis*, and *Schoenoplectus tabernaemontani*. The dominating plants are interspersed with
126 open water bodies that are colonized by *Ceratophyllum demersum* in summer (Koch et al. 2017).
127 Intense draining and land amelioration practices began in the 1970s, which lowered the water level
128 to 1.6 m below ground surface and caused aerobic decomposition and concomitant degradation of
129 the peat (Voigtländer et al. 1996). The upper peat layer varies in depth between 0.6 and 3 m and
130 is highly degraded, reaching up to H10 on the von Post humification scale (Hahn et al. 2015).
131 Active draining ended in 1992, but dry conditions during summertime kept the water table well
132 below ground surface (Schönfeld-Bockholt et al. 2005, Koebisch et al. 2013) until concerns of
133 prolonged aerobic peat decomposition prompted the installation of a weir in 2009 at the outflow
134 of the catchment (Weisner and Schernewski 2013). After installation of the weir, the site has been
135 fully flooded year-round with an average water level of 0.6 m above the peat surface, and annual
136 average CH₄ flux increased ~186-fold from 0.0014 ± 0.0006 kg CH₄ m⁻² a⁻¹ to 0.26 ± 0.06 kg CH₄
137 m⁻² a⁻¹ (Hahn et al. 2015).

138 The study site polder Zarnekow ('Zarnekow' in the following, approx. 500 ha, 53°52'31.10" N,
139 12°53'19.60" E) is situated in the valley of the River Peene in Mecklenburg-Vorpommern (NE
140 Germany, Fig. 1a and c). The climate is slightly more continental compared to the Hütelmoor, with
141 a mean annual precipitation of 544 mm and a mean annual temperature of 8.7 °C (German Weather
142 Service, meteorological station Teterow, 24 km southwest of the study site; reference period 1981–
143 2010). The fen can be classified as a river valley mire system consisting of spring mires, wider
144 percolation mires, and flood mires along the River Peene. Drainage and low-intensity agricultural
145 use began in the eighteenth century when land-use changed to pastures and grassland. This was
146 intensified by active pumping in the mid-1970s. Due to land subsidence of several decimeters,
147 after rewetting (October 2004) water table depth increased to 0.1–0.5 m above peat surface. The
148 upper horizon is highly decomposed (0–0.3 m), followed by moderately decomposed peat to a
149 depth of 1 m and a deep layer of slightly decomposed peat up to a maximum depth of 10 m. The
150 open water bodies are densely colonized by *Ceratophyllum* spp. and *Typha latifolia* is the dominant
151 emergent macrophyte (Steffenhagen et al. 2012). Following flooding, CH₄ flux rates increased to
152 ~0.21 kg m⁻² a⁻¹ (Augustin and Chojnicki 2008). No pre-rewetting CH₄ flux data were available
153 for the Zarnekow site, but published CH₄ flux rates of representative drained fens from the same
154 region have been shown to be negligible, and many were CH₄ sinks (Augustin et al. 1998).

155 **2.2 Collection and analysis of peat cores and porewater samples**

156 Peat and porewater samples were collected at four different locations (n=4) in Hütelmoor (October
157 2014) and at five locations (n=5) in Zarnekow (July 2015) and spanned a distance of 1,200 m and
158 250 m, respectively, to cover the whole lateral extension at each site (Fig. 1b and c). Sampling
159 depths in the Hütelmoor were 0-5, 5-10, 10-20, 20-30, 30-40, and 40-50 cm below the peat surface,
160 except for core numbers 1 and 4 where samples could only be obtained up to a depth of 10-20 and
161 30-40 cm, respectively. Sampling depths in Zarnekow were 0-5, 25-30, and 50-55 cm below the
162 peat surface. Previous work at Zarnekow has revealed little variation in peat properties with depth
163 (e.g., Zak and Gelbrecht 2007), hence, a lower depth resolution in Zarnekow cores was chosen for

164 this study. Peat cores were collected with a Perspex liner (ID: 60 mm, Hütelmoor) and a peat auger
165 (Zarnekow). In order to minimize oxygen contamination, the outer layer of the peat core was
166 omitted. Subsamples for molecular analysis were immediately packed in 50 ml sterile Falcon tubes
167 and stored at -80 °C until further processing.

168 Pore waters in the Hütelmoor were collected with a stainless-steel push-point sampler attached to
169 a plastic syringe to recover the samples from 10 cm depth intervals. Samples were immediately
170 filtered with 0.45 µm membrane sterile, disposable syringe filters. Pore waters in Zarnekow were
171 sampled with permanently installed dialysis samplers consisting of slotted polypropylene (PP)
172 pipes (length: 636 mm, ID: 34 mm) surrounded with 0.22 µm polyethersulfone membrane. The
173 PP pipes were fixed at distinct peat depths (surface level, 20 and 40 cm depth) and connected with
174 PP tubes (4x6 mm IDxAD). Water samples were drawn out from the dialysis sampler pipes with
175 a syringe through the PP tube. Due to practical restrictions in accessibility and sampling,
176 permanent dialysis samplers could not be installed at the desired locations in the Hütelmoor,
177 resulting in the different sampling techniques described above.

178 At both sites, electrical conductivity (EC), dissolved oxygen (DO), and pH were measured
179 immediately after sampling (Sentix 41 pH probe and a TetraCon 325 conductivity measuring cell
180 attached to a WTW multi 340i handheld; WTW, Weilheim). In this paper, EC is presented and
181 was not converted to salinity (i.e., psu), as a conversion would be imprecise for brackish waters.
182 A simplified equation for conversion can be found in Schemel (2001). Headspace CH₄
183 concentrations of porewater samples were measured with an Agilent 7890A gas chromatograph
184 (Agilent Technologies, Germany) equipped with a flame ionization detector and a Carboxen PLOT
185 Capillary Column or HP-Plot Q (Porapak-Q) column. The measured headspace CH₄ concentration
186 was then converted into a dissolved CH₄ concentration using the temperature-corrected solubility
187 coefficient (Wilhelm et al. 1977). Isotopic composition of dissolved CH₄ for Hütelmoor was
188 analyzed using the gas chromatography-combustion-technique (GC-C) and the gas
189 chromatography-high-temperature-conversion-technique (GC-HTC). The gas was directly

190 injected in a Gas Chromatograph Agilent 7890A, CH₄ was quantitatively converted to CO₂, and
191 the δ¹³C values were then measured with the isotope-ratio-mass-spectrometer MAT-253 (Thermo
192 Finnigan, Germany). The δ¹³C of dissolved CH₄ in Zarnekow was analyzed using a laser-based
193 isotope analyzer equipped with a small sample isotope module for analyses of discrete gas samples
194 (cavity ring down spectroscopy CRDS; Picarro G2201-I, Santa Clara, CA, USA). Calibration was
195 carried out before, during and after analyses using certified standards of known isotopic
196 composition (obtained from Isometric Instruments, Victoria, BC, Canada, and from Westfalen AG,
197 Münster, Germany). Reproducibility of results was typically +/- 1 ‰. In the presence of high
198 concentrations of hydrogen sulfide interfering with laser-based isotope analysis, samples were
199 treated with iron(III) sulfate to oxidize and/or precipitate sulfide. For both sites, sulfate and nitrate
200 concentrations were analyzed by ion chromatography (IC, Thermo Fisher Scientific Dionex) using
201 an Ion Pac AS-9-HC 4 column, partly after dilution of the sample. Dissolved metal concentrations
202 were analyzed by ICP-OES (iCAP 6300 DUO, Thermo Fisher Scientific). Accuracy and precision
203 were routinely checked with a certified CASS standard as previously described (Kowalski et al.
204 2012).

205 For the incubation experiments, peat cores were collected from Zarnekow in March 2012 using a
206 modified Kajak Corer with a plexiglass tube. The intact cores were placed in a cool box and
207 immediately transported to the Leibniz-Institute of Freshwater Ecology and Inland Fisheries in
208 Berlin where they were sectioned into a total of 12 samples. Fresh, surficial organic sediment (0-
209 10 cm depth, 6 individual samples) was separated from the bulk peat (10-20 cm depth, 6 individual
210 samples) and the samples were placed in 60 ml plastic cups. The cups were filled completely and
211 closed with air-tight caps to minimize oxygen contamination. The samples were then express-
212 shipped (< 24 hours) to the lab at the Netherlands Institute of Ecology for immediate processing
213 and analysis. For CH₄ production incubations, 5 g of material and 10 ml of nitrogen (N₂)-flushed
214 MilliQ water was weighed into three (n=3) 150 mL flasks for both surficial organic sediment and
215 bulk peat. The flasks were capped with rubber stoppers, flushed with N₂ for approximately one

216 hour, then incubated stationary at 20°C in the dark. For CH₄ oxidation incubations, 5 g of fresh
217 material and 10 ml of MilliQ water was weighed into three 150 mL flasks for both surficial organic
218 sediment and bulk peat. The flasks were capped with rubber stoppers and 1.4 ml of pure CH₄ was
219 added to obtain a headspace CH₄ concentration of approximately 10,000 ppm. Incubations were
220 performed in the dark at 20°C on a gyratory shaker (120 rpm). For all incubations, headspace CH₄
221 concentration was determined using a gas chromatograph equipped with a flame ionization
222 detector on days 1, 3, 5, and 8 of the incubation. Potential CH₄ production and oxidation rate were
223 determined by linear regression of CH₄ concentration over all sampling times.

224

225 **2.3 Gene amplification and phylogenetic analysis**

226 Genomic DNA was extracted from 0.2–0.3 g of duplicates of peat soil per sample using an EurX
227 Soil DNA Kit (Roboklon, Berlin, Germany). DNA concentrations were quantified with a
228 Nanophotometer P360 (Implen GmbH, München, DE) and Qubit 2.0 Fluorometer (Thermo Fisher
229 Scientific, Darmstadt, Germany). Polymerase chain reaction (PCR) amplification of bacterial and
230 archaeal 16S rRNA genes was performed using the primer combination of S-D-Bact-0341-b-S-
231 17/S-D-Bact-0785-a-A-21 (Herlemann et al. 2011) and S-D-Arch-0349-a-S-17/S-D-Arch-0786-a-
232 A-20 (Takai and Horikoshi 2000), respectively, with barcodes contained in the 5'-end. The PCR
233 mix contained 1x PCR buffer (Tris•Cl, KCl, (NH₄)₂SO₄, 15 mM MgCl₂; pH 8.7) (QIAGEN,
234 Hilden, Germany), 0.5 μM of each primer (Biomers, Ulm, Germany), 0.2 mM of each
235 deoxynucleoside (Thermo Fisher Scientific, Darmstadt, Germany), and 0.025 U μl⁻¹ hot start
236 polymerase (QIAGEN, Hilden, Germany). PCR samples were kept at 95 °C for 5 min to denature
237 the DNA, with amplification proceeding for 40 cycles at 95 °C for 1 min, 56 °C for 45 s and 72
238 °C for 90 s; a final extension of 10 min at 72 °C was added to ensure complete amplification. PCR
239 products were purified with a Hi Yield Gel/PCR DNA fragment extraction kit (Süd-Laborbedarf,
240 Gauting, Germany). To reduce amplification bias, PCR products of three individual runs per
241 sample were combined. PCR products of different samples were pooled in equimolar

242 concentrations and compressed to a final volume of 10 μl with a concentration of 200 $\text{ng } \mu\text{l}^{-1}$ in a
243 vacuum centrifuge Concentrator Plus (Eppendorf, Hamburg, Germany).

244 Illumina sequencing was performed by GATC Biotech AG using 300 bp paired-end mode and a
245 20% PhiX Control v3 library to counteract the effects of low-diversity sequence libraries. Raw
246 data was demultiplexed using an own script based on CutAdapt (Martin 2011). Ambiguous
247 nucleotides at sequence ends were trimmed and a 10% mismatch was allowed for primer
248 identification, whereas barcode sequences needed to be present without any mismatches and with
249 a minimum Phred-Score of Q25 for each nucleotide. After sorting, overlapping paired-end reads
250 were merged using PEAR [Q25, p 0.0001, v20] (Zhang et al. 2014). The orientation of the merged
251 sequences was standardized according to the barcode information obtained from demultiplexing.
252 Low-quality reads were removed using Trimmomatic [SE, LEADING Q25, TRAILING Q25,
253 SLIDINGWINDOW 5:25; MINLEN 200] (Bolger et al. 2014). Chimeric sequences were removed
254 using USEARCH 6.1 and the QIIME-script identify_chimeric_seqs.py (Caporaso et al. 2010). Pre-
255 processed sequences were taxonomically assigned to operational taxonomic units (OTUs) at a
256 nucleotide sequence identity of 97% using QIIME's pick_open_reference_otus.py script and the
257 GreenGenes database 13.05 (McDonald et al. 2012) as reference. The taxonomic assignment of
258 representative sequences was further checked for correct taxonomical classification by
259 phylogenetic tree calculations in the ARB environment referenced against the SILVA database
260 (<https://www.arb-silva.de>) version 119 (Quast et al. 2013). The resulting OTU table was filtered
261 for singletons, OTUs assigned to chloroplasts or mitochondria, and for low-abundance OTUs
262 (below 0.2% within each sample). Archaeal and bacterial samples were processed separately while
263 only OTUs that were assigned to the respective domain were considered for further analysis. For
264 archaea, a total of 6,844,177 valid sequences were obtained, ranging from 60,496 to 398,660 in
265 individual samples. These sequences were classified into 402 OTUs. For bacteria, a total of
266 2,586,148 valid sequences were obtained, ranging from 22,826 to 164,916 in individual samples.
267 These sequences were classified into 843 OTUs. The OTU tables were then collapsed at a higher

268 taxonomic level to generate the bubble plots. The 16S rRNA gene sequence data have been
269 deposited at NCBI under the Bioproject PRJNA356778. Hütelmoor sequence read archive
270 accession numbers are SRR5118134-SRR5118155 for bacterial and SRR5119428-SRR5119449
271 for archaeal sequences, respectively. Zarnekow accession numbers are SRR6854018-
272 SRR6854033 and SRR6854205-SRR6854220 for bacterial and archaeal sequences, respectively.

273 **2.4 qPCR analysis**

274 Quantitative polymerase chain reaction (qPCR) for the determination of methanotrophic and
275 methanogenic functional gene copy numbers and overall bacterial 16S rRNA gene copy numbers
276 was performed via SybrGreen assays on a Bio-Rad CFX instrument (Bio-Rad, Munich, Germany)
277 with slight modifications after Liebner et al. (2015). The functional methanotrophic *pmoA* gene
278 was amplified with the primer combination A189F/Mb661 (Kolb et al. 2003) suitable for detecting
279 all known aerobic methanotrophic Proteobacteria. Annealing was done at 55 °C after a 7-cycle-
280 step touchdown starting at 62 °C. The functional methanogenic *mcrA* gene was amplified with the
281 *mlas/mcrA*-rev primer pair (Steinberg and Regan 2009) with annealing at 57 °C. The bacterial 16S
282 rRNA gene was quantified with the primers Eub341F/Eub534R according to Degelmann et al.
283 (2010) with annealing at 58 °C. Different DNA template concentrations were tested prior to the
284 qPCR runs to determine optimal template concentration without inhibitions through co-extracts.
285 The 25 µl reactions contained 12.5 µl of iTaq universal Sybr Green supermix (Bio-Rad, Munich,
286 Germany), 0.25 µM concentrations of the primers, and 5 µl of DNA template. Data acquisition
287 was always done at 80 °C to avoid quantification of primer dimers. The specificity of each run
288 was verified through melt-curve analysis and gel electrophoresis. Only runs with efficiencies
289 between 82 and 105% were used for further analysis. Measurements were performed in duplicates.
290 The ratio of methanogens to methanotrophs was determined based on gene abundances of *mcrA*
291 and *pmoA*. The marker gene for the soluble monooxygenase, *mmoX*, was neglected due to the
292 absence of *Methylocella* in the sequencing data (Fig. 4).

293 **2.5 Data visualization and statistical analysis**

294 All data visualization and statistical analysis were done in R (R Core Team). The taxonomic
295 relative abundances across samples were visualized through bubble plots with the R package
296 ggplot2 (Wickham 2009). Differences in microbial community composition were visualized with
297 2-dimensional non-metric multidimensional scaling (NMDS) based on Bray-Curtis distances. The
298 NMDS ordinations were constructed using R package vegan (Oksanen et al. 2017). An
299 environmental fit was performed on the ordinations to determine the measured geochemical
300 parameters that may influence community composition. The geochemical data were fitted to the
301 ordinations as vectors with a significance of $p < 0.05$. Depth profiles were constructed with the
302 porewater geochemical data, as well as with the microbial abundances, to elucidate depthwise
303 trends and assess whether differences in microbial community and abundances among the two fens
304 are related to differences in their respective geochemistry.

305

306 **3 Results**

307 **3.1 Environmental characteristics and site geochemistry**

308 The two rewetted fens varied substantially in their environmental characteristics (e.g., proximity
309 to the sea) and porewater geochemistry (Fig. 2, Tables 1 and 2). EC was more than three times
310 higher in Hütelmoor than in Zarnekow, averaging 5.3 and 1.5 mS cm^{-1} , respectively. Mean values
311 of pH were approximately neutral (6.5 to 7.0) in the upper peat profile and comparable in both
312 fens until a depth of about 30 cm where pH decreased to ~ 6 in the Hütelmoor. Concentrations of
313 the TEAs nitrate and sulfate were lower in Zarnekow and near zero in the pore water at all depths,
314 while nitrate and sulfate were abundant in the upper and lower peat profile in Hütelmoor at ~ 1.5
315 to 3.0 mM and ~ 4 to 20 mM, respectively (Fig. 2). Iron concentrations were higher in the
316 Hütelmoor pore water, while manganese concentrations were higher in Zarnekow pore water.
317 Dissolved oxygen concentrations in the upper peat profile (i.e. 0 to 25 cm depths) were much
318 higher in Hütelmoor than in Zarnekow (Fig. 2). Here DO concentrations averaged ~ 0.25 mM until
319 a depth of 15 cm at which they dropped sharply, reaching concentrations slightly below 0.05 mM

320 at 25 cm. In Zarnekow, DO concentrations did not exceed 0.1 mM and varied little with depth.
321 Regarding geochemical conditions, Hütelmoor core (HC) 1 differed from all other Hütelmoor
322 cores and was more similar to Zarnekow cores. In HC 1 – the core taken nearest to potential
323 freshwater sources (Fig. 1b) – pore water EC and DO concentrations were lower while pH was
324 slightly higher than in all other Hütelmoor cores. Moreover, this was the only Hütelmoor core
325 where nitrate concentrations were below detection limit (0.001mM) (Fig. 2). In all cores we found
326 high concentrations of dissolved CH₄ that varied within and among fens and were slightly higher
327 in Zarnekow pore water. Stable isotope ratios of $\delta^{13}\text{C-CH}_4$ (Fig. 2) in the upper peat (approx.
328 -59‰) suggest a predominance of acetoclastic methanogenesis, with a shift to hydrogenotrophic
329 methanogenesis around -65‰ in the lower peat profile. Additionally, the observed shifts toward
330 less negative $\delta^{13}\text{C-CH}_4$ values in the upper peat layer, as in HC 1 and HC 2, could indicate partial
331 oxidation of CH₄ occurred (Chasar et al. 2000).

332 **3.2 Community composition of bacteria and archaea**

333 Bacterial sequences could be affiliated into a total of 30 bacterial phyla (Fig. 3). Among them,
334 Proteobacteria, Acidobacteria, Actinobacteria, Chloroflexi, Nitrospirae and Bacteroidetes were
335 present in all samples. With mean relative abundance of 48%, Proteobacteria was the most
336 abundant phylum. Some taxa (e.g., Verrucomicrobia, Atribacteria (OP9), and AD3) were present
337 only in Hütelmoor. Variation in community composition was larger in Hütelmoor samples than in
338 Zarnekow. Within Proteobacteria, the alpha subdivision was the most dominant group, having
339 contributed 26.7% to all the libraries on average (Fig. 4). The family *Hyphomicrobiaceae*
340 dominated the Alphaproteobacteria, and was distributed evenly across samples, but missing in the
341 surface and bottom peat layers in HC 2. In addition, methanotrophs were clearly in low abundance
342 across all samples, representing only 0.06% and 0.05% of the bacterial community in Hütelmoor
343 and Zarnekow, respectively. Of the few methanotrophs that were detected, type II methanotrophs
344 (mainly *Methylocystaceae*) outcompeted type I methanotrophs (mainly *Methylococcaceae*) in the
345 community, while members of the genus *Methylocella* were absent (Fig. 4).

346 Within the archaeal community, Bathyarchaeota were mostly dominating over Euryarchaeota (Fig.
347 5). The MCG group (mainly the order of pGrfC26) in Bathyarchaeota prevailed across all samples
348 but was especially abundant in HC 2 samples. In addition to Bathyarchaeota, methanogenic
349 archaea were important, and on average contributed 30.6% to the whole archaeal community.
350 Among the methanogens, acetoclastic methanogens were more abundant in most of the samples
351 and *Methanosaetaceae* (24.8%) were the major component. They were present in most samples
352 and much more dominant than *Methanosarcinaceae* (2.0%). Hydrogenotrophic methanogens, such
353 as *Methanomassiliicoccaceae* (1.6%), *Methanoregulaceae* (1.2%) and *Methanocellaceae* (0.6%),
354 albeit low in abundance, were detected in many samples. Hütelmoor samples displayed greater
355 variability in archaeal community composition compared to Zarnekow samples. The putative
356 anaerobic methanotrophs of the ANME-2D (Raghoebarsing et al. 2006) clade occurred in patchy
357 abundance with dominance in single spots of both sites. In HC 1 they represented a mean relative
358 abundance of 40.9% of total archaeal reads but were almost absent in all other Hütelmoor cores.
359 In Zarnekow core (ZC) 3, ANME-2D represented up to approximately 30% of all archaea but were
360 otherwise low in abundance.

361 **3.3 Environmental drivers of microbial community composition**

362 Bacterial and archaeal population at both peatland sites showed distinct clustering (Fig. 6) with
363 similarly high intra- and inter-site variations but greater overall variation in community
364 composition in the Hütelmoor. Community composition varied much more strongly in HC 2 than
365 in any other core (Fig. 6). Bacterial communities in HC 1 were more similar to communities in all
366 Zarnekow cores than in other Hütelmoor cores (Fig. 6a). The archaeal community in HC 1 was
367 more similar to Zarnekow cores as well (Fig. 6b). Environmental fit vectors suggest pH, oxygen
368 and alternative TEA availability as important factors influencing microbial community
369 composition. The EC vector suggests the importance of brackish conditions in shaping microbial
370 communities in the Hütelmoor (Fig. 6a - c).

371 **3.4 Total microbial and functional gene abundances**

372 Quantitative PCR results show that in both fens, *mcrA* abundance is up to two orders of magnitude
373 greater than *pmoA* abundance (Fig. 7, Tables 1 and 2). Gene copy numbers of *mcrA* are overall
374 higher and spatially more stable in Zarnekow than in Hütelmoor. Total microbial abundance
375 declined with depth more strongly in Hütelmoor than in Zarnekow (Fig. 7). There was a
376 pronounced decrease in microbial abundances at 20 cm depth in the Hütelmoor. For example, 16S
377 rRNA gene and *pmoA* gene copy numbers in deeper samples (below 20 cm depth) are one order
378 of magnitude lower than in upper samples on average, while the *mcrA* gene abundance are
379 approximately two orders of magnitude lower. Hütelmoor samples also exhibited larger
380 heterogeneity in terms of abundances than Zarnekow samples. Contrary to previous studies,
381 methanotroph abundance did not correlate with dissolved CH₄ or oxygen concentrations.

382

383 **4 Discussion**

384 **4.1 Fen geochemistry and relations to microbial community composition**

385 The rewetting of drained fens promotes elevated CH₄ production and emission, which can
386 potentially offset carbon sink benefits. Few studies have attempted to link microbial community
387 dynamics and site geochemistry with observed patterns in CH₄ production and/or emission in
388 rewetted fens, while such data are crucial for predicting long-term changes to CH₄ cycling (Galand
389 et al. 2002, Yrjälä et al. 2011, Juottonen et al. 2012). In this study, we show that CH₄-cycling
390 microbial community composition is related to patterns in site geochemistry in two rewetted fens
391 with high CH₄ emissions, high methanogen abundances, and low methanotroph abundances. Our
392 results suggest that high methanogen abundances concurrent with low methanotroph abundances
393 are characteristic of rewetted fens with ongoing high CH₄ emissions. Thus, we present microbial
394 evidence for sustained elevated CH₄ emissions in mostly inundated rewetted temperate fens.

395 The environmental conditions and associated geochemistry of the two rewetted fens were largely
396 different. Depth profiles of porewater geochemical parameters show the fens differed in EC
397 throughout the entire peat profile, while pH and concentrations of alternative TEAs differed at

398 certain depths. In general, concentrations of TEAs oxygen, sulfate, nitrate, and iron were higher
399 in the Hütelmoor. In Zarnekow, geochemical conditions varied little across the fen and along the
400 peat depth profiles (Fig. 2). As expected, the geochemical heterogeneity was reflected in microbial
401 community structure in both sites, suggesting the importance of environmental characteristics and
402 associated geochemical conditions as drivers of microbial community composition (Figs. 2, 3, 4,
403 6). The NMDS ordinations (Fig. 6) show large variation in archaeal and bacterial community
404 composition in the coastal brackish fen, and much less variation in the freshwater riparian fen.
405 Environmental fit vectors (Fig. 6) suggest that salinity (indicated by the EC vector), pH, oxygen
406 and alternative TEA availability are the most important measured factors influencing microbial
407 communities in the two fens. Patterns in microbial community composition have previously been
408 linked to salinity (e.g., Chambers et al. 2016), pH (e.g., Yrjälä et al. 2011), and TEA availability
409 in peatlands (e.g., He et al. 2015).

410 Comparing the geochemical depth profiles (Fig. 2) with the relative abundance of bacteria and
411 archaea (Figs. 3 and 4) provides a more complete picture of the relationships between microbial
412 communities and site geochemistry, particularly with respect to TEA utilization. While the
413 porewater depth profiles suggest there is little nitrate available for microbial use in HC 1, the
414 relative abundance plot for Archaea showed that this core was dominated by ANME-2D. ANME-
415 2D were recently discovered to be anaerobic methanotrophs that oxidize CH₄ performing reverse
416 methanogenesis using nitrate as an electron acceptor (Haroon et al. 2013). However, ANME-2D
417 has also been implicated in the iron-mediated anaerobic oxidation of methane (Ettwig et al. 2016),
418 and the HC 1 site showed slightly higher total iron concentrations. The relevance of ANME-2D as
419 CH₄ oxidizers in terrestrial habitats is still not clear. Rewetting converts the fens into widely
420 anaerobic conditions, thus providing conditions suitable for the establishment of anaerobic
421 oxidation of methane, but this has yet to be demonstrated in fens. The patchy yet locally high
422 abundance of ANME-2D both in Hütelmoor and in Zarnekow suggests an ecological relevance of
423 this group. Shifts towards less negative $\delta^{13}\text{C}$ -CH₄ signatures in the upper peat profile, for example,

424 from -65 to -60‰ in HC 1 (where ANME-2D was abundant), may indicate that partial oxidation
425 of CH₄ occurred, but we could only speculate whether or not ANME-2D are actively involved in
426 this CH₄ oxidation.

427 Although TEA input may be higher in the Hütelmoor, here, methanogenic conditions also
428 predominate. This finding contrasts the measured oxygen concentrations in the upper peat profile,
429 as methanogenesis under persistently oxygenated conditions is thermodynamically not possible.
430 However, seasonal analysis of oxygen concentrations in both sites suggests highly fluctuating
431 oxygen regimes both spatially and temporary (data not shown). Such non-uniform distribution of
432 redox processes has already been described elsewhere, in particular for methanogenesis (Hoehler
433 et al. 2001, Knorr et al. 2009). It is possible that oxygen levels in both fens are highly variable,
434 allowing for both aerobic and anaerobic carbon turnover processes. Recent studies from wetlands
435 also show that methanogenesis can occur in aerobic layers, driven mainly by *Methanosaeta*
436 (Narrowe et al. 2017, Wagner 2017), which were detected in a high abundance in this study (Fig.
437 5). Further, oxygen may not necessarily be available within aggregates entailing anaerobic
438 pathways and thus, the existence of anaerobic microenvironments may also partially explain the
439 seemingly contradictory co-occurrence of oxygen and the highly abundant methanogens.
440 Anaerobic conditions are also reflected by the extensive and stable occurrence of the strictly
441 anaerobic syntrophs (e.g., *Syntrophobacteraceae*, *Syntrophaceae*) in most samples, even in the top
442 centimeters. This suggests that syntrophic degradation of organic material is taking place in the
443 uppermost layer and the fermented substances are readily available for methanogens. As
444 geochemistry and microbial community composition differ among the sites in this study, it is thus
445 notable that a similarly high abundance of methanogens, and low abundance of methanotrophs
446 was detected in both fens. The dominance of methanogens implies that readily available substrates
447 and favorable geochemical conditions promote high anaerobic carbon turnover despite seasonally
448 fluctuating oxygen concentrations in the upper peat layer.

449 **4.2 Low methanotroph abundances in rewetted fens**

450 Methanogens (mainly *Methanosaetaceae*) dominated nearly all of the various niches detected in
451 this study, while methanotrophs were highly under-represented in both sites (Figs. 3 and 4).
452 Functional and ribosomal gene copy numbers not only show a high ratio of methanogen to
453 methanotroph abundance (Fig. 7), irrespective of site and time of sampling, but also a small
454 contribution of methanotrophs to total bacterial population in both sites. Methanotrophs constitute
455 only ~0.06% of the total bacterial population in the Hütelmoor and ~0.05% at Zarnekow. It should
456 be noted that in this study we measured only gene abundances and not transcript abundances, and
457 the pool both of active methanogens and methanotrophs was likely smaller than the numbers
458 presented here (Freitag and Prosser 2009, Freitag et al. 2010, Cheema et al. 2015, Franchini et al.
459 2015). Also, as we were unable to obtain microbial samples from before rewetting, a direct
460 comparison of microbial abundances was not possible. This was therefore, not a study of rewetting
461 effects. For this reason, we performed an exhaustive literature search on relevant studies of pristine
462 fens. Compared to pristine fens, we detected a low abundance of methanotrophs. Liebner et al.
463 (2015), for example, found methanotrophs represented 0.5% of the total bacterial community in a
464 pristine, subarctic transitional bog/fen palsa, while *mcrA* and *pmoA* abundances were nearly
465 identical. In a pristine Swiss alpine fen, Liebner et al. (2012) found methanotrophs generally
466 outnumbered methanogens by an order of magnitude. Cheema et al. (2015) and Franchini et al.
467 (2015) reported *mcrA* abundances higher than *pmoA* abundances by only one order of magnitude
468 in a separate Swiss alpine fen. In the rewetted fens in our study, *mcrA* gene abundance was up to
469 two orders of magnitude higher than *pmoA* abundance (Fig. 7). Due to inevitable differences in
470 methodology and equipment, direct comparisons of absolute gene abundances are limited.
471 Therefore, only the abundances of methanotrophs relative to methanogens and relative to the total
472 bacterial community were compared, rather than absolute abundances. We are confident that this
473 kind of ‘normalization’ can mitigate the bias of different experiments and allows a comparison of
474 sites. Further, all primers and equipment used in this study were identical to those used by Liebner
475 et al. (2012, 2015), making the comparison more reliable.

476 As most methanotrophs live along the oxic-anoxic boundary of the peat surface and plant roots
477 therein (Le Mer and Roger 2001), the low methanotroph abundances in both fens could be
478 explained by disturbances to this boundary zone and associated geochemical pathways following
479 inundation. In rewetted fens, a massive plant dieback has been observed along with strong changes
480 in surface peat geochemistry (Hahn-Schöfl et al. 2011, Hahn et al. 2015). In addition to substrate
481 (i.e. CH₄) availability, oxygen availability is the most important factor governing the activity of
482 most methanotrophs (Le Mer and Roger 2001, Hernandez et al. 2015). The anoxic conditions at
483 the peat surface caused by inundation may have disturbed existing methanotrophic niches, either
484 directly by habitat destruction, and/or indirectly by promoting the growth of organisms that are
485 able to outcompete methanotrophs for oxygen. Heterotrophic organisms, for example, have been
486 shown to outcompete methanotrophs for oxygen when oxygen concentrations are greater than 5
487 μM (van Bodegom et al. 2001). Our microbial data support this conclusion, as
488 *Hyphomicrobiaceae*, most of which are aerobic heterotrophs, was the most abundant bacterial
489 family in both fens. Incubation data from Zarnekow (Fig. S1) show that the CH₄ oxidation potential
490 is high, however incubations provide ideal conditions for methanotrophs and thus only potential
491 rates. It is likely that, *in situ*, the activity of methanotrophs is overprinted by the activity of
492 competitive organisms such as heterotrophs. It is also possible that methane oxidation may occur
493 in the water column above the peat surface, but this was beyond the scope of this study.
494 Nevertheless, it is low enough that methane production and emissions remain high, as
495 demonstrated by the high dissolved CH₄ concentrations and ongoing high fluxes.

496 Comparable studies have so far been conducted in nutrient-poor or mesotrophic fens where post-
497 rewetting CH₄ emissions, though higher than pre-rewetting, did not exceed those of similar pristine
498 sites (e.g., Yrjälä et al. 2011, Juottonen et al. 2005, Juottonen et al. 2012). Nevertheless, there is
499 mounting evidence linking CH₄-cycling microbe abundances to CH₄ dynamics in rewetted fens.
500 Juottonen et al. (2012), for example, compared *pmoA* gene abundances in three natural and three
501 rewetted fens and found them to be lower in rewetted sites. The same study also measured a lower

502 abundance of *mcrA* genes in rewetted sites, which was attributed to a lack of available labile
503 organic carbon compounds. In peatlands, and especially fens, litter and root exudates from vascular
504 plants can stimulate CH₄ emissions (Megonigal et al. 2005, Bridgham et al. 2013, Agethen and
505 Knorr 2018), and excess labile substrate has been proposed as one reason for substantial increases
506 in CH₄ emissions in rewetted fens (Hahn-Schöfl et al. 2011). Future studies should compare pre-
507 and post-rewetting microbial abundances along with changes in CH₄ emissions, plant
508 communities, and peat geochemistry to better assess the effect rewetting has on the CH₄-cycling
509 microbial community.

510

511 **5 Conclusion**

512 Despite a recent increase in the number of rewetting projects in Northern Europe, few studies have
513 characterized CH₄-cycling microbes in restored peatlands, especially fens. In this study, we show
514 that rewetted fens differing in geochemical conditions and microbial community composition have
515 a similarly low abundance of methanotrophs, a high abundance of methanogens, and an established
516 anaerobic carbon cycling microbial community. Comparing these data to pristine wetlands with
517 lower CH₄ emission rates, we found that pristine wetlands have a higher abundance of
518 methanotrophs than measured in the fens in this study, suggesting the inundation and associated
519 anoxia caused by flooding may disturb methanotrophic niches and negatively affect the ability of
520 methanotrophic communities to establish. The abundances of methane producers and consumers
521 are thus suggested as indicators of continued elevated CH₄ emissions following the rewetting of
522 drained fens. Management decisions regarding rewetting processes should consider that
523 disturbances to methanotrophic niches is possible if rewetting leads to long-term inundation of the
524 peat surface.

525

526 **6 Data availability**

527 The 16S rRNA gene sequence data have been deposited at NCBI under the Bioproject
528 PRJNA356778. Hütelmoor sequence read archive accession numbers are SRR5118134-
529 SRR5118155 for bacterial and SRR5119428-SRR5119449 for archaeal sequences, respectively.
530 Zarnekow accession numbers are SRR6854018-SRR6854033 and SRR6854205-SRR6854220 for
531 bacterial and archaeal sequences, respectively.

532

533 **7 Author contribution**

534 FK, SL, GJ, XW, and VU formulated the research questions and study design. FK, MW, SL and
535 TS performed field work. SL, GJ, XW, VU and TS visualized the data and prepared maps. XW
536 and VU prepared the original draft. SL, MEB, KK, GL, and DZ provided analytical data
537 measurements. FH conducted the bioinformatics analyses. PB provided the incubation data and
538 prepared the supplemental figure. All authors contributed to the discussion and interpretation of
539 the data and the writing of the paper.

540

541 **8 Competing interests**

542 The authors declare that they have no conflict of interest.

543

544 **9 Acknowledgements**

545 This study was conducted within the framework of the Research Training Group 'Baltic
546 TRANSCOAST' funded by the DFG (Deutsche Forschungsgemeinschaft) under grant number
547 GRK 2000. This is Baltic TRANSCOAST publication no. GRK2000/000X. The financial support
548 to Xi Wen (Grant No. 201408620031 to X.W.) provided by the China Scholarship Council (CSC),
549 and to Matthias Winkel (ARCSS-1500931) provided by the National Science Foundation (NSF),
550 is gratefully acknowledged. This study was supported by the Helmholtz Gemeinschaft (HGF) by
551 funding the Helmholtz Young Investigators Group of S.L. (VH-NG-919) and T.S. (Grant VH-NG-
552 821), a Helmholtz Postdoc Programme grant to F.K. (Grant PD-129), and further supported by the

553 Terrestrial Environmental Observatories (TERENO) Network. The Leibniz Institute for Baltic Sea
554 Research (IOW) is also acknowledged for funding the lab work in this study. The European Social
555 Fund (ESF) and the Ministry of Education, Science and Culture of Mecklenburg-Western
556 Pomerania funded this work within the scope of the project WETSCAPES (ESF/14-BM-A55-
557 0030/16). Dr. Matthias Gehre, head of the Laboratory of Stable Isotopes at the Helmholtz Centre
558 for Environmental Research, is acknowledged for providing carbon isotope measurements for this
559 study. Anke Saborowski and Anne Köhler are also acknowledged for support in the laboratory.

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585 **10 References**

- 586
- 587 Abdalla, M., Hastings, A., Truu, J., Espenberg, M., Mander, U., and Smith, P.: Emissions of
588 methane from northern peatlands: a review of management impacts and implications for future
589 management options, *Ecology and Evolution*, 6, 7080-7102, doi:10.1002/ece3.2469, 2016.
- 590
- 591 Agethen, S. and Knorr, K.-H.: *Juncus effusus* mono-stands in restored cutover peat bogs – Analysis
592 of litter quality, controls of anaerobic decomposition, and the risk of secondary carbon loss. *Soil*
593 *Biology and Biochemistry*, 117, 139-152, doi:10.1016/j.soilbio.2017.11.020, 2018.
- 594
- 595 Augustin, J., Merbach, W., and Rogasik, J.: Factors influencing nitrous oxide and methane
596 emissions from minerotrophic fens in northeast Germany, *Biology and Fertility of Soils*, 28(1), 1-
597 4, doi:10.1007/s003740050455, 1998.
- 598
- 599 Augustin, J. and Chojnicki, B.: Austausch von klimarelevanten Spurengasen, Klimawirkung und
600 Kohlenstoffdynamik in den ersten Jahren nach der Wiedervernässung von degradiertem
601 Niedermoorgrünland (Exchange of climate relevant trace gases, climate effect and carbon
602 dynamics in the first years after re-wetting of degraded fen grassland), In: Gelbrecht, J., Zak, D.,
603 and Augustin, J. (eds.), *Phosphor- und Kohlenstoff- Dynamik und Vegetationsentwicklung in*
604 *wiedervernässten Mooren des Peenetales in Mecklenburg-Vorpommern (Phosphorus and carbon*
605 *dynamics and vegetation development in re-wetted peatland of the Peene valley in Mecklenburg-*
606 *Western Pomerania)*, Leibniz-Institut für Gewässerökologie und und Binnenfischerei, Berlin, pp.
607 50–67 (in German), 2008.
- 608
- 609 Blodau, C.: Thermodynamic control on terminal electron transfer and methanogenesis. In:
610 Tratnyek, P. G., Grundl, T. J., and Haderlein, S. B. (eds.), *Aquatic Redox Chemistry*, ACS
611 *Symposium Series*, American Chemical Society, distributed in print by Oxford University Press
612 Inc., Washington, DC, pp. 65-82., doi:10.1021/bk-2011-1071.ch004, 2011.
- 613
- 614 Bolger, A. M., Lohse, M., and Usadel, B.: Trimmomatic: a flexible trimmer for Illumina sequence
615 data, *Bioinformatics*, 30, 2114-2120, doi:10.1093/bioinformatics/btu170., 2014.
- 616
- 617 Bridgman, S. D., Cadillo-Quiroz, H., Keller, J. K., and Zhuang, Q.: Methane emissions from
618 wetlands: biogeochemical, microbial, and modeling perspectives from local to global scales,
619 *Global Change Biology* 19, 1325-1356, doi:10.1111/gcb.12131, 2013.
- 620
- 621 Cadillo-Quiroz, H., Yashiro, E., Yavitt, J. B., and Zinder, S. H.: Characterization of the archaeal
622 community in a minerotrophic fen and terminal restriction fragment length polymorphism-directed

623 isolation of a novel hydrogenotrophic methanogen, *Applied and Environmental Microbiology*
624 74(7), 2059-2068, doi:10.1128/AEM.02222-07, 2008.
625

626 Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K.,
627 Fierer, N., Peña, A. G., Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T., Knights, D.,
628 Koenig, J. E., Ley, R. E., Lozupone, C. A., McDonald, D., Muegge, B. D., Pirrung, M., Reeder,
629 J., Sevinsky, J. R., Turnbaugh, P. J., Walters, W. A., Widmann, J., Yatsunenko, T., Zaneveld, J.,
630 and Knight, R.: QIIME allows analysis of high-throughput community sequencing data, *Nature*
631 *Methods*, 7(5), 335-336, doi:10.1038/nmeth.f.303, 2010.
632

633 Chambers, L. G., Guevara, R., Boyer, J. N., Troxler, T. G., and Davis, S. E.: Effects of salinity
634 and inundation on microbial community structure and function in a mangrove peat soil, *Wetlands*,
635 36(2), 361-371, doi:10.1007/s13157-016-0745-8, 2016.
636

637 Chasar, L. S., Chanton, J. P., Glaser, P. H., and Siegel, D. I.: Methane concentration and stable
638 isotope distribution as evidence of rhizospheric processes: comparison of a fen and bog in the
639 Glacial Lake Agassiz Peatland complex, *Annals of Botany*, 86, 655-663,
640 doi:10.1006/anbo.2000.1172, 2000.
641

642 Cheema, S., Zeyer, J., and Henneberger, R.: Methanotrophic and methanogenic communities in
643 Swiss alpine fens dominated by *Carex rostrata* and *Eriophorum angustifolium*, *Applied and*
644 *Environmental Microbiology*, 81(17), 5832-5844, doi:10.1128/AEM.01519-15, 2015.
645

646 Conrad, R.: Soil microorganisms as controllers of atmospheric trace gases (H₂, CO, CH₄, OCS,
647 N₂O, and NO), *Microbiological Reviews*, 60(4), 609-640, 1996.
648

649 Degelmann, D. M., Borken, W., Drake, H. L., and Kolb, S.: Different atmospheric methane-
650 oxidizing communities in European beech and Norway spruce soils, *Applied and Environmental*
651 *Microbiology*, 76, 3228-3235, doi:10.1128/AEM.02730-09, 2010.
652

653 Emsens, W.-J., Aggenbach, C. J. S., Schoutens, K., Smolders, A. J. P., Zak, D., and van Diggelen,
654 R.: Soil iron content as a predictor of carbon and nutrient mobilization in rewetted fens, *PLoS*
655 *ONE*, 11(4), e0153166, doi:10.1371/journal.pone.0153166, 2016.
656

657 Franchini, A. G., Henneberger, R., Aeppli, M., and Zeyer, J.: Methane dynamics in an alpine fen:
658 a field-based study on methanogenic and methanotrophic microbial communities, *FEMS*
659 *Microbiology Ecology*, 91(3), 1-13, doi:10.1093/femsec/fiu032, 2015.
660

661 Franz, D., Koebisch, F., Larmanou, E., Augustin, J., and Sachs, T.: High net CO₂ and CH₄ release
662 at a eutrophic shallow lake on a formerly drained fen, *Biogeosciences*, 13, 3051-3070,
663 doi:10.5194/bg-13-3051-2016, 2016.
664

665 Freitag, T. E. and Prosser, J. I.: Correlation of methane production and functional gene
666 transcription activity in a peat soil, *Applied and Environmental Microbiology*, 75(21), 6679-6687,
667 doi:10.1128/AEM.01021-09, 2009.
668

669 Freitag, T. E., Toet, S., Ineson, P., and Prosser, J. I.: Links between methane flux and
670 transcriptional activities of methanogens and methane oxidizers in a blanket peat bog, *FEMS*
671 *Microbiology Ecology*, 73, 157-165, doi:10.1111/j.1574-6941.2010.00871.x, 2010.
672

673 Galand, P. E., Saarnio, S., Fritze, H., and Yrjälä, K.: Depth related diversity of methanogen
674 Archaea in Finnish oligotrophic fen, *FEMS Microbiology Ecology*, 42, 441-449,
675 doi:10.1111/j.1574-6941.2002.tb01033.x., 2002.
676

677 Galand, P. E., Fritze, H., Conrad, R., and Yrjälä, K.: Pathways for methanogenesis and diversity
678 of methanogenic archaea in three boreal peatland ecosystems, *Applied and Environmental*
679 *Microbiology*, 71(4), 2195-2198, doi:10.1128/AEM.71.4.2195-2198.2005, 2005.
680

681 Hahn, J., Köhler, S., Glatzel, S., and Jurasinski, G.: Methane exchange in a coastal fen the first
682 year after flooding – a systems shift, *PLOS ONE*, 10(10), e0140657, doi:10.1371/journal.
683 pone.0140657, 2015.
684

685 Hahn-Schöfl, M., Zak, D., Mincke, M., Gelbrecht, J., Augustin, J., and Freibauer, A.: Organic
686 sediment formed during inundation of a degraded fen grassland emits large fluxes of CH₄ and CO₂,
687 *Biogeosciences*, 8, 1539-1550, doi:10.5194/bg-8-1539-2011, 2011.
688

689 Haroon, M. F., Hu, S., Shi, Y., Imelfort, M., Keller, J., Hugenholtz, P., Yuan, Z., and Tyson, G.:
690 Anaerobic oxidation of methane couple to nitrate reduction in a novel archaeal lineage, *Nature*,
691 500, 567-570, doi:10.1038/nature12375, 2013.
692

693 He, S., Malfatti, S. A., McFarland, J. W., Anderson, F. E., Pati, A., Huntemann, M., Tremblay, J.,
694 del Rio, T. G., Waldrop, M. P., Windham-Myers, L., and Tringe, S. G.: Patterns in wetland
695 microbial community composition and functional gene repertoire associated with methane
696 emissions, *mBio*, 6(3), 1-15, doi:10.1128/mBio.00066-15, 2015.
697

698 Herlemann, D. P., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J. J., and Andersson, A.F.:
699 Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea, *ISME*
700 *J*, 5(10), 1571-1579, doi:10.1038/ismej.2011.41., 2011.
701

702 Hernandez, M. E., Beck, D. A., Lidstrom, M. E., and Chistoserdova, L.: Oxygen availability is a
703 major factor in determining the composition of microbial communities involved in methane
704 oxidation, *PeerJ*, 3, e801, doi:10.7717/peerj.801, 2015.
705

706 Hoehler, T. M., Alperin, M. J., Albert, D. B., and Martens, C. S.: Apparent minimum free energy
707 requirements for methanogenic Archaea and sulfate-reducing bacteria in an anoxic marine
708 sediment, *FEMS Microbiology Ecology*, 38, 33-41, doi:10.1016/S0168-6496(01)00175-1, 2001.
709

710 Jerman, V., Metje, M., Mandic-Mulec, I., and Frenzel, P.: Wetland restoration and
711 methanogenesis: the activity of microbial populations and competition for substrates at different
712 temperatures, *Biogeosciences*, 6, 1127-1138, doi:10.5194/bg-6-1127-2009, 2009.
713

714 Joosten, H., Brust, K., Couwenberg, J., Gerner, A., Holsten, B., Permien, T., Schäfer, A.,
715 Tanneberger, F., Trepel, M., and Wahren, A.: MoorFutures® Integration of additional ecosystem
716 services (including biodiversity) into carbon credits – standard, methodology and transferability
717 to other regions, Bundesamt für Naturschutz (Federal Ministry for the Environment, BfN), BfN-
718 Skripten 407, Bonn, Germany, 2015.
719

720 Juottonen, H., Galand, P. E., Tuittila, E.-S., Laine, J., Fritze, H., Yrjälä, K.: Methanogen
721 communities and bacteria along an ecohydrological gradient in a northern raised bog complex,
722 *Environmental microbiology*, 7(10), 1547-1557, doi: 10.3389/fmicb.2015.00356, 2015.
723

724 Juottonen, H., Hynninen, A., Nieminen, M., Tuomivirta, T. T., Tuittila, E.-S., Nousiainen, H.,
725 Kell, D. K., Yrjälä, K., Tervahauta, A., and Fritze, H.: Methane-cycling microbial communities
726 and methane emission in natural and restored peatlands, *Applied and Environmental*
727 *Microbiology*, 78(17), 6386-6389, doi:10.1128/AEM.00261-12, 2012.
728

729 Jurasinski, G., Glatzel, S., Hahn, J., Koch, S., Koch, M., and Koebisch, F.: Turn on, fade out –
730 Methane exchange in a coastal fen over a period of six years after rewetting, *Geophysical Research*
731 *Abstracts*, 18, EGU2016-14899, 2016.
732

733 Keller, J. K. and Bridgman, S. D.: Pathways of anaerobic carbon cycling across an ombrotrophic-
734 minerotrophic peatland gradient, *Limnology and Oceanography*, 52(1), 96-107,
735 doi:10.4319/lo.2007.52.1.0096, 2007.
736

737 Kelly, C. A., Dice, N. B. and Martens, C. S.: Temporal variations in the stable carbon isotopic
738 composition of methane emitted from Minnesota peatlands, *Global Biogeochemical Cycles*, 6(3),
739 263-269, doi:10.1029/92GB01478, 1992.
740

741 Knorr, K.-H., Lischeid, G., and Blodau, C.: Dynamics of redox processes in a minerotrophic fen
742 exposed to a water table manipulation, *Geoderma*, 153, 379-392,
743 doi:10.1016/j.geoderma.2009.08.023, 2009.
744

745 Knorr, K.-H. and Blodau, C.: Impact of experimental drought and rewetting on redox
746 transformations and methanogenesis in mesocosms of a northern fen soil, *Soil Biology and*
747 *Biochemistry*, 1187-1198, doi:10.1016/j.soilbio.2009.02.030, 2009.
748

749 Koch, M., Koebisch, F., Hahn, J., and Jurasinski, G.: From meadow to shallow lake: Monitoring
750 secondary succession in a coastal fen after rewetting by flooding based on aerial imagery and plot
751 data, *Mires and Peat*, 19(11), 1-17, doi:10.19189/MaP.2015.OMB.188, 2017.
752

753 Koebisch, F., Glatzel, S., and Jurasinski, G.: Vegetation controls emissions in a coastal brackish
754 fen, *Wetlands Ecology and Management*, 21(5), 323-337, doi:10.1007/s11273-013-9304-8, 2013.
755

756 Koebisch, F., Jurasinski, G., Koch, M., Hofmann, J., and Glatzel, S.: Controls for multi-scale
757 temporal variation in ecosystem methane exchange during the growing season of a permanently
758 inundated fen, *Agricultural and Forest Meteorology*, 204, 94-105,
759 doi:10.1016/j.agrformet.2015.02.002, 2015.
760

761 Kolb, S., Knief, C., Stubner, S., and Conrad, R.: Quantitative detection of methanotrophs in soil
762 by novel *pmoA*-targeted real-time PCR assays, *Applied and Environmental Microbiology*, 69,
763 2423-2429, doi:10.1128/AEM.69.5.2423-2429.2003, 2003.
764

765 Komulainen, V.-M., Nykanen, H., Martikainen, P. J., and Laine, J.: Short-term effect of restoration
766 on vegetation change and methane emissions from peatlands drained for forestry in southern
767 Finland, *Canadian Journal of Forest Research*, 28, 402-411, doi:10.1139/x98-011, 1998.
768

769 Kowalski, N., Dellwig, O., Beck, M., Grunwald, M., Dürselen, C.-D., Badewien, T. H., Brumsack,
770 H.-J., van Beusekom, J. E. E., and Böttcher, M. E.: A comparative study of manganese dynamics
771 in the water column and sediments of intertidal systems of the North Sea, *Estuarine, Coastal and*
772 *Shelf Science*, 100, 3-17, doi:10.1016/j.ecss.2011.03.011, 2012.
773

774 Le Mer, J. and Roger. P.: Production, oxidation, emission, and consumption of methane by soils:
775 a review, *European Journal of Soil Biology*, 37, 25-50, doi:10.1016/S1164-5563(01)01067-6,
776 2001.
777

778 Liebner, S., Schwarzenbach, S. P., and Zeyer, J.: Methane emissions from an alpine fen in central
779 Switzerland, *Biogeochemistry*, 109, 287-299, doi:10.1007/s10533-011-9629-4, 2012.
780

781 Liebner, S., Ganzert, L., Kiss, A., Yang, S., Wagner, D., and Svenning, M. M.: Shifts in
782 methanogenic community composition and methane fluxes along the degradation of discontinuous
783 permafrost, *Frontiers in Microbiology*, 6(356), 1-10, doi:10.3389/fmicb.2015.00356, 2015.
784

785 Liu, D. Y., Ding, W. X., Jia, Z. J., and Cai, Z. C.: Relation between methanogenic archaea and
786 methane production potential in selected and natural wetland ecosystems across China,
787 *Biogeosciences*, 8, 329-338, doi:10.5194/bg-8-329-2011, 2011.
788

789 Martin, M.: Cutadapt removes adapter sequences from high-throughput sequencing reads.
790 *EMBnet. Journal*, 17(1), 10-12, doi:10.14806/ej.17.1.200, 2011.
791

792 McDonald, D., Price, M. N., Goodrich, J., Nawrocki, E. P., DeSantis, T. Z., Probst, A., Andersen,
793 G. L., Knight, R., and Hugenholtz, P.: An improved Greengenes taxonomy with explicit ranks for
794 ecological and evolutionary analyses of bacteria and archaea, *ISME J*, 6(3), 610-618,
795 doi:10.1038%2Fismej.2011.139, 2012.
796

797 Megonigal, J. P., Mines, M. E., and Visscher, P. T.: Anaerobic metabolism: linkages to trace gases
798 and aerobic processes, In: Schlesinger, W.H. (ed.), *Biogeochemistry*, Elsevier, Oxford, UK, pp.
799 350-362, 2005.
800

801 Minke, M., Augustin, J., Burlo, A., Yarmashuk, T., Chuvashova, H., Thiele, A., Freibauer, A.,
802 Tikhonov, V., and Hoffman, M.: Water level, vegetation composition, and plant productivity
803 explain greenhouse gas fluxes in temperate cutover fens after inundation, *Biogeosciences*, 13,
804 3945-3970. doi:10.5194/bg-13-3945-2016, 2016.
805

806 Myhre, G., Shindell, D., Breon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque,
807 J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Rotstayn, L., Stephens, G., and Zhang, H.:
808 Anthropogenic and natural radiative forcing. Chapter 8. In: Stocker, T. F., Qin, D., Plattner, G.-
809 K., Tignor, M., Allen, D., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M. (eds.),
810 *Climate Change 2013. The Physical Science Basis. Contribution of Working Group I to the Fifth*
811 *Assessment Report of the Intergovernmental Panel on Climate Change.*, Cambridge University
812 Press, Cambridge, UK and New York, USA, pp. 659–740, 2013.

813
814 Narrowe, A. B., Angle, J. C., Daly, R. A., Stefanik, K. C., Wrighton, K. C., and Miller, C. S.:
815 High-resolution sequencing reveals unexplored archaeal diversity in freshwater wetland soils,
816 *Environmental Microbiology*, 19(6), 2192-2209, doi: 10.1111/1462-2920.13703, 2017.
817
818 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P. R.,
819 O'Hara, R. B., Simpson, G. L., Solymos, Stevens, M. H. H., Szoecs, E., and Wagner, H.: *vegan*:
820 *Community Ecology Package*. R package version 2.4-5. [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)
821 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan), 2017.
822
823 Putkinen, A., Tuittila, E.-S., Siljanen, H. M .P, Bodrossy, L., and Fritze, H.: Recovery of methane
824 turnover and the associated microbial communities in restored cutover peatlands is strongly linked
825 with increasing sphagnum abundance, *Soil Biology and Biochemistry*, 116, 110-119, doi:
826 10.1016/j.soilbio.2017.10.005, 2018.
827
828 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner,
829 F.O.: The SILVA ribosomal RNA gene database project: improved data processing and web-based
830 tools, *Nucleic Acids Research*, 41, D590-596, doi:10.1093/nar/gks1219., 2013.
831
832 Raghoebarsing, A. A., Pol, A., van de Pas-Schoonen, K. T., Smolders, A. J. P., Ettwig, K. F.,
833 Rijpstra, W. I. C., Schouten, S., Damste, J. S. S., Op den Camp, H. J. M., Jetten, M. S. M., and
834 Strous, M.: A microbial consortium couples anaerobic methane oxidation to denitrification,
835 *Nature*, 440, 918–921, doi:10.1038/nature04617, 2006.
836
837 R Core Team: *R: A language and environment for statistical computing*. R Foundation for
838 *Statistical Computing*, Vienna, Austria, <https://www.R-project.org/>, 2017.
839
840 Schemel, L.: Simplified conversions between specific conductance and salinity units for use with
841 data from monitoring stations, *Interagency Ecological Program Newsletter*, 14(1), 17-18, 2001.
842
843 Schönfeld-Bockholt, R., Roth, D., and Dittmann, L. *Friends of the natural history in Mecklenburg*.
844 *Ch. Teilflächenbezogene ökologische und futterwirtschaftliche Beurteilung des Grünlandes im*
845 *Naturschutzgebiet Heiligensee und Hütelmoor*, 2005.
846
847 Segers, R.: Methane production and methane consumption: a review of processes underlying
848 wetland methane fluxes, *Biogeochemistry*, 41, 23-51, doi:10.1023/A:1005929032764, 1998.
849

850 Steffenhagen, P., Zak, D., Schulz, K., Timmerman, T., and Zerbe, S.: Biomass and nutrient stock
851 of submersed and floating macrophytes in shallow lakes formed by rewetting of degraded fens,
852 *Hydrobiologia*, 692(1), 99-109, doi:10.1007/s10750-011-0833-y, 2012.
853

854 Steinberg, L. M. and Regan, J. M.: *mcrA*-targeted real-time quantitative PCR method to examine
855 methanogen communities, *Applied and Environmental Microbiology*, 75, 4435-4442,
856 doi:10.1128%2FAEM.02858-08, 2009.
857

858 Sun, C. L., Brauer, S. L., Cadillo-Quiroz, H., Zinder, S. H., and Yavitt, J. B.: Seasonal changes in
859 methanogenesis and methanogenic community in three peatlands, New York State, *Frontiers in*
860 *Microbiology*, 3(81), 1-8, doi:10.3389/fmicb.2012.00081, 2012.
861

862 Takai, K., and Horikoshi, K.: Rapid detection and quantification of members of the archaeal
863 community by quantitative PCR using fluorogenic probes, *Applied and Environmental*
864 *Microbiology*, 66, 5066-5072, 2000.
865

866 Tuittila, E.-S., Komulainen, V. M., Vasander, H., Nykänen, H., Martikainen, P. J., and Laine, J.:
867 Methane dynamics of a restored cut-away peatland, *Global Change Biology*, 6, 569-581,
868 doi:10.1046/j.1365-2486.2000.00341.x, 2000.
869

870 Urbanová, Z., Picek, T., and Bárta, J.: Effect of re-wetting on carbon and nutrient fluxes,
871 greenhouse gas production, and diversity of methanogenic archaeal community, *Ecological*
872 *Engineering*, 37, 1017-1026, doi:10.1016/j.ecoleng.2010.07.012, 2011.
873

874 Urbanová, Z., Bárta, J., and Picek, T.: Methane emissions and methanogenic archaea on pristine,
875 drained and restored mountain peatlands, central Europe, *Ecosystems*, 16(4), 664-677,
876 doi:10.1007/s10021-013-9637-4, 2013.
877


878 Vanselow-Algan, M., Schmidt, S. R., Greven, M., Fiencke, C., Kutzbach, L., and Pfeiffer, E.-M.:
879 High methane emissions dominated annual greenhouse gas balances 30 years after bog rewetting,
880 *Biogeosciences*, 12, 4361-4371, doi:10.5194/bg-12-4361-2015, 2015.
881

882 Voigtländer, U., Schmidt, J., and Scheller, W.: *Pflege-und Entwicklungsplan NSG Heiligensee*
883 *und Hütelmoor*, 1996.
884

885 Wagner, D.: Effect of varying soil water potentials on methanogenesis in aerated marshland soils,
886 *Scientific Reports*, 7(14706), doi:10.1038/s41598-017-14980-y, 2017.
887

888 Weisner, E. and Schernewski, G.: Adaptation to climate change: a combined coastal protection
889 and re-alignment scheme in a Baltic tourism region, *Journal of Coastal Research*, Special Issue 65,
890 1963-1968, doi:10.2112/SI65-332.1, 2013.
891
892 Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer New York, 2009.
893
894 Wilhelm, E., Batino, R., and Wilcock, R. J.: Low-pressure solubility of gases in liquid water,
895 *Chemical Reviews*, 77(2), 219-262, doi:10.1021/cr60306a003, 1977.
896
897 Wilson, D., Blain, D., Couwenburg, J., Evans, C. D., Murdiyarso, D., Page, S. E., Renou-Wilson,
898 F., Rieley, J. O., Sirin, A., Strack, M., and Tuittila, E.-S.: Greenhouse gas emission factors
899 associated with rewetting of organic soils, *Mires and Peat*, 17(4), 1-28,
900 doi:10.19189/MaP.2016.OMB.222, 2016.
901
902 Yrjälä, K., Tuomivirta, T. T., Juottonen, H., Putkinen, A., Lappi, K., Tuittila, E.-S., Penttilä, T.,
903 Minkkinen, K., Laines, J., Peltoniemi, K., and Fritze, H.: CH₄ production and oxidation processes
904 in a boreal fen ecosystem after long-term water table drawdown, *Global Change Biology*, 17,
905 1311-1320, doi:10.1111/j.1365-2486.2010.02290.x, 2011.
906
907 Zak, D. and Gelbrecht, J.: The mobilisation of phosphorus, organic carbon, and ammonium in the
908 initial stage of fen rewetting (a case study from NE Germany), *Biogeochemistry*, 85, 141-151,
909 doi:10.1007/s10533-007-9122-2, 2007.
910
911 Zak, D., Reuter, H., Augustin, J., Shatwell, T., Barth, M., Gelbrecht, J., and McInnes, R. J.:
912 Changes of the CO₂ and CH₄ production potential of rewetted fens in the perspective of temporal
913 vegetation shifts, *Biogeosciences*, 12, 2455-2468, doi:10.5194/bg-12-2455-2015, 2015.
914
915 Zhang, G., Haiyang, Y., Xianfang, F., Jing, M., and Hua, X.: Carbon isotope fractionation reveals
916 distinct process of CH₄ emission from different compartments of paddy ecosystem, *Scientific*
917 *Reports*, 6(27065), doi:10.1038/srep27065, 2016.
918
919 Zhang, J., Kobert, K., Flouri, T., and Stamatakis, A.: PEAR: a fast and accurate Illumina Paired-
920 End reAd merger, *Bioinformatics*, 30(5), 614-620, doi:10.1093/bioinformatics/btt593, 2014.
921
922 Zhou, X., Zhang, Z., Tian, L., Li, X., and Tian, C.: Microbial communities in peatlands along a
923 chronosequence on the Sanjiang Plain, China, *Nature Scientific Reports*, 7(9567), 1-11,
924 doi:10.1038/s41598-017-10436-5, 2017.
925
926


927



928


929 **Figure 1:** Location of study sites in northeastern Germany (a) and sampling locations within sites (b) Hütelmoor and (c) Zarnekow.
930 Maps b) and c) are drawn to the same scale. Image source: (a) QGIS, (b) and (c) Google Earth via QGIS OpenLayer Plugin. Imagery
931 date: August 9, 2015.

932




933
934
935
936
937
938


Figure 2: Depth profiles of oxygen, nitrate, total iron, manganese, and sulfate (upper panels), and profiles of pH, EC, dissolved methane, and the isotopic signature of methane-bound carbon (lower panels) in both study sites. Solid lines connect the respective means of individual wetlands (n=4 for Hütelmoor and n=5 for Zarnekow).



939
 940 **Figure 3:** Relative abundances of different bacterial lineages in the study sites. Along the horizontal axis samples are arranged
 941 according to site and depth. The rank order along the vertical axis is shown for the phylum level.




942
 943 **Figure 4:** Relative abundances of Proteobacteria phyla in the study sites. Along the horizontal axis samples are arranged according
 944 to site and depth. The rank order along the vertical axis is shown for the family level. If an assignment to the family level was not
 945 possible the next higher assignable taxonomic level was used.
 946



948
949
950
951
952
953

Figure 5: Relative abundances of different archaeal lineages in the study sites. Along the horizontal axis samples are arranged according to site and depth. The rank order along the vertical axis is shown for the family level. If an assignment to the family level was not possible, the next higher assignable taxonomic level was used.


954



955
956
957
958
959
960
961
962
963

Figure 6: NMDS plots showing (a) bacterial, (b) archaeal, and (c) microbial (bacterial plus archaeal) community composition across the nine peat cores. The point positions represent distinct microbial communities, with the border colors of the symbols referring to the study sites and their shapes representing the core number. HC 2 symbols are highlighted with red fill to emphasize the large variation in microbial community within the core. Environmental fit vectors with a significance of $p < 0.05$ are shown in green.

964



965
966
967
968
969
970
971
972

Figure 7: Depth distribution of qPCR abundances for total microbial (16S), methanogen (*mcrA*), methanotroph (*pmoA*), and ratio of *mcrA* to *pmoA* gene copy numbers in both sites. Microbial abundances were designated as numbers of gene copies per gram of dry peat soil. Duplicate measurements per depth section are shown against sampling depth using log-transformed values. Solid lines indicate mean abundances for individual wetlands (n=4 for Hütelmoor and n=5 for Zarnekow). Note that the plot at the right was split into two plots to capture very high *mcrA/pmoA* ratios in the upper peat layer.

973
974
975
976
977
978
979

Table 1: Environmental conditions, geochemical conditions, and microbial abundances in peat cores from the Hütelmoor, a coastal minerotrophic fen in northeastern Germany. Environmental conditions are described by pH and EC (electrical conductivity). Geochemical parameters shown are dissolved methane (CH₄) concentrations, the isotopic signature of methane-bound carbon ($\delta^{13}\text{C}-\text{CH}_4$), and concentrations of terminal electron acceptors which are denoted with their respective chemical abbreviations. Microbial abundances here represent the mean value of subsamples for each depth section (n=2). nd = not detected.

Core, depth	pH	EC	$\delta^{13}\text{C}-\text{CH}_4$	Dissolved CH ₄	O ₂	NO ₃ ⁻	Fe	Mn	SO ₄ ²⁻	16S	<i>mcrA</i>	<i>pmoA</i>	<i>mcrA/pmoA</i>
cm		mS cm ⁻¹				mM					gene copies g dry peat ⁻¹		
HC 1, 0-5	7.2	1.79	-60.2	0.14	0.30	nd	0.10	0.03	0.03	2.04x10 ¹⁰	1.15x10 ⁰⁸	6.60x10 ⁰⁶	17.7
5-10	7.0	1.80	-60.7	0.31	0.18	nd	0.31	0.02	0.01	3.25 x10 ¹⁰	3.36x10 ⁰⁷	6.68x10 ⁰⁷	0.51
10-15	7.0	2.35	-65.1	0.23	0.05	nd	0.60	0.03	nd	2.11x10 ¹⁰	8.12x10 ⁰⁷	1.76x10 ⁰⁷	6.12
15-20	7.1	2.94	-66.1	0.11	nd	0.03	1.34	0.06	nd	3.08x10 ¹⁰	1.21x10 ⁰⁸	2.76x10 ⁰⁷	4.41
HC 2, 0-5	6.9	3.01	-57.8	0.46	0.05	0.03	0.03	0.01	nd	1.10x10 ¹¹	1.13x10 ¹⁰	1.03x10 ⁰⁷	1,170
5-10	6.7	2.60	-63.2	0.34	0.17	2.63	0.10	0.01	0.01	5.51x10 ¹⁰	7.27x10 ⁰⁷	1.69x10 ⁰⁷	4.73
10-20	7.2	5.73	-60.4	0.06	0.29	3.00	1.41	0.02	nd	3.13x10 ¹⁰	4.47x10 ⁰⁶	7.32x10 ⁰⁶	0.74
20-30	7.0	7.29	-61.8	0.08	0.08	nd	1.51	0.02	0.29	4.71x10 ⁰⁹	6.41x10 ⁰⁵	4.50x10 ⁰⁵	3.75
30-40	6.5	9.66	-64.2	0.64	nd	nd	1.68	0.02	3.66	2.09x10 ⁰⁹	6.21x10 ⁰⁵	3.90x10 ⁰⁴	18.3
40-50	6.4	9.71	-64.5	0.20	nd	nd	5.35	0.03	17.1	4.09x10 ⁰⁹	2.47x10 ⁰⁶	2.75x10 ⁰⁵	10.7
HC 3, 0-5	6.6	2.93	-57.7	0.23	0.29	2.77	0.11	0.01	0.04	1.10x10 ¹¹	1.34x10 ⁰⁹	3.51x10 ⁰⁸	3.86
5-10	6.6	3.00	-57.4	0.19	0.27	2.69	0.01	0.01	0.03	8.72x10 ¹⁰	1.40x10 ⁰⁹	3.42x10 ⁰⁷	46.6
10-20	6.4	3.77	-57.3	0.49	0.24	3.08	0.05	nd	nd	6.08x10 ¹⁰	5.86x10 ⁰⁸	9.35x10 ⁰⁶	63.6
20-30	6.1	6.77	-57.4	0.42	0.11	nd	0.20	nd	nd	4.26x10 ¹⁰	3.48x10 ⁰⁸	1.92x10 ⁰⁷	18.2
30-40	6.5	8.56	-59.4	0.08	0.03	nd	0.16	nd	nd	1.05x10 ¹⁰	3.20x10 ⁰⁶	1.17x10 ⁰⁶	2.74
40-50	5.6	9.36	-59.5	0.12	0.01	nd	0.02	nd	0.08	3.18x10 ⁰⁹	2.16x10 ⁰⁶	2.58x10 ⁰⁵	8.39
HC 4, 0-5	6.6	2.93	-61.2	0.25	0.30	2.72	0.02	0.01	0.04	1.17x10 ¹¹	3.63x10 ⁰⁹	3.09x10 ⁰⁸	11.7
5-10	6.7	2.65	-59.2	0.13	0.30	2.87	0.01	nd	0.05	4.87x10 ¹⁰	1.09x10 ⁰⁹	7.51x10 ⁰⁷	14.5
10-20	6.6	5.20	-60.5	0.05	0.30	3.05	0.14	nd	nd	4.85x10 ¹⁰	8.71x10 ⁰⁸	2.15x10 ⁰⁷	40.8
20-30	7.2	6.06	-59.1	0.05	0.01	nd	0.06	nd	0.02	9.78x10 ⁰⁹	5.82x10 ⁰⁷	7.91x10 ⁰⁶	7.36
30-40	6.6	8.11	-60.6	0.29	nd	nd	0.09	nd	0.67	1.60x10 ⁰⁹	1.58x10 ⁰⁶	1.25x10 ⁰⁶	1.27

980
981
982
983
984
985
986
987
988
989
990
991
992
993
994


1001
1002
1003
1004
1005
1006
1007

Table 2: Environmental conditions, geochemical conditions, and microbial abundances in peat cores from Zarnekow, a freshwater minerotrophic fen in northeastern Germany. Environmental conditions are described by pH and EC (electrical conductivity). Geochemical parameters shown are dissolved methane (CH₄) concentrations, the isotopic signature of methane-bound carbon ($\delta^{13}\text{C}-\text{CH}_4$), and concentrations of terminal electron acceptors which are denoted with their respective chemical abbreviations. Microbial abundances here represent the mean value of subsamples for each depth section (n=2). nd = not detected.

Core, depth	pH	EC	$\delta^{13}\text{C}-\text{CH}_4$	Dissolved CH ₄	O ₂	NO ₃ ⁻	Fe	Mn	SO ₄ ²⁻	16S	<i>mcrA</i>	<i>pmoA</i>	<i>mcrA/pmoA</i>
cm		mS cm ⁻¹				mM					gene copies g dry peat ⁻¹		
ZC 1, 0-5	6.64	1.03	-64.5	0.51	0.07	0.001	0.007	0.002	0.002	6.33x10 ¹⁰	1.02x10 ⁰⁹	1.49x10 ⁰⁷	69.7
25-30	6.67	1.14	-62.0	0.64	0.08	0.001	0.087	0.028	0.003	4.25x10 ¹⁰	8.96x10 ⁰⁸	9.14x10 ⁰⁶	98.0
50-55	6.66	1.31	-62.5	0.63	0.09	0.005	0.310	0.037	0.002	3.40x10 ¹⁰	3.97x10 ⁰⁸	6.85x10 ⁰⁶	58.1
ZC 2, 0-5	6.91	1.00	-59.2	0.17	0.08	0.004	0.012	0.069	0.007	1.43x10 ¹¹	1.14x10 ¹⁰	4.35x10 ⁰⁷	261
25-30	6.76	1.29	-51.3	0.15	0.10	0.001	0.215	0.033	0.013	6.44x10 ¹⁰	1.45x10 ⁰⁹	2.34x10 ⁰⁷	61.8
50-55	6.64	1.52	-61.1	0.62	0.04	nd	0.410	0.054	0.003	5.64x10 ¹⁰	5.10x10 ⁰⁸	1.50x10 ⁰⁷	34.0
ZC 3, 0-5	6.88	1.17	-60.5	0.50	0.10	0.001	0.073	0.074	0.032	7.86x10 ¹⁰	2.78x10 ⁰⁹	3.26x10 ⁰⁷	85.7
25-30	7.04	3.39	-61.9	0.10	0.03	0.002	1.046	0.188	0.003	5.79x10 ¹⁰	7.81x10 ⁰⁸	1.55x10 ⁰⁷	51.8
50-55	6.92	3.82	-68.7	0.59	0.02	nd	0.779	0.123	0.003	3.41x10 ¹⁰	2.21x10 ⁰⁸	5.41x10 ⁰⁶	40.9
ZC 4, 0-5	7.3	1.06	-61.5	0.14	0.12	0.010	0.013	0.024	0.035	7.19x10 ¹⁰	1.28x10 ⁰⁹	6.53x10 ⁰⁷	19.6
25-30	7.13	1.58	-65.1	0.12	0.11	0.002	0.301	0.049	0.002	7.19x10 ¹⁰	nd	4.60x10 ⁰⁷	-
50-55	6.89	1.51	-67.6	0.17	0.11	0.002	0.366	0.048	0.002	5.42x10 ¹⁰	9.47x10 ⁰⁸	4.50x10 ⁰⁷	21.0
ZC 5, 0-5	6.81	0.83	-63.7	0.57	0.01	0.002	0.005	0.035	0.005	8.73x10 ¹⁰	8.73x10 ⁰⁸	4.97x10 ⁰⁷	17.6
25-30	6.72	0.86	-63.5	0.53	0.06	0.002	0.139	0.043	0.001	8.94x10 ¹⁰	5.21x10 ⁰⁸	5.57x10 ⁰⁷	93.4
50-55	6.58	1.00	-63.8	0.37	0.06	0.002	0.275	0.045	0.002	8.00x10 ¹⁰	2.14x10 ⁰⁸	1.44x10 ⁰⁸	14.9

1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032

1040 **Supplemental Material**



1041
1042

1043 Figure S1: Incubation data from Zarnekow, a freshwater minerotrophic fen in Northeastern Germany. Rates of methane production
1044 (n=3) and methane oxidation (n=3) are shown for both fresh (surficial) organic sediment and the bulk peat.