

# 1 Ecophysiological characteristics of red, green and brown strains of 2 the Baltic picocyanobacterium *Synechococcus* sp. – a laboratory 3 study

4  
5 S. Śliwińska-Wilczewska<sup>1</sup>, A. Cieszyńska<sup>2</sup>, and A. Latała<sup>1</sup>

6  
7 <sup>1</sup>University of Gdańsk, Institute of Oceanography, Laboratory of Marine Plant Ecophysiology, Gdynia, Poland

8 <sup>2</sup>Institute of Oceanology Polish Academy of Sciences, Department of Marine Physics, Marine Biophysics Laboratory, Sopot,  
9 Poland

10 Correspondence to: A. Cieszyńska (acieszynska@iopan.gda.pl, cieszynska.agata@gmail.com)

11  
12 **Abstract.** The contribution of picocyanobacteria (PCY) to summer phytoplankton blooms, accompanied by an ecological  
13 crisis, is a new phenomenon in Europe. This issue requires careful investigation. Therefore, the work, which examines the  
14 response of *Synechococcus* sp. physiology to different environmental conditions was conducted. Three strains of  
15 *Synechococcus* sp. (red BA-120, green BA-124 and brown BA-132) were cultivated in a laboratory under previously  
16 determined environmental conditions. These conditions were as follows: temperature (T) from 10 by 5 to 25°C, salinity from  
17 3 by 5 to 18 PSU and Photosynthetically Active Radiation (PAR) from 10 by 90 to 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ , which gave 64  
18 combinations of synthetic, though real environmental conditions. Scenarios reflecting all possible combinations were applied  
19 in the laboratory experiments. Results pointed to differences in final number of cells between strains. However, there was  
20 also a similar pattern for BA-124 and BA-132, which showed the highest concentrations of picocyanobacteria cells at higher  
21 T and PAR. This was also the case for BA-120, but only to a certain degree as the number of cells started to decrease above  
22 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  PAR. Pigmentation, chlorophyll *a* (Chl *a*), fluorescence and rate of photosynthesis presented both  
23 similarities and differences between the strains. In this context, more consistent features were observed for brown and red  
24 strains when compared to the green. In this paper are defined the ecophysiological responses of PCY.

## 25 26 1 Introduction

27  
28 The presence of picoplankton and its contribution to marine biomass were ignored in environmental studies conducted  
29 before 1970. This was related to the poor accuracy of research equipment, which did not enable the recording and  
30 recognition of such small organisms. Before the discovery of picocyanobacteria (PCY) in the oceans by Johnson and  
31 Sieburth (1979) and Waterbury et al. (1979) there only existed incidental reports of this fraction of cyanobacteria occurrence  
32 in aquatic ecosystems. Since then, the number of PCY observations has rapidly increased, and currently they are known to be  
33 present in many marine, brackish and freshwater ecosystems of the world (e.g., Callieri, 2010; Sorokin and Zakuskina, 2010;  
34 Flombaum et al., 2013; Jodłowska and Śliwińska, 2014; Jasser and Callieri, 2017). Additionally, recent works showed that  
35 many aquatic ecosystems have been experiencing super-dense, long-term blooms of picocyanobacteria (Sorokin et al., 2004;  
36 Sorokin and Zakuskina, 2010), whilst in the past PCY were often described as a non-blooming group (Stockner et al., 1988).  
37 Sorokin and Zakuskina (2010) found that the picocyanobacteria blooms were accompanied by great changes in the benthic  
38 habitats.

39 Picocyanobacteria of the *Synechococcus* genus are extremely important organisms in the world's oceans. This is the  
40 smallest fraction of plankton ranked by the size of cells, which ranges from 0.2 to 2.0  $\mu\text{m}$  (Sieburth et al., 1978).  
41 Chroococcoid genus of the *Synechococcus* are ubiquitous components of the natural plankton communities in aquatic  
42 environments. Picocyanobacteria of the *Synechococcus* group span a range of different colors, depending on their pigment

43 composition (Stomp et al., 2007; Haverkamp et al., 2008). Baltic strains of *Synechococcus* sp. are classified as red strains  
44 with phycoerythrin (PE), green strains rich in phycocyanin (PC) and the brown strains containing two different bilin  
45 pigments known as phycoerythrobilin (PEB) and phycourobilin (PUB), which both bind to the apoprotein PE (Six et al.,  
46 2007a, b; Haverkamp et al., 2008; 2009). The three strains of *Synechococcus* sp.: BA-120 (red), BA-124 (green), and BA-  
47 132 (brown) examined in this work (Fig. S1 in Supplement) are different morphotypes representatives. The existence of  
48 these different colors picocyanobacteria is commonly found in the Baltic Sea (Andersson et al., 1996; Hajdu et al., 2007;  
49 Stomp et al., 2007; Haverkamp et al., 2009; Mazur-Marzec et al., 2013; Larsson et al., 2014; Paczkowska et al., 2017).

50 Picocyanobacterial species are phylogenetically divided into several major clusters. These clusters have been  
51 identified, based on photosynthetic pigmentation, nitrogen requirements, motility and salinity preferences (Herdman et al.,  
52 2001). Picocyanobacteria that are often found and isolated from marine, brackish and freshwater environments are related to  
53 *Synechococcus* cluster 5 (Herdman et al., 2001). *Synechococcus* cluster 5 is divided in two sub-clusters: 5.1 and 5.2. The  
54 members of cluster 5.1 typically produce PE as their main photosynthetic pigment. In contrast, members of cluster 5.2 have a  
55 green coloration because they produce PC (Herdman et al., 2001; Larsson et al., 2014). The diversity of picocyanobacteria  
56 has been investigated mainly by analysis of the 16S rRNA gene. However, the phylogenetic tree of *Synechococcus* sp. is not  
57 always consistent with their pigmentation type (Haverkamp et al., 2008). Thus, the actual taxonomic position may be  
58 incorrectly defined due to the morphological plasticity of these organisms (Callieri, 2010).

59 Despite its association with open ocean systems, it has become increasingly evident in recent years that *Synechococcus*  
60 sp. is a significant contributor to cyanobacterial blooms (Beardall, 2008). Surprisingly, this species may also comprise 80%  
61 and more of the total cyanobacterial biomass during cyanobacterial blooms in the Baltic Sea (Stal et al., 2003; Mazur-  
62 Marzec et al., 2013).

63 Recently, it has been confirmed that PCY are able to excrete harmful and allelopathic substances (e.g., Jakubowska and  
64 Szelaż-Wasilewska, 2015; Jasser and Callieri, 2017; Śliwińska-Wilczewska et al., 2017; Barreiro Felpeto et al., 2018). Many  
65 different factors, including physical parameters, availability and competition for resources, selective grazing and allelopathic  
66 interactions can affect the occurrence of harmful blooms in aquatic ecosystems. The development of massive algal blooming  
67 is a consequence of the interaction between many favorable factors. *Synechococcus* sp. greatly contributes to these massive  
68 blooms, but so far the characteristics of the life cycle of Baltic PCY has not been sufficiently studied. This knowledge needs  
69 to be expanded and improved, especially because of bloom toxicity and negative impacts on ecosystems (Jasser and Callieri,  
70 2017; Śliwińska-Wilczewska et al., 2018a).

71 According to the above all, phytoplankton is of great interest to scientists in understanding its life cycles and impact on  
72 the ecosystem in different parts of the world's oceans and within diverse environmental conditions. In order to investigate it,  
73 scientists use various types of research methodology: in-situ measurements, laboratory experiments and numerical  
74 estimations. All of these approaches are necessary and essential in marine phytoplankton examination. Some laboratory and  
75 field studies of ecophysiological responses of picocyanobacteria to different growth conditions have already been completed  
76 for typical oceanic mediums, semi-closed seas and lakes (e.g., Glover et al., 1986; Kuosa, 1988; Stal et al., 1999; Agawin et  
77 al., 2000; Callieri and Stockner, 2002; Hajdu et al., 2007; Sánchez-Baracaldo et al., 2008; Cai and Kong, 2013; Motwani et  
78 al., 2013; Jodłowska and Śliwińska, 2014; Stawiarski et al., 2016). However, there is still a need to provide more systematic  
79 information about these organisms. **What is more, the need is amplified by the fact that there are only few research papers on**  
80 **the brown strain of Baltic *Synechococcus* sp.** (Stal et al., 2003; Haverkamp et al., 2008; 2009; Jodłowska and Śliwińska,  
81 2014). **This gives limited knowledge of PCY and their life cycle in the Baltic Sea, as brown form also contributes to total**  
82 **pico- and phytoplankton biomass in the area of interest (Stal et al., 2003). The above strengthens the motivation to conduct**  
83 **studies on the brown strain of *Synechococcus* sp.**

84 The overall goal of this paper is to determine the most favorable and unfavorable environmental conditions for PCY to  
85 grow on the basis of three different strains of *Synechococcus* sp. ecophysiological analysis. The initial step of these works  
86 was to carry out laboratory experiments with *Synechococcus* sp. cultures. In order to create different environmental

87 conditions in the Baltic Sea range, combinations of physical quantities were determined. These quantities were as follows:  
88 scalar irradiance in Photosynthetically Active Radiation (PAR) spectrum range (10, 100, 190, 280  $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ),  
89 salinity (3, 8, 13, 18 PSU), and temperature (T) (10, 15, 20, 25°C). In total, 64 combinations (environmental scenarios) were  
90 generated. The second step was to plot and analyze all results after seven days of incubations. For the results, the number of  
91 cells, pigmentation, Chl *a* fluorescence parameters, and rate of photosynthesis were collected. The third step was to extract  
92 any significant relations between the results and specific physical factors by using a statistical analysis, which included the  
93 variance method analysis (two-way ANOVA) and Tukey's HSD post-hoc test. Derived laboratory results will help to  
94 develop the knowledge of the picocyanobacteria life cycle. Moreover, the PCY experiments underlie the improved numerical  
95 approach to phytoplankton modeling development. On the basis of derived results, the algorithm for picocyanobacterium  
96 growth will be created in a separate study.

## 98 2 Material and methods

### 100 2.1 Material and culture conditions

102 Three different phenotypes of picocyanobacteria strains from the genus *Synechococcus* were examined: BA-120 (red), BA-  
103 124 (green), and BA-132 (brown). The cultures preparation was carried out as follows. The *Synechococcus* sp. strains were  
104 isolated from the coastal zone of the Gulf of Gdansk (southern Baltic Sea) and maintained as unialgal cultures in the Culture  
105 Collection of Baltic Algae (CCBA) at the Institute of Oceanography, University of Gdańsk, Poland (Latała et al., 2006). The  
106 experiments on the 'batch cultures' were carried out in 25 mL glass Erlenmeyer flasks containing sterilized f/2 medium  
107 (Guillard, 1975). In order to develop the media, the appropriate amount of Tropic Marine Synthetic Sea Salt was dissolved in  
108 distilled water. The final salinity was 3, 8, 13 and 18 PSU, measured with salinometer (inoLab Cond Level 1, Weilheim in  
109 Oberbayern, Germany). Salinity of the media was measured in PSU (practical salinity units). The major nutrients,  
110 microelements and vitamin concentrations were added according to a method proposed by Guillard (1975) (any of the  
111 components in f/2 media were not replaced by Tropic Marine Synthetic Sea Salt). Culture media was prepared with artificial  
112 seawater filtered through a 0.45- $\mu\text{m}$  filters (Macherey-Nagel MN GF-5) using a vacuum pump (600 mbar) and autoclaved.  
113 Into 25 mL Erlenmeyer glass flasks, the cells of specific strains were inoculated. The picocyanobacteria cultures were  
114 acclimated to the various synthetic environmental conditions for two days. The conditions were the combinations of different  
115 values of: scalar irradiance in Photosynthetically Active Radiation (PAR) spectrum (10, 100, 190 and 280  $\mu\text{mol photons m}^{-2}$   
116  $\text{s}^{-1}$ ), temperature (T) (10, 15, 20 and 25°C), and salinity (3, 8, 13 and 18 PSU). Values of quantities representing each  
117 environmental condition were applied at the fixed intervals, i.e.: PAR, interval 90; T, interval 5; salinity, interval 5. The  
118 synthetic environmental conditions of salinity and T applied in the laboratory are representative for the Baltic Sea area  
119 (Feistelet al., 2008; 2009; Siegel and Gerth, 2017). Regarding PAR, its levels has been generated the highest possible to be  
120 achieved in the laboratory. These values are generally lower than mean PAR intensities being observed in the summertime in  
121 the Baltic (Leppäranta and Myrberg, 2009). Moreover, the values of environmental conditions variables (salinity,  
122 temperature, PAR) were also specified in certain ranges to make this study comparable with other laboratory cultures  
123 experiments available in literature. The combination of the quantities of environmental variables is called a scenario in the  
124 present paper. The intensity of PAR was measured using a LI-COR spherical quantum-meter. Fluorescent lamps (Cool  
125 White 40W, Sylvania, USA) were used as source of irradiance and combined with halogen lamps (100W, Sylvania, USA) to  
126 obtain more intensive light. After acclimation time (2 d), the picocyanobacteria cells served as inoculum for the right test  
127 cultures with the initial number of cells equal to  $10^6 \text{ cells mL}^{-1}$ . The flasks with picocyanobacteria were shaken (once a day)  
128 during the experiment. In order to achieve the most reliable results, test cultures were grown in three replicas and were  
129 incubated for one week at each combination of light, temperature and salinity. On the last day of incubation the number of

130 cells, pigment content, Chl *a* fluorescence, and rate of photosynthesis were measured in each replica. Results were reported  
131 as mean values  $\pm$  standard deviation (SD).

## 133 2.2 Determination of the number of cells

134  
135 The number of cells (N) in cultures was counted with flow cytometer BD Accuri™ C6 Plus (BD Biosciences, San Jose, CA,  
136 USA) according to the procedure proposed by Śliwińska-Wilczewska et al. (2018b). Events were recorded in list form.  
137 Samples were run at a flow rate of approximately 14  $\mu\text{L min}^{-1}$ . Selection of this flow rate was based on previous  
138 introductory experiments to determine the most relevant effectiveness. Choosing an adequate discriminator and thresholds  
139 plays a key role in recording the cells correctly. The most reasonable solution to record chlorophyll fluorescing  
140 cyanobacteria and microalgae is to choose the red fluorescence as the discriminator (Fig. S1) and to select a high threshold,  
141 enough to eliminate optical and electronic noise (Marie et al., 2005). Concerning this, the discriminator was set on the red  
142 (chlorophyll) fluorescence with a standard threshold of 80,000 on FSC-H. Flow was daily calibrated with Spherotech 6- and  
143 8- Peak Validation Beads (BD, San Jose, USA). This ensures that the cytometer is working properly before running  
144 experimental samples. FITC, PE, and PE-Cy5 detectors were daily calibrated with SPHERO™ Rainbow Calibration  
145 Particles (BD, San Jose, USA), and the APC channel was calibrated with SPHERO 6-peaks Allophycocyanin Calibration  
146 Particles (APC). Detectors FL1, FL2, and FL3 read fluorescence emissions excited by the blue laser (480 nm), while detector  
147 FL4 reads emissions excited by the red laser (640 nm). The flow cytometry was used to establish the initial number of  
148 picocyanobacteria cells and to measure the final cells concentration after the incubation period.

## 150 2.3 Determination of the pigments content

151  
152 The concentration of photosynthetic pigments of analyzed picocyanobacteria was measured by the spectrophotometric  
153 method (Strickland and Parsons, 1972). The analysis of mL-specific (pigment content per mL) and cell-specific (pigment  
154 content per cell) pigmentation was conducted. After seven days of incubation, 4 mL of culture was filtered in order to  
155 separate the picocyanobacteria cells from the medium. Chl *a* and carotenoids (Car) were extracted from the  
156 picocyanobacteria cells with cold 90% acetone (5 mL). To improve extraction, the cells were disintegrated for two minutes  
157 by ultrasonication. Then, the test-tube with the extract was held in the dark for three hours at  $-60^{\circ}\text{C}$ . To remove cell debris  
158 and filter out the particles, the extracts were centrifuged at 10,000 rpm ( $8496 \times g$ ) for 5 min (Sigma 2-16P, Osterode am  
159 Harz, Germany). The absorbance of pigments was estimated on the basis of Beckman spectrophotometer UV-VIS DU 530  
160 measurements at specific wavelengths (750, 665 and 480 nm), using 1 cm quartz cuvette. Pigment concentration was  
161 calculated according to Strickland and Parsons (1972). The following formulas have been used: Chl *a* ( $\mu\text{g mL}^{-1}$ ) =  
162  $11.236(A_{665}-A_{750})V_a/V_b$ , Car ( $\mu\text{g mL}^{-1}$ ) =  $4(A_{480}-A_{750})V_a/V_b$ , where:  $V_a$  - extract volume (in this study 5 mL),  $V_b$  - sample  
163 volume (in this study 4 mL), and  $A_x$  - absorbance estimated at wavelength  $x$  in a 1-cm cuvette.

## 165 2.4 Chlorophyll fluorescence analyses

166  
167 Chl *a* fluorescence was measured with a Pulse Amplitude Modulation (PAM) fluorometer (FMS1, Hansatech, King's Lynn,  
168 Norfolk, UK). The FMS1 uses a 594 nm amber modulating beam with 4-step frequency control as a measuring light and is  
169 equipped with a dual-purpose halogen light source providing actinic light ( $0 - 3000 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  in 50 steps) and a  
170 saturating pulse ( $0 - 20000 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  in 100 steps). FMS1 also has a 735 nm far-red LED source for preferential  
171 PSI excitation allowing accurate determination of the  $F_o'$  parameter. Samples were filtered through 13-mm glass fiber filters  
172 (Whatman GF/C, pore size = 1.2  $\mu\text{m}$ ). Before measurement, the filtered sample was kept in the dark for 10 min. The  
173 maximum photochemical efficiency of photosystem II (PSII) at dark-adapted state ( $F_v/F_m$ ) and the photochemical efficiency

174 of PSII under actinic light intensity ( $\Phi$ PSII) were estimated. The actinic light was different for different cultures, the same as  
175 the PAR level was for each incubation. The above is similar to the method used by Campbell et al. (1998).

## 177 2.5 Measurements of photosynthesis rate

178  
179 The measurements of oxygen evolution were carried out on the day seventh of the experiment using a Clark-type oxygen  
180 electrode (Chlorolab 2, Hansatech). Temperature was controlled with a cooling system LAUDA (E100, Germany).  
181 Illumination was provided by a high intensity probe-type light array with 11 red LED's centered on 650 nm. Irradiance was  
182 measured with a quantum sensor (Quantitherm, Hansatech, King's Lynn, Norfolk, UK). Dark respiration was estimated from  
183 O<sub>2</sub> uptake by cells incubated in the dark. Experimental data was fitted to the photosynthesis irradiance response ( $P-E$ )  
184 curves using equation (Jassby and Platt, 1976) and Statistica® 13.1 software and photosynthetic parameters, i.e., the  
185 photosynthetic capacity ( $P_m$ ), the initial slope of  $P-E$  curve ( $\alpha$ ) and the dark respiration ( $R_d$ ) (Sakshaug et al., 1997).

## 187 2.6 Statistical analyses

188  
189 The effect of light and temperature separately and then their combinations impact on growth, pigments content, fluorescence  
190 and photosynthesis performance of examined strains were analyzed using two-way variance analysis (ANOVA). A post hoc  
191 test (Tukey's HSD) was used to show which results differed under varied conditions over the experimental period (Sheskin  
192 2000). The confident levels in the statistical analysis were: 95% ( $*p < 0.05$ ), 99% ( $**p < 0.01$ ), 99.9% ( $***p < 0.001$ ). The  
193 statistical analyses were performed using Statistica® 13.1 and Matlab 2012b software. According to the literature, light and  
194 temperature are major factors controlling the growth and distribution of picocyanobacteria (e.g.: Jasser and Arvola, 2003),  
195 and they may have considerable significance on the abundance of the *Synechococcus* community (Glover, 1985; Glover et  
196 al., 1985; 1986, Joint and Pomroy, 1986; Jasser and Arvola, 2003; Jasser, 2006; Jodłowska and Śliwińska, 2014), as a result  
197 it was decided that light and temperature would be the independent variables in ANOVA and post-hoc test analysis. The  
198 dependent variable was always the parameter, which had been measured.

## 200 3 Results

### 202 3.1 Number of cells

203  
204 For all three picoplankton strains, ANOVA analysis indicated that in each scenario the independent variable (temperature or  
205 PAR) significantly influenced the dependent variable. What is more, post-hoc tests indicated that multiple factors (T and  
206 PAR together) had an impact on the PCY growth.

207 According to post-hoc tests, 2008 multiple comparisons (70%) out of all 2880 completed for three strains, indicated the  
208 highest statistical significance (Tukey HSD,  $*** p < 0.001$ ), 160 multiple comparisons (6%) pointed to the statistical  
209 significance of  $0.001 < ** p < 0.01$ , and 114 (4%) stated for the significance of  $0.01 < * p < 0.05$ . The rest of the multiple  
210 comparisons (598, 20%) indicated no statistically significance differences (Tukey HSD,  $p \geq 0.05$ ).

211 Both PAR and T affected the number of *Synechococcus* sp. BA-120 cells significantly (ANOVA,  $F_{9,32} = 42.3$ ,  $*** p <$   
212  $0.001$ , ANOVA,  $F_{9,32} = 22.7$ ,  $*** p < 0.001$ , ANOVA,  $F_{9,32} = 9.6$ ,  $*** p < 0.001$  and ANOVA,  $F_{9,32} = 12.2$ ,  $*** p < 0.001$ ,  
213 for salinity 3, 8, 13, 18 PSU, respectively). For BA-120, the number of cells increased with T in each medium (salinities 3, 8,  
214 13, 18 PSU) (Fig. 1A, a-d). The minimum number of cells was estimated in salinity 3 PSU, T 10°C and PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$   
215 ( $1.6 \times 10^6 \text{ cell mL}^{-1}$ , Fig. 1A, a), whilst the maximum in salinity 18 PSU, T 25°C, PAR 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$   
216 ( $11.5 \times 10^6 \text{ cell mL}^{-1}$ , Fig. 1A, d). The decrease in number of cells was observed from PAR 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$   
217 onwards. This can likely be related to the photo-inhibition of photosystem II (PSII). The above was the case in each salinity

218 (Figs. 1A, a-d). It seemed that the most important environmental factor for BA-120 number of cells was T, with PAR playing  
219 an additional role, for instance in the context of photo-inhibition. Multiple comparisons tests pointed to the strong  
220 significance of PAR and T combined in influencing the number of *Synechococcus* sp. BA-120 cells. According to the  
221 statistics, 82% of multiple comparisons were statistically significant (Tukey HSD, \*  $p < 0.05$ ) with 91% of them having the  
222 highest significance level (Tukey HSD, \*\*\*  $p < 0.001$ ).

223 Both PAR and T also significantly affected the number of *Synechococcus* sp. BA-124 cells (ANOVA,  $F_{9,32} = 7.9$ , \*\*\*  $p$   
224  $< 0.001$ , ANOVA,  $F_{9,32} = 13.6$ , \*\*\*  $p < 0.001$ , ANOVA,  $F_{9,32} = 8.4$ , \*\*\*  $p < 0.001$  and ANOVA,  $F_{9,32} = 2.8$ , \*\*  $p < 0.01$ , for  
225 salinity 3, 8, 13, 18 PSU, respectively). For BA-124, number of cells increased with T and PAR in all salinities (Figs. 1B, a-  
226 d). The lowest number of cells was calculated in salinity 3 PSU, T 10°C and PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  ( $2.0 \times 10^6 \text{ cell mL}^{-1}$ ,  
227 Fig. 1B, a) and the highest number of cells was reached in salinity 18 PSU, T 25°C, PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$   
228 ( $43.6 \times 10^6 \text{ cell mL}^{-1}$ , Fig. 1B, d). Intensive cell division was also estimated under the highest T and PAR conditions in  
229 salinity 13 PSU with a number of cells of  $41.1 \times 10^6 \text{ cell mL}^{-1}$  (Fig. 1B, c). Generally, the number of cells was the highest  
230 when compared to BA-120 and BA-132 in respective scenarios. One of the observations was the difference in BA-124  
231 number of cells between lower and higher PAR and T scenarios. BA-124 seemed to be more sensitive to changes in PAR  
232 and T in their lower rather than in higher ranges. The highest number of *Synechococcus* sp. BA-124 cells were noted in  
233 moderate- and high-salinity mediums. However, optimum salinities for strain BA-124 were 8 and 13 PSU. Due to post-hoc  
234 analysis, salinity 13 PSU differentiated the conditions for cell abundances under different PAR and T at a lower degree when  
235 compared to other salinities under respective PAR and T (Fig. 1B, c). Another feature of BA-124 was the number of cells in  
236 low T and high PAR scenarios were nearly equal to cell abundances in high T and low PAR scenarios. This was not the case  
237 for BA-120 and BA-132 strains. The observation was supported by Tukey's tests, where only few statistically significant  
238 differences in number of cells were observed between scenarios with elevated PAR (280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ), low T (10,  
239 15°C) and those with high T (25°C) and low PAR (10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ). These differences were observed between 15°C  
240 and 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and 25°C and 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  in salinities 3 and 8 PSU (Tukey HSD, \*\*  $p < 0.05$  in  
241 both cases, Figs. 1B, a-b). Multiple comparisons tests showed high significance of combinations of PAR and T in affecting  
242 the number of cells. According to Tukey HSD tests, 72% of multiple comparisons were statistically significant (\*  $p < 0.05$ )  
243 with 82% of them with the highest significance level (\*\*\*  $p < 0.001$ ).

244 Similarly to BA-120 and BA-124, it was found that PAR and T significantly affected the number of *Synechococcus* sp.  
245 BA-132 cells (ANOVA,  $F_{9,32} = 6.8$ , \*\*\*  $p < 0.001$ , ANOVA,  $F_{9,32} = 5.4$ , \*\*\*  $p < 0.001$ , ANOVA,  $F_{9,32} = 5.6$ , \*\*\*  $p < 0.001$   
246 and ANOVA,  $F_{9,32} = 12.5$ , \*\*  $p < 0.01$ , for salinity 3, 8, 13, 18 PSU, respectively). For BA-132, the positive impact of T and  
247 PAR on number of cells (Figs. 1C, a-d) was observed in each medium. Note that positive impact means the increasing  
248 (positive) dependency, whilst negative impact means decreasing (negative) dependency between the independent and  
249 dependent variable, e.g.: between T and abundance. Salinity played a more significant role here than when compared to BA-  
250 124. It was found that the higher the salinity, the higher the number of cells of BA-132. What is more, according to the  
251 statistical analysis, salinity 18 PSU differentiated the number of cells the most (Fig. 1C, d). In salinity 18 PSU, the cell  
252 abundances could be described as a linear increasing function of ambient T and PAR. This was also observed in other  
253 salinities but not as intensively pronounced as in the highest-saline medium. Moreover, in high salinity, the sensitivity of  
254 number of cells to T changes was much lower than in low salinities. PAR did not determine the number of cells as strongly  
255 as T, which was quite consistent to the observation noted for BA-120. The minimum number of cells was observed in 3 PSU,  
256 10°C and 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  ( $1.4 \times 10^6 \text{ cell mL}^{-1}$ , Fig. 1C, a), whilst the maximum in 18 PSU, 25°C, 280  $\mu\text{mol photons}$   
257  $\text{m}^{-2} \text{s}^{-1}$  ( $16.1 \times 10^6 \text{ cell mL}^{-1}$ , Fig. 1C, d). In addition, the lowest values of BA-132 number of cells were calculated for the  
258 lowest T and PAR condition in each salinity. Tukey HSD post hoc tests indicated high significance of the combination of  
259 PAR and T in affecting the cell abundances. According those tests, 84% of multiple comparisons were statistically  
260 significant (\*  $p < 0.05$ ) with 90% of them with the highest significance (\*\*\*  $p < 0.001$ ).

261 Concerning all three strains, high salinity generally had a positive impact on number of *Synechococcus* sp. cells. What is  
262 more, the relations between salinity and number of cells for all strains, especially red and brown were positive with almost  
263 linearly increasing dependency. The highest average increase was noted for BA-132.

### 264 3.2 Pigment content

265 The results showed that for all strains, cell-specific pigment composition (pigment content per cell) was environmentally  
266 driven (Figs. 2, 3). The analysis of mL-specific pigmentation (pigment content per mL) was also done (Figs. S2 and S3 in  
267 Supplement), however, the mL-specific pigment content is another way to illustrate the biomass and that is why it is not  
268 described in this section in detail.

269 It was estimated, that PAR and T significantly affected the Chl *a* cell-specific content of *Synechococcus* sp. BA-120  
270 (ANOVA,  $F_{9,32} = 33.7$ , \*\*\*  $p < 0.001$ , ANOVA,  $F_{9,32} = 5.3$ , \*\*\*  $p < 0.001$ , ANOVA,  $F_{9,32} = 15.6$ , \*\*\*  $p < 0.001$  and  
271 ANOVA,  $F_{9,32} = 5.7$ , \*\*\*  $p < 0.001$ , for salinity 3, 8, 13, 18 PSU, respectively). Both PAR and T also affected the Car  
272 content in the BA-120 strain cells significantly (ANOVA,  $F_{9,32} = 25.8$ , \*\*\*  $p < 0.001$ , ANOVA,  $F_{9,32} = 7.5$ , \*\*\*  $p < 0.001$ ,  
273 ANOVA,  $F_{9,32} = 7.3$ , \*\*\*  $p < 0.001$ , and ANOVA,  $F_{9,32} = 12.0$ , \*\*\*  $p < 0.001$ , for salinity 3, 8, 13, 18 PSU, respectively). It  
274 was found that cell-specific Chl *a* and Car concentrations decreased with the increase of salinity (Figs. 2A, 3A). On average,  
275 the cell content of pigments for BA-120 was the highest when compared to the other strains. Chl *a* concentration dominated  
276 over Car concentration in each scenario. What is more, there were very high cell-specific concentrations of Chl *a* observed  
277 for the whole T domain at low PAR. Maximum Chl *a* content was measured under T 25°C and PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$   
278 <sup>1</sup>. This was the case in each salinity. The highest Chl *a* concentration within all scenarios was reached in BA-120 cells in  
279 salinity 3 PSU and was equal to 0.339  $\text{pg cell}^{-1}$  (Fig. 2A, a). For other salinities these maximums were as follows: 0.233  $\text{pg}$   
280  $\text{cell}^{-1}$  (8 PSU, Fig. 2A, b), 0.164  $\text{pg cell}^{-1}$  (13 PSU, Fig. 2A, c), 0.100  $\text{pg cell}^{-1}$  (18 PSU, Fig. 2A, d). The highest Car content  
281 was measured in salinity 3 PSU under T of 20°C and PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and reached 0.160  $\text{pg cell}^{-1}$  (Fig. 3A, a).  
282 The lowest concentrations of Chl *a* (0.038  $\text{pg cell}^{-1}$ ) and Car (0.031  $\text{pg cell}^{-1}$ ) were measured in salinity 18 PSU, T 25°C,  
283 PAR 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. 2A, d) and salinity 18 PSU, T 15°C, PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. 3A, d),  
284 respectively. Multiple comparisons tests indicated the significance of PAR and T combined in shaping the pigmentation. Due  
285 to those tests, 52% and 55% of multiple comparisons in Chl *a* and Car content analysis, respectively, were statistically  
286 significant (Tukey HSD, \*  $p < 0.05$ ) with 80% (for Chl *a*) and 74% (for Car) of them with the highest significance (Tukey  
287 HSD, \*\*\*  $p < 0.001$ ).

288 Both PAR and T affected the Chl *a* cell-specific content (ANOVA,  $F_{9,32} = 3.3$ , \*\*  $p < 0.01$ , ANOVA,  $F_{9,32} = 8.3$ , \*\*\*  $p <$   
289 0.001, ANOVA,  $F_{9,32} = 69.8$ , \*\*\*  $p < 0.001$  and ANOVA,  $F_{9,32} = 17.5$ , \*\*\*  $p < 0.001$ , for salinity 3, 8, 13, 18 PSU,  
290 respectively) and Car cell-specific content (ANOVA,  $F_{9,32} = 4.6$ , \*\*\*  $p < 0.001$ , ANOVA,  $F_{9,32} = 65.5$ , \*\*\*  $p < 0.001$ ,  
291 ANOVA,  $F_{9,32} = 83.1$ , \*\*\*  $p < 0.001$  and ANOVA,  $F_{9,32} = 43.2$ , \*\*\*  $p < 0.001$ , for salinity 3, 8, 13, 18 PSU, respectively) of  
292 *Synechococcus* sp. BA-124 significantly. Generally, PAR and high T increase had a negative impact on pigmentation (Figs.  
293 2B, 3B). Maximum values of cell-specific Chl *a* and Car concentrations were measured under 10°C and 10  $\mu\text{mol photons m}^{-2}$   
294  $\text{s}^{-1}$  in each salinity medium. These values, concerning salinities from the lowest to the highest, were as follows: 0.095,  
295 0.102, 0.176, 0.148  $\text{pg cell}^{-1}$  for Chl *a* (Figs. 2B, a-d) and 0.051, 0.067, 0.087, 0.079  $\text{pg cell}^{-1}$  for Car (Figs. 3B, a-d).  
296 Nonetheless, there were also some exceptions. In salinity 3 PSU, high Car contents were calculated under 280  $\mu\text{mol photons}$   
297  $\text{m}^{-2} \text{s}^{-1}$  and T: 15, 20°C and equaled to 0.042  $\text{pg cell}^{-1}$  and 0.041  $\text{pg cell}^{-1}$ , respectively (Fig. 3B, a). On average, salinity  
298 increase had a negative impact on pigmentation. The lowest cell-specific concentrations of Chl *a* and Car in BA-124 cells  
299 were estimated in the same scenario: salinity 18 PSU, T 10°C, PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and were equal to 0.013  $\text{pg}$   
300  $\text{cell}^{-1}$  (Fig. 2B, d) and 0.009  $\text{pg cell}^{-1}$  (Fig. 3B, d), for Chl *a* and Car, respectively. Multiple comparisons tests pointed to the  
301 significance of PAR and T combined in influencing the pigmentation. According to the statistics, 47% and 54% of multiple  
302  
303

304 comparisons in Chl *a* and Car content analysis, were statistically significant (Tukey HSD, \*  $p < 0.05$ ) with 83% (for Chl *a*)  
305 and 79% (for Car) of them with the highest significance level (Tukey HSD, \*\*\*  $p < 0.001$ ).

306 It was also examined that PAR and T affected the Chl *a* cell-specific content (ANOVA,  $F_{9,32} = 6.5$ ,  $p < 0.001$ , ANOVA,  
307  $F_{9,32} = 11.1$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 21.5$ ,  $p < 0.001$  and ANOVA,  $F_{9,32} = 6.5$ ,  $p < 0.001$ , for salinity 3, 8, 13, 18 PSU,  
308 respectively) and Car cell-specific content (ANOVA,  $F_{9,32} = 8.6$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 9.6$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} =$   
309  $4.6$ ,  $p < 0.001$  and ANOVA,  $F_{9,32} = 26.8$ ,  $p < 0.001$ , for salinity 3, 8, 13, 18 PSU, respectively) of *Synechococcus* sp. BA-132  
310 significantly. It was found that salinity increase had a negative impact on cell-specific Chl *a* and Car concentrations. BA-132  
311 was richer in cell-specific pigments than BA-124 (Figs. 2C, 3C). Along with PAR increase, the Chl *a* concentration  
312 decreased significantly. The maximum Chl *a* cell-specific content was measured in moderate or high T (20°C in salinity 13  
313 PSU and 25°C in salinity 3, 8, 18 PSU) under the lowest PAR (10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ). These maximums were 0.299 pg  
314  $\text{cell}^{-1}$  in salinity 3 PSU (Fig. 2C, a), 0.248 pg  $\text{cell}^{-1}$  in salinity 8 PSU (Fig. 2C, b), 0.151 pg  $\text{cell}^{-1}$  in salinity 13 PSU (Fig. 2C,  
315 c) and 0.073 pg  $\text{cell}^{-1}$  in salinity 18 PSU (Fig. 2C, d). Consistently with Chl *a*, Car cell-specific content maximums were also  
316 measured under the lowest PAR (10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) but contrary to Chl *a*, at the lowest T (10°C). These maximums  
317 were: 0.194 pg  $\text{cell}^{-1}$  in salinity 3 PSU (Fig. 3C, a), 0.131 pg  $\text{cell}^{-1}$  in salinity 8 PSU (Fig. 3C, b), 0.097 pg  $\text{cell}^{-1}$  in salinity 13  
318 PSU (Fig. 3C, c), 0.062 pg  $\text{cell}^{-1}$  in salinity 18 PSU (Fig. 3C, d). Minimums of Chl *a* and Car cell-specific contents within all  
319 scenarios were estimated in salinity 18 PSU, T 15°C and PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  being equal to 0.020 pg  $\text{cell}^{-1}$  (Fig.  
320 2C, d) and 0.19 pg  $\text{cell}^{-1}$  (Fig. 3C, d), for Chl *a* and Car, respectively. Regarding Chl *a* for minimum content per cell the  
321 same concentration as above mentioned (0.020 pg  $\text{cell}^{-1}$ ) was also estimated in salinity 13 PSU for the same conditions of T  
322 and PAR (Fig. 2C, c). Tukey HSD tests pointed to the significance of PAR and T combined in impacting the pigmentation.  
323 According to those tests, 66% and 61% of multiple comparisons in Chl *a* and Car content analysis, respectively, were  
324 statistically significant (Tukey HSD, \*  $p < 0.05$ ), with 81% (for Chl *a*) and 75% (for Car) of them with the highest  
325 significance (Tukey HSD, \*\*\*  $p < 0.001$ ).

326

### 327 3.3 Chl *a* fluorescence

328

329 The parameters of Chl *a* fluorescence were depicted as two-factor-dependent graphs, where the values in between the  
330 specific measurements were interpolated (Figs. 4, 5). For all strains, Chl *a* fluorescence parameters were measured and  
331 examined. These parameters were: the maximum photochemical efficiency of photosystem II (PSII) at dark-adapted state  
332 ( $F_v/F_m$ ) and the photochemical efficiency of PSII under actinic light intensity ( $\Phi\text{PSII}$ ).

333 The results showed that PAR and T affected  $F_v/F_m$  (ANOVA,  $F_{9,32} = 5.2$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 5.7$ ,  $p < 0.001$ ,  
334 ANOVA,  $F_{9,32} = 4.8$ ,  $p < 0.001$  and ANOVA,  $F_{9,32} = 33.9$ ,  $p < 0.001$ , for salinity 3, 8, 13, 18 PSU, respectively) and  $\Phi\text{PSII}$   
335 (ANOVA,  $F_{9,32} = 4.5$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 5.7$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 6.3$ ,  $p < 0.001$  and ANOVA,  $F_{9,32} = 2.3$ ,  $p$   
336  $< 0.05$ , for salinity 3, 8, 13, 18 PSU, respectively) of *Synechococcus* sp. BA-120 significantly. For this strain, especially in  
337 low T scenarios and in all scenarios with the lowest salinity, higher  $F_v/F_m$  was observed for 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  when  
338 compared to 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. 4A, a). Generally, strong fluctuations were noticeable in  $F_v/F_m$  values, which  
339 disabled the fixed environmentally driven pattern determination. However, there was a constant relation noted between T  
340 and PAR and  $\Phi\text{PSII}$ . PAR and T increase had a negative impact on  $\Phi\text{PSII}$ . The impact was the strongest in low salinity  
341 (Figs. 5A, a-b). Nonetheless, in each salinity, the lowest  $\Phi\text{PSII}$  were observed under the highest T and elevated PAR (190 or  
342 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ). On the contrary, the highest  $\Phi\text{PSII}$  values were calculated in the lowest T and PAR conditions in  
343 every salinity. The highest  $F_v/F_m$ , for all BA-120 experiments equaled 0.804 and was estimated for scenario: salinity 18  
344 PSU, T 10°C, PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. 4A, d). Generally, maximum values of  $F_v/F_m$  in each medium were  
345 associated with the lowest temperature. Minimum  $F_v/F_m$  within all scenarios was estimated for salinity 3 PSU, T 25°C and  
346 PAR 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (0.409, Fig. 4A, a). Concerning  $\Phi\text{PSII}$ , the greatest value was 0.768 estimated in salinity 18  
347 PSU, T 10°C and PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. 5A, d). Minimum  $\Phi\text{PSII}$  was measured in salinity 3 PSU, T 25°C and

348 PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (0.241, Fig. 5A, a). Multiple comparisons tests pointed to a strong environmental influence  
349 on Chl *a* fluorescence parameters. Regarding  $F_v/F_m$ , 65% of all comparisons were statistically significant (Tukey HSD, \*  $p < 0.05$ ) with 78% of them having the highest significance (Tukey, HSD, \*\*\*  $p < 0.001$ ). For  $\Phi\text{PSII}$  the percentages were as  
350 follows: 80% of all comparisons were statistically significant (Tukey HSD, \*  $p < 0.05$ ) and 87% of them had the highest  
351 significance (\*\*\*  $p < 0.001$ ).  
352

353 Both PAR and T significantly affected  $F_v/F_m$  (ANOVA,  $F_{9,32} = 46.2$ , \*\*\*  $p < 0.001$ , ANOVA,  $F_{9,32} = 5.1$ , \*\*\*  $p < 0.001$ ,  
354 ANOVA,  $F_{9,32} = 5.0$ , \*\*\*  $p < 0.001$  and ANOVA,  $F_{9,32} = 20.6$ , \*\*\*  $p < 0.001$ , for 3, 8, 13, 18 PSU, respectively) and  $\Phi\text{PSII}$   
355 (ANOVA,  $F_{9,32} = 25.0$ , \*\*\*  $p < 0.001$ , ANOVA,  $F_{9,32} = 11.6$ , \*\*\*  $p < 0.001$ , ANOVA,  $F_{9,32} = 15.4$ ,  $p < 0.001$  and ANOVA,  
356  $F_{9,32} = 5.2$ ,  $p < 0.001$ , for 3, 8, 13, 18 PSU, respectively) of *Synechococcus* sp. BA-124. For this strain,  $F_v/F_m$  reached the  
357 lowest values when compared to the respective incubations of other strains. The values of  $F_v/F_m$  generally decreased along  
358 with PAR and T increases but with some exceptions. Generally,  $\Phi\text{PSII}$  environmentally driven characteristics were similar to  
359  $F_v/F_m$  characteristics. The  $F_v/F_m$  minimums were measured under the lowest T and highest PAR in each salinity (Figs. 4B, a-  
360 d). The lowest value within all scenarios was 0.124 and was observed in salinity 3 PSU, T 10°C and PAR 280  $\mu\text{mol photons}$   
361  $\text{m}^{-2} \text{s}^{-1}$  (Fig. 4B, a). The  $F_v/F_m$  maximums were estimated for the highest T and the lowest PAR in each salinity. The highest  
362  $F_v/F_m$  equaled 0.560 for salinity 3 PSU, T 25°C and PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. 4B, a). Minimums of  $\Phi\text{PSII}$ ,  
363 consistently with  $F_v/F_m$ , were noted under the lowest T and highest PAR. The lowest  $\Phi\text{PSII}$  within all BA-124 experiments  
364 was 0.114 (followed by the minimum in salinity 3 PSU being equal to 0.116, Fig. 5B, a) and was measured in salinity 13  
365 PSU (Fig. 5B, c). Maximums of  $\Phi\text{PSII}$  were observed in the highest T and lowest PAR in each medium, similarly to  $F_v/F_m$ .  
366 The greatest value of  $\Phi\text{PSII}$  was 0.542 and was measured in salinity 3 PSU, T 25°C and PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig.  
367 5B, a). Tukey HSD post hoc test showed that PAR and T combined influenced Chl *a* fluorescence parameters significantly.  
368 Concerning  $F_v/F_m$ , 77% of all comparisons were statistically significant (\*  $p < 0.05$ ) with 88% of them having the highest  
369 significance (\*\*\*  $p < 0.001$ ). For  $\Phi\text{PSII}$  the percentages were as follows: 79% of all comparisons were statistically  
370 significant (\*  $p < 0.05$ ) and 89% of them had the highest significance (\*\*\*  $p < 0.001$ ).  
371

372 It was found that both PAR and T affected  $F_v/F_m$  (ANOVA,  $F_{9,32} = 4.3$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 4.8$ ,  $p < 0.001$ ,  
373 ANOVA,  $F_{9,32} = 4.5$ ,  $p < 0.001$  and ANOVA,  $F_{9,32} = 5.7$ ,  $p < 0.001$ , for salinity 3, 8, 13, 18 PSU, respectively) and  $\Phi\text{PSII}$   
374 (ANOVA,  $F_{9,32} = 10.1$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 7.7$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 4.7$ ,  $p < 0.001$  and ANOVA,  $F_{9,32} = 7.0$ ,  
375  $p < 0.001$ , for salinity 3, 8, 13, 18 PSU, respectively) of *Synechococcus* sp. BA-132, significantly. For this strain,  $F_v/F_m$   
376 decreased along with the PAR increase but was positively affected by T in each salinity (Figs. 4C, a-d). Minimum values of  
377  $F_v/F_m$  were measured in the highest PAR and the lowest T in each salinity. The lowest  $F_v/F_m$  within all experiments on BA-  
378 132 stated for salinity 13 PSU ( $F_v/F_m = 0.155$ , Fig. 4C, c). In salinity 3 PSU, under aforementioned conditions of T and  
379 PAR, the  $F_v/F_m$  value was also low compared to the others and equaled 0.160 (Fig. 4C, a). The maximums of  $F_v/F_m$  were  
380 measured in T 25°C and PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ . This was the case for all mediums. The highest  $F_v/F_m$  were noted in  
381 salinities 13 and 18 PSU and equaled 0.742 (Fig. 4C, c) and 0.733 (Fig. 4C, d), respectively. The lowest  $\Phi\text{PSII}$  were noted  
382 under the highest PAR and T conditions in every salinity (Figs. 5C, a-d). The minimum  $\Phi\text{PSII}$ , within all gathered results,  
383 was obtained in salinity 3 PSU and equaled 0.281 (Fig. 5C, a). Maximums of  $\Phi\text{PSII}$  were measured under completely  
384 opposite conditions to the ones stating for minimums, i.e. the lowest PAR and T. The highest  $\Phi\text{PSII}$ , 0.786, was noted in  
385 salinity 8 PSU, T 10°C and PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. 5C, b). The  $\Phi\text{PSII}$  reached generally higher values than  
386  $F_v/F_m$  in BA-132 experiments.  $\Phi\text{PSII}$  reached lower values than  $\Phi\text{PSII}$  measured under respective conditions for two other  
387 strains. Multiple comparisons tests point to a strong environmental influence on Chl *a* fluorescence parameters. For  $F_v/F_m$ ,  
388 78% of all comparisons were statistically significant (Tukey HSD, \*  $p < 0.05$ ) with 89% of them with the highest  
389 significance (Tukey, HSD, \*\*\*  $p < 0.001$ ). For  $\Phi\text{PSII}$ , 82% of all comparisons were statistically significant (Tukey HSD, \*  $p$   
389  $< 0.05$ ), with 89% of them having the highest significance level (Tukey, HSD, \*\*\*  $p < 0.001$ ).

390 Generally, for the BA-120 strain,  $F_v/F_m$  was affected negatively by T increase, while BA-124 and BA-132 strains were  
391 affected positively. T increase had a positive impact on  $\Phi$ PSII for BA-124 and a negative impact for BA-120 and BA-132.  
392 On average,  $\Phi$ PSII decreased along with PAR increase in all cultures.

### 394 3.4 Photosynthesis

395  
396 Net photosynthetic light-response curves for three PCY strains were analyzed. For all cultures, the photosynthesis  
397 parameters were: maximum of photosynthesis, photosynthesis efficiency at low irradiance, and dark respiration ( $P_m$ ,  $\alpha$ ,  $R_d$ ,  
398 respectively) and these were estimated for Chl *a*-specific and cell-specific domains (Figs. S4-S6 in Supplement). It should be  
399 noted that dark respiration values were negative (less oxygen than carbon dioxide (CO<sub>2</sub>)), which meant the lower  $R_d$ , the less  
400 net oxygen concentration was. This, in turn, indicated higher respiration rate.

401 For BA-120 statistical study showed significant dependence of PAR and T on Chl *a*-specific  $P_m$  in salinities 3, 8 and 18  
402 PSU (ANOVA,  $F_{9,32} = 2.4$ ,  $p < 0.05$ ,  $F_{9,32} = 3.2$ ,  $p < 0.05$  and  $F_{9,32} = 5.2$ ,  $p < 0.001$ , respectively) and pointed to no  
403 statistically significant dependence of ecological conditions on  $P_m$  in salinity 13 PSU (ANOVA,  $p \geq 0.05$ ). Regarding cell-  
404 specific  $P_m$  there was no statistically significant influence of PAR and T on this parameter in salinity 18 PSU (ANOVA,  $p \geq$   
405  $0.05$ ) but was in salinity 3 PSU (ANOVA,  $F_{9,32} = 3.5$ ,  $p < 0.05$ ), 8 PSU (ANOVA,  $F_{9,32} = 2.6$ ,  $p < 0.05$ ), and 13 PSU  
406 (ANOVA,  $F_{9,32} = 3.0$ ,  $p < 0.05$ ). For Chl *a*-specific  $\alpha$ , statistical study indicated no environmental impacts in salinities 3, 8  
407 and 13 PSU but an impact of PAR and T in salinity 18 PSU (ANOVA,  $F_{9,32} = 2.7$ ,  $p < 0.05$ ), while for cell-specific  $\alpha$   
408 statistical significance of PAR and T influence was obtained for all salinities (ANOVA,  $F_{9,32} = 5.1$ ,  $p < 0.001$ , ANOVA,  $F_{9,32}$   
409  $= 2.9$ ,  $p < 0.05$ , ANOVA,  $F_{9,32} = 2.5$ ,  $p < 0.05$  and ANOVA,  $F_{9,32} = 4.8$ ,  $p < 0.001$ , for salinity 3, 8, 13 and 18 PSU,  
410 respectively). Regarding  $R_d$ , two-way ANOVA pointed to no environmental determination of Chl *a*-specific  $R_d$  values  
411 (ANOVA,  $p > 0.05$ ) but it showed the influence of PAR and T on cell-specific  $R_d$  (ANOVA,  $F_{9,32} = 9.2$ ,  $p < 0.001$ , ANOVA,  
412  $F_{9,32} = 3.8$ ,  $p < 0.01$ , ANOVA,  $F_{9,32} = 3.8$ ,  $p < 0.01$ , ANOVA,  $F_{9,32} = 4.5$ ,  $p < 0.001$ , in salinities 3, 8, 13, 18 PSU,  
413 respectively). Tukey HSD tests pointed to some statistically significant multiple comparisons but showed a weak influence  
414 of PAR and T combined on Chl *a*-specific parameters. Regarding  $\alpha$ , only 3% of all multiple comparisons were statistically  
415 significant (\*  $p < 0.05$ ) with 7% of them at the highest statistical significance level (\*\*\*)  $p < 0.001$ . For  $P_m$ , 36% of all  
416 multiple comparisons were statistically significant (\*  $p < 0.05$ ) with 64% of them with the highest significance (\*\*\*)  $p <$   
417  $0.001$ . Regarding  $R_d$ , as mentioned above, no statistically significant analysis of variance was indicated. Due to that, no post  
418 hoc tests were proceeded. Note that in this section, to make it more concise, the notation for all statistically significance  
419 multiple comparisons percentage (\*  $p < 0.05$ ) and the percentage of the multiple comparisons of the highest significance  
420 within the significant ones (\*\*\*)  $p < 0.001 \times (* p < 0.05)^{-1}$  were written in parenthesis, one by one, separated with comma.  
421 Similarly to Chl *a*-specific calculations, Tukey HSD test pointed to a selective influence of PAR and T combined on cell-  
422 specific parameters. However, this dependence was stronger when compared to Chl *a*-specific estimations ( $P_m$  (16%, 52%),  
423  $\alpha$  (19%, 43%),  $R_d$  (28%, 56%)). Nonetheless, there were also some fixed relations noted for both calculation domains. For  
424 Chl *a*-specific photosynthesis,  $P_m$  increased along with PAR up to PAR of 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Figs. S4, a, c). Above  
425 this level  $P_m$  value started to decrease slightly. This was the case in all salinities. Minimum  $P_m$  was measured for cells grown  
426 in scenario: salinity 3 PSU, T 15°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and it was 0.12  $\mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{h}^{-1}$  (Fig. S4, a),  
427 whilst the maximum equalled 1.31  $\mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{h}^{-1}$  and was reached in salinity 18 PSU, T 25°C, 190  $\mu\text{mol photons}$   
428  $\text{m}^{-2} \text{s}^{-1}$  (Fig. S4, c). Dark respiration rate ( $R_d$ ) increased with T increase and decreased with PAR increase (Figs. S5, a, c).  
429 Minimum  $R_d$  (-0.31  $\mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{h}^{-1}$ ) was measured in salinity 18 PSU, T 10°C, PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$   
430 (Fig. S5, c), while maximum (-0.02  $\mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{h}^{-1}$ ) was estimated in salinity 3 PSU, T 25°C, PAR 10  $\mu\text{mol}$   
431  $\text{photons m}^{-2} \text{s}^{-1}$  (Fig. S5, a). On the contrary, it was more difficult to determine a fixed pattern of  $\alpha$  changes unequivocally.  
432 The most fixed tendency of  $\alpha$  changes was observed between all temperature-differenced scenarios in 18 PSU salinity  
433 medium (Figs. S6, a, c). Under those conditions, it was noticeable that  $\alpha$  decreased with PAR and T increase till it reached

434 PAR level of 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ . Then,  $\alpha$  started to rise slowly. Regarding all gathered results (all mediums together),  
435 minimum  $\alpha$  was measured in salinity 3 PSU, T 25°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and equalled 0.002  $\mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{ h}^{-1}$  [ $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ] $^{-1}$  (Fig. S6, a), whilst maximum was 0.013  $\mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{ h}^{-1}$  [ $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ] $^{-1}$  in  
436 salinity 13 PSU, T 10°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ . On the other hand, for cell-specific domain,  $P_m$  increased along with  
437 T and it was more pronounced in higher salinities. Concerning all results, minimum  $P_m$  was 28.58  $\mu\text{mol O}_2 \text{ cell } 10^{-9} \text{ h}^{-1}$  and,  
438 similarly to Chl  $a$ -specific  $P_m$  was measured in scenario: salinity 13 PSU, T 10°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ , whilst  
439 maximum  $P_m$  equalled 55.16  $\mu\text{mol O}_2 \text{ cell } 10^{-9} \text{ h}^{-1}$  and was reached in salinity 8 PSU, T 25°C, 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$   
440 (data not shown). Regarding  $\alpha$ , this parameter was generally negatively affected by PAR and T up to PAR of 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ . However minimum value was obtained for cells growing in moderate T (salinity 8 PSU, T 20°C, PAR 10  
441  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) and equalled 0.81  $\mu\text{mol O}_2 \text{ cell } 10^{-9} \text{ h}^{-1}$  [ $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ] $^{-1}$ . Maximum  $\alpha$  equalled 1.57  $\mu\text{mol O}_2$   
442  $\text{cell } 10^{-9} \text{ h}^{-1}$  [ $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ] $^{-1}$  and was measured in salinity 18 PSU, T 10°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. S6,  
443 d). Generally, T and PAR had a positive impact on  $R_d$  for cultures grown in PAR range up to 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ . For  
444 cultures grown under elevated PAR conditions,  $R_d$  was lower (more intensive respiration) when compared to low PAR  
445 scenarios. The lowest  $R_d$  within all BA-120 experiments results was -16.97  $\mu\text{mol O}_2 \text{ cell } 10^{-9} \text{ h}^{-1}$  and noted in salinity 3  
446 PSU, T 10°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. S5, b), whilst the highest  $R_d$  was measured in salinity 18 PSU, T 25°C,  
447 PAR 100  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and equalled -2.06  $\mu\text{mol O}_2 \text{ cell } 10^{-9} \text{ h}^{-1}$  (Fig. S5, d).

450 For BA-124, statistical study showed significant dependence of ecological conditions on photosynthesis parameters,  
451 excluding Chl  $a$ -specific  $\alpha$  (ANOVA,  $p \geq 0.05$ ) and cell-specific  $P_m$  (ANOVA,  $p \geq 0.05$ ). For the rest parameters ANOVA  
452 results were as follows: Chl  $a$ -specific  $P_m$  (ANOVA,  $F_{9,32} = 4.8$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 19.7$ ,  $p < 0.001$ , ANOVA,  $F_{9,32}$   
453  $= 9.14$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 6.5$ ,  $p < 0.001$  in salinity 3, 8, 13, 18 PSU, respectively); cell-specific  $P_m$  (ANOVA,  $F_{9,32}$   
454  $= 7.5$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 6.1$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 4.3$ ,  $p < 0.001$  in salinity 8, 13 and 18 PSU, respectively);  
455 Chl  $a$ -specific  $\alpha$  (ANOVA,  $F_{9,32} = 5.0$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 3.3$ ,  $p < 0.01$ , ANOVA,  $F_{9,32} = 3.8$ ,  $p < 0.01$  in salinity 3,  
456 8 and 18 PSU, respectively); cell-specific  $\alpha$  (ANOVA,  $F_{9,32} = 6.6$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 17.9$ ,  $p < 0.001$ , ANOVA,  
457  $F_{9,32} = 18.9$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 3.1$ ,  $p < 0.01$ , in salinity 3, 8, 13, 18 PSU, respectively); Chl  $a$ -specific  $R_d$  (ANOVA,  
458  $F_{9,32} = 10.0$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 4.9$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 3.8$ ,  $p < 0.01$ , ANOVA,  $F_{9,32} = 2.6$ ,  $p < 0.05$ , in  
459 salinity 3, 8, 13, 18 PSU, respectively); cell-specific  $R_d$  (ANOVA,  $F_{9,32} = 13.0$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 2.2$ ,  $p < 0.05$ ,  
460 ANOVA,  $F_{9,32} = 40.4$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 3.1$ ,  $p < 0.01$ ). Post-hoc tests showed there must have been other factors,  
461 which affected the whole process of photosynthesis as there were many not statistically significant multiple comparisons  
462 defined. Generally, Tukey HSD tests pointed to only few statistically significant multiple comparisons, in both Chl  $a$ -  
463 specific, especially for  $P_m$ , ( $P_m$  (60%, 76%),  $\alpha$  (9%, 29%),  $R_d$  (30%, 47%) and cell-specific ( $P_m$  (22%, 56%),  $\alpha$  (34%, 63%),  
464  $R_d$  (30%, 74%)) estimations. Nonetheless, for  $P_m$  there was a tendency noted, which suggested that on average, the maximum  
465 of photosynthesis was higher at elevated PAR. This was the case in both estimations, Chl  $a$ -specific and cell-specific.  
466 Maximum Chl  $a$ -specific  $P_m$  was 3.0 and minimum 0.16  $\mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{ h}^{-1}$ . These values were measured in salinity 18  
467 PSU in T 25°C, PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and T 10°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ , respectively (Fig. S4, g).  
468 Maximum cell-specific  $P_m$  was obtained in salinity 8 PSU, T 25°C, PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and minimum in salinity  
469 13 PSU, T 20°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (data not shown here). These extreme values were 53.41 and 19.17  $\mu\text{mol O}_2$   
470  $\text{cell} \cdot 10^{-9} \text{ h}^{-1}$ , respectively. It was difficult to determine a fixed relation between ecological state and  $\alpha$  changes in both  
471 domains, which was supported by the post-hoc test (more than 91% of multiple comparisons were not statistically significant  
472 ( $p \geq 0.05$ ) in Chl  $a$ -specific and more than 35% in cell-specific estimations). Maximum Chl  $a$ -specific  $\alpha$  was 0.02  $\mu\text{mol O}_2$   
473  $(\mu\text{g Chl } a)^{-1} \text{ h}^{-1}$  [ $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ] $^{-1}$  and was measured in salinity 3 PSU, T 15°C, PAR 100  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig.  
474 S6, e), while maximum cell-specific  $\alpha$  (1.77  $\mu\text{mol O}_2 \text{ cell } 10^{-9} \text{ h}^{-1}$  [ $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ] $^{-1}$ ) was obtained in salinity 13 PSU,  
475 T 10°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ . Minimum Chl  $a$ -specific  $\alpha$  was 0.003  $\mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{ h}^{-1}$  [ $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ] $^{-1}$   
476 and was measured in two scenarios: salinity 3 PSU, T 10°C, PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. S6, e) and salinity 18  
477 PSU, T 15°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. S6, g). Minimum cell-specific  $\alpha$  equalled 0.08  $\mu\text{mol O}_2 \text{ cell } 10^{-9} \text{ h}^{-1}$  [ $\mu\text{mol}$

478 photons  $\text{m}^{-2} \text{s}^{-1}$  and was measured in salinity 18 PSU, T 15°C, PAR 190  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. S6, h). Similarly to  $\alpha$ , it  
479 was difficult to determine fixed relations between PAR and T and  $R_d$ , which was supported by statistics (about 70% of  
480 multiple comparisons for both Chl *a*-specific and cell-specific  $R_d$  were not statistically significant (Tukey HSD,  $p \geq 0.05$ )).  
481 Nonetheless, it was observed that, generally,  $R_d$  decreased along with PAR increase in cell-specific estimations. Maximum  
482 Chl *a*-specific and cell-specific  $R_d$  was  $-0.03 \mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{h}^{-1}$  and  $-1.52 \mu\text{mol O}_2 \text{ cell } 10^{-9}$ , respectively. These  
483 values were obtained in salinity 13 PSU, T 20°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and salinity 18 PSU, T 20°C, PAR 190  $\mu\text{mol}$   
484  $\text{photons m}^{-2} \text{s}^{-1}$ , respectively for Chl *a*- and cell-specific calculations. Minimum Chl *a*-specific  $R_d$  was measured in salinity  
485 13 PSU, T 10°C, PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and was  $-0.27 \mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{h}^{-1}$ , whilst minimum cell-specific  $R_d$   
486 was measured in salinity 13 PSU, T 10°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and equalled  $-12.19 \mu\text{mol O}_2 \text{ cell } 10^{-9} \text{h}^{-1}$  (data not  
487 shown here).

488 For BA-132, statistical study showed significant dependence of PAR and T on Chl *a*- and cell-specific  $P_m$  (for Chl *a*-  
489 specific: ANOVA,  $F_{9,32} = 6.2$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 23.1$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 25.2$ ,  $p < 0.001$ , ANOVA,  $F_{9,32}$   
490  $= 16.0$ ,  $p < 0.001$ ; for cell-specific: ANOVA,  $F_{9,32} = 4.8$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 24.3$ ,  $p < 0.001$ , ANOVA,  $F_{9,32} = 24.3$ ,  
491  $p < 0.001$ , ANOVA,  $F_{9,32} = 21.2$ ,  $p < 0.001$ ; all numbers given for salinities 3, 8, 13, 18 PSU, respectively). Regarding other  
492 Chl *a*-specific parameters, there were no statistically significant impacts of PAR and T on  $\alpha$  in salinities 3, 13, 18 PSU  
493 (ANOVA,  $p \geq 0.05$ ) but were in salinity 8 PSU (ANOVA,  $F_{9,32} = 2.7$ ,  $p < 0.05$ ) and no impacts on Chl *a*-specific  $R_d$  in  
494 salinities 3, 8, 18 PSU (ANOVA,  $p \geq 0.05$ ) but were in salinity 13 PSU (ANOVA,  $F_{9,32} = 2.8$ ,  $p < 0.05$ ). Regarding other  
495 than  $P_m$  cell-specific parameters, there was no ecological determination of  $\alpha$  noted in salinities 3 and 8 PSU and of  $R_d$  in  
496 salinity 13 PSU (ANOVA,  $p \geq 0.05$ ), while there were statistically significant environmental impacts calculated for  $\alpha$  in  
497 salinity 13 PSU (ANOVA,  $F_{9,32} = 3.2$ ,  $p < 0.01$ ) and 18 PSU (ANOVA,  $F_{9,32} = 2.9$ ,  $p < 0.05$ ) and for  $R_d$  in salinities 3, 8 and  
498 18 PSU (ANOVA,  $F_{9,32} = 3.2$ ,  $p < 0.05$ , ANOVA,  $F_{9,32} = 3.1$ ,  $p < 0.01$ , ANOVA,  $F_{9,32} = 2.4$ ,  $p < 0.05$ , respectively). Tukey  
499 HSD tests pointed to statistically significant multiple comparisons, in both Chl *a*-specific and cell-specific maximum of  
500 photosynthesis ( $P_m$  (68%, 85%),  $P_m$  (62%, 76%), respectively). Post hoc tests indicated no significant multiple comparisons  
501 for Chl *a*-specific  $\alpha$  (>1%, >1%), a few significant multiple comparisons for Chl *a*-specific  $R_d$  (8%, 38%), cell-specific  $\alpha$   
502 (18%, 67%) and cell-specific  $R_d$  (6%, 20%). It was observed, that in cell-specific estimations,  $P_m$  increased along with PAR  
503 increase, while  $\alpha$  decreased at elevated PAR. It was the most difficult to determine a fixed tendency for the  $R_d$  response to  
504 changing environmental conditions. This was supported by statistical tests (Tukey HSD, more than 93% of multiple  
505 comparisons were not statistically significant ( $p \geq 0.05$ )). Maximum cell-specific  $P_m$  was  $158.94 \mu\text{mol O}_2 \text{ cell } 10^{-9} \text{h}^{-1}$  and  
506 was reached in salinity 8 PSU, T 25°C, PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ , whilst minimum equalled  $28.04 \mu\text{mol O}_2 \text{ cell } 10^{-9} \text{h}^{-1}$   
507 in salinity 18 PSU, T 15°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. S4, l). Maximum cell-specific  $\alpha$  was  $1.78 \mu\text{mol O}_2 \text{ cell } 10^{-9}$   
508  $\text{h}^{-1} \mu\text{mol photons m}^{-2} \text{s}^{-1}$  and was measured in salinity 13 PSU, T 20°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ , while minimum was  
509 reached in salinity 18 PSU, T 20°C, PAR 100  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and equalled  $0.19 \mu\text{mol O}_2 \text{ cell } 10^{-9} \text{h}^{-1}$  [ $\mu\text{mol photons}$   
510  $\text{m}^{-2} \text{s}^{-1}$ ]<sup>-1</sup> (Fig. S6, l). Regarding cell-specific  $R_d$ , maximum was measured in salinity 18 PSU, T 15°C, PAR 100  $\mu\text{mol}$   
511  $\text{photons m}^{-2} \text{s}^{-1}$  and equalled  $-3.17 \mu\text{mol O}_2 \text{ cell } 10^{-9} \text{h}^{-1}$  (Fig. S5, l), whilst minimum was  $-15.55 \mu\text{mol O}_2 \text{ cell } 10^{-9} \text{h}^{-1}$  and  
512 was obtained in salinity 3 PSU, T 10°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. S5, j). For Chl *a*-specific  $P_m$ , the increases along  
513 with T and salinity was observed, whilst  $\alpha$  presented strong changing characteristics between scenarios. The fixed influence  
514 of PAR and T on  $\alpha$  values was difficult to determine, which was supported by statistics (ANOVA,  $p \geq 0.05$ ). Contrary to the  
515 above, it was plainly evident that PAR increase had a negative impact on Chl *a*-specific  $R_d$ . Maximum Chl *a*-specific  $P_m$  was  
516  $6.22 \mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{h}^{-1}$  and was reached in salinity 18 PSU, T 25°C, PAR 280  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. S4, k),  
517 whilst minimum equalled  $0.12 \mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{h}^{-1}$  in salinity 3 PSU, T 25°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. S4, i).  
518 Maximum Chl *a*-specific  $\alpha$  was  $0.02 \mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{h}^{-1}$  [ $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ]<sup>-1</sup> and was measured in salinity 18 PSU,  
519 T 15°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Fig. S6, k), while minimum was reached in salinity 3 PSU, T 15°C, PAR 10  $\mu\text{mol}$   
520  $\text{photons m}^{-2} \text{s}^{-1}$  and equalled  $0.003 \mu\text{mol O}_2 (\mu\text{g Chl } a)^{-1} \text{h}^{-1}$  [ $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ]<sup>-1</sup> (Fig. S6, i). Concerning Chl *a*-specific  
521  $R_d$ , maximum was measured in salinity 3 PSU, T 20°C, PAR 10  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  and equalled  $-0.02 \mu\text{mol O}_2 \text{ cell } 10^{-9}$

522  $\text{h}^{-1}$  (Fig. S5, i), whilst minimum was  $-0.39 \mu\text{mol O}_2 \text{ cell}^{-1} \text{ h}^{-1}$  and was obtained in salinity 13 PSU, T  $25^\circ\text{C}$ , PAR 280  
523  $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ . Generally, in both domains, photosynthesis parameters were the highest for BA-132 when compared  
524 to other strains.

525 The analysis of photosynthesis characteristics enabled examining and defining the photoacclimation process of all three  
526 strains of *Synechococcus* sp. This was done on the basis of the photosynthetic parameters (Figs. S4-S6) and Photosynthesis-  
527 Irradiance (*P-E*) curves (exemplification shown in Fig. 6). The curves were plotted on the basis of laboratory results (Clark  
528 oxygen electrode measurements) using the equation of Jassby and Platt (1976). According to a photoacclimation model  
529 description (Prezelin, 1981; Prezelin and Sweeney, 1979; Ramus, 1981; Richardson et al., 1983; Pniewski et al., 2016), the  
530 results of the present study indicated changes in Photosynthetic Units (PSU) sizes as the photoacclimation mechanism,  
531 which occurred most frequently (Table 1). There were also *P-E* curves pointing to some changes in enzymatic reactions and  
532 the altering of accessory pigments activity. Changes in PSU numbers were noted as well, but these observations were  
533 episodic. In this paper the term 'OTHER' states for changes in enzymatic reactions and the altering of accessory pigments  
534 activity and concerns photoacclimation mechanisms other than changes in PSU sizes (PSUsize) or changes in PSU number  
535 (PSUno.). In general, photoacclimation did not occur in low-saline medium (salinity 3). According to our results, the process  
536 was observed in only four cases in low salinity: BA-120  $25^\circ\text{C}$  salinity 3 PSU, BA-124  $25^\circ\text{C}$  salinity 3 PSU, BA-132  $10^\circ\text{C}$   
537 salinity 3 PSU, and BA-132  $25^\circ\text{C}$  salinity 3 PSU. For BA-120, photoacclimation occurred more frequently at higher T (20  
538 and  $25^\circ\text{C}$ ) than lower T (10 and  $15^\circ\text{C}$ ). However, if it had been observed in low T conditions, it usually stated for OTHER,  
539 not for PSUsize or PSUno. For BA-124 and BA-132 photoacclimation was noted in the whole T range. All photoacclimation  
540 mechanisms observed for different strains are listed in Table 1.

#### 542 4 Discussion

543  
544 Picoplanktonic organisms show a lot of adaptations, which enable them to spread in aquatic environments. What is more,  
545 picocyanobacteria often dominate and occupy the niches, which are inaccessible for other photoautotrophs. Owing to the fact  
546 that PCY are small-sized cells and consequently possess an advantageous surface area to volume ratio, they can assimilate  
547 trace amount of nutrients and effectively absorb light. Therefore, in oligotrophic regions of seas and oceans PCY compete  
548 with other cyanobacteria and microalgae and it can determine primary production of the whole marine ecosystem (Six et al.,  
549 2007a; Richardson and Jackson, 2007; Worden and Wilken, 2016). This is also true for eutrophic basins (Stal et al., 2003;  
550 Haverkamp et al., 2008; 2009; Callieri, 2010; Mazur-Marzec et al., 2013).

551 The distribution of PCY are determined by their optimal ecological requirements for light and temperature. Due to the  
552 presented results, PAR and T had positive effects on the number of cells for two out of the three studied strains of  
553 *Synechococcus* sp. The highest cell concentrations were noted at the highest T ( $25^\circ\text{C}$ ) and the highest PAR level ( $280 \mu\text{mol}$   
554  $\text{photons m}^{-2} \text{ s}^{-1}$ ) for BA-124 and BA-132. The BA-120 strain behaved differently when compared to the other two in high  
555 PAR conditions. The decrease in number of cells appeared then, i.e. cell abundances for cultures grown under the most  
556 elevated PAR were lower than the number of cells measured for *Synechococcus* sp. BA-120 cells grown under  $190 \mu\text{mol}$   
557  $\text{photons m}^{-2} \text{ s}^{-1}$ . According to the results derived from pigmentation, Chl *a* fluorescence and photosynthesis sections of the  
558 present study, the decrease in number of cells under the elevated PAR could have likely been associated with Photosystem II  
559 photo-inhibition. This was a result of a few observations, which are described as follows. Firstly, there was a higher cell-  
560 specific Car content observed for  $280 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$  when compared to  $190 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ . Secondly, higher  
561  $F_v/F_m$  values were observed for  $280 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$  when compared to  $190 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ , especially for low T  
562 scenarios and for all scenarios in the lowest salinity medium. Thirdly, for Chl *a*-specific photosynthesis,  $P_m$  increased along  
563 with PAR until  $190 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ , above which the values started to decrease slightly in all salinity mediums.  
564 According to the above, a PAR level of  $190 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$  could be defined as the PSII photo-inhibition point for the  
565 red strain. This implies BA-120 did not lead as effective photosynthesis being grown in PAR of more than  $190 \mu\text{mol photons}$

566  $\text{m}^{-2} \text{s}^{-1}$  as the cells grown in PAR levels equal or are beneath the above mentioned value. The results showed that in all  
567 synthetically created environmental scenarios, BA-124 was the strain of the highest cell abundance. This is consistent with  
568 the Baltic Sea field studies (Mazur-Marzec et al., 2013).

569 Cyanobacteria are generally recognized to prefer low light intensity for growth (Fogg and Thake, 1987; Ibelings, 1996).  
570 Some picoplanktonic organisms demonstrated the ability to survive and resume growth after periods of total darkness. Such  
571 a pronounced capacity for survival in the dark would enable these organisms to outlive the seasonal rhythm of winter  
572 darkness and sinking into the aphotic zone (Antia, 1976). The investigated strains of *Synechococcus* sp. were found to be  
573 well acclimated to relatively low and high PAR levels. The latter was especially evident at the high treatment T. This  
574 conclusion is consistent with the observations of picocyanobacteria maximum abundance at the euphotic zone in coastal and  
575 offshore marine waters (Stal et al., 2003; Callieri, 2010). Moreover, Kana and Glibert (1987a,b) showed that *Synechococcus*  
576 sp. could grow at irradiance as high as  $2000 \mu\text{mol photon m}^{-2} \text{s}^{-1}$ .

577 Surface and near-surface populations experience extremely variable light and temperature conditions (Millie et al.,  
578 1990), and these factors are the ones that affect the composition of photosynthetic pigments and photosynthesis performance  
579 of picocyanobacteria (Jodłowska and Śliwińska, 2014). Picocyanobacteria with a high concentration of PC are chromatically  
580 better adapted to harvest longer wavelengths of PAR than those with PE as a dominating pigment. Therefore, such  
581 picocyanobacteria, such as the BA-124 strain, usually dominate in surface euphotic waters (Stal et al., 2003; Haverkamp et  
582 al., 2008; 2009). On the other hand, the strains rich in PE (BA-120 and BA-132), usually occurred deeper (Fahnenstiel et al.,  
583 1991; Hauschild et al., 1991; Vörös et al., 1991). Nonetheless, generally PCY, thanks to their high concentration of  
584 photosynthetic pigments, may occur in waters under low light intensity (Stal et al., 2003). Carotenoids have a dual role in the  
585 cell: to maintain a high capacity for photosynthetic light absorption and to provide protection against photooxidation  
586 (Siefermann-Harms, 1987). This feature additionally explains why picoplanktonic *Synechococcus* is able to grow  
587 successfully both in the surface layer of the sea and also in deeper waters (Stal and Walsby, 2000; Stal et al., 2003). This  
588 research showed that for BA-120 cell-specific pigments content, there were very high concentrations of Chl *a* observed in  
589 the whole T domain under low PAR. This could have implied the photoacclimation type, which was the change in PSU  
590 number. This mechanism was also observed in *P-E* curves for scenario with salinity 8 PSU and temperature  $20^{\circ}\text{C}$ .

591 PAR and T were also the main factors influencing the changes in Chl *a* fluorescence in three strains of *Synechococcus*  
592 sp. and may be of a great importance in the context of PCY domination in many aquatic ecosystems during the summer  
593 period. Due to Chl *a* fluorescence parameters results, it should be noted that PAR increase always had a negative impact on  
594  $\Phi\text{PSII}$ , which implied that cells, previously acclimated to high light conditions, had lower PSII photosystem efficiency under  
595 actinic light.

596 The results showed that T, PAR and salinity influenced the photosynthesis parameters only to a certain degree. There  
597 were many not statistically significant multiple comparisons pointed by post hoc tests. However, it was found that generally,  
598 in cell-specific estimations, elevated PAR had a negative effect on  $\alpha$  and PAR increase influenced the respiration negatively.  
599 For each of the studied strains of *Synechococcus* sp., the highest  $\alpha$  and the lowest  $R_d$  were noted for the cells grown under the  
600 lowest PAR ( $10 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ ). On the other hand, the highest values of  $P_m$  were noted at the highest PAR. It pointed  
601 to inability for the cells incubated in low PAR conditions to be as effective in photosynthesis as the cells grown under high  
602 irradiances. According to our results, on the basis of *P-E* curves, three types of photoacclimation mechanisms of  
603 *Synechococcus* sp. were observed: change in PSU size, change in PSU number and altering accessory pigments activity and  
604 changes in enzymatic reactions. This was a striking observation because in the literature results predominantly derive the two  
605 first aforementioned types of recognition (Stal et al., 2003; Jodłowska and Śliwińska, 2014). The present study showed that  
606 changes in PSU size occur most frequently (Table 1). The second, ranked by frequency of occurrence, was the altering of  
607 accessory pigment activity. PSU number changes in *Synechococcus* sp. rarely occurred, which is consistent with literature  
608 (Jodłowska and Śliwińska, 2014). Moreover, in our study, photoacclimation mechanisms occurred less frequently in the  
609 scenarios with salinity 3 PSU. The changes of photosynthesis parameters ( $P_m$ ,  $\alpha$ ,  $R_d$ ) under different environmental

610 conditions explains the occurrence of different photoacclimation mechanisms. According to our results, *Synechococcus*  
611 strains present different ecophysiological characteristics, however, they all demonstrate their tolerance to elevated PAR (for  
612 BA-120 to a certain degree) and T levels and could have effectively acclimated to varied water conditions. These strains  
613 were able to change the composition of photosynthetic pigments in order to use light quanta better. The ability of  
614 *Synechococcus* to sustain their growth in low light conditions and their low photo-inhibition in exposure to high light  
615 intensities could give picocyanobacteria an advantage in optically changeable waters (Jasser, 2006).

616 Due to occurrence of extremes in salinity and other environmental conditions in the Baltic Sea area, the Baltic  
617 inhabitants are highly adapted to different regions and often reach their physiological limits (Snoeijis-Leijonmalm and  
618 Andrén, 2017). The changing environmental conditions the cultures were grown in during the experiments were salinity, T  
619 and PAR. Daily mean sea surface temperature (Leppäranta and Myrberg, 2009) presents strongly pronounced annual cycles  
620 in the Baltic Sea area. Sea surface temperature (SST) range between about 10 and 20°C may be timed in the Baltic between  
621 June and September with some inter-annual changes (Siegel and Gerth, 2017). SSTs reaching and exceeding 20°C are also  
622 observed in the Baltic basin. For instance, according to Siegel and Gerth (2017), SSTs higher than 20°C were recorded in  
623 almost whole Baltic area beyond Danish Straits, Bothnian Bay and northern Bothnian Basin in the warmest week of 2016, in  
624 July. According to above, the temperatures, under which the picocyanobacterium cultures were grown in the present study  
625 (10 – 25°C) can be defined as representative for the Baltic Sea. Furthermore, the salinity ranges applied in the experiment are  
626 also Baltic's representatives. The Baltic Sea horizontal salinity gradient is high and different sub-basins are characterized by  
627 different mean salinity values. The gradient decreases North towards. The highest salinity is observed in the Baltic Sea  
628 boundary to the North Sea (Skagerrak, mean salinity ranges between 28.34 and 32.71), while the lowest mean salinity is  
629 observed in the Baltic northernmost regions (around 2.35 – 3.96 in Bothnian Basin). These numbers were determined on the  
630 basis of climatological data from the Baltic Atlas of Long-Term Inventory and Climatology (Feistel et al., 2008; 2010).  
631 Thus, the presented analysis may derive accurate assumptions regarding the regional distribution of *Synechococcus* sp.  
632 strains in the Baltic Sea. For instance, a salinity horizontal gradient can be one of the factors determining the abundance of a  
633 certain strain in the basin. More saline waters are most preferred by BA-132. On that basis, one can assume the concentration  
634 of this strain will be higher near the Baltic Sea entrance (Danish Straits) than in Bothnian Bay. Additionally, it was observed  
635 that despite elevated PAR conditions being more suitable for BA-124 and BA-132 to grow intensively, all analyzed strains  
636 were able to survive and grow in low PAR conditions. This is consistent with other previously published Baltic studies (Stal  
637 et al., 2003; Jodłowska and Śliwińska, 2014) stating that this is caused by phycobilisomes, which are structural components  
638 of picocyanobacteria PSII photosystem. The presence of PCY cells throughout the whole euphotic water column was also  
639 reported in limnological studies (Becker et al., 2004, Callieri, 2007).

640 The discrepancies between the strains ecophysiology derived in this study amplified the need for in-depth investigation  
641 of three strains separately. What is more, according to the author's best knowledge, Baltic brown strain (BA-132) is the least  
642 recognized strain out of three analyzed *Synechococcus* sp. strains, so far. Stal et al. (2003) and Haverkamp et al. (2008)  
643 pointed to its inhabitation in the Baltic Sea but did not give its characteristics in detail. In more recent research new  
644 information has appeared, which has provided a more detailed examination of BA-132 (Jodłowska and Śliwińska, 2014).  
645 Nonetheless, this strain still required careful studies. The present paper derives the new knowledge on the BA-132 responses  
646 to changing ecological conditions. What is more, the study places BA-132 among the other *Synechococcus* sp. strains and  
647 compares their ecophysiology pointing to significant differences between these organisms.

648 The study of Baltic picoplankton ecophysiology is also of a great importance in the context of climate change.  
649 According to Belkin (2009), the Baltic Sea is among the Large Marine Ecosystems (LME), where the most rapid warming is  
650 being observed (the increase in SST between 1982 and 2006 > 0.9°C). Moreover, there are studies pointing to an increase of  
651 average winter temperatures in northern Europe by several degrees by the year 2100 (Meier, 2002). These along with the  
652 presented results, which suggest that all analyzed strains of *Synechococcus* sp. were positively affected by T can be a strong  
653 argument for further numerical research on examining the effect of long-term positive temperature trend on the abundance of

654 PCY in the Baltic Sea (the need for picoplankton model representation). What is more, the feedback relation, which is the  
655 surface most layer being warmed up by irradiance trapped in the cells of phytoplankton may derive interesting conclusions  
656 on the functioning of the ecosystem and the living organisms being the internal source of heat in the marine medium.

657 The observed feature that T increase had a positive impact on all strains' number of cells is also consistent with field  
658 studies, which indicate the seasonal cycle of PCY maximal abundances (Flombaum et al., 2013; Dutkiewicz et al., 2015;  
659 Worden and Wilken, 2016). Hajdu et al. (2007) showed that during the decline phase of Baltic cyanobacterial blooms in late  
660 summer, unicellular and colony-forming picocyanobacteria increased in abundance. Mazur-Marzec et al. (2013) indicated  
661 that in summer cyanobacterial biomass was usually high and ranged from 20% at the beginning of July to 97% in late July  
662 and August. Moreover, Paczkowska et al. (2017) pointed to the abundance of 40-90% in the summertime in the Baltic Sea  
663 and to PCY being a dominant size group in all Baltic basins. Stal et al. (1999) reported that 65% of the phytoplankton-  
664 associated Chl *a* concentration in the Baltic Proper during late summer belonged to picoplankton, while the second most  
665 dominant group was nitrogen-fixing cyanobacteria (*Aphanizomenon* sp., *Dolichospermum* sp. and *Nodularia* sp.). Contrary  
666 to that, there were also some reports regarding high PCY abundance in the wintertime. For instance, during the winter-  
667 spring period, picocyanobacteria was the second most dominant fraction in the Baltic Sea (Paczkowska et al., 2017). The  
668 present study showed that PCY can survive and grow also in low T and PAR conditions, which is consistent to the above  
669 cited field research of Paczkowska et al. (2017).

670 The studies of autecology of the PCY community and an understanding of its response to main environmental factors  
671 could be an important step in recognizing the phenomenon of PCY blooms in marine environments. Additionally, the  
672 laboratory experiments became a foundation in developing a new approach to Baltic Sea phytoplankton modeling -  
673 development of pico-bioalgorithm describing PCY growth, which will enable long-term numerical studies on the response of  
674 PCY to changes in environmental conditions.

## 676 5 Conclusions

677  
678 This study showed both differences and similarities in three strains of Baltic *Synechococcus* sp. ecophysiology.  
679 Discrepancies in number of cells, pigmentation changes, Chl *a* fluorescence and photosynthesis characteristics implied that  
680 BA-120, BA-124 and BA-132 should be studied and examined separately. Nonetheless, there were also fixed features  
681 similar for all analyzed strains, offering a reason to associate these features with *Synechococcus* sp. as a species, in general.  
682 For instance, according to the derived results, PAR and T played a key role in the life cycle of all three strains. Additionally,  
683 the positive impact of salinity on the number of cells was observed in each culture. Another similarity was the prevalence of  
684 the one mechanism of photoacclimation, which was the change in size of PSU. This second most frequent type was altering  
685 of accessory pigments and the least frequent was the change in PSU number.

686 Contrary to that, the main differences were: different responses of number of cells to different environmental conditions  
687 in different cultures; various photoacclimation mechanisms observed; and changes in pigmentation. According to the latest  
688 research, PCY are a great contributor to total primary production in the Baltic Sea and may contribute to summer  
689 cyanobacteria bloom at a high degree. This explains the authors' motivation to lead an in-depth investigation on Baltic PCY  
690 response to a changing environment. The present research is a first step on the way to deriving new knowledge on  
691 *Synechococcus* sp. ecophysiology and is a foundation for further studies.

## 693 Acknowledgments

694  
695 The authors would like to thank the Reviewers and Editor for their valuable comments and suggestions to improve the  
696 quality of the paper. The authors would like to thank Simon Bretherton for English language support and Proof Reading  
697 Service company for professionally proofread. The authors gratefully thank Jakub Maculewicz (IO UG), for his excellent

698 and professional technical assistance. The author SSW was financially supported by BMN grants, Poland, no. 538-G245-  
699 B568-17. This work has been funded by the Polish National Science Centre project (contract number:  
700 2012/07/N/ST10/03485) entitled: “Improved understanding of phytoplankton blooms in the Baltic Sea based on numerical  
701 models and existing data sets”. The author (AC) received funding from Polish National Science Centre in a doctoral  
702 scholarship program (contract number: 2016/20/T/ST10/00214). AC contribution was also supported by the statutory  
703 funding of IO PAS.

## 704 705 **References**

- 706  
707 Agawin, N. S., Duarte, C. M., and Agustí, S.: Nutrient and temperature control of the contribution of picoplankton to  
708 phytoplankton biomass and production, *Limn. Oceanogr.*, 45(3), 591–600, <https://doi.org/10.4319/lo.2000.45.3.0591>,  
709 2000.
- 710 Antia, N. J.: Effects of temperature on the darkness survival of marine microplanktonic algae, *Microb. Ecol.*, 3, 41–54, 1976.
- 711 Barreiro Felpeto, A., Śliwińska-Wilczewska, S., Złoch, I., and Vasconcelos, V.: Light-dependent cytotoxicity in the allelopathic  
712 interaction between picoplanktic and filamentous cyanobacteria, *J. Plankton Res.*,  
713 <https://doi.org/10.1093/plankt/fby004>, 2018.
- 714 Beardall, J.: *Blooms of Synechococcus: An analysis of the problem worldwide and possible causative factors in relation to*  
715 *nuisance blooms in the Gippsland Lakes; Monash University: Clayton, VIC, Australia, 2008; pp. 1–8, 2008.*
- 716 Becker, S., Singh, A. K., Postius, C., Böger, P., and Ernst, A.: Genetic diversity and distribution of periphytic *Synechococcus*  
717 spp. in biofilms and picoplankton of Lake Constance, *FEMS Microbiol. Ecol.*, 49, 181–190, 2004.
- 718 Belkin, I. M.: Rapid warming of large marine ecosystems, *Prog Oceanogr.*, 81 (1-4), 207–213,  
719 <https://doi.org/10.1016/j.pocean.2009.04.011>, 2009.
- 720 Cai, Y., and Kong, F.: Diversity and dynamics of picocyanobacteria and bloom-forming cyanobacteria in a large shallow  
721 eutrophic lake (lake Chaohu, China), *J. Limnol.*, 72(3), 473–484, doi:10.4081/jlimnol.2013.e38, 2013.
- 722 Callieri, C.: Picophytoplankton in freshwater ecosystems: The importance of small-sized phototrophs, *Freshw. Rev.*, 1, 1–28,  
723 <https://doi.org/10.1608/FRJ-1.1.1>, 2007.
- 724 Callieri, C.: Single cells and microcolonies of freshwater picocyanobacteria: A common ecology, *J. Limnol.*, 69, 257–277,  
725 <https://doi.org/10.4081/jlimnol.2010.257>, 2010.
- 726 Callieri, C., and Stockner, J. G.: Freshwater autotrophic picoplankton: A review, *J. Limnol.*, 61, 1–14,  
727 <https://doi.org/10.4081/jlimnol.2002.1>, 2002.
- 728 Campbell, D., Hurry, V., Clarke, A. K., Gustafsson, P., and Öquist, G.: Chlorophyll fluorescence analysis of cyanobacterial  
729 photosynthesis and acclimation, *Microbiol. Mol. Biol. Rev.*, 62(3), 667–683, 1998.
- 730 Dutkiewicz, S., Morris, J. J., Follows, M. J., Scott, J., Levitan, O., Dyhrman, S. T., and Berman-Frank, I.: Impact of ocean  
731 acidification on the structure of future phytoplankton communities. *Nat. Clim. Change.*, 5(11), 1002–1006,  
732 <https://doi.org/10.1038/nclimate2722>, 2015.
- 733 Fahnenstiel, G. L., Carrick, H. J., Rogers, C. E., and Sicko-Goad, L.: Red fluorescing phototrophic picoplankton in the  
734 Laurentian Great Lakes: What are they and what are they doing?, *Int. Rev. Ges. Hydrobiol.*, 76(4), 603–616,  
735 <https://doi.org/10.1002/iroh.19910760411>, 1991.
- 736 Feistel, R., Feistel, S., Nausch, G., Szaron, J., Lysiak-Pastuszek, E., and Ærtebjerg, G.: *BALTIC: Monthly time series 1900–*  
737 *2005*, edited by: Feistel, R., Nausch, G., and Wasmund, N., *State and Evolution of the Baltic Sea, 1952–2005, A*  
738 *Detailed 50-Year Survey of Meteorology and Climate, Physics, Chemistry, Biology, and Marine Environment*, John  
739 *Wiley & Sons, Inc., Hoboken, 311–336, 2008.*
- 740 Feistel, R., Weinreb, S., Wolf, H., Seitz, S., Spitzer, P., Adel, B., Nausch, G., Schneider, B., and Wright, D. G.: Density  
741 and absolute salinity of the Baltic Sea 2006–2009, *Ocean Sci.*, 6, 3–24, [www.ocean-sci.net/6/3/2010/](http://www.ocean-sci.net/6/3/2010/), 2010.

742 Flombaum, P., Gallegos, J. L., Gordillo, R. A., Rincón, J., Zabala, L. L., Jiao, N., Karl, D. M., Li, W. K. W., Lomas, M. W.,  
743 Veneziano, D., Vera, C. S., Vrugt J. A., and Martiny A. C.: Present and future global distributions of the marine  
744 Cyanobacteria *Prochlorococcus* and *Synechococcus*, Proc. Natl. Acad. Sci., 110(24), 9824–9829,  
745 <https://doi.org/10.1073/pnas.1307701110>, 2013.

746 Fogg, G. E., and Thake, B. (Eds.): Algal Cultures and Phytoplankton Ecology, University of Wisconsin Press, Madison and  
747 Milwaukee, 1987.

748 Glover, H. E.: The physiology and ecology of marine Cyanobacteria, *Synechococcus* spp., in: Advances in Aquatic  
749 Microbiology, Vol. 3, Jannasch, H. W., and Williams Leb, P. J., (Eds.), New York, Academic Press, 49–107, 1985.

750 Glover, H. E., Phinney, D. A., and Yentsch, C. S.: Photosynthetic characteristics of picoplankton compared with those of  
751 larger phytoplankton populations, in various water masses in the Gulf of Maine, Biol. Oceanogr., 3, 223–248, 1985.

752 Glover, H. E., Campbell, L., and Prézelin, B. B.: Contribution of *Synechococcus* spp. to size-fraction primary productivity in  
753 three waters masses in the Northwest Atlantic Ocean, Mar. Biol., 91, 193–203, 1986.

754 Guillard, R. R. L.: Culture of phytoplankton for feeding marine invertebrates, in: Culture of Marine Invertebrate Animals,  
755 Smith, W. L., and Chanley, M. H. (Eds.), Plenum Press, New York, USA, 26–60, 1975.

756 Hajdu, S., Högländer, H., and Larsson, U.: Phytoplankton vertical distributions and composition in Baltic Sea cyanobacterial  
757 blooms, Harmful Algae, 6(2), 189–205, <https://doi.org/10.1016/j.hal.2006.07.006>, 2007.

758 Hauschild, C. A., McMurter, H. J. G., and Pick, F. R.: Effect of spectral quality on growth and pigmentation of  
759 picocyanobacteria, J. Phycol. 27, 698–702, <https://doi.org/10.1111/j.0022-3646.1991.00698.x>, 1991.

760 Haverkamp, T., Acinas, S. G., Doeleman, M., Stomp, M., Huisman, J., and Stal, L. J.: Diversity and phylogeny of Baltic Sea  
761 picocyanobacteria inferred from their ITS and phycobiliprotein operons, Environ. Microbiol. 10(1), 174–188,  
762 <https://doi.org/10.1111/j.1462-2920.2007.01442.x>, 2008.

763 Haverkamp, T. H., Schouten, D., Doeleman, M., Wollenzien, U., Huisman, J., and Stal, L. J.: Colorful microdiversity of  
764 *Synechococcus* strains (picocyanobacteria) isolated from the Baltic Sea, The ISME Journal, 3(4), 397–408, 2009.

765 Herdman, M., Castenholz, R. W., Iteman, I., Waterbury, J. B., and Rippka, R.: The Archaea and the deeply branching and  
766 phototrophic bacteria, in: Boone, D. R., Castenholz, R. W. (Eds.), Bergey's Manual of Systematic Bacteriology, 2nd  
767 edn. Springer Verlag: Heidelberg, 493–514, 2001.

768 Ibelings, B. W.: Changes in photosynthesis in response to combined irradiance and temperature stress in cyanobacterial  
769 surface waterblooms, J. Phycol., 32, 549–557, <https://doi.org/10.1111/j.0022-3646.1996.00549.x>, 1996.

770 Jakubowska, N., and Szelaǳ-Wasilewska, E.: Toxic Picoplanktonic Cyanobacteria – Review, Mar. Drugs., 13, 1497–1518,  
771 <https://doi.org/10.3390/md13031497>, 2015.

772 Jassby, A. D., and Platt, T.: Mathematical formulation of the relationship between photosynthesis and light for  
773 phytoplankton, Limnol. Oceanogr., 21, 540–547, <https://doi.org/10.4319/lo.1976.21.4.0540>, 1976.

774 Jasser, I.: The relationship between autotrophic picoplankton (APP) – The smallest autotrophic component of food web and  
775 the trophic status and depth of lakes, Ecohydrol. and Hydrobiol., 6(1-4), 69–77, [https://doi.org/10.1016/S1642-3593\(06\)70128-8](https://doi.org/10.1016/S1642-3593(06)70128-8), 2006.

777 Jasser, I., and Arvola, L.: Potential effects of abiotic factors on the abundance of autotrophic picoplankton in four boreal  
778 lakes, J. Plankton Res., 25(8), 873–883, <https://doi.org/10.1093/plankt/25.8.873>, 2003.

779 Jasser, I., and Callieri, C.: Picocyanobacteria: The smallest cell-size cyanobacteria, in: Handbook on Cyanobacterial  
780 Monitoring and Cyanotoxin Analysis, Meriluoto, J., Spoof, L., and Codd G. A. (Eds.), John Wiley & Sons, Ltd,  
781 Chichester, UK, 19–27, <https://doi.org/10.1002/9781119068761.ch3>, 2017.

782 Jodłowska, S., and Śliwińska, S.: Effects of light intensity and temperature on the photosynthetic irradiance response curves  
783 and chlorophyll fluorescence in three picocyanobacterial strains of *Synechococcus*, Photosynthetica, 52(2), 223–232,  
784 <https://doi.org/10.1007/s11099-014-0024-y>, 2014.

785 Johnson, P. W., and Sieburth, J. M.: Chroococcoid cyanobacteria in the sea: A ubiquitous and diverse phototrophic biomass,  
786 *Limnol. Oceanogr.*, 24(5), 928–935, <https://doi.org/10.4319/lo.1979.24.5.0928>, 1979.

787 Joint, I. R., and Pomroy, A. J.: Photosynthetic characteristics of nanoplankton and picoplankton from the surface mixed  
788 layer, *Mar. Biol.*, 92, 465–474, 1986.

789 Kana, T. M., and Glibert, P. M.: Effect of irradiances up to 2000  $\mu\text{mol E m}^{-2} \text{ s}^{-1}$  on marine *Synechococcus* WH7803-I.  
790 Growth, pigmentation, and cell composition, *Deep-Sea Res.*, 34(4), 479–495, [https://doi.org/10.1016/0198-](https://doi.org/10.1016/0198-0149(87)90001-X)  
791 [0149\(87\)90001-X](https://doi.org/10.1016/0198-0149(87)90001-X), 1987a.

792 Kana, T. M., and Glibert, P. M.: Effect of irradiances up to 2000  $\mu\text{mol E m}^{-2} \text{ s}^{-1}$  on marine *Synechococcus* WH7803-II.  
793 Photosynthetic responses and mechanisms, *Deep-Sea Res.*, 34(4), 497–516, [https://doi.org/10.1016/0198-](https://doi.org/10.1016/0198-0149(87)90002-1)  
794 [0149\(87\)90002-1](https://doi.org/10.1016/0198-0149(87)90002-1), 1987b.

795 Kuosa, H.: Occurrence of autotrophic picoplankton along an open sea-inner archipelago gradient in the Gulf of Finland,  
796 Baltic Sea, *Ophelia*, 28, 85–93, 1988.

797 Latała, A., Jodłowska, S., and Pniewski, F.: Culture collection of Baltic Algae (CCBA) and characteristic of some strains by  
798 factorial experiment approach, *Algol. Stud.*, 122, 137–154, <https://doi.org/10.1127/1864-1318/2006/0122-0137>, 2006.

799 Larsson, J., Celepli, N., Ininbergs, K., Dupont, C.L., Yooseph, S., Bergman, B., and Ekman, M.: Picocyanobacteria  
800 containing a novel pigment gene cluster dominate the brackish water Baltic Sea, *The ISME Journal*, 8, 1892–1903,  
801 <https://doi.org/10.1038/ismej.2014.35>, 2014.

802 Leppäranta M., and Myrberg K., *Physical Oceanography of the Baltic Sea*, Springer, Berlin, pp. 378, 2009.

803 Marie, D., Simon, N., and Vaulot, D.: Phytoplankton cell counting by flow cytometry, *Algal Culturing Techniques*, 1., 253–  
804 267, 2005.

805 Mazur-Marzec, H., Sutryk, K., Kobos, J., Hebel, A., Hohlfeld, N., Błaszczuk, A., Toruńska, A., Kaczkowska, M.J., Łysiak-  
806 Pastuszek, E., Kraśniewski, W., and Jasser, I.: Occurrence of cyanobacteria and cyanotoxins in the Southern Baltic  
807 Proper. Filamentous cyanobacteria vs. single-celled picocyanobacteria, *Hydrobiologia*, 701, 235–252,  
808 <https://doi.org/10.1007/s10750-012-1278-7>, 2013.

809 Meier, H. E.: Regional ocean climate simulations with a 3D ice-ocean model for the Baltic Sea. Part 2: Results for sea ice,  
810 *Clim Dyn.*, 19, 255–266, 2002.

811 Millie, D. F., Ingram, D. A., and Dionigi, C. P.: Pigment and photosynthetic responses of *Oscillatoria agardhii*  
812 (Cyanophyta) to photon flux density and spectral quality, *J. Phycol.*, 26, 660–666, [https://doi.org/10.1111/j.0022-](https://doi.org/10.1111/j.0022-3646.1990.00660.x)  
813 [3646.1990.00660.x](https://doi.org/10.1111/j.0022-3646.1990.00660.x), 1990.

814 Motwani, N. H., and Gorokhova, E.: Mesozooplankton grazing on picocyanobacteria in the Baltic Sea as inferred from  
815 molecular diet analysis, *PLoS One*, 8(11), e79230, <https://doi.org/10.1371/journal.pone.0079230>, 2013.

816 Neumann, T.: Climate-change effects on the Baltic Sea ecosystem: A model study, *J. Marine Syst.*, 81(3), 213–224.  
817 <https://doi.org/10.1016/j.jmarsys.2009.12.001>, 2010.

818 Paczkowska J., Rowe O., Schlüter L., Legrand C., Karlson B., and Andersson A.: Allochthonous matter: An important factor  
819 shaping the phytoplankton community in the Baltic Sea, *J. Plankton Res.*, 39(1), 23–34,  
820 <https://doi.org/10.1093/plankt/fbw081>, 2017.


821 Pniewski, F. F., Biskup, P., Bubak, I., Richard, P., Latała, A., and Blanchard, G.: Photo-regulation in microphytobenthos  
822 from intertidal mudflats and non-tidal coastal shallows, *Estuar. Coast. Shelf S.* 152, 153–161,  
823 <https://doi.org/10.1016/j.ecss.2014.11.022>, 2015.

824 Prezelin, B. B.: Light reactions in photosynthesis, in: *Physiological Bases of phytoplankton Ecology*, Platt, T., (Ed.), Ottawa,  
825 *Canadian Bulletin of Fisheries and Aquatic Sciences*, no. 210, 1–46, 1981.

826 Prezelin, B. B., and Sweeney, B. M.: Photoadaptation of photosynthesis in two bloom-forming dinoflagellates, in: *Toxic*  
827 *Dinoflagellate Blooms*, Taylor, D., Seliger, H., (Eds.), Elsevier North Holland, Inc., 101–106, 1979.


- 828 Ramus, J.: The capture and transduction of light energy, in: *The Biology of Seaweeds*, Lobban, C. S., Wynne, M. J., (Eds.).  
829 Botanical Monographs, vo. 17, Oxford, Blackwell Scientific Publications, 458–492, 1981.
- 830 Richardson, K., Beardall, J., and Raven, J. A.: Adaptation of unicellular algae to irradiance: An analysis of strategies, *New*  
831 *Phytol.*, 93, 157–191, <https://doi.org/10.1111/j.1469-8137.1983.tb03422.x>, 1983.
- 832 Richardson, T. L., and Jackson, G. A.: Small phytoplankton and carbon export from the surface ocean, *Science*, 315, 838–  
833 840, <https://doi.org/10.1126/science.1133471>, 2007.
- 834 Sakshaug, E., Bricaud, A., Dandonneau, Y., Falkowski, P. G., Kiefer, D. A., Legendre, L. L., Morel, A., Parslow, J., and  
835 Takahashi, M.: Parameters of photosynthesis: Definitions, theory and interpretation of results, *J. Plankton Res.*, 19,  
836 1637–1670, <https://doi.org/10.1093/plankt/19.11.1637>, 1997.
- 837 Sánchez-Baracaldo, P., Handley, B. A., and Hayes, P. K.: Picocyanobacterial community structure of freshwater lakes and  
838 the Baltic Sea revealed by phylogenetic analyses and clade-specific quantitative PCR, *Microbiol.*, 154(11), 3347–  
839 3357, <https://doi.org/10.1099/mic.0.2008/019836-0>, 2008.
- 840 Sheskin D. J.: *Handbook of Parametric and Nonparametric Statistical Procedures: Third Edition*, CRC Press Company,  
841 London and New York, 867–980, 2000.
- 842 Sieburth J. M. N., Smatacek V., and Lenz J.: Pelagic ecosystem structure: Heterotrophic compartments of the plankton and  
843 their relationship to plankton size fractions, *Limnol Oceanogr.*, 23, 1256–126,  
844 <https://doi.org/10.4319/lo.1978.23.6.1256>, 1978.
- 845 Siefermann-Harms, D.: The light-harvesting and protective functions of carotenoids in photosynthetic membranes, *Physiol.*  
846 *Plant.*, 69, 561–568, <https://doi.org/10.1111/j.1399-3054.1987.tb09240.x>, 1987.
- 847 Siegel, H., and Gerth M.: *Sea surface temperature in the Baltic Sea in 2016, HELCOM Baltic Sea Environment Fact Sheets*  
848 *2017*, Online [Date Viewed: March 15, 2018], <http://www.helcom.fi/baltic-sea-trends/environment-fact-sheets/>.
- 849 Six, C., Finkel, Z. V., Irwin, A. J., and Campbell, D. A.: Light variability illuminates niche-partitioning among marine  
850 picocyanobacteria, *PLoS One* 2(12), e1341, <https://doi.org/10.1371/journal.pone.0001341>, 2007a.
- 851 Six, C., Thomas, J. C., Garczarek, L., Ostrowski, M., Dufresne, A., Blot, N., Scanlan, D. J., and Partensky, F.: Diversity and  
852 evolution of phycobilisomes in marine *Synechococcus* spp.: a comparative genomics study, *Genome Biol.*, 8(12),  
853 R259, <https://doi.org/10.1186/gb-2007-8-12-r259>, 2007b.
- 854 Śliwińska-Wilczewska, S., Maculewicz, J., Barreiro Felpeto, A., Vasconcelos, V., and Latała, A.: Allelopathic activity of the  
855 picocyanobacterium *Synechococcus* sp. on filamentous cyanobacteria, *J. Exp. Mar. Biol. Ecol.*, 496, 16–21,  
856 <https://doi.org/10.1016/j.jembe.2017.07.008>, 2017.
- 857 Śliwińska-Wilczewska, S., Maculewicz, J., Barreiro Felpeto, A., and Latała, A.: Allelopathic and bloom-forming  
858 picocyanobacteria in a changing world, *Toxins*, 10, 48; <https://doi.org/10.3390/toxins10010048>, 2018a.
- 859 Śliwińska-Wilczewska, S., Barreiro Felpeto, A., Maculewicz, J., Sobczyk, A., Vasconcelos, V., and Latała A.: Allelopathic  
860 activity of the picocyanobacterium *Synechococcus* sp. on unicellular eukaryote planktonic microalgae, *Mar.*  
861 *Freshwater Res.*, (in press), 2018b.
- 862 Snoeijjs-Leijonmalm, P., and Andrén, E.: Why is the Baltic Sea so special to live in?, in: *Biological Oceanography of the*  
863 *Baltic Sea*, Snoeijjs-Leijonmalm, P., Schubert, H., and Radziejewska, T. (Eds.), Springer, Dordrecht, 23–84, 2017.
- 864 Sorokin, P. Y., Sorokin, Y. I., Boscolo, R., and Giovanardi, O.: Bloom of picocyanobacteria in the Venice lagoon during  
865 summer–autumn 2001: ecological sequences, *Hydrobiologia*, 523(1-3), 71–85, 2004.
- 866 Sorokin, Y. I., and Zakuskina, O. Y.: Features of the Comacchio ecosystem transformed during persistent bloom of  
867 picocyanobacteria, *J. Oceanogr.*, 66, 373–387, 2010.
- 868 Stal, L. J., Albertano, P., Bergman, B., Bröckel, K., Gallon, J. R., Hayes, P. K., Sivonen, K., and Walsby,  
869 A. E.: BASIC: Baltic Sea cyanobacteria. An investigation of the structure and dynamics of water blooms of  
870 cyanobacteria in the Baltic Sea - Responses to a changing environment, *Cont. Shelf Res.*, 23, 1695–1714,  
871 <https://doi.org/10.1016/j.csr.2003.06.001>, 2003.

- 872 Stal, L. J., Staal, M., and Villbrandt, M.: Nutrient control of cyanobacterial blooms in the Baltic Sea, *Aquat. Microb. Ecol.*,  
873 18, 165–173, 1999.
- 874 Stal, L. J., and Walsby, A. E.: Photosynthesis and nitrogen fixation in a cyanobacterial bloom in the Baltic Sea, *Eur. J.*  
875 *Phycol.*, 35, 97–108, <https://doi.org/10.1080/09670260010001735681>, 2000.
- 876 Stawiarski, B., Buitenhuis, E. T., and Le Quèrè, C.: The physiological response of picophytoplankton to temperature and its  
877 model representation, *Front. Mar. Sci.*, 3, 164, <https://doi.org/10.3389/fmars.2016.00164>, 2016.
- 878 Stockner, J. G.: Phototrophic picoplankton: An overview from marine and freshwater ecosystems, *Limnol. Oceanogr.*, 33,  
879 765–775, <https://doi.org/10.4319/lo.1988.33.4part2.0765>, 1988.
- 880 Stomp, M., Huisman, J., Vörös, L., Pick, F. R., Laamanen, M., Haverkamp, T., and Stal, L. J.: Colourful coexistence of red  
881 and green picocyanobacteria in lakes and seas, *Ecol. Lett.*, 10, 290–298, [https://doi.org/10.1111/j.1461-](https://doi.org/10.1111/j.1461-0248.2007.01026.x)  
882 [0248.2007.01026.x](https://doi.org/10.1111/j.1461-0248.2007.01026.x), 2007.
- 883 Strickland, I. D. H., and Parsons T. R.: A practical handbook of seawater analysis, *J. Fish Res. Board Can.*, 167, 1–310,  
884 1972.
- 885 Vörös, L., Gulyas, P., and Nemeth, J.: Occurrence, dynamics and production of picoplankton in Hungarian shallow lakes,  
886 *Int. Rev. Ges. Hydrobiol.*, 76, 617–629, <https://doi.org/10.1002/iroh.19910760412>, 1991.
- 887 Waterbury, J. B., Watson, S. W., Guillard, R. R., and Brand, L. E.: Widespread occurrence of a unicellular, marine,  
888 planktonic, cyanobacterium, *Nature*, 277(5694), 293–294, <https://doi.org/10.1038/277293a0>, 1979.
- 889 Worden, A. Z., and Wilken, S.: A plankton bloom shifts as the ocean warms, *Science*, 354(6310), 287–288,  
890 <https://doi.org/10.1126/science.aaj1751>, 2016.
- 891  
892




893  
894  
895  
896

**Figure 1.** Cell number ( $10^6 \text{ cell mL}^{-1}$ ) for three *Synechococcus* sp. strains: BA-120 (A), BA-124 (B) and BA-132 (C) under different PAR and temperature conditions in 4 salinity mediums: 3 PSU (a), 8 PSU (b), 13 PSU (c) and 18 PSU (d).




898  
899  
900  
901

**Figure 2.** Cell-specific Chl *a* ( $\text{pg cell}^{-1}$ ) changes for three *Synechococcus* sp. strains: BA-120 (A), BA-124 (B) and BA-132 (C) under different PAR and temperature conditions in 4 salinity media : 3 PSU (a), 8 PSU (b), 13 PSU (c) and 18 PSU (d).



902  
 903  
 904  
 905  
 906  
 907

**Figure 3.** Cell-specific Car ( $\text{pg cell}^{-1}$ ) changes for three *Synechococcus* sp. strains: BA-120 (A), BA-124 (B) and BA-132 (C) under different PAR and temperature conditions in 4 salinity media: 3 PSU (a), 8 PSU (b), 13 PSU (c) and 18 PSU (d).




908

909

910


911

**Figure 4.** The maximum photochemical efficiency of PSII in the dark-adapted state ( $F_v/F_m$ ) for three *Synechococcus* sp. strains: BA-120 (A), BA-124 (B) and BA-132 (C) under different PAR and temperature conditions in 4 salinity mediums: 3 PSU (a), 8 PSU (b), 13 PSU (c) and 18 PSU (d).



912  
 913  
 914  
 915  
 916  
 917  
 918  
 919

**Figure 5.** The photochemical efficiency of PSII under actinic light intensity ( $\Phi_{PSII}$ ) for three *Synechococcus* sp. strains: BA-120 (A), BA-124 (B) and BA-132 (C) under different PAR and temperature conditions in 4 salinity media: 3 PSU (a), 8 PSU (b), 13 PSU (c) and 18 PSU (d).



920  
 921 **Figure 6.** Selected Chl *a* - specific and cell-specific (right side and left side panel, respectively) net photosynthetic–light  
 922 response curves for three *Synechococcus* sp. strains: BA-120 (a, b), BA-124 (c, d) and BA-132 (e, f) strains. Curves present  
 923 examples of three types of photoacclimation observed for *Synechococcus* sp. and these are as follows: change in number of  
 924 photosynthesis units (PSU) (a, b), change in size of PSU (c, d) and change in accessory pigments activity (e, f).  
 925  
 926  
 927  
 928  
 929  
 930  
 931  
 932  
 933  
 934  
 935  
 936

937 **Table 1.** Photoacclimation types (mechanisms) for three *Synechococcus* sp. strains: BA-120, BA-124 and BA-132 at  
 938 different ecological conditions. OTHER states for altering of accessory pigments activity or changes in enzymatic reactions;  
 939 PSUsizes states for the change in PSU sizes; PSUno. states for the change in PSU number. The symbols of labels indicate the  
 940 strain for which the mechanism is observed and are as follows: <sup>red</sup> for BA-120, <sup>green</sup> for BA-124 and <sup>brown</sup> for BA-132.

941

CONDITIONS	Salinity 3 PSU	Salinity 8 PSU	Salinity 13 PSU	Salinity 18 PSU
10°C	PSUsizes <sup>brown</sup>	OTHER <sup>red</sup>	PSUsizes <sup>red</sup>	OTHER <sup>red</sup>
		PSUsizes <sup>green</sup>	OTHER <sup>red</sup>	PSUsizes <sup>green</sup>
		PSUsizes <sup>brown</sup>	OTHER <sup>green</sup>	PSUsizes <sup>brown</sup>
15°C	-	PSUsizes <sup>green</sup>	OTHER <sup>red</sup>	PSUsizes <sup>brown</sup>
			PSUsizes <sup>green</sup>	
20°C	-	PSUsizes <sup>red</sup>	PSUsizes (or	PSUsizes <sup>green</sup>
		OTHER <sup>green</sup>	OTHER) <sup>red</sup>	
		PSUsizes	OTHER <sup>green</sup>	
25°C	OTHER <sup>red</sup> PSUsizes <sup>brown</sup>	PSUsizes <sup>red</sup>	PSUsizes <sup>red</sup>	PSUsizes <sup>green</sup>
		PSUsizes <sup>green</sup>	PSUsizes <sup>green</sup>	PSUsizes <sup>brown</sup>
		OTHER <sup>brown</sup>	PSUsizes <sup>brown</sup>	

942