

## ***Interactive comment on “Seasonal and spatial variability of methane emissions from a subtropical reservoir in Eastern China” by Le Yang et al.***

**Le Yang et al.**

yangboshi@live.cn

Received and published: 15 July 2018

The paper presents measurements of CH<sub>4</sub> fluxes between water and atmosphere at a large Chinese reservoir. Monthly measurements at several sites (including the river above and below the reservoir) were performed using floating chambers and bubble traps. The title is somewhat misleading because the main point of the paper is the comparison between the emissions from the (upstream and downstream) rivers with emissions from the reservoir itself. This is comparable to the study about CO<sub>2</sub> emissions done by Halbedel and Koschorreck (Biogeosciences, 2013, 10 (11), 7539-7551). I recommend to use the comparison of river and reservoir emissions as storyline in this

[Printer-friendly version](#)

[Discussion paper](#)



paper. Hypotheses should be developed along this storyline. Response: According to the reviewer's suggestions, the title of manuscript is changed into "Contrasting methane emissions from upstream and downstream rivers of an associated subtropical reservoir in eastern China" (lines 1-3). we referred the literature of Halbedel and Koschorreck (2013), the comparison of CH<sub>4</sub> emissions from (upstream and downstream) rivers and the associated reservoir was used as storyline in the revised manuscript, and the related hypotheses also proposed in lines 98-100.

Is the message of the paper that the river emits more CH<sub>4</sub> than the reservoir? That would mean that constructing a reservoir has the potential to reduce CH<sub>4</sub> emissions. The explanation of this somewhat surprising conclusion by the authors is that the deep oxalic waterbody slows down emissions by offering more options for methane oxidation. Response: According to our data, the upstream and downstream rivers emit much more CH<sub>4</sub> than the reservoir surfaces (lines 19-21), so reservoir construction indeed has the potential to reduce CH<sub>4</sub> emission if compared with CH<sub>4</sub> emission from the original lotic river (lines 22-24). On the other hand, constructing a reservoir flooded large quantities of soils in the watershed near the original rivers, which would transfer a CH<sub>4</sub> sink into a CH<sub>4</sub> source in the new flooded areas. The original hydrology conditions of natural river are changed after impoundment of reservoir, and the increased water depth, decreased water velocity, and thermal stratification in the large, deep reservoir like Xin'anjiang would had a potential influence to decline CH<sub>4</sub> emission compared with the upstream inflow rivers. Turning to the downstream river, the dissolved CH<sub>4</sub> in the hypolimnion before the dam tend to release to the atmosphere in the downstream river because of the differences in pressure and temperature. So the downstream river would has more CH<sub>4</sub> emission than the reservoir surfaces.

Do you have an idea were the CH<sub>4</sub> in the river comes from? Were the emissions in the streams higher because of the CH<sub>4</sub> construction or because of a higher gas transfer coefficient? Was the water released by the dam taken from the hypolimnion? Response: According to the thin boundary layer model, CH<sub>4</sub> flux (F) was determined by the gas

[Printer-friendly version](#)[Discussion paper](#)

transfer coefficient ( $k$ ) and the air-water  $\text{CH}_4$  concentration gradient ( $\Delta C = C_{\text{water}} - C_{\text{sat}}$ ), i.e.,  $F = k \times \Delta C$ . Thus, enhancement of them can increase the  $\text{CH}_4$  flux at the air-water interface. However, the dissolved  $\text{CH}_4$  concentrations in the water were not measured in our experiments, so we lack of the direct evidences of the impact of  $k$ ,  $\Delta C$  on  $F$ . The viewpoint stands from the  $\text{CH}_4$  emission at the air-water interface, actually, the diffusive  $\text{CH}_4$  flux was determined by the  $\text{CH}_4$  production in the sediments and the  $\text{CH}_4$  consumption (i.e, oxidation) during the vertical transport. The high  $\text{CH}_4$  flux in the upstream river was supposed to derived from the high rate of  $\text{CH}_4$  production in the sediments, because soil erosion and other man-made pollution enhance the organic carbon input in the upstream river (lines 361-363). The standardized Schmidt number of 600 ( $k_{600}$ ) was often reported to be positively correlated with wind speed. However, the average wind speeds in the upstream and the downstream rivers (3.39 m/s, 3.85 m/s, respectively) are close to those in NE, SE, and SW (4.43, 2.92, and 3.06 m/s respectively). So the spatial variability of  $\text{CH}_4$  flux probably had a weak relationship with wind speeds (Table 2). But water velocity, similar to wind speeds, also bring turbulent mixing on the air-water interface in a large river like Xin'anjiang River, which could see the literature (Beaulieu J, Shuster W, Rebolz J, Controls on gas transfer velocities in a large river, *Journal of Geophysical Research*, 2015). Xin'anjiang Reservoir is clam during the no or weak wind periods, because the main body of reservoir likes a huge clam lake with almost no water currents, but the (upstream and downstream) rivers had obviously water currents. The water velocity in the upstream river is high in June, while relatively low in other months, but in the downstream river, the water velocity was controlled by the dam operation, with a high water velocity during the daytime and low during the night.

Throughout the manuscript it is often not clear if a particular site is a river, a lake, or a reservoir (e.g. l.117). Lentic and lotic waterbodies, however, are different with respect to GHG emissions. This also has consequences regarding the methods. In a stream you cannot use an anchored chamber (Lorke et al. *Biogeosci.* 12 (23), 7013-7024) while in a lentic waterbody, drifting chambers are problematic because the wind drift

[Printer-friendly version](#)[Discussion paper](#)

might create artificial turbulence. Response: The particular site (Jiekou) in NW lake is located in a river. And the water velocity was fast in June, while kept a low level in other months. During our campaign, we avoid the flooded periods in June, and the chambers were tethered to a drifting boat (line ??). Although strong wind speeds create artificial turbulence, wind speeds was relatively low ( $<3 \text{ m s}^{-1}$ ) most time when the gas sampling was collected in our study.

The discussion about the reasons for the observed pattern is often speculative since important data were not measured (e.g. vertical concentration gradients in the reservoir, k600). Response: Just as the reviewer's comments, some important data was not measured in our experiments, and it is difficult to discuss the results in the discussion section. Despite some uncertainties, the mechanisms will be studied further, i.e. the effect of thermal stratification on CH<sub>4</sub> emission from reservoir surfaces.

Considering that the paper presents a rather limited dataset, it is rather long. I think the whole manuscript can be shorten considerably without lossing too much information. Response: The unimportant information was deleted from manuscript, according to the following suggestions (lines 116-120, 124-134, 177-184).

The method section lacks important information on the field procedures while at the same time somewhat trivial calculations are explained in great detail. Calculation details could be moved to the supplemental material. Response: According to the reviewer's suggestions, the important information was supplemented (lines 158-159) and three equations (i.e., Eq. S1, Eq. S2, Eq. S3) were deleted from the revised manuscript, which was moved into the supplemental material.

The language would be benefit from a check by a native English speaker. Response: The language was polished by a professional company.

Detailed remarks I.13: Remove "Seasonal variability showed that" Response: As the reviewer's suggestion, "Seasonal variability showed that" was deleted from the revised manuscript.

[Printer-friendly version](#)[Discussion paper](#)

I.15: “flat” is not the right term here. Response: “was flat” was replaced by “low” (line 15).

I.16/17: What does “interrupted by the bubbles” mean? Response: “interrupted by the bubbles” was replaced by “due to bubble activity” (line 16), because the occurrence of bubbles had a large impact on the fluctuation of CH<sub>4</sub> emission dynamics in the second half of year.

I. 22-24: Not a really new finding. Response: The original sentence was deleted and a new one instead in lines 24-26.

I.29: Are reservoirs wetlands? I.29: Replace “used to be often” by “are”. I.29: Logic of the sentence is strange. Reservoirs are not Energy. Response: To avoid unnecessary mistakes, the first sentence in the introduction was changed into “Hydropower has previously been regarded as a clean source of energy; however, this view is challenged by a growing body of research that has reported are carbon sources” (lines 31-33).

I.36: China has 98002 dams. Response: China has over 98, 000 dams (line 39).

I.40: Rules is not correct term here. Response: “changing pattern” was used here (Line 40).

I.41: “plant mediated” instead of “Plant-medium” Response: “Plant-medium” was changed into “plant-mediated” in line 46.

I.45 Reservoirs typically do not have littoral vegetation. Response: Different reservoirs had distinct distribution of littoral vegetation. In some reservoirs, emergent plant, e.g., reed, was distributed in the littoral zone, which has a large contribution to methane emission.

I.46-51 Sentence rather long. Response: The original sentence was divided into 2 short sentences (lines 47-51).

I.57-58 Why probably? Is that not stated in those papers? Response: Bubbles were

[Printer-friendly version](#)[Discussion paper](#)

not measured by the inverted funnels (bubble traps) in those three reservoirs, and only floating chambers was used to collect gas samples to measure the diffusive CH<sub>4</sub> emission, occasionally bubbles were captured in the chambers (lines 54-58).

I.61-63 What is the logic of this sentence? Response: The sentences are changed in lines 61-65, which show the consequences of river impoundment and their further impact on spatial variability of CH<sub>4</sub> emissions.

I.77-80: I do not understand what exactly you want to express? Response: The sentence is a summary one to show the environment factors have an impact on CH<sub>4</sub> emission from reservoirs, so the spatiotemporal variability should be stressed to avoid the error of estimating the CH<sub>4</sub> emission incorrectly (lines 77-80).

I.88: Should be included in a CH<sub>4</sub> budget of a hydroelectric reservoir. Response: CH<sub>4</sub> emission from the downstream river should be included in the CH<sub>4</sub> budget of a hydroelectric reservoir system. The last sentence of the paragraph was deleted in the revised manuscript, because it is implicit in the preceding text.

I.91-94: This is not really a hypothesis because it is long known that temperature has an influence on CH<sub>4</sub> emissions. Furthermore, in a deep reservoir the temperature at the bottom is rather constant over the season. The major seasonal difference is the different Corg and maybe O<sub>2</sub> supply to the sediment. Response: According to the reviewer's suggestions, the hypothesis was deleted from the revised manuscript.

I.103: Replace "dynamics of " by "monthly". I.104: "... region in 2015." Response: Accepted in the revised manuscript (lines 113-114).

I.107: Mean annual air temperature Response: The expression was revised according to the reviewer's suggestion (line 105).

I.107: "a total" Response: Deleted. Because the express was changed in lines 105-106.

I.108: How was evaporation measured? Is this the evaporation from the water surface?

[Printer-friendly version](#)[Discussion paper](#)

Response: The evaporation data was provided by a local meteorological station. I don't know how to measure evaporation.

I.109: Remove "which". Response: Deleted the word in line 107.

I.116: Do you mean "contributes" instead of "occupy"? Response: "contributes" was accepted in the revised manuscript (line 118).

I.117: "inflow" instead of "surface runoff". Response: Accepted in line 119.

I.117: Is it a river, a lake or a reservoir? Response: Actually, the NW lake is a river, because most water of reservoir derived from the NW lake.

I.113-120: this can be shortened Response: These sentences were shortened in lines 116-120.

I.125-140: This paragraph can be shorten. Can you also give the water depth at the sampling points (or the elevation of the bottom mean by morning? Give a range of hours) Response: The paragraph has been shorten (lines 124-133), and Table 1 was given to show the depths of sampling points (line 135).

I.146: Remove article before bubble. Response: The article was deleted.

I.149: Remove the last part of the sentence Response: The last part of the sentence was removed (line 144). The measurement of wind speed and temperatures was added in lines 144-146.

I.164: More details about the chamber method are required. Especially: Were the chambers fixed or drifting? 0.5 l sample volume is a lot. Were the measured data corrected for dilution with ambient air during sampling? Response: Three chambers were deployed in the water surface near the sampling points, but the chambers were tethered to a boat, which was drifted in the reservoir surface (lines 158-159). In addition, the total volume of a chamber is about 117L (line 150), and the maximal volume of a sampling bag is 0.5 L. However, only about 0.2-0.3L gas was collected into the

[Printer-friendly version](#)[Discussion paper](#)

bags once, so the total collected gas was about 1L, which was less than 1% of the total volume, so such influence could be ignored.

I.180: At what depth were the traps installed? How long were they developed? Were the bubble traps installed at all sites? Response: Water depth ranged between 5m and 25m (line 168). The traps were deployed about 24 h (line 174). The bubble traps were only deployed in the NW transect, because almost no bubble was captured by the static chambers in the other transects.

I.185. Remove “to reach”. Response: Accepted as the reviewer’s suggestion (line 177).

I.187-197: Paragraph is too long Response: Some trivial information was deleted in the paragraph (lines 177-184).

I.198-208: Despite the long description it is not fully clear to me which model was used to fit the data. Please provide the equation of the fit. How was the concentration calculated? Provide equation. What is the unit of  $dc/dt$ ? Response: CH<sub>4</sub> concentrations in the sampling bags were measured by a gas chromatograph instead of calculation, and its unit is ppm. The CH<sub>4</sub> flux (equation 1) is determined from the change of the gas concentration ( $\Delta c$ ) in the chamber headspace over time ( $\Delta t$ ). According to the suggestions by the reviewer #2, the linear regression is used to fit the measured CH<sub>4</sub> concentration data instead of para-curve model (line 187). Actually, the equation of the fit is  $d\Delta c/d\Delta t$ , i.e., the increased CH<sub>4</sub> concentration at the given time, and there is no unit for  $dc/dt$ .

I.214-217: How was the diffusive flux calculated when ebullition occurred? Equation 2. I wonder whether it makes sense to average over the transects, especially if single sites show extreme values. Response: From our data sets by the floating chambers, bubbles occurred in the NW transect. Bubbles were considered to be trapped in the chambers, when the CH<sub>4</sub> concentrations increased significantly and abruptly in the chambers. The ebullitive CH<sub>4</sub> flux in the bubble-trapped chamber was close to 2 mag-

[Printer-friendly version](#)[Discussion paper](#)



nitude higher than the diffusive CH<sub>4</sub> flux of the bubble-free chamber nearby. Thus, the diffusive flux was ignored when the bubble enter the chambers. The extreme values of CH<sub>4</sub> flux mainly caused by the bubbles, and the ebullitive CH<sub>4</sub> fluxes were calculated separately from the diffusive ones. In the NW transect, the frequency of bubble occurrence was 16.2% during our measurement periods, and the calculation of average CH<sub>4</sub> emission flux in the NW transect was calculated by the following equation (Eq. S2):  $F_{\text{average}} = 16.2\% \times F_{\text{ebullition}} + 83.8\% \times F_{\text{diffusion}}$  In the other transects with almost no bubble, the average CH<sub>4</sub> flux was calculated by the geometric mean of all the measured CH<sub>4</sub> fluxes by the 3 chambers in all sampling sites during the measurement periods.

I.224 add "... flux measured by the bubble traps)" Response: The phrase was added in lines 199-200.

Equation 3: Using equation 3 I get the unit  $\mu\text{g CH}_4 \text{ m}^{-2} \text{ h}^{-1}$ , not  $\text{mg CH}_4 \text{ m}^{-2} \text{ h}^{-1}$ . Response: The unit of F<sub>2</sub> in Eq. 2 is  $\text{mg CH}_4 \text{ m}^{-2} \text{ h}^{-1}$ , the transfer coefficient (1/1000) was added in equation 3 (line 202). However, there is no error in the data in the results, because the bubble gas samples were diluted 1000 times by pure N<sub>2</sub>, which offset the transfer coefficient (1/1000).

I.251: Add "in the flowing river". Response: The caption of Fig. 3 was changed in lines 223-224.

I.268-278: Given the large standard deviation the second digit after the decimal point is not significant and should be removed. Response: The suggestion is accepted in the results section (lines 226-233).

I.276: That is no surprise since the ebullition flux was calculated from the ebullition rate. More instructive would be to show the relation between ebullition flux and CH<sub>4</sub> concentration. Response: The Eq. 2 and Eq. S3 are as follows: (4) (5) Actually, , here C is a constant ( $7.16 \times 10^{-4}$ ), i.e., That is to say, the ebullitive flux (F<sub>2</sub>) is determined by the ebullition rate (ER) and the CH<sub>4</sub> concentrations (CCH<sub>4</sub>), so we easily found the ebulli-

[Printer-friendly version](#)[Discussion paper](#)

tive flux was positively correlated with ER and CCH<sub>4</sub> (Fig. S1a). The measured CH<sub>4</sub> concentrations in the bubble traps and its corresponding CH<sub>4</sub> fluxes were positively correlated in Fig. S1b (model:  $y=0.52x-7.20$ ,  $R^2=0.36$ ,  $p<0.001$ ,  $n=456$ ), although with low  $R^2$  compared with the regression model between CH<sub>4</sub> flux and ebullition rate ( $y=0.61x-1.10$ ,  $R^2=0.87$ ,  $p<0.001$ ,  $n=456$ ).

I.290: What is a “fluctuated upward pattern”? Response: The phase was deleted in the revised manuscript, and the expression was changed in lines 238-239.

I.302: Remove “average”. Response: The expression of the sentence was changed in lines 250-251.

I.335: This is surprising. Is it because there was no ebullition and the CH<sub>4</sub> concentration did not show a lateral gradient? In my experience the CH<sub>4</sub> flux from the open water depends strongly on turbulence. Was there a relation between flux and wind speed? Response: There was no significant difference in the lateral variability of CH<sub>4</sub> emission in the NE, SW, and SE transects after one-way ANOVA test (Fig. S3c, d, e). In the 3 transects of the main body, there was no bubble trapped in our chambers in 2015. According to the reviewer’s suggestions, CH<sub>4</sub> fluxes and wind speed was correlated at each transect, and the results is listed in Table 2 of the revised manuscript. Although the correlations are significant ( $p<0.05$ ), wind speed has a small account for the variability of CH<sub>4</sub> flux (a low value of  $R^2$ ). However, some high CH<sub>4</sub> fluxes were indeed caused by the turbulence of strong wind ( $> 10 \text{ m s}^{-1}$ ), e.g., the high CH<sub>4</sub> fluxes in SW transect on 8 February (SWP2:  $0.23 \text{ mg m}^{-2} \text{ h}^{-1}$ , SWP4:  $0.20 \text{ mg m}^{-2} \text{ h}^{-1}$ ).

I. 345-353: This section is not about seasonal variation – move it to 4.2 Response: The section was moved to 4.2 (lines 345-351).

I.350: What is the definition of the pelagic zone in a river? Response: Pelagic zone here is the deep water zone, e.g. the central zone.

I.347-348: Does that mean you have to reject hypothesis 2b? Response: From our

[Printer-friendly version](#)[Discussion paper](#)

results, bubbles have an obviously contribution to the high CH<sub>4</sub> emission from the upstream river, however, it don't mean the shallow water depth and fast water velocity are not important for CH<sub>4</sub> emission there, because many environmental factors have interaction effects. So we don't rejected the hypothesis.

I.360-361: How would the O<sub>2</sub> concentration at the surface would effect the CH<sub>4</sub> flux? I do not belief that. Response: According to reviewer's suggestion, the related content was deleted from revised manuscript. Thermal and dissolved oxygen stratification become weak in the second half of year, which probably related with the increased diffusive CH<sub>4</sub> flux, because Vertical transport of CH<sub>4</sub> in the water column is usually limited by the slow diffusion through the thermocline (or oxycline).

I.398: How would streamflow affect CH<sub>4</sub> emissions? What is the mechanism behind this dependency? Response: Streamflow at the outlet of dam had close relationship with water velocity in the downstream river, and a high water current velocity would enhance gas transfer velocity (k) at the air-water interface (lines 85-87) (Beaulieu et al., Controls on gas transfer velocities in a large river, Journal of Geophysical Research, 2012). In addition, Fearnside and Pueyo (2012) considered that downstream emissions are proportional to the streamflow after comparing the results between Tucuruí and Balbina in Brazil.

I.417: How fast was the water flow? If you used a fixed chamber, the high flux was probably an artefact because the chamber would create artificial turbulence which accelerates the flux (as shown by Lorke et al.). Response: Water flow was not measured in our experiments. The monitored transect (NW transect) is the main upstream inlet of the reservoir, with a fast water inflow in June because of the heavy rains in its upstream watershed. We collect the gas samples during the intermission of floods (24 June). The chambers were freely drifting while followed with our boat, the measured CH<sub>4</sub> flux in such conditions was more reliable than the anchored chambers (Lorke et al., 2015), although there is turbulence more or less.

[Printer-friendly version](#)[Discussion paper](#)

I.422: Vertical transport of CH<sub>4</sub> in the water column is usually limited by the slow diffusion through the thermocline. Thus, the thickness of the hypolimnion or epilimnion is not so important. Response: According to the reviewer's suggestion, the relevant content was deleted from the revised manuscript.

I.428-445: Paragraph too long Response: The paragraph was shortened in lines 366-373.

I.469: Where the emissions from the river high because of the CH<sub>4</sub> concentration or because of a high k<sub>600</sub>? Response: We lacked of direct evidences to explain the high CH<sub>4</sub> emissions from the downstream river, however, plenty of CH<sub>4</sub> emitted in the downstream river when the hypolimnion water passed through turbines in other dams (Kemenes et al., 2007; Deshmukh et al., 2016), because of the differences in temperature and pressure promote the high dissolved hypolimnion CH<sub>4</sub> concentrations release in the downstream river (lines 83-85, 401-405). In addition, the remainder of the paragraph are deleted from the revised manuscript, because CH<sub>4</sub> flux don't show the decrease trend with the distance to the dam after the amendment of CH<sub>4</sub> fluxes in Eq. 1 (Figure S3 a).

I. 474: You cannot simply transfer the results from another, completely different, reservoir. Response: According to the reviewer's suggestion, the sentence was deleted from our revised manuscript.

I.476-478: Sentence sounds strange. Response: It contains some uncertainty on the strong turbulence on CH<sub>4</sub> emission from the downstream river, so the sentence was deleted from original manuscript.

I.495-498: I thought that exactly this was the purpose of the study. Response: The sentences was moved to the abstract (lines 24-26).

Fig. 4: How can the CH<sub>4</sub> bubble concentration be higher than 100%? Response: The CH<sub>4</sub> bubble concentration values did not exceed 100%, and the y coordinate axis

[Printer-friendly version](#)[Discussion paper](#)

ranged from 0 to 400% to avoid to overlap with other data. To avoid the misunderstanding proposal by the reviewer, the range of y coordinate axis of bubble CH<sub>4</sub> concentration changes from -30% to 120% to show the complete error bars in the revised manuscript (Figure 4).

Fig. 6: Move to supplemental material. Response: As the reviewer's suggestion, the original Fig. 6 was moved into supplemental material (Figure S2).

Please also note the supplement to this comment:

<https://www.biogeosciences-discuss.net/bg-2018-195/bg-2018-195-AC1-supplement.pdf>

---

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2018-195>, 2018.


BGD

Interactive  
comment

Printer-friendly version

Discussion paper





**Fig. 1.** Monthly precipitation, evaporation, air temperature, and water level in the Xin'anjiang Reservoir in 2015.





Fig. 2. Location of transects and sampling points.




**Fig. 3.** Mean CH<sub>4</sub> emissions from the upstream river of the reservoir between December 2014 to January 2016.

Printer-friendly version


Discussion paper









**Fig. 4.** Mean ebullition rate, bubble CH<sub>4</sub> emission flux, and bubble CH<sub>4</sub> concentration recorded from the inflow river.




**Fig. 5.** Mean monthly diffusive CH<sub>4</sub> emission from the areas of the reservoir.



**Fig. 6.** Mean diffusive CH<sub>4</sub> emissions from different distances downstream of the reservoir.



**Fig. 7.** Mean CH<sub>4</sub> emissions from the reservoir and the upstream and downstream rivers.



**Fig. 8.** Discharge flow at 9:00 hrs in the downstream river below the dam during the measurement period.