

Table 3.

Species	Porosity at 10°C	Porosity at 20°C	Q_{10} Porosity	Symbiont Type
<i>Globigerinoides conglobatus</i>	-0.1757	-0.0657	2.674	Dinoflagellate ¹
<i>Neogloboquadrina dutertrei</i>	-0.1017	-0.0397	2.562	Pelagophytes ²
<i>Orbulina universa</i>	-0.195	-0.084	2.321	Dinoflagellate ¹
<i>Globigerinoides sacculifer</i>	-0.1698	-0.0858	1.979	Dinoflagellate ¹
<i>Globigerinella siphonifera</i>	-0.1625	-0.0965	1.684	Chrysophytes ¹
<i>Globigerinoides ruber</i>	-0.1104	-0.0834	1.324	Dinoflagellate ¹
<i>Globorotalia inflata</i>	-0.0628	-0.0898	0.699	Chrysophytes ¹
<i>Globorotalia truncatulinoides</i>	-0.0499	-0.0869	0.574	Asymbiotic ¹

¹ Ezard, T. H., et al. (2015). Environmental and biological controls on size-specific $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ in recent planktonic foraminifera. *Paleoceanography*, 30(3), 151-173.

² Bird et al., (2018). 16S rRNA gene metabarcoding and TEM reveals different ecological strategies within the genus *Neogloboquadrina* (planktonic foraminifer). *PloS one*, 13:1.