



1 Riverine particulate C and N generated at the permafrost thaw front:
2 case study of western Siberian rivers across a 1700-km latitudinal transect

3

4 Ivan V. KRICKOV¹, Artem G. LIM¹, Rinat M. MANASYPOV^{1,2},
5 Sergey V. LOIKO¹, Liudmila S. SHIROKOVA^{2,3}, Sergey N. KIRPOTIN¹,
6 Jan KARLSSON⁴, Oleg S. POKROVSKY^{3*}

7

8 ¹ *BIO-GEO-CLIM Laboratory, Tomsk State University, Tomsk, Russia*

9 ² *N. Laverov Federal Center for Integrated Arctic Research, Russian Academy of Sciences,*
10 *Arkhangelsk, Russia*

11 ³ *Geosciences and Environment Toulouse, UMR 5563 CNRS, 14 Avenue Edouard Belin 31400*
12 *Toulouse, France*

13 ⁴ *Climate Impacts Research Centre, Department of Ecology and Environmental Science, Umeå*
14 *University, 901 87 Umeå, Sweden*

15

16 **Email: oleg.pokrovsky@get.omp.eu*

17 *Key words: nutrient, particulate, suspended, landscape, bog, lake, forest, thaw, Siberia*

18

19 Submitted to *Biogeosciences*, May 2018

20

21

22

23

24

25

26

27

28

29



30 Abstract

31 In contrast to numerous studies on the dynamics of dissolved ($< 0.45 \mu\text{m}$) elements in
32 permafrost-affected high latitude rivers, very little is known of the behavior of river suspended
33 ($> 0.45 \mu\text{m}$) matter (RSM) in these regions. In order to test the effect of climate, permafrost and
34 physio-geographical landscape parameters (bogs, forest and lake coverage of the watershed) on
35 RSM and particulate C, N and P concentration in river water, we sampled 33 small and medium
36 size rivers (10 – 100,000 km² watershed) along a 1700 km N - S transect including both
37 permafrost-affected and permafrost-free zones of Western Siberian Lowland (WSL). The
38 concentration of C and N in RSM decreased with the increase in river watershed size, illustrating
39 *i*) the importance of organic debris in small rivers which drain peatlands and *ii*) the role of
40 mineral matter from bank abrasion in larger rivers. The presence of lakes in the watershed
41 increased C and N but decreased P concentrations in RSM. The C:N ratio in the RSM reflected
42 the source from deep rather than surface soil horizon, similar to that of other Arctic rivers. This
43 suggests the export of peat and mineral particles through suprapermafrost flow occurring at the
44 base of the active layer. There was a maximum of particulate C and N concentration at the
45 beginning of permafrost appearance (a sporadic and discontinuous zone, 62-64°N). This
46 presumably reflected the organic matter mobilization from newly thawed organic horizons in
47 soils at the active latitudinal thawing front. The results suggest that a northward shift of
48 permafrost boundaries and an increase in active layer thickness may increase particulate C and N
49 export by WSL rivers to the Arctic Ocean by a factor of 2, while P export may remain
50 unchanged. In contrast, within a long-term climate warming scenario, the disappearance of
51 permafrost in the north, the drainage of lakes and transformation of bogs to forest may decrease
52 C and N concentration in RSM by 2 to 3 times.

53

54



55 **1. Introduction**

56 High-latitude rivers are most vulnerable to the change of particulate nutrient regime due
57 to on-going climate change via altering their hydrological regime (Bring et al., 2016) and
58 widespread permafrost thaw that stimulates nutrient release (Vonk et al., 2015). For carbon (C),
59 the particulate fraction (POC) contributes substantially to the total organic C export from the
60 continent to the ocean (Schlesinger and Melack, 1981; Lal, 2003; Ludwig and Probst, 1996;
61 Galy et al., 2015; Li et al., 2017); a two-fold increase of Arctic rivers POC fluxes by 2100 is
62 predicted (Gordeev and Kravchishina, 2009). Although the reasons for strong variations of POC
63 in freshwaters are not yet fully understood (Tiang et al., 2015; Lee et al., 2015; Yang et al.,
64 2016), the temperature (Hilton, 2016) and runoff (Goni et al., 2015) combined with local storm
65 events (Jeong et al., 2012; Wiegner et al., 2009) are widely recognized as the most important
66 driving factors. This may be especially true for northern aquatic systems, being highly sensitive
67 to flood events, due to shallow water paths and short transit time in watersheds.

68 Of special interest to POC of the Arctic rivers is that, if soil organic C escapes
69 degradation during river transport and thus buried in marine sediments, it can contribute to a
70 geological carbon dioxide sink (e.g., Hilton et al., 2015). Further, potentially increased transport
71 of P and N may significantly change primary productivity in riverine (Wrona et al. 2016;
72 McClelland et al. 2007), estuarine (Emmerton et al. 2008b; McClelland et al. 2007) and ocean
73 Arctic ecosystems (Yool et al. 2015) thereby impeding rigorous predictions of climate change
74 impact on Arctic terrestrial-aquatic ecosystems.

75 Despite significant efforts in characterizing the fluxes, chemistry, and origin of
76 particulate organic matter (POM) in large Arctic Rivers (Lobbes et al., 2000; Dittmar and
77 Kattner, 2003; Unger et al., 2005; Guo et al., 2004, Guo and Macdonald, 2006; Gladyshev et al.,
78 2015; Emmerton et al., 2008a; McClelland et al., 2016; Gareis and Lesack, 2017), these studies
79 do not allow for assessment of mechanisms of POM generation in the watershed. In particular,



80 the role of size of the river watershed and its landscape (physio-geographical) parameters is still
81 poorly known. Thus, although detailed studies of particulate nutrients in small Arctic rivers
82 helped to constrain seasonal features of export fluxes (Cai et al., 2008; Dornblaser and Striegl,
83 2007; Lamoureux and Lafrenière, 2014; McClelland et al., 2014), the key environmental driving
84 factors of particulate nutrient concentration and stoichiometry in Arctic rivers—permafrost
85 coverage and lakes and forest proportion on the watershed—remain poorly resolved.

86 In this regard, large continental plains such as the western Siberia Lowland (WSL),
87 which contains sizeable reservoirs of frozen and thawed organic carbon, N, P and inorganic
88 nutrients (Sheng et al. 2004; Stepanova et al., 2015; Raudina et al., 2017), may be especially
89 useful in assessing environmental control on particulate nutrient transport to the Arctic Ocean. A
90 vast amount of frozen peat in this region can strongly affect the coastal Arctic system in the
91 event of permafrost thaw and enhanced RSM export from the watersheds. Due to the high
92 homogeneity of the WSL landscape, lithology, and topography, one can use the natural north-
93 south gradient of the permafrost zone distribution to assess the direct impact of permafrost
94 conditions on river water chemistry.

95 Detailed studies of the dissolved fraction of WSL river water demonstrated several
96 typical features occurring over a sizeable gradient of climate and permafrost. In pioneering
97 works of Frey and co-workers it was shown that southern permafrost-free regions export 3 to 4
98 times greater amounts of dissolved C, N and P (Frey and Smith, 2005; Frey et al., 2007a, b; Frey
99 and McClelland, 2009) and that wetlands exert a significant positive effect on carbon and
100 nutrient concentration in small rivers (Frey et al., 2007a; Frey and McClelland, 2009). Although
101 the majority of these features were confirmed by a more recent study of dissolved carbon and
102 nutrients in WSL rivers over main hydrological seasons (Pokrovsky et al., 2015 and Vorobyev et
103 al., 2017, respectively), an assessment of particulate load transport in WSL rivers has not yet



104 been performed and the main mechanisms controlling particulate C, N and P mobilization from
105 WSL rivers to the Arctic Ocean remain unknown.

106 To improve current understanding of magnitude and seasonality of riverine particulate
107 nutrient export, we quantified concentrations of C and macro- (N, P) nutrients across a vast
108 latitudinal gradient (1700 km) with special emphasis on permafrost-bearing zone during three
109 main hydrological regimes: 1) the peak of spring flood (early June 2016), 2) the summer base
110 flow (August 2016), and 3) the autumn high flow before the ice (October 2016). We aimed at
111 quantifying the effect of latitude, permafrost coverage and fundamental landscape features
112 (proportion of bogs, lakes and forest in the watershed) as well as the size of the river itself on
113 particulate C, N and P concentration and the relative fraction of particulate versus total
114 (particulate + dissolved) nutrient transport. We further used acquired knowledge to infer the
115 basic mechanisms of particulate nutrient mobilization from soils to rivers and applied these
116 mechanisms to prediction change in particulate nutrient concentration under climate warming,
117 landscape evolution and progressive permafrost thaw in the largest frozen peatland province in
118 the world.

119

120 **2. Study Site and Methods**

121 The rivers were sampled in the Western Siberia Lowland (WSL), a huge (> 2 million km²),
122 peatland and forest zone situated in the taiga forest, forest-tundra and tundra zone. The position
123 of biomes follows the decrease of mean annual air temperature (MAAT) from -0.5°C in the
124 south to -9.5°C in the north. The permafrost distribution also follows the latitudinal gradient of
125 MAAT and changes from absent, isolated and sporadic in the south to discontinuous and
126 continuous in the north. Further details of WSL physio-geographical settings, peat and
127 lithological description of the territory are provided elsewhere (Kremenetski et al., 2003;
128 Stepanova et al., 2015; Pokrovsky et al., 2015; Raudina et al., 2017). For each biome (taiga,



129 forest-tundra and tundra) several rivers with different watershed sizes were chosen and WSL
130 river dissolved load sampling was performed along a latitudinal transect following previous
131 strategies by Pokrovsky et al. (2015, 2016) and Vorobyev et al. (2017).

132 Altogether, we sampled 33 rivers that belong to watersheds of Ob, Pur and Taz including
133 these large rivers as well (**Fig. 1**). The landscape parameters of sampled catchments were
134 determined by digitizing available soil, vegetation, lithological and geocryological maps (**Table**
135 **S1** and Vorobyev et al., 2017). There was no covariation between river size and other landscape
136 parameters including permafrost coverage. Sampling was performed during three main
137 hydrological seasons: 1) spring flood (17 May – 15 June 2016), 2) summer baseflow (1 – 29
138 August 2016), and 3) autumn baseflow before ice (24 September – 13 October 2016). Note that
139 the most interesting period—in terms of soil connection to the rivers—occurred in late autumn
140 when the active layer depth was at its maximum. This period has not been covered in previous
141 studies of dissolved WSL river load.

142 The sampling strategy consisted of moving from south to north in spring and autumn
143 over a 2-3 week period, following the natural change of seasons. This allowed us to sample all
144 rivers of the transect at approximately the same time after ice off and before ice on. The year
145 2016 was normal for western Siberia in terms of spring, summer and autumn precipitation but
146 temperature were 4 and 2.7 °C higher than normal spring and summer, respectively, and not
147 different from the average T in autumn (Rosgidromet, 2017). For assessing inter-annual
148 variations in RSM concentrations, we analyzed the RSM samples collected in WSL rivers across
149 the same transect during a previous campaign in the spring of 2014 and 2015 and the summer
150 and autumn of 2014 and 2015.

151 Large water samples were collected from the middle of the river at 0.5 m depth in pre-
152 cleaned polypropylene jars (30 to 50 L) and were allowed to decantate over 2-3 days. The water
153 of the bottom layer of the barrels (approx. 30% of the initial volume) was centrifuged on-site for



154 20 min at 3500 rpm using 50-mL Nalgene tubes; sediment was frozen at -18°C and freeze-dried
155 later in the laboratory. In addition to decantation and centrifugation, RSM was collected via
156 direct filtration of large volumes (20 to 30 L) of river water with an Inox (AISI 304) Teflon®
157 PTFE-coated filtration unit (Fisher Bioblock) equipped with 142 mm acetate cellulose Sartorius
158 membranes ($0.45\ \mu\text{m}$) and operated at 5-7 bars. An average flow rate of 1-2 L/h was created by
159 a peristaltic pump (MasterFlex B/T) with Teflon tubing. For determination of total concentration
160 of suspended material, smaller volumes of freshly collected river water (1-2 L) were filtered on-
161 site (at the river bank or in the boat) with pre-weighted acetate cellulose filters (47 mm, 0.45
162 μm) and Nalgene 250-mL polystyrene filtration units using a Mityvac® manual vacuum pump.

163 There was reasonably good agreement, typically within 10%, between the concentration
164 of RSM collected in large barrels via decantation followed by centrifugation, a direct high-
165 pressure filtration using 142-mm membranes and vacuum filtration using Nalgene 250-mL unit.
166 The agreement was below 10% for large rivers in summer and autumn when the mineral
167 component dominated the RSM. Agreement was also between 10 and 20% for small organic-
168 rich rivers containing peat and plant debris especially in spring.

169 The C and N concentration in RSM collected from large-volume separation procedure
170 was measured using catalytic combustion with Cu-O at 900°C with an uncertainty of $\leq 0.5\%$
171 using Thermo Flash 2000 CN Analyzer at Tomsk University. The samples were analyzed before
172 and after 1:1 HCl treatment to distinguish between total and inorganic C; however the ratio of
173 $C_{\text{organic}} : C_{\text{carbonate}}$ in RSM was always above 20 and the contribution of carbonate C to total C in
174 the RSM was equal in average $0.3 \pm 0.3\%$ (2 s.d., $n = 30$). In addition to RSM, we compared total
175 and HCl-treated C analysis in peat soil column (organic part and 3 separate mineral horizons)
176 sampled from the middle part of river transect. The $C_{\text{carbonate}}$ share was below 2 % of total C
177 content for both the mineral and organic part of soil columns. The analyses we performed could
178 not distinguish mineral N linked to clays (NH_4^+ cation) and organic N in the RSM. For P, the



179 RSM samples were subjected to full acid leaching in a clean room following ICP-MS (Agilent
180 7500 ce) analyses using methods for C_{org}-rich natural samples described by Stepanova et al.
181 (2015). Water samples for DOC and total dissolved phosphorus (P_{tot}) were filtered on-site
182 through 0.45 µm acetate cellulose filters (Millipore, Sartorius) and analyzed following methods
183 previously described by Pokrovsky et al. (2015, 2016).

184 A regression analysis was used to quantify the relationship between C, N and P
185 concentration in RSM and the % of permafrost, wetlands, lake and forest coverage of the
186 watershed as well as the surface area of the watershed (S_{watershed}). In order to assess a general
187 impact of the permafrost on RSM nutrient concentration we separated all sampled rivers into
188 five categories according to the permafrost distribution on their watersheds: 1) permafrost-free
189 (south of 61°N), 2) isolated (61 to 63.5°N); 3) sporadic (63.5 to 65°N); 4) discontinuous (65 to
190 66°N), and 5) continuous permafrost zones (north of 66°N). The non-parametric statistics were
191 used because, based on Shapiro-Wilk test of the normality of variables, the data on C, N, P
192 concentration in RSM and the % of element in suspended form were not normally distributed.
193 For these reasons, we used the median, 1st and 3rd quartiles to trace dependence of nutrient
194 concentration to the type of permafrost distribution. The differences in suspended C, N and P
195 concentration between different seasons and between each two adjacent permafrost zones were
196 tested using a Mann-Whitney U test for a paired data set with significance level at 0.05. For
197 unpaired data, a non-parametric H-criterion Kruskal-Wallis test was performed for all watershed
198 sizes and all permafrost zones.

199

200 3. Results

201 3.1. C, N and P concentrations in RSM and their link to seasons and watershed size

202 Mean bulk RSM concentration in the WSL river waters did not depend on the open water
203 seasons and was equal to 7.1±3.9, 8.1±4.1, and 7.0±3.7 mg/L in spring, summer and autumn,



204 respectively (**Table 1**). The RSM concentrations weakly depended on the size of the watersheds
205 ($S_{\text{watershed}}$) with a negative relationship in autumn ($R^2 = 0.33$, $p < 0.05$, Fig. **S1 A**). Further, the
206 RSM concentration increased with permafrost coverage and latitude ($R^2 = 0.56$ and 0.41),
207 although this was visible only in autumn (**Fig. S1 B, C, Table S2**). The sporadic permafrost
208 zone exhibited the highest RSM concentration in summer (**Fig. S1 D**). Finally, there was no
209 correlation ($p > 0.05$) between lake, bog or forest coverage and the RSM concentration ($R^2 <$
210 0.2 , see also **Table S2**). For RSM concentration, statistically significant difference between
211 different permafrost zones, notably between permafrost-free and permafrost-bearing regions,
212 were evidenced in summer and autumn using Kruskal-Wallis and Mann-Whitney tests (**Table**
213 **S3**).

214 The concentrations of C, N and P in WSL rivers averaged over 3 seasons were equal to
215 $15.3 \pm 9.7\%$, $1.2 \pm 0.9\%$, and $0.49 \pm 0.42\%$ in mass of RSM (1.05 ± 0.805 , 0.083 ± 0.066 , and
216 0.035 ± 0.036 mg/L in the riverwater). The watershed size sizably affected the C concentration:
217 there was a power-law decrease of C with the size of watershed ($R^2 = 0.28$, 0.47 , and 0.25 in
218 spring, summer and autumn, respectively **Fig. 2A**) but there was no relationship with the N and
219 P concentrations in RSM ($R^2 < 0.2$, **Fig. 2 B, C**). Generally, a 2 to 3-fold increase in C_{org} , from
220 ca. 20-30% in rivers with $S_{\text{watershed}} < 100 \text{ km}^2$ to $C_{\text{org}} = 5-10\%$ in rivers with $S_{\text{watershed}} > 10,000$
221 km^2 was observed. The C:N ratio of RSM was independent on the watershed size in spring but
222 decreased 2-3 times with $S_{\text{watershed}}$ increase ($R^2 = 0.4$) in summer and autumn (**Fig. 2D**).

223 Finally, the inter-annual variations of suspended nutrient concentration in WSL rivers
224 were of secondary order importance when compared to season and watershed size control. We
225 did not find any inter-annual differences (at $p < 0.05$) in RSM concentration and P concentration
226 in RSM collected in June and August in 2014, 2015, and 2016 for the same 7 rivers.

227



228 3.2. Role of permafrost distribution and landscape parameters for C, N, and P
229 concentration and fraction of particulate nutrients

230 There was a local maximum of C and N concentration in isolated and sporadic
231 permafrost zone (**Fig. 3 A, B, D, E**), which was not seen for P (**Fig. 4 C, F**). Overall, the
232 differences in C and N concentrations in RSM among different permafrost zones were
233 significant as verified by the non-parametric Kruskal-Wallis H-test ($0.005 < p < 0.05$), while the
234 difference in P concentration between permafrost zones was not significant ($p > 0.05$, see **Table**
235 **S3 C, D**). Specifically, the C demonstrated a maximum concentration (significant at $p < 0.02$
236 during all three seasons) at 62-64°N (**Fig. S2 A**). The latitude generally did not impact N and P
237 concentration in RSM (**Fig. S2 B, C**). The differences between adjacent permafrost zones were
238 evidenced by C and N in summer and autumn (**Table S3 D**).

239 The landscape parameters of the watershed (bogs, lakes and forest coverage) sizably
240 affected ($p < 0.05$) suspended C and N. Bogs and lakes at the watershed increased the
241 concentration of C and N in RSM whereas forest generally decreased C in RSM (**Fig. 4 A-B-C**
242 **for C, and Fig. S3 A-B-C for N**). This increase in C and N % with bogs and lakes coverage and
243 a decrease with forest presence was mostly visible in summer and autumn. The increase in lake
244 coverage of the watershed led to a decrease in P concentration in RSM in summer and autumn
245 ($R^2 = 0.31$ and 0.22 , respectively, **Fig. S3 D-E-F**) that was especially visible in autumn in the
246 permafrost-free zone ($R = -0.88$, **Table S2**). During this period, the P concentration in RSM
247 positively correlated with the presence of forest in the permafrost zone ($R = 0.60$, **Table S2**).

248 The Mann Whitney U-test for the impact of watershed parameters demonstrated
249 significant differences in C and N concentration (all seasons) and P concentration (summer
250 baseflow) between watersheds having $< 10\%$ and $> 10\%$ lake coverage, **Table S3-E**. The
251 differences were also observed among watershed with $< 50\%$ and $> 50\%$ of bogs for C (all
252 seasons) and N (summer and autumn), **Table S3-F**. Finally, the forest coverage ($< 30\%$ and $>$



253 30%) exhibited significant effect on C and N (all seasons) and P (autumn baseflow), **Table S3-**
254 **G.**

255 The share of particulate carbon versus total (dissolved + particulate C) did not
256 demonstrate any significant dependence on $S_{\text{watershed}}$, bogs, forest and permafrost proportions on
257 the watershed ($R^2 < 0.3$, not shown). However, there was a localized maximum of particulate
258 carbon fraction around 64°N within the isolated to sporadic permafrost zone (**Fig. 5 A and C**).
259 The presence of lakes sizably increased the particulate over total transport of C in rivers ($R^2 =$
260 0.52 and 0.32 in spring and summer, respectively, **Fig. 5 B**). The P fraction in the RSM ranges
261 from 10 to 90% of its total (suspended + dissolved) amount without any link to size of river
262 watershed, % of forest and bogs, and type of permafrost distribution (not shown).

263

264

265 **4. Discussion**

266 *4.1. Concentrations of nutrients and impact of the watershed size*

267 The RSM values in WSL rivers (2 to 18 mg/L) are similar to other boreal rivers of low
268 runoff which drain peatlands such as Severnaya Dvina (2.3 to 16 mg/L; Pokrovsky et al., 2010)
269 but lower than the Ob River itself (around 30 mg/L; Gebhardt et al., 2004) and other big rivers
270 of the Kara Sea basin (average 22 mg/L; Gordeev et al., 1996). The POC values of the WSL
271 rivers (0.5 to 3.0 mg/L POC) are consistent with recent data on WSL river transects sampled in
272 2015 (Vorobyev et al., 2017) and are in agreement with those of the Ob-Taz River confluence
273 measured in June (1.3 mg/L; Gebhardt et al., 2004), the Ob River at Salekhard in May through
274 October (0.8 to 2.4 mg POC/L; Le Fouest et al., 2013), the low reaches of the Ob River (1.2 to 2.4
275 mg POC/L; McClelland et al., 2016), the mean multi-annual values of POC in subarctic rivers of
276 Northern Eurasia draining peatlands (3.2, 0.3, 0.9 mg POC/L for S. Dvina, Pechora and Ob as
277 compiled in Gordeev et al., 1996) and the Lena River basin (0.5 mg/L; Kutscher et al., 2015).



278 However, the C_{org} concentrations in RSM of WSL rivers (5 to 40%), notably in small and
279 medium size ($< 10,000\text{-}100,000 \text{ km}^2$) ones, are an order of magnitude higher than that in other
280 world rivers which drain mineral substrates (typically 1% C_{org} in RSM; Meybeck, 1993) and
281 significantly higher than the values of the Siberian rivers (2.3, 3.6, 5.8, 3.0% for Ob, Yenisey,
282 Lena and Kolyma, respectively; Gordeev and Kravchishina, 2009). Thus, typical concentration
283 of C_{org} in RSM of large ($S_{\text{watershed}} > 100,000 \text{ km}^2$) Central Siberian rivers that drain larch forest is
284 only 0.4 to 0.5 % (Pokrovsky et al., 2005). The C_{org} concentration in the RSM of Severnaya
285 Dvina River (which has sizeable proportion of bogs and lakes within its watershed compared to
286 WSL rivers) is $2.7\pm 0.7\%$ in May and $4.8\pm 1.1\%$ in August (Savenko et al., 2004). The N_{org}
287 content in RSM ranges from 0.3 to 1.8 % (0.05 to 0.2 mg particulate N_{tot}/L) which is much
288 higher than that in sedimentary rocks (0.05 to 0.06 %; Houlton et al., 2018) but is comparable
289 with the value reported for the freshwater part of Ob river estuary (0.16 mg N/L; Gebhardt et al.,
290 2004), the Ob River at Salekhard in May to October (0.1 to 0.3 mg PON/L; Le Fouest et al.,
291 2013), the Yukon River (0.14 ± 0.09 mg particulate N/L; Guo and MacDonald, 2006), and small
292 rivers of the North slope of Alaska (0.05 to 0.6 mg PON/L; McClelland et al., 2014).

293 High concentrations of C (and N) in the RSM of WSL rivers may stem from the organic
294 nature of soils that prevail on river watersheds. The Histosols, one of the dominant soil groups of
295 WSL, are capable of providing a sizeable amount of organic particles given the higher
296 susceptibility of peat to physical disintegration compared to mineral soils. The enrichment of the
297 river water in C-rich particles may occur at both the river bank (especially in small rivers
298 flowing through the wetlands) and within the extensive floodplains via remobilization of
299 organic-rich sediments during high flow periods.

300 The concentration of C and N in RSM decreased with increased river watershed size,
301 thereby illustrating the importance of organic particles in small rivers draining peatlands and the
302 role of mineral matter from bank abrasion in larger rivers. The impact of watershed size is more



303 significant for C than for N. Presumably this is because N is more affected by autochthonous
304 processes and that particulate N may partly be generated from phytoplankton and macrophytes
305 in the river. Small rivers ($S_{\text{watershed}} < 100\text{-}1000 \text{ km}^2$) exhibited the largest scatter in particulate C,
306 N (and P) concentrations. This is probably due to multiple sources of POM and the very short
307 transit time in the watershed that results in fast responses of river particulate load to minor
308 variations in surface hydrology including high sensitivity to local storm events.

309 The decrease of C:N in the RSM from small to large rivers likely reflected a shift in main
310 origin of suspended matter, from peat in small rivers to more lithogenic (deep soil) in large
311 rivers. This was mostly visible in summer and autumn; in spring the rivers exhibit a very
312 homogeneous C:N signature which may be linked to a dominant source of RSM from bank
313 abrasion and sediment transport as well as deposition within the riparian zone. In fact, the flood
314 plain of the Ob river and other rivers of the WSL extend more than 10 times the width of the
315 main channel (Vorobyev et al., 2015). Note that the C:N ratio in large rivers ($>100,000 \text{ km}^2$)
316 approach that of average sedimentary rocks (8.1; Houlton et al., 2018). In this regard, highly
317 homogeneous C:N ratios in particulate load of Arctic rivers (7 to 18 for Mackenzie, Yukon,
318 Kolyma, Lena, Yenisey and Ob regardless of season; McClelland et al., 2016) are interpreted as
319 the mixture of deep soil sources where $\text{C:N} < 10$ (Schädel et al., 2014) and upper organic-rich
320 horizons of soils with elevated C:N (Gentsch et al., 2015). The Ob River demonstrates the
321 youngest POC of all Arctic Rivers (-203 to $-220 \text{ ‰ } \Delta^{14}\text{C}$; McClelland et al., 2016) which
322 certainly indicates a relatively fresh (ca. 1,000-2,000 years old) origin of particulate carbon that
323 is presumably from intermediate peat horizons.

324 We believe that the variation in C:N in RSM may reflect different sources of organic
325 material feeding the river depending on seasons and latitudes. A compilation of C:N ratios in
326 peat and mineral horizons as well as in thermokarst lake sediments for four main sites of
327 latitudinal transect considered in this study is given in **Fig. S4 of Supplement**. The range of C:N



328 values in RSM rivers (10 to 20) is closer to that in sediments of thermokarst lakes (20 to 30).
329 Note that the resuspension of sediments may be an important source of water column POC
330 (Yang et al., 2016). The minerotrophic bogs, which are mostly linked to rivers via hydrological
331 networks, have a C:N ratio in upper peat horizons ranging from 24 to 28. In mineral soils of the
332 region, the C:N range is between 10 and 15 regardless of latitude, from the tundra situated Taz
333 River riparian zone to the taiga situated middle channel of the Ob River. For upper organic
334 horizons the C:N is always higher than the bottom mineral horizons. The old alluvial deposits of
335 the Pyakopur River (discontinuous permafrost zone) had only 0.2% of POC with C:N equal to 6.
336 Overall, there is an enrichment in N relative to C in the course of water transport of organic and
337 organo-mineral solid particles from soils and riparian deposits to the river water.

338

339

340 *4.2. A maximum of C and N in the isolated/sporadic permafrost zone and the impact of*
341 *river watershed characteristics*

342 Complementary to previous results on dissolved ($< 0.45 \mu\text{m}$) C and N concentrations in
343 WSL rivers acquired by Frey et al (2007a) and Vorobyev et al. (2017) that demonstrated weak or
344 no impact of permafrost on DOC and DON, the particulate C and N were affected by the presence
345 of permafrost in summer and autumn but not affected by its presence in spring. Moreover, during
346 freshet the permafrost distribution did not influence the bulk RSM concentration in WSL rivers.
347 This strongly implies that the delivery of RSM in rivers, and its chemical composition, are tightly
348 linked to the thickness of the active layer and limited by transport of soil particles over the
349 suprapermafrost flow to the river channel. This thickness is highest in September at the end of the
350 active season. In agreement with this, the C demonstrated a maximum concentration at 62-64°N,
351 in the sporadic to isolated permafrost zone and was most visible during summer and autumn
352 (**Fig. 3 A**). This latitudinal belt can be considered as a large-scale thawing front for the frozen



353 peat which corresponds to the southern boundary of permafrost persistence. Furthermore, a
354 maximum percentage of particulate C over total C (suspended + dissolved) was also in isolated
355 and sporadic permafrost zones in spring; this maximum shifted to the sporadic permafrost zone
356 in summer and moved northward to the discontinuous permafrost zone in autumn (**Fig. 5 C**). We
357 believe that this corresponds to a progressive increase in the thickness of the active layer which
358 controls the degree of peat and mineral particles leaching from the soil profile to the river. The
359 thickness of this layer increases from spring to autumn and more importantly it moves northward
360 during this period (Trofimova and Balybina, 2014). Enhanced mobilization of nutrients at the
361 “hot spot” of permafrost thaw in frozen peat landscapes was recently demonstrated on a local
362 scale in western Siberia (Loiko et al., 2017).

363 The impact of watershed characteristics on particulate C and N was clearly pronounced
364 with increased C and N concentration in RSM where there were increased bog and lake
365 proportions and decreased C and N concentration where there was increasing forest coverage.
366 The stronger impact of lakes compared to bogs on C concentration in RSM suggests that the
367 generation of C-rich particles occurs more efficiently in large water bodies than in stagnant
368 shallow water bodies. Several mechanisms are likely to operate in this regard. First,
369 photodegradation of DOM in large and shallow lakes of WSL is expected to be quite strong
370 similar to shallow Canadian thaw ponds (Laurion and Mladenov, 2013). Additionally, given the
371 very short transit time of water from the surrounding peat to the lakes via suprapermfrost flow
372 (Ala-aho et al., 2018a, b; Raudina et al., 2018), the allochthonous chromophoric DOM-rich
373 material that arrives to the lakes is subjected to fast degradation and coagulation such as that
374 shown in Scandinavian lakes (Kortelainen et al., 2006b; von Wachenfeldt and Tranvik, 2008).
375 Second, the peat abrasion at the border of the thermokarst lakes and thaw ponds, which are
376 highly abundant in the territory (Polishchuk et al., 2017, 2018), occurs due to wave erosion and
377 thermo-abrasion (Shirokova et al., 2013; Manasyrov et al., 2015). Physical disintegration of peat



378 at the lake coast likely generates a large amount of suspended organic-rich material that can be
379 exported to hydrological networks during, for example, lake drainage or through already existing
380 connecting channels (Kirpotin et al., 2008, 2011). Note that the maximal lake coverage of the
381 WSL territory is in the 63°N to 64°N latitudinal belt (Polishchuk et al., 2017) where maximum
382 C and N concentration in RSM also occurs. Because the majority of thermokarst lakes are
383 isolated water bodies without inlet and outlet, this connectivity is achieved via water movement
384 along the permafrost table in the thawed active layer (Raudina et al., 2018) in the form of so-
385 called suprapermfrost flow between peat bogs, lakes, and rivers.

386 Finally, for particulate P, neither its concentration nor the particulate fraction were
387 affected by permafrost distribution, probably due to the various biological uptake and mineral
388 precipitation processes controlling P removal both in soil profile and in the river water. For
389 example, lakes and bogs retained particulate P, similar to that of dissolved P, which is in
390 agreement with global assessments (Bouwman et al., 2013), P behavior in European northern
391 wetlands and lakes (Lidman et al., 2014), and recent results on dissolved P in the WSL rivers
392 (Vorobyev et al., 2017).

393

394 *4.3. Mechanisms of RSM generation and prospective for climate warming in western* 395 *Siberia*

396 A framework of particulate C, N and P generation in WSL rivers across the permafrost
397 gradient is shown in **Fig. 6**. We suggest that the concentration of suspended particles depends on
398 both the supply and losses in the catchments. The sources of suspended particles in WSL rivers
399 include: (i) vegetation litter which is washed by surficial flow to the river, especially in spring;
400 (ii) surface (peat) soil horizons, which are also most active in spring, especially in the north; (iii)
401 deep peat and mineral horizons which provide the particles via bank abrasion in spring and via
402 suprapermfrost flow in summer and autumn, (iv) lake and bog open water sediments formed



403 either by flocculation of DOM via photo- and bio-degradation processes or via lake coastal
404 abrasion due to wave erosion, and finally, (v) autochthonous organic debris of macrophytes,
405 periphyton and phytoplankton, whose contribution is maximal in summer and autumn. A non-
406 steady-state physical erosion of peat soils in WSL provides maximum particulate nutrients
407 within the most fragile zone of actively thawing permafrost between 62 and 64°N of the
408 sporadic to isolated permafrost zone. The maximal thickness of the active layer progressively
409 moves north during the active season thereby leading to maximal export of particulate C, N, and
410 P at the thawing front. However, we also suggest that part of the differences in mobilized
411 particulates is masked by retention in recipient waters. The transit time of water and particles in
412 the southern WSL rivers is much longer than that in northern rivers (Ala-aho et al., 2017, 2018a, b)
413 hence the biological uptake mechanisms together with physio-chemical processes such as photo-
414 degradation of POC (Mayer et al., 2006; Riggsbee et al., 2008) or cryocoagulation, (Pokrovsky et
415 al., 2018) have sufficient time to act on suspended matter of soil and shallow subsurface waters and
416 to remove the nutrients from the river water as well. In rivers of the continuous permafrost zone, a
417 relatively small stock of nutrient-rich particles within the soil profile and on soil surface (as plant
418 litter) is largely compensated for by a more rapid flushing and shorter travel time through soils and
419 rivers and also lower microbial and phytoplankton activity. As a result, the zone of sporadic to
420 isolated permafrost exhibits both maximal release of soil particles and minimal uptake by in-stream
421 processes. Further to the north, shallow unfrozen peat depth and low biomass cannot supply
422 sufficiently high suspended nutrients and the particulate transport of C and N decreases. In contrast,
423 for P, opposite gradients in supply versus in stream removal may cancel out the net effect of
424 temperature and permafrost on suspended P in the river water.

425 Based on these results we can also speculate on the conditions following warming and
426 permafrost thaw. On a short-term prospective (10-50 years), assuming a soil temperature rise
427 rate of 0.15 to 0.3 degree per 10 years in WSL (Pavlov et al., 2009; Anisimov et al., 2012), the



428 northern part of the WSL (discontinuous and continuous permafrost zones) will transform into
429 sporadic and isolated permafrost zones (Anisimov and Reneva, 2006). This will lead to increase
430 in C and N in RSM and overall increase in particulate versus dissolved transport of C and P.
431 Given the contemporary maximum of C and N at the permafrost thawing front this increase may
432 be two-fold. However, on a longer prospective (50-100 years), even the continuous permafrost
433 zone may disappear (Romanovsky et al., 2008; Nadyozhina et al., 2008) and this will decrease
434 the particulate C and N concentration in the northern rivers and, consequently, their export to the
435 coastal zone of the Kara Sea. Judging from the actual difference in nutrient concentrations
436 among adjusting permafrost zones, this decrease may be around a factor of 2 to 3. Furthermore,
437 on the same long-term prospective, the drainage of lakes and disappearance of bogs due to
438 colonization of northern tundra by forests (Anisimov et al., 2011; Anisimov and Sherstiukov,
439 2016) should lead to a further decrease in particulate nutrient load of WSL rivers.

440

441 **Conclusions:**

442 Relatively low bulk RSM concentration in WSL rivers stems from low runoff in this flat
443 peatland province of boreal and subarctic zone. High concentrations of C and N in the RSM of
444 WSL rivers reflect the essentially organic nature of soils across the WSL. At the
445 isolated/sporadic permafrost zone, we observed a maximum concentration of C and N in the
446 RSM and maximal fraction of particulate OC relative to total (dissolved + particulate). This
447 suggests the enhanced generation of C,N-rich RSM and a thawing front of permafrost, where
448 thickness of the active layer is maximal. The C and N concentrations in particulate load of WSL
449 rivers decrease with forest coverage of the watershed and increase with the proportion of lakes
450 and bogs; however, the bulk concentration of RSM did not depend on landscape parameters of
451 the watersheds. This implies generation of CN-rich particles via coastal peat abrasion, sediment
452 resuspension of photo- and bio-coagulation of DOM in lentic surface waters which are



453 hydrologically connected to rivers. To assess a northward permafrost boundary and forest line
454 shifting with increase in air and soil temperature we used a substituting space for time scenario
455 of climate warming in the WSL that was well developed for the dissolved fraction of C and
456 nutrients. From a short-term climate warming prospective, the effect of a northward shift of
457 permafrost boundary may produce about a two-fold increase in particulate C and N
458 concentration in rivers of the discontinuous and continuous permafrost zones, and thus may
459 enhance the export of these nutrients by the most northern WSL rivers to the Arctic Ocean. On a
460 long-term prospective, the disappearance of permafrost in the northern part of WSL will
461 decrease the concentrations of these nutrients to their current level. The P is unlikely to be
462 significantly affected by permafrost change. Moreover, within a long-term climate warming
463 scenario, the drainage of lakes and transformation of bogs to forest may decrease nutrient
464 concentration in RSM and corresponding export flux to the Arctic Ocean.

465

466 **Acknowledgements:**

467 This work was supported by RSCF No 18-17-00237 “Mechanisms of hydrochemical runoff of the
468 Ob river...” (analyses, modeling); RFBR project № 18-35-00563\18 , Ministry of Education and
469 Science of the Russian Federation № 6.7515.2017/9.1, and by VR (the Swedish Research
470 Council) grant no. 325-2014-6898.

471

472 **References**

473

- 474 Ala-aho, P., Soulsby, C., Pokrovsky, O.S., Kirpotin, S.N., Karlsson, J., Serikova, S.,
475 Manasypov, R.M., Krickov, I., Lim, A., and Tetzlaff D.: Permafrost and lakes control river
476 isotope composition across a boreal-arctic transect in the western Siberia lowland, Environ.
477 Res. Lett., 13 (3), 034028, <https://doi.org/10.1088/1748-9326/aaa4fe>, 2018a.
- 478 Ala-Aho, P., Soulsby, C., Pokrovsky, O.S., Kirpotin, S.N., Karlsson, J., Serikova, S., Vorobyev,
479 S.N., Manasypov, R.M., Loiko, S., and Tetzlaff D.: Using stable isotopes to assess surface
480 water source dynamics and hydrological connectivity in a high-latitude wetland and
481 permafrost influenced landscape, J. Hydrol., 556, 279–293,
482 <https://doi.org/10.1016/j.jhydrol.2017.11.024>, 2018b.
- 483 Anisimov, O. A., Anokhin, A., Lavrov, S. A., Malkova, G. V., Pavlov A.V., Romanovskiy,



- 484 V.E., Streletskiy, D. A., Kholodov, A.L., and Shiklomanov, N. I.: Continental multiyear
485 permafrost // Methods of study the sequences of climate changes for nature systems. Ed.
486 S.M. Semenov. Moscow: VNIIGMI, 2012. P. 268–328. (In Russian)
- 487 Anisimov, O.A., and Sherstiukov, A.B.: Evaluating the effect of climatic and environmental
488 factors on permafrost in Russia, Earth's Cryosphere, XX (№ 2), 78–86, 2016.
- 489 Anisimov, O.A., Zhiltsova, E.L., and Reneva, S.A.: Estimation of critical levels of climate
490 change impact on natural land ecosystems of Russian territory, Meteorology and
491 Hydrology, 11, 723-730, <https://doi.org/10.3103/S1068373911110033>, 2011.
- 492 Anisimov, O., and Reneva, S.: Permafrost and Changing Climate: The Russian Perspective,
493 AMBIO, 35(4), 169–175, <https://doi.org/10.1579/0044-7447>, 2006.
- 494 Battin, T. J., Kaplan, L. A., Findlay, S., Hopkinson, C. S., Marti, E., Packman A. I., Newbold, J.
495 D., and Sabater, F.: Biophysical controls on organic carbon fluxes in fluvial networks, Nat.
496 Geosci., 1, 95–100, <https://doi.org/10.1038/ngeo602>, 2008.
- 497 Blois, J. L., Williams, J. W., Fitzpatrick, M. C., Jackson, S. T., and Ferrier, S.: Space can
498 substitute for time in predicting climate-change effects on biodiversity, PNAS, 110 (23),
499 9374-9379, <https://doi.org/10.1073/pnas.1220228110>, 2013.
- 500 Bring, A., Fedorova, I., Dibike, Y., Hinzman, L., Mård, J., Mernild, S. H., Prowse, T.,
501 Semenova, O., Stuefer, S. L., and Woo, M.-K.: Arctic terrestrial hydrology: A synthesis of
502 processes, regional effects, and research challenges, J. Geophys. Res.-Biogeo., 121(3),
503 621–649, <https://doi.org/10.1002/2015JG003131>, 2016.
- 504 Bouwman, A. F., Bierkens, M. F. P., Griffioen, J., Hefting, M. M., Middelburg, J. J., Middelkoop,
505 H., and Slomp, C. P.: Nutrient dynamics, transfer and retention along the aquatic continuum
506 from land to ocean: towards integration of ecological and biogeochemical models,
507 Biogeosciences 10, 1-23, <https://doi.org/10.5194/bg-10-1-2013>, 2013.
- 508 Cai, Y., L. Guo, T. A. Douglas, and T. E. Whitedge: Seasonal variations in nutrient concentrations
509 and speciation in the Chena River, Alaska, J. Geophys. Res.-Biogeo., 113, G03035,
510 <https://doi:10.1029/2008JG000733>, 2008.
- 511 Cole, J. J., Prairie, Y. T., Caraco, N. F., McDowell, W. H., Tranvik, L. J., Striegl, R. G., Duarte, C.
512 M., Kortelainen, J. P., Downing, A., Middelburg, J. J., Melack, J.: Plumbing the Global
513 Carbon Cycle: Integrating Inland Waters into the Terrestrial Carbon Budget, Ecosystems,
514 10(1), 172–185, <https://doi.org/10.1007/s10021-006-9013-8>, 2007.
- 515 Dittmar, Th., and Kattner, G.: The biogeochemistry of the river and shelf ecosystem of the Arctic
516 Ocean: a review, Mar. Chem., 83, 103-120, [https://doi.org/10.1016/S0304-4203\(03\)00105-1](https://doi.org/10.1016/S0304-4203(03)00105-1),
517 2003.
- 518 Dornblaser, M. M., and R. G. Striegl: Nutrient (N, P) loads and yields at multiple scales and
519 subbasin types in the Yukon River basin, Alaska, J. Geophys. Res.-Biogeo., 112, G04S57,
520 <https://doi.org/10.1029/2006JG000366>, 2007.
- 521 Emmerton, C. A., Lesack, L. F. W., and Vincent, W. F.: Mackenzie River nutrient delivery to the
522 Arctic Ocean and effects of the Mackenzie Delta during open water conditions, Global
523 Biogeochem. Cy., 22, GB1024, <https://doi.org/10.1029/2006GB002856>, 2008a.
- 524 Emmerton, C. A., Lesack, L. F. W., and Vincent, W. F.: Nutrient and organic matter patterns across
525 the Mackenzie River, estuary and shelf during the seasonal recession of sea-ice, J. Marine
526 Syst., 74(3–4), 741–755, <https://doi.org/10.1016/j.jmarsys.2007.10.001>, 2008b.
- 527 Frey, K. E., McClelland, J. W., Holmes, R. M., and Smith, L. C.: Impacts of climate warming and
528 permafrost thaw on the riverine transport of nitrogen and phosphorus to the Kara Sea, J.
529 Geophys. Res.-Biogeo., 112, G04S58, doi:10.1029/2006JG000369, 2007a.
- 530 Frey, K. E., Siegel, D. I., and Smith, L.C.: Geochemistry of west Siberian streams and their
531 potential response to permafrost degradation, Water Resour. Res., 43(3), W03406,
532 <https://doi.org/10.1029/2006WR004902>, 2007b.
- 533 Frey, K. E., and Smith, L.C.: Amplified carbon release from vast West Siberian peatlands by 2100,



- 534 Geophys. Res. Lett., 32, L09401, <https://doi.org/10.1029/2004GL022025>, 2005.
- 535 Frey, K. E., and McClelland, J. W.: Impacts of permafrost degradation on arctic river
536 biogeochemistry, *Hydrol. Process.*, 23(1), 169–182, <https://doi.org/10.1002/hyp.7196>, 2009.
- 537 Galy, V., Peucker-Ehrenbrink, B. and Eglinton, T.: Global carbon export from the terrestrial
538 biosphere controlled by erosion, *Nature*, 521, 204–207, <http://dx.doi.org/10.1038/nature14400>,
539 2015.
- 540 Gareis, J. A. L., and Lesack, L. F. W.: Fluxes of particulates and nutrients during hydrologically
541 defined seasonal periods in an ice-affected great Arctic river, the Mackenzie, *Water Resour.*
542 *Res.*, 53(7), 6109–6132, <https://doi.org/10.1002/2017WR020623>, 2017.
- 543 Gebhardt, A. C., Gaye-Haake, C., Unger, D., Lahajnar, N., and Ittekkot V.: Recent particulate
544 organic carbon and total suspended matter fluxes from the Ob and Yenisei Rivers into the
545 Kara Sea (Siberia), *Mar. Geol.*, 207, 225– 245, <https://doi.org/10.1016/j.margeo.2004.03.010>,
546 2004.
- 547 Gentsch, N., Mikutta, R., Alves, R. J. E., Barta, J., Čapek, P., Gittel, A., Hugelius, G., Kuhry, P.,
548 Lashchinskiy, N., Palmtag, J., Richter, A., Šantručková, H., Schneckner, J., Shibistova, O.,
549 Urich, T., Wild, B., and Guggenberger, G.: Storage and transformation of organic matter
550 fractions in cryoturbated permafrost soils across the Siberian Arctic, *Biogeosciences*, 12(14),
551 4525–4542, <http://dx.doi.org/10.5194/bg-12-4525-2015>, 2015.
- 552 Gladyshev, M. I., Kolmakova, O. V., Tolomeev, A. P., Anishchenko, O. V., Makhutova, O. N.,
553 Kolmakova, A. A., Kravchuk, E. S., Glushchenko, L. A., Kolmakov, V. I., and Sushchik,
554 N.N.: Differences in organic matter and bacterioplankton between sections of the largest
555 Arctic river: Mosaic or continuum?, *Limnol. Oceanogr.*, 60, 1314–1331,
556 <https://doi.org/10.1002/lno.10097>, 2015.
- 557 Goñi, M. A., Hatten, J. A., Wheatcroft, R. A., and Borgeld, J. C.: Particulate organicmatter export
558 by two contrasting small mountainous rivers from the Pacific Northwest, U.S.A., *J. Geophys.*
559 *Res.-Biogeo.*, 118, 1–23, <https://doi.org/10.1002/jgrg.20024>, 2013.
- 560 Gordeev, V. V., Martin, J. M., Sidorov, I. S., and Sidorova, M. V.: A reassessment of the
561 Eurasian river input of water, sediment, major elements, and nutrients to the Arctic Ocean,
562 *Am. J. Sci.*, 296(6), 664–691, 1996.
- 563 Gordeev, V. V., and Kravchishina, M. D.: River flux of dissolved organic carbon (DOC) and
564 particulate organic carbon (POC) to the Arctic Ocean: what are the consequences of the
565 global changes?, in: *Influence of Climate Change on Changing Arctic and Sub-Arctic*
566 *Conditions*, Springer, 145–160, 2009.
- 567 Grosse, G., Goetz, S., McGuire, D., Romanovsky, V.E., and Schuur E. A. G.: Changing permafrost
568 in a warming world and feedbacks to the Earth system, *Environ. Res. Lett.*, 11, 040201,
569 <https://doi:10.1088/1748-9326/11/4/040201>, 2016.
- 570 Guo, L., and Macdonald, R. W.: Source and transport of terrigenous organic matter in the upper
571 Yukon River: Evidence from isotope ($\delta^{13}C$, $\delta^{14}C$, and $\delta^{15}N$) composition of dissolved,
572 colloidal, and particulate phases, *Global Biogeochem. Cy.*, 20, GB2011,
573 <https://doi.org/10.1029/2005GB002593>, 2006.
- 574 Guo, L., Zhang, J.-Z., and Guéguen, C.: Speciation and fluxes of nutrients (N, P, Si) from the upper
575 Yukon River, *Global Biogeochem. Cy.*, 18, GB1038, <https://doi.org/10.1029/2003GB002152>,
576 2004.
- 577 Hilton, R. G., Galy, V., Gaillardet, J., Dellinger, M., Bryant, C., O'Regan, M., Gröcke, D.R.,
578 Coxall, H., Bouchez, J., and Calmels, D.: Erosion of organic carbon in the Arctic as a
579 geological carbon dioxide sink, *Nature*, 524, 84–87, <https://doi:10.1038/nature14653>, 2015.
- 580 Hilton, R.G.: Climate regulates the erosional carbon export from the terrestrial biosphere,
581 *Geomorphology*, 277, 118–132, <https://doi.org/10.1016/j.geomorph.2016.03.028>, 2017.
- 582 Holmes, R. M., Peterson, B. J., Gordeev, V. V., Zhulidov, A. V., Meybeck, M., Lammers, R. B.,
583 and Vorosmarty, C. J.: Flux of nutrients from Russian rivers to the Arctic Ocean: Can we



- 584 establish a baseline against which to judge future changes?, *Water Resour. Res.*, 36(8), 2309–
585 2320, <https://doi.org/10.1029/2000WR900099>, 2000.
- 586 Jeong, J. J., Bartsch, S., Fleckenstein, J. H., Matzner, E., Tenhunen, J. D., Lee, S. D., Park, S. K.,
587 and Park, J.-H.: Differential storm responses of dissolved and particulate organic carbon in a
588 mountainous headwater stream, investigated by high-frequency, in situ optical measurements,
589 *J. Geophys. Res.-Biogeo.*, 117(3), 1–13, <https://doi.org/10.1029/2012JG001999>, 2012.
- 590 Kirpotin, S., Polishchuk, Y., Zakharova, E., Shirokova, L., Pokrovsky, O., Kolmakova, M., and
591 Dupre, B.: One of possible mechanisms of thermokarst lakes drainage in West-Siberian North,
592 *Int. J. Environ. Stud.*, 65(5), 631–635, <https://doi.org/10.1080/00207230802525208>, 2008.
- 593 Kirpotin, S., Polishchuk, Y., Bryksina, N., Sugaipova, A., Kouraev, A., Zakharova, E., Pokrovsky,
594 O.S., Shirokova, L., Kolmakova, M., Manassypov, R., and Dupre B.: West Siberian palsa
595 peatlands: distribution, typology, hydrology, cyclic development, present-day climate-driven
596 changes and impact on CO₂ cycle, *Int. J. Environ. Stud.*, 68(5), 603–623,
597 <https://doi.org/10.1080/00207233.2011.593901>, 2011.
- 598 Kortelainen, P., Mattsson, T., Finér, L., Ahtainen, M., Saukkonen, S., and Sallantausta, T.: Controls
599 on the export of C, N, P and Fe from undisturbed boreal catchments, Finland, *Aquat. Sci.*,
600 68(4), 453–468, <https://doi.org/10.1007/s00027-006-0833-6>, 2006a.
- 601 Kortelainen, P., Rantakari, M., Huttunen, J. T., Mattsson, T., Alm, J., Juutinen, S., Larmola, T.,
602 Silvola, J., and Martikainen, P. J.: Sediment respiration and lake trophic state are important
603 predictors of large CO₂ evasion from small boreal lakes, *Glob. Change Biol.*, 12, 1554–1567,
604 <https://doi.org/10.1111/j.1365-2486.2006.01167.x>, 2006b.
- 605 Kremenetski, K. V., Velichko, A. A., Borisova, O. K., MacDonald, G. M., Smith, L. C., Frey, K.
606 E., and Orlova, L. A.: Peatlands of the West Siberian Lowlands: Current knowledge on
607 zonation, carbon content, and Late Quaternary history, *Quaternary Sci. Rev.*, 22, 703–723,
608 [https://doi.org/10.1016/S0277-3791\(02\)00196-8](https://doi.org/10.1016/S0277-3791(02)00196-8), 2003.
- 609 Kutscher, L., Mörth, C.-M., Porcelli, D., Hirst, C., Maximov, T. C., Petrov, R. E., and Andersson,
610 P. S.: Spatial variation in concentration and sources of organic carbon in the Lena River,
611 Siberia, *J. Geophys. Res.-Biogeo.*, 122, 1999–2016, <https://doi.org/10.1002/2017JG003858>,
612 2017.
- 613 Lal, R.: Soil erosion and the global carbon budget, *Environ. Int.*, 29, 437–450,
614 [http://dx.doi.org/10.1016/S0160-4120\(02\)00192-7](http://dx.doi.org/10.1016/S0160-4120(02)00192-7), 2003.
- 615 Lamoureux, S. F., and Lafrenière, M. J.: Seasonal fluxes and age of particulate organic carbon
616 exported from Arctic catchments impacted by localized permafrost slope disturbances,
617 *Environ. Res. Lett.*, 9, 045002, <https://doi.org/10.1088/1748-9326/9/4/045002>, 2014.
- 618 Laurion, I., and Mladenov, N.: Dissolved organic matter photolysis in Canadian Arctic thaw ponds,
619 *Environ. Res. Lett.*, 8, 035026, <https://doi.org/10.1088/1748-9326/8/3/035026>, 2013.
- 620 Lee, M. H., Payeur-Poirier, J. L., Park, J. H., and Matzner, E.: Variability in runoff fluxes of
621 dissolved and particulate carbon and nitrogen from two watersheds of different tree species
622 during intense storm events, *Biogeosciences*, 13(18), 5421–5432, <https://doi.org/10.5194/bg-13-5421-2016>, 2016.
- 624 Le Fouest, V., Babin, M., and Tremblay, J.-É.: The fate of riverine nutrients on Arctic shelves,
625 *Biogeosciences*, 10, 3661–3677, <https://doi.org/10.5194/bg-10-3661-2013>, 2013.
- 626 Leonov, A. B., and Chicherina, O. V.: Export of biogenic components of the riverine flux to the
627 White Sea, *Water Resour.*, 31(2), 170–192, 2004.
- 628 Li, M., Peng, C., Wang, M., Xue, W., Zhang, K., Wang, K., Shi, G., and Zhu, Q.: The carbon flux
629 of global rivers: A re-evaluation of amount and spatial patterns, *Ecol. Indic.*, 80, 40–51,
630 <https://doi.org/10.1016/j.ecolind.2017.04.049>, 2017.
- 631 Lidman, F., Kohler, S. J., Morth, C.-M., and Laudon, H.: Metal transport in the boreal landscape –
632 the role of wetlands and the affinity for organic matter, *Environ. Sci. Technol.*, 48, 3783–3790,
633 <https://doi.org/10.1021/es4045506>, 2014.



- 634 Lobbes, J. M., Fitznar, H. P., and Kattner, G.: Biogeochemical characteristics of dissolved and
635 particulate organic matter in Russian rivers entering the Arctic Ocean, *Geochim. Cosmochim.*
636 *Ac.*, 64, 2973–2983, [https://doi.org/10.1016/S0016-7037\(00\)00409-9](https://doi.org/10.1016/S0016-7037(00)00409-9), 2000.
- 637 Loiko, S. V., Pokrovsky, O. S., Raudina, T., Lim, A., Kolesnichenko, L. G., Shirokova, L. S.,
638 Vorobyev, S. N., and Kirpotin, S. N.: Abrupt permafrost collapse enhances organic carbon,
639 CO₂, nutrient, and metal release into surface waters, *Chem. Geol.*, 471, 153–165,
640 <https://doi.org/10.1016/j.chemgeo.2017.10.002>, 2017.
- 641 Ludwig, W., and Probst, J. L.: Predicting the oceanic input of organic carbon by continental
642 erosion, *Global Biogeochemical Cy.*, 10(1), 23–41, <https://doi.org/10.1029/95GB02925>, 1996.
- 643 Manasypov, R. M., Vorobyev, S. N., Loiko, S. V., Kritzkov, I. V., Shirokova, L. S., Shevchenko,
644 V. P., Kirpotin, S. N., Kulizhsky, S. P., Kolesnichenko, L. G., Zemtzov, V. A., Sinkinov, V.
645 V., and Pokrovsky, O. S.: Seasonal dynamics of organic carbon and metals in thermokarst
646 lakes from the discontinuous permafrost zone of western Siberia, *Biogeosciences*, 12, 3009–
647 3028, <https://doi.org/10.5194/bg-12-3009-2015>, 2015.
- 648 Mayer, L. M., Schick, L. L., Skorko, K., and Boss, E.: Photodissolution of particulate organic
649 matter from sediments, *Limnol. Oceanogr.*, 51(2), 1064–1071,
650 <https://doi.org/10.4319/lo.2006.51.2.1064>, 2006.
- 651 McClelland, J. W., Stieglitz, M., Pan, F., Holmes, R. M., and Peterson, B. J.: Recent changes in
652 nitrate and dissolved organic carbon export from the upper Kuparuk River, North Slope,
653 Alaska, *J. Geophys. Res.-Bioge.*, 112, G04S60,
654 <http://doi.wiley.com/10.1029/2006JG000371>, 2007.
- 655 McClelland, J. W., Townsend-Small, A., Holmes, R. M., Pan, F., Stieglitz, M., Khosh, M., and
656 Peterson B. J.: River export of nutrients and organic matter from the North Slope of Alaska to
657 the Beaufort Sea, *Water Resour. Res.*, 50(2), 1823–1839,
658 <https://doi.org/10.1002/2013WR014722>, 2014.
- 659 McClelland, J. W., Holmes, R. M., Raymond, P. A., Striegl, R. G., Zhulidov, A. V., Zimov, S. A.,
660 Zimov, N., Tank, S. E., Spencer, R. G. M., Staples, R., Gurtovaya, T. Y., and Grif, C. G.:
661 Particulate organic carbon and nitrogen export from major Arctic rivers, *Global Biogeochem.*
662 *Cy.*, 30, 629–643, <https://doi.org/10.1002/2015GB005351>, 2016.
- 663 Meybeck, M.: C, N, P and S in rivers: from sources to global inputs, in: Wollast, R., Mackenzie, F.,
664 and Chou, L., editors, *Interaction of C, N, P, and S biogeochemical cycles and global change*,
665 *NATO ASI Series: Berlin, Heidelberg, Springer-Verlag*, 163–193, 1993.
- 666 Nadyozhina, E. D., Shkolnik, I. M., Pavlova, T. V., Mol Kentin, E. K., and Semioshina, A. A.:
667 Permafrost response to the climate warming as simulated by the regional climate model of
668 the main geophysical observatory, *Kriosfera Zemli*, XII(3), 3–11, 2008. (In Russian)
- 669 Pavlov, A. V., and Malkova, G. V.: Mapping of trends of the contemporary ground temperature
670 changes in the Russian north, *Kriosfera Zemli*, XIII(4), 32–39, 2009. (In Russian)
- 671 Pokrovsky, O. S., Manasypov, R. M., Shirokova, L. S., Loiko, S. V., Krickov, I. V., Kopysov S. G.,
672 Zemtzov, V. A., Kulizhsky, S. P., Vorobyev, S. N., and Kirpotin, S. N.: Permafrost coverage,
673 watershed area and season control of dissolved carbon and major elements in western Siberia
674 rivers, *Biogeosciences*, 12, 6301–6320, <https://doi.org/10.5194/bg-12-6301-2015>, 2015.
- 675 Pokrovsky, O. S., Manasypov, R. M., Loiko, S. V., Krickov, I. A., Kopysov, S. G., Kolesnichenko,
676 L. G., Vorobyev, S. N., and Kirpotin, S. N.: Trace element transport in western Siberia rivers
677 across a permafrost gradient, *Biogeosciences*, 13, 1877–1900, <https://doi:10.5194/bgd-12-17857-2015>, 2016.
- 679 Pokrovsky, O. S., Schott, J., Kudryavtzev, D. I., and Dupre, B.: Basalts weathering in Central
680 Siberia under permafrost conditions, *Geochim. Cosmochim. Ac.*, 69, 5659–5680,
681 <https://doi.org/10.1016/j.gca.2005.07.018>, 2005.
- 682 Pokrovsky, O. S., Viers, J., Shirokova, L. S., Shevchenko, V. P., Filipov, A. S., and Dupré B.:
683 Dissolved, suspended, and colloidal fluxes of organic carbon, major and trace elements in



- 684 Severnaya Dvina River and its tributary, *Chem. Geol.*, 273, 136–149,
685 <https://doi.org/10.1016/j.chemgeo.2010.02.018>, 2010.
- 686 Polishchuk, Y. M., Bogdanov, A. N., Polishchuk, V. Y., Manasypov, R. M., Shiorkova, L. S.,
687 Kirpotin, S. N., and Pokrovsky, O. S.: Size-distribution, surface coverage, water, carbon and
688 metal storage of thermokarst lakes (> 0.5 ha) in permafrost zone of the Western Siberia
689 Lowland, *Water*, 9 (3), 228, <https://doi.org/10.3390/w9030228>, 2017.
- 690 Polishchuk, Y. M., Bogdanov, A. N., Muratov, I. N., Polishchuk, V. Y., Lim, A., Manasypov, R.
691 M., Shirokova, L. S., and Pokrovsky, O. S.: Minor contribution of small thaw ponds to the
692 pools of carbon and methane in the inland waters of the permafrost-affected part of western
693 Siberian Lowland, *Environ. Res. Lett.*, 13(4), 045002, <https://doi.org/10.1088/1748-9326/aab046>, 2018.
- 695 Raudina, T. V., Loiko, S. V., Lim, A. G., Krickov, I. V., Shirokova, L. S., Istignichev, G. I.,
696 Kuzmina, D. M., Kulizhsky, S. P., Vorobyev, S. N., and Pokrovsky, O. S.: Dissolved organic
697 carbon and major and trace elements in peat porewater of sporadic, discontinuous, and
698 continuous permafrost zones of western Siberia, *Biogeosciences*, 14, 3561–3584,
699 <https://doi.org/10.5194/bg-14-3561-2017>, 2017.
- 700 Raudina, T.V., Loiko, S. V., Lim, A., Manasypov, R. M., Shirokova, L. S., Istigechev, G. I.,
701 Kuzmina, D. M., Kulizhsky, S. P., Vorobyev, S. N., and Pokrovsky, O. S.: Permafrost thaw
702 and climate warming may decrease the CO₂, carbon, and metal concentration in peat soil
703 waters of the Western Siberia Lowland, *Sci. Total. Environ.*, 634, 1004–1023,
704 <https://doi.org/10.1016/j.scitotenv.2018.04.059>, 2018.
- 705 Richardson, D. C., Newbold, J. D., Aufdenkampe, A. K., Taylor, P. G., and Kaplan, L. A.:
706 Measuring heterotrophic respiration rates of suspended particulate organic carbon from stream
707 ecosystems, *Limnol. Oceanogr.-Meth.*, 11(5), 247–261,
708 <https://doi.org/10.4319/lom.2013.11.247>, 2013.
- 709 Riggsbee, J., Orr, C., Leech, D., Doyle, M., and Wetzel, R.: Suspended sediments in river
710 ecosystems: photochemical sources of dissolved organic carbon, dissolved organic nitrogen,
711 and adsorptive removal of dissolved iron, *J. Geophys. Res. Biogeosci.*, 113, G03019,
712 <https://doi.org/10.1029/2007JG000654>, 2008.
- 713 Romanovsky, V. E., Kholodov, A. L., Marchenko, S. S., Oberman, N. G., Drozdov, D. S.,
714 Malkova, G. V., Moskalenko, N. G., Vasiliev, A. A., Sergeev, D. O., Zheleznyak, M. N.:
715 Thermal State and Fate of Permafrost in Russia: First Results of IPY, in: Proceedings of the
716 9th International Conference on Permafrost, University of Alaska, Fairbanks, June 29 –
717 July 3, 2008 / Kane D.L., Hinkel K.M. (eds.). vol. 2, 1511–1518.
- 718 Rosgidromet, R. F.: *Doklad ob osobennostyakh klimata na territorii Rossiyskoy Federatsii za 2016*
719 *god*, M.: Federal'naya sluzhba po gidrometeorologii i monitoringu okruzhayushchey sredy,
720 Rosgidromet, Moscow, 2017. (In Russian)
- 721 Savenko, V. S., Pokrovskii, O. S., Dupre, B., Baturin, G. N.: *Dokl. Akad. Nauk* 398, 97–101, 2004
722 [Dokl. Earth Sci. 398, 938, 2004].
- 723 Sheng, Y., Smith, L. C., MacDonald, G. M., Kremenetski, K. V., Frey, K. E., Velichko, A. A., Lee,
724 M., Beilman, D. W., and Dubinin, P.: A high-resolution GIS-based inventory of the west
725 Siberian peat carbon pool, *Global Biogeochem. Cy.*, 18, GB3004,
726 <https://doi.org/10.1029/2003GB002190>, 2004.
- 727 Schlesinger, W. H., and Melack, J. M.: Transport of organic carbon in the world's rivers, *Tellus*,
728 33, 172–187, <https://doi.org/10.3402/tellusa.v33i2.10706>, 1981.
- 729 Schädel, C., Schuur, E. A. G., Bracho, R., Elberling, B., Knoblauch, C., Lee, H., Luo, Y., Shaver,
730 G. R., and Turetsky, M. R.: Circumpolar assessment of permafrost C quality and its
731 vulnerability over time using long-term incubation data, *Global Change Biol.*, 20, 641–652,
732 <https://doi.org/10.1111/gcb.12417>, 2014.




- 733 Shirokova, L. S., Pokrovsky, O. S., Kirpotin, S. N., Desmukh, C., Pokrovsky, B. G., Audry, S.,
734 Viers, J.: Biogeochemistry of organic carbon, CO₂, CH₄, and trace elements in thermokarst
735 water bodies in discontinuous permafrost zones of Western Siberia, *Biogeochemistry*, 113,
736 573–593, <https://doi.org/10.1007/s10533-012-9790-4>, 2013.
- 737 Stepanova, V. M., Pokrovsky, O. S., Viers, J., Mironycheva-Tokareva, N. P., Kosykh, N. P., and
738 Vishnyakova, E. K.: Elemental composition of peat profiles in western Siberia: Effect of the
739 micro-landscape, latitude position and permafrost coverage, *Appl. Geochem.*, 53, 53–70,
740 <https://doi.org/10.1016/j.apgeochem.2014.12.004>, 2015.
- 741 Tian, H., Yang, Q., Najjar, R., Ren, W., Friedrichs, M. A. M., Hopkinson, C. S., and Pan, S.:
742 Anthropogenic and climatic influences on carbon fluxes from Eastern North America to the
743 Atlantic ocean: a process-based modeling study, *J. Geophys. Res.-Biogeo.*, 752–772,
744 <http://dx.doi.org/10.1002/2014JG002760>, 2015.
- 745 Trofimova, I. E., and Balybina, A. S.: Classification of climates and climatic regionalization of the
746 West-Siberian plain, *Geography and Natural Resources*, 35(2), 114–122,
747 <https://doi.org/10.1134/S1875372814020024>, 2014.
- 748 Vonk, J. E., Tank, S. E., Bowden, W. B., Laurion, I., Vincent, W. F., Alekseychik, P., Amyot, M.,
749 Billet, M. F., Canário, J., Cory R. M., Deshpande, B. N., Helbig, M., Jammet, M., Karlsson, J.,
750 Larouche, J., MacMillan, G., Rautio, M., Walter Anthony, K. M., and Wickland, K. P.:
751 Reviews and syntheses: Effects of permafrost thaw on Arctic aquatic ecosystems,
752 *Biogeosciences*, 12(23), 7129–7167, <https://doi.org/10.5194/bg-12-7129-2015>, 2015.
- 753 von Wachenfeldt, E., and Tranvik, L. J.: Sedimentation in boreal lakes - The role of flocculation of
754 allochthonous dissolved organic matter in the water column, *Ecosystems*, 11(5), 803–814,
755 <https://doi.org/10.1007/s10021-008-9162-z>, 2008.
- 756 Vorobyev, S. N., Pokrovsky, O. S., Serikova, S., Manasypov, R. M., Krickov, I. V., Shirokova, L.
757 S., Lim, A., Kolesnichenko, L. G., Kirpotin, S. N., and Karlsson, J.: Permafrost boundary shift
758 in western Siberia may not modify dissolved nutrient concentrations in rivers, *Water*, 9, 985,
759 <https://doi.org/10.3390/w9120985>, 2017.
- 760 Wiegner, T. N., Tubal, R. L., and MacKenzie, R. A.: Bioavailability and export of dissolved
761 organic matter from a tropical river during base- and stormflow conditions, *Limnol.*
762 *Oceanogr.*, 54(4), 1233–1242, <https://doi.org/10.4319/lo.2009.54.4.1233>, 2009.
- 763 Wrona, F. J., Johansson M., Culp J. M., Jenkins A., Mård J., Myers-Smith, I. H., Prowse, T. D.,
764 Vincent, W. F., and Wookey, P. A.: Transitions in Arctic ecosystems: Ecological implications
765 of a changing hydrological regime, *J. Geophys. Res.-Biogeo.*, 121(3), 650–674,
766 <https://doi.org/10.1002/2015JG003133>, 2016.
- 767 Unger, D., Gaye-Haake, B., Neumann, K., Gebhardt, A. C., and Ittekkot, V.: Biogeochemistry of
768 suspended and sedimentary material in the Ob and Yenisei rivers and Kara Sea: amino acids
769 and amino sugars, *Cont. Shelf. Res.*, 25(4), 437–460, <https://doi.org/10.1016/j.csr.2004.09.014>,
770 2005.
- 771 Yool, A., Popova, E. E., and Coward, A. C.: Future change in ocean productivity: Is the Arctic the
772 new Atlantic?, *J. Geophys. Res.-Oceans*, 120(12), 7771–7790,
773 <https://doi.org/10.1002/2015JC011167>, 2015.
- 774 Yang, Q., Zhang, X., Xu, X., Asrar, G. R., Smith, R. A., Shih, J. S., and Duan, S.: Spatial patterns
775 and environmental controls of particulate organic carbon in surface waters in the conterminous
776 United States, *Sci. Total. Environ.*, 554–555, 266–275,
777 <https://doi.org/10.1016/j.scitotenv.2016.02.164>, 2016.

778

779

780

781



782


783 **Fig. 1.** Sampling sites and physio-geographical context of WSL territory investigated in this
784 work. The sampling numbers are explained in Table S1.

785



786

787



788

789


790

791 **Fig. 2.** Particulate ($> 0.45 \mu\text{m}$) C (A), N (B), P (C) concentration in the RMS (%) and C: N ratio
792 (D) in RSM as a function of river watershed size. The solid lines represent 2nd degree
793 polynomial fitting of the data with regression coefficients shown for each season in
794 corresponding panels. Only the curves with $R^2 > 0.3$ are depicted.

795

796

797



798

799

800

801 **Fig. 3.** Box plot of first and third quartiles (25 and 75%) of C (A), N (B) and P (C) concentration

802 in RSM (mass %) in five permafrost zones over three seasons. The C, N and P concentrations in

803 the river water are shown in panels D, E and F, respectively.

804

805

806

807


808

809

810



811



812


813

814 **Fig. 4.** The dependence of C concentration in RSM (%) on the coverage of watershed by bogs

815 (A), lakes (B) and forest (C). The solid lines represent 2nd degree polynomial fitting of the data

816 with regression coefficients shown for each season in corresponding panels.

817




818

819

820

821 **Fig. 5.** Fraction of particulate OC of total (dissolved + particulate) form plotted as a function of
822 latitude (A), lake fraction on the watershed (B) and a box plot of fractions for 5 permafrost zones
823 (C). The solid lines in A and B represent 2nd degree polynomial (A, autumn) and linear (B,
824 spring) fitting of the data with regression coefficients equal to 0.32 and 0.52, respectively.



825

826 **Fig. 6.** A cartoon of spatial and temporal partitioning of particulate nutrients in WSL rivers
 827 across the permafrost gradient. The panels **A**, **B**, **C** and **D** represent from main sources (**A**, lakes
 828 and bogs in summer and **B**, alluvial deposits in spring) and sinks (**C**, photo- and bio-degradation)
 829 and **D**, uptake by taiga forest) of particulate nutrients in WSL rivers. The panel **E** depicts the
 830 spatial gradient of C and N in RSM occurring in spring (blue line) and autumn (red line). A non
 831 steady-state physical erosion of peat soils in WSL provides the maximum of particulate nutrients
 832 within the zone of most “fragile”, actively thawing permafrost. The maximal thickness of active
 833 layer progressively moves to the north during the active season thus leading to the maximal
 834 removal of particulate C, N, and P at the thawing front.



835 **Table 1.** Mean (\pm SD) values of RSM, C, N, P concentration (mass %) and relative proportion
 836 of suspended C and P overall total concentration for 5 permafrost zones and 3 seasons across the
 837 WSL transect.

Season	Variable	Permafrost				
		Absent	Isolated	Sporadic	Discontinuous	Continuous
Spring	RSM, mg/l	6.2 \pm 4.9	4.9 \pm 1.5	7.2 \pm 3.0	7.7 \pm 2.5	10.2 \pm 4.9
	C, %	12.7 \pm 13.0	17.5 \pm 6.5	21 \pm 14	7.4 \pm 8.5	3.6 \pm 3.2
	N, %	1.4 \pm 1.5	1.3 \pm 0.8	1.8 \pm 1.8	0.6 \pm 0.7	0.3 \pm 0.3
	P, %	0.32 \pm 0.28	0.33 \pm 0.26	0.30 \pm 0.25	0.11 \pm 0.004	0.21 \pm 0.18
	% C _{RSM} of total C	3.5 \pm 2.4	8.4 \pm 6.7	13.2 \pm 7.9	4.9 \pm 5.0	3.1 \pm 2.2
	% P _{RSM} of total P	30.0 \pm 21.5	59.2 \pm 18.7	55.6 \pm 21.9	40.2 \pm 36.2	44.5 \pm 30.4
Summer	RSM, mg/l	10.0 \pm 4.6	7.5 \pm 2.9	10.2 \pm 3.7	5.8 \pm 1.5	3.6 \pm 2.5
	C, %	10.7 \pm 4.6	24.7 \pm 8.9	20.0 \pm 6.0	12.6 \pm 5.9	13.5 \pm 2.1
	N, %	0.9 \pm 0.3	1.9 \pm 0.6	1.6 \pm 0.7	1.2 \pm 0.6	1.2 \pm 0.2
	P, %	0.86 \pm 0.68	0.39 \pm 0.34	0.45 \pm 0.27	0.48 \pm 0.46	0.72 \pm 0.34
	% C _{RSM} of total C	10.7 \pm 10.1	15.6 \pm 4.4	21.0 \pm 4.2	12.2 \pm 5.3	5.6 \pm 3.0
	% P _{RSM} of P total	57.0 \pm 25.2	53.5 \pm 21.8	67.9 \pm 17.8	55.1 \pm 28.7	32.6 \pm 18.7
Autumn	RSM, mg/l	3.4 \pm 2.4	5.1 \pm 1.4	8.7 \pm 3.3	10.7 \pm 2.6	8.9 \pm 3.4
	C, %	11.0 \pm 6.0	25.7 \pm 8.0	17.4 \pm 6.5	13.6 \pm 6.9	7.3 \pm 3.5
	N, %	0.9 \pm 0.5	1.7 \pm 0.4	1.2 \pm 0.5	1.1 \pm 0.5	0.7 \pm 0.4
	P, %	0.93 \pm 0.64	0.33 \pm 0.15	0.57 \pm 0.21	0.70 \pm 0.45	0.30 \pm 0.21
	% C _{RSM} of total C	4.35 \pm 3.9	12.4 \pm 4.8	17.2 \pm 7.5	18.9 \pm 11.4	4.8 \pm 2.8
	% P _{RSM} of P total	42.8 \pm 32.7	71.9 \pm 9.9	82.8 \pm 11.4	76.9 \pm 14.0	40.8 \pm 8.6

838

839

840

841