

# Development of nascent autotrophic carbon fixation systems in various redox conditions of the fluid degassing in early Earth

Sergey A. Marakushev, Olga V. Belonogova

5 Institute of Problems of Chemical Physics, Russian Academy of Sciences, Academician Semenov avenue 1, Chernogolovka, Moscow region, 142432 Russian Federation

**Correspondence:** Sergey A. Marakushev ([shukaram@yandex.ru](mailto:shukaram@yandex.ru); [marak@cat.icp.ac.ru](mailto:marak@cat.icp.ac.ru))

10 **Abstract.** The origin and development of primary autotrophic metabolism on early Earth were influenced by the two main regimes of degassing of the Earth – reducing (predominance CH<sub>4</sub>) and oxidative (CO<sub>2</sub>). Among the existing theories of the autotrophic origin of life in hydrothermal environments, CO<sub>2</sub> is usually considered the carbon source for nascent autotrophic metabolism. However, the ancestral carbon used in metabolism may have been  
15 derived from CH<sub>4</sub> if the outflow of magma fluid to the surface of the Earth consisted mainly of methane. In such an environment, the primary autotrophic metabolic systems had to be methanotrophic. Due to the absence of molecular oxygen in the Archean conditions, this metabolism would have been anaerobic, i.e., oxidation of methane must be realized by inorganic high-potential electron acceptors. In light of the primacy and prevalence of CH<sub>4</sub>-  
20 dependent metabolism in hydrothermal systems of the ancient Earth, we propose a model of carbon fixation where the methane is fixed/transformed in a sequence of reactions in an autocatalytic methane-fumarate cycle. Nitrogen oxides are thermodynamically most favorable among possible oxidants of methane; however, even the activity of oxygen created by mineral buffers of iron in hydrothermal conditions is sufficient for methanotrophic acetogenesis. The  
25 hydrothermal system model is considered in the form of a phase diagram, which demonstrates the area of redox and P, T conditions favorable for the development of primary methanotrophic metabolism.

## 1 Deep methane degassing of the early Earth

30

The deep hydrocarbon generation in seismically active satellites manifested as the significant concentrations of hydrocarbons, including methane on their surface. For example, there is a prevalence of methane on Titan and Enceladus (the satellites of Saturn) (Tobie et al., 2006;

Bouquet et al., 2015) and on Europa (the satellite of Jupiter) (e.g. Russell et al., 2017).  
35 Additionally, high concentrations of methane are assumed to be present on early Mars (Oehler  
and Etiope, 2017).


On Earth, methane and other hydrocarbons are generated in magma chambers and are carried  
by fluids to the surface through volcanic processes, and are trapped in the gas-liquid  
inclusions during the minerals formation. This has been observed in the quartz–methane  
40 amygdaline inclusions that occur in the form of relics present in metamorphic-basaltic rocks  
of Greenland that were dated to 3.8 billion years (Touret, 2003). The inclusion of  
hydrocarbons and reduced organic compounds in Archean quartz (Touret, 2003, Schreiber et  
al., 2017) indicates a sufficiently reductive environment at this time. There is evidence that  
45 Archean atmosphere was enriched in hydrogen and methane (Tian et al, 2005; Zahnle et al,  
2019), but the oxidation state of magma sources apparently has changed (Aulbach et al,  
2017.). According to the trace-elements data of igneous zircons crustal origin (mainly Ce-  
based oxybarometer), it was shown that the Hadean continental crust was significantly more  
reduced than its modern counterpart and experienced progressive oxidation ~3.6 billion years  
ago (Yang et al., 2014). In addition, the oxygen activity (log units) in the earth's crust  
50 periodically changed with regards to the quartz-magnetite-fayalite redox buffer from -8 to +4  
in Hadean and from -7 to +7 in the early Archean. Significant fluctuations in the redox state  
of Archean and Hadean zircons indicate a pulsed regime of earth degassing during this period  
of time, which, in our opinion, is related to impulses in the geomagnetic field (Allredge,  
1984; Larson and Olson, 1991; Aubert et al., 2010). Thus, the evolution of the Earth over a  
55 period of 4.6 billion years is determined by the impulsive degassing of its liquid core along  
the structures of the dislocation of its solid silicate shells (mantle and crust).

Of all the magmatic formations of the world, the alkaline magmatism is the deepest and  
controlled by mantle cycles (Balashov and Glaznev, 2006; Marakushev and Marakushev,  
2010) and in its magma chambers hydrocarbon substances arises. Thermodynamic  
60 calculations show the preference of deep formation and stability of hydrocarbons, which are  
carried in fluids to the Earth surface (temperature and pressure decreasing), are transformed to  
methane (Marakushev and Marakushev, 2006). This is confirmed by the massive production  
of abiotic methane at ~40 km depth (Brovarone et al., 2017) and the discovered bubbles of  
hydrocarbons trapped in eclogite, a metamorphic rock that forms at high pressure at a depth of  
65 at least 80 km (Tao et al., 2018). The gas-liquid inclusions of methane in diamond,  
geochemical analyzes and P-T experiments on the synthesis of hydrocarbons (Scott et al.,

2004; Schulte et al., 2006; Kolesnikov, et al., 2009; Etiope and Sherwood Lollar, 2013; McCollom, 2013; Smit et al., 2016) also prove its deep origin.

The global mid-ocean ridge system represents a major site for outgassing of volatiles from the earth's mantle. Methane, which was believed to have a surface origin (low-temperature serpentinization  $\sim 100^\circ\text{C}$ ), apparently is formed at depth, at temperatures of ca.  $400^\circ\text{C}$  under redox conditions characterizing intrusive rocks derived from sub-ridge melts (Mével, 2003; Wang et al., 2018). Thus, deep, alkaline-basalt magmatism (elevated alkali content, especially  $\text{K}_2\text{O}$ ), in contrast to basalt-andesitic one, is mainly responsible for methane degassing on the earth's surface. With increasing alkalinity (alkaline slope) in the fluid inclusions of igneous rocks invariably appear with different hydrocarbons (Potter and Konnerup-Madsen, 2003; Nivin et al., 2005). The high content of potassium in the high-silica Hadean crust (Boehnke et al., 2018) indicates the depth of magmatism and its hydrocarbon specificity.

The model of two-stage development of fluids (**I**  $\leftrightarrow$  **II**) generated by the Earth's core via mantle magma chambers is presented in the phase diagram of the C–H–O compositions, Fig. 1. Fluids ejected from the liquid core were initially saturated with hydrogen, with oxygenic components being of minor importance. However, during the process of the earth's silicate shells (mantle and crust) extension (associated, in our opinion, with a Hadean to Paleoproterozoic geodynamo (Tarduno et al., 2015)), an increase of fluid permeability stimulates the selective migration of hydrogen (the most mobile component) from it. This process is responsible for hydrogen losing its leading position in ejected fluids and being fundamental to the evolution of low and normal alkalinity magmatism (Marakushev and Marakushev, 2008, 2010). In this scenario, the fractionation of chemical components in the fluid would result in rich acidic  $\text{CO}_2$  solutions (for example,  $\text{H}_2 + 2\text{CO} = \text{H}_2\text{O} + 0.5\text{CO}_2 + 1.5\text{C}$  and  $\text{H}_2\text{O} + \text{CO}_2 = \text{H}_2\text{CO}_3$ ). These solutions are widely observed in the composition of fluid inclusions in minerals of all igneous rocks of low and normal alkalinity. The water-carbonic regime of hydrogen fluid evolution is represented in the phase diagram, Fig. 1, as an area of thermodynamic stability (facies **I**) limited by C– $\text{CO}_2$ – $\text{H}_2\text{O}$  paragenesis.



95

**Figure 1.** Two regimes of evolution of the C–H–O system on the phase diagram of its compositions. Roman numerals denote various regimes of hydrogen fluid evolution: **(I)** water-carbonic and **(II)** water-hydrocarbon solutions, separated by H<sub>2</sub>O–C equilibrium (red). Parageneses (assemblages) of the initial substances (H<sub>2</sub>, CO, CO<sub>2</sub>) are denoted by dashed sub-  
 100 lines, while dash-dotted lines indicate the parageneses (C<sub>2</sub>H<sub>6</sub>–H<sub>2</sub>O, C<sub>2</sub>H<sub>4</sub>–H<sub>2</sub>O, etc) of hydrocarbons (black triangles) with water. Black squares denote organic substances within the two component (H<sub>2</sub>–CO) subsystem: methanol (1), ethylene glycol (2), acetate (3), succinate (4), pyruvate (5), and fumarate (6).

105 The transition to compression of silicate shells prevents hydrogen migration from fluids and stimulates the production of hydrocarbons within them; for example, consider the reactions:  $3\text{H}_2 + \text{CO} = \text{H}_2\text{O} + \text{CH}_4$ ,  $5\text{H}_2 + 2\text{CO} = 2\text{H}_2\text{O} + \text{C}_2\text{H}_6$  (Fig.1, facies **II**, reducing conditions). The hydrogen in the reaction like  $4\text{H}_2 + \text{H}_2\text{CO}_3 = 3\text{H}_2\text{O} + \text{CH}_4$  destroys the acid components in fluids, and this determines the alkaline slope in the development of  
 110 magmatism. This is a two-stage model of the development of the C–H–O system (**I** ↔ **II**), which depends on the composition of earth's core fluids, and their transformations in magma chambers.

The existing theories on the origin of autotrophic life mainly identify carbon dioxide as the unique carbon source for metabolism. This autotrophic metabolism should have originated  
 115 at a high partial pressure of CO<sub>2</sub> in the environment (paragenesis CO<sub>2</sub> + H<sub>2</sub>O, Fig. 1, facies **I**). We assume that in geodynamic regime **II** (CH<sub>4</sub> + H<sub>2</sub>O paragenesis), carbon ancestral

metabolism could use methane as a carbon source if the flow of free energy from the geochemical environment was coupled with biomass formation reactions. Perhaps, these different regimes of fluid degassing determined the physicochemical conditions of the ambient environment, which, in turn, provided an opportunity for the emergence and development of various systems of ancient autotrophic metabolism. In regime **II** (Fig. 1), methane and other hydrocarbons could be substrates of the emerging autotrophic metabolism.

The above geochemical and petrological data indicate highly heterogeneous redox conditions between the present-day Earth and conditions that periodically arose in the early Earth. We consider that the anaerobic reductive geochemical conditions of the Archean played a decisive role in the origin and development of carbon and energy metabolism, which were vastly different from those observed in the tops of the branches of the modern phylogenetic tree of prokaryotes. Most metabolically-anaerobic chemoautotrophic organisms are either extinct or strongly limited to narrow anoxic ecological niches. Lateral gene transfer and subsequent phylogenetic divergence erased most evolutionary information recorded in ancestral prokaryotic genomes (Martin et al., 2016).

## 2 Anaerobic oxidation of methane

The study of anaerobic oxidation of methane (AOM) in modern oxygen-free environments (marine sedimentary rocks, gas-hydrates, mud volcanoes, black smokers, hydrocarbon seeps) has increased in recent years. This direction was sparked by the discovery of anaerobic methanotrophic archaea (Hinrichs et al., 1999) and, subsequently, their structural consortia with sulfate-reducing bacteria (Knittel and Boetius, 2009). A similar consortia was later discovered in archaea species that function in chemical conjunction with the bacterium *Candidatus Methylophilus oxyfera*, which itself can independently couple AOM to denitrification (Ettwig et al., 2010; Haroon et al. 2013). Furthermore, the microbiological AOM was recently shown to be directly associated with the reduction of iron and manganese compounds and minerals (Beal, 2009; Ettwig et al., 2016; Oni and Friedrich, 2017; He et al., 2018), as, for example, in the reaction  $\text{CH}_4 + 8\text{Fe}^{3+} + 2\text{H}_2\text{O} \rightarrow \text{CO}_2 + 8\text{Fe}^{2+} + 8\text{H}^+$  ( $\Delta G^{0'} = -454$  kJ/mol  $\text{CH}_4$ ).

Recent studies have suggested that both archaea (ANME-2d) (Haroon et al., 2013) and bacteria (*Methylobacter*) (Martinez-Cruz et al., 2017), without partners, may themselves be versatile methanotrophs capable of using different oxidants as electron acceptors under different environmental conditions. AOM occur by the reversal canonical methanogenesis

pathway (Timmers et al., 2017) and, perhaps, the evolution of life periodically includes forward or reverse pathways depending on the substrate (methanogen-methanotroph “switch back” (McGlynn, 2017). For example, nickel enzyme purified from methanogenic archaea can catalyze the oxidation of methane to methyl coenzyme M ( $\text{CH}_4 + \text{CoM-S-S-CoB} \rightarrow$   
155  $\text{CH}_3\text{-S-CoM} + \text{HS-CoB}$ ;  $\Delta G^\circ = 30 \pm 10$  kJ), that is the reverse reaction of methyl coenzyme M reductase (Scheller et al., 2010). In general, methano- and methylotrophs use different but often interrelated pathways of carbon fixation (Smejkalová et al., 2010). Newly described methanotrophic anaerobic prokaryotes are frequently discovered in various extreme environmental conditions (Semrau et al., 2008), underscoring the functional and phylogenetic  
160 diversity of this group. The search for relict forms of anaerobic methanotrophic metabolism continues.

In 2013, Wolfgang Nitschke and Michael Russell described the possibility of methane assimilation as the sole source of carbon for primordial metabolism (Nitschke and Russell, 2013). They suggested that methanotrophy and not methanogenesis may have been the  
165 founding metabolism in the first protocells and presented a model of methanotrophic acetogenesis in which methane, as the carbon source, is assimilated into the biomimetic analogue of the modern reverse acetyl-CoA pathway. The proposed methane oxidant in this pathway of  $\text{CH}_4$  fixation was nitric oxide (NO), formed via nitrate/nitrite transformation (‘denitrifying methanotrophic acetogenesis’) (Russell and Nitschke, 2017). The authors  
170 consider the process of low-temperature harzburgite (ophiolites) hydrothermal serpentinization in the presence of carbon oxides served as the main source of methane. Nevertheless, Wang et al. (2018) argue that there is another unified deep high-temperature process of methane-making for these hydrothermal areas.

In the absence of oxygen, the methane oxidation requires electron acceptors with a high  
175 redox potential (such as nitrate, manganese (IV), iron (III), and sulfate). Thermodynamic calculations of anaerobic methanotrophic acetogenesis reactions in aqueous hydrothermal conditions that require oxidized compounds such as sulfur, nitrogen, and iron are considered in Table 1. For example, the free energy of the reaction  $\text{CH}_4 + 6\text{Fe}_2\text{O}_3 = 0.5\text{CH}_3\text{COOH} + \text{H}_2\text{O} + 4\text{Fe}_3\text{O}_4$  at 473 K is equal to the sum of the free energy of products  
180 formation minus the sum of free energy of the reactants formation at the same temperature ( $\Delta G_{473}^0 = (0.5\Delta G_{\text{CH}_3\text{COOH}}^0 + \Delta G_{\text{H}_2\text{O}}^0 + 4\Delta G_{\text{Fe}_3\text{O}_4}^0) - (\Delta G_{\text{CH}_4}^0 + 6\Delta G_{\text{Fe}_2\text{O}_3}^0) = -6.49$  kJ/mol).

**Table 1.** Free Gibbs energy of aqueous reactions of anaerobic methanotrophic acetogenesis at 298 and 473 K at the saturated vapor pressure ( $P_{SAT}$ ). The oxidized and reduced states of oxidant in the reaction are conditionally called the redox pairs. The oxidation of methane to fully ionized and non-ionized forms of acetate is presented. Value of  $\Delta G^0_T$  at 298 and 473 K indicates the advantage of the reactions at low (L) or high (H) temperatures. Free energies of aqueous substances formation at  $P_{SAT}$  (Amend and Shock, 2001) were used in calculations.

<b>Redox pairs of nitrogen</b>	$\Delta G^0_{298}$ kJ/mol CH <sub>4</sub>	$\Delta G^0_{473}$ kJ/mol CH <sub>4</sub>	
CH <sub>4</sub> + 2NO = 0.5CH <sub>3</sub> COOH + H <sub>2</sub> O + N <sub>2</sub>	-586.78	-563.18	L
CH <sub>4</sub> + 2NO = 0.5CH <sub>3</sub> COO <sup>-</sup> + 0,5 H <sup>+</sup> + H <sub>2</sub> O + N <sub>2</sub>	-573.39	-538.33	
CH <sub>4</sub> + 4NO = 0.5CH <sub>3</sub> COOH + H <sub>2</sub> O + 2N <sub>2</sub> O	-582.32	-535.87	L
CH <sub>4</sub> + 4NO = 0.5CH <sub>3</sub> COO <sup>-</sup> + 0,5 H <sup>+</sup> + H <sub>2</sub> O + 2N <sub>2</sub> O	-568.93	-511.02	
CH <sub>4</sub> + 4HNO <sub>2</sub> = 0.5CH <sub>3</sub> COOH + 3H <sub>2</sub> O + 4NO	-264.46	-320.79	H
CH <sub>4</sub> + 4NO <sub>2</sub> <sup>-</sup> + 3,5H <sup>+</sup> = 0.5CH <sub>3</sub> COO <sup>-</sup> + 4NO + 3H <sub>2</sub> O	-324.71	-398.7	
CH <sub>4</sub> + 2HNO <sub>3</sub> = 0.5CH <sub>3</sub> COOH + H <sub>2</sub> O + 2HNO <sub>2</sub>	-295.16	-286.35	L
CH <sub>4</sub> + 2NO <sub>3</sub> <sup>-</sup> = 0.5CH <sub>3</sub> COO <sup>-</sup> + 2NO <sub>2</sub> <sup>-</sup> + H <sub>2</sub> O + 0,5H <sup>+</sup>	-230.07	-217.58	
CH <sub>4</sub> + 1.33HNO <sub>3</sub> = 0.5CH <sub>3</sub> COOH + 1.67H <sub>2</sub> O + 1.33NO	-286.41	-299.38	H
CH <sub>4</sub> + 1.33NO <sub>3</sub> <sup>-</sup> + 0,83H <sup>+</sup> = 0.5CH <sub>3</sub> COO <sup>-</sup> + 1.33NO + 1.67H <sub>2</sub> O	-263.12	-304.34	
CH <sub>4</sub> + 0,8HNO <sub>3</sub> = 0.5CH <sub>3</sub> COOH + 1.4H <sub>2</sub> O + 0,4N <sub>2</sub>	-405.67	-403.98	L
CH <sub>4</sub> + 0,8NO <sub>3</sub> <sup>-</sup> + 0,3H <sup>+</sup> = 0.5CH <sub>3</sub> COO <sup>-</sup> + 0,4N <sub>2</sub> + 1,4H <sub>2</sub> O	-386.33	-382.11	
<b>Redox pairs of iron (mineral buffers)</b>			
CH <sub>4</sub> + 6Fe <sub>2</sub> O <sub>3</sub> = 0.5CH <sub>3</sub> COOH + H <sub>2</sub> O + 4Fe <sub>3</sub> O <sub>4</sub>	11.84	-6.49	H
CH <sub>4</sub> + 6Fe <sub>2</sub> O <sub>3</sub> = 0.5CH <sub>3</sub> COO <sup>-</sup> + 0,5 H <sup>+</sup> + H <sub>2</sub> O + 4Fe <sub>3</sub> O <sub>4</sub>	25.23	18.36	H
CH <sub>4</sub> + 1,5FeS <sub>2</sub> + 0,5Fe <sub>3</sub> O <sub>4</sub> = 0.5CH <sub>3</sub> COOH + H <sub>2</sub> O + 3FeS	44.65	28.85	H
CH <sub>4</sub> + 1,5FeS <sub>2</sub> + 0,5Fe <sub>3</sub> O <sub>4</sub> = 0.5CH <sub>3</sub> COO <sup>-</sup> + 0,5 H <sup>+</sup> + H <sub>2</sub> O + 3FeS	58.04	53.7	H
CH <sub>4</sub> + 2Fe <sub>3</sub> O <sub>4</sub> + 3SiO <sub>2</sub> = 0.5CH <sub>3</sub> COOH + H <sub>2</sub> O + 3Fe <sub>2</sub> SiO <sub>4</sub>	57.23	16.19	H
CH <sub>4</sub> + 2Fe <sub>3</sub> O <sub>4</sub> + 3SiO <sub>2</sub> = 0.5CH <sub>3</sub> COO <sup>-</sup> + 0,5 H <sup>+</sup> + H <sub>2</sub> O + 3Fe <sub>2</sub> SiO <sub>4</sub>	70.62	41.04	H
<b>Redox pair of sulphur</b>			
CH <sub>4</sub> + 0,5H <sub>2</sub> SO <sub>4</sub> = 0.5CH <sub>3</sub> COOH + H <sub>2</sub> O + 0,5H <sub>2</sub> S	-42.57	-69.93	H
CH <sub>4</sub> + 0,5SO <sub>4</sub> <sup>2-</sup> + 0,5 H <sup>+</sup> = 0.5CH <sub>3</sub> COO <sup>-</sup> + H <sub>2</sub> O + 0,5H <sub>2</sub> S	-29.18	-45.07	
<b>Carboxy-methano acetogenesis</b>			
CH <sub>4</sub> + CO <sub>2</sub> + 2NO + 2H <sub>2</sub> = CH <sub>3</sub> COOH + N <sub>2</sub> + 2H <sub>2</sub> O	-671,53	-620,83	L
CH <sub>4</sub> + HCO <sub>3</sub> <sup>-</sup> + 2NO + 2H <sub>2</sub> = CH <sub>3</sub> COO <sup>-</sup> + N <sub>2</sub> + 3H <sub>2</sub> O	-680.97	-636.35	
CH <sub>4</sub> + 0,5CO <sub>2</sub> + 6Fe <sub>2</sub> O <sub>3</sub> + H <sub>2</sub> = 0,75CH <sub>3</sub> COOH + 4Fe <sub>3</sub> O <sub>4</sub> + 1,5H <sub>2</sub> O	-30.54	-35.34	H
CH <sub>4</sub> + 0,5HCO <sub>3</sub> <sup>-</sup> + 6Fe <sub>2</sub> O <sub>3</sub> + H <sub>2</sub> = 0,75CH <sub>3</sub> COO <sup>-</sup> + 4Fe <sub>3</sub> O <sub>4</sub> + 2H <sub>2</sub> O + 0,25H <sup>+</sup>	-28.56	-30.64	
CH <sub>4</sub> + CO <sub>2</sub> = CH <sub>3</sub> COOH	24.27	39.89	L
CH <sub>4</sub> + HCO <sub>3</sub> <sup>-</sup> = CH <sub>3</sub> COO <sup>-</sup> + H <sub>2</sub> O	14.83	24.41	
<b>Carboxy- acetogenesis</b>			
CO <sub>2</sub> + 2H <sub>2</sub> = 0,5CH <sub>3</sub> COOH + H <sub>2</sub> O	-84,75	-57,65	H
HCO <sub>3</sub> <sup>-</sup> + 2H <sub>2</sub> + 0,5H <sup>+</sup> = 0,5CH <sub>3</sub> COO <sup>-</sup> + 2H <sub>2</sub> O	-107,58	-98.01	

190 It is obvious that methane oxidation with nitrogen oxide compounds is thermodynamically very favorable, compared to oxidants such as sulfate, magnetite and hematite. The acetogenesis reactions are energetically more preferable under acidic hydrothermal conditions (the protonated compounds). The change in the free energy with temperature change indicates whether the reaction displays a thermodynamic preference for  
195 low-temperature (L) or high-temperature (H) conditions. The reactions of methane with

sulfate and iron-oxides is the most thermodynamically favorable with increasing temperature (decreasing  $\Delta G^0_r$ ), whereas the reactions with nitrogen-oxides have different directions. The methane fixation is an energetically more favorable process than CO<sub>2</sub> fixation. For example, in aerobic acetogenesis ( $\text{CH}_4 + \text{O}_2 = 0,5\text{CH}_3\text{COOH} + \text{H}_2\text{O}$ ), more free energy is released in the methane fixation reaction ( $\Delta G^0_{298} = -417.35$  kJ/mol under standard conditions; aqueous constants from (Amend and Shock, 2001)) than in the process of CO<sub>2</sub> fixation ( $\text{CO}_2 + 2\text{H}_2 = 0,5\text{CH}_3\text{COOH} + \text{H}_2\text{O}$ ;  $\Delta G^0_{298} = -84.75$  kJ).

The most favorable reaction  $\text{CH}_4 + 2\text{NO} = 0,5\text{CH}_3\text{COOH} + \text{H}_2\text{O} + \text{N}_2$  (Table 1) can be represented as a model of methanotrophic acetogenesis, which is part of the reverse acetyl-CoA pathway. The second part of this path is the reaction of CO<sub>2</sub> reduction:  $\text{CO}_2 + 2\text{H}_2 = 0,5\text{CH}_3\text{COOH} + \text{H}_2\text{O}$ . In sum, this is a very thermodynamically favorable pathway of carbon fixation in the form CH<sub>4</sub> and CO<sub>2</sub>:  $\text{CH}_4 + \text{CO}_2 + 2\text{NO} + 2\text{H}_2 = \text{CH}_3\text{COOH} + \text{N}_2 + 2\text{H}_2\text{O}$ . The different stoichiometry of acetogenesis was observed in the archaean Methanosarcina acetivorans, when methane oxidation was associated with the reduction of iron (III) [Soo et al, 2016]. A reaction is proposed in which four methane molecules are oxidized and two CO<sub>2</sub> molecules are reduced to form three acetate molecules. Increasing the ratio of CH<sub>4</sub> to CO<sub>2</sub> ( $4\text{CH}_4 + 2\text{CO}_2 + 24\text{Fe}_2\text{O}_3 + 4\text{H}_2 = 3\text{CH}_3\text{COOH} + 6\text{H}_2\text{O} + 16\text{Fe}_3\text{O}_4$ ) makes the process of anaerobic acetogenesis more thermodynamically favorable (Table. 1, carboxy-methano acetogenesis).


The LUCA era apparently proceeded in an environment with high CO<sub>2</sub> partial pressure, whereas the pre-LUCA period proceeded in a reducing environment with a significant availability of methane. The question thus arises: was this ancestral reverse acetyl-CoA relic pathway the only metabolic CH<sub>4</sub> fixation system, or were there other proto-biochemical mechanisms for the assimilation of carbon?

In addition to the acetyl-CoA pathway, autocatalytic CO<sub>2</sub> fixation cycles have been suggested as the first metabolic autocatalytic systems on early Earth (Wächtershäuser, 1990; Smith and Morowitz, 2004; Marakushev and Belonogova, 2009, 2013; Fuchs, 2011; Braakman and Smith, 2012, 2013). These include an autocatalytic reductive tricarboxylic acids (rTCA) cycle (reverse citrate cycle, Arnon cycle), a 3-hydroxypropionate cycle, a 3-hydroxypropionate/4-hydroxybutyrate cycle, a reductive dicarboxylate/4-hydroxybutyrate cycle, and a reducing pentose phosphate (Calvin–Benson–Bassham) cycle. The defined sequences of intermediates of these cycles are modules such that their combination can create a variety of metabolic systems, including the specific pathways of carbon fixation (Lorenz et al., 2011; Marakushev and Belonogova, 2010, 2015; Braakman and Smith, 2012, 2013). To

230 be considered a possible metabolic alternative, assimilating methane through autocatalytic cycle intermediates must satisfy the fundamental requirements of thermodynamics.

### 3 The proposed methane-fumarate cycle

235 Based on the hypothesis of primordial anaerobic methanotrophic metabolism origin, we assume that some components and modules of the metabolic cycles (carboxylic and keto acids, and their associations (parageneses)) may also be relicts of ancient methanotrophic metabolism. One of the few known reactions of CH<sub>4</sub> fixation is the formation of 2-methylsuccinate as a result of the reaction: fumarate+CH<sub>4</sub> → 2-methylsuccinate (Thauer and Shima, 2008; Haynes and Gonzalez, 2014), and fumarate addition has been widely proposed as an initial step in the anaerobic oxidation of both aromatic and aliphatic hydrocarbons (Musat, 2015). The reaction of methane with fumarate satisfies the “minimal energy requirements” for autotrophic growth (Beasley and Nanny, 2012), and we consider the possibility of its participation in nascent autotrophic metabolism. We propose a simplified model of the methane-fumarate (MF) cycle, Fig. 2, which could have originated in the reductive Archean hydrothermal systems, at a high partial pressure of endogenous methane (facies II, Figure 1). The cycle is initiated by the reaction of fumarate + methane → 2-methylsuccinate. In the hydration and dehydrogenation or anaerobic oxidation reactions, 2-methylsuccinate is converted to citramalate, which is disproportionated to acetate and pyruvate with cleavage of a carbon-carbon bond. Pyruvate is an important “hub” metabolite that is a precursor for amino acids, carbohydrates, cofactors, and lipids in an extant metabolic network. The following carbon assimilation reaction in the form of CO<sub>2</sub> with the formation of oxaloacetate is a biomimetic analogue of the reductive tricarboxylic acid (rTCA) cycle reaction. An α-carboxylation of pyruvate is a critical anabolic pathway in modern biochemistry, which resupplies rTCA cycle intermediates. Oxaloacetate is transformed into fumarate in the reactions of the citrate cycle intermediates. The resulting fumarate assimilates methane and begins a new MF cycle, in one turnover of which an acetate molecule is formed from methane and carbon dioxide molecules: CH<sub>4</sub>+CO<sub>2</sub> = CH<sub>3</sub>COOH, Table. 1. The nonenzymatic flow of some reaction sequences of the rTCA cycle, such as oxaloacetate → malate → fumarate → succinate has been recently experimentally confirmed (Muchowska et al., 2017).



260

265

270

**Figure 2.** The scheme of the proposed methane-fumarate (MF) cycle. Carbon from methane is introduced into the fumarate (marked by red), and from  $\text{CO}_2$  into the pyruvate (marked by blue) with the formation of a C–C bond. The reaction sequence pyruvate  $\rightarrow$  oxaloacetate  $\rightarrow$  malate  $\rightarrow$  fumarate  $\rightarrow$  succinate is part of the reductive tricarboxylic acid (rTCA) cycle, an enzyme-free archaic version of which is proposed as the basis of ancient autotrophic metabolism (Wächtershäuser 1990; Smith and Morowitz 2004). The inset shows options for the oxidative transformation of 2-methylsuccinate to citramalate with hematite ( $\text{Fe}_2\text{O}_3$ ) and nitric oxide (NO) as oxidants. The chemical potential of hydrogen in the environment determines the equilibrium shift in the reactions succinate  $\leftrightarrow$  fumarate and 2-methyl succinate  $\leftrightarrow$  mesaconate.

275

Transformation of fumarate into 2-methylsuccinate introduces into the cycle five-carbon intermediates, such as citramalate and mesaconate, functioning, for example, in the reductive 3-hydroxypropionate  $\text{CO}_2$  fixation cycle. The autocatalytic nature of the cycle derives from the branching point associated with citramalate cleavage and can be shown by the example of doubling the intermediate as in the reaction:  $\text{C}_4\text{H}_6\text{O}_5$  (malate) + 1,5 $\text{CH}_4$  + 2,5 $\text{CO}_2$  = 2 $\text{C}_4\text{H}_6\text{O}_5$

(two malates). This type of autotrophic metabolism, as in the case of the acetyl-CoA pathway, can be defined as carboxy-methanotrophic acetogenesis. As it happened with the archaean WL pathway (eg, Soo et al., 2016), methanotrophy probably should be combined with the carboxylation process, therefore nature had to look for all possible sources of carbon dioxide. The problem of the most energetically unfavorable reaction of 2-methylsuccinate transformation into citramalate ( $\Delta G^0_{298} = 96.57$  kJ), Table 2a, can be solved by using oxidants, such as oxides of nitrogen and iron (Fig. 2, inset). Nitric oxide (NO) is the strongest oxidant, but the reaction with  $Fe_2O_3$  is also favorable at physiological temperatures.

**Table 2.** The free energy of reactions of MF cycle (a) and reactions of its anaerobic oxidative branch (b) with different oxidants at 298 and 473 K and  $P_{SAT}$ . For comparison, the reaction of methane oxidation with molecular oxygen is considered. Free energies of aqueous substances formation at  $P_{SAT}$  (Amend and Shock, 2001; Marakushev and Belonogova, 2012, 2013 (El. Suppl. Mat.)) were used in calculations.

<b>a. Reactions of cycle</b>	$\Delta G^0_{298}$ kJ/mol	$\Delta G^0_{473}$ kJ/mol	
1. $(CH_2)_2(COOH)_2$ ( <b>fumarate</b> ) + $CH_4 = (CH_2)(CH_3CH)(COOH)_2$ ( <b>2-methylsuccinate</b> )	-44.95	-29.97	L
2. $(CH_2)(CH_3CH)(COOH)_2 + H_2O = (CH_3CH)CH(OH)(COOH)_2$ ( <b>citramalate</b> ) + $H_2$	96.57	94.14	H
3. $(CH_3CH)CH(OH)(COOH)_2 = CH_3COOH$ ( <b>acetate</b> ) + $CH_3(CO)COOH$ ( <b>pyruvate</b> )	19.35	1.73	H
4. $CH_3(CO)COOH + CO_2 = CH_2CO(COOH)_2$ ( <b>oxaloacetate</b> )	13.11	35.03	L
5. $CH_2CO(COOH)_2 + H_2 = CH_2CH(OH)(COOH)_2$ ( <b>malate</b> )	-65.49	-55.78	L
6. $CH_2CH(OH)(COOH)_2 = (CH_2)_2(COOH)_2 + H_2O$	5.68	-5.26	H
7. $(CH_2)_2(COOH)_2$ ( <b>fumarate</b> ) + $H_2 = (CH_2)_2(COOH)_2$ ( <b>succinate</b> )	-102.24	-88.88	L
8. $(CH_3C=CH)(COOH)_2$ ( <b>mesaconate</b> ) + $H_2 = (CH_2)(CH_3CH)(COOH)_2$	-66.53	-54.85	L
<b>b. Oxidative reactions of cycle methane branch</b>	$\Delta G^0_{298}$ kJ/mol	$\Delta G^0_{473}$ kJ/mol	
$(CH_2)_2(COOH)_2$ ( <b>fumarate</b> ) + $CH_4 + H_2O = CH_3COOH$ ( <b>acetate</b> ) + $CH_3(CO)COOH$ ( <b>pyruvate</b> ) + $H_2$	70.97	65.9	H
$(CH_2)_2(COOH)_2 + CH_4 + 0.5O_2 = CH_3COOH + CH_3(CO)COOH$	-192.22	-182.53	L
$(CH_2)_2(COOH)_2 + CH_4 + Fe_3O_4 + 1.5SiO_2 = CH_3COOH + CH_3(CO)COOH + 1.5Fe_2SiO_4$	45.08	25.21	H
$(CH_2)_2(COOH)_2 + CH_4 + 3Fe_2O_3 = CH_3COOH + CH_3(CO)COOH + 2Fe_3O_4$	22.38	13.87	H
$(CH_2)_2(COOH)_2 + CH_4 + 0.75FeS_2 + 0.25Fe_3O_4 = CH_3COOH + CH_3(CO)COOH + 1.5FeS$	38.78	31.54	H
$(CH_2)_2(COOH)_2 + CH_4 + 2HNO_2 = CH_3COOH + CH_3(CO)COOH + H_2O + 2NO$	-115.77	-143.28	H
$(CH_2)_2(COOH)_2 + CH_4 + 2NO = CH_3COOH + CH_3(CO)COOH + N_2O$	-274.7	-250.83	L
$(CH_2)_2(COOH)_2 + CH_4 + NO = CH_3COOH + CH_3(CO)COOH + 0.5N_2$	-276.93	-264.48	L

The reversibility of the reactions in the citrate cycle is mainly determined by the equilibrium of fumarate+ $H_2 =$  succinate ( $\Delta G^0_{298} = -102.24$ ,  $\Delta G^0_{473} = -88.88$  kJ), Table 2a. The

change in the direction of electron flow therein is determined by the chemical potential of  
295 hydrogen (Marakushev and Belonogova, 2009), and therefore, different proto-metabolic  
cycles could be formed, for example, the oxidative citrate and reductive 3-hydroxypropionate  
cycles (succinate → fumarate), the rTCA cycle (fumarate → succinate), or the proposed CH<sub>4</sub>  
fixation cycle (succinate → fumarate → 2-methylsuccinate).

Anaerobic methane-oxidizing branch of cycle represents the transformation of fumarate  
300 to pyruvate and acetate. The free energies of these reactions with various inorganic oxidizing  
agents at 298 and 473 K are given in Table 2b. Assimilation of CH<sub>4</sub> in the oxidizing branch of  
the cycle, Fig. 2a, demonstrates highly favorable thermodynamics with all redox pairs of  
nitrogen. The same reaction with the iron redox pair becomes more favorable at an increasing  
temperature. Reactions with iron mineral buffers, Table 2a, are closer to the equilibrium state,  
305 which ultimately determines the possibilities of primordial cycle flow in the forward or  
reverse directions (development of methanogenesis or methanotrophic acetogenesis). An  
analysis of the oldest known association of microfossils suggests that methane-cycling  
methanogen-methanotroph communities were a significant component of the earth's early  
biosphere (Schopf et al., 2017). It is possible that a high methane partial pressure which  
310 existed in a geodynamic regime of high endogenous methane flow in ancient Earth, promoted  
the formation of various versions of carbon assimilating systems.

Methyl group formation by the oxidation of methane is limited by kinetics, because the  
dissociation energy of the C–H bond in methane (439 kJ) exceeds that of the X–H bond in  
other organic molecules, with the exception of the O–H bond in H<sub>2</sub>O (497 kJ) and other  
315 oxygen-derived molecular species. In the field of alkane oxidation, enzymatic metal-oxo  
species, promote C–H activation through a metallo-radical pathway. This involves hydrogen  
radical abstraction from the alkane by the oxo species, followed by a rapid rebound of the  
radical species onto the metal hydroxo intermediate (Roudesly et al., 2017). The calculation  
of the potential energy surface showed the thermodynamic possibility of anaerobic oxidation  
320 of methane via fumarate addition, in a reaction catalyzed by the glycy radical (Beasley and  
Nanny, 2012). The reaction mechanism fumarate+CH<sub>4</sub> → 2-methylsuccinate, fig. 2, seems to  
be similar to the radical mechanism of breaking the C–H bond with the formation of the C–C  
bond, catalyzed by benzylsuccinate synthase (Buckel and Golding, 2006; Austin et al., 2011)  
during microbiological fixation of toluene by fumarate. Radicals of amino acids and  
325 dipeptides may be the possible catalysts of methane activation with the formation of methyl  
radical as an attacking agent. The formation of pyruvate and oxaloacetate in MF cycle, Fig. 2,  
indicates a very likely formation of amino acids in simple aqueous synthesis, for example:

330  $C_3H_4O_3$  (pyruvate)+ $NH_3 = C_3H_7O_3N$  (serine),  $\Delta G^0_{298} = -10,10$  kJ or pyruvate+ $NH_3+H_2 = C_3H_7NO_2$ (alanine)+ $H_2O$ ,  $\Delta G^0_{298} = -124,8$  kJ. Barge et al., 2019 show that pyruvate can form the alanine in hydrothermal systems in the presence of mixed-valence iron oxyhydroxides. Moreover, the generation of reactive oxygen species  $H_2O_2$  and  $OH^\bullet$  from minerals and  $H_2O$  in anaerobic environments (eg. Xian et al., 2019) creates the possibility of various radical mechanisms for the oxidation of substrates in a hydrothermal environment. According to (Weiss et al., 2016), LUCA metabolism had an excess of radical reaction mechanisms, which, 335 in our opinion, could also participate in the reaction of  $CH_4$  fixation in the cycle, overcoming the activation barriers of kinetically unfavorable reaction steps.


Our understanding of the emergence of methanotrophic metabolism is within the framework of the hydrothermal theory for the origin of life (eg, Martin et al., 2008) with all its advantages (continuous flow of energy and matter, the temperature gradient, great 340 possibilities of homogeneous and heterogeneous metal catalysis). Before the occurrence of cellular structures, the primary autotrophic metabolism on the surface of minerals created the chemical space of competing autocatalytic carbon fixation cycles. The accumulation of “biomass” probably led to the emergence of heterotrophic protometabolism and the creation of a certain matrix of the organo-mineral system in which a cascade of proto-biochemical 345 redox reactions could occur, such as in the modern soils (K eraval et al., 2016). Regardless of the specific mechanism of the functioning of the precellular autotrophic metabolism (“reductive surface pyrite world” (W achtersh user, 1988), “hydrothermal reactor” (Russell and Martin, 2004), “organo-mineral matrix” (K eraval et al., 2016), and others) its origin and development was subject to the laws of aqueous thermodynamics.

350

#### **4 Anaerobic methane oxidation in the hydrothermal systems**

We represent the hydrothermal system in the form of a phase diagram which displays the chemical potential of oxygen vs. temperature at saturated vapor pressure ( $P_{SAT}$ ), where 355 temperatures and pressures are below critical thresholds (647,3 K and 22,1 MPa) (Fig. 3). The chemical potentials ( $\mu_i$ ) of components representing its partial energy, the value  $\mu_i$  is expressed through activity  $\alpha_i$  or fugacity  $f_i$  as follows:  $\mu_i = (\mu^0_i)_{T,p} + RT \ln \alpha_i = (\mu^0_i)_{T,p} + RT \ln f_i$ . Here numerical values depend on conventional standard states. For activity, the state of pure crystalline substance or unit molal concentration is usually considered as a standard state at a 360 given temperature and pressure. In this state  $\alpha_i = 1$  and, hence,  $\mu_i = (\mu^0_i)_{T,p}$ . The diagram is a two-component system (extensive parameters: C and H), since oxygen become intensive

parameters, like the temperature, and pressure. Oxygen is represented by the chemical potential of  $O_2$  in hydrothermal solution ( $\mu^P_{O_2} = RT \ln a_{O_2}$ , where  $a_{O_2}$  denotes the chemical activity of oxygen). According to the Gibbs' phase rule, at arbitrary pressure, the nonvariant equilibria in the diagram (points) consist of four phases, and the three-phase equilibria (lines) divide the divariant stability fields (facies) of the two-phase equilibria.



**Figure 3.** Diagram of the chemical potential of oxygen ( $\mu^P_{O_2}$ ) - temperature (T, K) at saturated vapor pressure ( $P_{SAT}$ ). The areas of thermodynamic stability of substances and their parageneses were calculated according to the method described in Korzhinskii, 1959 and Marakushev and Belonogova (2009). Points (indicated by temperature values) and lines represent four-phase and three-phase equilibria, separating the two-phase divariant fields of substance stabilities. The bold black line represents the equilibrium of  $CO_2 \leftrightarrow CH_4$  and separates their areas of thermodynamic stability (I and II). The dashed lines are equilibria of mineral buffers: hematite-magnetite,  $Fe_2O_3/Fe_3O_4$  (HM), pyrite-pyrrhotite-magnetite,  $FeS_2+Fe_3O_4/FeS$  (PPM), and quartz-magnetite-fayalite,  $SiO_2+Fe_3O_4/Fe_2SiO_4$  (QMF). The isolines of the activity of  $O_2$  ( $10^{-n}$  M, dot lines) are drawn. The acetate and succinate facies contoured with red and blue equilibria, respectively. We provide only two linear diagrams of the two-component C-H system in the  $CO_2$  and  $CH_4$  facies. The transition between this facies with the change of chemical potential of oxygen is indicated by a green arrow. Abbreviation: Succinate – Suc, Fumarate – Fum, Acetate –Acet.

The equilibrium  $\text{CH}_4 + 2\text{O}_2 = \text{CO}_2 + 2\text{H}_2\text{O}$  (bold black line) divides the diagram into the facies of  $\text{CO}_2$  (**I**) and  $\text{CH}_4$  (**II**) (oxic and anoxic areas of the hydrothermal system) and illustrates the two main possibilities for the development of the C–H–O system in the facies carbon dioxide or methane. Intermediates of the MF cycle are acetate, succinate, and fumarate, and we considered their metastable equilibria and parageneses. In all phase space under consideration, there are fumarate facies. The equilibrium of  $5\text{Fum} + 4\text{CH}_4 + 2\text{O}_2 = 6\text{Suc}$  at low-temperature (Fig. 3), is located in the region of very low partial pressures of oxygen, whereas the equilibrium of  $\text{Fum} + 2\text{CH}_4 + \text{O}_2 = 3\text{Acet}$  at high-temperature occurs in facies of high pressures. Acetate and succinate facies (contoured with red and blue equilibria, respectively) completely encompass the equilibrium of  $\text{CH}_4 + \text{O}_2 = \text{CO}_2 + \text{H}_2\text{O}$ . That is, in hydrothermal solution, the parageneses of some components within the MF cycle are stable in both the  $\text{CO}_2$  and the  $\text{CH}_4$  facies. The whole system can develop in either of two directions as the chemical potential of oxygen changes: 1. the formation of low-temperature (Suc- $\text{H}_2\text{O}$ ) and high-temperature (Fum- $\text{H}_2\text{O}$ ) paragenesis in  $\text{CO}_2$  facies (**I**) and 2. the formation of low-temperature (Suc- $\text{CH}_4$ ) and high-temperature (Fum- $\text{CH}_4$ ) paragenesis in  $\text{CH}_4$  facies (**II**). Thus, within these facies, protobiochemical systems supporting carbon fixation in the form of  $\text{CO}_2$  or  $\text{CH}_4$  can develop, and methane facies (**II**) represent a broad area of  $\text{CH}_4$  assimilation by carboxylic acids in an aqueous environment. The high stability of the succinate-fumarate-acetate paragenesis in hydrothermal systems at  $200^\circ\text{C}$  (473 K) was experimentally shown (Estrada et al., 2017).

Mineral buffers up to 549 K are located in the facies of succinate, but the equilibrium of HM remains in the area of thermodynamic  $\text{CO}_2$  stability (facies **I**), and PPM and QMF equilibria occur in methane facies **II** and intersect the fundamental equilibrium of  $2\text{Suc} + 2\text{CH}_4 + \text{O}_2 = 5\text{Acet}$ . Magnetite ( $\text{Fe}_3\text{O}_4$ ) facies (between HM and QMF equilibria) encompass  $\text{CH}_4/\text{CO}_2$  equilibrium in nearly the entire temperature range of the hydrothermal system under consideration. Thus, the redox areas of magnetite stability correspond to the formation conditions both  $\text{CO}_2$  and  $\text{CH}_4$  assimilating systems. The presence of magnetite in the early Archean ocean was shown by (Li et al., 2017]. Shibuya et al. (2016) also conclude that iron redox reactions probably played an important role in the early evolution of methanotrophic metabolisms in the Hadean alkaline hydrothermal system. The QMF buffer equilibrium is completely located in the methane facies (**I**), which, according to data (Yang et al., 2014), corresponds on average to the redox conditions of Hadean mantle and crust. Up to the 3.6 billion years ago and maybe even to the great oxidative event of 2.2-2.4 billion years

ago on, the earth's surface the oxidation potential of the magnetite redox pairs, apparently, determined the chemical potential of environmental oxygen.

## 5 Conclusion

420

The cyclic planetary fluid flows (outgassing of volatiles from mantle) drive earth's chemical evolution, leading to the formation of different geobiochemical systems of carbon fixation. Impulses of CO<sub>2</sub> and CH<sub>4</sub> degassing on our planet must have determined the preference of specific autotrophic carbon fixation metabolism development. It is widely accepted that autotrophic metabolism is the fixation of inorganic carbon solely in the form of CO<sub>2</sub>, but the origin of methane, both on the ancient Earth, and on the planets and satellites (for example, on Titan) is clearly inorganic; therefore, carbon fixation from methane is also a manifestation of autotrophic metabolism (formation of organic compounds from inorganic precursors).

425

The variety of modern autotrophic carbon fixation seems to be created by the association of the different metabolic associations and modules that, apparently, could function in the ancestral systems of the anaerobic fixation of CH<sub>4</sub>. When approximately ~ 3.6 billion years (Yang et al., 2014), a CO<sub>2</sub> degassing regime became dominant on our planet, the relic methanotrophy systems were forced to die out or be thrown into unusual and extreme ecological niches. If we consider LUCA as a relatively recent player in the evolution of life (Cornish-Bowden and Cárdenas, 2017), the ancestral metabolic systems of carbon fixation in putative pre-LUCA could differ appreciably from modern ones.

430

435

In the process of development of CO<sub>2</sub> fixation systems on the Earth, the main problem was the presence of electron donors (therefore, evolution created selective reducing agents: NADH, NADPH, FADH), whereas the fixation of CH<sub>4</sub> essentially depended on the presence of electron acceptors. The oxygen-containing nitrogen compounds are the best oxidants in the hydrothermal systems, but their presence there is unlikely. Nevertheless, the redox pairs of hematite-magnetite and quartz-magnetite-fayalite create a specific area of chemical potential of oxygen that satisfies the thermodynamic requirements of oxidation and assimilation of methane by protometabolic pathways. Hydrothermal systems of ancient Earth may have been very similar to those that currently exist on some extraterrestrial cosmic bodies, such as Europa or Enceladus. The degassing of these cosmic bodies can currently support methane metabolism, but the problem is to know if there are electron acceptors there (Russell et al., 2017).

440

445

450 **Author Disclosure Statement**

No competing financial interests exist.

**References**

- 455 Alldredge, L.R.: A discussion of impulses and jerks in the geomagnetic field, *J. Geophys. Res.*, 89, 4403–4412, doi: 10.1029/JB089iB06p04403, 1984.
- Amend, J.P., and Shock, E.L.: Energetics of overall metabolic reactions of thermophilic and hyperthermophilic Archaea and Bacteria, *FEMS Microbiol. Rev.*, 25, 175–243, doi: 10.1016/S0168-6445(00)00062-0, 2001.
- 460 Aubert, J., Tarduno, J.A., and Johnson, C.L.: Observations and models of the long-term evolution of earth's magnetic field, *Space Sci. Rev.*, 155, 337–370, doi: 10.1007/s11214-010-9684-5, 2010.
- Aulbach, S., Woodland, A. B., Vasilyev, P., Galvez, M. E. and Viljoen K. S.: Effects of low-pressure igneous processes and subduction on Fe<sup>3+</sup>/ZFe and redox state of mantle eclogites from Lace (Kaarvaal craton), *Earth Planet. Sci. Lett.*, 474, 283–295, 2017.
- 465 Austin, R.N., and Groves, J.T.: Alkane-oxidizing metallo enzymes in the carbon cycle, *Metallomics*, 3, 775–787, doi: 10.1039/c1mt00048a, 2011.
- Balashov, Y.A., and Glaznev, V.N.: Cycles of alkaline magmatism, *Geochem. Int.*, 44, 274–285, doi: 10.1134/S0016702906030050, 2006.
- 470 Barge, L.M., Flores, E., Baum, M.M., VanderVelde, D.G. and Russell, M.J.: Redox and pH gradients drive amino acid synthesis in iron oxyhydroxide mineral systems, *PNAS USA*, doi: 10.1073/pnas.1812098116, 2019.
- Beal, E.J., House, C.H., and Orphan, V.J.: Manganese- and iron-dependent marine methane oxidation, *Science*, 325, 184–187, doi: 10.1126/science.1169984, 2009.
- 475 Beasley, K.K., and Nanny M.A.: Potential energy surface for anaerobic oxidation of methane via fumarate addition, *Environ. Sci. Technol.*, 46, 8244–8252, doi: 10.1021/es3009503, 2012.
- Boehnke, P., Bell, E.A., Stephana, T., Trappitscha, R., Keller, C. B., Pardo, O.S., Davis, A.M., Harrison, T. M. and Pellina, M.: Potassic, high-silica Hadean crust, *Proc. Natl. Acad. Sci.* 115, 6353–6356, doi: 10.1073/pnas.1720880115, 2018.
- 480 Bouquet, A., Mousis, O., Waite, J.H., and Picaud, S.: Possible evidence for a methane source in Enceladus' ocean, *Geophys. Res. Lett.*, 42, 1334–1339, doi: 10.1002/2014GL063013, 2015.

- 485 Braakman, R., and Smith, E.: The emergence and early evolution of biological carbon-  
fixation, *PLOS Comput. Biol.*, 8, 1–16, doi: 10.1371/journal.pcbi.1002455, 2012.
- Braakman, R., and Smith, E.: The compositional and evolutionary logic of metabolism, *Phys.  
Biol.*, 10, 1–63, doi: 10.1088/1478-3975/10/1/011001, 2013.
- 490 Brovarone, A.V., Martinez I., Elmaleh A., Compagnoni, R., Chaduteau, C., Ferraris, C., and  
Esteve, I.: Massive production of abiotic methane during subduction evidenced in  
metamorphosed ophicarbonates from the Italian Alps, *Nat. Commun.*, 8, 14134, doi:  
10.1038/ncomms14134, 2017.
- Buckel, W., and Golding, B.T.: Radical enzymes in anaerobes, *Ann. Rev. Microbiol.*, 60, 27-  
49, doi: 10.1146/annurev.micro.60.080805.142216, 2006.
- Cornish-Bowden, A., and Cárdenas, M.L.: “Life Before LUCA,” *J. Theor. Biol.*, 434, 68–74,  
495 doi: 10.1016/j.jtbi.2017.05.023, 2017.
- Estrada, C.F., Mamajanov, I., Hao, J., Sverjensky, D.A., Cody, G.D., and Hazen, R.M.:  
Aspartate transformation at 200 °C with brucite [Mg(OH)<sub>2</sub>], NH<sub>3</sub>, and H<sub>2</sub>: implications  
for prebiotic molecules in hydrothermal systems, *Chem. Geol.*, 457, 162–172, doi:  
10.1016/j.chemgeo.2017.03.025, 2017.
- 500 Etiope, G., and Sherwood Lollar, B.: Abiotic methane on Earth, *Rev. Geophys.*, 51, 276–299,  
doi:10.1002/rog.20011, 2013 and references therein.
- Ettwig, K.F., Butler, M.K., Le Paslier, D., Pelletier, E., Mangenot, S., Kuypers, M.M.,  
Schreiber, F., Dutilh, B.E., Zedelius, J., de Beer, D., Gloerich, J., Wessels HJ, van Alen  
T, Luesken F, Wu ML, van de Pas-Schoonen KT, Op den Camp HJ, Janssen-Megens,  
505 E.M., Francoijs, K.J., Stunnenberg, H., Weissenbach, J., Jetten, M.S., and Strous, M.:  
Nitrite-driven anaerobic methane oxidation by oxygenic bacteria, *Nature*, 464, 543–548,  
doi: 10.1038/nature08883, 2010.
- Ettwig, K.F., Zhu, B., Speth, D., Keltjens, J.T., Jetten, M.S.M., and Kartal, B.: Archaea  
catalyze iron-dependent anaerobic oxidation of methane, *Proc. Natl. Acad. Sci.*, 113,  
510 12792–12796, doi: 10.1073/pnas.1609534113, 2016.
- Fuchs, G.: Alternative pathways of carbon dioxide fixation: Insights into the early evolution  
of life? *Ann. Rev. Microbiol.*, 65, 631–658, doi: 10.1146/annurev-micro-090110-  
102801, 2011.
- 515 Haroon, M.F., Hu, S., Shi, Y., Imelfort, M., Keller, J., Hugenholtz, P., Yuan, Z., and Tyson,  
G.W.: Anaerobic oxidation of methane coupled to nitrate reduction in a novel archaeal  
lineage, *Nature*, 500, 567–570. DOI: 10.1038/nature12375, 2013.

- Haynes, C.A., and Gonzalez, R.: Rethinking biological activation of methane and conversion to liquid fuels, *Nature Chem. Biol.*, 10, 331–339, doi: 10.1038/nchembio.1509, 2014.
- 520 He, Z., Zhang, Q., Feng, Y., Luo, H., Pan, X., and Gadd, G.M.: Microbiological and environmental significance of metal-dependent anaerobic oxidation of methane, *Sci. Total Environ.*, 610–611, 759–768, doi: 10.1016/j.scitotenv.2017.08.140, 2018.
- Hinrichs, K.U., Hayes, J.M., Sylva, S.P., Brewer, P.G., and DeLong, E.F.: Methane-consuming archaeobacteria in marine sediments, *Nature*, 398, 802–805. doi :10.1016/S0146-6380(00)00106-6, 1999.
- 525 Kéraval, B., Lehours, A.C., Colombet, J., Amblard, C., Alvarez, G. and Fontaine, S.: Soil carbon dioxide emissions controlled by an extracellular oxidative metabolism identifiable by its isotope signature, *Biogeosciences*, 13, 6353–6362, doi:10.5194/bg-13-6353-2016, 2016.
- 530 Knittel, K., and Boetius, A.: Anaerobic oxidation of methane: progress with an unknown process. *Ann. Rev. Microbiol.*, 63, 311–334, doi: 10.1146/annurev.micro.61.080706.093130, 2009.
- Kolesnikov, A., Kutcherov, V. G., and Goncharov, A. F.: Methane-derived hydrocarbons produced under upper-mantle conditions, *Nat. Geosci.* 2, 566–570, doi:10.1038/ngeo591, 2009.
- 535 Korzhinskii, D.S.: Physicochemical basis of the analysis of the paragenesis of minerals, Consultants Bureau, Inc. (New York), and Chapman & Hall (London), 1959.
- Larson, R.L., and Olson, P.: Mantle plumes control magnetic reversal frequency, *Earth Planet. Sci. Lett.*, 107, 437–447, doi: 10.1016/0012-821X(91)90091-U, 1991.
- 540 Li, Yi-L., Konhauser, K.O., and Zhai, M.: The formation of magnetite in the early Archean oceans, *Earth Planet. Sci. Lett.*, 466, 103–114, doi: 10.1016/j.epsl.2017.03.013, 2017.
- Lorenz, D.M., Jeng, A., and Deem, M.W.: The emergence of modularity in biological systems, *Phys. Life Rev.*, 8, 129–160, doi: 10.1016/j.pprev.2011.02.003, 2011.
- 545 Marakushev, A.A., and Marakushev, S.A.: PT facies of elementary, hydrocarbon, and organic substances, *Dokl. Earth Sci.*, 406, 141–147, doi: 10.1134/S1028334X0601034X, 2006.
- Marakushev, A.A., and Marakushev, S.A.: Formation of oil and gas fields, *Lithol. Miner. Resour.*, 43, 454–469, doi: 10.1134/S0024490208050039, 2008.

- Marakushev, A.A., and Marakushev, S.A.: Fluid evolution of the Earth and origin of the biosphere, in: *Man and the Geosphere*, edited by: Florinsky, I.V., Nova Science Publishers Inc, New York, 3–31, 2010.
- 550 Marakushev, S.A., and Belonogova, O.V.: The parageneses thermodynamic analysis of chemoautotrophic CO<sub>2</sub> fixation archaic cycle components, their stability and self-organization in hydrothermal systems, *J. Theor. Biol.*, 257, 588–597, doi: 10.1016/j.jtbi.2008.11.032, 2009.
- 555 Marakushev, S.A., and Belonogova, O.V.: Metabolic design and biomimetic catalysis of the archaic chemoautotrophic CO<sub>2</sub> fixation cycle, *Mos. Univer. Chem. Bull.*, 65, 212–218, doi: 10.3103/S0027131410030211, 2010.
- Marakushev, S.A., and Belonogova, O.V.: Thermodynamic factors of natural selection in autocatalytic chemical systems, *Dokl. Biochem. Biophys.*, 444, 131–136, doi: 560 10.1134/S1607672912030015, 2012.
- Marakushev, S.A., and Belonogova, O.V.: The divergence and natural selection of autocatalytic primordial metabolic systems, *Orig. Life Evol. Biosph.*, 43, 263–281, doi: 10.1007/s11084-013-9340-7, 2013.
- Marakushev, S.A., and Belonogova, O.V.: The Chemical Potentials of Hydrothermal Systems and the Formation of Coupled Modular Metabolic Pathways, *Biophysics*, 60, 542–552, 565 doi: 10.1134/S0006350915040168, 2015.
- Martin, W., Baross, J., Kelley, D., and Russell, M.J.: Hydrothermal vents and the origin of life, *Nat. Rev. Microbiol.*, 6, 805–814, doi: 10.1038/nrmicro1991, 2008.
- Martin, W.F., Weiss, M.C., Neukirchen, S., Nelson-Sathi, S., and Sousa, F.L.: Physiology, 570 phylogeny, and LUCA, *Microbial Cell*, 3, 582–587, doi: 10.15698/mic2016.12.545, 2016.
- Martinez-Cruz, K., Leewis, M.-C., Herriott, I.C., Sepulveda-Jauregui, A., Anthony, K.W., Thalasso, F., and Leigh, M.B.: Anaerobic oxidation of methane by aerobic methanotrophs in sub-Arctic lake sediments, *Sci. Total Environ.*, 607–608, 23-31, doi: 575 10.1016/j.scitotenv.2017.06.187, 2017.
- McCollom, T. M.: Laboratory simulations of abiotic hydrocarbon formation in Earth’s deep subsurface, *Rev. Mineral. Geochem.*, 75, 467–494, doi: 10.2138/rmg.2013.75.15, 2013.
- McGlynn, S.E.: Energy metabolism during anaerobic methane oxidation in ANME Archaea, *Microbes and Environments*, 32, 5-13, doi: 10.1264/jsme2.ME16166, 2017.**
- 580 Mével, C.: Serpentinization of abyssal peridotites at mid-ocean ridges, *(C. R.) Geosci.*, 335, 825–852, doi: 10.1016/j.crte.2003.08.006, 2003.

- Muchowska, K. B., Varma, S.J., Chevallot-Beroux, E., Lethuillier-Karl, L., Li G., and Moran, J.: Metals promote sequences of the reverse Krebs cycle, *Nat. Ecol. Evol.*, 1, 1716–1721, doi: 10.1038/s41559-017-0311-7, 2017.
- 585 Musat, F.: The anaerobic degradation of gaseous, nonmethane alkanes - From in situ processes to microorganisms, *Comput. Struct. Biotechn. J.*, 13 222–228, doi: 10.1016/j.csbj.2015.03.002, 2015.
- Nitschke, W., and Russell, M.J.: Beating the acetyl-CoA pathway to the origin of life, *Phil. Trans. Royal Soc. London, Series B*, 368, 20120258, doi: 10.1098/rstb.2012.0258,  
590 2013.
- Nivin, V.A., Treloar, P.J., Konopleva, N.G., and Ikorsky, S.V.: A review of the occurrence, form and origin of C-bearing species in the Khibiny alkaline igneous complex, Kola Peninsula, NW Russia, *Lithos*, 85, 93–112, doi: 10.1016/j.lithos.2005.03.021, 2005.
- Oehler, D.Z., and Etiope, G.: Methane seepage on Mars: where to look and why, *Astrobiol.*,  
595 17, 1233–1264, doi: 10.1089/ast.2017.1657, 2017.
- Oni, O.E. and Friedrich, M.W.: Metal oxide reduction linked to anaerobic methane oxidation, *Trends Microbiol.*, 25, 88–90, doi: 10.1016/j.tim.2016.12.001, 2017.
- Potter, J., and Konnerup-Madsen, J.: A review of the occurrence and origin of abiogenic hydrocarbons in igneous rocks, in: *Hydrocarbons in Crystalline Rocks*, edited by: Petford, N., and McCaffrey, K.J.W., Special Publications, 214, Geological Society,  
600 London, 151–173, doi:10.1144/GSL.SP.2003.214.01.10, 2003.
- Roudesly, F., Oble, J., and Poli, G.: Metal-catalyzed C H activation/functionalization: The fundamentals, *J. Molec. Cat. A: Chem.*, 426, 275–296, doi: 10.1016/j.molcata.2016.06.020, 2017.
- 605 Russell, M.J., and Martin, W.: The rocky roots of the acetyl-CoA pathway, *Trends Biochem. Sci.* 29, 358-363, doi: 10.1016/j.tibs.2004.05.007, 2004.
- Russell, M.J., Murray, A.E., and Kevin, P.H.: The possible emergence of life and differentiation of a shallow biosphere on irradiated icy worlds: The example of Europa, *Astrobiol.*, 17, 1265–1273, doi: 10.1089/ast.2016.1600, 2017.
- 610 Russell, M.J., and Nitschke, W.: Methane: Fuel or Exhaust at the Emergence of Life? *Astrobiol.*, 17, 1053–1066, doi: 10.1089/ast.2016.1599, 2017.
- Schopf, J.W., Kitajima, K., Spicuzza M.J., Kudryavtsev A.B., and Valley, J.W.: SIMS analyses of the oldest known assemblage of microfossils document their taxon-correlated carbon isotope compositions, *Proc. Natl. Acad. Sci. USA*, 115, 53–58, doi:  
615 10.1073/pnas.1718063115, 2017.

- Scheller, S., Goenrich, M., Boecher, R., Thauer, R.K., and Jaun, B.: The key nickel enzyme of methanogenesis catalyses the anaerobic oxidation of methane, *Nature*, 465, 606–609, doi: 10.1038/nature09015, 2010.
- Schreiber, U., Mayer, C., Schmitz, O.J., Rosendahl, P., Bronja A, Greule, M., Keppler, F., I. Mulder, I., Sattler, T., and Schöler, H.F.: Organic compounds in fluid inclusions of Archean quartz—Analogues of prebiotic chemistry on early Earth, *PLoS ONE*, 12, e0177570, doi.org/10.1371/journal.pone.0177570, 2017.
- 620
- Scott, H. P., Hemley, R. J., and Mao, H.: Generation of methane in the Earth’s mantle: in situ high pressure–temperature measurements of carbonate reduction, *Proc. Natl Acad. Sci. USA*, 101, 14023–14026, doi: 10.1073/pnas.0405930101, 2004.
- 625
- Semrau, J.D., DiSpirito, A.A., and Murrell, J.C.: Life in the extreme: thermoacidophilic methanotrophy, *Trends Microbiol.*, 16, 190–193, doi: 10.1016/j.tim.2008.02.004, 2008.
- Shibuya, T., Russell, M.J., and Takai, K.: Free energy distribution and hydrothermal mineral precipitation in Hadean submarine alkaline vent systems: Importance of iron redox reactions under anoxic conditions, *Geochim. Cosmochim. Acta*, 175, 1–19, doi: 10.1016/j.gca.2015.11.021, 2016.
- 630
- Schulte, M., Blake, D., Hoehler, T., and McCollom, T.: Serpentinization and its implications for life on the early Earth and Mars, *Astrobiology* 6, 364–376, doi:10.1089/ast.2006.6.364, 2006.
- 635
- Smejkalová, H., Erb, T.J., and Fuchs, G.: Methanol assimilation in *Methylobacterium extorquens* AM1: demonstration of all enzymes and their regulation, *PLoS ONE*, 5, e13001, doi: 10.1371/journal.pone.0013001, 2010.
- Smith, E. and Morowitz, H.G.: Universality in intermediary metabolism, *Proc. Natl. Acad. Sci. USA*, 101, 13168–13173, doi: 10.1073/pnas.0404922101, 2004.
- 640
- Smit, K.V., Shirey, S.B., Stern, R.A., Steele, A., and Wang, W.: Diamond growth from C–H–N–O recycled fluids in the lithosphere: Evidence from CH<sub>4</sub> micro-inclusions and  $\delta^{13}\text{C}$ – $\delta^{15}\text{N}$ –N content in Marange mixed-habit diamonds, *Lithos*, 265, 68–81, doi:10.1016/j.lithos.2016.03.015, 2016.
- Soo, V.W.C., McAnulty, M.J., Tripathi, A., Zhu, F., Zhang, L., Smith, P.B., Hatzakis, E., Agrawal, S., Nazem-Bokaei, H., Gopalakrishnan, S., Salis, H.S., Ferry, J.G., Maranas, C.D., Patterson, A.D., and Wood, T.K.: Reversing methanogenesis to capture methane for liquid biofuel precursors, *Microb. Cell Factories*, 15, 11–25, doi: 10.1186/s12934-015-0397-z, 2016.
- 645

- 650 Tao, R., Zhang, L., Tian, M., Zhu, J., Liu, X., Liu, J., Höfer, H.E., Stagno, V., and Fei, Y.:  
Formation of abiotic hydrocarbon from reduction of carbonate in subduction zones:  
Constraints from petrological observation and experimental simulation, *Geochim.  
Cosmochim. Acta*, 239, 390-408, doi: 10.1016/j.gca.2018.08.008, 2018.
- 655 Tarduno, J.A., Cottrell, R.D., Davis, W.J., Nimmo, F. and Bono, R.K.: A Hadean to  
Paleoarchean geodynamo recorded by single zircon crystals, *Science*, 349, 521-524,  
doi: 10.1126/science.aaa9114, 2015.
- Thauer, R.K., and Shima, S.: Methane as fuel for anaerobic microorganisms, *Ann. NY Acad.  
Sci.*, 1125, 158–170, doi: 10.1196/annals.1419.000 90, 2008.
- 660 Timmers, P.H.A., Welte, C.U., Koehorst, J.J., Plugge, C.M., Jetten, M.S.M., and Stams,  
A.J.M.: Reverse methanogenesis and respiration in methanotrophic archaea, *Archaea*,  
2017, 1–22, doi: 10.1155/2017/1654237, 2017.
- Tian, F., Toon, O. B., Pavlov, A. A. and De Sterck, H.: A hydrogen-rich early Earth  
atmosphere, *Science*, 308, 1014–1017, doi: 10.1126/science.1106983, 2005.
- Tobie, G., Lunine, J.I., and Sotin, C.: Episodic outgassing as the origin of atmospheric  
methane on Titan, *Nature*, 440, 61–64, doi: 10.1038/nature04497, 2006.
- 665 Touret, J.L.R.: Remnants of early Archaean hydrothermal methane and brines in pillow-  
breccia from the Isua-Greenstone Belt, West Greenland, *Precambrian Res.*, 126, 219–  
233, doi: 10.1016/S0301-9268(03)00096-2, 2003.
- Wächtershäuser, G.: Before enzymes and templates: theory of surface metabolism, *Microbiol.  
Rev.* 52, 452-484, doi:10.1101/cshperspect.a002162, 1988.
- 670 Wächtershäuser, G.: Evolution of the first metabolic cycles, *Proc. Natl. Acad. Sci. USA*, 87,  
200–204, doi: 10.1073/pnas.87.1.200, 1990.
- Wang, D.T., Reeves, E.P., McDermott, J.M., Seewald, J.S., and Ono, S.: Clumped  
isotopologue constraints on the origin of methane at seafloor hot springs, *Geochim  
Cosmochim Acta*, 223, 141–158, doi: 10.1016/j.gca.2017.11.030, 2018.
- 675 Weiss, M.C., Sousa, F.L., Mrnjavac, N., Neukirchen, S., Roettger, M., Nelson-Sathi, S., and  
Martin W.F.: The physiology and habitat of the last universal common ancestor, *Nat.  
Microbiol.*, 1, 1–8 , doi: 10.1038/NMICROBIOL.2016.116, 2016.
- 680 Xian, H., Zhu, J., Tan W., Tang, H., Liu, P., Zhu, R., Liang, X., Wei, J,b, He, H., and Teng,  
H. H.: The mechanism of defect induced hydroxylation on pyrite surfaces and  
implications for hydroxyl radical generation in prebiotic chemistry, *Geochim.  
Cosmochim. Acta*, 244, 163–172, doi: 10.1016/j.gca.2018.10.009, 2019.

- Yang, X., Gaillard, F., and Scaillet, B.: A relatively reduced Hadean continental crust and implications for the early atmosphere and crustal rheology, *Earth Planet. Sci. Lett.*, 393, 210–219, doi:10.1016/j.epsl.2014.02.056, 2014.
- 685 Zahnle, K.J., Gacesa, M., and Catling, D.C.: Strange messenger: A new history of hydrogen on Earth, as told by Xenon, *Geochim. Cosmochim. Acta* 244, 56–85, doi: 10.1016/j.gca.2018.09.017, 2019.