

1 **MS No.:** bg-2018-30  
2 **Title:** Basin-scale variability of microbial methanol uptake in the Atlantic Ocean  
3 **Author(s):** Stephanie L. Sargeant et al.  
4 **MS Type:** Research article  
5

6 Dear Silvio Pantoja,

7 Please see below for a point by point reply to all the reviewer's comments and your own minor  
8 corrections. We also include a revised manuscript showing all the track changes. Please note that the  
9 references have been amended to comply with Biogeosciences format in the finalised version only.

10 Yours sincerely,

11 Joanna Dixon, Stephanie Sargeant & Colin Murrell

12

13 **Response to Reviewers Comments** (Authors responses are in bold)

14 **L. Chistoserdova (milachis@uw.edu)**

15 Received and published: 8 March 2018

16 This study provides an inventory of measurements relevant to methanol consumption by  
17 microbial communities across the Atlantic, a rare basin-wide evaluation. The description is  
18 somewhat monotonous, but it is what it is. A great variability is uncovered across provinces  
19 and across depths, but little correlation is found of methanol oxidation/assimilation with  
20 respect to where it happens. In general, dissimilation is somewhat correlated with the  
21 presence of SAR11, and, in general, assimilation is two orders of magnitude lower than  
22 dissimilation. Which means SAR11 uses some other carbon source(s) for building biomass,  
23 and these remain unknown. In general, I think, even if many questions remain unanswered,  
24 this is a useful benchmark study. – **We thank L. Chistoserdova for their comments.**

25 Improvements that I would like to suggest: – **Each of L. Chistoserdova's comments has  
26 been addressed individually as follows.**

27 Page 14, line 15, please say tetrahydrofolate-linked C1 transfer pathway, there are various  
28 oxidation levels and none of them are methyl- level after methanol oxidation. - **“methyl-THF**

1 linked oxidation pathway” will be changed to “tetrahydrofolate-linked C1 transfer  
2 pathway” (Page 14, line 15).

3 Same page, lines 15-18. You do not see any bona fide methylotrophs in your 16S libraries.  
4 How can you conclude that they are active, along with SAR11? Either elaborate or remove  
5 this statement. PCR amplification of specific genes does not compare with 16S analysis, and  
6 you do not do any in this study anyway. – **Specific gene amplification, using *mxoF***  
7 **functional gene primers, has been conducted previously on the same samples as the**  
8 **current study looking at 16S rRNA analysis. The *mxoF* functional gene analysis**  
9 **identified classic methylotrophic bacteria from these samples, these results are**  
10 **published in a previous manuscript Dixon et al. (2013). To clarify this in the text we**  
11 **have amended “Methylotrophic bacteria such as *Methylophaga* sp., *Methylococcaceae***  
12 **sp. and *Hyphomicrobium* sp. have been previously identified, using *mxoF* functional**  
13 **gene primers (which encode for the classical methanol dehydrogenase), from the upper**  
14 **water column of Atlantic Ocean provinces (Dixon et al., 2013)” to “Previously**  
15 **methylotrophic bacteria such as *Methylophaga* sp., *Methylococcaceae* sp. and**  
16 ***Hyphomicrobium* sp. have been identified, using *mxoF* functional gene primers (which**  
17 **encode for the classical methanol dehydrogenase), from the same DNA samples**  
18 **analysed for 16S rRNA genes in this study, from the upper water column of Atlantic**  
19 **Ocean provinces (Dixon et al., 2013)” (Page 14, lines 7-10).**

20

21 Meantime, an interesting question: while true methylotrophs do inhabit marine waters, why  
22 are they so sparse and apparently uncompetitive compared to SAR11? Can you elaborate? –  
23 **The authors agree this is an interesting question and more work is needed to unpick**  
24 **this. We don’t have an answer for this with the published literature and knowledge**

1 currently available, however we can speculate that it may be down to the shear  
2 abundance and evolutionary strategy of SAR11 in comparison to true methylotrophs.  
3 SAR11 are the most abundant, free living, heterotrophic bacteria in open ocean systems  
4 and are often the most abundant organisms in oligotrophic waters. The competitiveness  
5 and high abundance of SAR11 cells in open ocean waters could be one part of a reason  
6 why true methylotrophs are relatively sparse in comparison. SAR11 have been shown  
7 to have one of the smallest genome sizes of any replicating cell and Giovannoni et al.,  
8 (2005) suggest that the streamlining hypothesis may provide an explanation for this.  
9 The streamline hypothesis, assumption that selection reduces genome size due to the  
10 metabolic burden of replicating DNA without adaptive value, could be the strategy  
11 responsible for the dominance and success of the SAR11 clade in oligotrophic waters.  
12 (this has not been added to the manuscript)

13 Fig. 5 would greatly benefit from introducing colors, would be so much easier to compare  
14 guild distribution. Also, please order the taxa in a uniform way, i.e. use the same taxon order  
15 in each panel. – **We have made the appropriate changes as suggested.**

16 Table 1. Specify that you show ranges below averages/means, specify which. Specify what  
17 NA means. – **A comment has been added to the end of the Table caption to clarify what  
18 these values are “Values given are average  $\pm$  standard deviation (range). NA denotes  
19 that data is not available.”**

20

21 **S. Giovannoni (Referee) [steve.giovannoni@oregonstate.edu](mailto:steve.giovannoni@oregonstate.edu)**

22 Received and published: 11 March 2018

23

1 This is a valuable study that adds significantly to our understanding of methanol oxidation in  
2 the oceans. The authors report seawater methanol oxidation rates obtained with <sup>14</sup>C tracer  
3 methods, and microbial diversity measurements, from a latitudinal transect between 40S and  
4 50N. They find that methanol oxidation rates are correlated with SAR11 relative abundance.  
5 Overall, the reported rates of methanol oxidation are in good agreement with previous  
6 measurements, but this study is exceptional in geographical scope and exploration of  
7 variables such as community composition and depth. Interestingly, the manuscript reports an  
8 inverse correlation between bacterial production estimated by the <sup>3</sup>H leucine method and  
9 methanol oxidation. Although there have been a number of reports previously on methanol  
10 cycling in the oceans, I see the subject coming of age with this report, which confirms what  
11 we knew and also shows us new trends that could only have been observed by making  
12 extensive measurements across a latitudinal transect. - **We thank S. Giovannoni for his**  
13 **comments.**

14 A couple of comments follow about aspects of the paper that could be improved. - **Each of S.**  
15 **Giovannoni's comments has been addressed individually as follows.**

16

17 1. I recommend commenting on the abundance of methylophaga and OM43 in the 454 data,  
18 or indicate they were not detected if that is the case. It may be that the relatively low  
19 coverage obtained in this study (386 seqs/sample) led to these taxa being undetectable. If this  
20 is the case, that should be explained so that readers new to this topic understand the issues.  
21 OM43 is not mentioned at all, but perhaps it should be, since it has been shown to be an  
22 obligate methylotroph, is one of the dominant taxa in some coastal environments, and has  
23 been shown to be a source of abundant XoxF peptides in a coastal ocean metaproteome. –  
24 **The authors recognise this omission in the manuscript and acknowledge that these taxa**

1 should be included to reflect our current understanding of marine methanol utilisation.

2 Therefore, we have added the following;

3 “Although numerically very rare (1-11 16S rRNA gene sequences per sample), 16S  
4 rRNA gene sequences identified as *Methylophaga* spp., *Methylophaga* sp. DMS021  
5 (EU001861) and uncultured *Methylophaga* sp. (EU031899), were found in each of the  
6 Atlantic Ocean provinces in this study (at 97% PAR or 200m depth), consistent with  
7 previous identification of *Methylophaga* spp. in these Atlantic provinces using *mxoF* gene  
8 cloning in (Dixon et al., 2013).” (Page 14, lines 14-19).

9 “Members of *Betaproteobacteria*, OM43, have been shown to be obligate methylotrophs,  
10 with cultivated cells of strain HTCC2181 dissimilating 3.5 times more methanol than  
11 was assimilated (Halsey et al., 2012). OM43 were not successfully identified in the 16S  
12 rRNA sequences in this study, which could be an artefact of the relatively low sequence  
13 coverage (386 sequences per sample) leading to this taxon not being detectable. During  
14 a previous coastal study in the western English Channel (16S rRNA pyrosequence data,  
15 Sargeant et al., 2016) only a single sequence of the OM43 clade, HTCC2181, was  
16 identified. This is a limitation of this type of environmental sequencing effort and  
17 should be a consideration in planning any future projects aiming to understand  
18 microbial function through process measurements alongside the generation of  
19 metagenomic datasets.” (Page 14, line 28 – Page 15, line 6).

20 We have also added the additional reference: Halsey KH, Carter AE, Giovannoni SJ  
21 (2012) Synergistic metabolism of a broad range of C1 compounds in the marine  
22 methylotrophic bacterium HTCC2181. *Environmental Microbiology* 14:630-640.

23 2. Amplicon ratios are not as powerful as cell numbers for identifying correlations between  
24 taxa and rates, although they are much easier to obtain. So, the correlations with SAR11 are

1 not with SAR11 cells per unit volume, which would be best, but rather a correlation between  
2 the relative success of SAR11 in the community and rates of MeOH oxidation. I suggest the  
3 authors revisit the manuscript and choose wording that conveys these issues to oceanographer  
4 readers, who often misunderstand this aspect of relative abundance data. – **The authors**  
5 **recognise that this is a limitation and clarity should be provided. We have added;**

6 **“It should be noted that this correlation has been made with amplicon ratios, relating to**  
7 **the relative success of SAR11 in the community, rather than with SAR11 cell numbers**  
8 **specifically.” (Page 13, lines 10-12).**

9 **“More work is required to add clarity and understanding to the role that SAR11 cells**  
10 **play in marine community methanol dissimilation.” (Page 14, lines 3-4).**

11

12 **Associate Editor (minor comments)**

13 Page 10, line 20. Replace “calculated” with “found” or “detected”, or something like that.

14 **Done.**

15

16 Legend Fig. 1 replace “circles” with “ovals” **Done.**

17 **We hope the manuscript is now acceptable,**

18 **Best wishes**

19

20 **Jo**

21 **Dr Joanna L Dixon**

# 1 Basin-scale variability of microbial methanol uptake in the 2 Atlantic Ocean

3 Stephanie L. Sargeant<sup>1\*</sup>, J. Colin Murrell<sup>2</sup>, Philip D. Nightingale<sup>1</sup>, Joanna L. Dixon<sup>1</sup>

4 <sup>1</sup>Plymouth Marine Laboratory, Prospect Place, West Hoe, Plymouth, Devon, PL1 3DH, UK

5 <sup>2</sup>School of Environmental Science, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK

6 \*Now at: Centre for Research in Biosciences, University of the West of England, Frenchay Campus,  
7 Coldharbour Lane, Bristol, BS16 1QY, UK.

8 Correspondence to: Joanna L. Dixon (jod@pml.ac.uk)

9 **Abstract.** Methanol is a climate active gas and the most abundant oxygenated volatile organic compound  
10 (OVOC) in the atmosphere and seawater. Marine methylotrophs are aerobic bacteria that utilise methanol from  
11 seawater as a source of carbon (assimilation) and/or energy (dissimilation). A few spatially limited studies have  
12 previously reported methanol oxidation rates in seawater; however the basin-wide ubiquity of marine microbial  
13 methanol utilisation remains unknown. This study uniquely combines seawater <sup>14</sup>C labelled methanol tracer  
14 studies with 16S rRNA pyrosequencing to investigate variability in microbial methanol dissimilation and known  
15 methanol utilising bacteria throughout a meridional transect of the Atlantic Ocean between 47° N to 39° S.  
16 Microbial methanol dissimilation varied between 0.05–1.68 nmol l<sup>-1</sup> h<sup>-1</sup> in the top 200 m of the Atlantic Ocean  
17 and showed significant variability between biogeochemical provinces. The highest rates of methanol  
18 dissimilation were found in the northern subtropical gyre (average 0.99±0.41 nmol l<sup>-1</sup> h<sup>-1</sup>), which were up to  
19 eight times greater than other Atlantic regions. Microbial methanol dissimilation rates displayed a significant  
20 inverse correlation with heterotrophic bacterial production (determined using <sup>3</sup>H-leucine). Despite significant  
21 depth stratification of bacterial communities, methanol dissimilation rates showed much greater variability  
22 between oceanic provinces compared to depth. There were no significant differences in rates between samples  
23 collected under light and dark environmental conditions. The variability in the numbers of SAR11 (16S rRNA  
24 gene sequences) were estimated to explain approximately 50% of the changes in microbial methanol  
25 dissimilation rates. We estimate that SAR11 cells in the Atlantic Ocean account for between 0.3-59 % of the  
26 rates of methanol dissimilation in Atlantic waters, compared to <0.01-2.3 % for temperate coastal waters. These  
27 results make a substantial contribution to our current knowledge and understanding of the utilisation of  
28 methanol by marine microbial communities, but highlight the lack of understanding of *in situ* methanol  
29 production mechanisms.

30

31

## 32 1. Introduction

33 Methanol is the most abundant oxygenated volatile organic compound (OVOC) in the  
34 background troposphere where it acts as a climate active gas, influencing the oxidative  
35 capacity of the atmosphere, concentrations of ozone and hydroxyl radicals (Carpenter et al.,  
36 2012). Methanol has been shown to be ubiquitous in waters of the Atlantic Ocean ranging

1 between <math>27\text{--}361\text{ nM}</math> (Beale et al., 2013; Williams et al., 2004; Yang et al., 2013; Yang et  
2 al., 2014). Our knowledge of the sources and sinks of methanol is limited and often lacks  
3 consensus. For example, recent eddy covariance flux estimates demonstrated a consistent flux  
4 of atmospheric methanol into the surface waters of a meridional transect of the Atlantic  
5 Ocean (Yang et al., 2013). However, along a similar transect, 12 months earlier, Beale et al.  
6 (2013) calculated that the Atlantic Ocean represents an overall source of methanol to the  
7 atmosphere ( $3\text{ Tg yr}^{-1}$ ), which was largely attributable to an efflux from the North Atlantic  
8 gyre; where surface concentrations were as high as  $361\text{ nM}$ . Wet deposition from rainwater  
9 has also recently been suggested to represent a supply of methanol to the ocean (Felix et al.,  
10 2014).

11 Although *in situ* marine photochemical production of methanol has previously been found to  
12 be insignificant (Dixon et al., 2013), there is thought to be a substantial unidentified  
13 biological source of methanol in seawater (Dixon et al., 2011a). Biological production by  
14 phytoplankton and during the breakdown of marine algal cells are possible sources (Heikes et  
15 al., 2002; Nightingale, 1991; Sieburth and Keller, 1989). Recent laboratory culture  
16 experiments suggest that methanol is produced by a wide variety of phytoplankton including  
17 cyanobacteria (*Prochlorococcus marinus*, *Synechococcus* sp. and *Trichodesmium*  
18 *erythraeum*) and Eukarya (*Emiliana huxleyi*, *Phaeodactylum tricorutum* and  
19 *Nannochloropsis oculata*, *Dunaliella tertiolecta*) (Mincer and Aicher, 2016, Halsey et al.,  
20 2017). The mechanisms of *in situ* methanol production and their regulation remains largely  
21 unknown, although Halsey et al. (2017) reported light-dependent rates of methanol  
22 production in cultures of the marine green flagellate *Dunaliella tertiolecta* (cell size of  $10\text{--}12$   
23  $\mu\text{m}$ ).

24 Methylophilic bacteria are capable of utilising one-carbon compounds including methanol  
25 as their sole source of energy (methanol dissimilation) and carbon (methanol assimilation).  
26 Methylophilic are widespread in terrestrial and aquatic systems (Kolb, 2009), but research  
27 into these bacteria in marine environments is still at an early stage. Traditionally,  
28 methylophilic were thought to utilise methanol dehydrogenase (MDH encoded by *mxhF*,  
29 McDonald and Murrell, 1997) to metabolise methanol to formaldehyde, with further  
30 oxidation to  $\text{CO}_2$  or incorporation of carbon into biomass (Chistoserdova, 2011;  
31 Chistoserdova et al., 2009). However, recent progress in this field has resulted in the  
32 discovery of the *xoxF* gene, encoding an alternative MDH (Wilson et al., 2008) and  
33 seemingly present in all known gram-negative methylophilic to date (Chistoserdova, 2011;



1 Chistoserdova et al., 2009). The presence of methylotrophs in seawater has been confirmed  
2 using a range of molecular approaches including functional gene primers, stable isotope  
3 probing and metaproteomics (Dixon et al., 2013; Grob et al., 2015; Neufeld et al., 2008;  
4 Neufeld et al., 2007; Taubert et al., 2015). There are also bacterial cells that utilise methanol  
5 and other C<sub>1</sub> compounds for the production of energy but not biomass e.g. SAR11 for which  
6 Sun et al. (2011) proposed the new term ‘methylvores’, distinct from true methylotrophs  
7 which use C<sub>1</sub> compounds as sources of carbon and energy.

8 Limited studies of microbial methanol assimilation in the Atlantic Ocean have previously  
9 shown rates up to 0.42 nmol l<sup>-1</sup> h<sup>-1</sup> in recently upwelled coastal waters of the Mauritanian  
10 Upwelling (Dixon et al., 2013). However, open ocean waters of the Atlantic were  
11 substantially lower ranging between 0.002–0.028 nmol l<sup>-1</sup> h<sup>-1</sup> (Dixon et al., 2013). Microbial  
12 methanol dissimilation rates are generally up to 1000-fold higher than rates of assimilation;  
13 ranging between 0.70–11.2 and <0.001–0.026 nmol l<sup>-1</sup> h<sup>-1</sup> respectively for coastal waters  
14 (Sargeant et al., 2016; Dixon et al., 2011b). Methanol dissimilation rates ranging between  
15 0.08–6.1 nmol l<sup>-1</sup> h<sup>-1</sup> have also been found in open ocean Atlantic waters (Dixon et al.,  
16 2011a). However, despite the ubiquity of methanol in seawater, the spatial extent or  
17 quantification of microbial methanol utilisation for energy production on a basin scale has not  
18 been previously investigated. Therefore, the objective of this research was to simultaneously  
19 characterise the spatial variability in microbial methanol dissimilation rates (at depths to 200  
20 m) and in microbial community groups throughout contrasting biogeochemical regions of the  
21 Atlantic Ocean. This study represents the first basin-wide approach to investigating methanol  
22 as a source of reducing power and energy for microbes.

23

## 24 **2. Materials and Methods**

### 25 *2.1. Sampling strategy*

26 Sampling was carried out during an Atlantic Meridional Transect (AMT) ([http://www.amt-](http://www.amt-uk.org)  
27 [uk.org](http://www.amt-uk.org)). The research cruise (JC039, RRS James Cook, 13/10/09–01/12/09) departed from  
28 Falmouth, UK (50.15° N, 05.07° W) and arrived in Punta Arenas, Chile (53.14° S, 70.92°  
29 W). Water samples were collected daily from pre-dawn (97, 33, 14 and 1 %  
30 photosynthetically active radiation (PAR) equivalent depths and 200 m) and solar noon (97  
31 %) conductivity-temperature-depth (CTD) casts. The PAR equivalent depths were 5 m, 10–31

1 m, 15-54 m and 38-127 m for the 97, 33, 14, 1 % light levels respectively and typically varied  
2 with oceanic province. The pre-dawn and solar noon sampling periods were approximately  
3 45-65 nautical miles apart (sampling locations are shown in Fig. 1). The Atlantic Ocean was  
4 divided into five oceanic provinces, following the approach of Dixon et al. (2013), according  
5 broadly to chlorophyll *a* concentrations (<0.15 mg m<sup>-3</sup> gyre regions, >0.15 mg m<sup>-3</sup> temperate  
6 or upwelling regions, Fig. 1) with the northern gyre sub-divided into northern subtropical  
7 gyre (NSG) and northern tropical gyre (NTG). Measurements of the concentration of  
8 methanol in seawater (Beale et al. 2013) and of methanol assimilation rates (Dixon et al.  
9 2013) made during this transect have been reported previously.

10

## 11 2.2. Microbial methanol uptake

12 The oxidation of methanol to CO<sub>2</sub> (dissimilation) was determined using <sup>14</sup>C-labelled  
13 methanol (American Radiolabelled Chemicals Inc, Saint Louis, MO, USA) seawater  
14 incubations as previously described in Dixon et al. (2011b). Seawater samples of 1 ml were  
15 incubated with ~10 nM (final concentration) <sup>14</sup>C-labelled methanol to measure rates of  
16 microbial methanol dissimilation. Seawater methanol concentrations ranged between 48-361  
17 nM (Beale et al., 2013) thus the radiotracer additions represent 3-21 % of *in situ*  
18 concentrations in Atlantic waters. Incubations were conducted in triplicate, with 'killed'  
19 controls (5 % trichloroacetic acid, TCA, final concentration), at *in situ* temperatures and in  
20 the dark. Incubation temperatures were determined by the sea surface temperature recorded  
21 by the corresponding CTD casts. Sample counts of <sup>14</sup>CO<sub>2</sub>, captured in the precipitate as  
22 Sr<sup>14</sup>CO<sub>3</sub> (nCi ml<sup>-1</sup> h<sup>-1</sup>), were divided by the total <sup>14</sup>CH<sub>3</sub>OH added to the sample (nCi ml<sup>-1</sup>) to  
23 calculate the apparent rate constants, *k* (h<sup>-1</sup>).

24

25 The incorporation of methanol carbon into microbial biomass (assimilation) was determined  
26 using sample volumes of 320 ml to increase the total sample counts (Dixon et al., 2011b)  
27 following procedures outlined in Dixon et al. (2011b, 2013). Filter sample counts were  
28 divided by the total <sup>14</sup>CH<sub>3</sub>OH added to the sample (nCi ml<sup>-1</sup>) to calculate the apparent rate  
29 constants, *k* (h<sup>-1</sup>). For both methanol assimilation and dissimilation, the specific activity of  
30 <sup>14</sup>C-labelled methanol (57.1 mCi mmol<sup>-1</sup>) was multiplied by the apparent rate constants to  
31 calculate rates of microbial methanol uptake (nmol l<sup>-1</sup> h<sup>-1</sup>) following the approach of Dixon et  
32 al. (2013). Evaluation of control samples suggests that ≤0.3 % of the added <sup>14</sup>CH<sub>3</sub>OH is

1 recovered on the filters and  $\leq 2$  % in the resultant precipitate for methanol assimilation and  
2 dissimilation respectively.

3

### 4 2.3. Bacterial leucine incorporation

5 Rates of bacterial leucine incorporation were measured using the incorporation of  $^3\text{H}$ -leucine  
6 into bacterial protein in seawater samples using the method described by Smith and Azam  
7 (1992). A final concentration of 25 nM (6.8  $\mu\text{l}$ ) of  $^3\text{H}$ -leucine (calculated using the specific  
8 activity of 161 Ci  $\text{mmol}^{-1}$ , concentrations 1 mCi  $\text{ml}^{-1}$ , American Radiolabelled Chemicals Inc,  
9 Saint Louis, MO, USA) was incubated with 1.7 ml seawater samples. Incubations were  
10 conducted in triplicate with 'killed' controls (5 % TCA, final concentrations), at *in situ*  
11 temperature and in the dark.

12

### 13 2.4. Bacterial community composition

14 Seawater samples of approximately twenty litres were collected for bacterial DNA analysis  
15 from 97, 33, 1 and  $<1$  % (200 m) PAR equivalent depths during pre-dawn CTD casts only.  
16 Samples were filtered through 0.22  $\mu\text{m}$  Sterivex polyethersulfone filters (Millipore, Watford,  
17 UK) using a peristaltic pump. Filters were incubated with 1.6 ml of RNA Later (Life  
18 Technologies, to preserve samples during shipment) overnight at 4° C, after which the RNA  
19 Later was removed. Filters were stored immediately at -80° C before being shipped back to  
20 the UK on dry ice and subsequently stored at -20 °C.

21 Bacterial DNA was extracted from filters using a modified phenol:chloroform:isoamyl  
22 alcohol extraction method as previously described in Neufeld et al. (2007). Extracted DNA  
23 was cleaned using Amicon ultra-0.5 centrifugal filter devices (Millipore) to remove any RNA  
24 Later residue. The 16S rRNA gene primers 341F (Muyzer et al., 1993) and 907R (Muyzer et  
25 al., 1998) were used for PCR amplification (32 cycles) with an annealing temperature of 55°  
26 C. Purification of PCR products from agarose gels was conducted using the QIAquick gel  
27 extraction kit (Qiagen, Crawley, UK) before being sent to Molecular Research LP (MR DNA,  
28 <http://www.mrdnalab.com>) for 454 pyrosequencing using the GS-flx platform.

29 The 16S rRNA gene sequences were depleted of barcodes and primers, and then sequences  
30 less than 200 bp, with ambiguous bases or with homopolymer runs exceeding 6 bp, were

1 removed. Sequences were de-noised and chimeras removed. After the removal of singleton  
2 sequences, operational taxonomic units (OTUs) were defined at 97 % 16S rRNA gene  
3 identity using Quantitative Insights Into Microbial Ecology (QIIME, <http://qiime.org>,  
4 *Caporaso et al.* 2010). The OTUs were assigned taxonomically using BLASTn (Basic Local  
5 Alignment Search Tool, NCBI) against the Silva database (<http://www.arb-silva.de>).  
6 Sequences were randomly re-sampled to the lowest number of sequences per sample (386  
7 sequences per DNA sample) to standardise the sequencing effort.

8

### 9 **3. Results**

#### 10 *3.1. Microbial methanol dissimilation*

##### 11 *3.1.1 Surface*

12 Pre-dawn surface rates of microbial methanol dissimilation ranged between 0.05–1.49 nmol l<sup>-1</sup> h<sup>-1</sup>  
13 throughout the transect of the Atlantic Ocean (Fig. 2a). Maximum variability in surface  
14 rates of methanol dissimilation (average of  $0.96 \pm 0.45$  nmol l<sup>-1</sup> h<sup>-1</sup>, n=10) were observed  
15 north of 25° N in NT and NSG regions. At the southern limit of the NSG, rates of methanol  
16 dissimilation decreased sharply from 1.48 to 0.34 nmol l<sup>-1</sup> h<sup>-1</sup>. Generally, surface rates  
17 continued to decrease in a southward direction throughout the NTG and EQU regions,  
18 reaching a minimum of 0.05 nmol l<sup>-1</sup> h<sup>-1</sup> in Equatorial upwelling waters. Interestingly, surface  
19 rates started to gradually increase to 0.39 nmol l<sup>-1</sup> h<sup>-1</sup> in waters of the oligotrophic SG, before  
20 declining to 0.18 nmol l<sup>-1</sup> h<sup>-1</sup> in the ST area. Methanol dissimilation rates determined at pre-  
21 dawn (dark) generally exhibited a similar latitudinal pattern to those from solar noon (light).  
22 Rates south of 25° N (NTG, EQU, SG, ST) showed a significant, almost 1:1 relationship,  
23 between light (solar noon, y) and dark (pre-dawn, x) *in situ* sampling conditions ( $y=1.06x$ ,  
24  $r=0.6240$ , n=13,  $P<0.05$ ), with most variability between results from light versus dark  
25 sampling occurring north of 25° N in NT and NSG provinces. This is most likely a reflection  
26 of these waters exhibiting the greatest spatial variability, as the pre-dawn and midday stations  
27 were typically 55 nautical miles apart.

##### 28 *3.1.2 Depth distributions*

29 The average rates of methanol dissimilation with depth are shown in Fig. 3a for each oceanic  
30 province. Rates varied between 0.05–1.68 nmol l<sup>-1</sup> h<sup>-1</sup>, but showed no consistent statistically

1 significant trend with depth. However, clear differences were observed in microbial methanol  
2 dissimilation in the top 200 m between contrasting provinces in the Atlantic Ocean; where  
3 NSG>NT>SG>ST>NTG>EQU. The highest rates of methanol dissimilation in the top 200 m  
4 were observed in the most northern latitudes ( $0.22\text{--}1.50$  and  $0.15\text{--}1.68$   $\text{nmol l}^{-1} \text{h}^{-1}$  for NT  
5 and NSG respectively), consistent with surface trends (Fig. 2a). A strong decrease was  
6 observed between the NSG ( $0.99 \pm 0.41$   $\text{nmol l}^{-1} \text{h}^{-1}$ ) and the NTG ( $0.18 \pm 0.04$   $\text{nmol l}^{-1} \text{h}^{-1}$ )  
7 regions. However, rates of microbial methanol dissimilation determined in the oligotrophic  
8 waters of the NTG ( $0.18 \pm 0.04$   $\text{nmol l}^{-1} \text{h}^{-1}$ ) and SG ( $0.24 \pm 0.12$   $\text{nmol l}^{-1} \text{h}^{-1}$ ) regions were  
9 comparable with rates in the ST region ( $0.20 \pm 0.05$   $\text{nmol l}^{-1} \text{h}^{-1}$ ), with the EQU exhibiting the  
10 lowest average rates of  $0.11 \pm 0.03$   $\text{nmol l}^{-1} \text{h}^{-1}$ .

11 Overall, latitudinal trends in depth profiles for methanol dissimilation rates mirrored those  
12 found in surface waters. Surface microbial methanol dissimilation rates determined from pre-  
13 dawn (x) water were compared to those from 200 m (y), which are in permanent darkness  
14 (the deepest 1 % PAR equivalent depth of 175 m was found was in the SG at  $\sim 19.50^\circ\text{S}$ ) and  
15 also showed a  $\sim 1:1$  relationship ( $y=0.967x$ ,  $r=0.9237$ ,  $n=19$ ,  $P<0.001$ ).

16

### 17 3.2. Bacterial leucine incorporation rates

#### 18 3.2.1 Surface

19 Rates of bacterial leucine incorporation (BLI) varied between  $2.9\text{--}25.2$   $\text{pmol l}^{-1} \text{h}^{-1}$  in the pre-  
20 dawn surface waters of the Atlantic transect (Fig. 2b). On average, surface rates of BLI were  
21 highest in the relatively more productive EQU upwelling region ( $18.3 \pm 4.8$   $\text{pmol l}^{-1} \text{h}^{-1}$ ), and  
22 lowest in the northern sub-tropical gyre (NSG,  $5.2 \pm 2.3$   $\text{pmol l}^{-1} \text{h}^{-1}$ ). Surface rates of BLI  
23 averaged  $7.8 \pm 2.3$   $\text{pmol l}^{-1} \text{h}^{-1}$  and  $7.7 \pm 2.4$   $\text{pmol l}^{-1} \text{h}^{-1}$  in the NTG and SG regions  
24 respectively. The one measurement of BLI in the ST suggested much higher rates ( $25.2$   $\text{pmol}$   
25  $\text{l}^{-1} \text{h}^{-1}$ ) than previously determined during the transect, even when compared to the NT region  
26 ( $9.9 \pm 3.9$   $\text{pmol l}^{-1} \text{h}^{-1}$ ). Pre-dawn (dark) rates of BLI generally exhibited a similar latitudinal  
27 pattern to those from solar noon (light), with more variability between light and dark  
28 sampling observed in the waters of the productive EQU region. Bacterial rates of leucine  
29 incorporation determined from samples collected at solar noon (y) were approximately 20%  
30 less than those determined at pre-dawn ( $y=0.7815x$ ,  $r=0.7288$ ,  $n=22$ ,  $P<0.001$ ), perhaps  
31 reflecting a degree of light inhibition of heterotrophic bacterial production.

1

### 2 3.2.2 Depth profiles

3 Rates of bacterial leucine incorporation varied between 0.5–60.2 pmol l<sup>-1</sup> h<sup>-1</sup> throughout the  
4 top 200 m of the water column. In the sunlit depths (97-1 % PAR) generally BLI rates  
5 followed the pattern EQU>NTG≈SG>NT>NSG (excluding the outliers of 60.2 and 31.3 pmol  
6 l<sup>-1</sup> h<sup>-1</sup> observed for the NSG at 14 % PAR from two depth profiles in this province). This  
7 trend differs slightly from that observed for surface only data due to sub-surface (1-14 %  
8 PAR) maxima observed in both the north and south oligotrophic gyres (NSG, NTG, SG). In  
9 the NT, NTG and EQU provinces, BLI rates were generally higher in sunlit depths compared  
10 to the dark at 200 m (Fig. 3b). However, there were no statistical differences between the  
11 provinces for rates of BLI determined at 200 m.

12

### 13 3.3. Bacterial community composition

#### 14 3.3.1 Surface

15 The total number of operational taxonomic units (OTUs) sequenced throughout the Atlantic  
16 Ocean varied between 91–207. Overall, the largest contributors to surface bacterial  
17 communities were *Prochlorococcus* and SAR11 16S rRNA gene sequences (Fig. 5a); which  
18 together accounted for between 21-60 % of all OTUs (21% in the SG and 60% in the NSG).  
19 These bacteria typically numerically dominate surface waters of nutrient depleted oceanic  
20 regions e.g. Gomez-Pereira et al. (2013). The numbers of *Prochlorococcus*, determined via  
21 flow cytometry, for the same surface samples from which 16S rRNA genes were amplified  
22 range between 0.81 x 10<sup>5</sup> for the NTG region and 3.10 x 10<sup>5</sup> cells ml<sup>-1</sup> for the EQU region  
23 (see Table 2 for summary). *Prochlorococcus* 16S rRNA gene sequences contributed an  
24 average of 28 ± 12 % of the community composition of surface samples throughout the  
25 surface Atlantic Ocean. Numbers of SAR11 16S rRNA gene sequences contributed a  
26 maximum of 24 % to the total 16S rRNA gene sequences for the NSG region, and overall  
27 contributed an average of 11 ± 3 % to the bacterial community in surface waters of the  
28 Atlantic Ocean. There was a clear shift between surface bacterial communities in the two  
29 northern gyre provinces with *Prochlorococcus* and SAR11 16S rRNA gene sequences  
30 decreasing from the NSG to the NTG region (59 and 33 % of total 16S rRNA gene sequences  
31 respectively). *Oceanspirillales* and *Flavobacteriales* 16S rRNA gene sequences contributed

1 approximately double the amount (compared to the total 16S rRNA sequences) in the NTG  
2 compared to the NSG region (25 and 12 % respectively).

3  
4 Microbial communities of the surface waters of the NT, NSG and EQU provinces were  
5 dominated by *Prochlorococcus*, *Alteromonadales* and SAR11, together representing between  
6 64–72 % of 16S rRNA gene sequences. These orders were less dominant in the more  
7 oligotrophic waters of the NTG and SG, accounting for 43 % and 34 % of 16S rRNA gene  
8 sequences respectively. In these oligotrophic regions (NTG and SG) microbial communities  
9 appear less dominated by a few orders, with a more even spread of bacterial orders  
10 contributing to the community composition (Fig. 5a).

### 11 12 3.3.2 Depth profiles

13 The largest contributors to bacterial communities at the 33 % PAR depths were, like surface  
14 communities, *Prochlorococcus* and SAR11 16S rRNA gene sequences (Fig. 5b). Together  
15 they accounted for between 47-70 % of all OTUs, with the minimum and maximum  
16 contributions in the SG and EQU provinces respectively. If the proportion of sequences  
17 contributing individually <5% were included then collectively they accounted for between  
18 69-91 % of all 16S rRNA gene sequences. The main differences between the surface and  
19 33% PAR equivalent depth (14-31 m) are the increasing dominance of the cyanobacteria  
20 *Prochlorococcus*, and the decrease in relative contribution of *Alteromonadales* at 33% PAR  
21 depths, particularly in the NT region.

22  
23 In the darker 1 % PAR depths (59-127 m) *Prochlorococcus* and SAR11 16S rRNA gene  
24 sequences (Fig. 5c) still accounted for between 32-65 % of all OTUs, with the minimum and  
25 maximum contributions in the SG and EQU respectively. With the addition of sequences for  
26 each Order contributing <5 % to the total 16SrRNA gene sequences, these three categories  
27 accounted for 60-81% of all 16S rRNA gene sequences retrieved throughout each of the  
28 regions sampled. Two notable differences at this light level in the SG region compared to the  
29 other provinces are the 12 % contribution made by the Order III *Incertae Sedis* which belongs  
30 to the *Bacteroidetes* class, and the relative reduction in contribution made by  
31 *Prochlorococcus* (11 % compared to an Atlantic average of  $27\pm 15$  % at 1 % PAR). However,  
32 the latter trend is not confirmed in the cell numbers of *Prochlorococcus* determined via flow  
33 cytometry (Table 2).

34

1 In the permanent dark of 200 m, SAR11 bacteria contributed between 14-29 % in northern  
2 regions, which contrasted to only 4-5 % in the EQU and SG provinces. The SAR324 clade  
3 contributed 8-11 % in the northern gyre. Both uncultivated bacteria and those that  
4 individually comprised <5 % contributed relatively highly to the OTUs (10-36 % and 21-33  
5 % respectively). These two groupings together with the SAR11 and SAR324 make up 83-89  
6 % in northern regions and between 37-56 % in the SG and EQU provinces respectively. For  
7 the EQU region the *Alteromonadales* order is also significant at 25 % (which collectively  
8 comprise 81 % of all OTUs for EQU), whilst for the SG the cyanobacteria *Prochlorococcus*  
9 and *Synechococcus* comprise 52 % (which collectively comprise 89 % of all OTUs for SG).

10

#### 11 **4. Discussion**

##### 12 *4.1. Basin scale variability in biological methanol uptake*

13 Maximum rates of methanol dissimilation in the Atlantic Ocean were recorded in the NSG  
14 province at 33 % PAR light depth (25 m, 1.68 nmol l<sup>-1</sup> h<sup>-1</sup>, Fig. 2 and Fig. 4a). An overview  
15 of the variation in rates of methanol dissimilation to CO<sub>2</sub> throughout the top 200 m of the  
16 water column in the Atlantic Ocean is shown in Fig. 4a, which illustrates sub-surface maxima  
17 in northerly latitudes. However, no statistically significant differences were ~~calculated~~found  
18 between rates of methanol dissimilation in the euphotic zone (97-1 % PAR) compared to the  
19 aphotic zone (samples from 200 m) in the NSG ( $t_{NSG}=2.63$ ,  $t_{20}=2.85$  for  $P<0.01$ ), NTG ( $t_{NTG}=  
20 0.02$ ,  $t_{12}=3.05$  for  $P<0.01$ ), EQU ( $t_{EQU}=1.01$ ,  $t_{18}=2.88$  for  $P<0.01$ ) and SG regions ( $t_{SG}=0.88$ ,  
21  $t_{19}=2.88$  for  $P<0.01$ ). This is consistent with a previous study in the north east Atlantic Ocean,  
22 which similarly reported no significant variability in methanol dissimilation rates with depth  
23 (Dixon and Nightingale, 2012). Nevertheless, greater variability with depth was observed for  
24 methanol dissimilation rates from the northern gyre ( $F_{NSG}=3.22$  where  $F_{3,17}=3.20$ ,  $P<=0.05$   
25 and  $F_{NTG}=5.14$  where  $F_{2,10}=4.10$ ,  $P<0.05$ ). Variability in rates from the euphotic zone were  
26 found to be significantly higher than those from 200 m in northern ( $t_{NT}=3.17$ ,  $t_{20}=2.85$  for  
27  $P<0.01$ ) and southern temperate regions ( $t_{ST}=5.03$ ,  $t_{10}=3.17$  for  $P<0.01$ ).

28

29 Although the highest rates of methanol dissimilation were determined in the NSG, these  
30 values were approximately seven times lower than the maxima determined during a seasonal  
31 study of the temperate western English Channel (0.5-11.2 nmol l<sup>-1</sup> h<sup>-1</sup>, Sargeant et al., 2016).  
32 Rates determined in the temperate waters of the south Atlantic (0.11-0.45 nmol l<sup>-1</sup> h<sup>-1</sup>) are  
33 most comparable to the lowest rates determined during late spring and early summer of ~0.50



1 nmol l<sup>-1</sup> h<sup>-1</sup> in temperate northern coastal waters (Sargeant et al., 2016). The seasonal study  
2 in the western English Channel showed maximum rates of up to 11.2 nmol l<sup>-1</sup> h<sup>-1</sup> during  
3 autumn and winter months (Sargeant et al., 2016). The differences in methanol dissimilation  
4 rates between the temperate waters of the North (0.83±0.42 nmol l<sup>-1</sup> h<sup>-1</sup>) and South  
5 (0.27±0.13 nmol l<sup>-1</sup> h<sup>-1</sup>) Atlantic may therefore reflect seasonal differences between  
6 hemispheres i.e. sampling in the NT region occurred during late autumn compared to late  
7 spring in the ST region.

8  
9 Methanol assimilation rates were generally two orders of magnitude lower than dissimilation  
10 rates, reaching a maximum of 0.028 nmol l<sup>-1</sup> h<sup>-1</sup> in the top 200m throughout the Atlantic  
11 Ocean (Fig. 4b). Rates of methanol assimilation exhibited sub-surface maxima (at 33% PAR  
12 equivalent depth) which were particularly evident just north of the Equator (EQU) and in the  
13 northern gyre (NSG) of 0.015±0.004 nmol l<sup>-1</sup> h<sup>-1</sup>. These subsurface rates were on average  
14 higher than surface values (0.004±0.004 nmol l<sup>-1</sup> h<sup>-1</sup>). Results are similar to findings by  
15 Dixon and Nightingale (2012) who also demonstrated sub-surface maxima between 20–30 m  
16 in the north east Atlantic. The methanol assimilation rates are shown for direct comparison to  
17 dissimilation, but have been previously discussed in more detail in Dixon et al. (2012).

#### 18 19 4.2. Bacterial community and productivity

20 In contrast to microbial methanol dissimilation, rates of bacterial leucine incorporation were  
21 lowest in the northern oligotrophic gyre (NSG 5.2 ± 2.3 pmol l<sup>-1</sup> h<sup>-1</sup>, NTG 7.8 ± 2.3 pmol l<sup>-1</sup>  
22 h<sup>-1</sup>) reflecting lower microbial activity in these regions of the Atlantic. Surface microbial  
23 methanol dissimilation rates exhibited a statistically significant inverse correlation with  
24 bacterial leucine incorporation, ( $r = -0.351$ ,  $n = 36$ ,  $P \leq 0.05$ ). This is consistent with  
25 findings from a seasonal study in the western English Channel, where surface rates of  
26 methanol dissimilation were also inversely correlated to bacterial production (Sargeant et al.,  
27 2016). For all the depth data a negative correlation was also found in the NTG, EQU and SG  
28 regions ( $r = -0.372$ ,  $n = 52$ ,  $P \leq 0.01$ ), but NT and NSG areas showed methanol dissimilation  
29 rates independent of BLI. The productivity of heterotrophic bacteria is generally associated  
30 with the concentrations of phytoplankton-derived dissolved organic matter (DOM) e.g.  
31 proteins, lipids and carbohydrates which are utilised as sources of energy and carbon (Benner  
32 and Herndl, 2011; Nagata, 2008; Ogawa and Tanoue, 2003). Results from this present study  
33 indicate that in regions of low heterotrophic bacterial production i.e. in the northern Atlantic

1 Gyre (minimum rate of bacterial leucine incorporation of  $3 \text{ pmol l}^{-1} \text{ h}^{-1}$ ) rates of methanol  
2 dissimilation were relatively higher. In oligotrophic regions, phytoplankton-derived DOM is  
3 scarce, suggesting that those bacteria able to metabolise methanol are using the carbon from  
4 methanol as an alternative source of energy (and to a lesser extent carbon).

5 Although the bacterial community 16S rRNA gene sequence data did not display any clear  
6 patterns with changing biogeochemical province (in contrast to microbial methanol  
7 dissimilation rates), the bacterial community was shown to be depth-stratified throughout the  
8 Atlantic Ocean (Fig. 6a). A non-metric multi-dimensional scale (MDS) plot of a Bray-Curtis  
9 similarity matrix of 16S rRNA gene sequences (Fig. 6a) found bacterial community samples  
10 to cluster into three distinct groupings possibly reflecting light levels: sunlit (97 and 33 %  
11 PAR), minimal light (1 % PAR) and dark (200 m). Bacterial community samples from the  
12 same PAR equivalent depths were found to group together regardless of biogeochemical  
13 province. A larger cluster formed of samples from 97 and 33% PAR is likely to be formed of  
14 bacterial communities originating from the well-mixed surface layer of the water column,  
15 accounting for their similarity in composition. When all environmental parameters were  
16 considered together (including bacterial numbers and BLI) a Euclidean distance matrix non-  
17 metric MDS also demonstrated photic waters (97-1 % PAR) clustered together, and were  
18 significantly different to dark waters from 200m (Fig. 6b). However, no significant  
19 differences were observed between rates of methanol dissimilation determined from the  
20 euphotic zone (samples from 97-1 % PAR equivalent depths) compared to the aphotic zone  
21 (samples from 200 m depth) for gyre and equatorial regions (NSG  $t_{\text{NSG}}=2.63$  ( $t_{20}=2.85$  for  
22  $P<0.01$ ), NTG  $t_{\text{NTG}}=0.02$  ( $t_{12}=3.05$  for  $P<0.01$ ), EQU  $t_{\text{EQU}}=1.01$  ( $t_{18}=2.88$  for  $P<0.01$ ) and SG  
23  $t_{\text{SG}}=0.88$  ( $t_{19}=2.88$  for  $P<0.01$ )) although, clear differences between provinces were evident  
24 (Fig. 6c). This is consistent with results from Dixon and Nightingale (2012) who also found  
25 no significant variation of methanol dissimilation with depth in the north east Atlantic Ocean.  
26 These data suggest that light levels do not have a strong role to play in microbial methanol  
27 dissimilation in waters of the Atlantic, despite the overall bacterial community showing  
28 strong variability with depth (or incident light). Depth-stratification of microbial communities  
29 has been observed previously by Carlson et al. [2004], DeLong et al. [2006] and between  
30 euphotic and aphotic zones in the north western Sargasso Sea (Carlson et al., 2004).  
31 Heywood et al. (2006) suggested that the physical separation of low nutrient surface waters in  
32 gyre regions from mixing with more nutrient rich waters below a defined pycnocline, in  
33 combination with differing levels of light availability, could partially explain changes in

1 bacterial community composition throughout the water column. Therefore, these results could  
2 indicate that methanol dissimilation is limited to specific microbial groups that are present  
3 relatively uniformly between the surface and 200m, although more depth variability is shown  
4 north of 25°N where rates of methanol dissimilation are the highest and most variable.

5

#### 6 4.3. Methanol dissimilation and SAR11

7 SAR11 cells have been shown to utilise methanol, but only as a source of energy (Sun et al.,  
8 2011). The numbers of SAR11 16S rRNA gene sequences exhibited a statistically significant  
9 correlation with rates of microbial methanol dissimilation throughout the Atlantic basin ( $r =$   
10  $0.477$ ,  $n = 20$ ,  $P < 0.05$ ), where the number of SAR11 16S rRNA gene sequences explained  
11 approximately half of the spatial variability in rates of methanol dissimilation. It should be  
12 noted that this correlation has been made with amplicon ratios, relating to the relative success  
13 of SAR11 in the community, rather than with SAR11 cell numbers specifically. In culture,  
14 SAR11 cells (strain HTCC1062) have previously been shown to utilise methanol as a source  
15 of energy at a rate of  $\sim 5 \times 10^{-20}$  moles cell<sup>-1</sup> h<sup>-1</sup> (Sun et al., 2011), which equates to 2 nmol l<sup>-1</sup>  
16 h<sup>-1</sup> (using a culture cell abundance of  $4 \times 10^7$  cells mL<sup>-1</sup>, Sun et al., 2011). SAR11 cells  
17 dominate ( $59 \pm 4\%$ ) the low nucleic acid (LNA) fraction of bacterioplankton consistently  
18 across the Atlantic Ocean, where typically numbers of LNA range between  $0.2$ - $1.0 \times 10^9$  cells  
19 l<sup>-1</sup> (Mary et al., 2006a). Thus estimates of *in situ* SAR11 numbers range between  $0.12$ - $0.59 \times$   
20  $10^9$  cells l<sup>-1</sup>. This is consistent with estimates from the Sargasso Sea of  $\sim 0.1 \times 10^9$  cells l<sup>-1</sup>  
21 (where they are reported to contribute  $\sim 25\%$  of total prokaryotic abundance of  $0.4 \times 10^6$  cells  
22 mL<sup>-1</sup>, Malmstrom et al., 2004). Thus, we estimate that SAR11 cells of the Atlantic Ocean  
23 could be oxidising methanol at rates between  $5$ - $29.5$  pmol l<sup>-1</sup> h<sup>-1</sup>, which could account for  
24 between  $0.3$ - $59\%$  of the rates of methanol dissimilation in surface Atlantic waters.

25 A seasonal investigation in the western English Channel reported bacterial numbers ranging  
26 between  $2.0$ - $15.8 \times 10^5$  cells ml<sup>-1</sup> (Sargeant et al., 2016) which agrees well with data from  
27 Mary et al. (2006b,  $2.0$ - $16.0 \times 10^5$  cells ml<sup>-1</sup>). Assuming that SAR11 contribute between  $9$ -  
28  $20\%$  of total bacterioplankton (Mary et al., 2006b) suggests SAR11 numbers range between  
29  $0.18$ - $3.16 \times 10^5$  cells ml<sup>-1</sup> at this coastal site. Using the above estimate of  $\sim 5 \times 10^{-20}$  moles cell<sup>-1</sup>  
30 h<sup>-1</sup> for rates of methanol dissimilation in cultured SAR11 cells suggests that SAR11 could  
31 oxidise methanol at rates ranging between  $0.9$ - $15.8$  pmol l<sup>-1</sup> h<sup>-1</sup> in temperate coastal regions.  
32 This equates to  $<0.01$ - $2.3\%$  of microbial community methanol dissimilation rates ( $0.7$ - $11.2$

1 nmol l<sup>-1</sup> h<sup>-1</sup>, Sargeant et al., 2016). Therefore, we suggest that cells of the SAR11 clade are  
2 more likely to make a larger contribution to marine microbial methanol dissimilation in open  
3 ocean environments, where alternative sources of carbon are more limited relative to  
4 temperate coastal waters. More work is required to add clarity and understanding to the role  
5 that SAR11 cells play in marine community methanol dissimilation.

6  
7 Previously methylotrophic bacteria such as *Methylophaga* sp., *Methylococcaceae* sp. and  
8 *Hyphomicrobium* sp. have been identified, using *mxoF* functional gene primers (which  
9 encode for the classical methanol dehydrogenase), from the same DNA samples analysed for  
10 16S rRNA genes in this study, from the upper water column of Atlantic Ocean provinces  
11 Methylotrophic bacteria such as *Methylophaga* sp., *Methylococcaceae* sp. and  
12 *Hyphomicrobium* sp. have been previously identified, using *mxoF* functional gene primers  
13 (which encode for the classical methanol dehydrogenase), from the upper water column of  
14 Atlantic Ocean provinces (Dixon et al., 2013). Although numerically very rare (1-11 16S  
15 rRNA gene sequences per sample), 16S rRNA gene sequences identified as *Methylophaga*  
16 spp., *Methylophaga* sp. DMS021 (EU001861) and uncultured *Methylophaga* sp. (EU031899),  
17 were found in each of the Atlantic Ocean provinces in this study (at 97% PAR or 200m  
18 depth), consistent with previous identification of *Methylophaga* spp. in these Atlantic  
19 provinces using *mxoF* gene cloning in (Dixon et al., 2013). More recently the *mxoF* gene,  
20 which encodes an alternative methanol dehydrogenase, has also been found to be widespread  
21 in coastal marine environments (Taubert et al., 2015). SAR11 bacteria are thought to contain  
22 an Fe alcohol dehydrogenase, which although not specific for methanol, can oxidise methanol  
23 (and other short chain alcohols) to formaldehyde which is then thought to be converted to  
24 CO<sub>2</sub> by a methyl-THF linked oxidation pathway-tetrahydrofolate-linked C1 transfer pathway  
25 to produce energy (Sun et al., 2012). Thus it seems likely that both methylotrophic bacteria  
26 possessing *mxoF* and/or *mxoF*, together with microbes such as SAR11 (Sun et al., 2011), are  
27 largely responsible for the turnover of methanol in seawater.

28 Members of *Betaproteobacteria*, OM43, have been shown to be potentially important  
29 obligate methylotrophs, with cultivated cells of strain HTCC2181 dissimilating 3.5 times  
30 more methanol than was assimilated (Halsey et al., 2012). OM43 were not successfully  
31 identified in the 16S rRNA sequences in this study, which could be an artefact of the  
32 relatively low sequence coverage (386 sequences per sample) leading to this taxon not being

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

1 detectable. During a previous coastal study, also analysing 16S rRNA pyrosequence data, in  
2 the western English Channel (Sargeant et al., 2016) only a single sequence of the OM43  
3 clade, HTCC2181, was identified. This is a limitation of this type of environmental  
4 sequencing effort and should be a consideration in planning any future projects aiming to  
5 understand microbial function through process measurements alongside the generation of  
6 metagenomic datasets.”

#### 8 4.4. Marine methanol cycling

9 Data from this study substantially add to the measurements of microbial methanol  
10 dissimilation rates in seawater. This extended spatial coverage clearly demonstrates that  
11 methanol dissimilation is a widespread microbial process taking place in light and dark  
12 environments throughout the Atlantic Ocean. Dissimilation rates are typically two orders of  
13 magnitude greater than assimilation rates across most of the Atlantic Basin. These data  
14 suggest that methanol is an important source of energy for microbes. This is particularly true  
15 in the northern oligotrophic waters of the Atlantic Ocean, where corresponding *in situ*  
16 methanol concentrations range between 148-281 nM (Table 1). What is not clear is the source  
17 of methanol in open ocean waters, which is suspected to be biological in nature (Dixon et al.,  
18 2011a). Although direct flux estimates suggest that the atmosphere could also act as a source  
19 to the ocean (Yang et al, 2013), the magnitude of this flux is insufficient to support the  
20 observed rates of microbial methanol consumed by bacteria, and hence is suspected to be a  
21 minor contribution (Dixon et al., 2011a). Recent culture studies indicate that  
22 *Prochlorococcus sp.*, *Synechococcus sp.* and *Trichodesmium sp.* could produce methanol  
23 (Mincer and Aicher, 2016, Halsey et al., 2017), but *in situ* production mechanisms are  
24 unknown. Further work is needed to fully elucidate and quantify the sources of methanol in  
25 marine waters.

#### 27 5. Conclusions

28 This study reports the first basin-wide understanding of microbial methanol dissimilation  
29 rates in seawater. Radiochemical assays have demonstrated active metabolism throughout the  
30 top 200 m of the water column, with rates being substantially higher in the northern  
31 subtropical Atlantic gyre. Microbial methanol dissimilation rates showed a positive

1 correlation with the numbers of SAR11 16S rRNA gene sequences, and an inverse  
2 relationship with bacterial leucine incorporation. Future work should determine marine  
3 methanol sources and understand the relative contribution of various microbial orders to  
4 methanol loss processes.

5

## 6 **Acknowledgements**

7 We thank all scientists, officers and crew of the RRS James Cook during JC029. We also  
8 thank Rachael Beale for determining methanol concentrations, Glen Tarran for flow  
9 cytometry data, Karen Tait, Yin Chen and Michael Cunliffe for advice with molecular  
10 biology work. Satellite data were processed by the NERC Earth Observation Data  
11 Acquisition and Analysis Service (NEODAAS) at Plymouth Marine Laboratory  
12 (<http://www.neodaas.ac.uk>). This work was funded by the UK Natural Environmental  
13 Research Council (NERC) and the Earth and Life Systems Alliance, Norwich Research Park.  
14 This study is also a contribution to the international IMBeR project and was also supported  
15 by the UK Natural Environment Research Council National Capability funding to Plymouth  
16 Marine Laboratory and the National Oceanography Centre, Southampton. This is contribution  
17 number 323 of the AMT programme.

18

## 19 **Conflict of Interest Statement**

20 The Authors declare no conflict of interest with this manuscript.

21

## 22 **References**

- 23 Beale, R., J. L. Dixon, S. R. Arnold, P. S. Liss, and P. D. Nightingale (2013), Methanol,  
24 acetaldehyde, and acetone in the surface waters of the Atlantic Ocean, *Journal of*  
25 *Geophysical Research: Oceans*, 118(10), 5412-5425, doi: 10.1002/jgrc.20322.
- 26 Benner, R., and G. Herndl (2011), Bacterially derived dissolved organic matter in the  
27 microbial carbon pump, in *Microbial carbon pump in the ocean* edited by N. Jiao, F.  
28 Azam and S. Sanders, pp. 46-48, Science/AAAS, Washington.

- 1 Caporaso, J. G., J. Kuczynski, J. Stombaugh, K. Bittinger, *et al.* (2010), QIIME allows  
2 analysis of high-throughput community sequencing data, *Nature Methods*, 7, 335-336,  
3 doi: 10.1038/nmeth.f.303.
- 4 Carlson, C. A., S. J. Giovannoni, D. A. Hansell, S. J. Goldberg, R. Parsons, and K. Vergin  
5 (2004), Interactions among dissolved organic carbon, microbial processes, and  
6 community structure in the mesopelagic zone of the northwestern Sargasso Sea,  
7 *Limnology and Oceanography* 49, 1073-1083, doi: 10.4319/lo.2004.49.4.1073.
- 8 Carpenter, L. J., S. D. Archer and R. Beale (2012), Ocean-atmosphere trace gas exchange,  
9 *Chem. Soc. Rev.*, 41, 6473-6506, doi:10.1039/c2cs35121h
- 10 Chistoserdova, L. (2011), Modularity of methylotrophy, revisited, *Environmental*  
11 *Microbiology*, 13(10), 2603-2622, doi: 10.1111/j.1462-2920.2011.02464.x.
- 12 Chistoserdova, L., M. G. Kalyuzhnaya, and M. E. Lidstrom (2009), The expanding world of  
13 methylotrophic metabolism, *Annual Review of Microbiology*, 63, 477, doi:  
14 10.1146/annurev.micro.091208.073600.
- 15 DeLong, E. F., C. M. Preston, T. Mincer, V. Rich, S. J. Hallam, N.-U. Frigaard, A. Martinez,  
16 M. B. Sullivan, R. Edwards, and B. R. Brito (2006), Community genomics among  
17 stratified microbial assemblages in the ocean's interior, *Science* 311, 496-503, doi:  
18 10.1126/science.1120250.
- 19 Dixon, J. L., and P. D. Nightingale (2012), Fine scale variability in methanol uptake and  
20 oxidation in the micro-layer and near-surface waters of the Atlantic, *Biogeosciences*,  
21 9, 2961-2972, doi: 10.5194/bg-9-2961-2012.
- 22 Dixon, J. L., R. Beale, and P. D. Nightingale (2011a), Rapid biological oxidation of methanol  
23 in the tropical Atlantic: significance as a microbial carbon source, *Biogeosciences*, 8,  
24 2707-2716, doi: 10.5194/bg-8-2707-2011.
- 25 Dixon, J. L., R. Beale, and P. D. Nightingale (2011b), Microbial methanol uptake in northeast  
26 Atlantic waters, *The ISME Journal*, 5, 704 - 716, doi: 10.1038/ismej.2010.169.
- 27 Dixon, J. L., S. Sargeant, P. D. Nightingale, and J. C. Murrell (2013), Gradients in microbial  
28 methanol uptake: productive coastal upwelling waters to oligotrophic gyres in the  
29 Atlantic Ocean, *The ISME Journal*, 7, 568-580, doi: 10.1038/ismej.2012.130.
- 30 Felix, J. D., S. B. Jones, G. B. Avery, J. D. Willey, R. N. Mead, and R. J. Kieber (2014),  
31 Temporal and spatial variations in rainwater methanol, *Atmospheric Chemistry and*  
32 *Physics Discussions*, 14, 1375 - 1398, doi: 10.5194/acpd-14-1375-2014.
- 33 Gómez-Pereira, P. R., M. Hartmann, C. Grob, G. A. Tarran, A. P. Martin, B. M. Fuchs, D. J.  
34 Scanlan and M. V. Zubkov (2013), Comparable light stimulation of organic nutrient

1 uptake by SAR11 and Prochlorococcus in the North Atlantic subtropical gyre, *The*  
2 *ISME Journal* 7, 603–614; doi:10.1038/ismej.2012.126

3 Grob, C., M. Taubert, A. M. Howat, O. J. Burns, J. L. Dixon, H. H. Richnow, N. Jehmlich,  
4 M. Bergen, Y. Chen, and J. C. Murrell (2015), Combining metagenomics with  
5 metaproteomics and stable isotope probing reveals metabolic pathways used by a  
6 naturally occurring marine methylotroph, *Environmental Microbiology* 17, 4007-  
7 47018; doi: 10.1111/1462-2920.12935.

8 Halsey, KH, Giovannoni, SJ, Graus, M, Zhao, Y, Landry, Z, Thrash, JC, Vergin KL, de  
9 Gouw J (2017), Biological cycling of volatile organic carbon by phytoplankton and  
10 bacterioplankton. *Limnology and Oceanography* 62, 2650-2661: doi:  
11 10.1002/lno.10596.

12 [Halsey, KH, Carter, AE, Giovannoni, SJ \(2012\), Synergistic metabolism of a broad range of Cl](#)  
13 [compounds in the marine methylotrophic bacterium HTCC2181. \*Environmental\*](#)  
14 [Microbiology](#) 14, 630-640: doi:10.1111/j.1462-2920.2011.02605.x.

15 Heikes, B., W. Chang, M. Pilson, E. Swift, *et al.* (2002), Atmospheric methanol budget and  
16 ocean implication, *Global Biogeochemical Cycles*, 16, 1133, doi:  
17 10.1029/2002GB001895.

18 Heywood, J., M. Zubkov, G. Tarran, B. Fuchs, and P. Holligan (2006), Prokaryoplankton  
19 standing stocks in oligotrophic gyre and equatorial provinces of the Atlantic Ocean:  
20 evaluation of inter-annual variability, *Deep Sea Research Part II: Topical Studies in*  
21 *Oceanography*, 53, 1530-1547, doi: 10.1016/j.dsr2.2006.05.005.

22 Kolb, S. (2009), Aerobic methanol-oxidising bacteria in soil, *FEMS Microbiology Letters*, 1 -  
23 10, doi: 10.1111/j.1574-6968.2009.01681.x.

24 Malmstrom, R. R., Kiene, R. P., Cottrell, M. T., & Kirchman, D. L. (2004). Contribution of  
25 SAR11 bacteria to dissolved dimethylsulfoniopropionate and amino acid uptake in the  
26 North Atlantic Ocean. *Applied and Environmental Microbiology*, 70, 4129-4135.

27 Mary, I., J. L. Heywood, B. M. Fuchs, R. Amann, G. A. Tarran, P. H. Burkill, and M. V.  
28 Zubkov (2006a), SAR11 dominance among metabolically active low nucleic acid  
29 bacterioplankton in surface waters along an Atlantic meridional transect, *Aquatic*  
30 *microbial ecology*, 45, 107-113, doi: 10.3354/ame045107.

31 Mary, I., D. Cummings, I. Biegala, P. Burkill, S. Archer, and M. Zubkov (2006b), Seasonal  
32 dynamics of bacterioplankton community structure at a coastal station in the western  
33 English Channel, *Aquatic Microbial Ecology*, 42, 119-126, doi: 10.3354/ame042119.



- 1 Mincer, T.J., and A. C. Aicher (2016), Methanol production by a broad phylogenetic array of  
2 marine phytolankton, *Plos One*, 11, doi: 10.1371/journal.pone.0150820.
- 3 McDonald, I. R., and J. C. Murrell (1997), The methanol dehydrogenase structural gene *mxoF*  
4 and its use as a functional gene probe for methanotrophs and methylotrophs, *Applied*  
5 *and Environmental Microbiology*, 63, 3218-3224.
- 6 Muyzer, G., E. C. De Waal, and A. G. Uitierlinden (1993), Profiling of complex microbial  
7 populations by Denaturing Gradient Gel Electrophoresis analysis of Polymerase  
8 Chain Reaction-amplified genes coding for 16S rRNA, *Applied and Environmental*  
9 *Microbiology*, 59, 695-700.
- 10 Muyzer, G., T. Brinkhoff, U. Nubel, C. Santegoeds, H. Schäfer, and C. Waver (1998),  
11 Denaturing gradient gel electrophoresis (DGGE) in microbial ecology, in *Molecular*  
12 *Microbial Ecology Manual*, edited by J. D. van Elsas and F. J. de Bruijn, pp. 1–27,  
13 Kluwer Academic Publishers, Dordrecht, The Netherlands.
- 14 Nagata, T. (2008), Organic matter - bacteria interactions in seawater, in *Microbial Ecology of*  
15 *the Oceans*, edited by D. L. Kirchman, John Wiley and Sons, New Jersey.
- 16 Neufeld, J. D., Y. Chen, M. G. Dumont, and J. C. Murrell (2008), Marine methylotrophs  
17 revealed by stable-isotope probing, multiple displacement amplification and  
18 metagenomics, *Environmental Microbiology*, 10, 1526-1535, doi: 10.1111/j.1462-  
19 2920.2008.01568.x.
- 20 Neufeld, J. D., H. Schafer, M. J. Cox, R. Boden, I. R. McDonald, and J. C. Murrell (2007),  
21 Stable-isotope probing implicates *Methylophaga* spp and novel  
22 *Gammaproteobacteria* in marine methanol and methylamine metabolism., *The ISME*  
23 *Journal*, 1, 480 - 491, doi: 10.1038/ismej.2007.65.
- 24 Nightingale, P. D. (1991), Low molecular weight halocarbons in seawater, Ph.D. thesis,  
25 *University of East Anglia, Norwich, UK.*
- 26 Ogawa, H., and E. Tanoue (2003), Dissolved organic matter in oceanic waters, *Journal of*  
27 *Oceanography*, 59, 129-147.
- 28 Sargeant, S. L., J. C. Murrell, P. D. Nightingale, and J. L. Dixon (2016), Seasonal variability  
29 in microbial methanol utilisation in coastal waters of the western English Channel.,  
30 *Marine Ecology Progress Series*, 550, 53 - 64.
- 31 Sieburth, J. M., and M. D. Keller (1989), Methylaminotrophic bacteria in xenic nanoalgal  
32 cultures: Incidence, significance, and role of methylated algal osmoprotectants,  
33 *Biological Oceanography*, 6(5-6), 383-395, doi: 10.1080/01965581.1988.10749541.

- 1 Smith, D. C., and F. Azam (1992), A simple, economical method for measuring bacterial  
2 protein synthesis rates in seawater using  $^3\text{H}$ -leucine, *Marine Microbial Food Webs*, 6,  
3 107-114.
- 4 Sun, J., L. Steindler, J. C. Thrash, K. H. Halsey, D. P. Smith, A. E. Carter, Z. C. Landry, and  
5 S. J. Giovannoni (2011), One carbon metabolism in SAR11 pelagic marine bacteria,  
6 *Plos One*, 6, doi: 10.1371/journal.pone.0023973.
- 7 Taubert, M., C. Grob, A. M. Howat, O. J. Burns, J. L. Dixon, Y. Chen, and J. C. Murrell  
8 (2015), *coxF* encoding an alternative methanol dehydrogenase is widespread in  
9 coastal marine environments, *Environmental Microbiology*, 17, 3937 - 3948, doi:  
10 10.1111/1462-2920.12896.
- 11 Williams, J., R. Holzinger, V. Gros, X. Xu, E. Atlas, and D. W. R. Wallace (2004),  
12 Measurements of organic species in air and seawater from the tropical Atlantic,  
13 *Geophysical Research Letters*, 31, doi: 10.1029/2004GL020012.
- 14 Wilson, S. M., M. P. Gleisten, and T. J. Donohue (2008), Identification of proteins involved  
15 in formaldehyde metabolism by *Rhodobacter sphaeroides*, *Microbiology*, 154, 296-  
16 305, doi: 10.1099/mic.0.2007/011346-0.
- 17 Yang, M., P. D. Nightingale, R. Beale, P. S. Liss, B. Blomquist, and C. Fairall (2013),  
18 Atmospheric deposition of methanol over the Atlantic Ocean, *Proceedings of the*  
19 *National Academy of Sciences USA*, 110, 20034 - 20039, doi:  
20 10.1073/pnas.1317840110.
- 21 Yang, M., R. Beale, P. Liss, M. Johnson, B. Blomquist, and P. D. Nightingale (2014), Air-sea  
22 fluxes of oxygenated volatile organic compounds accross the Atlantic Ocean,  
23 *Atmospheric Chemistry and Physics*, 14, 7499-7517, doi: 10.5194/acpd-14-8015-  
24 2014.

1 **Figure and Table legends.**

2 **Figure 1.** Remotely sensed MODIS-Aqua chlorophyll *a* composite image of the Atlantic  
3 Ocean from November 2009 (image courtesy of NEODAAS). White squares represent  
4 sampling points and [circles-ovals](#) indicate samples within different oceanic provinces,  
5 labelled with province names NT (northern temperate), NSG (northern subtropical gyre),  
6 NTG (northern tropical gyre), EQU (equatorial upwelling), SG (southern gyre), ST (southern  
7 temperate).

8  
9 **Figure 2.** Variability in a) microbial methanol dissimilation rates (using the specific activity  
10 of  $^{14}\text{CH}_3\text{OH}$ ) and b) bacterial leucine incorporation (BLI), in surface waters of the Atlantic  
11 Ocean. Rates were determined from pre-dawn ( $\blacklozenge$  solid line) and solar noon ( $\blacklozenge$  dashed line)  
12 CTD casts. Error bars represent  $\pm 1$  s.d. of triplicate samples, dashed vertical lines indicate  
13 Atlantic province boundaries.

14  
15 **Figure 3.** Average depth profiles in Atlantic provinces for a) microbial methanol  
16 dissimilation (using the specific activity of  $^{14}\text{CH}_3\text{OH}$ ) and b) bacterial leucine incorporation  
17 (BLI) in pre-dawn waters. Error bars represent  $\pm 1$  s.d. of variation within the province,  
18 province averages derived from NT (n = 5), NSG (n = 5), NTG (n = 3), EQU (n = 4), SG (n =  
19 5) and ST (n = 3), except for BLI where there is no data from the ST.

20  
21 **Figure 4.** Microbial methanol (a) dissimilation and (b) assimilation rates ( $\text{nmol l}^{-1} \text{h}^{-1}$ ) in the  
22 top 200 m of an Atlantic Meridional transect (contour plots).

23  
24 **Figure 5.** Changes in bacterial community composition (Order, identified using 16S rRNA  
25 gene sequencing) for a) 97 % PAR surface 5m, b) 33 % PAR 10-31m, c) 1 % PAR 15-54m  
26 and d) 200 m for different provinces (NT, NSG, NTG, EQU and SG) of the Atlantic Ocean.  
27 Analysis is based on a rarefied sample of 386 sequences per sample. Bacterial Orders  
28 individually contributing to less than 5% of the total sample sequences were pooled together  
29 into 'Others (<5%)' for clarity. Where  $\blacksquare$  *Prochlorococcus*,  $\blacksquare$  *Alteromonadales*,  $\square$   
30 SAR11 clade,  $\blacksquare$  *Oceanospirillales*,  $\blacksquare$  *Rhodospirillales*,  $\blacksquare$  *Flavobacteriales*,  $\blacksquare$   
31 *Rhodobacterales*,  $\blacksquare$  *Sphingomonadales*,  $\blacksquare$  *Synechococcus*,  $\blacksquare$  *Acidimicrobiales*,  $\blacksquare$   
32 Order III *Incertae Sedis*,  $\blacksquare$  SAR324 clade (Marine group B),  $\blacksquare$  uncultivated bacterium,  $\blacksquare$   
33 other bacteria individually comprising <5%.

1  
2 **Figure 6.** Non-metric multi-dimensional scale (MDS) plots of (a) a Bray-Curtis similarity  
3 matrix of the 16S rRNA gene sequences of the bacterial community, (b) a Euclidean distance  
4 matrix of environmental parameters (salinity, temperature, chl. a, primary productivity,  
5 inorganic nutrients, flow cytometry cell numbers, BLI) and (c) a Euclidean distance matrix of  
6 rates of methanol dissimilation. Dashed lines highlight significant sample grouping. Plots  
7 generated using PRIMER-E ([www.primers-e.com](http://www.primers-e.com)). For (a) and (b) ■ represents samples from  
8 200 m i.e. 0 % PAR.

9  
10  
11  
12 **Table 1.** Summary of rates of methanol uptake (dissimilation and assimilation), methanol  
13 concentrations, bacterial leucine incorporation (BLI) and production (BP), numbers of  
14 heterotrophic bacteria (BN), *Prochlorococcus* (Pros) and *Synechococcus* (Syns). “Values  
15 given are average ± standard deviation (range). NA denotes that data is not available.”

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

1  
2

Figure 1.






Figure 2.







Figure 3.

Figure 4.





**Figure 5.**







Figure 6.



(a)



(b)



(c)

**Table 1.** Summary of rates of methanol uptake (dissimilation and assimilation), methanol concentrations, bacterial leucine incorporation (BLI) and production (BP), numbers of heterotrophic bacteria (BN), *Prochlorococcus* (Pros) and *Synechococcus* (Syns). Values given are average  $\pm$  standard deviation (range). NA denotes that data is not available.

|  | Overall                        | Atlantic province              |                                |                                |                                |                                |                                |
|--|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|
|  |                                | NT                             | NSG                            | NTG                            | EQU                            | SG                             | ST                             |
| Methanol dissimilation<br>(nmol L <sup>-1</sup> h <sup>-1</sup> )                      | 0.45 $\pm$ 0.42<br>(0.01–1.68) | 0.69 $\pm$ 0.35<br>(0.22–1.50) | 0.99 $\pm$ 0.41<br>(0.15–1.68) | 0.18 $\pm$ 0.04<br>(0.10–0.25) | 0.11 $\pm$ 0.03<br>(0.07–0.17) | 0.24 $\pm$ 0.12<br>(0.01–0.45) | 0.20 $\pm$ 0.05<br>(0.11–0.27) |
| Methanol assimilation (x 10 <sup>-2</sup> )<br>(nmol L <sup>-1</sup> h <sup>-1</sup> ) | 0.51 $\pm$ 0.54<br>(0.00–2.24) | 0.54 $\pm$ 0.53<br>(0.00–2.23) | 0.53 $\pm$ 0.56<br>(0.17–1.51) | NA                             | 0.67 $\pm$ 0.66<br>(0.00–2.24) | 0.19 $\pm$ 0.16<br>(0.00–0.57) | NA                             |
| BLI (pmol L <sup>-1</sup> h <sup>-1</sup> )  | 9.4 $\pm$ 8.9<br>(0.5–60.2)    | 7.7 $\pm$ 4.0<br>(0.9–14.2)    | 9.7 $\pm$ 14.2<br>(1.0–60.2)   | 8.0 $\pm$ 4.3<br>(2.0–17.0)    | 13.7 $\pm$ 7.9<br>(0.6–26.4)   | 8.2 $\pm$ 9.5<br>(0.5–41.5)    | NA                             |
| <sup>a</sup> BP(TCF) (ng C L <sup>-1</sup> h <sup>-1</sup> )                           | 14.6 $\pm$ 13.8<br>(0.8–96.1)  | 11.9 $\pm$ 6.1<br>(1.5–22.0)   | 15.0 $\pm$ 21.9<br>(1.5–96.1)  | 12.4 $\pm$ 6.6<br>(3.2–26.3)   | 21.2 $\pm$ 12.2<br>(1.0–41.0)  | 12.7 $\pm$ 14.8<br>(0.8–64.3)  | NA                             |
| <sup>b</sup> BP (ECF) (ng C L <sup>-1</sup> h <sup>-1</sup> )                          | 4.8 $\pm$ 4.6<br>(0.3–31.6)    | 3.9 $\pm$ 2.0<br>(0.5–7.2)     | 4.9 $\pm$ 7.2<br>(0.5–31.6)    | 4.1 $\pm$ 2.2<br>(1.0–8.7)     | 7.0 $\pm$ 4.0<br>(0.3–13.5)    | 4.2 $\pm$ 4.9<br>(0.3–21.1)    | NA                             |
| Numbers of heterotrophic bacteria<br>(x10 <sup>5</sup> ) (cells mL <sup>-1</sup> )     | 6.5 $\pm$ 6.3<br>(1.4–82.6)    | NA                             | NA                             | 5.8 $\pm$ 2.0<br>(1.6–9.8)     | 8.8 $\pm$ 10.3<br>(1.4–82.6)   | 5.4 $\pm$ 4.4<br>(1.5–35.8)    | NA                             |
| Numbers of <i>Prochlorococcus</i> sp.<br>(x10 <sup>5</sup> cells mL <sup>-1</sup> )    | 1.12 $\pm$ 4.62<br>(0.0–4.62)  | 0.91 $\pm$ 0.07<br>(0.0–2.56)  | 0.89 $\pm$ 0.71<br>(0.0–4.21)  | 1.52 $\pm$ 1.23<br>(0.0–4.19)  | 1.67 $\pm$ 0.2<br>(0.0–4.62)   | 1.20 $\pm$ 0.01<br>(0.0–2.45)  | 0.35 $\pm$ 0.22<br>(0.0–2.33)  |
| Numbers of <i>Synechococcus</i> sp.<br>(x10 <sup>4</sup> cells mL <sup>-1</sup> )      | 1.64 $\pm$ 31.4<br>(0.0–31.4)  | 1.96 $\pm$ 3.61<br>(0.0–12.7)  | 0.18 $\pm$ 0.21<br>(0.0–0.93)  | 0.15 $\pm$ 0.17<br>(0.0–0.73)  | 1.34 $\pm$ 2.69<br>(0.0–12.8)  | 0.14 $\pm$ 0.13<br>(0.0–0.79)  | 8.30 $\pm$ 10.3<br>(0.02–31.4) |
| <sup>c</sup> Methanol (nM)   | 143 $\pm$ 82<br>(38–420)       | 110 $\pm$ 126<br>(38–420)      | 203 $\pm$ 38<br>(154–281)      | 193 $\pm$ 46<br>(148–278)      | 148 $\pm$ 37<br>(117–241)      | 110 $\pm$ 33<br>(58–176)       | 132                            |

<sup>a</sup>Theoretical conversion factor (TCF) 1.55 kg C mol leu<sup>-1</sup>, <sup>b</sup>empirical conversion factor (ECF) 0.51 kg C mol leu<sup>-1</sup>, <sup>c</sup>From Beale et al., (2013)