

Zero to moderate methane emissions in a densely rooted, pristine Patagonian bog - biogeochemical controls as revealed from isotopic evidence

Wiebke Münchberger^{1, 2}, Klaus-Holger Knorr¹, Christian Blodau^{1, †}, Verónica A. Pancotto^{3, 4}, Till
5 Kleinebecker^{2, 5}

¹Ecohydrology and Biogeochemistry Research Group, Institute of Landscape Ecology, University of Muenster, Heisenbergstraße 2, 48149 Muenster, Germany

²Biodiversity and Ecosystem Research Group, Institute of Landscape Ecology, University of Muenster, Heisenbergstraße 2, 48149 Muenster, Germany

10 ³Centro Austral de Investigaciones Científicas (CADIC-CONICET), B. Houssay 200, 9410 Ushuaia, Tierra del Fuego, Argentina

⁴Instituto de Ciencias Polares y Ambiente (ICPA-UNTDF), Fuegia Basket, 9410 Ushuaia, Tierra del Fuego, Argentina

⁵Institute of Landscape Ecology and Resources Management, Giessen University, Heinrich-Buff-Ring 26, 35392 Gießen, Germany

15 [†]deceased, July 2016

Correspondence to: Wiebke Münchberger (wiebke.muenchberger@uni-muenster.de), Klaus-Holger Knorr (kh.knorr@uni-muenster.de)

Abstract. Peatlands are significant global methane (CH₄) sources, but processes governing CH₄ dynamics have been predominantly studied on the northern hemisphere. Southern hemispheric and tropical bogs can be dominated by cushion-
20 forming vascular plants (e.g. *Astelia pumila*, *Donatia fascicularis*). These cushion bogs are found in many (mostly southern) parts of the world but could also serve as extreme examples for densely rooted northern hemispheric bogs dominated by rushes and sedges. We report highly variable summer CH₄ emissions from different microforms in a Patagonian cushion bog as determined by chamber measurements. Driving biogeochemical processes were identified from pore water profiles and carbon isotopic signatures. An intensive root activity throughout a rhizosphere stretching over 2 m depth accompanied by molecular
25 oxygen release created aerobic microsities in water-saturated peat leading to a thorough CH₄ oxidation (< 0.003 mmol L⁻¹ pore water CH₄, enriched δ¹³C-CH₄ by up to 10‰) and negligible emissions (0.09 ± 0.16 mmol CH₄ m⁻² d⁻¹) from *Astelia* lawns. In sparsely or even non-rooted peat below adjacent pools similar pore water profile patterns as obtained under *Astelia* lawns were found which emitted very few amounts of CH₄ (0.23 ± 0.25 mmol m⁻² d⁻¹). Below the *A. pumila* rhizosphere pore water concentrations increased sharply to 0.40 ± 0.25 mmol CH₄ L⁻¹ and CH₄ was predominantly produced by hydrogenotrophic
30 methanogenesis. Few *Sphagnum* lawns and – surprisingly – one lawn dominated by cushion-forming *D. fascicularis* were found to be local CH₄ emission hot spots with up to 1.52 ± 1.10 mmol CH₄ m⁻² d⁻¹ presumably as root density and molecular oxygen release dropped below a certain threshold. The spatial distribution of root characteristics supposedly causing such pronounced CH₄ emission pattern was evaluated on a conceptual level aiming to exemplify scenarios in densely rooted bogs. We conclude that presence of cushion vegetation as a proxy for negligible CH₄ emissions from cushion bogs needs to be

interpreted with caution. Nevertheless, overall ecosystem CH₄ emissions at our study site were probably minute compared to bog ecosystems worldwide and widely decoupled from environmental controls due intensive root activity of e.g. *A. pumila*.

1 Introduction

5 Peatland ecosystems are significant natural methane (CH₄) sources on the global scale responsible for about 10% of global annual CH₄ emissions (Aselmann and Crutzen, 1989; Mikaloff Fletcher et al., 2004). CH₄ has a 28-fold higher global warming potential over a 100-year time horizon compared to carbon dioxide (CO₂) (IPCC, 2014) and, thus, processes governing CH₄ dynamics in peatlands receive much attention to estimate global greenhouse gas emissions.

The slowdown of decomposition and accumulation of C in rainwater-fed peatlands (bogs) results from e.g. a high water table, anoxic conditions, recalcitrant peat forming litter, low temperatures, inactivation of oxidative enzymes and an accumulation of decomposition end-products (Beer and Blodau, 2007; Limpens et al., 2008; Bonaiuti et al., 2017). Once methanogenic conditions are established, CH₄ production is mainly controlled by substrate supply of labile organic matter (Hornibrook et al., 1997; Whalen, 2005). Then, slow diffusive transport along the concentration gradient from deep anoxic peat layers to the atmosphere leads to CH₄ release. In upper, unsaturated and oxic peat zones that typically extent only about few decimetres (Whalen, 2005; Limpens et al., 2008), CH₄ might get consumed by methanotrophic microbes (Chasar et al., 2000; Whalen, 2005; Berger et al., 2018). Consequently, water table position and fluctuations strongly control the amount of emitted CH₄ (Blodau and Moore, 2003; Whalen, 2005). While CH₄ oxidation suppresses CH₄ emissions in diffusion dominated systems, ebullition by fast release of gas bubbles or plant-mediated transport by aerenchymatic roots can substantially increase CH₄ emissions (Fechner-Levy and Hemond, 1996; Joabsson et al., 1999 and references therein; Chasar et al., 2000; Colmer, 2003; Whalen, 2005; Knoblauch et al., 2015; Burger et al., 2016; Berger et al., 2018). With a high share of ebullitive fluxes, vegetated or open water pools in peatlands are considered as strong CH₄ emitters (Hamilton et al., 1994; Blodau 2002; Burger et al., 2016) that have, however, received less attention than the vegetated surfaces (Pelletier et al., 2014). Pools can even turn the peatlands' C balance into a source (Pelletier et al., 2014), but examples of low emission pools have been also reported (Knoblauch et al., 2015).

Carbon isotopic signatures ($\delta^{13}\text{C}$) are a valuable tool for identifying mechanisms and pathways of CH₄ dynamics. Carbon isotopic signatures in pore water of northern hemispheric bogs (hereafter termed northern bogs) vary typically between -80 to -50‰ and -25 to +10‰ for CH₄ and CO₂, respectively (e.g. Whiticar et al., 1986; Hornibrook et al., 1997; Chasar et al., 2000; Hornibrook et al., 2000; Beer et al., 2008; Steinmann et al., 2008; Corbett et al., 2013). Depending on the available substrate for methanogenesis, it was mostly observed that either the acetoclastic or hydrogenotrophic pathway predominates, resulting in distinctive carbon isotope signatures of CH₄ (acetate fermentation $\delta^{13}\text{C}\text{-CH}_4 \sim -65$ to -50% , CO₂ reduction $\delta^{13}\text{C}\text{-CH}_4 \sim -110$ to -60%) (Whiticar et al., 1986). Methane is prevalingly produced by acetoclastic methanogenesis when fresh, labile organic matter is available whereas the production pathway shifts towards hydrogenotrophic methanogenesis when organic matter becomes increasingly recalcitrant (Hornibrook et al., 1997; Popp et al., 1999; Chasar et al., 2000; Conrad, 2005). A predominance of the latter pathway is indicated by strong fractionation between ¹²C/¹³C resulting in relatively depleted $\delta^{13}\text{C}$ -

CH₄ corresponding to a fractionation factor α_c between 1.055 and 1.090 (Whiticar et al., 1986) and a $\delta^{13}\text{C}$ -CH₄ isotopic signature following that of $\delta^{13}\text{C}$ -CO₂ (Hornibrook et al., 2000). Subsequent methanotrophy alters the isotopic signature of CH₄ by discriminating against ¹³C and the residual CH₄ remains enriched while produced CO₂ becomes depleted in ¹³C (Chasar et al., 2000; Popp et al., 1999). As both, acetoclastic methanogenesis and methanotrophy, result in a $\delta^{13}\text{C}$ -CH₄ signature enriched in ¹³C and a small apparent fractionation factor (Whiticar et al., 1986; Conrad, 2005), the acetoclastic pathway cannot be clearly separated from methanotrophic conditions based on carbon isotopes only (Conrad, 2005).

The majority of studies elucidating CH₄ dynamics in peatlands have been conducted on the northern hemisphere (e.g. Blodau, 2002; Limpens et al., 2008; Yu, 2012 and references in these), whereas only little research deals with CH₄ dynamics in bogs on the southern hemisphere (hereafter termed southern bogs) (Broder et al., 2015). Contrary to northern bogs, southern bogs are not only vegetated by *Sphagnum* mosses, dwarf shrubs or few graminoids such as the rush-like species *Tetroncium magellanicum*, but also by vascular plants (Kleinebecker et al., 2007) that can densely root upper peat layers (Fritz et al., 2011; Knorr et al., 2015). Another feature that can be found in both systems are pools with sparse or no vegetation (Fritz et al., 2011; González Garraza et al., 2018). The vascular plants, for instance of the genus *Astelia* (Asteliaceae), *Donatia* (Stylidiaceae) or *Oreobolus* (Cyperaceae), independently developed a cushion life form to protect from harsh climate in cold environments (Gibson and Kirkpatrick, 1985; Boucher et al., 2016). Bogs dominated by cushion-forming vascular plants (hereafter termed cushion bogs) can be found in many (mostly southern) parts of the world, for instance all along the high Andes (Coombes and Ramsay, 2001; Benavides et al., 2013; Fonkén, 2014) down to southernmost Patagonia (Ruthsatz and Villagran, 1991; Heusser, 1995; Kleinebecker et al., 2007; Grootjans et al., 2014), in highlands of eastern Africa (Dullo et al., 2017) as well as in New Guinea (Hope, 2014), New Zealand and Tasmania (Gibson and Kirkpatrick, 1985; Ruthsatz and Villagran, 1991) and some sub-Antarctic islands (Ruthsatz and Villagran, 1991). These cushion bogs could also be regarded as extreme examples for densely rooted northern bogs that are dominated by rushes and sedges.

To our knowledge, only two studies reported CH₄ emissions from southern vascular-plant dominated bogs with quite inconsistent results. Over a New Zealand bog dominated by the evergreen “wire rush” *Empodisma robustum*, CH₄ emissions exceeded those commonly reported from northern bogs (Goodrich et al., 2015), while emissions from a Patagonian cushion bog dominated by *Astelia pumila* were negligible (Fritz et al., 2011). Both, high and negligible CH₄ emissions, were mainly explained by the extensive aerenchymous roots of the prevailing plant species: The aerenchyma might function as a conduit for CH₄ from deep peat layers to the atmosphere in the first case, whereas in the second case pronounced O₂ supply by roots might have resulted in a thorough oxidation of pore water CH₄ and enhanced organic matter decomposition. However, Fritz et al. (2011) were the first to investigate CH₄ dynamics below *Astelia* lawns and determined emissions only sporadically without considering all predominant microforms and plant communities. Thus, general conclusions about CH₄ emissions from this type of ecosystem are so far restricted. Furthermore, more knowledge on biogeochemical processes throughout the peat column of *Astelia* lawns as a model plant for cushion-forming, aerenchymous plants and in other microforms is needed to extent our limited understanding of these first insights into CH₄ dynamics in cushion bogs. Despite the variety of bogs dominated by densely rooted, aerenchymous plants on the northern and southern hemisphere, there is some evidence from latest research

that very low CH₄ pore water concentrations could be a more common phenomenon in those ecosystems (Dullo et al., 2017; Agethen et al., 2018; Knorr et al., 2015). In general, promotion or suppression of CH₄ production should be determined by the ratio of root density and activity associated with O₂ release versus presence of labile root organic matter or exudates accompanied with O₂ consumption (Blodau, 2002; Agethen et al., 2018).

5 Here, we quantified austral summer CH₄ emissions from dominant microforms in a Patagonian cushion bog and examined possible environmental and biogeochemical controls. We hypothesized that (I) emissions from cushion plant vegetation dominated by *Astelia pumila* or *Donatia fascicularis* are negligible while pools and *Sphagnum* lawns emit CH₄ in considerable amounts, and that (II) pore water CH₄ and DIC concentration profiles and its carbon isotopic signatures below densely-rooted *Astelia* lawns reflect a distinct CH₄ oxidation effect contrary to sparsely- or even non-rooted peat below pools, and that (III)
10 besides by methanotrophy, isotopic composition of pore water CH₄ could also be affected by a predominance of acetoclastic methanogenesis throughout the rhizosphere of *Astelia* lawns shifting towards CO₂ reduction with peat depth and in sparsely- or even non-rooted peat below pools. Biogeochemical controls of CH₄ emissions from *Sphagnum* lawns were not subject of this study, as they have been intensively studied in northern bogs (e.g. Hornibrook et al., 1997; Beer and Blodau, 2007; Steinmann et al., 2008; Corbett et al., 2013) and even southern bogs with mixed vegetation in absence of notable cushion plant
15 coverage (Broder et al., 2015).

2 Material and Methods

2.1 Description of the study site

The study was conducted at a cushion bog on the Peninsula Mitre in southernmost Patagonia, Tierra del Fuego (Moat, 54°58'S, 66°44'W, Fig. 1). Peatlands of the Peninsula Mitre provide the nowadays rare opportunity to study carbon dynamics under
20 pristine conditions as they can be considered as largely undisturbed by human activities such as drainage, agriculture or elevated atmospheric nitrogen deposition (Kleinebecker et al., 2008; Fritz et al., 2011; Grootjans et al., 2014; Paredes et al., 2014). About 45% of this area are covered by peatlands (Iturraspe, 2012) while little is known about these ecosystems because of their poor accessibility (Iturraspe, 2012). The study site at Moat belongs to a complex system of sloping mires, blanket bogs, fens and – in coastal areas - cushion bogs (Iturraspe, 2012; Borromei et al., 2014). The studied cushion bog is located in
25 exposed proximity to the Beagle channel with harsh winds (Grootjans et al., 2014) resulting in an oceanic climate with average daily temperatures of 6°C and annual precipitation of 500 mm (Fritz, 2012).

Large areas of the cushion bogs in Moat are characterized by a comparatively flat micro-relief and covered by the cushion-forming plants *Astelia pumila* and *Donatia fascicularis*. The vegetation of the studied bog is composed of a mosaic of different plant communities characterized by specific plant species with lawns of *A. pumila* being the prevailing microform. *Sphagnum*
30 *magellanicum* or *D. fascicularis* grow in small lawns (patches of few square meters) at pool margins where *S. magellanicum* supposedly benefits from protection against desiccation. *Tetroncium magellanicum*, a rush-like herb that does not form cushions but presumably establishes aerenchymous roots as it tolerates inundation (von Mehring, 2013), is associated

predominantly with *Sphagnum* lawns but also with *Astelia* and *Donatia* lawns or sporadically inhabits pools. These sparsely vegetated pools of different size (< 0.5 m² - ~10 m²) and ~0.5 m depth are embedded in *Astelia* lawns. Pool sediments are inhabited by cyanobacteria (Arsenault et al., 2018; González Garraza et al., 2018) and submerged *Sphagnum* mosses such as *Sphagnum cuspidatum* (Kleinebecker et al., 2007; Kip et al., 2012). Peat formation started ~11000 years ago (Borromei et al., 2014), and previously, the bog was dominated by *S. magellanicum* while *A. pumila* invaded the area around 2600 years before present as determined by pollen analyses (Heusser, 1995). The peat below *Astelia* lawns is densely rooted with on the average 2.15 g DW L of peat⁻¹ down to 1.7 m (Fritz et al., 2011). Mean root lifetimes of *A. pumila* has been estimated to be ~3-4 years indicating a high root turnover (Knorr et al., 2015).

2.2 Sampling and analysis of solid peat and root biomass

10 Peat coring in *Astelia* and *Sphagnum* lawns (data not shown) was done using a Russian peat corer (Eijkelkamp Agrisearch Equipment, Giesbeek, the Netherlands) down to a maximum depth of 7.5 m in *Astelia* lawns. Roots of *A. pumila* in *Astelia* lawn cores were sorted out and treated separately. The density of *D. fascicularis* root biomass was determined by cutting three 0.1 x 0.1 m sods down to a depth of 0.4 m. Peat and root biomass samples were oven dried at 70°C until constant weight to calculate peat bulk density and porosity as well as *D. fascicularis* root density. Total C and N contents together with abundance of ¹⁵N and ¹³C in the solid peat were determined using an elemental analyser (EA 3000, EuroVector, Redavalle, Italy) connected to an isotope ratio mass spectrometer (IRMS, NU instruments, Wrexham, UK).

2.3 Environmental variables

Environmental variables were determined during two measurement campaigns in 2015 and 2016 at half-hourly intervals. Photosynthetic active radiation (PAR, HOBO S-LIA-M003, Onset, USA, up to 2500 μmol m⁻² s⁻¹) and air temperature (HOBO S-TMB-M0x, Onset, USA) were recorded by a weather station (HOBO U30 NRC, Onset, USA) in both years during austral summer months. Water table fluctuations at two replicate sites in *Astelia* lawns (Levellogger Edge 3001 and Barologger Edge, Solinst, Canada; both installed in perforated PVC tubes) and soil temperature in four depths of 0.05, 0.1, 0.3 and 0.5 m (HOBO TMCx-HD and HOBO U-12-008, Onset, USA) were measured continuously.

2.4 Chamber measurements and analyses of soil-atmosphere CH₄ fluxes

25 2.4.1 Field measurements

The closed chamber technique was used to determine CH₄ fluxes during two measurement campaigns in austral summer from December 2014 to March 2015 and in February and March 2016. One week prior to measurements, PVC collars with a height of 0.2 m were permanently installed ~0.15 m into the peat of the dominant microforms characterized by a specific plant species

and a particular plant community (*Astelia* lawns: 2015 N = 3, 2016 N = 5; *Sphagnum* lawns: 2015 N = 3, 2016 N = 5; *Donatia* lawns: 2016 N = 3). The exact installation depth and the micro-relief of the surface within each collar was determined repeatedly to calculate the exact headspace volume for CH₄ flux estimation. The collars were equally arranged around three wooden platforms constructed in January 2014 to minimize disturbance during measurements and distributed over the study site to account for spatial variability. Different numbers of replicates between the two campaigns and the different microforms are the result of logistical constraints.

A cylindrical, transparent chamber with a basal area of 0.13 m² and a height of 0.4 m was used for soil-atmosphere flux measurements. The chamber was equipped with a fan to ensure mixing of the headspace air, a PAR sensor (HOBO S-LIA-M003, Onset, USA) and a temperature sensor (HOBO S-TMB-M0x, Onset, USA). A second temperature sensor recorded ambient temperature (HOBO S-TMB-M0x, Onset, USA) and all sensor data were logged in 1s-intervals to a data logger (HOBO U30 NRC, Onset, USA). This set-up allowed us to control air temperature inside the chamber within approximately 3°C deviation of the ambient air temperature. If necessary, ice packs were installed inside the chamber to avoid temperature increase. Due to strong wind conditions we decided against the installation of a vent tube. An opening in the top of the chamber avoided overpressure during chamber placement. This opening was closed immediately after the chamber was gently placed on a collar for at least 3 minutes to conduct measurements. Collars were equipped with a water-filled rim to ensure a gas-tight seal between chamber and collar during measurements. Pool fluxes were performed with a floating chamber of identical dimension and design on four pools located at each platform. The chamber wall extended approximately 0.04 m into the water during measurements.

All measurements were performed under a broad range of irradiance between 7:00 am to 22:00 pm (local time) to determine diurnal variations in CH₄ fluxes. At least once during each sampling period of two to three consecutive days, all selected microforms and pools were measured under dark conditions. Opaque, reflective material was used to cover the chamber for dark conditions. Measurements were taken every two weeks during the first campaign in 2015 (N = 405 measurements) and during two occasions in 2016 (N = 132 measurements). A more regular measurement routine throughout the whole season was constrained by harsh weather conditions and the remoteness of our study site.

The chamber was connected by a 2 mm inner diameter polyethylene tubing to a gas analyser (Los Gatos Ultraportable Greenhouse Gas Analyser 915-001, Los Gatos Research) to record the increase of CH₄ concentrations over time at a rate of 1 hz. Between measurements, atmospheric background concentrations were achieved inside the chamber. The gas analyser was equipped with an external pump providing a flow rate of 2 L min⁻¹ and pumping the analysed gas back to chamber creating a closed system. Prior to each campaign, the instrument was recalibrated.

We estimated the cover of plant species within each collar to the nearest 5% and counted the number of *T. magellanicum* shoots to further characterize the different microforms (Table 1). In 2015, vegetation surveys were conducted at the beginning and at the end of the measurement campaign. Since no change in vegetation was observed, surveys were conducted only once during the campaign in 2016. For each collar, the position above the water table was determined once during each sampling

period of consecutive days to account for different positions in the micro-relief and to determine a specific water table position. Additionally, we determined the area of each individual microform patch in which a collar had been installed.

2.4.2 CH₄ flux calculation and statistical analyses

5 Soil-atmosphere CH₄ fluxes were calculated from the gas concentration increase over time within the chamber using the software package (MATLAB Release R2015a) routine described in Eckhardt and Kutzbach (2016) and Kutzbach et al. (2007). Gas concentration was modelled either as a linear or an exponential function of time. Models performance was compared using Akaike's Information Criterion (AIC) as a measure of goodness-of-fit. This routine resulted in 90% of cases in a linear function of time (N = 485) and in 10% of cases in an exponential function of time (N = 52).

10 We visually inspected all concentration increases over time and none of the measurements showed a stepwise concentration increase indicative of ebullient events. Of the 3 minutes measurement time, 50 - 180 s (mostly around 90 s) were selected for the CH₄ flux calculation to exclude unstable conditions due to e.g. chamber placement particularly at the beginning of the measurement. Fluxes with a slope not significantly different from 0 (tested with an F-Test, $p > 0.05$) were set to zero (20% of cases, N = 105).

15 Normal distributions of CH₄ flux datasets were checked by a Kolmogorov-Smirnov test. Due to non-normality, the Kruskal-Wallis analysis of variance followed by multiple comparison U-tests adjusted by the Bonferroni method were applied to determine significant differences between CH₄ fluxes of different microforms and collars. The relationships between CH₄ fluxes and environmental variables (soil temperature, water table depth fluctuations) as well as additional features of individual collars were evaluated by Spearman's rank correlation (Table 2). A response of CH₄ emissions to the investigated environmental variables was identified only for a few individual collars. Therefore, no seasonal or annual ecosystem emissions were calculated. Statistical analyses were performed using Matlab (MATLAB Statistics Toolbox Release R2017a). All flux data were reported as mean \pm standard deviation.

2.5 Depth profiles of peat pore water concentrations

2.5.1 Sampling and field measurements

25 Multilevel piezometers (MLP), as described in Beer and Blodau (2007), were installed during both measurement campaigns in austral summer 2015 and 2016 to determine pore water concentrations in depth profiles. The MLPs provided a spatial sampling resolution of either 0.1 or 0.2 m and were equipped with diffusive equilibration samplers made of permeable silicone tubes providing 4 mL gas sample volume together with a 5 mL crimp-vial filled with deionized water and covered with a permeable membrane (Supor-200 0.2 μ , Pall corporation, Pall life sciences). To collect gas samples, MLPs were stepwise retrieved and the gas volume extracted from the equilibration sampler with a 3 mL syringe was transferred into nitrogen-flushed 5 mL crimp vials. The crimp vials were capped with a butyl stopper with an aluminium crimp seal. MLPs were installed 30 in *Astelia* lawns and pools in three replicates at each platform where closed chamber measurements were performed.

In January 2015, MLPs were installed in *Astelia* lawns (0.2 m resolution, 2 m depth) and pools (0.1 m resolution 0-1 m, 0.2 m resolution below down to a maximum depth of 4 m) for five weeks equilibration time. From gas samples pore water concentrations of CH₄, dissolved inorganic carbon (DIC) and hydrogen (H₂) were determined. To measure sulfate concentrations, a potential electron acceptor originating from sea spray and affecting competitiveness of methanogenesis (Broder et al., 2015), a pore water subsample obtained from each crimp-vial was transferred to a storage vial and stored frozen until analysis. An aliquot of each remaining pore water sample was measured in-situ with a pH/EC-meter (Combo HI 98129/130, Hannah instruments, Germany).

In January 2016, MLPs were installed to a depth of 3 m in pools (resolution as in 2015) for three weeks equilibration time and afterwards again in *Astelia* lawns (0.1 m resolution in transition zone from rooted to non-rooted peat in 1-2 m depth) for nine weeks of equilibration time. In 2016, equilibration samplers were used to collect CH₄ and CO₂ samples and determine their carbon stable isotopic signatures. Pore water samples obtained from crimp-vials were used to measure O₂ concentrations in-situ with a planar trace oxygen minisensor (Fibox 3, Presens, Germany).

2.5.2 Analytical procedures

Gas samples were transported to Germany and analysed within four weeks. Gaseous CH₄ and CO₂ concentrations were measured with a gas chromatograph (8610C, SRI Instruments, USA) equipped with a methaniser and flame ionization detector (FID). Hydrogen concentrations were analysed on a H₂-Analyzer (Ametek TA 3000 H₂ Analyser, Trace Analytical TA 3000r). Sulfate concentrations were obtained by ion chromatography (883 Basic IC plus, Metrohm, Herisau, Switzerland).

Stable carbon isotopic signatures of CH₄ and CO₂ were simultaneously determined by Cavity Ring-Down Spectroscopy (CRDS; Picarro G2201-I connected to a small sample isotope module (SSIM, model A0314), Picarro Inc., USA). As this set-up required a minimum sample volume of 20 mL, samples were diluted with nitrogen prior measurements. The instrument was calibrated in the beginning of every measurement day using two working standards of CH₄ (1000 ppm, -42.48‰) and CO₂ (1000 ppm, -31.07‰). Isotopic signatures are given in δ-notation relative to Vienna Peedee belemnite (VPDB). As samples were stored for several weeks and δ¹³C-CO₂ values were biased in case of high CH₄ concentrations in the sample, a correction procedure was applied following Berger et al. (2018).

2.5.3 Calculations and statistical analyses

Pore water concentrations of CH₄, dissolved inorganic carbon (DIC) and H₂ were re-calculated from gaseous concentrations obtained from equilibration samplers by applying Henry's law following Eq. (1):

$$c = K_H * p \tag{1}$$

where c is the concentration in mol L⁻¹, p the pressure in atm and K_H the Henry-constant that was corrected according to Sander (1999) for mean soil temperatures of 10°C in February 2015 and 2016 as well as 7°C in April at a depth of 0.5 m. DIC

concentrations at prevalent pH conditions were calculated with respect to carbonate speciation considering equilibrium constants following Stumm and Morgan (1996). All pore water data were reported as mean \pm standard deviation (N = 3).

Zones of CH₄ production and consumption in the peat column were visually identified based on observed pore water concentration gradients. For a quantitative evaluation of pore water concentration gradients, steady-state conditions as well as a dominance of diffusive gas transport would have to be assumed; this may only partly apply to the system under study here, given the large root biomass. Nevertheless, we additionally applied a software routine using inverse modelling (PROFILE, Berg et al., 1998) for zone identification. This modelling approach supported the results obtained visually, but provided further rough estimates for production and consumption zones due to the complex diffusivity in the rhizosphere of highly rooted peat and, thus, results are presented in the appendix only.

10 An apparent isotopic fractionation factor α_c was determined (Whiticar et al., 1986; Hornibrook et al., 2000) to assess the predominant methanogenic pathway and / or methanotrophic activity following Eq. (2):

$$\alpha_c = \frac{\delta^{13}C_{CO_2} + 1000}{\delta^{13}C_{CH_4} + 1000} \quad (2)$$

3 Results

3.1 Characteristics of solid peat and root biomass

15 Roots of *A. pumila* were present in the upper profile down to a depth of 1.8 m (rhizosphere) in *Astelia* lawn cores. Throughout the rhizosphere, we observed a highly decomposed and amorphous peat supposedly originating from *A. pumila* that developed above *Sphagnum* peat. Peat below *Sphagnum* lawns was continuously formed predominantly by *Sphagnum*. Total C and N contents in the peat together with natural abundance of ¹⁵N and ¹³C are given in Table 3. Root density of *D. fascicularis* was 0.014 \pm 0.010 g DW L of peat⁻¹ at 0.35 m depth while root density of *A. pumila* was not determined once again as it has been
20 already quantified with > 4 g DW L of peat⁻¹ in around 0.5 m depth (Fritz et al., 2011).

3.2 Environmental conditions and potential controls on CH₄ fluxes

The study site was characterized by daily mean PAR values up to ~700 $\mu\text{mol m}^{-2} \text{s}^{-1}$ with maximum values exceeding 2000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and daily mean air temperatures ranging from -0.5°C to 17°C during the measurement campaigns in austral summer from January to April (Fig. 2). Soil temperature in summer reached maximum daily average values of 15°C and 12°C
25 at 0.05 m depth and 0.5 m depth, respectively. Concurrently, the water table fluctuated close to the surface between -0.03 m and -0.23 m and differed consistently about few centimetres between the two water table measurement sites in *Astelia* lawns. CH₄ fluxes approached zero in many cases, but differed significantly from zero down to absolute values of 0.01 mmol CH₄ m⁻² d⁻¹. Fluxes below absolute values of 0.01 mmol CH₄ m⁻² d⁻¹ did not differ significantly from zero and were therefore set to zero. For collars exhibiting low magnitude fluxes neither seasonal trends (data not shown) nor a relationship to water table and

soil temperature fluctuations could be identified (Table 2). Yet, significant relationships between water table fluctuations or variations in soil temperatures were established for those few lawns with considerable emissions (Table 2).

3.3 Soil-atmosphere CH₄ fluxes and features of microforms possibly affecting CH₄ emissions

Summer CH₄ emissions in a South Patagonian cushion bog were highly variable among the dominant microforms *Astelia* 5 lawns, *Sphagnum* lawns and *Donatia* lawns characterized by specific plant species as well as pools and showed a pronounced spatial pattern even within microforms. Compared to all other microforms, only *Sphagnum* lawns showed considerable emissions with 1.52 ± 1.10 mmol CH₄ m⁻² d⁻¹ while *Astelia* lawns emitted nearly no CH₄ with 0.09 ± 0.16 mmol m⁻² d⁻¹ (Fig. 3). Methane emissions from pools were also low (0.23 ± 0.25 mmol m⁻² d⁻¹) and not significantly different from intermediate *Donatia* lawn fluxes (0.66 ± 0.96 mmol CH₄ m⁻² d⁻¹). During some occasions, we determined even significant, negative CH₄ 10 fluxes, mainly from *Astelia* lawns (Fig. 3a, Table 2). Most of these negative fluxes as well as the majority of fluxes (41%) that were set to zero were obtained from one individual collar throughout the whole measurement campaigns precluding the possibility of a measurement artefact.

As indicated by high standard deviations, CH₄ emissions between replicates of all microforms were not consistent. Field observations suggested that CH₄ emissions of a microform were not only controlled by the predominant plant species, but 15 might have been associated with additional features of the respective microform. Therefore additional microform features (i.e. number of *T. magellanicum* shoots, cover of *D. fascicularis*, extend of the microform in which a specific collar was installed) were assessed. However, as we did not expect such small-scale spatial variability between emissions of individual collars, the results of this survey given in Table 2 are of rather explorative, preliminary character as the low number of replicates (3 to 5 20 collars) in combination with several outliers barely allows any sound statistical analysis. Nevertheless, collars with elevated or comparatively high emissions had some features in common: Emissions from one *Astelia* lawn were with ~ 0.3 mmol CH₄ m⁻² d⁻¹ (significantly) higher in both sampling years compared to fluxes measured from the other *Astelia* lawns. This *Astelia* lawn collar with elevated emissions was characterized by (I) the presence of many *T. magellanicum* shoots together with a high share of *D. fascicularis* and (II) placed in a lawn with small extent surrounded by a mosaic of small pools and *D. fascicularis* patches (both smaller than 1 m²). Surprisingly, also one *Donatia* lawn was a substantial CH₄ source of 2.10 ± 0.14 25 mmol m⁻² d⁻¹, even exceeding the highest emissions observed from *Sphagnum* lawns on all measurement occasions in 2016. Similar to the *Astelia* lawn with elevated emissions, the collars of *Donatia* and *Sphagnum* lawns with high emissions were characterized by (I) a high amount of *T. magellanicum* shoots and (II) placed in lawns surrounded by small pools and other patches of *D. fascicularis* with no *A. pumila* nearby. Emissions obtained from one *Sphagnum* lawn were with less than 0.3 mmol CH₄ m⁻² d⁻¹ in both sampling years significantly lower compared to emissions from other *Sphagnum* lawns. Vice versa, 30 this collar with low *Sphagnum* emissions was characterized by (I) the lowest number of *T. magellanicum* shoots among all *Sphagnum* lawn collars and (II) installed in a small *Sphagnum* lawn surrounded by *A. pumila*.

3.4 Pore water CH₄ and DIC concentration profiles

Pore water profiles in peat columns below *Astelia* lawns and pools showed similar trends in pore water concentrations and during both sampling years. CH₄ concentrations were almost zero in the upper pore water profile below *Astelia* lawns (< 0.003 mmol L⁻¹) down to a depth of around 1.5 m, which corresponds to the zone where the rhizosphere was most pronounced (Fig. 4a). Below this depth, CH₄ concentrations sharply increased up to 0.25 ± 0.08 mmol L⁻¹ at 3 m depth. CH₄ concentrations in pore water profiles below pools resembled profiles obtained under *Astelia* lawns on elevated levels. Throughout the upper profile down to 1.5 m, CH₄ concentrations were ~0.03 mmol L⁻¹ with a peak around 0.3 m depth (0.06 ± 0.06 mmol L⁻¹) and increased steeply to 0.40 ± 0.25 mmol L⁻¹ in 3 m depth. Maximum concentrations in comparable depths reached similar levels in both sampling years below *Astelia* lawns, but maximum concentrations below pools were about 3 times higher in 2015 compared to 2016.

DIC predominantly occurred as dissolved CO₂ because of the low pH ranging from 3.36 to 4.77. Contrary to CH₄, DIC concentrations increased constantly with depth from around 1 mmol L⁻¹ near the surface to 2.60 ± 1.0 mmol L⁻¹ in 2 m depth below *Astelia* lawns and 2.94 ± 1.1 mmol L⁻¹ in 3 m depth below pools (Fig. 4b). With depth, DIC converged with CH₄ concentrations, which was reflected by DIC:CH₄ ratios that were extremely high in the rhizosphere below *Astelia* lawns exceeding 100 (Fig. 4c). Beneath the rhizosphere, ratios steeply approached values below 40. Under pools, ratios slightly decreased with depth but were mostly around 40 down to 1.5 m with two distinct peaks of very little CH₄ at the surface and around 0.8 m depth. In deep peat layers below the rhizosphere, DIC:CH₄ ratios detected under *Astelia* lawns and pools converged and reached lowest ratios of ~10 and ~5, respectively.

Visual inspection of concentration profiles but also modelling of CH₄ production and consumption rates indicated a predominance of CH₄ consumption throughout the whole rhizosphere below *Astelia* lawns and in roughly corresponding depths under pools (Fig. 4a, Fig. 6). Maximum CH₄ consumption was identified in the lower rhizosphere around 1.5-2 m where the increase in CH₄ concentration was most pronounced. Deep peat layers below the rhizosphere of *Astelia* lawns and similar depths below pools were considered as CH₄ sources. In these depths, the high concentration levels of CH₄ (up to 0.40 ± 0.25 mmol L⁻¹) sustain substantial upward diffusion of CH₄ following the concentration gradient into the consumption zone throughout the rhizosphere of *Astelia* lawns and in corresponding depths below pools.

3.5 Carbon isotopic values in pore water and apparent fractionation

Values of δ¹³C in CH₄ did not show a clear depth trend below *Astelia* lawns and ranged in the upper profile down to 2 m depth between -87.2 ± 10.1 to -72.1 ± 10.3‰ (Fig. 4d). Isotopic δ¹³C-CH₄ values below pools became less negative with depth from -93.4 ± 9.2 to -73.7 ± 1.2‰ with a minimum peak in upper peat layers around 0.3 m of -101.0 ± 5.6‰. In the upper meter of the profile, values of δ¹³C-CH₄ were thus up to 25‰ more negative below pools compared to *Astelia* lawns while the isotopic values of both profiles converged in deep peat layers below 2 m at -80‰. Values of δ¹³C in CO₂ below *Astelia* lawns became more negative with depth in the upper profile down to 1 m reaching values as low as -32.4 ± 0.8‰ and increased to -19.5 ±

1.0‰ in 3 m depth. Below pools, values became less negative with depth throughout the whole profile from -21.2 ± 0.4 to -6.0 ± 1.5 ‰ (Fig. 4e).

Apparent fractionation factors varied only slightly and showed opposite trends below the two microforms. Values increased slightly from 1.060 to 1.066 below *Astelia* lawns, peaking at 1.045 in 0.5 m depth. Below pools, apparent fractionation factors
5 decreased with depth from 1.079 to 1.073 with a maximum of 1.091 around 0.3 m depth (Fig. 4f).

3.6 Hydrogen, oxygen and sulfate concentrations in pore water profiles

Hydrogen concentrations were mostly below 3 nmol L^{-1} below *Astelia* lawns while they were elevated below pools, ranging between 3 and 10 nmol L^{-1} (Fig. 5a). Maximum concentrations were reached in the upper profile of *Astelia* lawns with up to $11.83 \pm 17.88 \text{ nmol L}^{-1}$ and, besides a peak at the surface, below pools from 1 to 2 m depth with up to $41.61 \pm 64.63 \text{ nmol L}^{-1}$.
10 Molecular oxygen concentrations were mostly below 5% saturation and tended to be higher in the upper profile down to 2 m below *Astelia* lawns compared to pools (Fig. 5b). Sulfate concentrations were similar below both, *Astelia* lawns and pools, and reached highest values near the surface with up to $16.43 \pm 11.85 \text{ } \mu\text{mol L}^{-1}$. With depth, concentrations decreased and approached zero below 1 m (Fig. 5c).

4 Discussion

15 This study is among the first dealing with CH_4 dynamics in an austral, vascular-plant dominated cushion bog. We aimed to reveal patterns of CH_4 emissions and their environmental as well as potential belowground biogeochemical controls. We furthermore attempted to elucidate the CH_4 dynamics in the peat below the four predominant microforms on a conceptual level based on the results presented here and in previous studies.

4.1 Environmental controls on CH_4 emissions

20 Summer CH_4 emissions of a South Patagonian cushion bog dominated by *A. pumila* were low albeit spatial heterogeneity was pronounced with small patches of *Sphagnum* or *Donatia* lawns being local emission hot spots. The variation of CH_4 fluxes was mainly controlled by the dominant plant species of each lawn microform which emitted significantly different amounts of CH_4 . We found only few significant responses of CH_4 fluxes to water table fluctuations or variations in soil temperatures (Table 2). Such weak coupling between environmental variables and CH_4 emissions is not surprising as fluxes from most
25 individual collars were relatively low. Furthermore, the water table fluctuated only slightly near the surface (Fig. 2) and may thus not serve as a primary control on CH_4 emissions. Stronger effects may only be expected when amplitude and duration of fluctuations affect peat redox conditions substantially (Blodau and Moore, 2003; Knorr et al., 2009). Under such stable moisture conditions and as long as methanogenesis is not limited by substrate supply, the sensitivity of methanogenic microbial consortia to temperature should become apparent (Whalen, 2005). Variations in landscape CH_4 fluxes have indeed been related

to soil temperature previously (Rinne et al., 2007; Jackowicz-Korczyński et al., 2010; Goodrich et al., 2015) which is in accordance with the present study. The temporal variability of CH₄ emissions over the two investigated seasons was rather low (data not shown) due to low seasonality in both, temperature and precipitation, in the study region. Measurement campaigns extending over shoulder seasons and comparing years of contrasting weather conditions would be necessary to reveal the impact of more pronounced water table fluctuations or temperature regimes on CH₄ fluxes in South Patagonian cushion bogs.

4.2 *Astelia* lawns – zero emission scenario

Emissions from *Astelia* lawns were minute ($0.09 \pm 0.16 \text{ mmol m}^{-2} \text{ d}^{-1}$) and significantly lower than from all other microforms (Fig. 3) verifying our first hypothesis. The near-zero emissions were well explained by near-zero CH₄ concentrations ($< 0.003 \text{ mmol L}^{-1}$) in the rhizosphere down to around 1.8 m. Only below, concentrations increased sharply to 0.2 mmol L^{-1} in 2 m depth (Fig. 4a). The low CH₄ concentrations even turned some *Astelia* lawns into a small sink for atmospheric CH₄ as shown by negative fluxes that were obtained sporadically throughout the whole measurement campaigns, confirming earlier results (Kip et al., 2012). These findings agree with previous research by Fritz et al. (2011) who observed CH₄ emissions from *Astelia* lawns in a comparably low magnitude as presented here and presented a similar CH₄ concentration depth profile. Extremely low CH₄ concentrations have also been obtained by Dullo et al. (2017) in upper peat layers of an Ethiopian cushion bog. Nevertheless, in deep peat layers below the rhizosphere, CH₄ concentrations of our study approached magnitudes reported for Chilean bogs with mixed vegetation consisting of *Sphagnum* mosses and *A. pumila* (Broder et al., 2015). This is in the lower range of concentrations described from northern bogs mostly around 0.5 mmol L^{-1} (e.g. Blodau and Moore, 2003; Beer and Blodau, 2007; Beer et al., 2008; Corbett et al., 2013) but even reaching up to 5 mmol L^{-1} in comparable depths (Steinmann et al., 2008).

A CH₄ concentration profile as observed at our site yet suggests that in deep peat layers below the rhizosphere ($> 2 \text{ m}$) CH₄ was produced at substantial rates and comparably low oxidation, as indicated by DIC:CH₄ ratios notably smaller than 40 (Fig. 4c). Subsequently, CH₄ got almost completely oxidized on its diffusive way to the atmosphere as visualized on a conceptual level (Fig. 6). Ratios of DIC:CH₄ under methanogenic conditions in northern bogs are typically around 5 (e.g. Hornibrook et al., 1997; Blodau and Moore, 2003; Corbett et al., 2013), but can be as high as 30 (Steinmann et al., 2008). Notably higher ratios throughout the rhizosphere suggested methanotrophic activity and availability of either alternative electron acceptors or O₂ from aerenchymous roots suppressing methanogenesis (Colmer, 2003; Mainiero and Kazda, 2005; Knorr et al., 2008a, Fritz et al., 2011; Dullo et al., 2017). Diffusion and plant-mediated transport as pathways for upward CH₄ transport would have caused higher emissions and ebullition requires supersaturation of at least $\sim 350 \text{ mmol CH}_4 \text{ L}^{-1}$ in the pore water (Fechner-Levy and Hemond, 1996; Beer et al., 2008). The sharp drop in CH₄ concentrations from -2 to -1.5 m depth characterized this peat layer as a zone of strongest methanotrophic activity, as supported by modelled consumption zones.

Carbon isotopic values obtained from pore water profiles reflected rhizosphere processes, although the $\delta^{13}\text{C}\text{-CH}_4$ signature alone did not provide a clear indication for oxidative effects in the rhizosphere as proposed in our second hypothesis. A strong effect of only methanotrophy should result in a shift of the mean $\delta^{13}\text{C}\text{-CH}_4$ signature to more enriched values and distinct changes in the fractionation factor from lower to upper horizons. Methane isotopic signatures were close to -80‰ in deep peat layers while above 2 m depth values ranged from -90 to -70‰ throughout the rhizosphere (Fig. 4d) while the fractionation factor increased slightly (Fig. 4f). These surprisingly negative $\delta^{13}\text{C}\text{-CH}_4$ values throughout the rhizosphere along with a comparatively small shift and a wider standard variation of replicate samples are an unexpected result that needs further explanation.

We can only explain the pronounced variation in replicate $\delta^{13}\text{C}\text{-CH}_4$ values to be associated with the presence of aerenchymous roots and the mean $\delta^{13}\text{C}\text{-CH}_4$ depth pattern by a co-existence of aerobic and anaerobic microsites that would both support such strong variation within short spatial scales. The occurrence of small-scale aerobic microsites attached to roots releasing O_2 into water-saturated (peat) layers has been previously suggested (Colmer, 2003), also for peatlands on the northern hemisphere (Popp et al., 1999; Knorr et al., 2008a; Corbett et al., 2013). Life time and activity of *A. pumila* roots are highly dynamic (Knorr et al., 2015) and supposedly governed temporal and spatial expansion of microsites. Within these microsites, labile organic matter and CH_4 were rapidly consumed by aerobic respiration and methanotrophy resulting in CH_4 enriched in ^{13}C by up to 10‰ (Fig. 4d). This $\delta^{13}\text{C}\text{-CH}_4$ signature and the apparently small $^{12}\text{C}/^{13}\text{C}$ fractionation observed throughout the rhizosphere (Fig 4f) revealed the influence of a strongly fractionating process such as CH_4 oxidation. Although based on ^{13}C isotope fractionation alone CH_4 oxidation cannot be clearly separated from acetoclastic methanogenesis (Whiticar et al., 1986; Conrad, 2005), oxidative effects in the rhizosphere were very likely since diffusion or plant-mediated transport were excluded as sinks for CH_4 . Methanotrophy leaves CH_4 in the pore water enriched by ~10‰ while produced CO_2 becomes depleted (Chasar et al., 2000; Popp et al., 1999) as observed here throughout the rhizosphere with less negative signatures occurring only at greater depths (Fig. 4e). Also values of $\delta^{13}\text{C}\text{-CO}_2$ of less than -30‰ that were substantially lower than the source organic matter material around -26‰ (Table 3) can only be explained by the influence of ^{13}C depleted carbon from oxidation of ^{13}C depleted CH_4 . Substantial rates of root respiration were a further source for depleted CO_2 in the pore water (Corbett et al., 2013) as suggested by high DIC: CH_4 ratios mostly exceeding 100 throughout the prevalingly water-saturated rhizosphere (Fig. 4c). Nevertheless, O_2 release by roots still exceeded consumption, as demonstrated by O_2 saturation reaching up to 10% (Fig 5b). This O_2 supply may thus not only fuel CH_4 oxidation but also heterotrophic respiration using molecular oxygen or other electron acceptors regenerated by this O_2 input (Colmer, 2003; Mainiero and Kazda; 2005). Likely further sources of CO_2 could have been dissimilatory sulfate reduction in anaerobic microsites of surface peat layers down to 0.4 m (Fig. 5c) with sulfate originating e.g. from sea spray (Kleinebecker et al., 2008; Broder et al., 2015). Taking together these results obtained from CH_4 and DIC concentration profiles and its carbon isotopic signatures, the second hypothesis stating that pore water concentrations below *Astelia* lawns reflect a distinct CH_4 oxidation effect was confirmed.

Enriched CH_4 being only a leftover from CH_4 oxidation should result in a $\delta^{13}\text{C}\text{-CH}_4$ depth pattern following that of the $\delta^{13}\text{C}\text{-CO}_2$ profile (Hornibrook et al., 2000), but such pattern was not observed in our study (Fig. 4d and e). The surprisingly negative

$\delta^{13}\text{C}\text{-CH}_4$ signature throughout the rhizosphere is likely explained by production of few amounts of CH_4 in anaerobic microsites in absence of living roots. Either hydrogenotrophic or acetogenesis and subsequent acetoclastic methanogenesis cause large fractionation (Steinmann et al., 2008) and could add depleted CH_4 to the rhizosphere. As we did not quantify other parameters such as labile organic root matter or root exudates, acetate concentrations or its carbon isotopic signatures, we cannot clearly
5 determine the relative importance of both pathways for our study. Elevated H_2 concentrations in surface peat layers (Fig. 5a) indicated high H_2 production by fermentation compared to consumption by e.g. sulfate reduction or hydrogenotrophic methanogenesis, presumably due to high root litter input and root exudates (Knorr et al., 2009; Estop-Aragonés et al., 2013). But as acetogens can outcompete hydrogenotrophic methanogens at low temperatures especially when H_2 is available, (Kotsyurbenko et al., 2001), we propose that acetoclastic methanogenesis was the more important pathway here. Together with
10 the probably high amounts of labile organic matter available to serve as substrate for methanogenesis throughout 2 m of rhizosphere, this explanation would support our third hypothesis. With depth, root density presumably decreased (Fritz et al., 2011) and a slight increase in $^{12}\text{C}/^{13}\text{C}$ fractionation indicated a shift to a predominance of hydrogenotrophic methanogenesis and, below 2 m depth, absence of microsites (Fig. 4f) as α_c -values of replicate samples varied less and increased towards 1.065 (Whiticar et al., 1986; Conrad, 2005).

15 **4.3 Pools – low emission scenario**

Pools exhibited with $0.23 \pm 0.25 \text{ mmol CH}_4 \text{ m}^{-2} \text{ d}^{-1}$ second lowest emissions of the microforms under study (Fig. 3). Such low emissions were contrary to our first hypothesis and somewhat surprising since they have been previously reported to be similar to *Sphagnum* lawns (Fritz et al., 2011) and were expected to be relevant CH_4 sources because of prevailing anoxic conditions (Blodau, 2002). Moreover, other studies identified pools as hot-spots of CH_4 emissions from peatlands (Hamilton et al., 1994;
20 Pelletier et al., 2014; Burger et al., 2016). This was clearly not the case here, as emissions were in the same order of magnitude as pool fluxes in South Patagonian *Sphagnum* bogs (Lehmann et al., 2016).

Low emissions were associated with CH_4 pore water concentrations surprisingly resembling those below *Astelia* lawns on slightly elevated levels (Fig. 4a). These low pore water concentrations in sparsely- or even non-rooted peat below pools were not supported by our second hypothesis and require another explanation. The $\delta^{13}\text{C}\text{-CH}_4$ signature was depleted compared to
25 *Astelia* lawns down to 2 m depth, followed largely that of $\delta^{13}\text{C}\text{-CO}_2$ (Fig. 4d, e) and, thus, did not indicate a methanotrophic effect. As upward diffusion of CH_4 against the concentration gradient is impossible, we can only explain low pore water concentrations by lateral exchange of CH_4 . We suggest that root O_2 release by *A. pumila* controlled even CH_4 dynamics below adjacent microforms as visualized on a conceptual level in Figure 6. The thorough CH_4 oxidation below *Astelia* lawns may establish a lateral concentration gradient of CH_4 . Thereby, CH_4 from adjacent microforms may diffuse to the rhizosphere of
30 *Astelia* lawns where it was consumed. As the micro-relief is not pronounced at our study site, we propose that there is a large depth zone throughout the 2 m rhizosphere of *Astelia* lawns and in corresponding depths below pools with negligible water flow and a predominance of diffusive transport. In this depth zone, we observed a highly decomposed and amorphous peat as

previously described for cushion bogs (Ruthsatz and Villagran, 1991; Fritz et al., 2011) indicating a very high water permeability that would together with a high water table position facilitate such lateral exchange (Baird et al., 2016). The assumption of lateral exchange was supported by a peak in DIC:CH₄ ratios accompanied by very low CH₄ concentrations around 0.8 m below pools (Fig. 4c) inhibiting upward diffusion. Lateral root ingrowth and O₂ supply from adjacent *Astelia* lawns could furthermore suppress methanogenesis and keep CH₄ concentrations below pools low, in particular if pools are small.

Patterns of carbon isotopic signatures, levels of H₂ concentrations ~10 nmol L⁻¹ sufficient to maintain hydrogenotrophic methanogenesis (Heimann et al., 2010), and low DIC:CH₄ ratios suggested methanogenic conditions over the whole profile (Fig. 4c, 5c). Methanogenesis was especially indicated by a slight peak of CH₄ concentrations associated with lower DIC:CH₄ ratios and elevated H₂ ratios around 0.3 m depth. However, even high activity of fermentation processes as indicated by peaking H₂ levels from 1 to 2 m depth did not translate to changes in fractionation factor (Fig. 4f, 5a), but suggest that CH₄ was produced only very locally if at all. Low substrate supply from highly decomposed organic matter and thus unfavourable thermodynamic conditions seemed to limit CH₄ production as suggested by very negative δ¹³C-CH₄ signatures (Fig. 4d) accompanied with peaking α_c-values in the upper profile (Fig. 4f) (Knorr et al., 2008b; Hornibrook et al., 1997). Instead, methanogenesis might have been outcompeted by other electron accepting processes, such as sulfate reduction (Fig. 5c). If produced at all, CH₄ was rapidly consumed around 0.3 m depth by methanotrophic bacteria inhabiting submerged *Sphagnum* mosses that provided recalcitrant organic matter on pool bottoms (Kleinebecker et al., 2007; Kip et al., 2012; Knoblauch et al., 2015), or associated with cyanobacteria (Arsenault et al., 2018; González Garraza et al., 2018). Within the open water of pools which was well-mixed by strong winds, CH₄ production and excess of H₂ were diminished due to electron acceptor availability for consumption (e.g. O₂, sulfate). Thus, the pools at our study site belonged to examples of low emission pools as described in Knoblauch et al. (2015).

4.4 *Sphagnum* and *Donatia* lawns – low to moderate emission scenario

CH₄ emissions from *Sphagnum* and *Donatia* lawns were most variable among microforms ranging from near-zero to local emission hotspots. *Sphagnum* lawns showed significantly higher CH₄ emissions of 1.52 ± 1.10 mmol m⁻² d⁻¹ than all other microforms (Fig. 3). Substantially higher fluxes from *Sphagnum* lawns at our study site have been previously described by Fritz et al. (2011) and emissions reported here were in the same order of magnitude as determined for *Sphagnum* lawns in South Patagonian bogs dominated by *Sphagnum* mosses (Broder et al., 2015; Lehmann et al., 2016). Highest emissions obtained in the present study from *Sphagnum* and *Donatia* lawns were at intermediate levels compared to northern bogs (Blodau, 2002; Laine et al., 2007; Limpens et al., 2008).

We present the first CH₄ emission data obtained from *Donatia* lawns. Contrary to our expectations, *Donatia* lawns emitted 0.66 ± 0.96 mmol CH₄ m⁻² d⁻¹ and had thus second highest CH₄ emissions of our study. One lawn even exceeded emissions of the strongest *Sphagnum* lawn source (Fig. 3c, Table 2) on all measurement occasions. As these comparatively high emissions

could be well reproduced, we conclude that an artefact can be excluded. Therefore, our first hypothesis, that cushion plants emit negligible amounts of CH₄, could retrospectively only be partly confirmed, as it depends on the specific cushion-forming plant species. We conclude that the presence of cushion plants as a proxy for negligible CH₄ emissions from cushion bogs needs to be carefully interpreted.

5 The spatial distribution of vascular plant root characteristics and its oxidative effects apparently controlled CH₄ dynamics in the study site as suggested by the pronounced spatial variability of emissions associated with our preliminary observations of microform features (Table 2). *D. fascicularis* established a deep and aerenchymous, but with < 0.05 g DW L of peat⁻¹ a substantially less dense rooting system compared to *A. pumila* with > 4 g DW L of peat⁻¹ in around 0.5 m depth (Fritz et al., 2011). The lower root density of *D. fascicularis* was probably below a specific threshold at which O₂ release did not suppress
10 emissions any more. Instead, the roots might have accelerated CH₄ production by presence of labile organic matter while facilitating emissions (e.g. Joabsson et al., 1999; Blodau, 2002; Colmer, 2003; Whalen, 2005; Agethen et al., 2018). As root O₂ release promotes nutrient mobilization (Colmer, 2003), this assumption of little if any O₂ release by *D. fascicularis* is in line with Schmidt et al. (2010) who found nutrient concentrations in *D. fascicularis* to be almost as low as in *S. magellanicum* while *A. pumila* contained significantly higher amounts of nutrients.

15 *Sphagnum* and *Donatia* lawns that showed substantial CH₄ emissions were beyond the sphere of influence by *A. pumila* roots. Although not measured here, pore water CH₄ concentrations below *Sphagnum* and *Donatia* lawns with remarkably higher emissions probably increased steeply below the water table (Fig. 6), like common depth profiles in northern (e.g. Beer and Blodau, 2007; Limpens et al., 2008; Corbett et al., 2013) or southern bogs (Broder et al., 2015). This explanation implies that the pore water below those lawns was not well-connected to pore water influenced by *A. pumila* roots. A spatial variability of
20 peat physical properties on the ecosystem scale was suggested by water table records (Fig. 2) demonstrating that one site responded faster than the other to, for instance, precipitation events.

We speculate that the biogeochemistry below *Sphagnum* and *Donatia* lawns with near-zero emissions was, likewise to pools, influenced by root activity of nearby *A. pumila* (as visualized in Fig. 6). This assumption was supported by previous research in our study site. Fritz et al. (2011) showed that below *Sphagnum* lawns pore water CH₄ concentrations can be substantially
25 lowered throughout rhizosphere-influenced peat layers. Further research should be undertaken to investigate why CH₄ emissions from *Sphagnum* and *Donatia* lawns appear to be highly variable in order to understand CH₄ dynamics in under-researched cushion bogs.

4.5 Implications for ecosystem CH₄ emissions from Patagonian cushion bogs

Highest summer emissions in the present study were considerably lower compared to maximum austral summer fluxes of ~150
30 mg CH₄ m⁻² d⁻¹ (9.4 mmol m⁻² d⁻¹) determined for a New Zealand bog dominated by the vascular evergreen “wire rush,” *Empodisma robustum* (Goodrich et al., 2015). The authors explained such high CH₄ emissions by comparatively wet conditions and a high density of aerenchymous vegetation providing a gas conduit for CH₄ transport in their study site. Compared to CH₄

emissions of northern bogs ranging from 3 to 53 mg m⁻² d⁻¹ at intermediate levels in wet bogs to up to 80 mg m⁻² d⁻¹ (Blodau, 2002; Laine et al., 2007; Limpens et al., 2008; 0.2 to 2.2 and up to 5 mmol m⁻² d⁻¹), highest summer emissions in the present study represent a pronounced CH₄ source. Nevertheless, overall ecosystem CH₄ emissions would probably be among the lowest described for pristine bog ecosystems worldwide when taking into account the high proportion of the surface area where *Astelia* lawns prevailed.

While *Donatia* lawns covered only small parts of our study site, we observed other parts of the complex bog system in the whole study area to be widely dominated by *D. fascicularis* instead of *A. pumila* which is in line with previous studies (Heusser, 1995; Grootjans et al., 2014) who described the cushion bogs in the Moat landscape to be covered by both, *A. pumila* and *D. fascicularis*. Thus, amounts of emitted CH₄ could be significant also on the ecosystem scale once *D. fascicularis* dominates. For instance, a predominance of *D. fascicularis* is typical for later vegetation succession stages in cushion bogs of Tierra del Fuego (Heusser, 1995) or Chile (Ruthsatz and Villagran, 1991; Kleinebecker et al., 2007).

5 Conclusion

We conclude from our study that the spatial distribution of cushion-forming *A. pumila* root density and activity and associated O₂ supply strongly controlled CH₄ production and consumption in a pristine Patagonian cushion bog. Thereby, CH₄ emission were reduced to near-zero levels and largely decoupled from environmental controls. The high root density even regulated CH₄ dynamics below adjacent microforms with less or non-rooted peat such as pools, *Sphagnum* lawns or lawns dominated by cushion-forming *D. fascicularis* by maintaining lateral concentration gradients in upper peat layers. Nevertheless, when root density dropped below a certain threshold, CH₄ production might have been accelerated by presence of labile root organic matter or exudates accompanied with O₂ consumption along with facilitated CH₄ transport by aerenchymatic roots of e.g. *D. fascicularis*. Under such circumstances, CH₄ emissions increased to intermediate levels compared to northern bogs. Therefore, the presence of cushion plants as a proxy for negligible CH₄ emissions from cushion bogs should be interpreted carefully. As cushion bogs can be found in many (mostly southern) parts of the world and only very limited knowledge about CH₄ dynamics from these systems exist, future research should take into account a possibly high spatial variability of CH₄ emission from bogs dominated by cushion plants. We demonstrated an extreme scenario for how a spatial distribution of root density in the peat can lead to a pronounced pattern of CH₄ emissions. Yet, the underlying ratio between root characteristics and O₂ supply determining this emission pattern should be applicable to other densely rooted peatlands in general. Further research should be undertaken to prove relationships that have been developed here on a conceptual level in order to extend our knowledge on CH₄ dynamics in under-researched cushion bogs.

Data availability. The data can be accessed by email request to the corresponding authors.

Author contributions. CB, TK and WM designed the study. WM, CB, TK and VAP conducted field work and sample analyses with the help of KHK. WM performed data analyses and prepared the manuscript with contributions from KHK, TK and VAP.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgement. This study was carried out within the research project CANDYbog which was funded by the Deutsche Forschungsgemeinschaft (German Research Foundation, DFG; Grant No. KL2265/3-1 and BL 563/19-1). We highly appreciated the Centro Austral de Investigaciones Científicas (CADIC-CONICET) providing facilities that enabled our fieldwork in Ushuaia. The Prefectura Naval Argentina is acknowledged for offering facilities supporting our fieldwork. Lucas Varela of the La Posta Hostel family is appreciated for offering working space and logistics. We wish to acknowledge the laborious field assistance of Isabella Nördemann, Carla Bockermann, Claudia Frank, Bettina Breuer, Lina Birkner, Juliane Kohlstruck, Laura Jansen, David Holl, Lars Kutzbach, María Florencia Castagnani and María Noel Szudruk Pascual. All analyses of this study were carried out in the laboratory of the institute of Landscape Ecology. The assistance of Ulrike Berning-
5
10 Mader, Sebastian R. Schmidt, Ronya Wallis, Sabrina Knaack, Sina Berger, Simona Bonaiuti and Leandra Praetzel is greatly acknowledged. We thank Peter Sulmann for technical assistance. We are grateful for the valuable comments of two anonymous referees. The support by Open Access Publication Fund of University of Muenster is acknowledged.

References

- Agethen, S., Sander, M., Waldemer, C., and Knorr, K.-H.: Plant rhizosphere oxidation reduces methane production and emission in rewetted peatlands, *Soil Biology and Biochemistry*, 125, 125-135, 2018.
- 5 Arsenault, J., Talbot, J., and Moore, T. R.: Environmental controls of C, N and P biogeochemistry in peatland pools, *Sci Total Environ*, 631-632, 714-722, 2018.
- Aselmann, I., and Crutzen, P. J.: Global distribution of natural freshwater wetlands and rice paddies, their net primary productivity, seasonality and possible methane emissions, *J Atmos Chem*, 8, 307-358, 1989.
- Baird, A. J., Milner, A. M., Blundell, A., Swindles, G. T., and Morris, P. J.: Microform-scale variations in peatland permeability and their ecohydrological implications, *J Ecol*, 104, 531-544, 2016.
- 10 Beer, J., and Blodau, C.: Transport and thermodynamics constrain belowground carbon turnover in a northern peatland, *Geochim Cosmochim Acta*, 71, 2989-3002, 2007.
- Beer, J., Lee, K., Whiticar, M., and Blodau, C.: Geochemical controls on anaerobic organic matter decomposition in a northern peatland, *Limnol Oceanogr*, 53, 1393-1407, 2008.
- 15 Benavides, J. C., Vitt, D. H., and Wieder, R. K.: The influence of climate change on recent peat accumulation patterns of *Distichia muscoides* cushion bogs in the high-elevation tropical Andes of Colombia, *Journal of Geophysical Research: Biogeosciences*, 118, 1627-1635, 2013.
- Berg, P., Risgaard-Petersen, N., and Rysgaard, S.: Interpretation of measured concentration profiles in sediment pore water, *Limnol Oceanogr*, 43, 1500-1510, 1998.
- 20 Berger, S., Praetzel, L. S. E., Goebel, M., Blodau, C., and Knorr, K. H.: Differential response of carbon cycling to long-term nutrient input and altered hydrological conditions in a continental Canadian peatland, *Biogeosciences*, 15, 885-903, 2018.
- Blodau, C.: Carbon cycling in peatlands - A review of processes and controls, *Environmental Reviews*, 10, 111-134, 2002.
- Blodau, C., and Moore, T. R.: Experimental response of peatland carbon dynamics to a water table fluctuation, *Aquat Sci*, 65, 47-62, 2003.
- 25 Bonaiuti, S., Blodau, C., and Knorr, K.-H.: Transport, anoxia and end-product accumulation control carbon dioxide and methane production and release in peat soils, *Biogeochemistry*, 133, 219-239, 2017.
- Borromei, A. M., Ponce, J. F., Coronato, A., Candel, M. S., Olivera, D., and Okuda, M.: Reconstrucción de la vegetación posglacial y su relación con el ascenso relativo del nivel del mar en el extremo este del canal Beagle, Tierra del Fuego, *Argentina Andean Geology*, 41, 362-379, 2014.
- 30 Boucher, F. C., Lavergne, S., Basile, M., Choler, P., and Aubert, S.: Evolution and biogeography of the cushion life form in angiosperms, *Perspectives in Plant Ecology, Evolution and Systematics*, 20, 22-31, 2016.

- Broder, T., Blodau, C., Biester, H., and Knorr, K. H.: Sea spray, trace elements, and decomposition patterns as possible constraints on the evolution of CH₄ and CO₂ concentrations and isotopic signatures in oceanic ombrotrophic bogs, *Biogeochemistry*, 122, 327-342, 2015.
- Burger, M., Berger, S., Spangenberg, I., and Blodau, C.: Summer fluxes of methane and carbon dioxide from a pond and
5 floating mat in a continental Canadian peatland, *Biogeosciences*, 13, 3777-3791, 2016.
- Chasar, L. S., Chanton, J. P., Glaser, P. H., and Siegel, D. I.: Methane concentration and stable isotope distribution as evidence of rhizospheric processes: Comparison of a fen and bog in the Glacial Lake Agassiz Peatland complex, *Annals of Botany*, 86, 655-663, 2000.
- Colmer, T. D.: Long-distance transport of gases in plants: a perspective on internal aeration and radial oxygen loss from roots,
10 *Plant, Cell & Environment*, 26, 17-36, 2003.
- Conrad, R.: Quantification of methanogenic pathways using stable carbon isotopic signatures: a review and a proposal, *Org Geochem*, 36, 739-752, 2005.
- Coombes, L., and Ramsay, P. M.: Vegetation of a cushion mire at 3600 m on Volcán Chiles, Ecuador, in: *The Ecology of Volcán Chiles. High altitude ecosystems on the Ecuador Colombia border*, edited by: Ramsay, P. M., Pebble & Shell
15 Publications, Plymouth, 47-54, 2001.
- Corbett, J. E., Tfaily, M. M., Burdige, D. J., Cooper, W. T., Glaser, P. H., and Chanton, J. P.: Partitioning pathways of CO₂ production in peatlands with stable carbon isotopes, *Biogeochemistry*, 114, 327-340, 2013.
- Dullo, B. W., Grootjans, A. P., Roelofs, J. G. M., Senbeta, A. F., Fritz, C., and Lamers, L. P. M.: Radial oxygen loss by the cushion plant *Eriocaulon schimperii* prevents methane emissions from an East-African mountain mire, *Plant Biology*, 19, 736-
20 741, 2017.
- Eckhardt, T. and Kutzbach, L.: MATLAB code to calculate gas fluxes from chamber based methods. Institute of Soil Science, Hamburg, Germany, <https://doi.pangaea.de/10.1594/PANGAEA.857799>, access: 17.07.2016.
- Estop-Aragónés, C., Knorr, K. H., and Blodau, C.: Belowground in situ redox dynamics and methanogenesis recovery in a degraded fen during dry-wet cycles and flooding, *Biogeosciences*, 10, 421-436, 2013.
- 25 Fechner-Levy, E. J., and Hemond, H. F.: Trapped methane volume and potential effects on methane ebullition in a northern peatland, *Limnol Oceanogr*, 41, 1375-1383, 1996.
- Fonkén, M. S. M.: An introduction to the bofedales of the Peruvian High Andes, *Mires and Peat*, 15, 2014.
- Fritz, C., Pancotto, V. A., Elzenga, J. T. M., Visser, E. J. W., Grootjans, A. P., Pol, A., Iturraspe, R., Roelofs, J. G. M., and Smolders, A. J. P.: Zero methane emission bogs: extreme rhizosphere oxygenation by cushion plants in Patagonia, *New Phytol*,
30 190, 398-408, 2011.
- Fritz, C.: Limits of Sphagnum bog growth in the New World: Biochemistry and ecohydrology of peatlands in South America and New Zealand, Ph.D thesis, Radboud University of Nijmegen / University of Groningen, Netherlands, 193 pp., 2012.
- Gibson, N., and Kirkpatrick, J. B.: A Comparison of the Cushion Plant-Communities of New-Zealand and Tasmania, *New Zeal J Bot*, 23, 549-566, 1985.

- González Garraza, G., Pancotto, V. A., Mansilla, R., and Escobar, J.: Importancia de la humedad en la caracterización de las comunidades microalgales en turberas compactas de Tierra del Fuego, in: Resúmenes de la Reunión Argentina de Ecología, XXVIII Reunión Argentina de Ecología, Mar del Plata, Argentina, 29 de octubre al 2 de noviembre de 2018, 353, 2018.
- Goodrich, J. P., Campbell, D. I., Roulet, N. T., Clearwater, M. J., and Schipper, L. A.: Overriding control of methane flux temporal variability by water table dynamics in a Southern Hemisphere, raised bog, *Journal of Geophysical Research: Biogeosciences*, 120, 819-831, 2015.
- Grootjans, J., Iturraspe, R., Fritz, C., Moen, A., and Joosten, H.: Mires and mire types of Peninsula Mitre, Tierra del Fuego, Argentina, *Mires and Peat*, 14, 1-20, 2014.
- Hamilton, J. D., Kelly, C. A., Rudd, J. W. M., Hesslein, R. H., and Roulet, N. T.: Flux to the atmosphere of CH₄ and CO₂ from wetland ponds on the Hudson Bay lowlands (HBLs), *Journal of Geophysical Research: Atmospheres*, 99, 1495-1510, 1994.
- Heimann, A., Jakobsen, R., and Blodau, C.: Energetic Constraints on H₂-Dependent Terminal Electron Accepting Processes in Anoxic Environments: A Review of Observations and Model Approaches, *Environ Sci Technol*, 44, 24-33, 2010.
- Heusser, C. J.: Paleocology of a *Donatia-Astelia* Cushion Bog, Magellanic Moorland-Sub-Antarctic Evergreen Forest Transition, Southern Tierra-Del-Fuego, Argentina, *Rev Palaeobot Palyno*, 89, 429-440, 1995.
- Hope, G. S.: Peat in the mountains of New Guinea, *Mires and Peat*, 15, 2014.
- Hornibrook, E. R. C., Longstaffe, F. J., and Fyfe, W. S.: Spatial distribution of microbial methane production pathways in temperate zone wetland soils: Stable carbon and hydrogen isotope evidence, *Geochim Cosmochim Acta*, 61, 745-753, 1997.
- Hornibrook, E. R. C., Longstaffe, F. J., and Fyfe, W. S.: Evolution of stable carbon isotope compositions for methane and carbon dioxide in freshwater wetlands and other anaerobic environments, *Geochim Cosmochim Acta*, 64, 1013-1027, 2000.
- IPCC: Anthropogenic and Natural Radiative Forcing, in: *Climate Change 2013 – The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Change, I. P. o. C., Cambridge University Press, Cambridge, 659-740, 2014.
- Iturraspe, R.: Spatial analysis and description of eastern peatlands of Tierra del Fuego, Argentina, in: *Mires from pole to pole*, edited by: Lindholm, T., and Heikkilä, R., Finnish Environment Institute (SYKE), Helsinki, Finland, 2012.
- Jackowicz-Korczyński, M., Christensen, T. R., Bäckstrand, K., Crill, P., Friborg, T., Mastepanov, M., and Ström, L.: Annual cycle of methane emission from a subarctic peatland, *Journal of Geophysical Research: Biogeosciences*, 115, 2010.
- Joabsson, A., Christensen, T. R., and Wallén, B.: Vascular plant controls on methane emissions from northern peatforming wetlands, *Trends in Ecology & Evolution*, 14, 385-388, 1999.
- Kip, N., Fritz, C., Langelaan, E. S., Pan, Y., Bodrossy, L., Pancotto, V., Jetten, M. S. M., Smolders, A. J. P., and den Camp, H. J. M. O.: Methanotrophic activity and diversity in different *Sphagnum magellanicum* dominated habitats in the southernmost peat bogs of Patagonia, *Biogeosciences*, 9, 47-55, 2012.
- Kleinebecker, T., Hölzel, N., and Vogel, A.: Gradients of continentality and moisture in South Patagonian ombrotrophic peatland vegetation, *Folia Geobot*, 42, 363-382, 2007.

- Kleinebecker, T., Hölzel, N., and Vogel, A.: South Patagonian ombrotrophic bog vegetation reflects biogeochemical gradients at the landscape level, *J Veg Sci*, 19, 151-160, 2008.
- Knoblauch, C., Spott, O., Evgrafova, S., Kutzbach, L., and Pfeiffer, E.-M.: Regulation of methane production, oxidation, and emission by vascular plants and bryophytes in ponds of the northeast Siberian polygonal tundra, *Journal of Geophysical Research: Biogeosciences*, 120, 2525-2541, 2015.
- Knorr, K. H., Oosterwoud, M. R., and Blodau, C.: Experimental drought alters rates of soil respiration and methanogenesis but not carbon exchange in soil of a temperate fen, *Soil Biol Biochem*, 40, 1781-1791, 2008a.
- Knorr, K. H., Glaser, B., and Blodau, C.: Fluxes and ^{13}C isotopic composition of dissolved carbon and pathways of methanogenesis in a fen soil exposed to experimental drought, *Biogeosciences*, 5, 1457-1473, 2008b.
- 10 Knorr, K.-H., Lischeid, G., and Blodau, C.: Dynamics of redox processes in a minerotrophic fen exposed to a water table manipulation, *Geoderma*, 153, 379-392, 2009.
- Knorr, K.-H., Horn, M. A., and Borken, W.: Significant nonsymbiotic nitrogen fixation in Patagonian ombrotrophic bogs, *Global Change Biol*, 21, 2357-2365, 2015.
- Kotsyurbenko, O. R., Glagolev, M. V., Nozhevnikova, A. N., and Conrad, R.: Competition between homoacetogenic bacteria and methanogenic archaea for hydrogen at low temperature, *Fems Microbiol Ecol*, 38, 153-159, 2001.
- 15 Kutzbach, L., Schneider, J., Sachs, T., Giebels, M., Nykanen, H., Shurpali, N. J., Martikainen, P. J., Alm, J., and Wilmking, M.: CO_2 flux determination by closed-chamber methods can be seriously biased by inappropriate application of linear regression, *Biogeosciences*, 4, 1005-1025, 2007.
- Laine, A., Wilson, D., Kiely, G., and Byrne, K. A.: Methane flux dynamics in an Irish lowland blanket bog, *Plant Soil*, 299, 181-193, 2007.
- 20 Lehmann, J., Münchberger, W., Knoth, C., Blodau, C., Nieberding, F., Prinz, T., Pancotto, V., and Kleinebecker, T.: High-Resolution Classification of South Patagonian Peat Bog Microforms Reveals Potential Gaps in Up-Scaled CH_4 Fluxes by use of Unmanned Aerial System (UAS) and CIR Imagery, *Remote Sens-Basel*, 8, 173, 2016.
- Limpens, J., Berendse, F., Blodau, C., Canadell, J. G., Freeman, C., Holden, J., Roulet, N., Rydin, H., and Schaepman-Strub, G.: Peatlands and the carbon cycle: from local processes to global implications - a synthesis, *Biogeosciences*, 5, 1475-1491, 2008.
- 25 Mainiero, R., and Kazda, M.: Effects of *Carex rostrata* on soil oxygen in relation to soil moisture, *Plant Soil*, 270, 311-320, 2005.
- Mikaloff Fletcher, S. E., Tans, P. P., Bruhwiler, L. M., Miller, J. B., and Heimann, M.: CH_4 sources estimated from atmospheric observations of CH_4 and its $^{13}\text{C}/^{12}\text{C}$ isotopic ratios: 1. Inverse modeling of source processes, *Global Biogeochem Cy*, 18, 2004.
- 30 Paredes, N. I., Consolo, V. F., Pancotto, V. A., Fritz, C., Barrera, M. D., Arambarri, A. M., and Salerno, G. L.: Microfungal composition in an *Astelia-Donatia* cushion peatland in Tierra del Fuego, Argentina. , *Darwinia*, 2, 2014.

- Pelletier, L., Strachan, I. B., Garneau, M., and Roulet, N. T.: Carbon release from boreal peatland open water pools: Implication for the contemporary C exchange, *Journal of Geophysical Research: Biogeosciences*, 119, 207-222, 2014.
- Popp, T. J., Chanton, J. P., Whiting, G. J., and Grant, N.: Methane stable isotope distribution at a *Carex* dominated fen in north central Alberta, *Global Biogeochem Cy*, 13, 1063-1077, 1999.
- 5 Rinne, J., Riutta, T., Pihlatie, M., Aurela, M., Haapanala, S., Tuovinen, J.-P., Tuittila, E.-S., and Vesala, T.: Annual cycle of methane emission from a boreal fen measured by the eddy covariance technique, *Tellus B*, 59, 449-457, 2007.
- Ruthsatz, B., and Villagran, C.: Vegetation Pattern and Soil Nutrients of a Magellanic Moorland on the Cordillera-De-Piuchue, Chiloe Island, Chile, *Rev Chil Hist Nat*, 64, 461-478, 1991.
- Sander, R.: Compilation of Henry's Law constants for inorganic and organic species of potential importance in environmental chemistry. Max Planck Institute of Chemistry, Mainz, Germany, available at: <http://www.henrys-law.org/henry-3.0.pdf> (last access: 1 June 2017), 1999
- 10 Schmidt, S. R., Kleinebecker, T., Vogel, A., and Holzner, N.: Interspecific and geographical differences of plant tissue nutrient concentrations along an environmental gradient in Southern Patagonia, Chile, *Aquat Bot*, 92, 149-156, 2010.
- Steinmann, P., Eilrich, B., Leuenberger, M., and Burns, S. J.: Stable carbon isotope composition and concentrations of CO₂ and CH₄ in the deep catotelm of a peat bog, *Geochim Cosmochim Acta*, 72, 6015-6026, 2008.
- 15 Stumm, W., and Morgan, J. J.: Aquatic chemistry. Chemical Equilibria and Rates in Natural Waters, 3rd ed., Wiley, Hoboken, N.J., USA, 1996.
- von Mering, S.: *Tetroncium* and its only species, *T. magellanicum* (Juncaginaceae): distribution, ecology and lectotypification, *Willdenowia*, 43, 13-24, 2013.
- 20 Whalen, S. C.: Biogeochemistry of Methane Exchange between Natural Wetlands and the Atmosphere, *Environmental Engineering Science*, 22, 73-94, 2005.
- Whiticar, M. J., Faber, E., and Schoell, M.: Biogenic methane formation in marine and freshwater environments: CO₂ reduction vs. acetate fermentation—Isotope evidence, *Geochim Cosmochim Acta*, 50, 693-709, 1986.
- Yu, Z. C.: Northern peatland carbon stocks and dynamics: a review, *Biogeosciences*, 9, 4071-4085, 2012.

Figures




Figure 1. Location of the study area in southernmost Patagonia. The investigated cushion bog in Moat is located 130 km eastern of Ushuaia (Argentina). The map was created at <http://www.simplemappr.net> accessed on 11.12.2017 and in Matlab 5 (MATLAB Mapping Toolbox Release R2017a).




Figure 2. Time series of environmental variables during the study period. Photosynthetic active radiation (PAR, a, b) and air temperature (c, d) are presented as half-hourly and daily means during two austral summer periods. Soil temperature (e) and water table position below surface (f) were continuously recorded and are shown as half-hourly means. Note the difference in response time between platform a and platform b where water table fluctuations were recorded.




Figure 3 a-c. Mean CH₄ emissions and their standard deviation determined from dominant microforms (a) and individual collars during the two measurement campaigns (b, c) in a Patagonian cushion bog. Different letter superscripts denote significantly different amounts of emitted CH₄ between microforms despite variability within microforms was substantial (a) or from collars within one microform (b, c) that were identified by Kruskal-Wallis analysis of variance and multiple comparison U-tests with Bonferroni adjustment.





Figure 4 a-f. Pore water composition in depth profiles obtained from MLPS (N = 3) installed in *Astelia* lawns and pools during two sampling campaigns in austral summer 2015 and 2016. Mean values and their standard deviation are presented. CH₄ (a) and DIC (b) concentrations as well as related DIC:CH₄ (c) ratios were determined during both sampling events while carbon isotope values of CH₄ (d) and DIC (e) and corresponding fractionation factors (f) were investigated only during the second

campaign. DIC:CH₄ ratios are shown on a logarithmic scale without negative standard deviation for *Astelia* lawns and pools. Displayed depths represent the centre of a MLP segment and had to be standardized because of different sampling resolution in both years. The water table position and approximate maximum rooting depth (rhizosphere) refer to *Astelia* lawns only. Note the different scale of the y-axis in the upper and lower three panels.



5

Figure 5 a-c. Profiles of H₂ (a), O₂ (b) and sulfate (c) pore water concentrations obtained from MLPs (N = 3) installed in *Astelia* lawns and pools during two sampling campaigns in austral summer 2015 and 2016. Mean values and their standard deviation are presented. Displayed depths represent the centre of a MLP segment and had to be standardized because of different sampling resolution in both years. The water table position and approximate maximum rooting depth (rhizosphere)

10 refer to *Astelia* lawns only.




Figure 6. Schematic CH_4 concentration profiles below the four dominant microforms in a Patagonian cushion bog. The shape of profiles was derived from data obtained in the present study together with those by Limpens et al. (2008), Fritz et al. (2011) and Broder et al. (2015). The size of arrows represents the relative magnitude of either a CH_4 or CO_2 flux following the diffusion gradient. The water table position is displayed by a dashed line near the surface. CH_4 oxidation by root O_2 release resulted in near-zero CH_4 pore water concentrations in densely rooted peat below *Astelia* lawns. With increasing distance from densely rooted peat, CH_4 oxidation decreased while CH_4 concentration increased until reaching a depth profile typical for those of northern peatlands with less if any roots in water-saturated peat layers. Exemplary sequence of microforms, derived from observed amounts of emitted CH_4 . Other sequences of microforms are reasonable as microforms occur in various combinations in the field.

Tables

Table 1. Characteristic plant species of dominant microforms in a Patagonian cushion bog. Species composition was determined during the second measurement campaign within collars where closed chamber measurements were conducted. Mean cover (%) and its standard deviation of characteristic were given and plant species with a mean cover < 5 % were listed

5 with +.

species	microform		
	<i>Astelia</i> lawn	<i>Sphagnum</i> lawn	<i>Donatia</i> lawn
<i>Astelia pumila</i>	84 ± 16		
<i>Donatia fascicularis</i>	8 ± 10	+	90 ± 9
<i>Sphagnum magellanicum</i>		94 ± 4	
<i>Tetroncium magellanicum</i>	+	7 ± 2	+
<i>Caltha dioneifolia</i>	+	+	6 ± 5
<i>Empetrum rubrum</i>	+		
<i>Gaultheria antarctica</i>	+	+	+
<i>Myrteola nummularia</i>	+		

Table 2. CH₄ fluxes determined from individual collars installed in dominant microforms of a Patagonian cushion bog. Number of flux measurements, their mean and standard deviation are presented. Significantly different fluxes from collars within one microform were identified by Kruskal-Wallis analysis of variance and multiple comparison U-tests with Bonferroni adjustment. Results of computing Spearman's rank correlation coefficient rho for the relationships between CH₄ fluxes and possible environmental controls, namely water table depth determined separately for each collar and soil temperature in 0.05 m depth, are given. Furthermore, characteristics of collars and results of computing Spearman's rank correlation for the relationships between mean CH₄ fluxes during the second campaign and each feature are shown. Microform size refers to the extent of a microform in which a specific collar was installed. P-values < 0.05 indicate a correlation significantly different from zero.

10

year	micro-form	Kruskal-Wallis analysis			water table depth, m		soil temperature, °C		presence <i>T. magellanicum</i>			presence <i>D. fascicularis</i>			microform size			
		N	mean flux, mmol CH ₄ m ⁻² d ⁻¹	p	rho	p	rho	p	number of shoots	rho	p	cover, %	rho	p	size, m ²	rho	p	
2015	<i>Astelia</i> lawn	1	60	0.32 ± 0.05	< 0.05	-0.22	n.s.	n.d.	n.d.									
		2	64	-0.02 ± 0.09	n.s.	-0.21	n.s.	0.17	n.s.									
		3	55	-0.01 ± 0.07	n.s.	0.03	n.s.	-0.08	n.s.									
	<i>Sphagnum</i> lawn	1	62	0.21 ± 0.05	< 0.05	-0.21	n.s.	n.d.	n.d.									
		2	64	1.96 ± 0.33	n.s.	0.41	< 0.05	0.52	< 0.05									
		3	54	2.99 ± 0.35	n.s.	0.2	n.s.	0.6	< 0.05									
	pool	1	13	0.05 ± 0.03	n.s.			n.d.	n.d.									
		2	12	0.14 ± 0.08	n.s.			0.4	n.s.									
		3	10	0.36 ± 0.2	n.s.			0.64	< 0.05									
		4	11	0.28 ± 0.2	n.s.			-0.01	n.s.									
	2016	<i>Astelia</i> lawn	1	7	0.29 ± 0.04	n.s.	0.27	n.s.	-0.13	n.s.	15			25				1.12
			2	11	-0.02 ± 0.1	n.s.	-0.22	n.s.	-0.16	n.s.	0			4				4.45
3			8	0.07 ± 0.05	n.s.	-0.68	n.s.	-0.31	n.s.	0	0.67 ^a	n.s.	0	0.01	n.s.	26.66	-0.10	n.s.
4			11	0.06 ± 0.15	n.s.	-0.09	n.s.	0.13	n.s.	0			8				3.38	
5			11	0.06 ± 0.04	n.s.	-0.44	n.s.	0.14	n.s.	6			2				4.55	
<i>Donatia</i> lawn		1	9	2.1 ± 0.14	< 0.05	-0.82	< 0.05	0.74	< 0.05	30			80				0.26	
		2	11	0.01 ± 0.09	n.s.	0.29	n.s.	-0.64	< 0.05	9	1.00 ^a	n.s.	97	-1.00 ^a	n.s.	0.15	0.50	n.s.
		3	10	0.08 ± 0.04	n.s.	-0.46	n.s.	-0.2	n.s.	18			92				0.96	
<i>Sphagnum</i> lawn		1	9	0.09 ± 0.05	< 0.05	0.68	n.s.	-0.78	< 0.05	66			0			0.43		
		2	11	1.01 ± 0.06	n.s.	0.15	n.s.	-0.42	n.s.	165	0.40	n.s.	0	0.71 ^a	n.s.	2.38	0.00	n.s.

	3	11	1.39 ± 0.1	n.s.	-0.05	n.s.	-0.05	n.s.	175	0	2.7
	4	9	1.9 ± 0.08	< 0.05	-0.44	n.s.	0.25	n.s.	108	4	0.38
	5	10	0.68 ± 0.03	n.s.	0.17	n.s.	-0.22	n.s.	120	0	0.65
pool	1	2	0.35 ± 0.11	n.s.			n.d.	n.d.			
	3	1	0.6 ± 0	n.s.			n.d.	n.d.			
	4	1	1.35 ± 0	n.s.			n.d.	n.d.			

n.s. = not significant, n.d. = not determined, ^a = despite a (relative) high absolute value of Spearman's rho, p-values did not indicate a significant correlation, probably because of outliers.

Table 3. Summary of peat characteristics obtained from two peat cores taken in *Astelia* lawns in a Patagonian cushion bog. Data were averaged over the respective depths sampled in 0.1 m resolution.

peat characteristics	depth, m	bulk density, g cm ⁻³	N, %	C, %	CN	δ ¹⁵ N	δ ¹³ C
amorphous, highly decomposed cushion peat with many living <i>Astelia</i> roots, only close to the surface (0-0.4 m) with recognizable plant material	0-1	0.03	1.42	49.31	37.62	-1.24	-25.63
	1-2	0.06	1.83	52.45	28.69	1.24	-26.18
<i>Sphagnum</i> peat, with depth increasingly amorphous and decomposed	2-3	0.06	1.52	51.37	34.38	1.12	-26.33
	3-4	0.06	1.22	50.49	41.98	0.22	-25.78
	4-5	0.11	1.17	47.86	41.45	-1.59	-26.43
	5-6	0.10	1.18	47.56	40.67	-1.62	-26.77
	6-7	0.15	1.28	50.27	44.34	-1.97	-27.08