



Rapid mineralization of biogenic volatile organic compounds in temperate and Arctic soils

Christian Nyrop Albers^{1,2}, Magnus Kramshøj^{1,2,3}, Riikka Rinnan^{2,3}

5 ¹Geological Survey of Denmark and Greenland (GEUS), Department of Geochemistry, Øster Voldgade 10, DK-1350 Copenhagen K, Denmark

²Center for Permafrost (CENPERM), Department of Geosciences and Natural Resource Management, University of Copenhagen, Øster Voldgade 10, DK-1350 Copenhagen K, Denmark

³Department of Biology, University of Copenhagen, Universitetsparken 15, DK-2100 Copenhagen E, Denmark

10 *Correspondence to:* Christian N Albers (cal@geus.dk)

Abstract. Biogenic volatile organic compounds (BVOCs) are produced by all life forms. Their release into the atmosphere is important with regards to a number of physical and chemical processes and great effort has been put into determining sources and sinks of these compounds in recent years. Soil microbes as a possible sink for BVOCs in the atmosphere has been suggested, however, experimental evidence for this sink is scarce despite its potentially high importance to both carbon cycling and atmospheric concentrations of these gases. We therefore conducted a study with a number of commonly occurring BVOCs labelled with ¹⁴C and modified existing methods to study mineralization of these compounds to ¹⁴CO₂ in four different top soils. Five of the six BVOCs were rapidly mineralized by microbes in all soils. However, great differences were observed with regards to speed of mineralization, extent of mineralization and variation between soil types. Methanol, benzaldehyde, acetophenone and the oxygenated monoterpene geraniol were mineralized within hours in all soils. The hydrocarbon monoterpene p-cymene was mineralized rapidly in soil from a coniferous forest but slower in soil from and adjacent beech stand while chloroform was mineralized slowly in all soils. From our study it is clear that soil microbes are able to degrade completely BVOCs released by aboveground vegetation as well as BVOCs released by soil microbes and plant roots. In addition to the possible atmospheric implications of this degradation the very fast mineralization rates are likely important in shaping the net BVOC emissions from soil and it is possible that BVOC formation and degradation may be an important but little recognized part of internal carbon cycling in soil.

15
20
25



1 Introduction

30 Non-methane biogenic volatile organic compounds (BVOCs) are produced by all life forms, with plants being the most important contributors to the atmospheric concentrations of BVOCs and also the most studied group of BVOC emitters (Laothawornkitkul et al., 2009; Peñuelas et al., 2014). Production of BVOCs in soil (McNeal and Herbert, 2009; Ramirez et al., 2010) and by isolated soil microorganisms (Insam and Seewald, 2010; Garbeva et al., 2014) has been shown as well, though.

35 BVOCs comprise a very high diversity concerning molecular size and chemical structures, which leads to a high compound-to-compound variation in life times and reactions in the environment. Chemical oxidation reactions are regarded as the dominant BVOC sink in air, with impacts on the concentrations of methane, ozone, secondary organic aerosol and even clouds (Peñuelas and Staudt, 2010; Glasius and Goldstein, 2016). In addition to chemical reactions in the atmosphere, an uptake or deposition of BVOCs into or onto soil has been observed (Ramirez et al., 2010; Spielmann
40 et al., 2017) and so has a bidirectional atmosphere/soil exchange of certain BVOCs (Asensio et al., 2007; Asensio et al., 2008; Gray et al., 2014). The mechanism behind the soil uptake has not been investigated, but it may owe to processes like adsorption to organic matter, dissolution in soil water and microbial degradation. Adsorption and dissolution may be predicted if the chemical characteristics of the BVOCs in question are known, and fate models for these parameters could then be set up. The microbial degradability of BVOCs - and especially the rate of degradation - are on the other
45 hand very difficult to predict.

It is known from lab experiments, that many BVOCs can be degraded by soil bacteria functioning as substrates for growth (e.g. Cripps, 1975; Misra et al., 1996; Kleinheinz, 1999; El Khawand et al., 2016). However, the studies on
50 BVOC degradation in soil or by isolated soil microorganisms, have typically used BVOC concentrations of 3-6 orders of magnitude higher than those present in the environment. Degradation experiments with such high concentrations are very well suited for selectively enriching BVOC degraders and showing the potential for use of a degrader organism in industrial processes. However, they do not serve to assess degradation at realistic environmental concentrations that would be too low to sustain bacterial growth singly due to degradation of a specific BVOC. Thus, we do not know if microbial BVOC degradation in soil is of environmental importance. An exception from this is isoprene degradation, of
55 which there is substantial evidence at environmental conditions (Cleveland and Yavitt, 1997; Gray et al., 2015).

Degradation experiments with BVOCs in soil are difficult to interpret as the same compounds may be produced and released by the soil while also being degraded. By using isotopically labelled compounds in degradation experiments it is possible to target degradation alone. Isotopic labelling also enables working with compounds at lower concentrations.
60 This is especially true for using radioactive ^{14}C -labelling, which furthermore enables one to determine complete mineralization to $^{14}\text{CO}_2$. Compared to compound removal over time, complete mineralization is the ultimate proof of degradation and is often used in pesticide fate studies. However, apart from three studies looking at mineralization of ^{14}C -labelled geraniol (Owen et al., 2007), methanol (Stacheter et al., 2013) and chloroform (Albers et al., 2011), we are unaware of such studies with BVOCs. Furthermore, so far no BVOC mineralization studies were done at concentrations
65 observed in natural environments.



The aim of this study was to assess microbial mineralization of different BVOCs in soils from contrasting environments. The microbial sink of BVOCs in soil would be of potentially high importance to both carbon cycling and atmospheric concentrations of these gases. We therefore purchased a number of commonly occurring BVOCs labelled with ^{14}C and modified existing methods to study mineralization of these compounds to $^{14}\text{CO}_2$ in four different soils.

2 Materials and Methods

2.1 Soil sampling and characterization

Soil was sampled at four sites representing common ecosystem types in the temperate and Arctic temperature zones.

75 From the temperate zone, we sampled a coniferous forest site ($12^\circ 03' 40''$ E, $56^\circ 02' 22''$ N) dominated by Norway spruce (*Picea abies*) and Scots pine (*Pinus sylvestris*) and a European beech (*Fagus sylvatica*) forest site ($12^\circ 04' 22''$ E, $56^\circ 02' 22''$ N). The two sites were located 750 m apart. At both sites the Aeolian sandy soil is 200 to 400 years old and has been forested for at least 150 years. Both sites lack underwood, forest floor vegetation and moss cover, and a 5-10 cm thick organic layer has accumulated on top of the sand. Loose litter was removed before sampling the organic layer.

80 From the Arctic, we sampled a tundra heath site ($53^\circ 27' 48''$ W, $69^\circ 15' 49''$) dominated by 5-15 cm tall dwarf shrubs *Empetrum nigrum*, *Betula nana* and *Cassiope tetragona*. A 4-8 cm thick organic layer has accumulated on top of a sandy parent soil. We sampled the organic layer in between individual plants. The second Arctic site was an area with bare ground without vegetation and with coarse soil particles ("Arctic bare soil", $53^\circ 27' 58''$ W, $69^\circ 15' 57''$), located 300 m from the heath site. Here, the top 5 cm was sampled.

85 From each site, 10-12 replicate samples were cored with a brass core (diameter 38 mm) from within a 25 m² area and pooled in a plastic bag. After arrival to the laboratory, the pooled samples were gently mixed by hand and larger roots were removed to get the final soil sample. The Arctic bare soil contained no roots and instead of mixing by hand, this soil was homogenized by sieving (5 mm). The mixed samples were stored at 3°C for a period of up to six weeks before mineralization experiments were initiated.

90

Water content was determined gravimetrically after drying at 105°C for 24 hours. Soil organic matter was determined as loss on ignition (LOI; 550°C, 2h). pH was determined with a pH electrode in slurries of soil:water (1:2.5) after 30 min shaking.

95 For each soil, triplicate DNA extractions were made from 0.25 g subsamples using the PowerLyzer PowerSoil DNA Isolation Kit (MoBio, Carlsbad, California). Total bacterial biomass was quantified as 16S gene copies by qPCR targeting the 16S rRNA sequence using forward primer 341F (5'-CCTACGGGAGGCAGCAG-3') and reverse primer 518R (5'-ATTACCGCGGCTGCTGG-3') and 1 µL DNA template, as previously described (Feld et al., 2016). Total fungal biomass was determined as ITS2 gene copies by targeting the fungal ITS2 nuclear ribosomal DNA region using

100



forward primer gITS7 (GTGARTCATCGARTCTTTG) and reverse primer ITS4 (TCCTCCGCTTATTGATATGC) as previously described (Christiansen et al., 2017). All qPCR was run in technical triplicates.

2.2 Atmospheric BVOC-concentrations

105 A snap-shot of the atmospheric concentration of a range of BVOCs was determined on the day of soil sampling in each of the two forest sites and in the Arctic sampling area. Triplicate 6 L air samples (12 L at the Arctic sites) were drawn through a sorbent cartridge 10 cm above soil surface (Coniferous, Beech and Arctic Bare sites) or 5 cm above the canopy (Arctic Heath site). Two types of sorbent cartridges were used in order to capture a range of BVOCs (Tenax TA/Carbograph 1TD as sorbent) and (Carbotrap B/Carboxen 1000/Carboxen 1003) to capture halogenated VOCs,
110 including the model compound chloroform. The sorbent cartridges were sampled and analyzed by GC-MS as previously described (Kramshøj et al., 2015; Johnsen et al., 2016)

2.3 Incubations for BVOC mineralization

Six ^{14}C -labeled BVOCs were used as model compounds representing different molecular weights and chemical classes
115 (Fig. 1, Table 2). ^{14}C -methanol (58 mCi millimole $^{-1}$), [ring- ^{14}C]-benzaldehyde (>99% radiochemical purity; 60 mCi millimole $^{-1}$) (trans)-[1- ^{14}C]-Geraniol (99% radiochemical purity; 55 mCi millimole $^{-1}$), [ring- ^{14}C]-acetophenone (99% radiochemical purity; 55 mCi millimole $^{-1}$) and ^{14}C -chloroform (>99% radiochemical purity; 2.25 mCi millimole $^{-1}$) were purchased from American Radiolabeled Chemicals Inc. (St. Louis, MO). [1-methyl- ^{14}C]-p-cymene (96% radiochemical purity; 57 mCi millimole $^{-1}$) was purchased from Moravek (Brea, Ca). Stock solutions (3×10^7 DPM mL $^{-1}$) were made in
120 sterile water (methanol and benzaldehyde), ethanol (acetophenone, geraniol and p-cymene) or acetonitrile (chloroform) and stored at -18°C until use.




Figure 1. Chemical structures of the used model compounds. Radiolabeled C is marked with an asterisk.


125

Incubations were carried out in 120 mL serum flasks. Into each flask, 5 (coniferous), 6 (beech and Arctic heath) or 10 (Arctic bare) g fresh weight (f.w.) soil with natural moisture was weighed and equilibrated overnight at 10°C. A small glass vial containing 2.5 mL 1M NaOH and 0.01M NaHCO₃ was placed in the flask to trap ¹⁴CO₂ liberated from ¹⁴C-BVOC mineralization. The NaHCO₃ was used in order to precipitate all trapped ¹⁴CO₂ with Ba²⁺ added during the following analysis procedure. 0.5 mL radiolabeled BVOCs dissolved in sterile water was then distributed across the soil with a pipette. The transfer of the BVOC had to be carried out fast in order not to loose it from the aqueous solution. For each BVOC, portions of the aqueous solution were transferred to scintillation vials containing HiSafe 3 liquid scintillation cocktail (Perkin Elmer, Waltham, MA) just before transferring to the first incubation flask and just after transferring to the last incubation flask to assure that all flasks had received similar ¹⁴C-BVOC-concentrations. The scintillation vials were then counted on a liquid scintillation counter (Tri-Carb 2810 TR, PerkinElmer, Waltham, MA) for 30 minutes or until 1% uncertainty (2S, 95% CL) was achieved. The BVOC concentrations used for incubation corresponded to 43-73 ppbv (64-504 ng L⁻¹), assuming all BVOC was present in the headspace of the flasks. Most of the BVOCs were, however likely dissolved in water or adsorbed to the soil, so recalculating to a soil basis (0.8-11 μg kg⁻¹ f.w. soil) may be more appropriate.

140

Immediately after the transfer of the BVOC solution, flasks were closed with crimp-caps containing an alumina-coated septum (Mikrolab Aarhus, Denmark) and incubated at 10°C in the dark. At several time points, the alkaline CO₂-trap was exchanged through a needle syringe permanently installed in the septum (to avoid losing BVOCs when exchanging the CO₂-trap). 1 mL was transferred to each of two 2 mL Eppendorf tubes with either 0.7 mL water or 0.7 mL BaCl₂ (1.5M, to precipitate trapped ¹⁴CO₂ as Ba¹⁴CO₃) to differentiate between trapped ¹⁴CO₂ and dissolved ¹⁴-BVOC (Fig. 2). After 5 h reaction time, the tubes were centrifuged (12000 g, 2 min) and 1 mL from each tube was counted by liquid scintillation. 1 mL 1M NaOH had been added to the scintillation liquid in the case of the tube added only water. This was done to increase pH in the liquid and thereby avoid losses of ¹⁴CO₂.

145



150 **Figure 2.** Sketch of method for capturing $^{14}\text{CO}_2$, separating it from dissolved ^{14}C -BVOC and analyzing by liquid scintillation counting (LSC).

155 After the last sampling point, 30 mL methanol were added to the soil in each flask through the permanently installed needle to extract any residual ^{14}C -BVOC. After 24 h shaking the supernatant was transferred to a 50 mL centrifuge tube and centrifuged (4000 g, 5 min). The ^{14}C -activity was then determined in 3 mL supernatant by liquid scintillation counting.

160 Incubations were made in three replicates. In addition to each BVOC/soil combination, a negative control was included in which the soil had been sterilized by autoclaving twice. Oxygen consumption during incubation was determined for each soil type by incubating an additional flask in which oxygen spot sensors (PreSens, Regensburg, Germany) readable through the glass of the bottles, had been installed.

165 The incubation method is a modification of previous methods for measuring mineralization of organic compounds. Suitability and limitations of the method are discussed in the supplementary information.

3 Results and Discussion

3.1 Soil characterization

170 The four top-soils used in the study showed clear differences with regards to major soil parameters like soil organic matter content, pH and microbial biomass (Table 1). The two forest top soils differed in soil organic matter content, but were both acidic (pH just below 4). The Arctic Heath top soil had an organic matter content in between the two forest soils but a higher pH of 5.3. As expected, the Arctic bare soil differed most, as it comprised the parent mineral soil while the others were dominated by organic matter accumulation on top of the parent soil. Nevertheless, the bare soil did contain 3.6% soil organic matter and some bacterial biomass (Table 1), which may be due to its close proximity
175 (meter-scale) to vegetated areas. However, despite its relatively high content of organic matter and bacteria, it showed very low oxygen consumption during incubation ($0.3 \mu\text{M g}^{-1}$ dry weight (d.w.) d^{-1}) compared to the three organic soils ($5\text{-}14 \mu\text{M g}^{-1}$ d.w. d^{-1}). This indicates that the soil organic matter is not very reactive and/or that the bacterial activity



per cell is low in this soil. It should be stressed, that the measured oxygen consumption is not necessarily the same that it would be in nature, as the soil was disturbed (homogenized by hand) which may increase bioavailability of soil organic matter. Fungal biomass (determined as ITS2 gene copies) was much higher in the coniferous soil compared to the other organic soil types, which may be expected as fungi are known to play a key role in degradation of needle litter (Boberg, 2009).

Table 1. Soil parameters determined from homogenized samples with major roots removed. Average of three replicate extractions and analyses for 16s and ITS2 (\pm standard deviation) and one for the other parameters. Sample depth corresponds roughly to the depth of the organic layer after removing litter from the top and is the average depth of 10 pooled soil cores. At the Arctic bare soil no organic layer was present. O₂-consumption is measured during mineralization experiments and may be different from that in nature. All parameters except moisture are on dry weight basis.

Soil	Depth (cm)	SOM (%)	pH _{H2O}	Moisture (weight %)	16s (copies g ⁻¹)	ITS2 (copies g ⁻¹)	O ₂ -consumption (μM g ⁻¹ d ⁻¹)
Temp. conif.	0-6	78	3.8	45	6.2·10 ¹⁰ ±2.8·10 ¹⁰	5.5·10 ⁸ ±6.9·10 ⁷	9
Temp. beech	0-5	20	3.9	46	4.2·10 ¹⁰ ±1.2·10 ¹⁰	3.7·10 ⁷ ±1.7·10 ⁷	5
Arctic heath	0-6	36	5.3	51	2.8·10 ¹⁰ ±4.5·10 ¹⁰	7.0·10 ⁷ ±1.5·10 ⁶	14
Arctic bare	0-5	3.6	7.3	8.1	2.2·10 ⁹ ±1.1·10 ⁹	2.4·10 ⁶ ±6.8·10 ⁵	0.3

190 3.2 BVOC mineralization

As model compounds we chose six BVOCs that have well described natural sources, are commonly detected in nature and have quite different molecular weights and physical/chemical properties (Table 2).



195 **Table 2.** Characteristics of the model compounds sorted by boiling point (BP). X means clear evidence for specified source in the environment. (X) means that some evidence exists. ∞ means unlimited solubility (miscible).

Name	Cas. No.	BP (°C)	Molecular weight	S _w (mg L ⁻¹)	Plant source	Soil/microbial source	Anthropo- genic source
Chloroform	67-66-3	61	119	8000	(X) ^k	X ^{i,j}	X
Methanol	67-56-1	65	32	∞	X ^f	X ^{a,b,e}	X
p-cymene	99-87-6	177	134	23	X ^{g,m}	(X) ^{a,e}	
Benzaldehyde	100-52-7	178	106	3000	X ^{f,m}	X ^{c,d}	X
Acetophenone	98-86-2	202	120	5500	X ^m	X ^{c,d}	X
Geraniol	106-24-1	230	154	686	X ^{l,h}	(X) ^{l,s}	

^aAsensio et al., 2007. ^bSchink and Zeikus, 1980. ^cGutiérrez-Luna et al., 2010. ^dMcNeal and Herbert, 2009. ^eBäck et al., 2010.

^fKesselmeier and Staudt, 1999. ^gOrtega et al., 2008. ^hChen and Viljoen, 2010. ⁱHoekstra et al., 1998. ^jAlbers et al., 2010. ^kLaternus and Matucha, 2008. ^lSchulz and Dickschat, 2007. ^mJardine et al., 2010.

200 ^{*}Limited evidence for a soil or microbial source of these two monoterpenoids, but clear evidence for a general monoterpene production in soil and by various microorganisms (e.g. Schulz and Dickschat, 2007; Leff and Fierer, 2008; McNeal and Herbert, 2009; Bäck et al., 2010).

Five of the six BVOCs were rapidly mineralized in all four soils included in the mineralization experiment, with chloroform showing somewhat slower mineralization (Fig. 3). None of the sterilized soil samples showed any detectable mineralization so the degradation of the BVOCs was in all cases microbially derived. However, great differences were observed with regards to speed of mineralization, extent of mineralization and variation between soil types. Methanol and benzaldehyde showed the highest mineralization rates. Especially for methanol (Fig. 3b), mineralization was so fast that the CO₂ transfer rate from soil to trap was most likely determining the shape of the mineralization curve rather than the speed of mineralization. For example, the theoretical initial (0-2 h) mineralization rate in the Arctic Heath soil that can be determined with the applied method would be 40% h⁻¹ as calculated from the curve in Supplementary Fig. S1, and the observed mineralization of methanol in that soil type was 39% h⁻¹ (Table 3). Also benzaldehyde mineralization was so fast that probably the CO₂ transfer rate influenced the shape of the mineralization curve (Fig. 3d). The fact that methanol is degraded quickly in soil is not a surprise, as many isolated soil bacteria have the capability to degrade this BVOC (Kolb, 2009), and different temperate grassland and forest soils have been found to contain at least 10⁶ bacteria with the capability to degrade methanol per gram of soil (Stacheter et al., 2013). However, our data are the first to demonstrate degradation of methanol within the range of observed atmospheric concentrations (less than 100 ng L⁻¹, Seco et al. (2007)). Degradation of benzaldehyde in soil or by soil microorganisms has not been demonstrated, but benzaldehyde mineralization by pure microbial cultures has been shown (Kamada et al., 2002).

Following methanol and benzaldehyde, geraniol and acetophenone had the highest mineralization rates with most of the mineralization occurring within the first 24 hours of incubation (Fig. 3e,f). These four rapidly degraded compounds had in common that no clear difference in mineralization rate was observed between the soil types (Table 3). For methanol and partly for benzaldehyde this observation could be influenced by method limitations (too fast mineralization to be kept in pace by transfer of CO₂ to the trap may have masked any differences), however for geraniol and acetophenone



this was not the case. In other words, Arctic soils mineralized these compounds as quickly as temperate forest soils and perhaps even more interestingly, the Arctic bare soil showed similar mineralization rates as the organic soil types. This is despite a much lower abundance of microorganisms as determined by qPCR and a much lower microbial
230 heterotrophic activity during incubation as determined by oxygen consumption (Table 1). Geraniol mineralization has previously been investigated in soil sampled underneath *Populus tremula* tree crowns (Owen et al., 2007). The mineralization observed in that study was different from the one we observed, with an initial lag phase with less than 5% mineralization in the first ~10 hours. The lag phase was followed by maximum mineralization rates of 1-3% h⁻¹ which is close to what we observed right after the start of incubation (Table 3). An extremely high geraniol
235 concentration of 600 mg kg⁻¹ soil was used in that study compared to 6-11 µg kg⁻¹ soil in ours, which is the most likely cause of this difference in mineralization. The geraniol concentration used by Owen et al. (2007) would allow growth with geraniol as substrate (hence the lag phase) while the concentrations we used would allow only very limited microbial growth. However, the two studies all in all demonstrate that oxygenated monoterpenes may be degraded within a very large concentration range in soil.

240 P-cymene mineralization showed as the only BVOC clear differences between the soil types (Fig. 3c). Initial mineralization rates were by far the highest in the coniferous forest soil (10% h⁻¹, Table 3) followed by the Arctic heath soil (2% h⁻¹), the beech forest soil (0.4% h⁻¹) and the Arctic bare soil (0.2% h⁻¹). In other words, the coniferous forest soil showed a 25 times higher initial mineralization rate compared to the beech forest soil sampled just 750 meters
245 away. In addition, the three soils with slowest mineralization showed a slightly s-shaped mineralization curve meaning that mineralization rate increased after an initial lag-phase with slower mineralization (Fig. 3c and Table 3). All in all it appears that the coniferous forest soil is especially adapted to degrade p-cymene. P-cymene is a hydrocarbon monoterpene (monoterpene without heteroatoms) and these are emitted in very high concentrations in coniferous forests (Guenther et al., 1994; Rinne et al., 2009). Our measurements also showed a much higher concentration of this BVOC
250 group in the atmosphere of the coniferous forest compared to the other sampling sites (Table 4). In addition, needle litter emits high amounts of hydrocarbon monoterpenes (Aaltonen et al., 2011; Faiola et al., 2014) exposing the soil to these compounds found in higher concentrations in soil under conifers than deciduous trees (Smolander et al., 2006). All in all it seems likely that the high adaptation for p-cymene mineralization in the coniferous forest soil is caused by a high natural input of hydrocarbon monoterpenes to this soil type.

255 Chloroform, which is a well-known pollutant but also a natural product in soil (Hoekstra et al., 1998; Albers et al., 2010; Johnsen et al., 2016), was mineralized in all four soils (Fig. 3a), but at much slower rates compared to the other BVOCs (Table 3). Chloroform mineralization was previously determined at 10°C in a spruce forest soil in which initial mineralization rates of 0.01-0.04 % h⁻¹ were observed (Albers et al., 2011). These rates are roughly ten times lower than
260 the ones observed in our study (0.2-0.5% h⁻¹, Table 3). The spruce forest soil was similar to the coniferous forest soil used in our study, but there was a difference in chloroform concentration, which in our case was 4-7 µg kg⁻¹ soil and in the previous study was 350 µg kg⁻¹ soil. This stresses that the concentration used during incubation may to a high degree determine how fast the compound is mineralized. On the other hand, if mineralization rates are recalculated to a mass-unit per time-unit, differences in the case of chloroform mineralization would be much smaller between the two
265 studies.




Figure 3. Mineralization curves for six BVOCs in soil from coniferous forest, beech forest, Arctic heath and Arctic bare soil. Initial BVOC concentrations varied from 0.8-11 $\mu\text{g kg soil}^{-1}$. "Ster." means that soil was sterilized twice by autoclaving. Error bars are standard deviation of triplicate incubations. Some error bars are smaller than the symbols.

270

The extent of mineralization (determined as $^{14}\text{CO}_2$ release at the termination of experiment) was in general similar between soils but differed greatly between the BVOCs. Methanol showed almost 100% $^{14}\text{CO}_2$ release while acetophenone and geraniol released only 40%. The 40% release should not be interpreted as if only 40% of the compound was degraded, but rather that 60% of the mineralized compound was used as a carbon source for microbial growth. This is a generally accepted interpretation of mineralization curves that often go to yields of only 40-50% with the remaining part incorporated into biomass that is only slowly mineralized along with microbial turnover (Nowak et

275



al., 2011; Glanville et al., 2016). Just after incubation, we extracted non-degraded or metabolized BVOCs with methanol, and only sterilized samples released a major pool of methanol-extractable ^{14}C (typically between 50 and 280 95%) while non-sterilized samples released less than 5%. This strongly supports the interpretation that all BVOC was degraded. Geraniol was an exception from this with 15-25% of added ^{14}C extracted in non-sterilized samples and 83-92% extracted in sterilized samples. While methanol, benzaldehyde, acetophenone and p-cymene were conclusively degraded completely within 140 hours (and presumably much faster) we therefore cannot exclude the possibility that some less degradable degradation products of geraniol have accumulated. It has been shown that some fungi have the 285 ability to metabolize geraniol into various derivatives (Demyttenaere et al., 2000).

Based on extent of mineralization, some compounds (e.g. methanol) were used only as a source of energy (as electron donor), while others (e.g. geraniol and acetophenone) were also used as a carbon source for growth. Recently, Gunina et al. (2017) suggested that the oxidation state of a C-atom determines how much is released as CO_2 , and how much is 290 incorporated into biomass. They found a positive relationship between carbon oxidation state and $^{14}\text{CO}_2$ -release for seven easily degradable low molecular weight sugars, acids and amino acids. However, the carbon atom in methanol (oxidation state -2) is more reduced than the labeled carbon atoms in geraniol (oxidation state 0), benzaldehyde and acetophenone (both -1), so the oxidation state does not determine mineralization extent of the model BVOCs.

295 P-cymene was an exception from the minor difference in mineralization extent between soil types. In soil from the coniferous forest and from the Arctic heath, more than 80% of the ^{14}C was liberated as $^{14}\text{CO}_2$ (Fig. 3c). In the Arctic bare soil only half of this release was measured, while the beech forest soil was in between. In all soils, p-cymene dissipation was complete at the end of the experiment. One possible explanation for this difference is that different 300 microorganisms degrade p-cymene in the studied soils and that these different organisms have different degradation strategies for the compound, i.e. different fractions used for energy and growth. Another, perhaps more likely explanation, is that p-cymene is used as a carbon source mainly when degradation is occurring along with microbial growth. This explanation is supported by the fact that the slower the initial degradation is and the more s-shaped the mineralization curves are (presence of lag phase, Table 3), the more carbon seems to be accumulated into biomass (less 305 $^{14}\text{CO}_2$ -release, Fig. 3c). This is also supported by the earlier observed higher mineralization extent of ^{14}C -geraniol at high concentration that supported growth (mineralization extent of 64-75%, Owen et al., 2007) compared to the mineralization extent we observed for this compound with no or very little growth (33-46%, Fig. 3f). In addition, the highest mineralization extent in the case of geraniol was observed in the coniferous forest soil, which was the only soil where a lag phase (though very weak) was observed (Table 3).



310 **Table 3.** Mineralization parameters calculated from the mineralization experiment shown in Fig. 3. Initial mineralization rate is calculated as the average rate during the first two hours of incubation. A lag phase is noted where the initial mineralization rate is not the highest. (Yes) denotes a very weak lag phase.

	Initial mineralization rate (% h ⁻¹)				Lag phase?			
	Conif.	Beech	Heath	Bare	Conif.	Beech	Heath	Bare
Chloroform	0.20	0.17	0.34	0.48	No	No	No	No
Methanol	38	35	39	25	No	No	No	No
p-Cymene	10	0.4	2.0	0.2	No	Yes	(Yes)	Yes
Benzaldehyde	14	13	16	12	No	No	No	No
Acetophenone	2.0	2.6	4.3	2.3	(Yes)	(Yes)	No	(Yes)
Geraniol	1.5	3.5	4.3	2.3	(Yes)	No	No	No

315 The potential for very fast mineralization of different BVOCs in different temperate and Arctic soils may have significant environmental implications. A few previous studies have shown deposition of BVOCs onto soil (Ramirez et al., 2010; Spielmann et al., 2017) or a bidirectional atmosphere/soil exchange of certain BVOCs (Asensio et al., 2007; Asensio et al., 2008; Gray et al., 2014), but the mechanism behind the uptake of BVOCs into or onto soil has been largely uninvestigated. Our results suggest that BVOCs will be taken up from the atmosphere by microorganisms that then mineralize the compounds. The concentration of BVOCs in the atmosphere is very low, also at the sites where we
 320 sampled soil (Table 4). Mineralization experiments cannot be carried out at such low concentrations but we used BVOC concentrations that are much more realistic than those used in previous degradation studies. Furthermore, similar atmospheric concentrations as we used for incubations have been observed in nature for methanol (Seco et al., 2007), chloroform (Albers et al., 2011) and monoterpenes (Barney et al., 2009).

325 It is therefore very likely that soil microorganisms also take up and mineralize BVOCs in the natural environment and most likely also in urban environments, where concentrations in the air can be much higher due to additional anthropogenic input (Seco et al., 2007). *In situ* uptake studies using e.g. PTR-MS technology should be carried out in order to provide quantitative estimates of the importance of BVOC uptake in soil. However, simultaneous formation and degradation of the compounds is a complicating aspect in such studies. The use of labeled compounds in the field to
 330 determine simultaneous formation and degradation, as previously done in laboratory studies with methane (von Fischer and Hedin, 2002) and methyl halides (Rhew et al., 2003), could be a great supplement to more conventional PTR-MS studies.



335 **Table 4.** Atmospheric concentrations of relevant BVOCs (mean \pm standard deviation, $n=3$) measured 10 cm above soil surface (coniferous, beech and Arctic bare sites) or 5 cm above the canopy (Arctic heath site) the day of soil sampling. Methanol could not be analyzed with the applied methods.

Name	Atmospheric concentration (ng L ⁻¹)		
	Coniferous*	Beech	Arctic**
Oxygenated monoterpenes	0.00 \pm 0.00	0.00 \pm 0.00	0.01 \pm 0.01
Hydrocarbon monoterpenes	3.36 ^a \pm 0.32	0.37 ^b \pm 0.12	0.71 ^c \pm 0.10
Benzaldehyde	1.01 \pm 0.03	1.14 \pm 0.08	0.00 \pm 0.00
Acetophenone	0.44 \pm 0.06	0.59 \pm 0.03	0.01 \pm 0.01
Chloroform	0.10 \pm 0.02	0.06 \pm 0.00	0.06 \pm 0.00

* $n=2$ due to loss of a sample, except for chloroform ($n=3$). **One sample from the bare soil, two from the Arctic Heath. ^aMainly pinenes, camphene, carene and p-cymene. ^bMainly camphene, α -pinene, δ -terpinene and carene. ^cMainly δ -terpinene.

340 In addition to the uptake from the atmosphere, the very fast mineralization rates are likely important in shaping the net BVOC emissions from soil. The net BVOC release from soil to the atmosphere in general is low compared to the plant emissions (Peñuelas et al., 2014), but emissions may represent a minor portion of the amount that was excreted by soil microbes (Insam and Seewald, 2010; Garbeva et al., 2014) or by roots (Lin et al., 2007; Delory et al., 2016), produced for example with the purpose of communication. It is thus possible that BVOCs are a significant source of carbon to soil

345 microbes and hence that BVOC formation and degradation may be an important but little recognized part of internal carbon cycling in soil. In addition, plant litter releases BVOCs from both abiotic and biotic processes (for example terpenoids (Faiola et al., 2013) and methanol (Gray et al., 2010)). These BVOCs may to a large degree never reach the atmosphere but rather be an input of degradable carbon to microorganisms in the top soil.

350 4 Conclusions

In conclusion, we have shown that six chemically very different BVOCs are all mineralized by microbes in Arctic and temperate soils at environmentally relevant concentrations. Five of the BVOCs were mineralized very quickly, but still we observed a relatively large compound-to-compound variation in mineralization rate as well as mineralization extent compared to a much lower soil-to-soil variation. P-cymene was an exception from this pattern with both mineralization

355 rate and extent differing between soils of different origin.

It is thus clear that soil microbes are able to degrade completely and quickly BVOCs released by aboveground vegetation, soil microbes and plant roots and additional studies should be carried out to quantify this process in nature. In addition to the possible atmospheric implications of BVOC degradation by soil microbes, BVOC formation and degradation may be an important but little recognized part of internal carbon cycling in soil.

360



Acknowledgments. We wish to thank Pia Bach Jacobsen, GEUS, for excellent technical assistance in the laboratory. We also thank Anders Priemé, Copenhagen University for advice on the qPCR targeting the fungal community and for providing the ITS2 standard. The study was financially supported by the Danish Council for Independent Research | Natural Sciences (DFR-4002-00495), the Villum foundation (VKR022589), and by the Danish National Research Foundation (CENPERM DNR100).

Competing interests. The authors declare that they have no conflict of interest.

Data availability. The data set related to Figure 3 has been provided as a supplement.

370 **References**

- Aaltonen, H., Pumpanen, J., Pihlatie, M., Hakola, H., Hellén, H., Kulmala, L., Vesala, T. and Bäck, J.: Boreal pine forest floor biogenic volatile organic compound emissions peak in early summer and autumn, *Agric. For. Meteorol.*, 151(6), 682–691, doi:10.1016/J.AGRFORMET.2010.12.010, 2011.
- Albers, C. N., Laier, T. and Jacobsen, O. S.: Formation, fate and leaching of chloroform in coniferous forest soils, *Appl. Geochemistry*, 25, 1525–1535, doi:10.1016/j.apgeochem.2010.08.003, 2010.
- Albers, C. N., Jacobsen, O. S., Flores, É. M. M., Pereira, J. S. F. and Laier, T.: Spatial variation in natural formation of chloroform in the soils of four coniferous forests, *Biogeochemistry*, 103, 317–334, doi:10.1007/s10533-010-9467-9, 2011.
- Asensio, D., Peñuelas, J., Filella, I. and Llusà, J.: On-line screening of soil VOCs exchange responses to moisture, temperature and root presence, *Plant Soil*, 291(1-2), 249–261, doi:10.1007/s11104-006-9190-4, 2007.
- Asensio, D., Peñuelas, J., Prieto, P., Estiarte, M., Filella, I. and Llusà, J.: Interannual and seasonal changes in the soil exchange rates of monoterpenes and other VOCs in a Mediterranean shrubland, *Eur. J. Soil Sci.*, 59(5), 878–891, doi:10.1111/j.1365-2389.2008.01057.x, 2008.
- Barney, J. N., Sparks, J. P., Greenberg, J., Whitlow, T. H. and Guenther, A.: Biogenic volatile organic compounds from an invasive species: impacts on plant–plant interactions, *Plant Ecol.*, 203(2), 195–205, doi:10.1007/s11258-008-9529-4, 2009.
- Boberg, J.: Litter decomposing fungi in boreal forests, 67 pp., Dept. of Forest Mycology and Pathology, Swedish University of Agricultural Sciences., 2009.
- Bäck, J., Aaltonen, H., Hellén, H., Kajos, M. K., Patokoski, J., Taipale, R., Pumpanen, J. and Heinonsalo, J.: Variable emissions of microbial volatile organic compounds (MVOCs) from root-associated fungi isolated from Scots pine, *Atmos. Environ.*, 44(30), 3651–3659, doi:10.1016/J.ATMOSENV.2010.06.042, 2010.



Chen, W. and Viljoen, A. M.: Geraniol — A review of a commercially important fragrance material, *South African J. Bot.*, 76(4), 643–651, doi:10.1016/J.SAJB.2010.05.008, 2010.

Christiansen, C. T., Haugwitz, M. S., Priemé, A., Nielsen, C. S., Elberling, B., Michelsen, A., Grogan, P. and Blok, D.:
395 Enhanced summer warming reduces fungal decomposer diversity and litter mass loss more strongly in dry than in wet
tundra, *Glob. Chang. Biol.*, 23(1), 406–420, doi:10.1111/gcb.13362, 2017.

Cleveland, C. C. and Yavitt, J. B.: Consumption of atmospheric isoprene in soil, *Geophys. Res. Lett.*, 24(19), 2379–
2382, doi:10.1029/97GL02451, 1997.

Cripps, R. E.: The microbial metabolism of acetophenone. Metabolism of acetophenone and some chloroacetophenones
400 by an *Arthrobacter* species., *Biochem. J.*, 152(2), 233–241 [online] Available from:
<http://www.ncbi.nlm.nih.gov/pubmed/4061> (Accessed 7 December 2017), 1975.

Delory, B. M., Delaplace, P., Fauconnier, M.-L. and du Jardin, P.: Root-emitted volatile organic compounds: can they
mediate belowground plant-plant interactions?, *Plant Soil*, 402(1-2), 1–26, doi:10.1007/s11104-016-2823-3, 2016.

Demyttenaere, J. C. R., del Carmen Herrera, M. and De Kimpe, N.: Biotransformation of geraniol, nerol and citral by
405 sporulated surface cultures of *Aspergillus niger* and *Penicillium* sp., *Phytochemistry*, 55(4), 363–373,
doi:10.1016/S0031-9422(00)00330-7, 2000.

Faiola, C. L., VanderSchelden, G. S., Wen, M., Elloy, F. C., Cobos, D. R., Watts, R. J., Jobson, B. T. and VanReken, T.
M.: SOA Formation Potential of Emissions from Soil and Leaf Litter, *Environ. Sci. Technol.*, 48(2), 938–946,
doi:10.1021/es4040045, 2014.

410 Feld, L., Nielsen, T. K., Hansen, L. H., Aamand, J. and Albers, C. N.: Establishment of Bacterial Herbicide Degraders
in a Rapid Sand Filter for Bioremediation of Phenoxypropionate-Polluted Groundwater., *Appl. Environ. Microbiol.*,
82(3), 878–887, doi:10.1128/AEM.02600-15, 2016.

Von Fischer, J. C. and Hedin, L. O.: Separating methane production and consumption with a field-based isotope pool
dilution technique, *Global Biogeochem. Cycles*, 16(3), 8–1 – 8–13, doi:10.1029/2001GB001448, 2002.

415 Garbeva, P., Hordijk, C., Gerards, S. and de Boer, W.: Volatile-mediated interactions between phylogenetically
different soil bacteria, *Front. Microbiol.*, 5, 289, doi:10.3389/fmicb.2014.00289, 2014.

Glanville, H. C., Hill, P. W., Schnepf, A., Oburger, E. and Jones, D. L.: Combined use of empirical data and
mathematical modelling to better estimate the microbial turnover of isotopically labelled carbon substrates in soil, *Soil
Biol. Biochem.*, 94, 154–168, doi:10.1016/j.soilbio.2015.11.016, 2016.

420 Glasius, M. and Goldstein, A. H.: Recent Discoveries and Future Challenges in Atmospheric Organic Chemistry,
Environ. Sci. Technol., 50(6), 2754–2764, doi:10.1021/acs.est.5b05105, 2016.

Gray, C. M., Monson, R. K. and Fierer, N.: Emissions of volatile organic compounds during the decomposition of plant
litter, *J. Geophys. Res.*, 115(G3), G03015, doi:10.1029/2010JG001291, 2010.



- 425 Gray, C. M., Monson, R. K. and Fierer, N.: Biotic and abiotic controls on biogenic volatile organic compound fluxes from a subalpine forest floor, *J. Geophys. Res. Biogeosciences*, 119(4), 547–556, doi:10.1002/2013JG002575, 2014.
- Gray, C. M., Helmig, D. and Fierer, N.: Bacteria and fungi associated with isoprene consumption in soil, *Elem. Sci. Anthr.*, 3(0), 000053, doi:10.12952/journal.elementa.000053, 2015.
- Guenther, A., Zimmerman, P. and Wildermuth, M.: Natural volatile organic compound emission rate estimates for U.S. woodland landscapes, *Atmos. Environ.*, 28(6), 1197–1210, doi:10.1016/1352-2310(94)90297-6, 1994.
- 430 Gunina, A., Smith, A. R., Kuzyakov, Y. and Jones, D. L.: Microbial uptake and utilization of low molecular weight organic substrates in soil depend on carbon oxidation state, *Biogeochemistry*, 133(1), 89–100, doi:10.1007/s10533-017-0313-1, 2017.
- Gutiérrez-Luna, F. M., López-Bucio, J., Altamirano-Hernández, J., Valencia-Cantero, E., de la Cruz, H. R. and Macías-Rodríguez, L.: Plant growth-promoting rhizobacteria modulate root-system architecture in *Arabidopsis thaliana* through volatile organic compound emission, *Symbiosis*, 51(1), 75–83, doi:10.1007/s13199-010-0066-2, 2010.
- 435 Hoekstra, E. J., de Leer, E. W. B. and Brinkman, U. A. T.: Natural Formation of Chloroform and Brominated Trihalomethanes in Soil, *Environ. Sci. Technol.*, 32(23), 3724–3729, doi:10.1021/es980127c, 1998.
- Insam, H. and Seewald, M. S. A.: Volatile organic compounds (VOCs) in soils, *Biol. Fertil. Soils*, 46(3), 199–213, doi:10.1007/s00374-010-0442-3, 2010.
- 440 Jardine, K., Abrell, L., Kurc, S. A., Huxman, T., Ortega, J. and Guenther, A.: Volatile organic compound emissions from *Larrea tridentata* (creosotebush), *Atmos. Chem. Phys.*, 10(24), 12191–12206, doi:10.5194/acp-10-12191-2010, 2010.
- Johnsen, A. R., Jacobsen, O. S., Gudmundsson, L. and Albers, C. N.: Chloroform emissions from arctic and subarctic ecosystems in Greenland and Northern Scandinavia, *Biogeochemistry*, 130(1-2), 53–65, doi:10.1007/s10533-016-0241-5, 2016.
- 445 Kamada, F., Abe, S., Hiratsuka, N., Wariishi, H. and Tanaka, H.: Mineralization of aromatic compounds by brown-rot basidiomycetes – mechanisms involved in initial attack on the aromatic ring, *Microbiology*, 148(6), 1939–1946, doi:10.1099/00221287-148-6-1939, 2002.
- Kesselmeier, J. and Staudt, M.: Biogenic Volatile Organic Compounds (VOC): An Overview on Emission, Physiology and Ecology, *J. Atmos. Chem.*, 33(1), 23–88, doi:10.1023/A:1006127516791, 1999.
- 450 El Khawand, M., Crombie, A. T., Johnston, A., Vavlline, D. V., McAuliffe, J. C., Latone, J. A., Primak, Y. A., Lee, S.-K., Whited, G. M., McGenity, T. J. and Murrell, J. C.: Isolation of isoprene degrading bacteria from soils, development of isoA gene probes and identification of the active isoprene-degrading soil community using DNA-stable isotope probing, *Environ. Microbiol.*, 18(8), 2743–2753, doi:10.1111/1462-2920.13345, 2016.



- 455 Kleinheinz, G. T., Bagley, S. T., John, W. P. St., Rughani, J. R. and McGinnis, G. D.: Characterization of Alpha-Pinene-Degrading Microorganisms and Application to a Bench-Scale Biofiltration System for VOC Degradation, *Arch. Environ. Contam. Toxicol.*, 37(2), 151–157, doi:10.1007/s002449900500, 1999.
- Kolb, S.: Aerobic methanol-oxidizing Bacteria in soil, *FEMS Microbiol. Lett.*, 300(1), 1–10, doi:10.1111/j.1574-6968.2009.01681.x, 2009.
- 460 Kramshøj, M., Vedel-Petersen, I., Schollert, M., Rinnan, Å., Nymand, J., Ro-Poulsen, H. and Rinnan, R.: Large increases in Arctic biogenic volatile emissions are a direct effect of warming, *Nat. Geosci.*, 9(5), 349–352, doi:10.1038/ngeo2692, 2016.
- Laothawornkitkul, J., Taylor, J. E., Paul, N. D. and Hewitt, C. N.: Biogenic volatile organic compounds in the Earth system, *New Phytol.*, 183(1), 27–51, doi:10.1111/j.1469-8137.2009.02859.x, 2009.
- 465 Laturmus, F. and Matucha, M.: Chloride - a precursor in the formation of volatile organochlorines by forest plants?, *J. Environ. Radioact.*, 99(1), 119–125, doi:10.1016/j.jenvrad.2007.07.008, 2008.
- Leff, J. W. and Fierer, N.: Volatile organic compound (VOC) emissions from soil and litter samples, *Soil Biol. Biochem.*, 40(7), 1629–1636, doi:10.1016/J.SOILBIO.2008.01.018, 2008.
- Lin, C., Owen, S. M. and Peñuelas, J.: Volatile organic compounds in the roots and rhizosphere of *Pinus* spp., *Soil Biol. Biochem.*, 39(4), 951–960, doi:10.1016/J.SOILBIO.2006.11.007, 2007.
- 470 McNeal, K. S. and Herbert, B. E.: Volatile Organic Metabolites as Indicators of Soil Microbial Activity and Community Composition Shifts, *Soil Sci. Soc. Am. J.*, 73(2), 579–588, doi:10.2136/sssaj2007.0245, 2009.
- Misra, G., Pavlostathis, S. G., Perdue, E. M. and Araujo, R.: Aerobic biodegradation of selected monoterpenes, *Appl. Microbiol. Biotechnol.*, 45(6), 831–838, doi:10.1007/s002530050770, 1996.
- 475 Nowak, K. M., Miltner, A., Gehre, M., Schäffer, A. and Kästner, M.: Formation and Fate of Bound Residues from Microbial Biomass during 2,4-D Degradation in Soil, *Environ. Sci. Technol.*, 45(3), 999–1006, doi:10.1021/es103097f, 2011.
- Ortega, J., Helmig, D., Daly, R. W., Tanner, D. M., Guenther, A. B. and Herrick, J. D.: Approaches for quantifying reactive and low-volatility biogenic organic compound emissions by vegetation enclosure techniques – Part B: Applications, *Chemosphere*, 72(3), 365–380, doi:10.1016/j.chemosphere.2008.02.054, 2008.
- 480 Owen, S. M., Clark, S., Pompe, M. and Semple, K. T.: Biogenic volatile organic compounds as potential carbon sources for microbial communities in soil from the rhizosphere of *Populus tremula*, *FEMS Microbiol. Lett.*, 268(1), 34–39, doi:10.1111/j.1574-6968.2006.00602.x, 2007.
- Peñuelas, J. and Staudt, M.: BVOCs and global change, *Trends Plant Sci.*, 15(3), 133–144, doi:10.1016/J.TPLANTS.2009.12.005, 2010.
- 485



- Peñuelas, J., Asensio, D., Tholl, D., Wenke, K., Rosenkranz, M., Piechulla, B. and Schnitzler, J. P.: Biogenic volatile emissions from the soil. *Plant. Cell Environ.*, 37(8), 1866–1891, doi:10.1111/pce.12340, 2014.
- Ramirez, K. S., Lauber, C. L. and Fierer, N.: Microbial consumption and production of volatile organic compounds at the soil-litter interface, *Biogeochemistry*, 99(1-3), 97–107, doi:10.1007/s10533-009-9393-x, 2010.
- 490 Rhew, R. C., M, A. and ES, S.: Measuring terrestrial fluxes of methyl chloride and methyl bromide using a stable isotope tracer technique, *Geophys. Res. Lett.*, 30(21), 2103, doi:10.1029/2003GL018160, 2003.
- Rinne, J., Back, J. and Hakola, H.: Biogenic volatile organic compound emissions from the Eurasian taiga: current knowledge and future directions, *Boreal Environ. Res.*, 14(4), 807–826 [online] Available from: https://apps.webofknowledge.com/full_record.do?product=WOS&search_mode=GeneralSearch&qid=2&SID=N2EGBwkHspWF9UQ7ty8&page=2&doc=12 (Accessed 24 October 2017), 2009.
- 495 Schink, B. and Zeikus, J. G.: Microbial methanol formation: A major end product of pectin metabolism, *Curr. Microbiol.*, 4(6), 387–389, doi:10.1007/BF02605383, 1980.
- Schulz, S. and Dickschat, J. S.: Bacterial volatiles: the smell of small organisms, *Nat. Prod. Rep.*, 24(4), 814–842, doi:10.1039/b507392h, 2007.
- 500 Seco, R., Peñuelas, J. and Filella, I.: Short-chain oxygenated VOCs: Emission and uptake by plants and atmospheric sources, sinks, and concentrations, *Atmos. Environ.*, 41(12), 2477–2499, doi:10.1016/j.atmosenv.2006.11.029, 2007.
- Smolander, A., Ketola, R. A., Kotiaho, T., Kanerva, S., Suominen, K. and Kitunen, V.: Volatile monoterpenes in soil atmosphere under birch and conifers: Effects on soil N transformations, *Soil Biol. Biochem.*, 38(12), 3436–3442, doi:10.1016/j.soilbio.2006.05.019, 2006.
- 505 Spielmann, F. M., Langebner, S., Ghirardo, A., Hansel, A., Schnitzler, J.-P. and Wohlfahrt, G.: Isoprene and α -pinene deposition to grassland mesocosms, *Plant Soil*, 410(1-2), 313–322, doi:10.1007/s11104-016-3009-8, 2017.
- Stacheter, A., Noll, M., Lee, C. K., Selzer, M., Glowik, B., Ebertsch, L., Mertel, R., Schulz, D., Lampert, N., Drake, H. L. and Kolb, S.: Methanol oxidation by temperate soils and environmental determinants of associated methylotrophs, *ISME J.*, 7(5), 1051–1064, doi:10.1038/ismej.2012.167, 2013.