

Dear reviewer,

Thank you for taking time to review our manuscript.

We studied your comments and revised our draft accordingly.

We also went through our manuscript and made corrections on the texts.

Changes to the texts were marked in the manuscript and were presented as follows.

Hope that our revised draft will meet with your approval.

Best wishes,

Qinchuan Xin

### **Anonymous Referee # 2**

I appreciate authors' efforts in revising their manuscript. However, I am sorry but I still find that the manuscript falls short of being ready for publication. I have new concerns after reading the manuscript and I summarize my primary concerns below. I am also attaching an annotated version of the manuscript in which authors can find other minor comments.

**Reply:** please find detailed responses to your annotated comments after our replies to your primary concerns.

At a number of places in the manuscript, the authors have stressed that their scheme can be used for modeling leaf phenology in land surface models like the community land model (CLM). However, one thing that is being neglected here is that models like CLM are able to simulate leaf area index as a function of atmospheric CO<sub>2</sub> concentration. In authors' approach the variable  $m=LAI/GPP$  is a constant and calculated using present-day LAI and GPP. As such then, it is not possible to use this approach in land surface models that are designed for application for future climate and CO<sub>2</sub> scenarios.

**Reply:** First, our approach is different from others in terms of the leaf phenology sub-model. Although our approach does not account for CO<sub>2</sub> directly, it does account for CO<sub>2</sub> indirectly if replacing the MOD17 approach with a canopy photosynthesis sub-model that accounts for CO<sub>2</sub>. If the canopy photosynthesis sub-model accounts for the impacts of CO<sub>2</sub>, the solved steady-state LAI would become a function of CO<sub>2</sub> and so does the modeled LAI. Whether our approach of leaf phenology modeling could account for the impacts of specific climate variables is largely dependent on the GPP models we used.

Second, for the current phenology models, there are nearly no direct considerations for CO<sub>2</sub>. The table 2 from Richardson et al. (2005) in Global Change Biology summarized the phenology models in 14 terrestrial biosphere model. Most of the them only account for the influence of temperature and none of them accounts for the influences of CO<sub>2</sub> directly. In the community land model (CLM), there are also no functions that directly links leaf phenology to CO<sub>2</sub>. In the phenology sub-model, CLM divided vegetation into 3 classes of the evergreen type, the seasonal-deciduous type, and the

stress-deciduous type. None of these 3 classes include considerations of leaf area index as a function of CO<sub>2</sub> directly. It is only able to simulate leaf area index as a function of atmospheric CO<sub>2</sub> concentration indirectly.

**Table 2** Summary of models used in this analysis and their representation of phenology and seasonality of leaf area index (LAI). For models with 'prognostic' phenology, the seasonality of LAI is predicted based on climatic drivers; for those with 'prescribed' phenology, an average seasonal LAI cycle, as derived on a site-by-site basis from satellite (AVHRR) data, was used. Models with semi-prescribed and semi-prognostic phenology represent a hybrid of these approaches. GDD is growing degree days; T is temperature; C is carbon; PFT is plant functional type

Model name	Resolution	Leaf onset	Control on LAI	Leaf loss	Source
BEPS	Daily	Satellite	Satellite	Satellite	Ju <i>et al.</i> (2006)
Biome-BGC	Daily	GDD and radiation sum	Dynamic C allocation	Daylength and low temperature	Thornton <i>et al.</i> (2002)
Can-IBIS	Half-hourly	T threshold	GDD and dynamic C	Prescribed	El Maayar <i>et al.</i> (2002)
CN-CLASS	Half-hourly	C balance	C balance	Daylength and low temperature	Arain <i>et al.</i> (2006)
DLEM	Daily	T <sub>7-day</sub> > threshold	GDD to PFT limit	Daylength and low temperature	Tian <i>et al.</i> (2010)
Ecosys	Hourly	Hours above T threshold	Dynamic C allocation	Hours below T threshold	Grant <i>et al.</i> (2009)
ED2	Half-hourly	Semi-prescribed	Dynamic C allocation	GDD and leaf turnover	Medvigy <i>et al.</i> (2009)
ISAM	Half-hourly	Prescribed	Prescribed	Prescribed	Jain & Yang (2005)
LoTEC	Half-hourly	GDD	GDD	T-dependent turnover	Hanson <i>et al.</i> (2004)
LPJ_wsl	Daily	GDD	GDD	Leaf longevity (prescribed)	Sitch <i>et al.</i> (2003)
ORCHIDEE	Half-hourly	GDD and chilling	Dynamic C allocation	Decreasing T and T threshold	Krinner <i>et al.</i> (2005)
SiB3	Half-hourly	Prescribed	Prescribed	Prescribed	Baker <i>et al.</i> (2008)
SIBCASA	10 min	Prescribed	Prescribed	Prescribed	Schaefer <i>et al.</i> (2008)
SSiB2	Half-hourly	Prescribed	Prescribed	Prescribed	Zhan <i>et al.</i> (2003)

Third, we agree that we need to be cautious about the statements for the model application in land surface models. We removed those inappropriate statements.

I am uncomfortable with authors' comparison of their results with those obtained using the moving average model. The moving average model cannot clearly be used in current generation of land surface models which determine leaf area index prognostically. I suspect the reason the authors have compared results from their new time stepping method to those obtained using the moving average method because this is what they did in the past and are more comfortable with their moving average method. However, this does not mean that the results from the moving average method should be used as a benchmark.

**Reply:** We believe that making comparisons between the time-stepping method and the moving average method could add values to the manuscript. Second, such comparisons show that the time-stepping method is an alternative way to account for the lagged responses of plants to climate variation. And to our knowledge, this has not been shown before in previous studies. Second, the Growing Season Index, which is based on the moving average method, is widely used in many studies and applications. Third, we do not try to use the moving average method as a benchmark. We use the Growing Season Index as a benchmark but there is a need to show that the time-stepping method can be similarly applied to the Growing Season Index approach such that the two methods can be compared directly.

The justification for developing a simple model is still not strong enough in the revised manuscript.

Reply: First, one purpose of this study were to develop a method that can be used within a time-stepping model framework.

Second, improving the modeling accuracies on leaf phenology is also important. We added the citation to Richardson's publication in Global Change Biology in 2005 as follows "evaluation on 14 land surface models in deciduous forests suggested that almost all models predicted start of the season earlier or end of the season later than observations and the model biases were typically 14 weeks or more". Note that the biases for our modeled results (the MBE values in Table 2) are less than 6 days.

Third, we believe developing a simple model is necessary for many applications. To date, it is still nearly impossible for complex land surface models to operate at fine spatial resolution at global scales because of the computational costs. In our preliminary experiments, it takes nearly two days to run the Common Land model version 2014 for only one modeling year at 1 km solution in Tianhe supercomputer (the second fastest supercomputer in the world). A simple model that could be easily operate at regional to global scales would obviously help us understanding the land surface processes. For the same reasons, although there are complex land surface models, the 1 km MODIS GPP product generated based on the simple MOD17 approach is still popular and has been widely used.

In Figure 6 something doesn't seem right with the GLASS LAI data. Notice that in all panels (a, b, and c) the interannual variability in SOS, EOS, and GSL is much larger before the year 2000.

Reply: We do not think there are anything wrong. Please note that the GLASS data were produced using two different satellite data. From 1982 to 1999, the LAI product was generated at the resolution of 0.05 ° from the AVHRR reflectance. From 2000 to 2014, the LAI product was derived from MODIS land surface reflectance (MOD09A1) at the 1 km resolution. It is not surprising that the interannual variability in GLASS reduce significantly after 2000 for the derived SOS, EOS, and GSL.

We added explanations in the texts as follows "Note that the 8-day GLASS LAI product was generated at the 0.05 ° resolution using the AVHRR data from 1982 to 1999 and at the 1000 m resolution using the MODIS from 2000 to 2012. The significantly reduced interannual variability for SOS, EOS, and GSL after 2000 in the GLASS data suggests that the use of the AVHRR and MODIS data in the GLASS dataset could contribute uncertainties in the satellite-derived phenological metrics."

In summary, this manuscript needs to be written in such a way so that its primary argument (that a simple prognostic LAI modeling scheme can be used in complex land surface models) can be brought out much more clearly. In its current form, this message is not brought out properly and leaves a reader confused. Perhaps, another reviewer is needed who can look at the manuscript with a fresh set of eyes.

Reply: We agree that we need to be cautious about the argument that the developed modeling scheme can be used in complex land surface models. We removed these statements in the manuscript and only made a few discussions for potential applications.

Page 1 Line 1: Remove the comparison with moving average scheme.

Reply: We believe that making comparisons between the time-stepping method and the moving average method could add values to the manuscript. Second, such comparisons show that the time-stepping method is an alternative way to account for the lagged responses of plants to climate variation. And to our knowledge, this has not been shown before in previous studies. Second, the Growing Season Index, which is based on the moving average method, is widely used in many studies and applications. Third, we do not try to use the moving average method as a benchmark. We use the Growing Season Index as a benchmark but there is a need to show that the time-stepping method can be similarly applied to the Growing Season Index approach such that the two methods can be compared directly.

Page 1 Line 1: This scheme cannot be used in LSMs because  $m=LAI/GPP$  is not a function of CO<sub>2</sub>.

Reply: First, our approach is different from others in terms of the leaf phenology sub-model. Although our approach does not account for CO<sub>2</sub> directly, it does account for CO<sub>2</sub> indirectly if the MOD17 approach is replaced with a canopy photosynthesis sub-model that account for CO<sub>2</sub>. If the canopy photosynthesis sub-model accounts for the impacts of CO<sub>2</sub>, the solved steady-state LAI would become a function of CO<sub>2</sub> and so does the modeled LAI. Whether our approach of leaf phenology modeling could account for the impacts of specific climate variables is largely dependent on the GPP models we used. In other words, for land surface models that include the canopy photosynthesis sub-models, the developed method can be embedded into these models as an alternative phenology model if replacing the MOD17 approach with the canopy photosynthesis model.

Second, for the current phenology models, there are nearly no direct considerations for CO<sub>2</sub>. The table 2 from Richardson et al. (2005) in Global Change Biology summarized the phenology models in 14 terrestrial biosphere model. Most of the them only account for the influence of temperature and none of them accounts for the influences of CO<sub>2</sub> directly. In the community land model (CLM), there are also no functions that directly links leaf phenology to CO<sub>2</sub>. In the phenology sub-model, CLM divided vegetation into 3 classes of the evergreen type, the seasonal-deciduous type, and the stress-deciduous type. None of these 3 classes include considerations of leaf area index as a function of CO<sub>2</sub> directly. It is only able to simulate leaf area index as a function of atmospheric CO<sub>2</sub> concentration indirectly.

**Table 2** Summary of models used in this analysis and their representation of phenology and seasonality of leaf area index (LAI). For models with ‘prognostic’ phenology, the seasonality of LAI is predicted based on climatic drivers; for those with ‘prescribed’ phenology, an average seasonal LAI cycle, as derived on a site-by-site basis from satellite (AVHRR) data, was used. Models with semi-prescribed and semi-prognostic phenology represent a hybrid of these approaches. GDD is growing degree days; T is temperature; C is carbon; PFT is plant functional type

Model name	Resolution	Leaf onset	Control on LAI	Leaf loss	Source
BEPS	Daily	Satellite	Satellite	Satellite	Ju <i>et al.</i> (2006)
Biome-BGC	Daily	GDD and radiation sum	Dynamic C allocation	Daylength and low temperature	Thornton <i>et al.</i> (2002)
Can-IBIS	Half-hourly	T threshold	GDD and dynamic C	Prescribed	El Maayar <i>et al.</i> (2002)
CN-CLASS	Half-hourly	C balance	C balance	Daylength and low temperature	Arain <i>et al.</i> (2006)
DLEM	Daily	$T_{7\text{-day}} > \text{threshold}$	GDD to PFT limit	Daylength and low temperature	Tian <i>et al.</i> (2010)
Ecosys	Hourly	Hours above T threshold	Dynamic C allocation	Hours below T threshold	Grant <i>et al.</i> (2009)
ED2	Half-hourly	Semi-prescribed	Dynamic C allocation	GDD and leaf turnover	Medvigy <i>et al.</i> (2009)
ISAM	Half-hourly	Prescribed	Prescribed	Prescribed	Jain & Yang (2005)
LoTEC	Half-hourly	GDD	GDD	T-dependent turnover	Hanson <i>et al.</i> (2004)
LPJ_wsl	Daily	GDD	GDD	Leaf longevity (prescribed)	Sitch <i>et al.</i> (2003)
ORCHIDEE	Half-hourly	GDD and chilling	Dynamic C allocation	Decreasing T and T threshold	Krinner <i>et al.</i> (2005)
SiB3	Half-hourly	Prescribed	Prescribed	Prescribed	Baker <i>et al.</i> (2008)
SiBCASA	10 min	Prescribed	Prescribed	Prescribed	Schaefer <i>et al.</i> (2008)
SSiB2	Half-hourly	Prescribed	Prescribed	Prescribed	Zhan <i>et al.</i> (2003)

Third, we agree that we need to be cautious about the statements for the model application in land surface models. We removed those inappropriate statements.

Page 1 Line 12: not sure what this means “which is used to track the suitability of environmental conditions for plant photosynthesis”.

Reply: we removed the texts to avoid confusion.

Page 1 Line 18 - 21: suggested changes to texts.

Reply: we agree with your suggestions. The revised texts now read as follows “The developed method is applied to deciduous broadleaf forests in eastern United States and is found to perform well for simulating canopy LAI and GPP at the site scale as evaluated using both flux tower and satellite data. The method also captures the spatiotemporal variation of vegetation LAI and phenology across eastern United States as compared with satellite observations. The developed time-stepping scheme provides a simplified and improved version of our previous modeling approach to simulate leaf phenology and can be potentially applied at regional to global scales in future studies.”

Page 1 Line 25 - 26: suggested changes to texts.

Reply: thanks for your suggestion. It now reads as follows “Terrestrial plants play a key role in regulating the exchange of energy and materials (e.g., radiation, heat and moisture, carbon, and trace gas) between the land surface and the atmosphere (Beer *et al.*, 2010; Zhu *et al.*, 2017). The canopy structure and characteristics govern solar radiation interception and absorption (Ni-Meister *et al.*, 2010; Yuan *et al.*, 2013).”

Page 2 Line 11 - 18: long sentence, break into two

Reply: We break the long sentence into two sentences. It now reads as follows:

“These methods include both the light use efficiency models (e.g., the Carnegie-Ames-Stanford Approach (CASA) model (Potter et al., 1993), the MOD17 algorithm (Running et al., 2004), the Vegetation Photosynthesis Model (VPM) (Xiao et al., 2004), the eddy covariance light use efficiency (EC-LUE) model (Yuan et al., 2010), and the two-leaf light use efficiency (TL-LUE) model (He et al., 2013)) and the process-based models (e.g., the boreal ecosystem productivity simulator (BEPS) model (Liu et al., 1997), the Breathing Earth System Simulator (BESS) model (Ryu et al., 2011), the Growing Production-Day (GPD) model (Xin, 2016), the revised Simple Biosphere (SiB2) model (Sellers et al., 1996b)). Despite different from each other on the representation of vegetation processes, these methods have been successfully used for applications from field to global scales.”

Page 2 Line 19 - 20: suggested changes to texts.

Reply: We revised the text as follows “While remotely sensed vegetation data perfectly complements the canopy process models, the ability to dynamically simulate vegetation LAI is fundamental to enhance our abilities on predicting terrestrial ecosystem processes.”

Page 3 Line 5 - 7: suggested changes to texts. Weak.

Reply: We agreed that the argument needed improvements and revised the texts as follows “While these studies have greatly benefitted the development of the leaf phenology models, evaluation on 14 land surface models in deciduous forests suggested that almost all models predicted start of the season earlier or end of the season later than observations and the model biases were typically 14 weeks or more. It is therefore necessary to improve the current phenology models.”

Page 3 Line 10 - 11: suggested changes to texts.

Reply: Based on your suggestion, we revised the texts as follows “The physiological processes of leaf phenology and canopy photosynthesis are interrelated. Plants absorb carbon dioxide to accumulate biomass through photosynthesis and then redistribute the photosynthetic gain to organs such as leaves, roots, and stems to optimize carbon gain.”

Page 3 Line 15 - 16: disconnected and vague.

Reply: We now revised the sentence as “In essence, new leaf phenology models may need to account for the processes of canopy photosynthesis more closely and explicitly than the current leaf phenology models.”

Page 3 Line 18: Arora & Boer (2005), Global Change Biology, used a similar approach.

Reply: we studied the paper of Arora & Boer (2005) in Global Change Biology and added it to our citations. It reads as follows “(Arora and Boer, 2005) developed a carbon-gain-based scheme that initiates leaf onset when environmental conditions are beneficial for the plant in carbon terms to produce new leaves and initiates leaf offset when environmental conditions are unfavourable and incur carbon losses for plants.”

Page 3 Line 27: suggested changes to texts.

Reply: we revised as you suggested.

Page 3 Line 32: suggested changes to texts.

Reply: We revised as you suggested. It now reads “The improved method allows modeling of LAI time series in addition to the timing of individual phenophases.”

Page 3 Line 35: suggested changes to texts.

Reply: We revised as you suggested. It now reads “There remain shortcomings to overcome for the broad applications of the GPD model. First, the simple moving average method, despite being widely used in many studies, is empirical and cannot be used within the framework of models that operate at incremental time steps. Second, the developed GPD model that includes many subtle vegetation processes, such as canopy radiative transfer, leaf stomatal conductance, leaf transpiration, leaf photosynthesis, and soil evaporation, is computationally intensive and requires various climate input data for regional to global applications.”

Page 4 Line 1: Justification for developing a simple model still not strong enough.

Reply: we agree with your argument. One main purpose of this study were to develop a method that can be used within a time-stepping model framework. Another reason is to improve the modeling accuracies. We therefore added the citation to Richardson’s publication in Global Change Biology in 2005 as follows “evaluation on 14 land surface models in deciduous forests suggested that almost all models predicted start of the season earlier or end of the season later than observations and the model biases were typically 14 weeks or more”. Note that the biases for our modeled results (the MBE values in Table 2) are less than 6 days.

Page 4 Line 1: This is not a limitation because almost all of these processes are already in land surface models.

Reply: thank you for your comments. We agree that this is not a limitation but the computational costs are expensive at fine spatial resolution for large-scale applications. For example, it is nearly impossible to perform 1 km resolution global simulations using the current land surface models even with supercomputing. We revised the related texts as “is computationally intensive for regional to global applications”.

Page 4 Line 8 - 10: suggested changes to texts.

Reply: We revised the texts based on your suggestions. It now reads as follows “this study choose to simulate leaf dynamics for the deciduous broadleaf forests across the eastern United States. If successful, such a method can be potentially used for future applications to other biomes.”

Page 4 Line 23: suggested changes to texts.

Reply: We revised the sentence and it now reads as follows “where  $LAI_s$  denotes the steady-state leaf area index;  $m$  denotes the constant ratio of steady state leaf area index to environmental capacity denoted by  $GPP_s$ , which is the steady-state gross primary production.”

Page 5 Line 1: Does not this precludes the MOD17 approach from being used in prognostic models for future scenarios. This is the justification you gave for developing your LAI model (Line 20, Page 2)

Reply: The MOD17 approach cannot be used in prognostic models for future scenarios because it requires the inputs of satellite-derived LAI data. Our developed method provides the simulated LAI input data for the MOD17 approach such that the MOD17 approach can then be used as prognostic models without observational satellite data.

Page 5 Line 4: suggested changes to texts

Reply: we revised as you suggested. It now reads “A brief description of the MOD17 algorithm is provided here and details can be found from the user guide of the MODIS GPP product (Running and Zhao, 2015).”

Page 5 Line 8: why is this steady state if PAR, T, & VPD are time varying?

Reply: For each day when PAR, T, and VPD are given, the vegetation GPP corresponding to the steady-state leaf area index can be modeled using the MOD17 algorithm. We revised the equations to be more clear.

Based on the MOD17 algorithm, vegetation GPP corresponding to the steady-state leaf area index can be modeled as follows:

$$GPP_s = PAR \times FPAR_s \times \varepsilon_{max} \times f(T) \times f(VPD) \quad (1)$$

where  $GPP_s$  denotes the gross primary production corresponding to the steady-state leaf area index; PAR denotes photosynthetically active radiation;  $FPAR_s$  denotes the fraction of photosynthetically active radiation corresponding to the steady-state leaf area index;  $\varepsilon_{max}$  denotes maximum light use efficiency; and  $f(T)$  and  $f(VPD)$  denote the scalar functions that account for the limitation of temperature and vapor pressure deficit, respectively, on canopy photosynthesis.

Page 5 Line 24: for given values of PAR, T, & VPD?

Reply: You are correct. We revised the sentence as follows “Given the environmental conditions (i.e., given daily values of photosynthetically active radiation, temperature, and vapor pressure deficit), Equations 1 and 2 together form simultaneous equations, meaning that there are two unknown variables (i.e., LAI and GPP at the steady state) and two different general equations.”

Page 5 Line 26: I am hoping later in the manuscript, I will be told what PAR, T, and VPD is used to calculate LAIs. PAR, T, and VPD vary throughout the year. Is LAIs calculated every time step?

Reply: LAIs is calculated every time step on the daily basis. We added a sentence to explain it as follows “For every day, daily photosynthetically active radiation, daily minimum air temperature, and daily vapor pressure deficit are used as the forcing data to calculate  $LAI_s$  for the corresponding day.”

Page 6 Line 1: suggested changes to texts

Reply: we revised as you suggested. It now reads as follows “To obtain the non-zero solution, the numerical approach starts with a guess value of  $\text{LAI}_s$  and then then iterates to obtain the approximated solution of  $\text{LAI}_s$  until converging.”

Page 6 Line 12: I am still unsure if  $\text{LAI}_s$  is calculated daily and is a function of time.

Reply:  $\text{LAI}_s$  is calculated every time step on the daily basis.  $\text{LAI}_s$  is not a function of time directly but is a function of PAR, T, and VPD. Because PAR, T, and VPD vary throughout the year, the calculated  $\text{LAI}_s$  vary from day to day.

We added sentences as follows “For every day, daily photosynthetically active radiation, daily minimum air temperature, and daily vapor pressure deficit are used as the forcing data to calculate  $\text{LAI}_s$  for the corresponding day. Because photosynthetically active radiation, minimum air temperature, and vapor pressure deficit vary throughout the year, the calculated  $\text{LAI}_s$  vary from day to day.”

Page 6 Line 13: suggested changes to the equation.

Reply: we now revised the equation as follows:

$$\text{LAI}(n + 1) = \frac{1}{n} \sum_{i=1}^n \text{LAI}_s(i) \quad (2)$$

where  $\text{LAI}(n + 1)$  denotes leaf area index at the  $n + 1$  day;  $n$  denotes the number of days;  $i$  denotes an index starting from 1 to  $n$ ; and  $\text{LAI}_s$  denotes the steady state leaf area index.

Page 6 Line 22: no it just reflects that photosynthesis does not instantaneously leads to new/big leaves

Reply: we revised as you suggested. It now reads as “ $k_l$  denotes a time constant that reflects that photosynthesis does not instantaneously lead to new or big leaves”.

Page 7 Line 12: what is the LAI on 1<sup>st</sup> day of spring? You have to use either an imaginary LAI to get GPP started or you need to push leaves out (like in the real world deciduous trees use non-structural carbohydrates from previous years). So I am still unsure how does GPP becomes on the very 1<sup>st</sup> day of spring.

Reply: Our modeling approach is different from your understanding. For the first day of spring, the modeled LAI is zero, the modeled fraction of photosynthetically active radiation (FPAR) is zero, and the modeled GPP is zero. When time moves forward, the modeled steady-state LAI (i.e.,  $\text{LAI}_s(n)$ ) is non-zero. Based on Equation 7, the modeled increases in LAI (i.e.,  $\frac{d\text{LAI}(n)}{dt}$ ) is positive. And then in Equation 8, the modeled LAI on the next time step is not zero anymore and starts to increase. To make it clear, we revised the texts and also added Equation 8 as follows.

$$\frac{d\text{LAI}(n)}{dt} = k_l[\text{LAI}_s(n) - \text{LAI}(n)] \quad (3)$$

$$\text{LAI}(n + 1) = \text{LAI}(n) + \frac{d\text{LAI}(n)}{dt} \quad (4)$$

where  $t$  denotes the time;  $k_l$  denotes a time constant that reflects that photosynthesis does not instantaneously lead to new or big leaves; and  $\text{LAI}(n)$  and  $\text{LAI}_s(n)$  denote the leaf area index and the steady state leaf area index at the  $n$  day, respectively.

Page 7 Line 25: suggested changes to texts

Reply: we revised the texts following your suggestion.

Page 9 Line 1:  $m = \text{LAI}/\text{GPP}$  is likely going to change as atmosphere CO<sub>2</sub> increases. This makes this approach somewhat not suitable for LSMs which need to be used for future scenarios.

Reply: It is difficult to know whether the relationship would change or not change as atmosphere CO<sub>2</sub> changes. We believe that there are no reasons to add some relationships or mechanisms that we do not know yet into models. One concern you have is that you thought our modeled LAI was not a function of CO<sub>2</sub>. First, although our approach does not account for CO<sub>2</sub> directly, it does account for CO<sub>2</sub> indirectly if the MOD17 approach is replaced with a canopy photosynthesis sub-model that account for CO<sub>2</sub>. If the canopy photosynthesis sub-model accounts for the impacts of CO<sub>2</sub>, the solved steady-state LAI would become a function of CO<sub>2</sub> and so does the modeled LAI. Whether our approach of leaf phenology modeling could account for the impacts of specific climate variables is largely dependent on the GPP models we used. Second, for the current phenology models, there are nearly no direct considerations for CO<sub>2</sub>. The table 2 from Richardson et al. (2005) in Global Change Biology summarized the phenology models in 14 terrestrial biosphere model. Most of the them only account for the influence of temperature and none of them accounts for the influences of CO<sub>2</sub> directly.

**Table 2** Summary of models used in this analysis and their representation of phenology and seasonality of leaf area index (LAI). For models with ‘prognostic’ phenology, the seasonality of LAI is predicted based on climatic drivers; for those with ‘prescribed’ phenology, an average seasonal LAI cycle, as derived on a site-by-site basis from satellite (AVHRR) data, was used. Models with semi-prescribed and semi-prognostic phenology represent a hybrid of these approaches. GDD is growing degree days; T is temperature; C is carbon; PFT is plant functional type

Model name	Resolution	Leaf onset	Control on LAI	Leaf loss	Source
BEPS	Daily	Satellite	Satellite	Satellite	Ju <i>et al.</i> (2006)
Biome-BGC	Daily	GDD and radiation sum	Dynamic C allocation	Daylength and low temperature	Thornton <i>et al.</i> (2002)
Can-IBIS	Half-hourly	T threshold	GDD and dynamic C	Prescribed	El Maayar <i>et al.</i> (2002)
CN-CLASS	Half-hourly	C balance	C balance	Daylength and low temperature	Arain <i>et al.</i> (2006)
DLEM	Daily	$T_{7\text{-day}} > \text{threshold}$	GDD to PFT limit	Daylength and low temperature	Tian <i>et al.</i> (2010)
Ecosys	Hourly	Hours above T threshold	Dynamic C allocation	Hours below T threshold	Grant <i>et al.</i> (2009)
ED2	Half-hourly	Semi-prescribed	Dynamic C allocation	GDD and leaf turnover	Medvige <i>et al.</i> (2009)
ISAM	Half-hourly	Prescribed	Prescribed	Prescribed	Jain & Yang (2005)
LoTEC	Half-hourly	GDD	GDD	T-dependent turnover	Hanson <i>et al.</i> (2004)
LPJ_wsl	Daily	GDD	GDD	Leaf longevity (prescribed)	Sitch <i>et al.</i> (2003)
ORCHIDEE	Half-hourly	GDD and chilling	Dynamic C allocation	Decreasing T and T threshold	Krinner <i>et al.</i> (2005)
SiB3	Half-hourly	Prescribed	Prescribed	Prescribed	Baker <i>et al.</i> (2008)
SiBCASA	10 min	Prescribed	Prescribed	Prescribed	Schaefer <i>et al.</i> (2008)
SSiB2	Half-hourly	Prescribed	Prescribed	Prescribed	Zhan <i>et al.</i> (2003)

Page 9 Line 29: suggested changes to texts

Reply: we revised the texts following your suggestion.

Page 14 Line 1: unclear what this means and suggested changes to texts.

Reply: we removed the words that are unclear and revised the texts according to your suggestion. It now reads “The SGPD-based models generally outperform the GSI-based models as the achieved correlation coefficients are higher and the RMSE are smaller. Both the GSI-SMA and GSI-TS models predict spring onsets earlier than observations by more than 30 days and predict autumn senescence later than observations by more than 20 days.”

Page 14 Line 11: what does this mean?

Reply: We revised the texts as “The modeled and measured GPP are compared in Figure 3 to understand the performance of GPP modeling.”

Page 15 Line 10-11: suggested changes to texts

Reply: we replaced the word “extents” with “distributions” as you suggested.

Page 17 Line 6: suggested changes to texts

Reply: we replaced the word “extents” with “distributions” as you suggested.

Page 20 Line 15: Fig 6 needs more discussion. The R<sup>2</sup> between MODIS and GLASS values for period 2000 – 2010 should also be shown and discussed to see how the two observation-based estimates compare to each other. This will be helpful before model can be compared to these

observation-based estimates.

Reply: we polished Fig 6 and added discussion to the results “The correlation coefficients between the GLASS data and the MODIS data for SOS, EOS, and GSL from 2001 to 2014 are 0.892, 0.412, and 0.288, respectively. There are only 14 years overlapping between these two different datasets and the correlations are insignificant for both the derived EOS and GSL.”

Page 21 Line 1: Something does not seem right in GLASS data. How come the interannual variability in GLASS reduce significantly after 2000 for SOS, EOS, and GSL? What is this range? Please clarify.

Reply: We do not think there are anything wrong. Please note that the GLASS data were produced using two different satellite data. From 1982 to 1999, the LAI product was generated at the resolution of 0.05 ° from the AVHRR reflectance. From 2000 to 2014, the LAI product was derived from MODIS land surface reflectance (MOD09A1) at the 1 km resolution. It is not surprising that the interannual variability in GLASS reduce significantly after 2000 for the derived SOS, EOS, and GSL.

We added explanations in the texts as follows “Note that the 8-day GLASS LAI product was generated at the 0.05 ° resolution using the AVHRR data from 1982 to 1999 and at the 1000 m resolution using the MODIS from 2000 to 2012. The significantly reduced interannual variability for SOS, EOS, and GSL after 2000 in the GLASS data suggests that the use of the AVHRR and MODIS data in the GLASS dataset could contribute uncertainties in the satellite-derived phenological metrics.”

In addition, to explain the range in the figure, we added one sentence to the figure caption as follows “The shaded areas denote the standard deviation of the corresponding phenophases across spaces.”

Page 23 Line 11: This sounds like a huge statement. But this gap is already been bridged in many land models.

Reply: We agree with you and revised the sentence as follows “Here we provide a simple time-stepping solution that allow for simulating canopy photosynthesis, leaf area index, and leaf phenology simultaneously.”

Page 24 Line 8: Of course, why do you need Fig 8 to tell this?

Reply: thank you for your comments. We revised the sentence as follows “Figure 8 suggests that LAI has stronger correlation with GPP than with temperature on the monthly basis.”

Page 24 Line 10: Is this subset of Fig 2a?

Reply: It is not a subset of Fig 2a. The analysis was performed at different time scales. We added a sentence to the figure caption “All data were averaged to the monthly time scale for analysis, making the point numbers different from the analysis at the weekly time scale in Figure 2..”

Page 25 Line 13: suggested changes to texts.

Reply: we revised the sentence based on your suggestions. It now reads “It implies that the LAI modeling in our developed method will likely benefit from improvements on the canopy photosynthesis model.”

Page 25 Line 14: Yes, of course, but not in LSMs for use in earth system models.

Reply: In this study we implement with the MOD17 approach to make predictions of LAI and GPP. If we replace the MOD17 approach with the canopy photosynthesis sub-model in the land surface model, the method could be applied to simulate both LAI and GPP as well. We agree that we have to be cautious with our statements before tests on applications in land surface models. We therefore removed the statements that are related to the applications for use in land surface models.

Page 25 Line 28: Yes but not using MOD17 GPP and what about your LAI is not a function of CO2.

Reply: we agree that it should be not used with the MOD17 approach and we revised the texts as “For land surface models that include the canopy photosynthesis sub-models, the developed method can be embedded into these models as an alternative phenology model if replacing the MOD17 approach with the canopy photosynthesis model.”

Another concern you have is that the modeled LAI is not a function of CO2. First, although our approach does not account for CO2 directly, it does account for CO2 indirectly if replacing the MOD17 approach with a canopy photosynthesis sub-model that accounts for CO2. If the canopy photosynthesis sub-model accounts for the impacts of CO2, the solved steady-state LAI would become a function of CO2 and so does the modeled LAI. Whether our approach of leaf phenology modeling could account for the impacts of specific climate variables is largely dependent on the GPP models we used. Second, for the current phenology models, there are nearly no direct considerations for CO2. The table 2 from Richardson et al. (2005) in Global Change Biology summarized the phenology models in 14 terrestrial biosphere model. Most of the them only account for the influence of temperature and none of them accounts for the influences of CO2 directly.

Page 25 Line 35: You are forgetting to mention explicitly that other than LAI, GPP is a function of a range of climate variables. So improvements in LAI do not lead to same amount of improvements in GPP.

Reply: We agree with your advice and added a sentence to the texts. It now reads “Because GPP is a function of a range of climate variables, improvements in modeling LAI do not lead to the same amount of improvements in the modeled GPP”.

# A simple time-stepping scheme to simulate leaf area index, phenology, and gross primary production across deciduous broadleaf forests in eastern United States

Qinchuan Xin<sup>1</sup>, Yongjiu Dai<sup>2</sup>, Xiaoping Liu<sup>1</sup>

5 <sup>1</sup>Guangdong Key Laboratory for Urbanization and Geo-simulation, Sun Yat-sen University, Guangzhou 510275, China

<sup>2</sup>School of Atmospheric Sciences, Sun Yat-sen University, Guangzhou 510275, China

Correspondence to: Qinchuan Xin (xinqinchuan@gmail.com); Yongjiu Dai (daiyj6@mail.sysu.edu.cn)

**Abstract.** Terrestrial plants play a key role in regulating the exchange of energy and materials between the land surface and the atmosphere. ~~Terrestrial biosphereRobust~~ models that simulate both leaf dynamics and canopy photosynthesis are required to understand the vegetation-climate interactions. This study proposes a simple time stepping scheme to simulate leaf area index (LAI), phenology, and gross primary production (GPP) when forced with climate variables. The method establishes a linear function between the steady-state LAI and the corresponding GPP, ~~which is used to track the suitability of environmental conditions for plant photosynthesis~~. The method applies the established function and the MOD17 algorithm to form simultaneous equations together, which can be solved numerically. To account for the time-lagged responses of plant growth to environmental conditions, a time stepping scheme is developed to simulate the LAI time series based on the solved steady-state LAI. The simulated LAI time series is then used to derive the timing of key phenophases and simulate canopy GPP with the MOD17 algorithm. The developed method is applied to deciduous broadleaf forests in eastern United States and ~~has-is~~ found to perform well ~~on-for~~ simulating canopy LAI and GPP at the site scale as evaluated using both flux tower and satellite data. The method ~~could~~ also capture~~s~~ the spatiotemporal variation of vegetation LAI and phenology across eastern United States as compared with satellite observations. The developed time-stepping scheme provides a simplified and improved version of our previous modeling approach to simulate leaf phenology and ~~forms a can be potentially applied at-basis for~~ regional to global applications-scales in future studies.

## 1 Introduction

Terrestrial plants play a key role in regulating the exchange of energy and materials (e.g., radiation, heat and moisture, 25 carbon, and trace gas ~~fluxes~~) between the land surface and the atmosphere (Beer et al., 2010; Zhu et al., 2017). The canopy structure~~s~~ and characteristics govern solar radiation interception and absorption (Ni-Meister et al., 2010; Yuan et al., 2013). Plants control water transpiration and photosynthetic carbon fixation through processes from transient changes in leaf stomatal conductance to seasonal variation in foliage dynamics (Eagleson, 2005). In turn, external environmental conditions, such as sunlight, temperature, and water and nutrient availability, selectively determine plant form and function (Bonan,

2008). Numerical ~~terrestrial biosphere~~ models that integrate multidisciplinary knowledge ~~of Earth sciences~~ allow to understand and predict the interactions between terrestrial ecosystems and the climate ~~under a changing global environment~~.

Developments on the terrestrial biosphere models essentially seek accurate solution to the simulation of energy and material 5 exchanging fluxes between ecosystems and the atmosphere. In terrestrial biosphere models, plant canopies are typically characterized using leaf area index (LAI; leaf area per unit ground area) because plant leaf is the basic organ that intercepts solar radiation for photosynthesis and transpiration (Li et al., 2018; Yuan et al., 2013). The exchanging fluxes of energy and materials over vegetation canopy can then be modeled as a function of environmental conditions (e.g., sunlight, soil moisture, temperature, and humidity) and vegetation LAI (Ding et al., 2014). The development of satellite remote sensing 10 technology offers large-scale observations for vegetation monitoring and a number of modeling approaches have been developed to quantify and simulate the land surface fluxes based on climate variables and satellite-derived LAI. These 15 methods, ~~including~~ include both the light use efficiency models (e.g., the Carnegie-Ames-Stanford Approach (CASA) model (Potter et al., 1993), the MOD17 algorithm (Running et al., 2004), the Vegetation Photosynthesis Model (VPM) (Xiao et al., 2004), the eddy covariance light use efficiency (EC-LUE) model (Yuan et al., 2010), and the two-leaf light use efficiency 20 (TL-LUE) model (He et al., 2013)) and the process-based models (e.g., the boreal ecosystem productivity simulator (BEPS) model (Liu et al., 1997), the Breathing Earth System Simulator (BESS) model (Ryu et al., 2011), the Growing Production-Day (GPD) model (Xin, 2016), the revised Simple Biosphere (SiB2) model (Sellers et al., 1996b)), ~~despite differing~~ Despite ~~different~~ from each other on the representation of vegetation processes, these methods have been successfully used for 25 applications from field to global scales. While ~~remotely sensing sensed vegetation~~ data ~~of vegetation activities~~ perfectly complements the canopy process models, ~~developing the sub-model ability that could to dynamically~~ simulate ~~the dynamics of~~ vegetation LAI is fundamental to enhance our abilities on predicting terrestrial ecosystem processes ~~under future scenarios~~.

Modeling vegetation leaf dynamics via climate variables requires in-depth understanding on plant phenological processes. 25 This modeling is still largely empirical to date and contributes considerable uncertainties to current terrestrial biosphere models (Richardson et al., 2012). One common method for simulating vegetation phenology is to predict the timing of key phenophases such as spring onset and autumn senescence in a growing season (Hufkens et al., 2018; Liu et al., 2018). For example, most phenology models originate from the Growing Degree Day (GDD) model, a method first proposed by De Réaumur dating back to 1735 (De Réaumur, 1735). The GDD model assumes that plant leaf onset begins when daily mean 30 temperatures accumulated from a fixed date reach a critical threshold. Studies have identified that various environmental factors other than temperature could affect plant phenology to certain degrees (Polgar and Primack, 2011), and therefore, efforts have been made to improve the GDD model by adding different influential factors, such as photoperiod, soil temperature, humidity, and soil moisture (Chuine et al., 1999; Hufkens et al., 2018; Liu et al., 2018; Melaas et al., 2013; Yang et al., 2012). Land surface models like the Community Land Model (Oleson et al., 2013) and the Biome-BGC model

(White et al., 2000) use a set of complicated and empirical equations to predict the timing of key phenophases across plant functional types. (Arora and Boer, 2005) developed a carbon-gain-based scheme that initiates leaf onset when environmental conditions are beneficial for the plant in carbon terms to produce new leaves and initiates leaf offset when environmental conditions are unfavourable and incur carbon losses for plants. Another method for vegetation phenology modeling is to 5 simulate the entire LAI time series over a growing season. For example, the DeNitrification DeComposition model uses an optimal seasonal growth curve of plant LAI and then calculates environmental stresses of water and nitrogen to limit daily carbon and nitrogen allocation to plant leaves (Yu et al., 2014). The Growing Season Index as proposed by (Jolly et al., 2005) is a widely used method that could simulate seasonal phenology curves using the climate variables of photoperiod, air 10 temperature, and vapor pressure deficit. While these studies have greatly benefitted the development of the leaf phenology models, evaluation on 14 land surface models in deciduous forests suggested that almost all models predicted start of the season earlier or end of the season later than observations and the model biases were typically 14 weeks or more. there is still a need-It is therefore necessary to improve the current phenology models.

The physiological processes of leaf phenology and canopy photosynthesis are interrelated. Plants absorb carbon dioxide to 15 accumulate biomasses through photosynthesis and then redistribute the photosynthetic gain to organs such as leaves, roots, and stems to optimize reproduction carbon gain. Given limited external resources, plants have evolved to effectively allocate photosynthate to organs in response to environment conditions so as to maximize photosynthetic carbon gain, the fundamental bioenergy for survival (Givnish, 1986). The strategy of biomass allocation among growth, maintenance, and reproduction in a continuously changing environment directly determines whether plants could persist under natural 20 competition pressures from both inter- and intra-species (Bonan, 2002). In essence, new leaf phenology models may need to account for the processes of canopy photosynthesis more closely and explicitly than the current leaf phenology models. there is a need to synthesize the analysis of both canopy photosynthesis and leaf phenology processes.

(Xin, 2016) proposed a parameterization scheme to simulate vegetation productivity and phenology simultaneously. The 25 method, named as the Growing Production Day (GPD) model, uses canopy gross primary production (GPP) instead of air temperature as an indicator that synthesizes various environmental factors on plant photosynthesis to track how the environment is suitable for vegetation growth. Analogous to the method that derives reference evapotranspiration, the developed method defines a hypothetic canopy with fixed LAI to model potential GPP under certain environment conditions. Similar to the GDD model, the GPD model predicts vegetation spring onset to occur when the accumulated reference GPP 30 reaches a critical threshold. The method has been successfully applied to the biomes of evergreen needleleaf forest, deciduous broadleaf forest, and grassland. To allow for predicting the entire LAI time series over a growing season, (Xin et al., 2018) further improved the GPD model by proposing a linear function between LAI and GPP at the steady state. The proposed function and the sophisticated-canopy GPP model (i.e., modeling GPP as a function of LAI and climate variables) together form a closed system of equations that includes both vegetation GPP and LAI. The improved GPD model uses the

numerical approach, a method that gives an initial value and then iterates to the convergence of the solution, to solve the closed system of equations and derives LAI in the steady state. The improved GPD model then applies the simple moving average method to the steady-state LAI to obtain the modeled LAI time series. The improved method ~~circumvents the need to empirically prescribe a fixed canopy and enables allows~~ modeling of LAI time series in addition to the timing of individual phenophases. There remain shortcomings to overcome for the broad applications of the GPD model. First, the simple moving average method, despite being widely used in many studies, is empirical and ~~does not match with the land surface models cannot be used within the framework of models~~ that ~~commonly~~ operate at incremental time steps. Second, the developed GPD model that includes many subtle vegetation processes, such as canopy radiative transfer, leaf stomatal conductance, leaf transpiration, leaf photosynthesis, and soil evaporation, ~~is computationally intensive and~~ requires various 10 climate input data and is computationally intensive for regional to global applications.

Aiming to solve the above-mentioned problems, the objectives of the study are to: 1) develop a time stepping scheme to simulate both leaf dynamics and vegetation productivity, and 2) simplify the GPD model to allow for long-term applications at a large scale. Given that the phenology modeling in deciduous broadleaf forest, a biome that have distinct seasonal 15 growing cycles, still has large uncertainties (Melaas et al., 2016), this study choose to simulate leaf dynamics for the deciduous broadleaf forests across the eastern United States, such that the developed method if successful, such a method can be potentially used could provide a potential basis for future applications to other biomes.

## 2 Methods and materials

### 2.1 Modeling steady-state leaf area index

20 One difficulty in vegetation phenology modeling is that the time scale associated with leaf allocation far exceeds that of many other vegetation processes. Unlike leaf photosynthesis that approaches equilibrium within a minute and stomatal functioning that reaches the steady state in minutes (Sellers et al., 1996a), leaf dynamics takes days or even months in response to weather variation (Zeng et al., 2013). (Xin et al., 2018) first put forward the concept of the steady-state leaf area index, i.e., canopy LAI when time approaches infinity while the environmental conditions remain unchanging. An alternative 25 biological explanation to the steady-state LAI is the maximum canopy LAI that an environment can sustain infinitely by its own photosynthetic activities. Supposing that the carrying capacity of canopy LAI is proportional to total canopy photosynthetic rate under a given environment, the steady-state LAI can be modeled as follows:

$$\text{LAI}_s = m\text{GPP}_s \quad (1)$$

where  $\text{LAI}_s$  denotes the steady-state leaf area index;  $m$  denotes the constant ratio of steady state leaf area index to environmental capacity; and denoted by  $\text{GPP}_s$  denotes, which is the steady-state gross primary production.

The above equation, despite having a simple form, provides a critical function that complements the canopy photosynthesis model. The only parameter  $m$  is dependent on plant functional type and can be quantified from field measurements as the average ratio of LAI to GPP at canopy closure (i.e., the time when both canopy LAI and GPP reach equilibrium). Studies have developed various canopy photosynthesis models, such as the light use efficiency models and the process-based models. Our previous studies (Xin, 2016; Xin et al., 2018) implemented a sophisticated canopy model that assembles the sub-models of canopy radiative transfer, leaf stomatal conductance, leaf transpiration, soil evaporation, and leaf photosynthesis. Although the method has been successfully applied to different biomes, the model structure is complicated for studies at the regional to global scales. To simulate canopy photosynthesis, this study implements the MOD17 algorithm, a big-leaf light use efficiency model that uses routine satellite products (Running et al., 2004). The use of the MOD17 algorithm could greatly simplify the modeling processes and reduce the required climate variables, thereby allowing for broad applications. A brief description ~~on~~of the MOD17 algorithm is provided here where\_and details can be found from the user guide of the MODIS GPP product (Running and Zhao, 2015).

Based on the MOD17 algorithm, vegetation GPP corresponding to the steady-state leaf area index can be modeled as follows:

$$GPP_s = PAR \times FPAR_s \times \varepsilon_{max} \times f(T) \times f(VPD) \quad (2)$$

where  $GPP_s$  denotes the steady state gross primary production corresponding to the steady-state leaf area index; PAR denotes photosynthetically active radiation;  $FPAR_s$  denotes the fraction of photosynthetically active radiation corresponding to the steady-state leaf area index;  $\varepsilon_{max}$  denotes maximum light use efficiency; and  $f(T)$  and  $f(VPD)$  denote the scalar functions that account for the limitation of temperature and vapor pressure deficit, respectively, on canopy photosynthesis.

20

The fraction of photosynthetically active radiation can be modeled as follows (Turner et al., 2006):

$$FPAR_s = 1 - \exp(-kLAI_s) \quad (3)$$

where  $k$  denotes the canopy light extinction coefficient and  $LAI_s$  denotes the steady-state leaf area index.

The environmental scalars can be modeled as follows:

$$f(T) = \max \left( \min \left( \frac{TMIN - TMIN_{min}}{TMIN_{max} - TMIN_{min}}, 1 \right), 0 \right) \quad (4)$$

$$f(VPD) = \max \left( \min \left( 1 - \frac{VPD - VPD_{min}}{VPD_{max} - VPD_{min}}, 1 \right), 0 \right) \quad (5)$$

25 where  $TMIN$  denotes daily minimum air temperature;  $TMIN_{min}$  and  $TMIN_{max}$  denote the lower and upper thresholds of daily minimum air temperature for vegetation photosynthetic activities, respectively;  $VPD$  denotes daily vapor pressure deficit;

and  $VPD_{min}$  and  $VPD_{max}$  denote the lower and upper thresholds of daily vapor pressure deficit for vegetation photosynthetic activities, respectively.

Given the environmental conditions (i.e., given daily values of photosynthetically active radiation, minimum air temperature, and vapor pressure deficit), Equations 1 and 2 together form simultaneous equations, meaning that there are two unknown variables (i.e., LAI and GPP at the steady state) and two different general equations. One may derive an analytic solution if both equations have simple forms. But because the dependence of GPP on LAI is non-linear, deriving the analytic solution is complicated and we could apply the numerical approach to obtain the solutions. Because  $LAI_s$  increases as a linear function of  $GPP_s$  in Equation 1 and  $GPP_s$  increases as a logarithmic function of  $LAI_s$  in Equation 2, the simultaneous equations have one and only one nonzero-non-zero solution of  $LAI_s$ . To obtain the nonzero-non-zero solution, the numerical approach starts with a guess value of  $LAI_s$  and then then iterates to obtain the approximated solution of  $LAI_s$  until converging. Note that the numerical approach is widely used in the land surface models. For example, as the stomatal resistance, the CO<sub>2</sub> partial pressure at the leaf surface, the internal leaf CO<sub>2</sub> partial pressure, and the leaf net photosynthesis are dependent on each other, the Community Land Model 4.5 uses the numerical approach to solve stomatal resistance and leaf photosynthesis iteratively until the internal leaf CO<sub>2</sub> partial pressure converges. For every day, daily photosynthetically active radiation, daily minimum air temperature, and daily vapor pressure deficit are used as the forcing data to calculate  $LAI_s$  for the corresponding day. Because photosynthetically active radiation, minimum air temperature, and vapor pressure deficit vary throughout the year, the calculated  $LAI_s$  vary from day to day.

## 2.2 Modeling leaf area index, phenology, and gross primary production

Because the physiological processes that plants allocate photosynthates to leaves do not respond instantaneously to climate variation, there is a need to simulate vegetation LAI as lagging behind the steady state. One method to account for the time lagging effect is to apply the simple moving average method to buffer abrupt changes from individual events in the time series. Our previous study applied the simple moving average method to model LAI as the unweighted mean of the previous  $LAI_s$  as follows (Xin et al., 2018):

$$LAI(n+1) = \frac{1}{n} \sum_{i=1}^n LAI_s(i) \quad (6)$$

where  $LAI(n+1)$  denotes leaf area index at the  $n+1$  day;  $n$  denotes the number of days;  $i$  denotes an index starting from 1 to  $n$ ; and  $LAI_s$  denotes the steady state leaf area index.

The simple moving average method, while showing useful in vegetation phenology modeling, is suitable for retrospective analysis rather than prediction, and importantly, it does not match with most land surface models that operate at incremental time steps. Analogous to the method used to simulate leaf stomatal conductance in response to environmental variation, this

study proposes a time stepping scheme to simulate LAI realistically as lagging behind the steady state by a simple restricted growth model (Sellers et al., 1996a) as follows:

$$\frac{d\text{LAI}(n)}{dt} = k_l [\text{LAI}_s(n) - \text{LAI}(n)] \quad (7)$$

$$\text{LAI}(n + 1) = \text{LAI}(n) + \frac{d\text{LAI}(n)}{dt} \quad (8)$$

where  $t$  denotes the time;  $k_l$  denotes a time constant that reflects ~~the responses of plant leaf allocation to climate variation that photosynthesis does not instantaneously lead to new or big leaves~~; and  $\text{LAI}(n)$  and  $\text{LAI}_s(n)$  denote the leaf area index and the steady state leaf area index at the n time step, respectively.

In the time stepping scheme, vegetation LAI does not change much during winter or summer as the current LAI is close to  $\text{LAI}_s$ , whereas vegetation LAI increases (or decreases) during spring (or autumn) as the current LAI is less (or greater) than  $\text{LAI}_s$ . For example, when the environment turns favorable for plant growth in spring,  $\text{LAI}_s$  exceeds LAI and  $d\text{LAI}/dt$  is

positive such that the modeled canopy LAI increases. Note that the method developed here essentially uses the canopy photosynthetic capacity (i.e., the steady-state gross primary production) instead of air temperature as a synthesized indicator to track the suitability of the environment to plant growth in time series, and therefore, the developed method is referred to as the Simplified Growing Production-Day (SGPD) model following our previous studies (Xin et al., 2018).

Given the modeled LAI time series, both vegetation phenology and canopy GPP can be easily modeled (Xin et al., 2018). Various approaches have already been developed to derive the timing of key phenophases such as spring onset and autumn senescence from seasonal LAI trajectories. This study models the phenological transition dates using a simple method that derives the first spring and last autumn dates at which LAI reaches 20%, 50%, and 80% of the seasonal amplitudes (Richardson et al., 2012). The selected relative amplitudes (20%, 50%, and 80%) are correspondent to different plant growth

stages over a growing season. Because the MOD17 algorithm only requires LAI, daily minimum temperature, daily vapor pressure deficit, and daily photosynthetically active radiation as model inputs, the canopy GPP is simply modeled by substituting the modeled LAI time series and the climate variables into the MOD17 algorithm. For the first day of spring when the modeled LAI is zero, the modeled fraction of photosynthetically active radiation is zero and the modeled GPP is zero. As times move forward, the modeled LAI increases and the modeled GPP increases as LAI increases but is still dependent on other climate variables such as solar radiation, temperature and vapor pressure deficit,

### 2.3 Comparative studies using Growing Season Index

The Growing Season Index (GSI), a widely used method in vegetation phenology modeling (Jolly et al., 2005), allows for modeling seasonal LAI time series rather than individual phenophases and is implemented to make direct comparisons with

the SGPD model. The GSI model performs comparably to or even outperforms other terrestrial biosphere models on predicting the timing of key phenophases for deciduous broadleaf forests (Melaas et al., 2013).

The instantaneous GSI is first derived based on the work of (Jolly et al., 2005) as follows:

$$iGSI = iTMIN \times iVPD \times iPhoto \quad (9)$$

5 where  $iGSI$  denotes instantaneous growing season index; and  $iTMIN$ ,  $iVPD$ , and  $iPhoto$  denote the instantaneous scalar functions that account for the constraints of daily minimum air temperature, vapor pressure deficit, and photoperiod, respectively, on vegetation growth.

10 The scalar functions for  $iTMIN$ ,  $iVPD$ , and  $iPhoto$  have the mathematic forms similar to Equations 4 and 5 and are ~~derived~~  
 obtained the same as defined in (Jolly et al., 2005) as follows:

$$iTMIN = \max \left( \min \left( \frac{TMIN - TMIN_{min}}{TMIN_{max} - TMIN_{min}}, 1 \right), 0 \right) \quad (10)$$

$$iVPD = \max \left( \min \left( 1 - \frac{VPD - VPD_{min}}{VPD_{max} - VPD_{min}}, 1 \right), 0 \right) \quad (11)$$

$$iPhoto = \max \left( \min \left( \frac{Photo - Photo_{min}}{Photo_{max} - Photo_{min}}, 1 \right), 0 \right) \quad (12)$$

where  $TMIN$  denotes daily minimum temperature;  $TMIN_{min}$  and  $TMIN_{max}$  denote the lower and upper thresholds of daily minimum air temperature for vegetation photosynthetic activities, respectively;  $VPD$  denotes daily vapor pressure deficit;  $VPD_{min}$  and  $VPD_{max}$  denote the lower and upper thresholds of daily vapor pressure deficit for vegetation photosynthetic activities, respectively;  $Photo$  denotes daily photoperiod; and  $Photo_{max}$  and  $Photo_{min}$  denote the lower and upper thresholds of daily photoperiod for vegetation photosynthetic activities, respectively.

15 LAI can be modeled as the simple moving average of the instantaneous GSI scaled using maximum LAI as follows:

$$GSI = \frac{1}{n_{day}} \sum_{i=0}^{n_{day}-1} iGSI \quad (13)$$

$$LAI = GSI \times LAI_{max} \quad (14)$$

where  $GSI$  denotes growing season index at the  $n$  day;  $n_{day}$  denotes the number of days;  $i$  denotes an index starting from 0 to the previous one day;  $iGSI$  denotes the instantaneous growing season index; and  $LAI_{max}$  denotes the maximum leaf area index at canopy closure.

It is noteworthy that the instantaneous GSI uses the product of the scalars of minimum temperature, vapor pressure deficit, and photoperiod as an indicator to track the potential canopy photosynthetic capacities on the daily basis. Both the GSI model and the SGPD model, despite having different forms, share the same modeling idea. To understand the differences between the simple moving average method and the time stepping method, the GSI model is also implemented with the 5 simple restricted growth model as follows:

$$\text{LAI}_s = \text{iGSI} \times \text{LAI}_{\max} \quad (15)$$

$$\frac{d\text{LAI}}{dt} = k_l(\text{LAI}_s - \text{LAI}) \quad (16)$$

where  $\text{iGSI}$  denotes the instantaneous growing season index;  $\text{LAI}_{\max}$  denotes the maximum leaf area index at canopy closure;  $k_l$  denotes a time constant that accounts for the lagged responses of plant leaf allocation to climate variation; and  $\text{LAI}$  and  $\text{LAI}_s$  denote the leaf area index and the steady state leaf area index, respectively.

10 With the modeled LAI time series, the phenological transition dates are then retrieved based on the seasonal amplitude ratio method, the same way as processing the LAI time series derived from the SGPD model. Vegetation GPP is modeled by substituting the modeled LAI time series into the MOD17 algorithm.

## 2.4 Model comparison and parameterization

This study compares four different modeling approaches, including the results simulated using both the SGPD model and the 15 simple moving average method (hereinafter referred to as SGPD-SMA), using both the SGPD model and the time stepping scheme (hereinafter referred to as SGPD-TS), using both the GSI model and the simple moving average method (hereinafter referred to as GSI-SMA), and using both the GSI model and the time stepping scheme (hereinafter referred to as GSI-TS). The commonly used metrics, including the Pearson correlation coefficient ( $R$ ), the coefficient of determination ( $R^2$ ), the root-mean-square error (RMSE), and the mean bias error (MBE), are derived for model assessment and comparison.

20 As the MOD17 algorithm is a well-parameterized model, this study applies the model parameters from literature directly. Following the user guide of the MODIS GPP product (Running and Zhao, 2015), key parameters in the MOD17 algorithm are set as  $\varepsilon_{\max} = 1.165 \text{ gC/MJ}$ ,  $\text{TMIN}_{\min} = -6.0 \text{ }^{\circ}\text{C}$ ,  $\text{TMIN}_{\max} = 9.94 \text{ }^{\circ}\text{C}$ ,  $\text{VPD}_{\min} = 0.65 \text{ kPa}$ , and  $\text{VPD}_{\max} = 1.65 \text{ kPa}$ . The light extinction coefficient of the canopy is 0.5. The parameter that defines the ratio of leaf area index to environmental 25 capacity is set as  $m = 0.58 \text{ m}^2(\text{leaf area})/\text{gC/day}$  as quantified using the average ratio of LAI to GPP at canopy closure using the flux tower data. The canopy maximum LAI is set as 5.80 based on the maximum 95th percentile of satellite-derived LAI across sites and years (Xin et al., 2018). The parameter  $n_{\text{day}}$  in the simple moving average method and the parameter  $k_l$  in the time stepping method control the response of plant leaf allocation to environmental variation. The parameter  $n_{\text{day}}$  is set as 21 days and the parameter  $k_l$  is calibrated as  $0.080 \text{ day}^{-1}$ .

## 2.5 Study materials and pre-processing

We evaluate our approach at the site scale using both the flux tower data and remote sensing data and at the regional scale using both the climate data and remote sensing data for deciduous broadleaf forests in eastern United States. For the site-scale studies, all the flux tower sites of deciduous broadleaf forests (Table 1) that are available in the AmeriFlux website

5 (<http://ameriflux.ornl.gov/>) were used for analysis. As the developed SGPD model is a simplified version of our previous modeling approach, the site-scale modeling studies only require daily incoming solar radiation, minimum air temperature, vapor pressure deficit, photoperiod, LAI, and GPP data. Daily incoming solar radiation, vapor pressure deficit, and GPP have already been provided in the Level 4 flux tower data, whereas daily minimum air temperature was processed from the half-hourly gap-filled Level 2 data and daily photoperiod as required by the GSI model was computed based on Equation 13  
10 17 as a function of geolocation and the day of year (Allen et al., 1998). As the MODIS LAI has been found to match field measurements well for deciduous broadleaf forests in eastern United States (Myneni et al., 2002), the 8-day 500 m MODIS LAI Version 6 products (MOD15A2H) that are downloaded from the Land Processes Distributed Active Archive Center  
15 (<https://lpdaac.usgs.gov/>) were used as the reference data. Canopy LAI at each site were extracted from MOD15A2H for the pixel that contains the corresponding site. The extracted 8-day MODIS LAI if identified as poor quality in MOD15A2H were replaced using the three-point median-value moving window technique. Spikes in the LAI time series were removed  
20 using the Hampel filter and then gap-filled using the autoregressive modeling approach (Akaike, 1969). The obtained 8-day LAI time series were further smoothed using the Savitzky-Golay filter and then linearly interpolated to generate daily time series. The phenological transition dates were retrieved from daily LAI time series using the method that derives the first spring and last autumn dates at which LAI reaches 20%, 50%, and 80% of the seasonal amplitudes, respectively (Richardson et al., 2012).

$$\text{Pho} = \frac{24}{\pi} \arccos \left( -\tan(\varphi) \tan \left( 0.409 \sin \left( \frac{2\pi}{365} \text{DOY} - 1.39 \right) \right) \right) \quad (17)$$

where Pho denotes daily photoperiod;  $\varphi$  denotes the latitude; and DOY denotes the day of year.

**Table 1: Site information for the studied flux towers of deciduous broadleaf forests.**

Site Code	Site Name	Lat (°N)	Lon (°W)	Elev (m)	Years	Reference
US-Bar	Bartlett Experimental Forest	44.0646	-71.2881	272	2004-2011	Jenkins et al. (2007)
US-ChR	Chestnut Ridge	35.9311	-84.3324	286	2006-2010	Hollinger et al. (2010)
US-Dk2	Duke Forest Hardwoods	35.9736	-79.1004	168	2007-2008	Oishi et al. (2008)
US-Ha1	Harvard Forest EMS Tower	42.5378	-72.1715	340	2000-2012	Urbanski et al. (2007)
US-MMS	Morgan Monroe State Forest	39.3231	-86.4131	275	2000-2014	Dragoni et al. (2011)
US-MOz	Missouri Ozark	38.7441	-92.2000	219	2005-2013	Gu et al. (2006)
US-Oho	Oak Openings	41.5545	-83.8438	230	2005-2011	Xie et al. (2014)
US-Slt	Silas Little Experimental Forest	39.9138	-74.5960	30	2005-2012	Clark et al. (2012)

US-UMB	Univ. of Mich. Biological Station	45.5598	-84.7138	234	2000-2012	Gough et al. (2013)
US-UMd	UMBS Disturbance	45.5625	-84.6975	239	2008-2012	Gough et al. (2013)
US-WBW	Walker Branch	35.9588	-84.2874	343	2000-2006	Miller et al. (2007)
US-WCr	Willow Creek	45.8060	-90.0798	515	2000-2013	Desai et al. (2008)


Our regional-scale studies used both the climate data and satellite remote sensing data from 1982 to 2016. The daily 1000 m Daymet Version 3 dataset (Thornton et al., 2012) was downloaded from the Oak Ridge National Laboratory Distributed Active Archive Center (<http://daymet.ornl.gov/>). The Daymet dataset provided daily incoming solar radiation, minimum 5 temperature, vapor pressure, and photoperiod data and we derived daily vapor pressure deficit as the difference between average saturated vapor pressure and vapor pressure. Two different satellite LAI products, including the Global Land Surface Satellite (GLASS) dataset (Xiao et al., 2014) spanning from 1982 to 2014 and the MODIS LAI dataset (Myneni et al., 2002) spanning from 2001 to 2016, were used for the regional studies. The 8-day GLASS LAI product was generated at the 0.05 ° resolution using the AVHRR data for the time period from 1982 to 1999 and at the 1000 m resolution using the 10 MODIS data for the time period from 2000 to 2012. The 8-day satellite LAI data across eastern United States were processed the same way as the processing of the site-scale data to obtain daily LAI time series. Because seasonal LAI amplitudes for each individual pixel could vary from year to year, the 2001-2010 average seasonal LAI amplitude were used as a baseline to derive the start of the season (SOS) and the end of the season (EOS) for each pixel for each year as the dates when seasonal LAI reaches 50% of the multi-year average seasonal LAI amplitude. The growing season length (GSL) was 15 derived as the difference between EOS and SOS. A 500 m MODIS-based land cover map was obtained from the USGS Land Cover Institute (<https://landcover.usgs.gov/>). The land cover map was generated by choosing the land cover classification with the highest overall confidence for each pixel in 10-year (2001-2010) Collection 5.1 MODIS land cover type (MCD12Q1) data (Broxton et al., 2014). The 500 m land cover map was resampled to 1000 m resolution using the majority resampling approach and was reprojected to the Lambert Conformal Conic projection to mask areas that are not covered by 20 deciduous broadleaf forests.

### 3 Results

#### 3.1 Site-scale modeling

Figure 1 shows an example for the simulated time series of LAI and GPP using data acquired at the US-UMB in 2004. The LAI time series simulated using both the SGPD-SMA and SGPD-TS methods are consistent with that obtained from 25 MODIS. The LAI simulated using both the GSI-SMA and GSI-TS methods could also capture the observed seasonal variation of LAI but the modeled phenophases obviously have a leading phase in spring and a lagging phase in autumn as compared with observations. For both the SGPD model and the GSI model, the results derived using the time stepping method are consistent with those derived using the simple moving average method, indicating that the time stepping method

is an effective way to reflect the lagging responses of plant leaf allocation to environmental conditions. By substituting the time series of LAI derived from different modeling approaches into the MOD17 algorithm, all the simulated GPP time series could match the flux tower measurements. Daily fluctuation in the observed GPP time series is largely due to variation in solar radiation from day to day. The GPP modeled using both the GSI-SMA and GSI-TS methods have slight overestimates 5 in the phenological transition periods like spring and autumn and match well with the flux tower observations in summer and winter.



10 **Figure 1: The modeled and measured daily time series of a) leaf area index and b) gross primary production are shown for the flux tower site of US-UMB in 2004. The reference LAI time series in Figure 1a are derived from the MODIS data and the reference GPP time series in Figure 1b are obtained from the flux tower measurements.**

Figure 2 shows the regression analysis between the modeled and satellite-derived LAI. Overall, the SGPD model outperforms the GSI model on modeling LAI. When evaluated against the MODIS LAI data, the SGPD-SMA and SGPD-TS models achieved the  $R^2$  of 0.887 and 0.890, respectively, and the RMSE of 0.804 and 0.778  $m^2/m^2$ , respectively, whereas the GSI-SMA and GSI-TS models achieved the  $R^2$  of 0.746 and 0.759, respectively, and the RMSE of 1.356 and 1.303  $m^2/m^2$ , respectively. Both the GSI-SMA and GSI-TS models simulate LAI reasonably in summer and winter but overestimate LAI in spring and autumn, and therefore, the strong correlations between the GSI-modeled and MODIS-derived LAI are largely due to the underlying seasonality of deciduous broadleaf forests. It is noteworthy that the time stepping method and the simple moving average method, despite having different mathematical expressions, generate nearly the same simulation results. The  $R^2$  values between the SGPD-TS model and the SGPD-SMA model and between the GSI-TS model and the GSI-SMA model are 0.989 and 0.994, respectively, and the regression lines are close to the lines of equity, indicating that the time stepping method is an alternative representation for the simple moving average method.




Figure 2: Scatter plots are shown for the comparisons a) between the SGPD-SMA LAI and the MODIS LAI, b) between the SGPD-TS LAI and the MODIS LAI, c) between the SGPD-TS LAI and the SGPD-SMA LAI, d) between the GSI-SMA LAI and the MODIS LAI, e) between the GSI-TS LAI and the MODIS LAI, and f) between the GSI-TS LAI and the GSI-SMA LAI on a

weekly basis. All available site-year flux tower data were included in the analysis. The solid lines denote the 1:1 lines and the dashed lines denote the regression lines.

Table 2 lists the statistical metrics that illustrate the model performance on predicting the timing of different phenophases.


5 As evaluated against satellite observations, the SGPD-SMA model could well retrieve the spring onset dates when LAI reaches 50% seasonal amplitude and the obtained correlation coefficient is 0.718 with RMSE of 13.04 days. The SGPD-TS model performs comparable to the SGPD-SMA model and the resulted correlation coefficients are all significant except for the dates that autumn LAI reaches 80% seasonal amplitudes. The SGPD-based models generally outperform the GSI-based models as the achieved correlation coefficients are higher and the RMSE are smaller ~~for more than 10 days~~. Both the GSI-SMA and GSI-TS models predict spring onsets earlier than observations ~~for by~~ more than 30 days and predict autumn senescence later than observations ~~for by~~ more than 20 days. By comparison, the SGPD-TS model predicts the dates that spring and autumn LAI reaches 50% seasonal amplitudes well with the MBE of only -2.56 and -2.86 days, respectively.

10 **Table 2: The performance of the modeled timings of phenophases as evaluated against satellite observations. The timings of phenophases were derived based on dates at which the leaf area index reaches 20%, 50%, 80% of seasonal amplitude. Positive mean bias error (MBE) indicates that the modeled spring onsets are earlier than the observed ones and negative MBE indicates the opposite.**

phenophases	SGPD-SMA			SGPD-TS			GSI-SMA			GSI-TS		
	R	RMSE	MBE									
		(days)	(days)									
Spring LAI 20%	0.790***	16.17	-10.85	0.824***	13.37	-8.34	0.763***	40.38	-38.30	0.770***	39.62	-37.58
Spring LAI 50%	0.718***	13.04	-1.97	0.691***	13.68	-2.56	0.653***	38.47	-34.92	0.657***	38.22	-34.63
Spring LAI 80%	0.432***	20.91	12.63	0.409***	21.19	12.41	0.560***	32.86	-28.00	0.565***	28.54	-23.55
Autumn LAI 80%	0.220	31.80	-25.56	0.164	27.90	-20.64	0.021	35.38	32.42	-0.004	35.23	32.27
Autumn LAI 50%	0.686***	9.80	-5.42	0.625***	9.48	-2.86	0.621***	24.20	23.07	0.616***	24.63	23.51
Autumn LAI 20%	0.703***	8.87	2.15	0.676***	10.91	6.37	0.689***	19.64	18.48	0.713***	22.93	22.00

20 The modeled and measured GPP are compared in Figure 3 to ~~address the key question that whether the simulated LAI could be applied to model canopy GPPunderstand the performance of GPP modeling~~. Compared with the flux tower measurements, the results modeled using the SGPD-SMA, SGPD-TS, GSI-SMA, and GSI-TS LAI could achieve the  $R^2$  values of 0.768, 0.773, 0.722, and 0.719, respectively, and the RMSE values of 2.273, 2.239, 2.577, 2.535 gC/m<sup>2</sup>day, respectively. The modeled results using the GSI-based LAI have higher errors, in terms of both RMSE and MBE, than those using the SGPD-based LAI. The accuracies of the modeled GPP using the SGPD-based LAI are only slightly lower than to that using the MODIS-based LAI directly. ~~Because GPP is also a function of a range of climate variables, improvements in modeling LAI do not lead to the same amount of improvements in the modeled GPP~~. The modeling results obtained based on the simple moving average method are nearly the same as those obtained based on the time stepping method. Given the

high degrees of consistency between the simple moving average method and the time stepping method on modeling LAI, phenology, and GPP, only the results obtained using the time stepping method are shown and discussed in the regional studies as presented in the following section.



5

**Figure 3: Scatter plots are shown for the comparisons a) between the GPP modeled using SGPD-SMA LAI and the flux tower GPP, b) between the GPP modeled using SGPD-TS LAI and the flux tower GPP, c) between the GPP modeled using MODIS LAI and the flux tower GPP, d) between the GPP modeled using GSI-SMA LAI and the flux tower GPP, and e) between the GPP modeled using GSI-TS LAI and the flux tower GPP on a weekly basis. All available site-year flux tower data were included in the analysis. All the modeled GPP were derived using the MOD17 algorithm. The solid lines denote the 1:1 lines and the dashed lines denote the regression lines.**

10

### 3.2 Regional-scale modeling

Figure 4 shows the spatial extents-distributions of the 10-year (2001-2010) mean LAI and associated errors as derived from remote sensing data and model simulations. The SGPD-TS method could well capture the spatial pattern of the satellite-

15

derived LAI, including the decreasing gradients from south to north and the decreases in mountain areas (Figure 4a and 4b). The 10-year mean LAI derived from the GSI-TS method (Figure 4c) also show a decreasing trend from south to north but the modeled LAI is much larger than the MODIS LAI. Because the GSI-TS method defines the maximum leaf area index for the growing season, the overestimation on the modeled 10-year mean LAI is primarily due to model overestimates in the 5 spring and autumn phenological transitions. As compared with the MODIS observations, RMSE and MBE obtained by the SGPD-TS method are much smaller than and distribute more evenly than those obtained by the GSI-TS method. RMSE for the GSI-TS LAI exhibit a decreasing north-south gradient, implying that the model accuracies are lower in southern areas lower than in northern areas. MBE for the GSI-TS model are greater than  $0.5 \text{ m}^2 \text{ m}^{-2}$  for most areas. When comparing SGPD-TS LAI with MODIS LAI, RMSE are less than  $0.5 \text{ m}^2 \text{ m}^{-2}$  and MBE are minor across the study region. The amplitudes of the 10 error metrics in the regional-scale studies are consistent with those in the site-scale studies. Note that some studies applied the multi-year mean LAI as derived from the remote sensing data to simulate the land surface processes, the results obtained here indicate that the SGPD-TS method can be used alternatively to provide multi-year mean LAI time series via climate variables for land surface studies.




Figure 4: The spatial extents-distributions are shown for a) the 2001-2010 mean MODIS LAI, b) the 2001-2010 mean SGPD-TS LAI, c) the 2001-2010 mean GSI-TS LAI, d) RMSE between SGPD-TS LAI and MODIS LAI, e) RMSE between GSI-TS LAI and MODIS LAI, f) MBE between SGPD-TS LAI and MODIS LAI, and g) MBE between GSI-TS LAI and MODIS LAI across eastern United States. The units for both RMSE and MBE are  $\text{m}^2$  (leaf area) per  $\text{m}^2$  (ground area).

5

The spatial extents-distributions for the 10-year mean phenological metrics including the start of the season (SOS), the end of the season (EOS), and the growing season length (GSL) are shown in Figure 5. The SGPD-TS method predicts lower SOS (i.e., earlier spring onset), higher EOS (i.e., later autumn senescence), and longer GSL in southern areas than in northern

areas. The spatial distributions of all phenological metrics derived using SGPD-TS LAI agree well with those derived using MODIS LAI. From the statistical analysis as shown in the subplots, the phenological metrics derived from the SGPD-TS method could achieve the correlation coefficient values of 0.879, 0.552, and 0.844, the RMSE values of 8.13, 7.54, and 13.73 days, and the MBE values of 0.71, -2.82, and -3.54 days, for SOS, EOS, and GSL, respectively, as compared to those derived from the MODIS data. Although the spatial distributions of the phenological metrics derived from the GSI-TS method match those derived from the satellite observations, the modeled results have considerable biases, where the RMSE values are 38.05, 14.37, and 51.58 days, and the MBE values are -36.33, 12.91, and 49.23 days, for SOS, EOS, and GSL, respectively. Consistent with the site-scale studies, the GSI-TS method predicts spring onset much earlier and autumn senescence later than the satellite-derived data, resulting in large overestimation of the growing season length.




Figure 5: The spatial extents\_distributions are shown for a) the start of the season (SOS) derived from MODIS LAI, b) SOS derived from SGPD-TS LAI, c) SOS derived from GSI-TS LAI, d) the end of the season (EOS) derived from MODIS LAI, e) EOS derived from SGPD-TS LAI, f) EOS derived from GSI-TS LAI, g) the growing season length (GSL) derived from MODIS LAI, h) GSL derived from SGPD-TS LAI, and i) GSL derived from GSI-TS LAI using the 10-year (2001-2010) mean data across eastern United States. The embedded subplots show the comparisons between modeled and MODIS-derived phenological metrics for SOS, EOS, and GSL, respectively.

Figure 6 displays the multi-year phenology anomalies that are spatially averaged for deciduous broadleaf forest across eastern United States. The use of phenology anomalies relative to the 2001-2010 average instead of absolute values makes the results directly comparable. The SGPD-TS method could capture the interannual variation of vegetation phenology retrieved from the remote sensing data. When comparing the SGPD-TS method with the MODIS (2001-2016) data, the 5 correlation coefficients are 0.896 ( $p<0.001$ ), 0.650 ( $p=0.006$ ), and 0.817 ( $p<0.001$ ), for SOS, EOS, and GSL, respectively. When comparing the SGPD-TS method with the GLASS (1982-2014) data, as derived from and the correlation coefficients are 0.554 ( $p=0.001$ ), 0.717 ( $p<0.001$ ), 0.637 ( $p<0.001$ ), for SOS, EOS, and GSL, respectively. The SGPD-TS method outperforms the GSI-TS method on capturing the long-term trends of vegetation phenophases, as the correlation coefficients obtained using the GSI-TS method are lower and sometimes insignificant. Yearly fluctuation in EOS derived using the GSI-10 TS method is smaller than those derived from both the SGPD-TS method and the satellite data. The correlation coefficients between the GLASS data and the MODIS data for SOS, EOS, and GSL from 2001 to 2014 are 0.892, 0.412, and 0.288, respectively. There are only 14 years overlapping between these two different datasets and the correlations are insignificant for both the derived EOS and GSL. The SOS and EOS derived from the GLASS data have much larger variation in 1982-2000 than in 2001-2010. Note that the 8-day GLASS LAI product was generated at the 0.05 °resolution using the AVHRR data from 1982 to 1999 and at the 1000 m resolution using the MODIS from 2000 to 2012. The significantly reduced interannual variability for SOS, EOS, and GSL after 2000 in the GLASS data, suggesting suggests that the use of the AVHRR and MODIS data in the GLASS dataset could contribute uncertainties in the satellite-derived phenological metrics. Both Figure 5 and 6 indicate that the SGPD-TS method is reliable on capturing the spatiotemporal patterns of regional vegetation phenophases.

20





Figure 6: The spatially-averaged phenology anomalies relative to the 2001-2010 average are shown for a) the start of the season (SOS), b) the end of the season (EOS), and c) the growing season length (GSL). SOS and EOS are derived as the date that LAI first and last reaches 50% of the seasonal amplitudes and GSL is derived as the difference between EOS and SOS. The shaded areas denote the standard deviation of the corresponding phenophases across spaces.

Figure 7 compares the simulated GPP using the MOD17 algorithm and LAI derived from different approaches. The 10-year average annual GPP obtained using SGPD-TS LAI has a similar spatial pattern with that obtained using MODIS LAI and has lower values than that obtained using GSI-TS LAI. Taking the GPP simulated using MODIS LAI as reference, the results simulated using SGPD-TS LAI achieve the correlation coefficient of 0.898 with RMSE of 78.78 gC/m<sup>2</sup>/year and MBE of 12.22 gC/m<sup>2</sup>/year, whereas the results simulated using GSI-TS LAI achieve the correlation coefficient of 0.898 with RMSE of 167.26 gC/m<sup>2</sup>/year and MBE of 144.41 gC/m<sup>2</sup>/year. Although the obtained correlation coefficients are close, the SGPD-TS method results in the regression lines closer to the 1:1 lines with smaller bias errors than the GSI-TS method. The zonally average profiles of the 2001-2010 average annual GPP as shown in Figure 7d suggest that the results obtained from the SGPD-TS method are close to those obtained using MODIS LAI, whereas the results obtained from the GSI-TS method have positive biases of approximately 120 - 180 gC/m<sup>2</sup>/year (roughly 10 - 15%) across latitudes. Note that the MOD17 algorithm has positive MBE of 0.247 gC/m<sup>2</sup>/day and 0.571 gC/m<sup>2</sup>/day when using SGPD-TS LAI and GSI-TS LAI, respectively, as model input data in the site-scale study. The differences in MBE between the two modeling methods are 0.324 gC/m<sup>2</sup>/day (or 118.26 gC/m<sup>2</sup>/year in equivalence) for the site-scale studies, which are consistent with the regional-scale studies.

15




5 | **Figure 7: Comparisons are shown for a) the spatial extent distribution of annual GPP modeled using MODIS LAI, b) the spatial extent distribution of annual GPP modeled using SGPD-TS LAI, c) the spatial extent distribution of annual GPP modeled using GSI-TS LAI, d) the zonally averaged profiles of annual gross primary production modeled using LAI derived from different approaches, e) the regression between GPP modeled using SGPD-TS LAI and MODIS LAI, and f) the regression between GPP modeled using GSI-TS LAI and MODIS LAI. The simulated daily GPP were first summed for each individual year and were then averaged across years to derive the 2001-2010 average annual GPP as shown in Figure 7a, 7b, and 7c. The shaded areas in Figure 7d mark the range of the standard deviation. All pixels of deciduous broadleaf forest across eastern United States are included in analysis in Figure 7e and 7f.**

#### 4 Discussion

10 | Here we provide a simple time-stepping solution that ~~bridges the canopy photosynthesis model and the leaf dynamics model~~  
~~allow for simulating canopy photosynthesis, leaf area index, and leaf phenology simultaneously~~. The developed method first  
proposes a linear function between the canopy photosynthetic capacity and the steady state LAI so as to complement the  
canopy photosynthesis model and then applies a simple restricted growth model to account for the lagged responses of plant  
leaf allocation to natural environment. In essence, the developed method, although having a simple form, has synthesized the  
15 | impacts of various climate factors on leaf dynamics because any climate variable that influences vegetation photosynthesis  
would affect the process of plant leaf allocation in the models as well. Consistent with field observations, the simulated LAI  
increases as the environmental conditions turn favorable for photosynthetic activities such as increases in photoperiod and  
temperature.

20 | Figure 8 further illustrate the relationship between mean LAI and different variables on a monthly basis. All data were  
averaged to the monthly time scale such that canopy LAI can be considered as nearly the steady state. On the monthly basis,  
mean LAI has a strong near-linear relationship with mean GPP ( $R \geq 0.888$ ) and the slope for the regression without intercept  
is 0.580, the same as we used in the model simulation. On the monthly basis, mean LAI is strongly correlated with mean  
temperature ( $R \geq 0.799$ ), indicating that temperature is the dominate factor that determines vegetation phenology. Factors like  
25 | vapor pressure deficit and photoperiod also have positive relationships with mean LAI on the monthly basis. Figure 8  
suggests that ~~LAI has stronger correlation with GPP than with temperature on the monthly basis~~  
~~the processes of leaf phenology and photosynthetic phenology for deciduous broadleaf forest are closely related~~. Our modeling approach that  
links canopy GPP with LAI reflects the empirical positive relationship found in Figure 8a.



5 | **Figure 8: Scatter plots are shown for the relationship a) between mean leaf area index and mean gross primary production, b) between mean leaf area index and mean temperature, c) between mean leaf area index and mean vapor pressure deficit, and d) mean leaf area index and mean photoperiod on a monthly basis. All available site-year flux tower data were included in the analysis. All data were averaged to the monthly time scale for analysis, making the point numbers different from the analysis at the weekly time scale in Figure 2. The dashed lines denote the regression lines. Figure 8a uses the regression without intercept.**

The performance of our developed method is largely dependent on the canopy photosynthesis model used. In our previous studies, we developed a process-based canopy photosynthesis model that synthesizes sub-models such as canopy radiative transfer, leaf transpiration, leaf stomatal conductance, leaf photosynthesis, and soil evaporation and applied it for modeling the LAI time series. When applying the simple moving average method, implementing the process-based model in Xin et al. 10 (2018) achieved higher accuracies than implementing the MOD17 algorithm on modeling canopy GPP and LAI as reflected by higher  $R^2$  and lower errors. The MOD17 algorithm only assumes the monotonic relationship between air temperature and

photosynthesis and between vapor pressure deficit and photosynthesis. It also does not account for the impacts of CO<sub>2</sub> on photosynthesis. The use of the MOD17 algorithm in this study thus has limitations in the model structure. It implies that the LAI modeling in our developed method will likely benefit from improvements on the canopy photosynthesis model. This study chooses the MOD17 algorithm instead of the sophisticated process-based model because the MOD17 algorithm is well parameterized across biomes and requires quite limited model inputs of climate variables. Successful implementation with the MOD17 algorithm allows for extending the developed method to applications across biomes at regional to global scales.

Land surface models that predict vegetation GPP require either satellite-derived LAI input data or the phenology sub-model. The main idea for this study is to improve the phenology modeling by providing time series of LAI simulated using climate variables, and hence enables to simulate GPP forced only by climate variables. Because we implement the MOD17 algorithm instead of the sophisticated process-based model for the purpose of simplicity, one should not expect that GPP simulated based on the model-simulated LAI could be more accurate than GPP simulated based on the satellite-derived LAI.

The time stepping scheme developed here is also an improvement over the simple moving average method as used in our previous studies. The results obtained using the time stepping method are consistent with the simple moving average method at the site scale and show to be reasonable at the regional scale. Compared to the simple moving average method, the time stepping method could fit seamlessly into the land surface be used in models that operate at incremental time steps such as the Community Land Model and the Common Land Model (Dai et al., 2003). Because the state-of-the-art For land surface models all that include the canopy photosynthesis sub-models, the developed method can then be easily embedded into these land surface models as an alternative phenology model if replacing the MOD17 approach with the canopy photosynthesis sub-model. Compared to the simple light use efficiency model like the MOD17 algorithm, implementation of the developed time-stepping scheme in the land surface models relies on supercomputing for global applications. To better understand the performance of the developed method, one study is now undertaken to implement the developed method with the Common Land Model for simulating multi-decadal LAI and GPP for global biomes forced only by climate variables.

25

Applying the developed method to other biomes and other regions still has issues to be solved appropriately. The time stepping method uses the parameter  $k_1$  to account for the time lags of leaf allocation in response to environmental changes. For the deciduous broadleaf forests, a biome with strong seasonality, the developed scheme achieved reasonable results with appropriate parameterization. Short vegetation like grasslands tends to respond much quickly to abrupt environment changes like precipitation and tropical ecosystems have strong resilience to short-term environmental variation (Levine et al., 2016; Shen et al., 2011). Another issue is to find the appropriate values of  $m$  for different biomes. One way to determine the values of  $m$  is to find the regression slope between leaf area index and gross primary production on a monthly basis. Model parameterization however still requires broad tests. These understandings from the observational studies imply that biomes have varied response speeds to the environment and proper model calibration and assessment are required for the developed

method. Using the observation data from remote sensing alone is inadequate for model development as satellite-derived LAI could have large uncertainties for some specific biomes other than deciduous broadleaf forests. Fortunately, global flux tower network and regional phenology observation networks are now established and offer abundant data for comprehensive model assessment.

## 5 5 Conclusions

~~Terrestrial biosphere-Numerical~~ models provide a basic tool for understanding the interactions between the land surface and the atmosphere. To provide a complete solution to the simulation of plant leaf dynamics and canopy photosynthesis, this study establishes a linear relationship between the steady state leaf area index and the corresponding canopy photosynthetic capacity. The proposed leaf allocation function complements the canopy photosynthesis model of the MOD17 algorithm to form simultaneous equations that can be solved using the numerical approach. To account for the time lagging of plant leaf allocation in response to climate variation, a time stepping scheme based on a simple restricted growth model is applied to the solved steady state leaf area index to obtain time series of leaf area index. The developed method could perform reasonably well on simulating leaf area index, phenology, and gross primary production for deciduous broadleaf forests across eastern United States over years as found in both the site-scale and regional-scale modeling studies. Compared to the simple moving average method, the time stepping scheme developed here is consistent with and can be ~~easily-potentially~~ embedded into ~~the state-of-the-art land surface~~ models that ~~typically~~ operate at incremental time steps. The developed method allows for simulating leaf area index and gross primary production simultaneously and provides a much simplified and improved version of our previous model as a basis for global applications in future studies.

### Author contributions

20 Qinchuan Xin designed the experiments and performed the simulations. All authors interpreted the results. Qinchuan Xin wrote the manuscript with contributions from all coauthors. The authors declare that they have no conflict of interest.

### Acknowledgments

We thank the researchers and investigators who are involved in collecting and sharing the AmeriFlux dataset. This research is supported by National Key R&D Program of China (grant nos. 2017YFA0604302 and 2017YFA0604402) and National 25 Natural Science Foundation of China (grant no. 41875122). We also thank anonymous reviewers for their constructive comments.

## References

Akaike, H.: Fitting autoregressive models for prediction, *Annals of the institute of Statistical Mathematics*, 21, 243-247, 1969.

5 Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56, FAO, Rome, 300, 6541, 1998.

Arora, V. K. and Boer, G. J.: A parameterization of leaf phenology for the terrestrial ecosystem component of climate models, *Global Change Biology*, 11, 39-59, 2005.

10 Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Roedenbeck, C., Arain, M. A., Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M., Luyssaert, S., Margolis, H., Oleson, K. W., Roupsard, O., Veenendaal, E., Viovy, N., Williams, C., Woodward, F. I., and Papale, D.: Terrestrial Gross Carbon Dioxide Uptake: Global Distribution and Covariation with Climate, *Science*, 329, 834-838, 2010.

Bonan, G. B.: *Ecological climatology: concepts and applications*, Cambridge University Press, 2002.

Bonan, G. B.: Forests and climate change: forcings, feedbacks, and the climate benefits of forests, *science*, 320, 1444-1449, 2008.

15 Broxton, P. D., Zeng, X., Sulla-Menashe, D., and Troch, P. A.: A global land cover climatology using MODIS data, *Journal of Applied Meteorology and Climatology*, 53, 1593-1605, 2014.

Chuine, I., Cour, P., and Rousseau, D. D.: Selecting models to predict the timing of flowering of temperate trees: implications for tree phenology modelling, *Plant Cell and Environment*, 22, 1-13, 1999.

20 Clark, K. L., Skowronski, N., Gallagher, M., Renninger, H., and Schäfer, K.: Effects of invasive insects and fire on forest energy exchange and evapotranspiration in the New Jersey pinelands, *Agricultural and Forest Meteorology*, 166, 50-61, 2012.

De Réaumur, R. A. F.: Observations du thermomètre, faites à Paris pendant l'année 1735, comparées avec celles qui ont été faites sous la ligne, à l'Isle de France, à Alger et en quelques-unes de nos îles de l'Amérique, *Mémoires de l'Académie des Sciences*, 1735. 545-584, 1735.

25 Desai, A. R., Noormets, A., Bolstad, P. V., Chen, J., Cook, B. D., Davis, K. J., Euskirchen, E. S., Gough, C., Martin, J. G., and Ricciuto, D. M.: Influence of vegetation and seasonal forcing on carbon dioxide fluxes across the Upper Midwest, USA: Implications for regional scaling, agricultural and forest meteorology, 148, 288-308, 2008.

Ding, R., Kang, S., Du, T., Hao, X., and Zhang, Y.: Scaling Up Stomatal Conductance from Leaf to Canopy Using a Dual-Leaf Model for Estimating Crop Evapotranspiration, *PloS one*, 9, e95584, 2014.

30 Dragoni, D., Schmid, H. P., Wayson, C. A., Potter, H., Grimmond, C. S. B., and Randolph, J. C.: Evidence of increased net ecosystem productivity associated with a longer vegetated season in a deciduous forest in south-central Indiana, USA, *Global Change Biology*, 17, 886-897, 2011.

Eagleson, P. S.: *Ecohydrology: Darwinian expression of vegetation form and function*, Cambridge University Press, 2005.

Givnish, T. J.: On the Economy of Plant Form and Function: Proceedings of the Sixth Maria Moors Cabot Symposium, Evolutionary Constraints on Primary Productivity, Adaptive Patterns of Energy Capture in Plants, Harvard Forest, August 1983, Cambridge University Press, 1986.

5 Gough, C. M., Hardiman, B. S., Nave, L. E., Bohrer, G., Maurer, K. D., Vogel, C. S., Nadelhoffer, K. J., and Curtis, P. S.: Sustained carbon uptake and storage following moderate disturbance in a Great Lakes forest, *Ecological Applications*, 23, 1202-1215, 2013.

10 Gu, L., Meyers, T., Pallardy, S. G., Hanson, P. J., Yang, B., Heuer, M., Hosman, K. P., Riggs, J. S., Sluss, D., and Wullschleger, S. D.: Direct and indirect effects of atmospheric conditions and soil moisture on surface energy partitioning revealed by a prolonged drought at a temperate forest site, *Journal of Geophysical Research: Atmospheres* (1984–2012), 111, 2006.

He, M., Ju, W., Zhou, Y., Chen, J., He, H., Wang, S., Wang, H., Guan, D., Yan, J., Li, Y., Hao, Y., and Zhao, F.: Development of a two-leaf light use efficiency model for improving the calculation of terrestrial gross primary productivity, *Agricultural and Forest Meteorology*, 173, 28-39, 2013.

15 Hollinger, D. Y., Ollinger, S., Richardson, A., Meyers, T., Dail, D., Martin, M., Scott, N., Arkebauer, T., Baldocchi, D., and Clark, K.: Albedo estimates for land surface models and support for a new paradigm based on foliage nitrogen concentration, *Global Change Biology*, 16, 696-710, 2010.

Hufkens, K., Basler, D., Milliman, T., Melaas, E. K., and Richardson, A. D.: An integrated phenology modelling framework in R, *Methods in Ecology and Evolution*, 9, 1276-1285, 2018.

20 Jenkins, J., Richardson, A. D., Braswell, B., Ollinger, S. V., Hollinger, D. Y., and Smith, M.-L.: Refining light-use efficiency calculations for a deciduous forest canopy using simultaneous tower-based carbon flux and radiometric measurements, *Agricultural and Forest Meteorology*, 143, 64-79, 2007.

Jolly, W. M., Nemani, R., and Running, S. W.: A generalized, bioclimatic index to predict foliar phenology in response to climate, *Global Change Biology*, 11, 619-632, 2005.

25 Levine, N. M., Zhang, K., Longo, M., Baccini, A., Phillips, O. L., Lewis, S. L., Alvarez-Dávila, E., de Andrade, A. C. S., Brienen, R. J., and Erwin, T. L.: Ecosystem heterogeneity determines the ecological resilience of the Amazon to climate change, *Proceedings of the National Academy of Sciences*, 113, 793-797, 2016.

Li, W., Guo, Q., Tao, S., and Su, Y.: VBRT: A novel voxel-based radiative transfer model for heterogeneous three-dimensional forest scenes, *Remote Sensing of Environment*, 206, 318-335, 2018.

30 Liu, J., Chen, J., Cihlar, J., and Park, W.: A process-based boreal ecosystem productivity simulator using remote sensing inputs, *Remote sensing of environment*, 62, 158-175, 1997.

Liu, Q., Fu, Y. H., Liu, Y., Janssens, I. A., and Piao, S.: Simulating the onset of spring vegetation growth across the Northern Hemisphere, *Global change biology*, 24, 1342-1356, 2018.

Melaas, E. K., Friedl, M. A., and Richardson, A. D.: Multiscale modeling of spring phenology across Deciduous Forests in the Eastern United States, *Glob. Change Biol.*, 22, 792-805, 2016.

35 Melaas, E. K., Richardson, A. D., Friedl, M. A., Dragoni, D., Gough, C. M., Herbst, M., Montagnani, L., and Moors, E.: Using FLUXNET data to improve models of springtime vegetation activity onset in forest ecosystems, *Agricultural and Forest Meteorology*, 171, 46-56, 2013.

Miller, G. R., Baldocchi, D. D., Law, B. E., and Meyers, T.: An analysis of soil moisture dynamics using multi-year data from a network of micrometeorological observation sites, *Advances in Water Resources*, 30, 1065-1081, 2007.

5 Myneni, R. B., Hoffman, S., Knyazikhin, Y., Privette, J. L., Glassy, J., Tian, Y., Wang, Y., Song, X., Zhang, Y., Smith, G. R., Lotsch, A., Friedl, M., Morisette, J. T., Votava, P., Nemani, R. R., and Running, S. W.: Global products of vegetation leaf area and fraction absorbed PAR from year one of MODIS data, *Remote Sensing of Environment*, 83, 214-231, 2002.

Ni-Meister, W., Yang, W., and Kiang, N. Y.: A clumped-foliage canopy radiative transfer model for a global dynamic terrestrial ecosystem model. I: Theory, *Agricultural and forest meteorology*, 150, 881-894, 2010.

Oishi, A. C., Oren, R., and Stoy, P. C.: Estimating components of forest evapotranspiration: a footprint approach for scaling sap flux measurements, *Agricultural and Forest Meteorology*, 148, 1719-1732, 2008.

10 Oleson, K., Lawrence, D., Bonan, G., Drewniak, B., Huang, M., Koven, C., Levis, S., Li, F., Riley, W., and Subin, Z.: Technical Description of version 4.5 of the Community Land Model (CLM), NCAR, National Center for Atmospheric Research (NCAR) Boulder, Colorado, 2013. 2013.

Polgar, C. A. and Primack, R. B.: Leaf-out phenology of temperate woody plants: from trees to ecosystems, *New Phytologist*, 191, 926-941, 2011.

15 Potter, C. S., Randerson, J. T., Field, C. B., Matson, P. A., Vitousek, P. M., Mooney, H. A., and Klooster, S. A.: Terrestrial ecosystem production: a process model based on global satellite and surface data, *Global Biogeochemical Cycles*, 7, 811-841, 1993.

Richardson, A. D., Anderson, R. S., Arain, M. A., Barr, A. G., Bohrer, G., Chen, G. S., Chen, J. M., Ciais, P., Davis, K. J., Desai, A. R., Dietze, M. C., Dragoni, D., Garrity, S. R., Gough, C. M., Grant, R., Hollinger, D. Y., Margolis, H. A., 20 McCaughey, H., Migliavacca, M., Monson, R. K., Munger, J. W., Poulter, B., Racsko, B. M., Ricciuto, D. M., Sahoo, A. K., Schaefer, K., Tian, H. Q., Vargas, R., Verbeeck, H., Xiao, J. F., and Xue, Y. K.: Terrestrial biosphere models need better representation of vegetation phenology: results from the North American Carbon Program Site Synthesis, *Global Change Biology*, 18, 566-584, 2012.

25 Running, S. W., Nemani, R. R., Heinsch, F. A., Zhao, M. S., Reeves, M., and Hashimoto, H.: A continuous satellite-derived measure of global terrestrial primary production, *Bioscience*, 54, 547-560, 2004.

Running, S. W. and Zhao, M.: Daily GPP and annual NPP (MOD17A2/A3) products NASA Earth Observing System MODIS land algorithm, *MOD17 User's Guide*, 2015. 2015.

30 Ryu, Y., Baldocchi, D. D., Kobayashi, H., Ingen, C., Li, J., Black, T. A., Beringer, J., Gorsel, E., Knohl, A., and Law, B. E.: Integration of MODIS land and atmosphere products with a coupled-process model to estimate gross primary productivity and evapotranspiration from 1 km to global scales, *Global Biogeochemical Cycles*, 25, GB4017, 2011.

Sellers, P., Randall, D., Collatz, G., Berry, J., Field, C., Dazlich, D., Zhang, C., Collelo, G., and Bounoua, L.: A revised land surface parameterization (SiB2) for atmospheric GCMs. Part I: Model formulation, *Journal of climate*, 9, 676-705, 1996a.

35 Sellers, P. J., Tucker, C. J., Collatz, G. J., Los, S. O., Justice, C. O., Dazlich, D. A., and Randall, D. A.: A revised land surface parameterization (SiB2) for atmospheric GCMs. Part II: The generation of global fields of terrestrial biophysical parameters from satellite data, *Journal of climate*, 9, 706-737, 1996b.

Shen, M., Tang, Y., Chen, J., Zhu, X., and Zheng, Y.: Influences of temperature and precipitation before the growing season on spring phenology in grasslands of the central and eastern Qinghai-Tibetan Plateau, *Agricultural and Forest Meteorology*, 151, 1711-1722, 2011.

Thornton, P., Thornton, M., Mayer, B., Wilhelmi, N., Wei, Y., and Cook, R.: Daymet: Daily surface weather on a 1 km grid for North America, 1980-2008, Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, USA., 2012. 2012.

Turner, D. P., Ritts, W. D., Cohen, W. B., Gower, S. T., Running, S. W., Zhao, M., Costa, M. H., Kirschbaum, A. A., Ham, J. M., and Saleska, S. R.: Evaluation of MODIS NPP and GPP products across multiple biomes, *Remote Sensing of Environment*, 102, 282-292, 2006.

10 Urbanski, S., Barford, C., Wofsy, S., Kucharik, C., Pyle, E., Budney, J., McKain, K., Fitzjarrald, D., Czikowsky, M., and Munger, J.: Factors controlling CO<sub>2</sub> exchange on timescales from hourly to decadal at Harvard Forest, *Journal of Geophysical Research: Biogeosciences*, 112, 2007.

White, M. A., Thornton, P. E., Running, S. W., and Nemani, R. R.: Parameterization and sensitivity analysis of the BIOME–BGC terrestrial ecosystem model: net primary production controls, *Earth interactions*, 4, 1-85, 2000.

15 Xiao, X., Zhang, Q., Braswell, B., Urbanski, S., Boles, S., Wofsy, S., Moore, B., and Ojima, D.: Modeling gross primary production of temperate deciduous broadleaf forest using satellite images and climate data, *Remote Sensing of Environment*, 91, 256-270, 2004.

20 Xiao, Z., Liang, S., Wang, J., Chen, P., Yin, X., Zhang, L., and Song, J.: Use of General Regression Neural Networks for Generating the GLASS Leaf Area Index Product From Time-Series MODIS Surface Reflectance, *IEEE Transactions on Geoscience and Remote Sensing*, 2014. 209-223, 2014.

Xie, J., Chen, J., Sun, G., Chu, H., Noormets, A., Ouyang, Z., John, R., Wan, S., and Guan, W.: Long-term variability and environmental control of the carbon cycle in an oak-dominated temperate forest, *Forest Ecology and Management*, 313, 319-328, 2014.

25 Xin, Q.: A risk-benefit model to simulate vegetation spring onset in response to multi-decadal climate variability: Theoretical basis and applications from the field to the Northern Hemisphere, *Agricultural and Forest Meteorology*, 228–229, 139-163, 2016.

Xin, Q., Dai, Y., Li, X., Liu, X., Gong, P., and Richardson, A. D.: A steady-state approximation approach to simulate seasonal leaf dynamics of deciduous broadleaf forests via climate variables, *Agricultural and Forest Meteorology*, 249, 44-56, 2018.

30 Yang, X., Mustard, J. F., Tang, J., and Xu, H.: Regional-scale phenology modeling based on meteorological records and remote sensing observations, *Journal of Geophysical Research: Biogeosciences*, 117, 2012.

Yu, C., Li, C., Xin, Q., Chen, H., Zhang, J., Zhang, F., Li, X., Clinton, N., Huang, X., Yue, Y., and Gong, P.: Dynamic assessment of the impact of drought on agricultural yield and scale-dependent return periods over large geographic regions, *Environmental Modelling & Software*, 62, 454-464, 2014.

35 Yuan, H., Dickinson, R. E., Dai, Y., Shaikh, M. J., Zhou, L., Shangguan, W., and Ji, D.: A 3D Canopy Radiative Transfer Model for Global Climate Modeling: Description, Validation, and Application, *Journal of Climate*, 27, 1168-1192, 2013.

Yuan, W., Liu, S., Yu, G., Bonnefond, J.-M., Chen, J., Davis, K., Desai, A. R., Goldstein, A. H., Gianelle, D., and Rossi, F.: Global estimates of evapotranspiration and gross primary production based on MODIS and global meteorology data, *Remote Sensing of Environment*, 114, 1416-1431, 2010.

5 Zeng, F., Collatz, G. J., Pinzon, J. E., and Ivanoff, A.: Evaluating and quantifying the climate-driven interannual variability in Global Inventory Modeling and Mapping Studies (GIMMS) Normalized Difference Vegetation Index (NDVI3g) at global scales, *Remote Sensing*, 5, 3918-3950, 2013.

Zhu, P., Zhuang, Q., Ciais, P., Welp, L., Li, W., and Xin, Q.: Elevated atmospheric CO<sub>2</sub> negatively impacts photosynthesis through radiative forcing and physiology-mediated climate feedback, *Geophysical Research Letters*, 44, 1956-1963, 2017.