

# Impact of peatlands on carbon dioxide (CO<sub>2</sub>) emissions from the Rajang River and Estuary, Malaysia

Denise Müller-Dum<sup>1</sup>, Thorsten Warneke<sup>1</sup>, Tim Rixen<sup>2,3</sup>, Moritz Müller<sup>4</sup>, Antje Baum<sup>2</sup>, Aliko Christodoulou<sup>1</sup>, Joanne Oakes<sup>5</sup>, Bradley D. Eyre<sup>5</sup>, and Justus Notholt<sup>1</sup>

- 5 <sup>1</sup> Institute of Environmental Physics, University of Bremen, Otto-Hahn-Allee 1, 28359 Bremen, Germany  
<sup>2</sup> Leibniz Center for Tropical Marine Research, Fahrenheitstr. 6, 28359 Bremen, Germany  
<sup>3</sup> Institute of Geology, University of Hamburg, Bundesstr. 55, 20146 Hamburg, Germany  
<sup>4</sup> Swinburne University of Technology, Faculty of Engineering, Computing and Science, Jalan Simpang Tiga, 93350 Kuching, Sarawak, Malaysia  
10 <sup>5</sup> Centre for Coastal Biogeochemistry, School of Environment, Science and Engineering, Southern Cross University, Lismore NSW 2480, Australia

*Correspondence to:* Denise Müller-Dum, [dmueller@iup.physik.uni-bremen.de](mailto:dmueller@iup.physik.uni-bremen.de)

**Abstract.** Tropical peat-draining rivers are known as potentially large sources of carbon dioxide (CO<sub>2</sub>) to the atmosphere due to high loads of carbon they receive from surrounding soils. However, not many seasonally resolved data are available, limiting our understanding of these systems. We report the first measurements of carbon dioxide partial pressure (*p*CO<sub>2</sub>) in the Rajang River and Estuary, the longest river in Malaysia. The Rajang River catchment is characterized by extensive peat deposits found in the delta region, and by human impact such as logging, land use and river damming. *p*CO<sub>2</sub> averaged 2540 ± 189 μatm during the wet season and 2350 ± 301 μatm during the dry season. Using three different parameterizations for the gas transfer velocity, calculated CO<sub>2</sub> fluxes to the atmosphere were 1.5 (0.5-2.0) g C m<sup>-2</sup> d<sup>-1</sup> (mean, minimum – maximum) during the wet season and 1.7 (0.6-2.6) g C m<sup>-2</sup> d<sup>-1</sup> during the dry season. This is at the low end of reported values for Southeast Asian peat-draining rivers, but similar to values reported for Southeast Asian rivers that do not flow through peat deposits. In the Rajang River, peatlands probably do not contribute much to the CO<sub>2</sub> flux due to the proximity of the peatlands to the coast, which limits the opportunity for degradation of organic C during transport. Thus, we suggest that peat coverage is, by itself, insufficient as sole predictor of CO<sub>2</sub> emissions from peat-draining rivers, and that other factors, like the spatial distribution of peat in the catchment and pH, also need to be considered.

## 1 Introduction

Tropical rivers transport large amounts of terrestrially derived carbon to the ocean (Dai et al., 2012) and the atmosphere (Aufdenkampe et al., 2011; Raymond et al., 2013). It has been estimated that 78% of riverine carbon dioxide (CO<sub>2</sub>) emissions occur in the tropics (Lauerwald et al., 2015). Tropical wetlands exert a particularly strong influence on the carbon budget of these rivers. Two regional studies independently showed that the partial pressure of CO<sub>2</sub> (*p*CO<sub>2</sub>) in rivers increases with increasing wetland coverage in the catchment. Borges et al. (2015) established a relationship between wetland extent and *p*CO<sub>2</sub> for African rivers. Wit et al. (2015) presented an analog synthesis for Southeast Asian rivers, which flow through peatlands. Peatlands are a special type of wetland, where organic matter accumulates at rates that make them the most effective terrestrial carbon store on a millennial timescale (Dommain et al., 2011). Southeast Asian peatlands store 68.5 Gt carbon (Page et al., 2011). The highest riverine dissolved organic carbon (DOC) concentrations reported so far were found in Southeast Asian peat-draining rivers (Alkhatib et al. 2007; Moore et al., 2011; Müller et al., 2015), with an annual average of 68 mg L<sup>-1</sup> DOC found in an undisturbed peat-draining river (Moore et al., 2013). Because of these high DOC concentrations, Indonesian rivers may account for 75 % of the DOC flux into the South China Sea (SCS) while accounting for 39 % of the discharge (Huang et al., 2017). Surprisingly, CO<sub>2</sub> emissions from these rivers are not exceptionally high (Müller et al., 2015; Wit et al., 2015). This is attributed to a short residence time of the organic matter in the river, allowing little time for decomposition, and the resistance of peat-derived carbon to bacterial degradation. Nevertheless, the CO<sub>2</sub> flux from peat-draining rivers to the atmosphere increases with increasing peat coverage in the river basin (Wit et al., 2015), showing that these ecosystems exert an important influence on a river's carbon budget.

However, most Southeast Asian peat-draining rivers are disturbed by human activities such as river damming, urbanization, deforestation (Milliman and Farnsworth, 2011) and discharge of untreated wastewater (Park et al., 2018). Anthropogenic change poses a new challenge to understanding carbon fluxes in Asian river systems, and more data are urgently needed to constrain the carbon budget for this important region (Park et al., 2018). In Malaysia, the country holding the second largest share of tropical peat (Page et al., 2011), river CO<sub>2</sub> emissions have only been studied in a small undisturbed peat-draining river (Müller et al., 2015), in estuaries (Chen et al., 2013; Müller et al., 2016) and in two river reaches which were not influenced by peat (Müller et al., 2016). In this study, the longest Malaysian river, the Rajang River on the island of Borneo, was investigated. This river flows through largely logged-over tropical rainforest (Gaveau et al., 2014), urban areas and disturbed peat swamps (Gaveau et al., 2016). The aim of this study was to assess the Rajang River and Estuary carbon load and to investigate the impact of peatlands on its CO<sub>2</sub> emissions. To this end, we surveyed longitudinal transects extending from river reaches that were not influenced by peat to the peat-covered delta.

## 2 Materials and Methods

### 2.1 Study area

The Rajang River is located in the Malaysian state of Sarawak in the northern part of the island of Borneo (Fig. 1a). Sarawak has a tropical climate with high temperatures (average 26.6°C, 1992-2016 in Sibul, DWD, 2018) and high precipitation (average 3,578 mm yr<sup>-1</sup>, 1992-2016 in Sibul, DWD, 2018). The region experiences two monsoonal periods: the northeastern monsoon with enhanced rainfall and frequent floods occurs between December and February (“wet season”, see Fig. 2a), while the southwestern monsoon from May until September is associated with relatively drier weather (“dry season”). However, despite the monsoon seasons, rainfall is high throughout the year (Sa’adi et al., 2017).

The Rajang River originates in the Iran mountains, a mountain range at the border between Malaysia and Indonesia (MacKinnon, 1996) with elevations of up to 1,800 m (Milliman and Farnsworth, 2011). It drains an area of approximately 52,010 km<sup>2</sup> (Lehner et al., 2006; DID 2017) whose geology is dominated by Cenozoic sedimentary and metamorphic rocks, consisting of siliciclastic rock with minor amounts of carbonates (Staub et al., 2000; Milliman and Farnsworth, 2011). The Rajang River flows approximately 530 km from east to west and discharges into the South China Sea (Milliman and Farnsworth, 2011). Main settlements along the river are the towns of Kapit, Kanowit and the city of Sibul (163,000 inhabitants) (see Fig. 1b). In addition, a large number of longhouses (traditional buildings inhabited by local communities) are located along the river and its tributaries (Ling et al., 2017). Hydroelectric power plants were built on two tributaries in the upper Rajang basin: The Bakun hydroelectric power plant commenced operation in 2011 and the Murum dam in 2015 (Sarawak Energy, 2013, see Fig. 1b). The construction of another hydroelectric power plant on a tributary in the southern Rajang basin is planned for the future (Sarawak Energy, 2013).

The Rajang delta system is comprehensively described in Staub and Gastaldo (2003). It is entirely surrounded by peatlands (Fig. 1b), which extend over an area that corresponds to approximately 11% of the catchment size (Nachtergaele et al., 2009). Most of these peatlands have been converted to industrial oil palm plantations (Gaveau et al., 2016, Fig. 1b). The main distributary channels forming the delta (from north to south) are the Igan, Hulu Seredeng (which splits up into Lassa and Paloh), Belawai and Rajang, which have a maximum tidal range (spring tide) of 3-6 m (Staub and Gastaldo, 2003). Saltwater intrudes into the estuary approximately as far as the point where the Rajang River splits up into its four southernmost distributaries, a few kilometers downstream of Sibul (Fig. 1b), depending on season. Tidal influence extends further inland approximately up to the town of Kanowit (Staub and Gastaldo, 2003).

Monthly discharge (Fig. 2a) was estimated from monthly precipitation (1992-2016; DWD, 2018) and an evapotranspiration rate of 1,545 mm yr<sup>-1</sup> (Kumagai et al., 2005) or 43.2%. Annual average discharge from 1992-2016 was 3,355 m<sup>3</sup> s<sup>-1</sup>, in good agreement with reported discharges of 3,490 m<sup>3</sup> s<sup>-1</sup> (Milliman and Farnsworth, 2011) and 3,372 m<sup>3</sup> s<sup>-1</sup> for the years 1991-2015 (Sa’adi et al, 2017).

## 2.2 Surveys

We sampled the Rajang River during two surveys, which were designed to get spatial coverage of both peat and non-peat areas during the wettest and driest period of one year. The first survey took place at the peak of the monsoon season in January 2016 (“wet season”). The second one was performed during the dry season in August 2016 (“dry season”). In January 2016, we entered the Rajang River through the Rajang river mouth (distributary 5 in Fig.1b), went upstream to the town of Kapit and back downstream to the town of Belawai at the Belawai river mouth (distributary 4 in Fig. 1b). In August 2016, we entered the Rajang River through the Rajang river mouth (5), went upstream to Kapit and back to Sibuluan. From there, we went out to the coast through the Lassa distributary (2), and back to Sibuluan through the Igan distributary (1). The last sampling stretch was from Sibuluan into the Paloh distributary (3) and back to Belawai (4). During this campaign, one stationary measurement was performed overnight in Sarikei in the Rajang distributary in order to assess tidal/diurnal variability.

## 2.3 CO<sub>2</sub> measurements

The setup on the boat was similar to the one described in Müller et al. (2016). Surface water was pumped through a shower-type equilibrator (Johnson, 1999) at a rate of approximately 15 L min<sup>-1</sup>. In the beginning, the equilibrator headspace was connected to an FTIR analyzer (Griffith et al., 2012), which allows for the simultaneous measurement of CO<sub>2</sub>, methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O) and carbon monoxide (CO). During the cruise in January 2016, a failure of the FTIR analyzer occurred and measurements were continued (also in August 2016) using an Li-820 non-dispersive infrared (NDIR) analyzer for the measurement of CO<sub>2</sub> (Licor, USA). For calibration and inter-calibration of the two instruments, a set of gravimetrically prepared gas mixtures (Deuste Steininger) in the range of the World Meteorological Organization (WMO) standard scale was measured, which were calibrated by the Max Planck Institute for Biogeochemistry in Jena, Germany. For the FTIR, spectra were averaged over 5 minutes and dry air mole fractions were retrieved using the software MALT5 (Griffith 1996). Li-820 data were stored with a temporal resolution of 1 minute. Gas partial pressure was determined using measurements of ambient pressure with a PTB110 barometer (Vaisala, Finland) and correction for removal of water according to Dickson et al. (2007). Water temperature was measured in the equilibrator and in the surface water and correction to water surface temperature was performed according to Dickson et al. (2007).

In August 2016, the internal pressure sensor of the Li-820 failed. Because the instrument performs an internal correction based on the cell pressure, this correction had to be reversed and recalculated with an assumed internal cell pressure. This procedure is described in the Supplement.

CO<sub>2</sub> fluxes ( $FCO_2$ , in gC m<sup>-2</sup> d<sup>-1</sup>) across the water-air interface were computed using the gas transfer equation

$$FCO_2 = kK_0(pCO_2^{water} - pCO_2^{air}) \cdot f_1 f_2 \quad (1),$$

where  $k$  is the gas transfer velocity (m s<sup>-1</sup>),  $K_0$  is the solubility (mol L<sup>-1</sup> atm<sup>-1</sup>) calculated according to Weiss (1974),  $pCO_2^{water}$  is the partial pressure of CO<sub>2</sub> in water,  $pCO_2^{air}$  is the partial pressure of CO<sub>2</sub> in the overlying air (both in  $\mu$ atm),  $f_1$  is a conversion

factor from  $L^{-1}$  to  $m^{-3}$ , and  $f_2$  is a conversion factor from  $\mu\text{mol s}^{-1}$  to  $\text{mg d}^{-1}$ . The atmospheric mole fractions of  $\text{CO}_2$  during the months of our measurements were derived from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network (Dlugokencky et al., 2018) for the closest station, which was Bukit Kototabang, Indonesia.

The gas transfer velocity ( $k$ ) is a critical, yet poorly constrained parameter. Many studies have attempted to relate  $k$  to the main drivers of turbulence, such as wind speed (e.g., Wanninkhof, 1992; Nightingale, 2000; Raymond and Cole, 2001) or, especially for rivers, catchment parameters like slope, water flow velocity and discharge (Raymond et al., 2012). We sampled mainly the downstream reaches of the Rajang River, which range in width from 271 m to several kilometers in the delta (Allen and Pavelsky, 2018). Therefore,  $k$ -parameterizations that were developed for estuaries or big rivers were considered the most appropriate. We compared three parameterizations to constrain the  $\text{CO}_2$  fluxes. The first parameterization is the one by Borges et al. (2004) for estuaries, which is driven both by wind speed and water flow velocity; the one by Alin et al. (2011), which was developed for rivers wider than 100 m and is driven by wind speed; and the one by Raymond and Cole (2001), which is driven by wind speed and was developed for big rivers and estuaries. Those parameterizations read:

$$k_{600,B04} = 1.0 + 1.719w^{0.5}h^{-0.5} + 2.58 u_{10} \quad (2),$$

$$k_{600,A11} = 4.46 + 7.11 \cdot u_{10} \quad (3),$$

$$15 \quad k_{600,R01} = 1.91 \cdot e^{0.35u_{10}} \quad (4),$$

where  $w$  is the water flow velocity ( $\text{cm s}^{-1}$ ),  $h$  is the depth (m) and  $u_{10}$  is the wind speed at 10 m ( $\text{m s}^{-1}$ ).  $h$  was taken from the bottom sounder recordings of our boat.  $w$  in the lower river reaches was measured by Staub and Esterle (1993) to be  $0.7 \text{ m s}^{-1}$ . A more recent study by Ling et al. (2017) reports  $w = 1.1 \text{ m s}^{-1}$  for the Rajang River upstream from Kapit. For the calculation of the gas exchange velocity  $k$ , we used the average of  $w = 0.9 \text{ m s}^{-1}$ . For  $u_{10}$ , on-site wind speed data was unfortunately not available. In such cases, other authors (e.g., Bouillon et al., 2012; some estuaries in Chen et al., 2013) have resorted to gridded wind data from the NOAA NCEP NCAR Reanalysis product (Kalnay et al., 1996). While we acknowledge the uncertainty introduced by using gridded data instead of in situ wind speed, we used this product as well, as the best one available for our study area. We retrieved daily wind speed at 10 m for the grid centered at  $2.85^\circ\text{N}$ ,  $112.5^\circ\text{E}$  for the time of our measurements. In-situ  $k$  is dependent on in-situ salinity and temperature and was calculated from  $k_{600}$ , exploiting its relationship with the Schmidt number (Wanninkhof, 1992).

The calculation of fluxes using  $k$ -parameterizations is associated with a large uncertainty, and it is difficult to determine the most suitable parameterization if none of them was developed in the study region. In addition, using input data from the literature (as for the water flow velocity) or gridded instead of measured wind data adds to this uncertainty (see Supplement). However, using three different parameterizations we are able to constrain the magnitude of  $\text{CO}_2$  emissions from the Rajang River. We will report the average  $\text{CO}_2$  fluxes from the three different parameterizations as well as minimum and maximum fluxes.

## 2.4 Ancillary measurements

In January 2016, individual water samples were taken at 15 stations between the river mouth and Kapit, including the distributary channels Rajang and Belawai. In August 2016, water samples were taken at 34 stations, with a higher sampling frequency and coverage in the delta (Rajang, Igan, Lassa, Paloh and Belawai, Fig. 1b). Water samples were taken from approximately 1 m below the surface using a Van Dorn water sampler. Particulate material was sampled on pre-weighed and pre-combusted glass fiber filters. From the net sample weight and the volume of filtered water, the amount of suspended particulate matter (SPM) was determined. For POC, 1N hydrochloric acid was added in order to remove inorganic carbon from the sample. For the determination of carbon, samples were catalytically combusted at 1050°C and combustion products were measured by thermal conductivity using a Euro EA3000 Elemental Analyzer. Repeatability for C content was 0.04 % (standard deviation).

In August 2016, water samples were also taken for the determination of dissolved inorganic carbon (DIC) and the isotopic composition ( $\delta^{13}\text{C}$ ) of DIC, because the isotopic composition of DIC can help in identifying its sources (Das et al., 2005; Campeau et al., 2017; 2018). Samples were poisoned with 200  $\mu\text{L}$  concentrated  $\text{HgCl}_2$  and filtered through Whatman glass fiber filters (GF/F, pore size 0.7  $\mu\text{m}$ ). 40 ml sampling vials were filled to the top, leaving no headspace, checked for the existence of bubbles, and stored refrigerated until analysis. Concentrations and  $\delta^{13}\text{C}$  of DIC were measured via continuous flow wet-oxidation isotope ratio mass spectrometry (CF-WO-IRMS) using an Aurora 1030W TOC analyzer coupled to a Thermo Delta V Plus IRMS (Oakes et al., 2010). Sodium bicarbonate (DIC) of known isotope composition dissolved in helium-purged milli-Q was used for drift correction and to verify concentrations and  $\delta^{13}\text{C}$  values. Reproducibility for DIC was  $\pm 10 \mu\text{mol L}^{-1}$  for concentrations and  $\pm 0.10\text{‰}$  for  $\delta^{13}\text{C}$  (standard deviations).

During both surveys, dissolved oxygen and water temperature were continuously measured with a temporal resolution of 5 minutes using an FDO 925 oxygen sensor and a WTW 3430 data logger (Xylem Inc., USA). The oxygen sensor was calibrated by the manufacturer, a routine function check was performed before the start of measurements using the check and calibration vessel (FDO © Check) provided by the company. The reported accuracy of a dissolved oxygen measurement at 20°C in air-saturated water is 1.5%, the precision of the accompanying temperature measurement is 0.2°C (WTW, 2012). pH, salinity and temperature were measured at the stations, using a SenTix 940 pH sensor (pH) and a Multiprobe (Aquaread AP-2000). The pH sensor was calibrated before the start of the measurements using NIST (National Institute of Standards and Technology) traceable buffers. Since salinity was only measured at the stations, we spatially interpolated salinity for the interpretation of  $p\text{CO}_2$  data. This procedure is described in the Supplement.

Different geographical extents of the river were covered during the two campaigns, and the salt intrusion limits were different during the two seasons. In order to keep the results from the two surveys comparable, we report results for three categories: non-peat (Kapit-Kanowit), peat (Kanowit-Sibu) and delta (downstream of Sibu). Their definition and properties are specified in Table 1. The non-peat and peat areas are directly comparable between seasons, because the same spatial extent was covered during both surveys and they were non-saline during both seasons.

## 2.5 Data analysis and export calculations

Data analysis was performed with Python 2.7.15 and ArcMap 10.5. Averages of measured parameters are reported  $\pm 1$  standard error unless stated otherwise. Errors for calculated parameters (e.g., river loads, see below) were determined with error propagation. For fluxes and derived quantities, we report the mean, minimum and maximum from the three  $k$ -parameterizations. Seasonal differences were tested for significance using the Mann-Whitney U-test from the Python Scipy Statistical Functions module. Data from the delta were excluded from the statistical tests due to the different geographical coverage achieved during the two surveys.

In order to calculate the total carbon export from the Rajang River for the months of our measurements, we derived DOC load, POC load, DIC load for the peat and non-peat area, as well as CO<sub>2</sub> outgassing as follows:

The river loads of DOC, POC and DIC were calculated for the peat and non-peat area combined, using

$$RIVER\ LOAD = C \cdot Q \cdot f_3 \quad (5),$$

where  $C$  is the average concentration of DOC/POC/DIC in mg L<sup>-1</sup>,  $Q$  is monthly discharge (m<sup>3</sup> s<sup>-1</sup>) and  $f_3$  is a conversion factor from s<sup>-1</sup> to month<sup>-1</sup>.

For DOC, we used the DOC concentrations reported by Martin et al. (2018). This data was acquired during 2017 downstream of Kanowit. Only freshwater values were considered (average for the wet and dry season: 2.0 mg L<sup>-1</sup> and 2.1 mg L<sup>-1</sup>). For POC, we used the area-weighted average concentration of peat and non-peat river reaches determined during our surveys.

For DIC, we used an area-weighted average concentration as well, which was determined from our measurements during the dry season. For the wet season survey, DIC was calculated from pH and  $p\text{CO}_2$  using the program CO<sub>2</sub>sys (Lewis and Wallace, 1998). Note that pH measurements were only available at the stations, and sometimes we did not have parallel  $p\text{CO}_2$  measurements. Therefore, the number of calculated DIC values for the peat and non-peat area is 6.

For the calculation of total CO<sub>2</sub> emissions from FCO<sub>2</sub>, the river surface area was required. River surface area was calculated from the GRWL (Global River Widths from Landsat) Database (Allen & Pavelsky, 2018) using Esri's ArcMap 10.5. Missing segments in the delta were manually delineated using a Landsat satellite image and their surface area was determined. This procedure is described in the Supplement. With the surface area of individual river segments at hand, CO<sub>2</sub> emissions were calculated for the non-peat area, peat area and the delta separately.



### 3 Results

#### 3.1 General characterization of the Rajang River

Measured salinity ranged between 0 and 18.6 during the wet season and 0 and 32.1 during the dry season. Saltwater was detected further upstream during the dry season than during the wet season (Fig. 3a and b). Saltwater penetrated further inland in the Rajang and Belawai distributaries than in the Igan distributary (Fig. 3b), suggesting that most freshwater is discharged via the Igan distributary.

The Rajang River was slightly acidic (6.7 (wet) and 6.8 (dry), area-weighted mean for the peat and non-peat area, see Table 2) and highly turbid, with area-weighted average SPM concentrations of  $187.2 \pm 75.7 \text{ mg L}^{-1}$  (wet season) and  $51.5 \pm 12.1 \text{ mg L}^{-1}$  (dry season, see Table 2). With higher SPM during the wet season ( $p < 0.001$ ), the organic carbon content of SPM was significantly decreased ( $1.5 \pm 0.4 \%$  on average,  $p = 0.01$ ) compared to the dry season ( $2.1 \pm 0.6 \%$  on average, see Table 2). POC ranged from  $0.7 \text{ mg L}^{-1}$  to  $9.1 \text{ mg L}^{-1}$  during the wet season (average  $2.6 \pm 0.6 \text{ mg L}^{-1}$ ) and from  $0.3 \text{ mg L}^{-1}$  to  $1.9 \text{ mg L}^{-1}$  during the dry season (average  $1.1 \pm 0.4 \text{ mg L}^{-1}$ , see Table 2). The seasonal difference was significant ( $p = 0.002$ ).

The river water was consistently undersaturated with oxygen with respect to the atmosphere. DO oversaturation was not observed. The area-weighted average DO was similar during the wet and dry seasons ( $81.1 \pm 5.5 \%$  and  $79.6 \pm 3.5 \%$ , wet/dry), with slightly lower DO in the delta ( $66.0 \pm 6.9 \%$  and  $71.2 \pm 11.1 \%$  (wet/dry)).

Measured DIC in the dry season ranged from  $153.7 \mu\text{mol L}^{-1}$  in the non-peat area to  $2399.2 \mu\text{mol L}^{-1}$  in the estuary and increased linearly with salinity (Fig. 4a,  $r = 0.98$ ,  $p < 0.001$ ). Concentrations averaged  $177.9 \pm 22.4 \mu\text{mol L}^{-1}$  in the peat and non-peat area and  $1302.3 \pm 749.2 \mu\text{mol L}^{-1}$  in the delta (Table 2). Calculated DIC for the wet season averaged  $289.8 \pm 32.1 \mu\text{mol L}^{-1}$  (area-weighted mean for the non-peat and peat area).  $\delta^{13}\text{C}$ -DIC ranged between  $-11.9 \text{ ‰}$  and  $-1.4 \text{ ‰}$  and averaged  $-7.0 \pm 1.5 \text{ ‰}$  in the peat and non-peat area and  $-5.9 \text{ ‰}$  in the delta (Table 2).  $\delta^{13}\text{C}$ -DIC was positively correlated with DIC for the delta ( $r = 0.81$ ,  $p < 0.001$ ) and negatively correlated with DIC for the freshwater part (peat and non-peat combined,  $r = -0.98$ ,  $p = 0.004$ , Fig. 4b). A Keeling plot revealed a linear relationship for freshwater samples (Fig. 4c) with a y-intercept ( $\pm$  SE) of  $-18.6 \pm 0.3 \text{ ‰}$ .

#### 3.2 Carbon dioxide

The Rajang River was oversaturated with  $\text{CO}_2$  with respect to the atmosphere, with an average  $p\text{CO}_2$  of  $2531 \pm 188 \mu\text{atm}$  (wet season) and  $2337 \pm 304 \mu\text{atm}$  (dry season) in the non-peat area.  $p\text{CO}_2$  was significantly higher in the peat area ( $p < 0.001$ , both seasons) with  $2990 \pm 239 \mu\text{atm}$  (wet season) and  $2994 \pm 141 \mu\text{atm}$  (dry season). The area-weighted means for the peat and non-peat area were  $2540 \pm 189 \mu\text{atm}$  (wet season) and  $2340 \pm 301 \mu\text{atm}$  (dry season). In the delta,  $p\text{CO}_2$  was more variable, and the average values of  $3005 \pm 1039 \mu\text{atm}$  (wet season) and  $2783 \pm 1437 \mu\text{atm}$  (dry season, see Table 2) were also significantly higher than in the non-peat area ( $p < 0.001$ , both seasons).  $p\text{CO}_2$  values were strikingly similar between wet and dry season, and so were the spatial patterns in  $p\text{CO}_2$  (Fig. 3c and d). Tidal variability of  $p\text{CO}_2$  was observed at an overnight

station in Sarikei in August 2016. During this time,  $p\text{CO}_2$  increased from approximately 3000  $\mu\text{atm}$  to almost 6000  $\mu\text{atm}$  during rising tide (see Supplement).

$p\text{CO}_2$  decreased with increasing salinity in the delta (Fig.3). However, a big spread of data in both the high-salinity region (during the tidal measurement described above) and the freshwater region was observed.  $p\text{CO}_2$  was correlated with DO (Fig. 5). An interesting pattern is consistently visible in both the wet and dry season data, by which main stem data can clearly be distinguished from those collected in the Belawai and Paloh distributaries. Unfortunately, our data did not allow identification of a diurnal signal for either  $p\text{CO}_2$  or DO. In the tidal part of the river, we had only one stationary measurement overnight, when a diurnal signal could not be identified due to the strong tidal signal. In the non-tidal part of the river, we had insufficient night-time data to make a statement about a day-night difference for  $p\text{CO}_2$  and DO.

10 Wind speed in the grid centered at 2.85°N, 112.5°E averaged 0.57  $\text{m s}^{-1}$  during our campaign in January 2016 and 1.09  $\text{m s}^{-1}$  during our campaign in August 2016. The calculated gas exchange velocities for a Schmidt number of 600 using the B04 model ( $k_{600, B04}$ ) were 8.23  $\text{cm h}^{-1}$  and 9.57  $\text{cm h}^{-1}$ , respectively. This compares to the A11 model with 8.51  $\text{cm h}^{-1}$  and 12.19  $\text{cm h}^{-1}$  and to the R01 model with 2.32  $\text{cm h}^{-1}$  and 2.79  $\text{cm h}^{-1}$  for the wet and dry season, respectively. The resultant  $\text{CO}_2$  fluxes ( $F\text{CO}_2$ ) to the atmosphere ranged between 0.5 and 2.4  $\text{gC m}^{-2} \text{d}^{-1}$  in the wet season and between 0.6 and 3.5  $\text{gC m}^{-2} \text{d}^{-1}$  in the  
15 dry season (per water surface unit area, see Table 2).

### 3.3 Carbon river load and $\text{CO}_2$ emissions

Discharge was above average during 2016 (Fig. 2a). Discharge during January 2016 (wet season) was in accordance with the long-term average, but discharge during August 2016 (dry season) was higher than usual. The Rajang River loads for the peat and non-peat area were 104 (82-123) GgC in January 2016 and 65 (45-83) GgC in August 2016; another 31 (12-41) (wet) and  
20 34 (12-51) Gg were emitted as  $\text{CO}_2$  from the delta (Table 3). Of the river loads of carbon, 91 (86-97) % (wet) and 82 (70-94) % (dry) were exported laterally by discharge. Approximately half of the laterally transported carbon was in the organic form ( $57 \pm 12\%$  and  $60 \pm 18\%$ , wet/dry). River loads were similar during both seasons, except that POC export was 3-fold higher in the wet season.  $\text{CO}_2$  emissions to the atmosphere accounted for 9 (3-14) % and 18 (6-30) % (wet/dry) of the total carbon load of the river.

## 25 4. Discussion

### 4.1 Sediment yield and organic carbon load

The Asia-Pacific region is known for its high sediment yields, especially where rivers drain Cenozoic sedimentary and volcanic rock (Milliman and Farnsworth, 2011). Therefore, the high suspended sediment load in the Rajang River is not surprising. However, SPM concentrations during our expeditions were substantially lower ( $187.2 \pm 75.7 \text{ mg L}^{-1}$  and  $51.5 \pm 12.1 \text{ mg L}^{-1}$ )  
30 than in July 1992 ( $613 \text{ mg L}^{-1}$ , Staub and Esterle, 1993). This could be an effect of upstream dams (operational since 2011 and

2015), which trap sediment in their reservoirs, thereby reducing downstream sediment loads (Vörösmarty et al., 2003, Snoussi et al., 2002). In support of this, SPM concentrations were intermediate in the upper Rajang River in 2014/2015 (218.3 mg L<sup>-1</sup>; Ling et al., 2017). These measurements were taken before, and the measurements in the current study were taken after, the Murum dam began full operation in the second quarter of 2015. Furthermore, SPM and POC concentrations (2.6 mg L<sup>-1</sup> and 5 1.1 mg L<sup>-1</sup>, wet/dry) in the Rajang River were similar to those in the Pearl River, China, (SPM: 70 mg L<sup>-1</sup>-247 mg L<sup>-1</sup>, POC: 1.0 mg L<sup>-1</sup>-3.8 mg L<sup>-1</sup>, Ni et al., 2008) and the Red River, Vietnam, (SPM: 294 ± 569 mg L<sup>-1</sup> (wet) and 113 ± 428 mg L<sup>-1</sup> (dry), POC: 3.7 ± 2.0 mg L<sup>-1</sup> (wet) and 1.1 ± 1.1 mg L<sup>-1</sup> (dry), Le et al., 2017), both of which are also affected by damming.

SPM was higher during the wet season than during the dry season in agreement with observations at the Kinabatangan River, Malaysia (Harun et al., 2014). This can be attributed to enhanced erosion during the wet season. In logged-over forest, as found 10 in most of the Rajang River basin, the energy impact of rain drops on the soil is higher than in densely vegetated areas, where rain is intercepted by the canopy before falling on the ground (Ling Lee et al., 2004). In agreement with this line of reasoning, Ling et al. (2016) showed that the amount of suspended solids in Malaysian streams draining areas with logging activities increased significantly after rain events. The decreased organic carbon content observed during the wet season further supports this, as it indicates a higher contribution of eroded mineral soil to SPM. This pattern is observed in many rivers in this region 15 (Huang et al., 2017). Despite the changing carbon content, most POC was still exported during the wet season, as in other Southeast Asian rivers (Ni et al., 2008; Moore et al., 2011). The proportion of laterally transported carbon in the Rajang River that is in organic form (57 ± 12 % and 60 ± 18%, wet/dry) is similar to that reported for the carbon flux to the South China Sea (50 ± 14%, Huang et al., 2017).

## 4.2 Inorganic carbon load

### 20 4.2.1 DIC concentrations and sources

DIC concentrations in the Rajang River were comparable to those reported by Huang et al. (2017) for the Rajang River (201 μmol L<sup>-1</sup> and 487 μmol L<sup>-1</sup>), but substantially lower than reported for the Mekong River (1173-2027 μmol L<sup>-1</sup>, Li et al., 2013) and the Pearl River (1740 μmol L<sup>-1</sup>, Huang et al., 2017). DIC concentrations in the Rajang are similar to those in the Musi River, Indonesia (250 μmol L<sup>-1</sup>, Huang et al., 2017), suggesting that the Rajang River compares better to the equatorial 25 Indonesian rivers than to rivers draining mainland Southeast Asia, probably because of the scarcity of carbonate rock, which has the highest weathering rate and is thus responsible for high DIC in rivers (Huang et al., 2012).

The source of DIC varied along the length of the Rajang River. In the estuarine part, the positive linear relationship between DIC and salinity (Fig. 4a) suggests that the main source of DIC in the estuary is marine. This is also supported by the relatively high δ<sup>13</sup>C-DIC of estuarine samples, as ocean DIC is more enriched, with δ<sup>13</sup>C-DIC between 0 and 2.5 ‰ (Rózanski et al., 30 2003). The positive correlation between DIC and δ<sup>13</sup>C in the delta thus implies an increasing contribution of marine DIC.

In the freshwater part of the Rajang River,  $\delta^{13}\text{C}$  values were more depleted ( $-7.0 \pm 1.5 \text{ ‰}$ , Table 2) than downstream, but were more enriched than  $\delta^{13}\text{C}$ -DIC values reported for other rivers in the region (Lupar and Saribas Rivers in Sarawak,  $-15.7 \text{ ‰}$  to  $-11.4 \text{ ‰}$ , Müller et al., 2016; Musi, Indragiri and Siak Rivers in Indonesia,  $-22.5 \text{ ‰}$  to  $-9.0 \text{ ‰}$ , Wit, 2017).

Multiple sources and processes are likely to influence  $\delta^{13}\text{C}$ -DIC in the Rajang River. To start with, the y-intercept of the Keeling plot for freshwater samples suggests that the initial freshwater source has a  $\delta^{13}\text{C}$ -DIC of  $-18.6 \text{ ‰}$ , which is consistent with  $\delta^{13}\text{C}$  values of bicarbonate from silicate weathering with soil  $\text{CO}_2$  from C3 plants ( $-22.1$  to  $-16.1 \text{ ‰}$ , Das et al., 2005). This is consistent with the assumption that only minor amounts of carbonates, which yield  $^{13}\text{C}$ -enriched DIC (Das et al., 2005), are present in the catchment (Staub et al., 2000). Another relevant source is rainwater DIC with a typical  $\delta^{13}\text{C}$  of  $-9.3 \text{ ‰}$  (Das et al., 2005). In a river with relatively low DIC ( $177.9 \mu\text{mol L}^{-1}$ ) and large surface runoff due to heavy rain, this source term is presumably non-negligible, and a significant contribution would partially explain the relatively high  $\delta^{13}\text{C}$ -DIC values.

With regard to in-stream processes, photosynthesis increases and respiration decreases  $\delta^{13}\text{C}$ -DIC (Campeau et al., 2017). Due to the high turbidity, it can be assumed that photosynthesis in the Rajang River is negligible. In contrast, the correlation of DO and  $p\text{CO}_2$  (Fig. 5) suggests that respiration is important. This assumption is supported by the negative correlation of  $\delta^{13}\text{C}$  and DIC for freshwater samples, because with increasing DIC,  $\delta^{13}\text{C}$  values get more depleted, suggesting that organic carbon (with a  $\delta^{13}\text{C}$  of around  $-26 \text{ ‰}$  for C3 plants, Rózanski et al., 2003) is respired to  $\text{CO}_2$  within the river. However, we observed overall relatively high  $\delta^{13}\text{C}$  values. Two processes are likely to be responsible for the downstream increase in  $\delta^{13}\text{C}$ : (1) Methanogenesis and (2) evasion of  $\text{CO}_2$ . (1) Campeau et al. (2018) observed a strong relationship between  $\text{CH}_4$  concentration and  $\delta^{13}\text{C}$ -DIC in a boreal stream draining a nutrient-poor fen, suggesting that fractionation during methanogenesis leads to an increase in  $\delta^{13}\text{C}$ -DIC. As the peat soils in the Rajang delta are also anaerobic and nutrient-poor, it is likely that methanogenesis plays a role there as well. This is consistent with high reported soil  $\text{CH}_4$  concentrations of up to 1465 ppm in a peat under an oil palm plantation in Sarawak (Melling et al., 2005). It would therefore be of high interest to investigate  $\text{CH}_4$  concentrations in the Rajang River in the future. (2)  $\text{CO}_2$  evasion is also known to lead to a gradual increase of  $\delta^{13}\text{C}$ -DIC values until in equilibrium with the atmosphere (with  $\delta^{13}\text{C}$ -DIC around  $+1 \text{ ‰}$ , Polsenare and Abril, 2012; Venkiteswaran et al., 2014; Campeau et al., 2018). Due to intracarbonate equilibrium fractionation, dissolved  $\text{CO}_2$  is more depleted in  $\delta^{13}\text{C}$  than the other carbonate species. Thus, if it is removed,  $\delta^{13}\text{C}$  of the remaining DIC increases. This effect depends on pH and is more pronounced in near-neutral waters and less strong in very acidic water (Campeau et al., 2018). We sampled the lower river reaches downstream of Kapit, which corresponds to approximately the last 200 kilometers of the river. In addition, the terrain is much steeper upstream of Kapit than in the lower river reaches, so that  $\text{CO}_2$  fluxes to the atmosphere are presumably much higher due to higher turbulence. This means that a large fraction of  $\text{CO}_2$  had likely already been emitted from the river surface before reaching Kapit, leading to the observed high  $\delta^{13}\text{C}$ -DIC values.

#### 4.2.2 $p\text{CO}_2$

$p\text{CO}_2$  in the Rajang River was higher than in most Southeast Asian rivers without peat influence, such as the Mekong River ( $p\text{CO}_2= 1090 \mu\text{atm}$ , Li et al., 2013), the Red River ( $p\text{CO}_2= 1589 \mu\text{atm}$ , Le et al., 2017) or the freshwater parts of the Lupar and Saribas Rivers ( $p\text{CO}_2= 1274 \mu\text{atm}$  and  $1159 \mu\text{atm}$ , Müller et al., 2016, Table 4, Fig. 6). This could be attributed to the peat influence. However, the Rajang River has a  $p\text{CO}_2$  at the low end of values reported for peat-draining rivers (Fig. 6): Wit et al. (2015) report values between  $2400 \mu\text{atm}$  in the Batang Hari (peat coverage = 5%) and  $8555 \mu\text{atm}$  in the Siak River (peat coverage = 22%).

A meaningful comparison is also the one between the Rajang River and the Indragiri River, Indonesia, because they have a similar peat coverage (Rajang: 11%, Indragiri: 12%) and peat coverage has previously been considered as a good predictor of river  $\text{CO}_2$  emissions (Wit et al., 2015). However,  $p\text{CO}_2$  in the Indragiri ( $5777 \mu\text{atm}$ ) was significantly higher than in the Rajang, which can be attributed to a lower pH (6.3, numbers from Wit et al., 2015). To illustrate this, we ran a simple exercise using  $\text{CO}_2\text{Sys}$ . At the given temperature, salinity and pH, the  $p\text{CO}_2$  of  $5777 \mu\text{atm}$  in the Indragiri corresponds to a DIC value of  $327 \mu\text{mol L}^{-1}$ . At a hypothetical pH of 6.8, as measured in the Rajang River, this DIC value corresponds, under otherwise unchanged conditions, to a  $p\text{CO}_2$  of  $2814 \mu\text{atm}$  – which is very close to the average values measured in the peat area of the Rajang River. This shows that pH is a major determinant for a river's  $p\text{CO}_2$  (Ruiz-Halpern et al., 2015), and that the peat coverage in a river basin is insufficient as sole predictor of  $\text{CO}_2$  fluxes. Rather, pH must be taken into account as well, and its drivers must be considered.  $\delta^{13}\text{C}$ -DIC in the Indragiri was lower ( $-16.8 \text{‰}$ , Wit, 2017) than in the Rajang ( $-7.0 \text{‰}$ ), implying that respiratory  $\text{CO}_2$  is more dominant in the Indragiri, while the Rajang might be more strongly influenced by weathering, which could explain the higher pH. Note also that peat coverage is usually reported for the entire catchment (e.g., Wit et al, 2015; Rixen et al., 2016) and does not reveal how much peat is found in estuarine vs. freshwater reaches, which makes comparisons more difficult.

While DOC and  $p\text{CO}_2$  are positively related to discharge in most rivers (e.g., Bouillon et al., 2012), this pattern is sometimes reversed in peat-draining rivers. This is due to dilution, when rainfall exceeds the infiltration capacity of the wet soil and water runs off at the surface (Clark et al., 2008; Rixen et al., 2016). In the Rajang River,  $p\text{CO}_2$  was slightly higher in the non-peat area during the wet season in agreement with many non-peat-draining tropical rivers (Bouillon et al., 2012; Teodoru et al., 2014; Scofield et al., 2016). However, the seasonality of  $p\text{CO}_2$  was very small, similar to other Malaysian rivers (Müller et al., 2016) and in line with the small seasonal variability of DOC concentrations in the Rajang River (Martin et al., 2018). Note that since our data was collected during two single surveys, they represent only a snapshot and do not allow strong claims about seasonality.

Due to an El Nino event, temperatures in Southeast Asia were unusually high in late 2015 and 2016, with a temperature extreme in April 2016 prevailing in most of Southeast Asia (Thirumalai et al., 2017) and unusually hot conditions also recorded in Sibul (Fig. 2b). Given that weathering rates increase at elevated temperatures, DIC from weathering could have been enhanced over other years, although this seems unlikely because DIC was relatively low and in agreement with previous studies. Another

factor to be taken into account is that decomposition in the dry upper soil layer is more intense at higher temperatures, and with incipient rainfall, all the resultant DOC is flushed out to the river. Therefore, it is possible that during the year of our measurements, DOC concentrations were higher than usual, and respiratory CO<sub>2</sub> may therefore have been enhanced compared to other years.

#### 5 4.2.3 Impact of the peatlands on the CO<sub>2</sub> emissions from the Rajang River

The fact that *p*CO<sub>2</sub> was significantly higher in the peat area than in the non-peat area implies, at first glance, that the peat areas are a source of CO<sub>2</sub> to the river. However, this difference has to be interpreted with caution, as the entire peat area is under tidal influence (Fig. 1b). In the following, we will present several arguments that suggest that the peatlands exert only a small influence on the CO<sub>2</sub> emissions from the Rajang River.

#### 10 a) DOC

One indicator of peatland influence on a river's carbon budget is DOC. DOC concentrations in the Rajang River delta were reported to range between 1.4 mg L<sup>-1</sup> and 3 mg L<sup>-1</sup> (Martin et al., 2018). This is at the low end of the range of DOC concentrations reported for peat-draining rivers in Indonesia: These range from 2.9 mg L<sup>-1</sup> in the Musi River (peat coverage = 3.5%) up to 21.9 mg L<sup>-1</sup> in the Siak River (peat coverage = 22%; Wit et al., 2015). Rivers whose catchment area is entirely covered by peat exhibit even higher DOC concentrations, with 52 mg L<sup>-1</sup> (wet season) and 44 mg L<sup>-1</sup> (dry season) in the Sebangau River, Indonesia (Moore et al., 2011), and 44 mg L<sup>-1</sup> in the Maludam River, Sarawak, Malaysia (Müller et al., 2015). The Rajang River compares rather to rivers like Lupar and Saribas, Malaysia, which exhibit DOC concentrations of 1.8 mg L<sup>-1</sup> and 3.7 mg L<sup>-1</sup> in their freshwater parts (no peat influence, Müller et al., 2016). Consequently, DOC concentrations imply that the peatlands' influence on the Rajang's DOC is rather small.

#### 20 b) Mixing model

An alternative approach is to theoretically calculate the increase in Rajang River *p*CO<sub>2</sub> that would result from the influx from peatlands. For this, we created a simple model to simulate the mixing of two water masses (see Supplement), one with a pH of 6.8 and a *p*CO<sub>2</sub> of 2434 µatm (designed to resemble the Rajang River, non-peat area) and the other with a pH of 3.8 and a *p*CO<sub>2</sub> of 8100 µatm (designed to resemble peat-draining tributaries, based on values for the peat-draining Maludam River in Sarawak, Müller et al., 2015). For simplicity, we assumed that mixing occurs at salinity = 0 and that the temperature in both water bodies is the same (28.4°C). From these values, DIC and total alkalinity (TA) of the two water bodies were calculated using CO<sub>2</sub>Sys. Since they can be assumed to be conservative, we simply calculated DIC and TA of the mixture as

$$DIC_{S=0} = (1 - pc) \cdot DIC_1 + pc \cdot DIC_2 \text{ and } TA_{S=0} = (1 - pc) \cdot TA_1 + pc \cdot TA_2 \quad (8) \text{ and } (9),$$

where *pc* is the peat coverage in the basin (*pc*=0.11). From DIC and TA, the *p*CO<sub>2</sub> of the mixture was computed (*p*CO<sub>2</sub> = 3058 µatm). This means that if all peat-draining tributaries in the Rajang River basin had a *p*CO<sub>2</sub> of 8100 µatm and a pH of 3.8, the

$p\text{CO}_2$  in the peat area would be enhanced by around 600  $\mu\text{atm}$ . However, this increase in  $p\text{CO}_2$  is obviously gradual. For example, at the city of Sibuluan, peat coverage was estimated at around 2%, for which the described mixing model yields a  $p\text{CO}_2$  of 2548  $\mu\text{atm}$  (Fig. S3). In most parts of the delta, dilution with sea water already plays a role. Therefore, the mixing model was extended, assuming that at  $p_c=3\%$ , salinity is still zero and then linearly increases until  $p_c=11\%$ ,  $S = 32$ ,  $\text{DIC}_{S=32} = 2347$   $\mu\text{mol L}^{-1}$  and  $\text{TA}_{S=32} = 2324$   $\mu\text{mol L}^{-1}$  (two end-member mixing model, see Supplement). As a result,  $p\text{CO}_2$  would theoretically not exceed 2605  $\mu\text{atm}$  if the peat-draining tributaries were the only source of  $\text{CO}_2$  downstream of Kanowit (Fig. S3).

However,  $p\text{CO}_2$  in the peat area was 2990  $\mu\text{atm}$  (wet) and 2994  $\mu\text{atm}$  (dry) and 3005  $\mu\text{atm}$  (wet) and 2783  $\mu\text{atm}$  (dry) in the delta, so there must be another source of  $\text{CO}_2$ . Since  $p\text{CO}_2$  in Sarikei varied 2-fold with the tidal cycle, it seems likely that a large part of the difference in  $p\text{CO}_2$  between non-peat area and peat area is actually a difference between river and tidal river. Tidal variability is often seen in estuaries (e.g., Bouillon et al., 2007, Oliveira et al., 2017), largely due to conservative mixing. However, during rising tide in Sarikei, we observed  $p\text{CO}_2$  values of almost 6000  $\mu\text{atm}$ , which is higher than the freshwater end-member, suggesting that other effects also play a role. Among those are decomposition of organic matter in intertidal sediments (Alongi et al., 1999, Cai et al., 1999) and subsequent transport of the produced  $\text{CO}_2$  to the river, as well as groundwater input (Rosentreter et al., 2018).

### 15 c) $\delta^{13}\text{C-DIC}$

If the peatlands acted as a significant source of  $\text{CO}_2$  to the Rajang River, this would also have to be visible in the  $\delta^{13}\text{C-DIC}$  values. In the peat-draining Maludam River,  $\delta^{13}\text{C-DIC}$  averaged -28.55 ‰ (Müller et al., 2015). Thus, it would be expected that the influx of peat-draining tributaries to the Rajang River would decrease  $\delta^{13}\text{C-DIC}$ . This was not observed. We were therefore unable to discern a large impact of peatlands on the DIC budget of the Rajang River. It is possible that, because the peatlands are located close to the coast in this system, mixing with sea water occurs before significant effects on the  $p\text{CO}_2$  are theoretically possible. This means that not only the peat coverage in the catchment is relevant, but also how much of this peat is found in estuarine reaches. These findings support the arguments of Müller et al. (2015) and Wit et al. (2015) that material derived from coastal peatlands is swiftly transported to the ocean, explaining why peat-draining rivers may not necessarily be strong sources of  $\text{CO}_2$  to the atmosphere.

## 25 5. Conclusions

The Rajang River is a typical Southeast Asian river, transporting large amounts of terrestrial material to the South China Sea. In contrast to other Southeast Asian rivers with similar peat coverage, the impact of the peatlands on the Rajang River's  $p\text{CO}_2$  appeared to be rather small, probably due to the proximity of the peatlands to the coast. As a consequence,  $\text{CO}_2$  emissions from the Rajang River were moderate compared to other Southeast Asian rivers and low if compared to Southeast Asian peat-draining rivers.

### **Data availability**

Calibrated data used in this manuscript are available as a Supplementary Table to this manuscript. Raw data are available at the Institute of Environmental Physics, University of Bremen, Bremen, Germany.

### **Author contribution**

- 5 DMD, TW, TR and MM designed this study. DMD performed all measurements and sample preparations during the first survey, AC, AB and MM performed the measurements and sample preparations during the second survey. JO and BE analyzed the DIC and isotopic data, DMD analyzed all other data. All co-authors contributed to the interpretation and discussion of the results. DMD prepared the manuscript with contributions from all co-authors.

### **Competing interests**

- 10 The authors declare that they have no conflict of interest.

### **Acknowledgements**

- We would like to thank the Sarawak Biodiversity Centre for permission to conduct research in Sarawak waters (permit no. SBC-RA-0097-MM and export permits SBC-EP-0040-MM and SBC-EP-0043-MM). We also thank Dr. Aazani Mujahid (University of Malaysia Sarawak, Kuching, Malaysia) for her extensive organizational help in preparing the campaigns.
- 15 Further, we would like to thank all students and scientists from Swinburne University of Technology and UNIMAS in Malaysia for their help during the sampling trips. Lukas Chin and the “SeaWonder” crew are acknowledged for their support during the cruises. The study was supported by the Central Research Development Fund of the University of Bremen, the MOHE FRGS Grant (FRGS/1/2015/WAB08/SWIN/02/1), SKLEC Open Research Fund (SKLEC-KF201610) and ARC Linkage Grant LP150100519.

20



## References

- Alin, S. R., de Fátima F. L. Rasera, M., Salimon, C., I., Richey, J. E., Holtgrieve, G. W., Krusche A. V., and Snidvongs, A. Physical controls on carbon dioxide transfer velocity and flux in low-gradient river systems and implications for regional carbon budgets. *J Geophys Res* 116, G01009, doi: 10.1029/2010JG001398, 2011.
- 5 Alkhatib, M., Jennerjahn, T. C., and Samiaji, J. Biogeochemistry of the Dumai River Estuary, Sumatra, Indonesia, a Tropical Black-Water River. *Limnol Oceanogr* 52(6): 2410-2417, 2007.
- Allen, G. H. and Pavelsky, T. M. Global extent of rivers and streams. *Science* 361, 585-588, doi: 10.1126/science.aat0636, 2018.
- Alongi, D. M., Tirendi, F., Dixon, P., Trott, L. A., and Brunskill, G. J. Mineralization of Organic Matter in Intertidal Sediments  
10 of a Tropical Semi-enclosed Delta. *Estuar Coast Shelf S* 48: 451-467, 1999.
- Aufdenkampe, A. K., Mayorga, E., Raymond, P. A., Melack, J. M., Doney, S. C., Alin, S. R., Aalto, R. E., and Yoo, K. Riverine coupling of biogeochemical cycles between land, oceans, and atmosphere. *Front Ecol Environ* 9(1): 53-60. doi: 10.1890/100014, 2011.
- Borges, A. V., Vanderborght, J.-P., Schiettecatte, L.-S., Gazeau, F., Ferrón-Smith, S., Delille, B., and Frankignoulle, M.  
15 Variability of the Gas Transfer Velocity of CO<sub>2</sub> in a Macrotidal Estuary (the Scheldt). *Estuaries* 27(4): 593-603, 2004.
- Borges, A. V., Abril, G., Darchambeau, F., Teodoru, C. R., Deborde, J., Vidal, L. O., Lambert, T., and Bouillon, S. Divergent biophysical controls of aquatic CO<sub>2</sub> and CH<sub>4</sub> in the World's two largest rivers. *Nature Sci Rep* 5:15614. doi: 10.1038/srep15614, 2015.
- Bouillon, S. Middelburg, J. J., Dehairs, F., Borges, A. V., Abril, G., Flindt, M. R., Ulomi, S. and Kristensen, E., Importance  
20 of intertidal sediment processes and porewater exchange on the water column biogeochemistry in a pristine mangrove creek (Ras Dege, Tanzania). *Biogeosciences* 4: 311-322, 2007.
- Bouillon, S., Yambélé, A., Spencer, R. G. M., Gilikin, D. P., Hernes, P. J., Six, J., Merckx, R., and Borges, A. V. Organic matter sources, fluxes and greenhouse gas exchange in the Oubangui River (Congo River basin). *Biogeosciences* 9, 2045-2062. doi: 10.5194/bg-9-2045-2012, 2012.
- 25 Cai, W.-J., Pomeroy, L.R., Moran, M. A., and Wang, Y. Oxygen and carbon dioxide mass balance for the estuarine-intertidal marsh complex of five rivers in the southeastern U.S. *Limnol Oceanogr* 44(3): 639-649, 1999.
- Campeau, A., Wallin, M. B., Giesler, R., löfgren, S., Mörth, C.-M., Schiff, S., Venkiteswaran, J. J., and Bishop, K. Multiple sources and sinks of dissolved inorganic carbon across Swedish streams, refocusing the lens of stable C isotopes. *Nature Sci Rep* 7:9158. doi: 10.1038/s41598-017-09049-9, 2017.

- Campeau, A., Bishop, K., Nilsson, M. B., Klemedtsson, L., Laudon, H., Leith, F. I., Öquist, M., and Wallin, M. B. Stable carbon isotopes reveal soil-stream DIC linkages in contrasting headwater catchments. *J Geophys Res-Bioge* 123: 149-167, doi: 10.1002/2017JG004083, 2018.
- Chen, C.-T. A., Huang, T.-H., Chen, Y.-C., Bai, Y., He, X. and Kang, Y. Air-sea exchanges of CO<sub>2</sub> in the world's coastal seas. *Biogeosciences* 10: 6509-6544. doi: 10.5194/bg-10-6509-2013, 2013.
- Clark, J. M., Lane, S. N., Chapman, P. J., and Adamson, J. K. Link between DOC in near surface peat and stream water in an upland catchment. *Sci Total Environ* 404(2-3): 308-315. doi: 0.1016/j.scitotenv.2007.11.002, 2008.
- Dai, M., Yin, Z., Meng, F., Liu, Q., and Cai, W.-J. Spatial distribution of riverine DOC inputs to the ocean: an updated global synthesis. *Curr Opin Env Sust* 4: 170-178. doi: 10.1016/j.cosust.2012.03.003, 2012.
- 10 Das, A., Krishnaswami, S., and Bhattacharya, S. K. Carbon isotope ratio of dissolved inorganic carbon (DIC) in rivers draining the Deccan Traps, India: Sources of DIC and their magnitudes. *Earth Planet Sc Lett* 236: 419-429, 2005.
- Dickson, A., Sabine, C., and Christian, G. Guide to best practices for ocean CO<sub>2</sub> measurements. North Pacific Marine Science Organization (PICES) Special Publication, 191 pp., 3<sup>rd</sup> edition. [http://cdiac.ess-dive.lbl.gov/ftp/oceans/Handbook\\_2007/Guide\\_all\\_in\\_one.pdf](http://cdiac.ess-dive.lbl.gov/ftp/oceans/Handbook_2007/Guide_all_in_one.pdf) (last access: 29<sup>th</sup> June 2018), 2007.
- 15 DID: Department of Irrigation and Drainage Sarawak: Resource Centre-IRBM 22 Basins. [http://www.did.sarawak.gov.my/modules/web/pages.php?mod=webpage&sub=page&id=315&menu\\_id=0&sub\\_id=314](http://www.did.sarawak.gov.my/modules/web/pages.php?mod=webpage&sub=page&id=315&menu_id=0&sub_id=314) (last access: 29<sup>th</sup> June 2018), 2017.
- Dlugokencky, E. J., Lang, P. M., Mund, J. W., Crotwell, A. M., Crotwell, M. J., and Thoning, K. W. Atmospheric carbon dioxide dry air mole fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1968-2016, 20 Version: 2017-07-28, Path: [ftp://aftp.cmdl.noaa.gov/data/trace\\_gases/co2/flask/surface/](ftp://aftp.cmdl.noaa.gov/data/trace_gases/co2/flask/surface/) (last access: 06/13/2018), 2018.
- Dommain, R., Couwenberg, J., and Joosten, H. Development and carbon sequestration of tropical peat domes in south-east Asia: links to post-glacial sea-level changes and Holocene climate variability. *Quaternary Sci Rev* 30: 999-1010. doi: 10.1016/j.quascirev.2011.01.018, 2011.
- DWD: German Weather Service: Climate data center, path: [ftp://ftp-cdc.dwd.de/pub/CDC/observations\\_global/CLIMAT/](ftp://ftp-cdc.dwd.de/pub/CDC/observations_global/CLIMAT/) (last access: 27<sup>th</sup> Aug, 2018), Station ID 96421, 2018.
- 25 Gaveau, D. L. A., Sloan, S., Molidena, E., Yaen, H., Sheil, D., Abram, N. K., Anrenaz, M., Nasi, R., Quinones, M., Wielaard, N., and Meijaard, E. Four Decades of Forest Persistence, Clearance and Logging on Borneo. *PlosOne* 9(7): e101654. doi: 10.1371/journal.pone.0101654, 2014.


- Gaveau, D.L.A.; Salim, M.; Arjasakusuma, S.: Deforestation and industrial plantations development in Borneo, Center for International Forestry Research (CIFOR), V2; REGBorneo\_PlantedIOPP\_1973to2016\_CIFOR, doi: 10.17528/CIFOR/DATA.00049, 2016.
- Griffith, D. W. T. Synthetic calibration and quantitative analysis of gas-phase FT-IR spectra. *Applied Spectroscopy* 50: 59-70, 1996.
- Griffith, D. W. T., Deutscher, N. M., Caldw, C., Kettlewell, G., Rigggenbach, M., and Hammer, S. A Fourier transform infrared trace gas and isotope analyser for atmospheric applications. *Atmos Meas Tech*, 5:2481–2498. doi: 10.5194/amt-5-2481-2012, 2012.
- Harun, S., Dambul, R., Abdullah, M. H., and Mohamed, M. Spatial and seasonal variations in surface water quality of the Lower Kinabatangan River Catchment, Sabah, Malaysia. *Journal of Tropical Biology and Conservation* 11: 117-131. ISSN 1823-3902, 2014.
- Huang, T. H., Fu, Y. H., Pan, P. Y., and Chen, C. T. A. Fluvial carbon fluxes in tropical rivers. *Curr Opin Env Sust* 4: 162-169, doi: 10.1016/j.cosust.2012.02.004, 2012.
- Huang, T. H., Chen, C. T. A, Tseng, H. C., Lou, J. Y., Wang, S. L., Yang, L., Kandasamy, S., Gao, X., Wang, J. T., Aldrian, E., Jacinto, G. S., Anshari, G. Z., Sompongchaiyakul, P., and Wang, B. J. Riverine carbon fluxes to the South China Sea. *Journal of Geophysical Research: Biogeosciences* 122: 1239-1259. doi: 10.1002/2016JG003701, 2017.
- Johnson, J. E. Evaluation of a seawater equilibrator for shipboard analysis of dissolved oceanic trace gases. *Anal Chim Acta* 395: 119-132, 1999.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R. Jenne, R., and Joseph, D. The NCEP/NCAR 40-year reanalysis project, *Bull. Amer. Meteor. Soc.*, 77, 437-470, 1996.
- Kumagai, T., Saitoh, T. M., Sato, Y., Takahashi, H., Manfroi, O. J., Morooka, T., Kuraji, K., Suzuki, M., Yasunari, T., and Komatsu, H. Annual water balance and seasonality of evapotranspiration in a Bornean tropical rainforest. *Agr Forest Meteorol*, 128: 81–92. doi: 10.1016/j.agrformet.2004.08.006, 2005.
- Lauerwald, R., Laruelle, G. G., Hartmann, J., Ciais, P., and Regnier, P. A. G. Spatial patterns in CO<sub>2</sub> evasion from the global river network. *Global Biogeochem Cy*, 29:1–21. doi: 10.1002/2014GB004941, 2015.
- Le, T. P. Q., Dao, V. N., Rochelle-Newall, E., Garnier, J., Lu, X., Billen, G., Duong, T. T., Ho, C. T., Etcheber, H., Nguyen, T. M. H., Nguyen, T. B. N., Nguyen, B. T., Le, N. D., and Pham, Q. L. Total organic carbon fluxes of the Red River System (Vietnam). *Earth Surf Proc Land* 42: 1329-1341. doi: 10.1002/esp.4107, 2017.

- Le, T. P. Q., Marchand, C., Ho, C. T., Duong, T. T., Nguyen, H. T. M., XiXi, L. Vu, D. A., Doan, P. K., and Le, N. D. CO<sub>2</sub> partial pressure and CO<sub>2</sub> emissions along the lower Red River (Vietnam). *Biogeosciences* 15: 4799-4814, doi: 10.5194/bg-15-4799-2018, 2018.
- Lehner, B, Verdin, K., and Jarvis, A. HydroSHEDS technical documentation. World Wildlife Funds US, Washington, D. C.,  
5 1 edition. <http://hydrosheds.cr.usgs.gov> (last access: 29<sup>th</sup> June 2018), 2006.
- Lewis, E., and D. W. R. Wallace. Program Developed for CO<sub>2</sub> System Calculations. ORNL/CDIAC-105. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee, 1998.
- Li, S., Lu, X. X., and Bush, R. T. CO<sub>2</sub> partial pressure and CO<sub>2</sub> emission in the Lower Mekong River. *J Hydrol* 504: 40-56. doi: 10.1016/j.jhydrol.2013.09.024, 2013.
- 10 Ling, T.-Y., Soo, C.-L., Sivalingam, J.-R., Nyanti, L., Sim, S.-F., and Grinang, J. Assessment of the water and sediment quality of tropical forest streams in upper reaches of the Baleh River, Sarawak, Malaysia, subjected to logging activities. *J Chem-NY*, Article ID 8503931. doi: 10.1155/2016/8503931, 2016.
- Ling, T.-Y., Soo, C.-L., Phan, T.-P., Nyanti, L., Sim, S.-F., and Grinang, J. Assessment of Water Quality of Batang Rajang at Pelagus Area, Sarawak, Malaysia. *Sains Malays* 46(3): 401-411. doi: 10.17576/jsm-2017-4603-07, 2017.
- 15 Ling Lee, H., Koh, H. L. and Al'Rabia'ah, H. A. Predicting soil loss from logging in Malaysia. *GIS and Remote Sensing in Hydrology, Water Resources and Environment. Proceedings of ICGRHWE held at the Three Gorges Dam, China, September 2003. IAHS Publ.* 289, 2004.
- MacKinnon, K.: The ecology of Kalimantan. In: *The ecology of Indonesia series, Vol. 3.* Oxford University Press, Oxford 1996.
- 20 Martin, P., Cherukuru, N., Tan, A. S. Y., Sanwlani, N., Mujahid, A., and Müller, M. Distribution and cycling of terrigenous dissolved organic carbon in peatland-draining rivers and coastal waters of Sarawak, Borneo. Manuscript submitted to *Biogeosciences*, 2018.
- Melling, L., Hatano, R., and Goh, K. J. Methane fluxes from three ecosystems in tropical peatland of Sarawak, Malaysia. *Soil Biol Biochem* 37: 1445-1453, doi:10.1016/j.soilbio.2005.01.001, 2005.
- 25 Milliman, J. D., and Farnsworth, K. L.: *River Discharge to the Coastal Ocean: A Global Synthesis.* Cambridge University Press, March 2011.
- Moore, S., Gauci, V., Evans, C. D., and Page, S. E. Fluvial organic carbon losses from a Bornean blackwater river. *Biogeosciences* 8: 901-909. doi: 10.5194/bg-8-901-2011, 2011.

- Moore, S., Evans, C. D., Page, S. E., Garnett, M. H., Jones, T. G., Freeman, C., Hooijer, A., Wiltshire, A. J., Limin, S. H., and Gauci, V. Deep instability of deforested tropical peatlands revealed by fluvial organic carbon fluxes. *Nature* 493: 660–663. Doi:10.1038/nature11818, 2013.
- Müller, D., Warneke, T., Rixen, T., Müller, M., Jamahari, S., Denise, N., Mujahid, A., and Notholt, J. Lateral carbon fluxes and CO<sub>2</sub> outgassing from a tropical peat-draining river. *Biogeosciences* 12: 5967-5979. doi: 10.5194/bg-12-5967-2015, 2015.
- Müller, D., Warneke, T., Rixen, T., Müller, M., Mujahid, A., Bange, H. W., and Notholt, J. Fate of terrestrial organic carbon and associated CO<sub>2</sub> and CO emissions from two Southeast Asian estuaries. *Biogeosciences* 13: 691-705. doi: 10.5194/bg-13-691-2016, 2016.
- Nachtergaele, F., van Velthuisen, H., and Verelst, L.: Harmonized World Soil Database. Path: 10 http://www.fao.org/docrep/018/aq361e/aq361e.pdf. (last access: 29<sup>th</sup> June 2018), 2009.
- Ni, H.-G., Lu, F.-H., Luo, X.-L., Tian, H.-Y. and Zeng, E. Y. Riverine inputs of total organic carbon and suspended particulate matter from the Pearl River Delta to the coastal ocean off South China. *Mar Pollut Bull* 56: 1150-1157, doi: 10.1016/j.marpolbul.2008.02.030, 2008.
- Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat, M. I., Boutin, J., and Upstill-Goddard, R. C. In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers. *Global Biogeochem Cy* 14(1): 373-387, 2000.
- Oakes, J. M., Eyre, B. D., Ross, D. J. and Turner, S. D. Stable isotopes trace estuarine transformations of carbon and nitrogen from primary- and secondary-treated paper and pulp mill effluent. *Environ Sci Technol* 44, 7411-7417, doi: 10.1021/es101789v, 2010.
- 20 Oliveira, A. P., Cabecadas, G., and Mateus, M. D. Inorganic carbon distribution and CO<sub>2</sub> fluxes in a large European estuary (Tagus, Portugal). *Nature Sci Rep* 7: 7376. doi:10.1038/s41598-017-06758-z, 2017.
- Page, S. E., Rieley, J. O., and Banks, C. J. Global and regional importance of the tropical peatland carbon pool. *Glob Change Biol* 17: 798-818. doi: 10.1111/j.1365-2486.2010.02279.x, 2011.
- Park, J.-H., Nayna, O. K., Begum, M. S., Chea, E., Hartmann, J., Keil, R. G., Kumar, S., Lu, X., Ran, L., Richey, J. E., Sarma, 25 V.V.S.S., Tareq, S. M., Xuan, D. T., and Yu, R. Reviews and syntheses: Anthropogenic perturbations to carbon fluxes in Asian river systems – concepts, emerging trends, and research challenges. *Biogeosciences* 15: 3049-3069. doi: 10.5194/bg-15-3049-2018, 2018.
- Polsenaere, P. and Abril, G. Modelling CO<sub>2</sub> degassing from small acidic rivers using water pCO<sub>2</sub>, DIC and δ<sup>13</sup>C-DIC data. *Geochim Cosmochim Acta* 91: 220-239, doi: 10.1016/j.gca.2012.05.030, 2012.


- Raymond, P. A., and Cole, J. J. Gas exchange in rivers and estuaries: Choosing a gas transfer velocity. *Estuaries* 24 (2): 312-317, 2001.
- Raymond, P. A., Zappa, C. J., Butman, D., Bott, T. L., Potter, J., Mulholland, P., Laursen, A. E., McDowell, W. H., and Newbold, D. Scaling the gas transfer velocity and hydraulic geometry in streams and small rivers. *Limnol Oceanogr Fluids and Environments* 2: 41-53, doi: 10.1215/21573689-1597669, 2012.
- Raymond, P. A., Hartmann, J., Lauerwald, R., Sobek, S., McDonald, C., Hoover, M., Butman, D., Striegl, R., Mayorga, E., Humborg, C., Kortelainen, P., Dürr, H., Meybeck, M., Ciais, P. and Guth, P. Global carbon dioxide emissions from inland waters. *Nature*, 503:355–359. doi: 10.1038/nature12760, 2013.
- Rixen, T., Baum, A., Wit, F., and Samiaji, J. Carbon leaching from tropical peat soils and consequences for carbon balances. *Front Earth Sci* 4:74. doi: 10.3389/feart.2016.00074, 2016.
- Rosentreter, J. A., Maher, D. T., Erler, D. V., Murray, R. And Eyre, B. D. Factors controlling seasonal CO<sub>2</sub> and CH<sub>4</sub> emissions in three tropical mangrove-dominated estuaries in Australia. Under review for *Estuar Coast Shelf S*, 2018.
- Rózanski, K., Froehlich, K., and Mook, W. G. *Environmental Isotopes in the Hydrological Cycle. Principles and Applications. Vol. 3: Surface Water.* International Atomic Energy Agency and United Nations Educational, Scientific and Cultural Organization, Krakow/Vienna/Groningen, 2003.
- Ruiz-Halpern, S., Maher, D., Santos, I. and B. D. Eyre. High CO<sub>2</sub> evasion during floods in an Australian subtropical estuary downstream from a modified acidic floodplain wetland. *Limnol Oceanogr* 60 (1), 42 – 56, doi: 10.1002/lno.10004, 2015.
- Sa'adi, Z., Shahid, S., Ismail, T., Chung, E.-S., and Wang, X.-J. Distributional changes in rainfall and river flow in Sarawak, Malaysia. *Asia-Pac. J. Atmos. Sci.* 53(4): 489-500. doi: 10.1007/s13143-017-0051-2, 2017.
- Sarawak Energy: About Hydropower. Path: <http://www.sarawakenergy.com.my/index.php/hydroelectric-projects> (last access: 29<sup>th</sup> June 2018), 2013.
- Scofield, V., Melack, J. M., Barbosa, P. M., Amaral, J. H. F., Forsberg, B. R., and Farjalla, V. F. Carbon dioxide outgassing from Amazonian aquatic ecosystems in the Negro River basin. *Biogeochemistry* 129: 77-91, doi: 10.1007/s10533-016-0220-x, 2016.
- Snoussi, M., Haida, S., and Imassi, S., Effects of the construction of dams on the water and sediment fluxes of the Moulouya and the Sebou Rivers, Morocco. *Reg Environ Change* 3: 5-12. doi: 10.1007/s10113-001-0035-7, 2002.
- Staub, J. R. and Esterle, J. S. Provenance and sediment dispersal in the Rajang River delta/ coastal plain system, Sarawak, East Malaysia. *Sedimentary Geology* 85: 191-201, 1993.
- Staub, J. R. and Gastaldo, R. A. Late Quarternary Sedimentation and peat development in the Rajang River Delta, Sarawak, East Malaysia. In: F. Hasan Sidi, Nummedal, D., Imbert, P., Darman, H., and Posamentier, H. W. *Tropical Deltas of Southeast*

- Asia – Sedimentology, Stratigraphy, and Petroleum Geology. SEPM Special Publication No. 76, p. 71-87, Tulsa, Oklahoma, USA, September 2003.
- Staub, J. R., Among, H. L., and Gastaldo, R. A. Seasonal sediment transport and deposition in the Rajang River delta, Sarawak, East Malaysia. *Sediment Geol* 133: 249-264, 2000.
- 5 Teodoru, C. R., Nyoni, F. C., Borges, A. V., Darchambeau, F., Nyambe, I., and Bouillon, S. Dynamics of greenhouse gases (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O) along the Zambezi River and major tributaries, and their importance in the riverine carbon budget. *Biogeosciences* 12: 2431-2453. doi: 10.5194/bg-12-2431-2015, 2015.
- Thirumalai, K., DiNezio, P. N., Okumura, Y, and Deser, C., Extreme temperatures in Southeast Asia caused by El Nino and worsened by global warming. *Nat Commun* 8: 15531. doi: 10.1038/ncomms15531, 2017.
- 10 Venkiteswaran, J. J., Schiff, S. L. And Wallin, M. B. Large carbon dioxide fluxes from headwater boreal and sub-boreal streams. *PLOS one*, 9(7): e101756, doi:10.1371/journal.pone.0101756, 2014.
- Vörösmarty, C. J., Meybeck, M., Fekete, B., Sharma, K., Green, P., and Syvitski, J. P. M. Anthropogenic sediment retention: major global impact from registered river impoundments. *Global Planet Change* 39: 169-190, 2003.
- Wanninkhof, R. Relationship between wind speed and gas exchange over the ocean. *J of Geophys Res* 97(C5): 7373-7382,  
15 1992
- Weiss, R. Carbon dioxide in water and seawater: The solubility of a non-ideal gas. *Mar Chem* 2: 203—215, 1974.
- White, J. W. C., Vaughn, B. H. and Michel S.E., Stable Isotopic Composition of Atmospheric Carbon Dioxide (<sup>13</sup>C and <sup>18</sup>O) from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1990-2014, Version: 2015-10-26, Path: ftp://afpt.cmdl.noaa.gov/data/trace\_gases/co2c13/flask/, University of Colorado, Institute of Arctic and Alpine Research  
20 (INSTAAR), 2015.
- Wit, F. Carbon footprints of peatland degradation. PhD Thesis, University of Bremen, Bremen, Germany, 2017.
- Wit, F., Müller, D., Baum, A., Warneke, T., Setiyo Pranowo, W., Müller, M., and Rixen, T. The impact of disturbed peatlands on river outgassing in Southeast Asia. *Nat Commun* 6: 10155. doi:10.1038/ncomms10155, 2015.
- WTW GmbH: FDO 925 and FDO 925-P Operating Manual, Weilheim, Germany, 11/2012.
- 25 Yao, G., Gao, Q., Wang, Z., Huang, X., He, T., Zhang, Y., Jiao, S. and Ding, J. Dynamics of CO<sub>2</sub> partial pressure and CO<sub>2</sub> outgassing in the lower reaches of the Xijiang River, a subtropical monsoon river in China. *Sci Total Environ* 376: 255-266, doi :10.1016/j.scitotenv.2007.01.080, 2007.



5 Figure 1: Map of the Rajang basin on the island of Borneo (a) and a close-up of the basin (b) including the location of the peatlands (Nachtergaele et al., 2009), industrial oil palm plantations (Gaveau et al., 2016) and the stations during the cruise. The distributaries are marked with numbers: 1-Igan, 2-Lassa, 3-Paloh, 4-Belawai, 5-Rajang.





**Figure 2: Monthly discharge calculated for the Rajang River (a) and average temperatures (b) for the city of Sibiu (2° 20' N, 111° 50' E) for the years 2015, 2016 and the long-term average from 1992-2016. Data from DWD (2018).**




Figure 3: Salinity (interpolated) and  $p\text{CO}_2$  distribution in the Rajang River and delta during the wet season survey (a,c) and dry season survey (b,d).






Figure 4: DIC versus salinity (a),  $\delta^{13}\text{C}$  versus DIC (b) and a Keeling plot of  $\delta^{13}\text{C}$  versus  $1/\text{DIC}$  for non-peat, peat and delta samples, respectively. Freshwater samples, including those from the delta region, are circled in in panel b and c. All data refer to dry season samples.

5





Figure 5: Correlation of  $p\text{CO}_2$  versus dissolved oxygen (DO) in the wet and dry season for individual distributaries.



**Figure 6: Comparison of average  $p\text{CO}_2$  values in Southeast Asian rivers. Colors distinguish peat-draining rivers from non-peat-draining rivers. The Rajang River (this study) is highlighted in red.**

**Table 1: Definition of the different areas considered in this study.**

	<b>Non-peat</b>	<b>Peat</b>	<b>Delta</b>
<b>Description of geographical extent</b>	Source to Kanowit	Kanowit to Sibul	Sibul to coast
<b>Definition of geographical extent</b>	lon >112.1000	112.1000 ≥lon ≥111.8193, lat ≤2.2831	lon <111.8193 or lat >2.2831
<b>Water surface area (km<sup>2</sup>)</b>	167.5	34.3	553.0
<b>Corresponding catchment area (km<sup>2</sup>)</b>	43550.5	899.6	7559.5
<b>Tidal influence</b>	None	Tidal	Tidal
<b>Influence of salinity</b>	Freshwater	Freshwater	Brackish

**Table 2: Average values of measured parameters for different river reaches (mean ± 1SE) for all parameters except FCO<sub>2</sub>, which is reported as mean (minimum – maximum).**

		Non-peat	Peat	Weighted mean peat/non-peat	Delta
<b>Average salinity</b>	<b>Wet</b>	0	0	0	7.1± 8.6 (n=566)
	<b>Dry</b>	0	0	0	16.0 ± 13.5 (n=2510)
<b>Temperature (°C)</b>	<b>Wet</b>	27.4 ± 0.4 (n=152)	28.7 ± 1.2 (n=89)	27.4 ± 0.4	28.6 ± 1.0 (n=126)
	<b>Dry</b>	28.4 ± 0.6 (n=628)	29.6 ± 0.2 (n=130)	28.4 ± 0.6	30.3 ± 2.6 (n=787)
<b>SPM (mg L<sup>-1</sup>)</b>	<b>Wet</b>	188.9 ± 76.9 (n=6)	104.5 ± 16.7 (n=2)	187.2 ± 75.7	162.4 ± 88.3 (n=7)
	<b>Dry</b>	51.9 ± 12.3 (n=7)	33.1 ± 2.5 (n=2)	51.5 ± 12.1	68.3 ± 31.6 (n=22)
<b>POC (mg L<sup>-1</sup>)</b>	<b>Wet</b>	2.6 ± 0.6 (n=6)	1.8 ± 0.0 (n=2)	2.6 ± 0.6	2.9 ± 2.8 (n=7)
	<b>Dry</b>	1.1 ± 0.4 (n=7)	0.8 ± 0.0 (n=2)	1.1 ± 0.4	1.0 ± 0.4 (n=22)
<b>OC content in SPM (%)</b>	<b>Wet</b>	1.5 ± 0.4 (n=6)	1.7 ± 0.3 (n=2)	1.5 ± 0.4	1.6 ± 0.6 (n=7)
	<b>Dry</b>	2.1 ± 0.6 (n=7)	2.5 ± 0.0 (n=2)	2.1 ± 0.6	1.5 ± 0.7 (n=22)
<b>O<sub>2</sub> (%)</b>	<b>Wet</b>	81.1 ± 5.4 (n=152)	80.8 ± 9.2 (n=89)	81.1 ± 5.5	66.0 ± 6.9 (n=126)
	<b>Dry</b>	79.8 ± 3.5 (n=628)	71.6 ± 1.3 (n=130)	79.6 ± 3.5	71.2 ± 11.1 (n=787)
<b>pH</b>	<b>Wet</b>	6.7 ± 0.1 (n=6)	6.6 ± 0.1 (n=2)	6.7 ± 0.1	6.9 ± 0.5 (n=7)
	<b>Dry</b>	6.8 ± 0.0 (n=7)	6.8 ± 0.0 (n=2)	6.8 ± 0.0	7.3 ± 0.5 (n=24)
<b>DIC (μmol L<sup>-1</sup>)</b>	<b>Wet*</b>	290.8 ± 32.8* (n=5)	243.6* (n=1)	289.8 ± 32.1	338.1 ± 41.9* (n=7)
	<b>Dry</b>	177.9 ± 22.4 (n=6)	177.6 (n=1)	177.9 ± 22.4	1302.3 ± 749.2 (n=21)
<b>δ<sup>13</sup>C-DIC (‰)</b>	<b>Dry</b>	-6.99 ± 1.47 (n=6)	-7.11 (n=1)	-7.0 ± 1.47	-5.90 ± 2.96 (n=21)
<b>pCO<sub>2</sub> (uatm)</b>	<b>Wet</b>	2531 ± 188 (n=703)	2990 ± 239 (n=170)	2540 ± 189	3005 ± 1039 (n=566)
	<b>Dry</b>	2337 ± 304 (n=1259)	2994 ± 141 (n=644)	2350 ± 301	2783 ± 1437 (n=2510)
<b>FCO<sub>2</sub> (gC m<sup>-2</sup> d<sup>-1</sup>)</b>	<b>Wet</b>	1.5 (0.5-2.0)	1.8 (0.7-2.4)	1.5 (0.5-2.0)	1.8 (0.7-2.4)
	<b>Dry</b>	1.7 (0.6-2.6)	2.3 (0.8-3.5)	1.7 (0.6-2.6)	2.0 (0.7-3.0)

**Table 3: Average discharge and calculated carbon loads and emissions to the atmosphere, estimated for the months of the two surveys and the whole year 2016.**

	<b>Jan 2016</b>	<b>August 2016</b>	<b>Annual estimate 2016</b>
	<b>Peat and non-peat area</b>		
<b>Discharge</b>	4347 m <sup>3</sup> s <sup>-1</sup>	3706 m <sup>3</sup> s <sup>-1</sup>	3942 m <sup>3</sup> s <sup>-1</sup>
<b>DOC load</b>	24 ± 4 GgC/month	21 ± 6 GgC/month	269 ± 60 GgC/year
<b>POC load</b>	30 ± 7 GgC/month	11 ± 4 GgC/month	246 ± 64 GgC/year
<b>DIC load</b>	41 ± 5 GgC/month	21 ± 3 GgC/month	371 ± 46 GgC/year
<b>CO<sub>2</sub> emissions</b>	10 (3-13) GgC/month	11 (4-17) GgC/month	126 (44-181) GgC/year
	<b>Delta</b>		
<b>Discharge (m<sup>3</sup> s<sup>-1</sup>)</b>	739 m <sup>3</sup> s <sup>-1</sup>	630 m <sup>3</sup> s <sup>-1</sup>	670 m <sup>3</sup> s <sup>-1</sup>
<b>CO<sub>2</sub> emissions</b>	31 (12-41) GgC/month	34 (12-51) GgC/month	391 (144-555) GgC/year



**Table 4:  $p\text{CO}_2$ , pH and  $\text{CO}_2$  evasion of several Southeast Asian rivers flowing into the South China Sea. \*The sampling points were located outside of the peat area, so the actual peat coverage at that point was zero.**

	<b>Country</b>	<b>Catchment size (km<sup>2</sup>)</b>	<b>Discharge (m<sup>3</sup> s<sup>-1</sup>)</b>	<b>Peat coverage (%)</b>	<b>pH</b>	<b><math>p\text{CO}_2</math> (<math>\mu\text{atm}</math>)</b>	<b><math>\text{CO}_2</math> evasion (g m<sup>-2</sup> d<sup>-1</sup>)</b>	<b>Reference</b>
<b>Mekong</b>	Vietnam/Myanmar/Laos/Thailand/Cambodia	795,000	15,000	-	7.4-7.9	1090	2.3	Li et al., 2013
<b>Xijiang</b>	China	350,000	7,290	-	7.6 ± 0.2	2600	2.2-4.2	Yao et al., 2007
<b>Rajang</b>	Malaysia	52,010	3,350	11	6.8 ± 0.1	2445 ± 245	1.6 (0.5-2.6)	This study
<b>Musi</b>	Indonesia	56,931	3,050	3.5	6.9 ± 0.3	4316 ± 928	7.6 ± 3.2	Wit et al., 2015
<b>Red</b>	China/Vietnam/Laos	156,450	2,640	-	8.1	1589 ± 885	6.4 ± 0.2	Le et al., 2018
<b>Batang Hari</b>	Indonesia	44,890	2,560	5	7.1	2400 ± 18	3.9 ± 0.8	Wit et al., 2015
<b>Indragiri</b>	Indonesia	17,968	1,180	11.9	6.3 ± 0.1	5777 ± 527	10.2 ± 2.7	Wit et al., 2015
<b>Siak</b>	Indonesia	10,423	684	21.9	5.1 ± 0.5	8555 ± 528	14.1 ± 2.7	Wit et al., 2015
<b>Lupar</b>	Malaysia	6,558	490	30.5*	6.9 ± 0.3	1274 ± 148	2.0 ± 0.5	Müller et al., 2016
<b>Saribas</b>	Malaysia	1,943	160	35.5*	7.3	1159 ± 29	1.1 ± 0.9	Müller et al., 2016
<b>Maludam</b>	Malaysia	91	4	100	3.8 ± 0.2	8100 ± 500	9.1 ± 4.7	Müller et al., 2015