

Fluvial organic carbon fluxes from oil palm plantations on tropical peatland

Sarah Cook^{1, 2*}, Mick J. Whelan², Chris D. Evans³, Vincent Gauci⁴, Mike Peacock⁵,
5 Mark H. Garnett⁶, Lip Khoon Kho⁷, Yit Arn Teh⁸, Susan E. Page²

¹ Department of Engineering, University of Warwick, Coventry, CV4 7AL, UK

² Centre for Landscape & Climate Research, School of Geography, Geology and the Environment, University of Leicester, LE1 7RH, UK

10 ³ Environment Centre Wales, Centre for Ecology and Hydrology, Bangor, LL57 2UW, UK

⁴ Faculty of STEM, School of Environment Earth and Ecosystems, The Open University, Milton Keynes, MK7 6AA, UK

⁵ Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences, 750 07, Uppsala, Sweden

15 ⁶ Natural Environment Research Council Radiocarbon Facility, Rankine Avenue, Scottish Enterprise Technology Park, East Kilbride, G75 0QF, UK

⁷ Tropical Peat Research Institute, Biological Research Division, Malaysian Palm Oil Board, Bandar Baru Bangi 43000, Kajang, Selangor, Malaysia

⁸ Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen AB24 3UU, UK

20 *Correspondence to:* Sarah Cook (sc606@le.ac.uk)

Abstract

Intact tropical peatlands are dense, long-term stores of carbon. However, the future security of these ecosystems is at risk from land conversion and extensive peatland drainage. This can enhance peat oxidation and convert long-term carbon sinks into significant carbon sources. In Southeast Asia, the largest land use on peatland is for oil palm plantation agriculture. Here, we present the first annual estimate of exported fluvial organic carbon in the drainage waters of four peatland oil palm plantation areas in Sarawak, Malaysia. Total organic carbon (TOC) fluxes from the plantation second and third order drains were dominated (91%) by dissolved organic carbon (DOC) and ranged from $34.4 \pm 9.7 \text{ C m}^{-2} \text{ yr}^{-1}$ to $57.7 \pm 16.3 \text{ g C m}^{-2} \text{ yr}^{-1}$ ($\pm 95\%$ confidence interval). These fluxes represent a single year survey which was strongly influenced by an El Niño event and therefore lower discharge than usual was observed. The magnitude of the flux was found to be influenced by water table depth, with higher TOC fluxes observed from more deeply drained sites. Radiocarbon dating on the DOC component indicated the presence of old (pre-1950s) carbon in all samples collected, with DOC at the most deeply drained site having a mean age of 735 years. Overall, our estimates suggest fluvial TOC contributes ~5% of total carbon losses from oil palm plantations on peat. Maintenance of high and stable water tables in oil palm plantations appears to be key to minimising TOC losses. This reinforces the importance of considering all carbon loss pathways, rather than just CO₂ emissions from the peat surface, in studies of tropical peatland land conversion.

1. Introduction

Tropical peat carbon stocks are estimated to be 105 Gt C (105×10^{15} g; Dargie et al., 2017), with over half (57 Gt C) stored within the peatlands of Southeast Asia (Page et al., 2011a; Dargie et al., 2017). Thus, Southeast Asian peatlands and tropical peatlands as a whole contain approximately 10% and 20% of the global peat carbon stocks respectively (Page et al., 2011a; Dargie et al., 2017). Disturbance, including burning, deforestation and drainage, often associated with land-use change, is common across the peatlands of Southeast Asia, driven by strong social and economic pressures to expand agricultural, palm oil and pulpwood production to support growing populations and economic development (Miettinen et al., 2012a). Consequently, only 6% of remaining peat swamp forest areas are considered pristine (Miettinen et al., 2016), whilst carbon emissions from peatlands converted to agriculture are globally significant and increasing (Wijedasa et al., 2017).

Oil palm has played a central role in land-use change within Indonesia and Malaysia over the last few decades, driven by global consumer demand for vegetable oil-based products, and the exceptionally high productivity of oil palm compared to other oil-producing crops (Wicke et al., 2008, 2011; Schrier-Uijl et al., 2013; Gandaseca et al., 2014; Cole et al., 2015; Wijedasa et al., 2017). Over the next 30 years around 50% of the remaining peat swamp forest in Indonesia is at risk of land conversion, predominately for oil palm cultivation, despite a recent moratorium on the issuing of new concession licences for agriculture or logging in peatlands (Wijedasa et al., 2018). Peatland oil palm expansion is also prevalent within the Malaysian state of Sarawak (SarVision, 2011; Cole et al., 2015). By early 2016, nearly half (46%) of the total peatland area in Sarawak was under industrial plantations, with 96% of this area used for the cultivation of oil palm (Wetlands International, 2016).

The conversion of peat swamp forest to oil palm plantation involves a sequence of major disturbances, principally in the form of deforestation and drainage to optimise soil moisture conditions for cultivation (Hooijer et al., 2010; Page et al., 2011b; Schrier-Uijl et al., 2013). Prior to planting, peat surfaces are typically compacted using caterpillar-tracked vehicles, in order to improve the rooting stability of the palms and to help with subsequent machinery movement during harvesting (Melling & Henson, 2011). These processes alter the peat's natural hydrological and biogeochemical functions, resulting in increased peat decomposition, loss of water storage and long-term subsidence (Hooijer et al., 2010; Tonks et al. 2017). This can give rise to oxidation of soil organic matter accumulated over millennia and to significant greenhouse gas (GHG) emissions (Couwenberg et al., 2010; Hirano et al., 2012). The result is often a reversal of the peatland carbon balance; from a net sink for

atmospheric carbon to a net source (Miettinen et al., 2017). Managed land-use types now contribute to approximately 78 % of Southeast Asia's total GHG emissions related to peat oxidation (146 Mt C yr⁻¹; Miettinen et al., 2017).

5 Previous research on the effects of peat swamp forest disturbance has predominately focused on direct atmospheric GHG emissions from the peat surface (Couwenberg et al., 2010; Hooijer et al., 2010; Page et al., 2011; Hirano et al., 2012; Matysek et al., 2017). Until fairly recently, fluvial carbon losses received less attention, but more recent data suggest that this flux can represent a substantial fraction of the tropical peatland carbon balance (Moore et al., 2013; Evans et al., 2014; Rixen et al., 2016; Yupi et al., 2017). Fluvial total
10 organic carbon (TOC) is typically dominated by dissolved organic carbon (DOC), with particulate organic carbon (POC) contributing < 10% of the total flux (Moore et al., 2013; Yupi et al., 2017). Dissolved organic matter (DOM) is composed of a complex mixture of aromatic and aliphatic organic compounds, which have varying susceptibility to a range of physico-chemical and biological processes including photochemical degradation, flocculation and microbial respiration, (Cory et al., 2014; Koehler et al., 2014; Catalán et al., 2015;
15 Logue et al., 2016; Evans et al., 2017). Over 50% of the organic carbon that is leached from tropical peat is believed to be subsequently mineralised and emitted to the atmosphere as CO₂ (Wit et al., 2015). These losses represent an important potential indirect contributor to GHG emissions. The riverine transport of TOC from land to ocean also represents a significant term in the global C budget (Ciais et al., 2013) and can have substantial impacts on the biogeochemistry and ecology of coastal waters (e.g. Frigstad et al., 2013).

20 Previous attempts to quantify fluvial carbon losses from tropical peatlands include Moore et al. (2013); Gandois et al. (2013); Wit et al. (2015); Rixen et al. (2016) and Yupi et al. (2016). Moore et al. (2013) reported that losses of DOC from disturbed tropical peatlands in Indonesia were around 50% greater than those from an adjacent intact peat swamp forest. However, this research was based on a limited number of field sites (three intact sites and five degraded sites, all of which had unregulated drainage). Additional data are needed to better
25 understand the dynamics of DOC in more intensively managed peatland environments with controlled drainage systems. This includes tropical peatland oil palm plantations where fluvial carbon losses remain unquantified. In addition, existing data demonstrate that the radiocarbon content of exported DOC (DO¹⁴C) from intact tropical peat swamp forests is consistently modern (Moore et al., 2013; Gandois et al., 2013; Muller et al., 2015). DO¹⁴C data for degraded tropical peatlands are more limited, particularly for peatland oil palm plantations. Moore et al.
30 (2013) reported DO¹⁴C data from five channels in drained and deforested peatlands in Indonesia, with mean

ages of 92 to 2,260 BP, and two measurements from oil palm plantations in Peninsular Malaysia which had mean ages of around 3,200 and 4,200 years BP. These limited data clearly suggest that tropical peatland drainage releases DOC from long-term carbon stores, but are insufficient to determine whether different forms of post drainage land-use (e.g. oil palm cultivation versus abandonment) or hydrological management (e.g. regulated versus unregulated drainage) lead to different rates or age of DOC export. There is a particular need to acquire additional data from oil palm plantations as the most extensive, but currently under-represented, post-clearance land-use on tropical peat.

In this paper, we quantify fluvial TOC concentrations, drainage channel discharge and fluvial TOC losses from four peatland oil palm plantations in Sarawak, Malaysia, over the course of one year. The main objectives of the study were: (i) to quantify DOC and POC transfers in channels draining peatland oil palm plantations; (ii) to derive annual area-specific TOC flux estimates from oil palm plantations for both the wet and dry seasons and (iii) to establish the age and quality of the DOC being lost.

2. Materials and Methods

2.1 Study site

The study was conducted in two adjacent oil palm estates: Sebungan (SE) and Sabaju (SA), situated in the Malaysian state of Sarawak, northern Borneo, east of the town of Bintulu (between 3°07.81' N and 3°14.91' N and 113°18.72' E and 113°32.19' E). The climate in this region is characterised by high temperatures (around 26°C) throughout the year. Annual precipitation is typically between 3000 and 3200 mm (Environmental Impact Assessment, 2006). The annual rainfall pattern is influenced by the Northeastern (October-January) and the Southwestern (May-August) Monsoons. The former is responsible for just under half of the annual rainfall, making this period the wettest, while the latter contributes around a quarter.

The estates cover a total area of 9,614 ha. The Sebungan estate is established on an elliptical peat dome (Fig 1.) formed between two rivers (Batang Kemena and Sungai Sebungan) which provide the main regional drainage for the plantation. The maximum peat depth is 5.6 m (Environmental Impact Assessment, 2006). Oil palm planting began in 2007 (making it the oldest plantation in this study) and the total planted area is 1,648 ha. Due to the age of the plantation, the semi-mature palms provide a partially closed canopy (70% closed) and some shade for the peat surface.

The Sabaju estate is divided into four individual oil palm plantations (Sabaju 1-4; Fig 1.). These are located on an irregularly shaped peat dome, with some small mineral soil hills protruding above the peat surface. Peat coring throughout the Sabaju estate has revealed very deep peat (8 m+) within Sabaju 4, moderately deep peat in Sabaju 3 (~4 m) and shallower peat in Sabaju 1 (~2 m). Sabaju 2 was excluded from this study due to the dominance of mineral soil. Of the three oil palm plantations under investigation (Sabaju 1, 3 and 4), the youngest palms were in Sabaju 4 (1,638 ha) which was planted in 2011, followed by Sabaju 3 (1,714 ha) planted in 2010. The oldest palms are located in Sabaju 1 (2,526 ha) planted in 2008.

Plantation management on both estates is typical of other peatland oil palm plantations in this region. Artificial drainage networks have been established to lower water tables (with a target range of -60 cm to -40 cm below the peat surface to optimise palm growth). The drainage networks consist of a grid of interconnecting ditches (Fig. 2a). The edges of the planting blocks (approximately 19 - 50 ha in area) are defined by a set of roads that provide access (Fig. 2a). Each planting block contains multiple parallel “first order” field drains (ditches approximately 0.5 m deep and 1m wide) at a spacing of every 4 planting rows which feed into a central “second order” collection drain (Fig. 2b). Second order drains subsequently feed into a system of main “third order” drains, which run parallel to the edge of the planting blocks. Water from the third order drains feeds into perimeter (ring) drains, which discharge into the adjacent river network (Fig. 2b). The hydrology of these agricultural landscapes is intensively managed throughout the year. Channel gradients are low, which means that water depths and flow directions can be controlled via channel alterations (e.g. using sand bags and boards to narrow channels and or obstruct water flow) and the installation of weirs. This allows some mitigation for the effects of extreme environmental conditions e.g. drought and flooding.

2.2 Water sample collection

A mixture of third order (main) and second order (collection) drains were monitored over an approximately one-year (54 week) period, from 3rd August 2015 to 8th August 2016, in the four main plantation study areas; Sebungan, Sabaju 1, Sabaju 3 and Sabaju 4. Sampling frequency was typically every 1-3 weeks. At each sampling, two water samples (1 x DOC; 1 x POC) were collected from each of the monitored channels. Samples for DOC determination were collected in pre-rinsed 60 ml Nalgene[®] wide-neck bottles. Water samples for POC determination were collected using 500 ml plastic bottles. Water temperature (°C), pH and electrical

conductivity (EC; $\mu\text{S cm}^{-1}$) were recorded in the unfiltered water samples *in situ*, using a portable pH and EC probe (Hanna HI 9813-6).

DOC samples were filtered in a field laboratory through 0.45 μm cellulose nitrate membrane filters which were pre-rinsed with sample, using a hand-held vacuum pump, within 24 hours of collection. After filtration, water samples were stored in the dark at 4°C (for approximately 12 weeks), to ensure DOC preservation (Cook et al., 2016), before being shipped back to the UK. On return to the UK samples were analysed on a Shimadzu Total Organic Carbon analyser as non-purgeable organic carbon (NPOC), to generate measured DOC concentrations. See Supplementary Information (Text S1) for further details.

POC concentrations were determined gravimetrically. Briefly, samples were filtered through 0.4 μm Advantec glass fibre filters which were dried for 3 hours at 105°C, weighed and combusted in a furnace for a further three hours at 550°C and reweighed. The particulate organic matter (POM) concentration was calculated from the difference in the filter mass between oven drying and combustion divided by the volume filtered. This was subsequently converted into POC assuming a 50% organic carbon content (Hope et al., 1994; Moore et al., 2011).

2.3 Carbon quality

The nature of the DOC sampled was investigated using $SUVA_{254}$ (Specific Ultra-Violet Absorption). This is the ultra-violet (UV) absorbance at 254 nm normalised to sample DOC concentration (Weishaar et al., 2003).

Absorbance at 254 nm is commonly used as a surrogate for DOC aromaticity, i.e. the fraction of DOC comprised of aromatic humic substances, which absorb light in this particular part of the electromagnetic spectrum. High- $SUVA_{254}$ compounds tend to be more photodegradable and low- $SUVA_{254}$ compounds more biodegradable (Jones et al., 2016).

UV-vis absorbance was measured using a Cole-Parmer UV/visible spectrophotometer (230 VAC, 50 Hz) at 254 nm. Samples were analysed immediately after filtration. $SUVA_{254}$ values ($\text{l mg}^{-1} \text{m}^{-1}$) were calculated from

$$SUVA_{254} = 100 \cdot \frac{A_{254}}{C_{\text{DOC}}} \quad (1)$$

where A_{254} is absorbance at 254nm and C_{DOC} is the DOC concentration (mg C l^{-1}), after Weishaar et al. (2003). Water samples with very high absorbance at 254nm (> 3.0) saturated the spectrophotometer and were therefore removed from the data set ($n=1$) prior to analysis.

2.4 Hydrology

5 Drainage channel discharge was determined periodically at a number of locations under different conditions using dilution gauging (Hongve, 1987; Hudson and Fraser, 2002). Briefly, a sodium chloride tracer solution was injected into the flow and the consequent concentration change was observed at a point downstream (~ 30 m; assuming full mixing has occurred). Electrical conductivity was used as a surrogate for concentration via site-specific concentration-conductivity calibrations. This technique is considered by us to be superior to methods
10 based on cross sectional area and velocity measurements in shallow, irregular channels. The measured discharge data were used to construct rating curves (stage-discharge relationships) for each location, where stage height was measured from stage boards which were calibrated against semi-continuous recordings of water level measured at 1-hour intervals using atmosphere-corrected pressure transducers (Mini-Divers, Schlumberger, D1501) installed in stilling wells. Whenever water samples were collected, stage was noted and converted to
15 discharge using the rating curve. The rating curve equations for all measured sites, along with the standard error of the estimate (SEE) derived from the regression equations, are presented in the Supplementary Information (Table S1; Figs. S1 to S3). The stage records from the Mini-Divers were used to reconstruct a continuous record of discharge for eight stations throughout the year from 15th September 2015 until 31st August 2016.

20 Peatland water table depths (below the peat surface) were determined using dip wells constructed from 32 mm diameter PVC tubes cut to 2 m lengths. 5 mm holes were drilled down each tube, spaced at 35 mm intervals. The top of each tube was fitted with a removable cap to allow access to the tube but preventing rain and debris from entering between measurements. The bottom of each tube was fitted with a glued PVC plug to prevent sediment encroachment. A cluster of three dip wells (inserted 1.5 m into the peat from the surface, at 0.5 m intervals from
25 one another) was installed in the centre of one planting block at each plantation study site. Measurements of water table depths were made using a dipmeter (*In-situ* Rugged Water Level TAPes) in each field sampling visit, at the same time as water samples were collected.

2.5 Flux calculation

30 The annual area-specific TOC flux (J : $\text{g C m}^{-2} \text{yr}^{-1}$) for each station was calculated from

$$J = C_W \cdot R_E = \frac{\sum(C_i \cdot Q_i)}{\sum Q_i} \cdot R_E \quad (2)$$

where C_W (g C m⁻³) is the annual flow-weighted concentration, C_i (g C m⁻³) is the instantaneous (sampled) concentration of TOC (DOC+POC) on sampling date i , Q_i is the corresponding discharge at time of sampling (m³ s⁻¹) and R_E is the annual runoff (m yr⁻¹).

R_E can be calculated from measured channel discharge and catchment area:

$$R_E = \frac{\sum Q_A}{A} = \frac{\sum_h^T Q_h}{A} \quad (3)$$

where Q_A is the measured mean annual discharge (m³ yr⁻¹) estimated as the sum of hourly discharge h (Q_h ; m³ h⁻¹) over T hours in the measurement period and A is the catchment area (m²).

In principle, the catchment area for a particular drain can be estimated from the topography of plantation block areas which it serves. However, because channel gradients in peatland landscapes are low, this is subject to a high level of uncertainty, particularly for lower order drains. Consequently, we applied a water balance approach to the calculation of discharge. To do this, R_E was assumed to be the same for all plantation sites, based on the assumption that all sites were hydrologically similar in terms of the annual water balance. While this is a simplistic approach all sites had similar soil properties, topography, vegetation and management and were sufficiently close together such that they experienced very similar rainfall. By also assuming no annual change in catchment water storage (the average difference between water table depth at the beginning and end of the monitoring period was ~ 30 mm, implying that any storage changes were in the order of 10s of mm), R_E can be calculated from climate data using

$$R_E = P - ET_a \quad (4)$$

where P is the annual rainfall (mm yr⁻¹) and ET_a is the annual actual evapotranspiration (mm yr⁻¹) which was calculated from

$$ET_a = k_c \cdot ET_0 \quad (5)$$

where k_c is the so-called ‘‘crop coefficient’’ and ET_0 is the annual reference evapotranspiration rate (a standardised ET rate which assumes soil moisture is not limiting).

ET_0 (mm yr⁻¹) was calculated from average daily values of temperature, relative humidity, wind speed and net radiation flux density using the Penman Monteith equation (e.g. Monteith, 1965). Meteorological data (including P) were available over the period 15th September 2015 to 6th August 2016 from an on-site automatic weather station (Davis Vantage Pro 2). Carr (2011) reports that k_c for oil palms typically varies between 0.8 and 1 when soil moisture is unlimited in the wet season. Lower k_c values have been reported in the dry season for other areas. However, although the near surface peat layer does dry out seasonally, palm oil roots generally extend much deeper (i.e. 50-70 cm; Othman et al., 2010; Veloo et al., 2015) – into or close to the saturated zone. We therefore assumed that soil moisture availability is rarely limiting, and adopted a value of 0.9 for k_c over the whole year. This assumption was supported by water balance modelling following the method of Whelan & Gandolfi (2002), which showed that measured discharge at all Sebunggan monitoring stations could be simulated well with $ET_a \approx 0.9 ET_0$ over the whole year (data not shown).

2.6 Flux uncertainty

TOC fluxes were subject to considerable uncertainties. Specifically, in (i) DOC concentration; (ii) the mean annual runoff (R_E) and (iii) channel discharge at the time of sampling (Q_i). These were accounted for in our overall flux estimates using a Monte Carlo Simulation approach as detailed below.

2.6.1 Monte Carlo Simulation

The second order drain with the most reliable catchment area was SE 1 (2,167,300 m²). This was defined topographically during a field reconnaissance in April 2015 in which water flow directions and the drainage layout were mapped manually and then digitised in ArcGIS. A value of R_E was then calculated by applying Equation (3) for the period for which complete meteorological data was available (15th September 2015 to 6th August 2016) and for the whole year (discharge measurements were available for the whole year). Their associated error was then estimated using standard errors derived using a Monte Carlo error propagation simulation (e.g. Iman and Conover, 1980; Farrar et al., 1989) in which the error in Q_h (± 15.14 m³ h⁻¹) was assumed to be the standard error of the estimate in the rating curve for SE 1 and in which an error of 25% was arbitrarily (and conservatively) assumed for A . Briefly, values for each variable were selected randomly from probability density functions (pdfs) and employed in Equation (3) in a large number (5000) of iterations. Gaussian pdfs were used based on the assumption that the estimate of a statistic is normally distributed about the

true value (central limit theorem) with the best-estimate value assumed for the mean and SEE assumed for the standard deviation. This is described in more detail in the Supplementary Information (Text S2; Fig.S4 to S7).

The following variables were sampled from their pdfs in calculating flux uncertainty in Equation (2) for each plantation area: C_i , Q_i and R_E . Variances for Q_i and C_i were derived, respectively, from (i) the SEE values given in the regression equations for the rating curves and (ii) the error associated with the DOC concentrations obtained using the TOC analyser (assumed to be the true value) with a precision of ~5% (Graneli et al., 1996; Bjorkvald et al. 2008; Shafer et al. 2010). This is detailed in the Supplementary Information (Text S1).

2.7 Radiocarbon dating ($DO^{14}C$)

Water samples for radiocarbon dating were acquired from three second order drains within both the Sebungan and Sabaju 3 plantations (6 samples in total). Sample collection from third order (main) drains was avoided to prevent pseudo-replication (i.e. nested catchments). All samples were collected over the course of 24 hours in the wet season (April 2016) in pre-rinsed (with sample) 500 ml polypropylene bottles and filtered using a 0.7 μm glass fibre filter. These filters were pre-combusted at high temperatures (450°C) to minimise the organic matter contamination risk. The water samples were stored at 4°C for approximately one year prior to analysis. Cold storage has been shown to be a viable method for the long-term preservation of carbon isotopic signatures (Gulliver et al., 2010). Samples were analysed by accelerator mass spectrometry (AMS) at the Natural Environment Research Council facility in East Kilbride, UK in 2017. Values were expressed as %modern (m) or conventional radiocarbon ages (in years BP, where 0 BP = AD 1950 = 100 %modern):

$$\text{Age (years BP)} = -8033 \cdot \text{Ln} (m/100) \quad (6)$$

$$m = \left(\frac{A_{\text{SAMPLE}}}{A_{\text{OXALIC}}} \right) \cdot 100 \quad (7)$$

where A_{SAMPLE} is the ^{13}C -normalised radioactivity in the sample and A_{OXALIC} is the ^{13}C -normalised radioactivity in the oxalic acid international radiocarbon standard with a radioactivity equivalent to the atmosphere in 1950 (i.e. 100 %modern = year AD 1950).

2.8 Age attribution model

DOC in water samples contains a mixture of organic matter from old (^{14}C depleted) peat and recently photosynthesised (^{14}C enriched) litter and plant material (Evans et al., 2014; Campeau et al., 2017). Ascribing a ‘mean age’ to carbon fixed post-1950 is complicated by nuclear bomb testing which released a pulse of enriched ^{14}C into the atmosphere. The ^{14}C isotopic signature is, therefore, likely to reflect a mixture of both old (pre-1950’s) and new (post bomb/ ^{14}C enriched) carbon, in varying amounts. Thus, no single definitive ‘mean age’ can be ascribed to the sample (Evans et al., 2007). To address this, the age attribution model previously described by Moore et al. (2013) and Evans et al. (2014) was used to infer an indicative age distribution for the DOC in the samples. The model assumes an exponential decrease in the amount of DOC produced with increasing depth (and therefore age) within the peat profile. Each year class in the profile was assigned a ^{14}C value based on estimated atmospheric CO_2 for that year (see Evans et al., 2014) and the maximum age was set to 4,300 years BP, based on ^{14}C basal ages recorded for this region by Dommain et al. (2011). The model can be expressed as:

$$DO^{14}C = \sum_{t=1}^{t=4,300} ({}^{14}\text{CO}_{2t} \cdot e^{(-k \cdot t)}) \quad (8)$$

where $DO^{14}C$ is the measured ^{14}C of the DOC sample, t is year prior to present day, ${}^{14}\text{CO}_{2t}$ is the ^{14}C level of atmospheric CO_2 in year t and k is an exponential decay constant with a value between 0 and 1. For each sample, k was adjusted to fit measured $DO^{14}C$ using an iterative optimisation routine (Microsoft Excel Goal Seek). Modelled age distributions were summarised by aggregating year classes into the age categories 0-9, 10-49, 50-99, 100-299, 300-499, 500-699, 700-999, 1000-2999 and 3000+ years.

2.9 Bulk density and aerated carbon stocks

Bulk density (BD) was determined on four peat cores (up to 4 m in length) per plantation, extracted using a Russian corer. Subsamples of 123 cm^3 volume were taken every 15 cm down to the water table (identified using a dipmeter) and every 50 cm thereafter. A total of 47 peat samples were oven dried at 105°C for up to 120 hours (until the dry weight of the sample had stabilised) and weighed to calculate the BD. The total aerated carbon stock SOC_{air} (kg m^{-2}) for each plantation was then derived (Tiemeyer et al., 2016) as:

$$SOC_{air} = \frac{z_{WT}}{n} \cdot \sum \rho_j \cdot SOC_j \quad (9)$$

where ρ_j (g cm^3) is bulk density for sample j , SOC_j is the soil organic carbon content (g kg^{-1}) of sample j (derived from loss on ignition assuming 50% carbon), n is the number of samples collected above the water table and z_{WT} is the average annual water table depth (m).

5 2.10 Statistical analysis

Statistical analysis was performed using GraphPad Prism v7. The threshold level of statistical significance was set at a probability of 0.05 but greater significance was also noted. For multiple comparisons, one-way ANOVAs were performed. The assumptions that the data adhered to normality and homogeneity of sample variance were checked *a priori* using the Shapiro-Wilk and Bartlett tests, respectively. If significant differences between the group means were identified then a *post-hoc* test was carried out. In cases where these assumptions were not valid, the non-parametric Kruskal-Wallis test was performed together with a *post-hoc* test as above. In addition, the relationship between variables was tested using linear regression models.

3. Results

3.1 Water table depths and aerated carbon stocks

15 Mean water table depths for the individual estates are displayed in Fig. 3 and Table 1. The lowest mean water table (i.e. furthest from the peat surface) was observed in the Sebungan estate (-55 cm; Fig. 3a). The highest mean water table (i.e. closest to the peat surface) was observed in Sabaju 1 (-31 cm; Fig. 3a). The SE site displayed the greatest degree of water table variability, with depths ranging from -128 cm to + 5 cm (i.e. above the peat surface). This was closely followed by Sabaju 3, where water depths ranged from -82 cm to + 4 cm.

20 The lowest variability was observed at Sabaju 1 where the water table depth varied between -53 cm and -6 cm. Seasonal variations in water table followed monthly rainfall (Fig. 3b). A relationship between the rainfall pattern and temporal variability in the water table depth could not be drawn due to differences in the data resolution (i.e. monthly data = rainfall; weekly data = water table depth).

25 BD measurements revealed higher compaction of the near-surface peat within the Sebungan estate compared to that in the Sabaju estate; the average BD for the upper 1 m of peat in the Sebungan estate was 0.18 ± 0.01 (standard error) g cm^{-3} , compared to an average BD value for the Sabaju estate of $0.10 \pm 0.01 \text{ g cm}^{-3}$ (Table 1). Overall, BD ranged from $0.06 \pm 0.01 \text{ g cm}^{-3}$ to $0.24 \pm 0.01 \text{ g cm}^{-3}$ across all samples. Similarly, the Sebungan

estate displayed a higher overall SOC_{air} value of 45.3 kg m⁻² which was more than double the SOC_{air} values calculated for the three Sabaju estates (14.2 to 18.9 kg m⁻²; Table 1).

3.2 Catchment hydrology

5 Mean calculated totals for P , ET_0 , ET_a and R_E over the period 15th September 2015 to 6th August 2016 were 2046 mm, 1135 mm, 1021 mm and 1025 mm respectively. The value for P was much lower than the typical annual value in the region (*ca* 3000mm yr⁻¹). This was due to the influence of the 2015/2016 El Niño event characterised by extended dry periods. The calculated R_E was 1022 ± 55 mm for the period for which complete meteorological data were available and 1090 ± 147 mm yr⁻¹ for the whole year. These values match the R_E
10 estimate derived from the water balance (1025 mm), lending confidence to our estimates of TOC fluxes.

3.3 Fluvial organic carbon

Mean annual TOC concentrations ranged from 29.3 to 51.2 mg l⁻¹ and 29.6 to 49.6 mg l⁻¹ in the second order collection drains and third order main drains, respectively (Table 2). The Sebungan plantation sites displayed the
15 highest TOC concentrations and those in Sabaju 3 the lowest (Table 2). Concentrations in both the main and collection drains at Sebungan were significantly different to all the other plantation sites ($p < 0.0001$, unpaired Kruskal-Wallis). DOC was always the dominant component of TOC, accounting for 84 – 95 % of TOC (Table 2). Overall, there were no significant seasonal temporal trends in TOC concentrations, which remained relatively stable for the duration of the investigation (~ 30 mg l⁻¹ to 50 mg l⁻¹; Fig. 4 and Fig. 5). This is seen in
20 contrast to the discharge (Fig. 5). There were no systematic differences in $SUVA_{254}$ between the different estates, or between second and third order drains.

TOC fluxes were determined for all individual plantations except for three second order drains (one in Sabaju 1 and two in Sabaju 3), which experienced significant changes in their hydrological regimes as a result of
25 plantation management actions. These actions included channel blocking and widening which led to complications in estimating discharge and therefore TOC fluxes. Mean annual TOC fluxes ranged from 34.4 ± 9.7 g C m⁻² yr⁻¹ to 57.7 ± 16.3 g C m⁻² yr⁻¹ (Fig. 6; Table 2; see Supplementary Table S2 for individual site fluxes), with significantly higher fluxes ($p < 0.05$, unpaired, one-way ANOVA) recorded in the Sebungan plantation and the lowest recorded in Sabaju 3 (Fig. 6). Mean annual TOC losses were 44.7 ± 12.6 g C m⁻² yr⁻¹
30 and 42.1 ± 11.9 g C m⁻² yr⁻¹ from the second and third order drains, respectively. The two components of TOC

(DOC and POC) contributed 93 % and 7 %, respectively, on average to annual TOC yields across all plantation sites, with a slightly higher contribution from DOC to TOC in the second order drains (91 % third order vs 94 % second order).

5 3.4 $DO^{14}C$

The greatest ^{14}C -enrichment was exhibited by site SA 3.6 (102.6 % modern) and the greatest ^{14}C -depletion by site SE 4 carbon (91.3 % modern; Table 3). Conventional mean $DO^{14}C$ age was positively correlated ($p < 0.05$) with the depth of the water table and drainage intensity (Table 3; Fig. 7), but this was strongly dependent on site SE 4, which had the deepest average drainage depth and greatest ^{14}C depletion, corresponding to a mean $DO^{14}C$ age of 735 ± 37 years BP (Table 3). The other five sites were all wetter (average water tables -21 to -50 cm) and clustered within a fairly narrow $DO^{14}C$ range (99.6 – 102.6 %modern).

The fitted age attribution model (Fig. 8) suggests that the majority of DOC in all samples, other than SE 4, originates from peat with a ^{14}C age of 100 - 500 years BP. Based on its lower measured ^{14}C value, the SE 4 sample is estimated to contain a larger proportion of older peat carbon, with > 35% estimated to be derived from material with a ^{14}C age exceeding 1000 years BP.

4. Discussion

Average DOC concentrations in water draining the Sabaju oil palm plantations were lower than those reported by Moore et al. (2013) for drained tropical peatlands (*ca* 52 mg L⁻¹), whilst those from the Sebungan plantation were in line with the values reported by Moore et al. (2013). The average annual TOC fluxes (Fig. 6) for the third order (main) and second order (collection) drains were also less than TOC flux estimates for drained tropical peat swamp forests reported elsewhere (94 to 108 g C m⁻² yr⁻¹; Moore et al., 2013; Muller et al., 2015) and those reported for intact peat swamp forests in Indonesia and Sarawak, Malaysia (63 - 64 g C m⁻² yr⁻¹) by Moore et al. (2013) and Muller et al. (2015). The annual Sebungan estate TOC fluxes were significantly higher than the fluxes from the Sabaju estate (Fig. 6). The lower annual fluxes (for drained tropical peat) reported here reflect a combination of relatively low TOC concentrations (e.g. compared to Moore et al, 2013) and relatively low total annual runoff during the study period. The latter was influenced by an El Niño-driven drought event (i.e. low rainfall recorded in 2016; Fig. 3b). Since discharge varies much more than DOC concentration (Fig. 5), the temporal pattern of DOC fluxes from the study catchments is primarily controlled by discharge (Clark et al., 2007). Initial plantation development on tropical peat is often associated with the release of large pulses of

carbon due to enhanced mineralisation during the first 5 years (Hooijer et al., 2010; Page et al., 2011b). Since the plantations which were sampled here were 6 to 9 years old, these initial responses to disturbance are unlikely to have been captured. Together with the low (El Niño related) rainfall and runoff rates experienced, our flux estimates are, therefore, probably relatively conservative of fluvial TOC losses from oil palm plantations overall. Reported peat surface CO₂ emissions from oil palm plantations are typically in the range of 900 g C m⁻² yr⁻¹ (Husnain et al., 2014) to 2700 g C m⁻² yr⁻¹ (Hooijer et al., 2012). As such, the TOC fluxes reported from this study could represent an additional carbon loss equal to ~2% to 5% of total carbon emissions from oil palm plantations on peat.

10 The spatial variations in the TOC flux across the four monitored oil palm plantations (Fig. 6) were principally controlled by DOC concentrations which, in turn, appeared to be related to water table depth, with higher DOC concentrations and fluxes from the deep-drained Sebungan site compared to the shallower-drained Sabaju sites (Figs. 3 and 5; Table 2 and 3). This is broadly consistent with the conclusions of previous analyses of peatland drainage impacts on DOC loss (Evans et al., 2016), but contrasts somewhat with the assessment by Moore et al. (2013), which recorded higher DOC fluxes (principally due to higher water losses) but not higher DOC concentrations. On the other hand, Yupi et al. (2017) did record higher DOC concentrations as well as fluxes from a small drainage-affected catchment, compared to a river draining a larger, relatively intact peat swamp forest. Differences between our results and those of Moore et al. may be explained by differences between the study areas; the Moore et al. study compared highly contrasting sites (drained and deforested versus undrained natural forest) with large resulting differences in water balance. In contrast, our study compared sites with differing drainage depths within a single land-use category and consequently a more uniform water balance. On the other hand, we observed variations in $DO^{14}C$ versus drainage depth that were consistent with those observed elsewhere in both tropical and high-latitude peatlands (Evans et al., 2014) (Fig. 7 and Table 3). This clearly suggests that deeper drainage leads to the mobilisation of older peat C into DOC and is also associated with the release of more humified (high $SUVA_{254}$) material, consistent with previous findings (Olefeldt et al., 2013). In general, our observed $SUVA_{254}$ values were higher than those reported previously in runoff from intact peat swamp forests (Moore et al., 2013; Gandois et al., 2013) which can be explained by a transition from plant-derived to peat-derived DOC sources following forest clearance and drainage (Kononen et al., 2016).

The differences in watertable depth (Fig. 3) between the Sabaju and Sebungan estates are further accentuated by the differences in bulk density, potentially leading to large differences in SOC_{air} (Table 1). This suggests a denser concentration of carbon in the peat above the water table (i.e. higher SOC_{air}) which could promote higher rates of organic matter decomposition and, hence, TOC production in the Sebungan estate. However, as peat carbon content was not measured the link between peat bulk density and the TOC fluxes cannot be fully established. The differences in these bulk densities could originate from differences in the site management, plantation age or intrinsic differences in the peat characteristics between the two peat domes.

Overall, our results suggest that the clearance and drainage of peat swamp forests for oil palm plantation leads to increased loss of carbon via fluvial pathways, in addition to recognised increases in CO_2 emissions. Further, our data suggest that deeper drainage within plantations leads to greater DOC export, while the greater ^{14}C -inferred DOC age and higher SUV_{A254} indicate that this C derives from deeper within the peat profile. Together with previous studies, our results suggest that the riverine export of DOC to coastal waters from peatland regions of Southeast Asia may have increased as a result of drainage and land-conversion, with potentially profound (but as yet uncertain) impacts on coastal marine ecosystems via altered energy and nutrient supply, pH and light regime (Durako et al., 2010; Frigstad et al., 2013; Traving et al., 2017).

With regard to the management of oil palm plantations on peatland, our measurements showed that all three Sabaju plantation sites had mean water table depths that were above the Roundtable on Sustainable Palm Oil's (RSPO) target range of 40 to 60 cm below the surface, whereas the Sebungan site fell within this range. While the Sabaju plantation had lower DOC fluxes, and all sites had lower rates of DOC loss than the highly degraded peatland sites studied by Moore et al. (2013), the ^{14}C -depleted DOC measurements (relative to previous data from undrained peat swamp forests) obtained from all study sites indicate release of older stored carbon. This suggests that even RSPO-compliant plantations may still be expected to experience elevated fluvial loss of previously stored peat carbon, which is also indicative of drainage-induced C loss more generally (Evans et al., 2014). Managing oil palm plantations on peat to minimise both gaseous and fluvial carbon losses thus remains a significant challenge, requiring coordination between governments, the plantation industry and academia (Wijedasa et al., 2016). However, our results suggesting relationships between drainage depth and DOC concentration, flux and ^{14}C -inferred source indicate that any measures that enable oil palm cultivation to be

maintained at higher water levels should lead to commensurate reductions in peat carbon losses and would, therefore, lower the broader environmental impacts of oil palm cultivation.

Acknowledgments

This work was supported by the Natural Environment Research Council (NERC; grant: X402NE53), the
5 Malaysian Oil Palm Board (grant: R010913000) and the AXA Research Fund. We are grateful to the University
of Aberdeen, the University of St. Andrews, Sarawak Oil Palms Berhad for additional financial support. We thank
the NERC Radiocarbon Facility (2049.0317) for assisting with the water sample radiocarbon dating. We also
thank L.K.K and the Tropical Peat Research Institute for field assistance and support. S.C., S.E.P., M.J.W., V.G.,
and C.D.E conceived, designed and implemented the study. S.C performed the Malaysian field data collection
10 and analysed the data along with M.P. All authors discussed the results and contributed to the writing of the
manuscript. Data is available in the accompanying supporting information.

References

- Bjorkvald, L., Buffan, I., Laudon, H., Morth, C-M.: Hydrogeochemistry of Fe and Mn in small boreal streams:
The role of seasonality, landscape type and scale, *Geochimica et Cosmochimica Acta.*, 72, 2789 –
15 2804, 2008
- Campeau, A., Bishop, K., Billett, M. F., Garnett, M. H., Laudon, H., Leach, J. A., Nilsson, M. B., Öquist, M. G.
and Wallin, M. B.: Aquatic export of young dissolved and gaseous carbon from a pristine boreal fen:
implications for peat carbon stock stability, *Global Change Biol.*, 23, 5523-5536, doi:
10.1111/gcb.13815, 2017.
- 20 Carr, M.K.V.: The Water Relations and Irrigation Requirements of Oil Palm (*Elaeis Guineensis*): A Review.
Experimental Agriculture., 47, 629-652, doi:10.1017/s0014479711000494, 2011.
- Catalan, N., Marce, R., Kothawala, D.N., Tranvik, L.J.: Organic carbon decomposition rates controlled by water
retention time across inland waters. *Nature Geosci.*, 9, 501-504, 2016.
- Clark, J.M., Lane, S.N., Chapman, P.J., Adamson, J.K.: Export of dissolved organic carbon from an upland
25 peatland during storm events: Implications for flux estimates. *J Hydrology.*, 347, 438-447, 2007.
- Cole, L.E.S., Bhagwat, S.A., Willis, K.J.: Long-term disturbance dynamics and resilience of tropical peat
swamp forests. *J Ecol.*, 103, 16-30, doi: 10.1111/1365-2745.12329, 2015.

- Cook, S., Peacock, M., Evans, C.D., Page, S.E., Whelan, M., Gauci, V., Khoon, K.L.: Cold storage as a method for the long-term preservation of tropical dissolved organic carbon (DOC). *Mires and Peat.*, 18, 1-8, doi: 10.19189/MaP.2016.OMB.249, 2016
- Cory, R.M., Ward, C.P., Crump, B.C., Kling, G.W.: Sunlight controls water column processing of carbon in arctic fresh waters. *Science*, 345., 925-928, doi:10.1126/science.1253119, 2014
- Couwenberg, J., Dommain, R., Joosten, H.: Greenhouse gas fluxes from tropical peatlands in south-east Asia. *Global Change Biol.*, 16, 1715-1732, doi: 10.1111/j.1365-2486.2009.02016.x, 2010
- Dargie, G.C., Lewis, S.L., Lawson, I.T., Mitchard, E.T.A., Page, S.E., Bocko, Y.E., Ifo, S.A.: Age, extent and carbon storage of the central Congo Basin peatland complex. *Nature.*, 542, 86-90, 2017
- Dommain, R., Couwenberg, J., Joosten, H.: Development and carbon sequestration of tropical peat domes in south-east Asia: links to post-glacial sea-level changes and Holocene climate variability. *Quaternary Sci Rev.*, 30, 999-1010, doi:10.1016/j.quascirev.2011.01.018, 2011.
- Durako, M. J., Kowalczyk, P., Mallin, M. A., Cooper, W. J., Souza, J. J., and Wells, D. H.: Interannual Variation in Photosynthetically Significant Optical Properties and Water Quality in a Coastal Blackwater River Plume, *Estuaries Coasts*, 33, 1430-1441, doi:10.1007/s12237-010-9302-5, 2010.
- Evans, C.D., Freeman, C., Cork, L.G., Thomas, D.N., Reynolds, B., Billett, M.F., Garnett, M.H., Norris, D.: Evidence against recent climate-induced destabilisation of soil carbon from C-14 analysis of riverine dissolved organic matter. *Geophys Res Lett.*, 34, doi:10.1029/2007GL029431, 2007.
- Evans, C.D., Futter, M.N., Moldan, F., Valinia, S., Frogbrook, Z., Kothawala, D.N.: Variability in organic carbon reactivity across lake residence time and trophic gradients. *Nature Geosci.*, 10, 832-835, doi: 10.1038/ngeo3051, 2017.
- Evans, C.D., Renou-Wilson, F., Strack, M.: The role of waterborne carbon in the greenhouse gas balance of drained and re-wetted peatlands. *Aquat Sci.*, 78, 573-590, 2016
- Evans, C.D., Page, S.E., Jones, T., Moore, S., Gauci, V., Laiho, R., Hruška, J., Allott, T.E.H., Billett, M.F., Tipping, E., Freeman, C., Garnett, M.H.: Contrasting vulnerability of drained tropical and high-latitude peatlands to fluvial loss of stored carbon. *Global Biogeochemical Cy.*, 28, 1215-1234, doi.org/10.1002/2013GB004782, 2014.
- Farrar, D., Allen, B., Crump, K., Shipp, A.: Evaluation of Uncertainty in Input Parameters to Pharmacokinetic Models and the Resulting Uncertainty in Output. *Toxicol Lett.*, 49, 371-385, 1989.

- Frigstad, H., Andersen, T., Hessen, D.O., Jeansson, E., Skogen, M., Naustvoll, L-J., Miles, M.W., Johannessen, T., Bellerby, R.G.J.: Long term trends in carbon, nutrients and stoichiometry in Norwegian coastal waters: Evidence of regime shift. *Prog Oceanogr.*, 111, 113-124, 2013.
- Gandaseca, S., Salimin, M.I., Ahmed, O.H.: Effect of cultivation in different age's oil palm plantation on selected chemical properties of peat swamp soils. *Agriculture, Forestry and Fisheries*, 3, 6-9, 2014.
- Gandois, L., Cobb, A.R., Hei, I.C., Lim, L.B.L., Abu Salim, K., Harvey, C.F.: Impact of deforestation on solid and dissolved organic matter characteristics of tropical peat forests: implications for carbon release. *Biogeochemistry.*, 114, 183-199, doi:10.1007/s10533-012-9799-8, 2013.
- Graneli, W., Lindell, M., Tranvik, L.: Photo-oxidative production of dissolved inorganic carbon in lakes of different humic content. *Limnol Oceanogr.*, 41, 698-706, 1996.
- Gulliver, P., Waldron, S., Scott, E.M., Bryant, C.L.: The Effect of Storage on the Radiocarbon, Stable Carbon and Nitrogen Isotopic Signatures and Concentrations of Riverine Dom. *Radiocarbon.*, 52, 1113-1122, 2010.
- Hirano, T., Segah, H., Kusin, K., Limin, S., Takahashi, H., Osaki, M.: Effects of disturbances on the carbon balance of tropical peat swamp forests. *Global Change Biol.*, 18, 3410-3422, doi: 10.1111/j.1365-2486.2012.02793.x, 2012.
- Hongve, D.: A Revised Procedure for Discharge Measurement by Means of the Salt Dilution Method. *Hydrological Processes*, 1, 267-270, 1987.
- Hooijer, A., Page, S., Canadell, J.G., Silvius, M., Kwadijk, J., Wosten, H., Jauhiainen, J.: Current and future CO₂ emissions from drained peatlands in Southeast Asia. *Biogeosciences.*, 7, 1505-1514, doi:10.5194/bg-7-1505-2010, 2010.
- Hooijer, A., Page, S., Jauhiainen, J., Lee, W.A., Lu, X.X., Idris, A., Anshari, G.: Subsidence and carbon loss in drained tropical peatlands. *Biogeosciences.*, 9, 1053-1071, doi:10.5194/bg-9-1053-2012, 2012.
- Hope, D., Billett, M.F., Cresser, M.S.: A Review of the Export of Carbon in River Water - Fluxes and Processes. *Environ Pollut*, 84, 301-324, 1994.
- Hribljan, J.A., Kane, E.S., Pypker, T.G., Chimner, R.A.: The effect of long-term water table manipulations on dissolved organic carbon dynamics in a poor fen peatland. *J Geophys Res-Biogeophys.*, 119, 577-595, doi: 10.1002/2013JG002527, 2014.
- Hudson, R., and Fraser, J.: Introduction to Salt Dilution Gauging for Streamflow Measurement Part IV: The Mass Balance (or Dry Injection) Method. *Streamline Watershed Management Bulletin.*, 9, 6-12, 2002.

- Husnain, H., Wigena, I.G.P., Dariah, A., Marwanto, S., Setyanto, P., Agus, F.: CO₂ emissions from tropical drained peat in Sumatra, Indonesia. *Mitig Adapt Strat Gl.*, 19, 845-862, 2014.
- Iman, R. L., and Conover, W. J.: Small Sample Sensitivity Analysis Techniques for Computer Models, with an Application to Risk Assessment. *Commun. Stat. Part A, Theory Meth.* 17, 749-1 842, 1980.
- 5 Jones, T.G., Evans, C.D., Jones, D.L., Hill, P.W., Freeman, C.: Transformations in DOC along a source to sea continuum; impacts of photo-degradation, biological processes and mixing. *Aquat Sci.* 78, 433-446, 2016.
- Koehler, B., Landelius, T., Weyhenmeyer, G.A., Machida, N., Tranvik, L.J.: Sunlight-induced carbon dioxide emissions from inland waters. *Global Biogeochemical Cy.*, 28, 696-711, 2014.
- 10 Kononen, M., Jauhiainen, J., Laiho, R., Spetz, P., Kusin, K., Limin, S., Vasander, H.: Land use increases the recalcitrance of tropical peat. *Wetl Ecol and Manag.*, 24, 717-731, 2016.
- Logue, J.B., Stedmon, C.A., Kellerman, A.M., Nielsen, N.J., Andersson, A.F., Laudon, H., Lindstrom, E.S., Kriberg, E.S.: Experimental insights into the importance of aquatic bacterial community composition to the degradation of dissolved organic matter. *Isme J.*, 10, 533-545, 2016.
- 15 Matysek, M., Evers, S., Samuel, M.K., Sjogersten, S.: High heterotrophic CO₂ emissions from a Malaysian oil palm plantations during dry-season. *Wetl Ecol and Manag.*, 26, 415-424, 2017.
- Melling, L., and Henson, I.E.: Greenhouse Gas Exchange of Tropical Peatlands - a Review. *J Oil Palm.*, 23, 1087-1095, 2011.
- Miettinen, J., Hooijer, A., Shi, C., Tollenaar, D., Vernimmen, R., Liew, S.C., Malins, C., Page, S.E.: Extent of
20 industrial plantations on Southeast Asian peatlands in 2010 with analysis of historical expansion and future projections. *GCB Bioenergy.*, 4, 908-918, 2012a.
- Miettinen, J., Hooijer, A., Tollenaar, D., Page, S., Malins, C., Vernimmen, R., Chi, C., Liew, S.C.: Historical Analysis and Projection of Oil Palm Plantation Expansion on Peatland in Southeast Asia, Washington DC, International Council on Clean Transportation, 2012b.
- 25 Miettinen, J., Hooijer, A., Vernimmen, R., Liew, S.C., Page, S.E.: From carbon sink to carbon source: extensive peat oxidation in insular Southeast Asia since 1990. *Environ Res Lett.*, 12, 2017.
- Miettinen, J., Shi, C.H., Liew, S.C.: Deforestation rates in insular Southeast Asia between 2000 and 2010. *Global Change Biol.*, 17, 2261-2270, doi: 10.1111/j.1365-2486.2011.02398x, 2011.

- Miettinen, J., Shi, C.H., Liew, S.C.: Land cover distribution in the peatlands of Peninsular Malaysia, Sumatra and Borneo in 2015 with changes since 1990. *Glob Ecol Conserv.*, 6, 67-78, doi: 10.1016/j.gecco.2016.02.004, 2016.
- 5 Monteith, J.L.: Evaporation and environment. *Proceedings of Symposium on Environmental Biology.*, 19, 205–234, 1965.
- Moore, S., Evans, C.D., Page, S.E., Garnett, M.H., Jones, T.H., Freeman, C., Hooijer, A., Wiltshire, A., Limin, S., Gauci, V.: Deep instability of deforested tropical peatlands revealed by fluvial organic carbon fluxes. *Nature.*, 493, 660-664, 2013.
- Moore, S., Gauci, V., Evans, C.D., Page, S.E.: Fluvial organic carbon losses from a Bornean blackwater river. *Biogeosci.*, 8, 901-909, 2011.
- 10 Muller, D., Warneke, T., Rixen, T., Muler, M., Jamahari, S., Denis, S., Mujahid, A., Notholt, J.: Lateral carbon fluxes and CO₂ outgassing from a tropical peat-draining river. *Biogeosci.*, 12, 5967-5979, 2015.
- Olefeldt, D., Roulet, N., Giesler, R., Persson, A.: Total waterborne carbon export and DOC composition from ten nested subarctic peatland catchments—importance of peatland cover, groundwater influence, and inter-annual variability of precipitation patterns. *Hydrol Process.*, 27, 2280-2294, 2013.
- 15 Othman, H., Mohammed, A.T., Harun, M.H., Darus, F.M., Mos, H.: Best management practises for oil palm planting on peat: optimum groundwater table. *MPOB Information Series.*, 528, 2010.
- Page, S.E., Rieley, J.O., Banks, C.J.: Global and regional importance of the tropical peatland carbon pool. *Global Change Biol.*, 17, 798-818, 2011a.
- 20 Page, S.E., Morrison, R., Malins, C., Hooijer, A., Rieley, J. O. & Jauhiainen, J.: Review of peat surface greenhouse gas emissions from oil palm plantations in Southeast Asia (ICCT White Paper 15). International Council on Clean Transportation, Washington, 2011b.
- Rixen, T., Baum, A., Wit, F., Samiaji, J.: Carbon Leaching from Tropical Peat Soils and Consequences for Carbon Balances. *Front Earth Sci.*, 4, 2016.
- 25 Shafer, M.M., Perkins, D.A., Antkiewicz, D.S., Stone, E.A., Qurasishi, T.A., Schauer, J.J.: Reactive oxygen species activity and chemical speciation of size-fractionated atmospheric particulate matter from Lahore, Pakistan: an important role for transition metals. *J Environ Monitor.*, 12, 704-715, 2010.
- Tiemeyer, B., Borraz, E.A., Augustin, J., Bechtold, M., Beetz, S., Beyer, C., Drosler, M., Elbi, M., Eickenscheidt, T., Fiedler, S., Forster, C., Freibauer, A., Giebels, M., Glatzel, S., Heinichen, J., Hoffman, M., Hoper, H., Jursinki, G., Leiber-Sauheitl, K., Peichl-Brak, M., Robkof, R., Sommer, M.,
- 30

- Zeitz, J.: High emissions of greenhouse gases from grasslands on peat and other organic soils. *Global Change Biol.*, 22, 4134-4149, 2016.
- 5 Tipping, E., Corbishley, H.T., Koprivnjak, J.F., Lapworth, D.J., Miller, M.P., Vincent, C.D., Hamilton-Taylor, J.: Quantification of natural DOM from UV absorption at two wavelengths. *Environ Chem.*, 6, 472-476, 2016.
- Tonks, A.J., Aplin, P., Beriro, D.J., Cooper, H., Evers, S., Vane, C.H., Sjogersten, S.: Impacts of conversion of tropical peat swamp forest to oil palm plantation on peat organic chemistry, physical properties and carbon stocks. *Geoderma.*, 289, 36–45, 2017.
- 10 Traving, S.J., Rowe, O., Jakobsen, N.M., Sorensen, H., Dinasquet, J., Stedmon, C.A., Andersson, A., Riemann, L.: The effect of increased loads of dissolved organic matter on estuarine microbial community composition and function. *Front. Microbiol.*, 8, 351, doi: 10.3389/fmicb.2017.00351, 2017.
- Veloo, R., van Ranst, E., Selliah, P.: Peat characteristics and its impact on oil palm yield. *NJAS – Wagen J Life Sc.*, 72-73, 33-40, 2015.
- 15 Weishaar, J.L., Aiken, G.R., Bergamaschi, B.A., Fram, M.S., Fujii, R., Mopper, K.: Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon. *Envir Sci Tech.*, 37, 4702-4708, 2003.
- Whelan, M.J. and Gandolfi, C.: Modelling of spatial controls on denitrification at the landscape scale. *Hydrol Process.*, 16, 1437 – 1450, 2002.
- 20 Wicke, B., Dornburg, V., Junginger, M., Faaij, A.: Different palm oil production systems for energy purposes and their greenhouse gas implications. *Biomass Bioenerg.*, 32, 1322-1337, 2008.
- Wijedasa, L.S., Page, S.E., Evans, C.D., Osaki, M.: Time for responsible peatland agriculture. *Science.*, 354, 562-562, 2016.
- Wijedasa, L.S., Jauhiainen, J., Kononen, M et al.: Denial of long-term issues with agriculture on tropical peatlands will have devastating consequences. *Global Change Biol.*, 23, 977-982, 2017.
- 25 Wijedasa, L.S., Sloan, S., Page, S.E., Clements, G.R., Lupascu, M., Evans, T.A: Carbon emissions from Southeast Asia peatlands will increase despite emission-reduction schemes. *Global Change Biol.*, doi: 10.1111/gcb.14340, 2018.
- Wit, F., Muller, D., Baum, A., Warneke, T., Pranowo, W.S., Muller, M., Rixen, T.: The impact of disturbed peatlands on river outgassing in Southeast Asia. *Nat Com.*, 6, 10155, 2017.

Yule, C.M.: Loss of biodiversity and ecosystem functioning in Indo-Malayan peat swamp forests. *Biodivers Conserv.*, 19, 393-409, 2010.

Yupi, H.M., Inoue, T., Bathgate, J., Putra, R.: Concentrations, loads and yields of organic carbon from two tropical peat swamp forest streams in Riau Province, Sumatra, Indonesia. *Mires and Peat*, 18, 2016.

Tables

Table 1. Mean water table, bulk density and SOC_{air} measurements for the four oil palm plantations; \pm represents the standard error of the mean.

Plantation	Mean water table depth (cm)	Average bulk density (g cm^{-3})		SOC_{air} (kg m^{-2})
		Whole core	Top 1m of peat	
Sabaju 1	31.3 \pm 1.1	0.095 \pm 0.008	0.102 \pm 0.009	14.7 \pm 1.6
Sabaju 3	35.1 \pm 1.8	0.081 \pm 0.005	0.098 \pm 0.007	14.2 \pm 2.3
Sabaju 4	40.9 \pm 1.7	0.092 \pm 0.005	0.103 \pm 0.012	18.9 \pm 1.2
Sebungan	55.3 \pm 3.3	0.165 \pm 0.008	0.182 \pm 0.007	45.3 \pm 2.1

Table 2. Fluvial organic carbon data for the monitored oil palm plantations. DOC, POC and TOC concentrations and $SUVA_{254}$ are shown as site mean \pm standard error of the mean. Annual TOC fluxes are shown as site mean \pm the 95% confidence interval (CI; standard error x 1.96) which encompasses the propagated error associated with uncertainty in the DOC concentration, discharge and annual runoff derived from the Monte Carlo Simulation. $SUVA_{254}$ values are means of samples collected for each drain type in each plantation throughout the sampling year.

Plantation	Drain type	No. of channels	Mean DOC concentration (mg l ⁻¹)	Mean POC concentration (mg l ⁻¹)	Mean TOC concentration (mg l ⁻¹)	Annual TOC flux (g C m ⁻² yr ⁻¹)	$SUVA_{254}$ (L mg-C ⁻¹ m ⁻¹)
Sabaju 1	Collection (2 nd order)	2	31.1 \pm 1.1	2.7 \pm 0.3	33.8 \pm 1.1	38.2 \pm 10.8	5.2 \pm 0.1
	Main (3 rd order)	1	29.7 \pm 0.9	3.9 \pm 0.5	33.6 \pm 0.8	36.8 \pm 10.5	5.2 \pm 0.1
Sabaju 3	Collection (2 nd order)	2	25.2 \pm 0.8	4.1 \pm 0.3	29.3 \pm 0.7	36.4 \pm 10.3	5.4 \pm 0.1
	Main (3 rd order)	1	25.0 \pm 1.0	4.6 \pm 0.5	29.6 \pm 1.1	34.5 \pm 9.7	5.1 \pm 0.1
Sabaju 4	Collection (2 nd order)	2	34.1 \pm 0.6	2.5 \pm 0.3	36.7 \pm 0.6	42.1 \pm 11.9	5.0 \pm 0.1
	Main (3 rd order)	2	35.3 \pm 0.6	2.7 \pm 0.3	38.0 \pm 0.7	43.1 \pm 12.2	5.6 \pm 0.1
Sebungan	Collection (2 nd order)	3	48.2 \pm 0.8	3.4 \pm 0.4	51.2 \pm 1.0	56.3 \pm 15.9	5.6 \pm 0.2
	Main (3 rd order)	1	47.1 \pm 0.8	2.5 \pm 0.2	49.6 \pm 0.8	53.0 \pm 15.0	6.4 \pm 0.1

Table 3. Mean and standard errors for radiocarbon $DO^{14}C$ expressed as %modern and in conventional radiocarbon years (years BP, relative to CE 1950), expressed at $\pm 1\sigma$ level, for individual sample sites across the Sebungan (SE 2, SE 3, SE 4) and Sabaju 3 (SA 3.1, SA 3.3, SA 3.6) estates. Mean ^{14}C levels $>100\%$ modern cannot be assigned an age and are subsequently referred to as ‘modern’. Mean annual water table depth data for each site are also presented, along with the maximum and minimum water table depths recorded and the % of time the water table was more than 60 cm from the peat surface. Negative numbers indicate distance below the peat surface.

Plantation	Sample site	^{14}C (%modern)	$DO^{14}C$ Age (years BP)	Water table depth (cm)			% of time watertable was below – 60 cm
				mean	maximum	minimum	
Sebungan	SE 2	100.10 \pm 0.46	modern	-21	5	-58	0 %
	SE 3	99.57 \pm 0.46	35 \pm 37	-52	-34	-89	27 %
	SE 4	91.26 \pm 0.42	735 \pm 37	-92	-65	-128	100 %
Sabaju 3	SA 3.1	102.37 \pm 0.47	modern	-30	4	-56	0 %
	SA 3.3	99.63 \pm 0.44	30 \pm 35	-39	4	-82	21 %
	SA 3.6	102.63 \pm 0.47	modern	-36	-14	-72	3 %

Figures




Fig. 1. Location of the Sebungan and Sabaju oil palm estates, in Sebauh Bintulu district Sarawak, Borneo. The estates are bordered by a network of rivers (grey and white lines) namely the Batang Kemena, Sungai Sebungan, S. Batang and S.Pandan. Arrows indicate direction of water flow.




Fig. 2. Example of **(a)** peatland oil palm plantation (Sebungan estate) layout with the planting blocks, drains and roads highlighted and **(b)** close up schematic of typical drainage set up on a peat oil palm planting block. Red box = planting area, black dashed line = first order field drains, solid black line = second order collection drain, solid blue line = third order main drain, blue dashed line = perimeter drain. Arrows show the prevailing direction of water flow




Fig. 3 Water table depths measured in dip wells in all monitored areas, presented as **(a)** box plots showing mean water table depths for all monitored plantations over the study period (August 2015-September 2016) (central horizontal line), along with the minimum and maximum depths recorded over the entire study period (bars). Box = 75th and 25th percentiles, positioned in sequence from highest to lowest watertable and **(b)** as a time-series of mean weekly water table depths. Negative values indicate that water table was below the peat surface, positive values indicate that there was standing water above the peat surface. Monthly rainfall data were obtained from the rainfall gauge at the Sebungan plantation base (August 2015 –August 2016).




Fig. 4 Weekly TOC concentration data for the study plantations, alongside monthly rainfall. Data presented are mean weekly TOC concentrations from all drains within each site. Monthly rainfall data were obtained from the rain gauge at the Sebungan plantation base (August 2015 –August 2016).




Fig. 5. Weekly TOC concentration data for the study plantations, alongside the mean discharge (Q , $m^3 s^{-1}$) for the monitored main and collection drains within each plantation estate (August 2015 –August 2016).




Fig. 6. Mean annual TOC fluxes for the plantation drains. Error bars represent \pm the 95% confidence interval (CI; standard error \times 1.96) which encompasses the propagated error associated with uncertainty in the DOC concentration, discharge and annual runoff derived from the Monte Carlo Simulation. Fluxes are separated into the different drain types; third order (main) and second order (collection). Horizontal bar lines represent contribution of DOC (bottom segment) and POC (top segment) to the overall TOC flux. Estates are presented in sequence of drainage intensity. Letters ‘a’ to ‘b’ denote significant differences ($p < 0.05$, unpaired, one-way ANOVA) across the study sites irrespective of drain type.




Fig. 7 Mean water table depths plotted against ^{14}C (%modern) for all six sites in the Sebungan (SE 2, SE 3, SE 4) and Sabaju 3 (SA 3.1, SA 3.3, SA 3.6) estates. Negative numbers denote distance below the peat surface.




Fig. 8. Modelled age distributions of $DO^{14}C$, as estimated from the age attribution model for all radiocarbon dated water samples ($n=6$) in the Sebungan (SE 2, SE 3, SE 4) and Sabaju 3 (SA 3.1, SA 3.3, SA 3.6) estates.