

Technical Note: Isotopic corrections for the radiocarbon composition of CO₂ in the soil gas environment must account for diffusion and diffusive mixing

Jocelyn E. Egan¹, David R. Bowling², David A. Risk³

¹Department of Earth Sciences, Dalhousie University, Halifax, Nova Scotia, B3H 4R2, Canada

²School of Biological Sciences, University of Utah, Salt Lake City, Utah, 84112, USA

³Department of Earth Sciences, St. Francis Xavier University, Antigonish, Nova Scotia, B2G 2W5, Canada

Correspondence to: Jocelyn E. Egan (jocelyn.egan@dal.ca)

Abstract. Earth system scientists working with radiocarbon in organic samples use a stable carbon isotope ($\delta^{13}\text{C}$) correction to account for mass-dependent fractionation, but it has not been evaluated for the soil gas environment, where both diffusive gas transport and diffusive mixing are important. Using theory and an analytical soil gas transport model, we demonstrate that the conventional correction is inappropriate for interpreting the radio-isotopic composition of CO₂ from biological production, because it does not account for important gas transport mechanisms. Based on theory used to interpret $\delta^{13}\text{C}$ of soil production from soil CO₂, we propose a new solution for radiocarbon applications in the soil gas environment that fully accounts for both mass-dependent diffusion and mass-independent diffusive mixing.

1 Introduction

Radiocarbon allows us to measure the age of *soil-respired CO₂* (CO₂ diffusing from the soil surface to the atmosphere, aka soil flux as in Cerling et al., 1991), but the traditional reporting convention for radiocarbon was not established for soil gas-phase sampling, but rather for solid (organic matter) sample analysis. The validity of this convention has never been explicitly tested for soil-respired CO₂.

The traditional radiocarbon reporting convention, $\Delta^{14}\text{C}$ (Stuiver and Polach, 1977), uses a mass-dependent correction based on the isotopic composition of wood. Its purpose is to correct for biochemical fractionation against the radiocarbon isotopologue (¹⁴CO₂) abundance during photosynthesis, which is assumed to be twice as strong as for ¹³CO₂ based on their respective departures in molecular mass from ¹²CO₂. The classical reference describing these conventional calculations is Stuiver and Polach (1977).

In the soil gas environment, researchers have different implicit expectations for fractionation processes. They generally assume that ¹⁴C of CO₂ is *not* biochemically fractionated in the gas phase, between the points of CO₂ *production* (biological production of CO₂ by soil organisms and roots) and measurement (subsurface or flux chamber samples). This assumption is

reasonable based on the short residence time of CO₂ (minutes to days) in the soil profile before emission to the atmosphere. However, soil gas isotopic signatures depart in predictable ways from the signature of production because of physical fractionation. It has been recognized for decades that δ¹³C of CO₂ at any point in the soil profile will never equal the isotopic signature of production, because of transport fractionations that alter produced CO₂ before it is measured (Cerling et al., 1991). This theory readily extends to ¹⁴C.

We argue here, that in the case of soil pore space ¹⁴C where mixing of air masses occurs, that the assumption that mass-dependent fractionation is twice as large for ¹⁴C as δ¹³C by biochemical and physical processes no longer holds. Using theory and physical modelling of soil gas transport to illustrate the issue with the current reporting convention correction, we propose an alternative approach for specific use cases.

2 Theory

To understand why the mass-dependent correction used in the Stuiver and Polach (1977) radiocarbon reporting convention may be a poor fit for soil gas studies, we can look at our current understanding of the stable isotopic composition, δ¹³C, of soil CO₂ (pore space CO₂, mole fraction with respect to dry air). We use delta notation to present the stable isotopic composition of CO₂:

$$\delta^{13}\text{C} = \left(\frac{R_s}{R_{\text{VPDB}}} - 1 \right) 1000, \quad (1)$$

where δ¹³C is the isotopic composition in ‰ (see Table 1 for a full list of abbreviations), R_s is the ¹³C/¹²C ratio of the sample, and R_{VPDB} is the ¹³C/¹²C ratio of the international standard, Vienna Pee Dee Belemnite.

From foundational work done by Cerling (1991) we know that the isotopic composition of soil CO₂ is different from that of soil-respired CO₂. Any change in δ¹³C of soil CO₂ with depth is influenced by 1) mixing of atmospheric and biological (or biogeochemical) sources of isotopically-distinct CO₂, which may occur via diffusion (no bulk gas flow; referred to as diffusive mixing for the remainder of the paper) or advection (bulk gas flow) and 2) kinetic fractionation by diffusion. The effect of these is illustrated in Fig. 1 using a simulated soil gas profile. In panel (a) two depth profiles of δ¹³C of CO₂ that were modelled in a steady-state environment are shown (the model will be described in Section 3). The profiles differ only in soil diffusivity; all other characteristics were held constant, including rates of production, and δ¹³C of CO₂ in the atmosphere (-8 ‰; circle) and biological production (-25 ‰; square with dashed line). In the resultant depth profile with higher soil diffusivity in panel (a), the δ¹³C of soil CO₂ ranges from -8 to -15.1 ‰. In the depth profile representing a soil with lower diffusivity, the δ¹³C of soil CO₂ ranges from -8 to -20.6 ‰. We stress again these two isotopic depth profiles differ only due to differences in transport as a result of their varying soil diffusivities. In the depth profile with lower soil diffusivity, atmospheric CO₂ does not penetrate downwards as readily, so the profile shape is much steeper near the soil-atmosphere

boundary, and is more reflective of the production source composition, -25 ‰, at depth. In the depth profile with higher soil diffusivity, atmospheric air of -8 ‰ more readily mixes from the surface downward by diffusion, so the near surface isotopic composition will be more reflective of the atmosphere due to diffusive mixing of these end-members near the soil surface.

Importantly, the soil CO₂ never equals the δ¹³C of production (-25 ‰) at any depth, in either profile in Fig. 1(a). It is not possible to directly measure δ¹³C of production *in situ*, because diffusion and diffusive mixing alter the character of CO₂ immediately after its production. From the site of production in the soil, ¹²CO₂ diffuses somewhat faster through the soil than ¹³CO₂, because the former has lower mass. This diffusive difference leads to isotopic fractionation, and results in depth profiles of δ¹³C of soil CO₂ that are isotopically enriched (less negative) as compared to the source of production. Work by Cerling (1984) and later by Cerling et al. (1991), demonstrated that the mass differences between the two isotopologues led to a difference in diffusion rate of each in air, amounting to a fractionation of 4.4 ‰ (note that this applies only to binary diffusion of CO₂ in air and will differ if CO₂ diffuses in other gases). As a result, the δ¹³C of soil CO₂ measured at any depth will be enriched by a minimum of 4.4 ‰ relative to the biological production CO₂ source. However, the δ¹³C of soil-respired CO₂ can be considerably more enriched than 4.4 ‰ relative to production due to diffusive mixing with the atmosphere as shown in Fig. 1(a).

A convenient theoretical formulation for correcting δ¹³C for both diffusion fractionation and diffusive mixing was introduced by Davidson (1995), following on the work of Cerling (1984) and Cerling et al. (1991). This approach allows one to combine measurements of CO₂ and its isotopic composition within the soil and the air above it, to infer the isotopic composition of CO₂ produced in the soil. This only applies when transport within the soil is purely by diffusion (no bulk air movement). The Davidson (Davidson, 1995) solution uses the difference between the diffusion coefficients for ¹²C and ¹³C as follows:

$$\delta_f^{13} = \frac{C_s(\delta_s^{13} - 4.4) - C_a(\delta_a^{13} - 4.4)}{1.0044(C_s - C_a)}, \quad (2)$$

where δ_f¹³ is the δ¹³C composition of CO₂ from soil production (biological respiration within the soil), C_s and δ_s¹³ are the mole fraction and isotopic composition of soil CO₂, and C_a and δ_a¹³ are the mole fraction and isotopic composition of CO₂ in the air just above the soil. In Fig. 2(a) the mole fraction and isotopic composition of soil CO₂ at a 40 cm depth and of the air just above the soil was “sampled” from model-generated soil depth profiles and the (unrounded) values were used to calculate the isotopic composition of production using Davidson’s equation (C_s = 14780 ppm, δ_s¹³ = -20.3832 ‰, C_a = 380 ppm and δ_a¹³ = -8 ‰). The resulting δ_f¹³ (e.g. Eq. (2)) at this depth equals the true isotopic composition of production (see inset box, 2(a)). However, because the Davidson approach accounts for diffusion and diffusive mixing, at any given soil depth, not just 40 cm, the modelled values of C_s and δ_s¹³ in Fig. 1(a) and 2(a) will always yield (via Eq. (2)) the true isotopic composition of production, δ_f¹³ = -25 ‰ (dashed line). If δ¹³C of soil CO₂ were (erroneously) interpreted to represent the

$\delta^{13}\text{C}$ of soil-respired CO_2 , the error could be as large as the absolute value of $(\delta_a - \delta_f^{13}) - 4.4 \text{ ‰}$. This Davidson (1995) δ_f^{13} approach has been shown to be robust when applied to field data from natural soils (Bowling et al., 2015; Breecker et al., 2012; Liang et al., 2016).

While ^{14}C is a radioactive isotope and thus decays with time, the half-life is sufficiently long so that $^{14}\text{CO}_2$ behaves similarly to stable isotopes on the timescales at which diffusion occurs in a soil gas system. In this way, $\delta^{13}\text{C}$ diffusive fractionation theory can be applied to the radiocarbon isotopic composition, $\delta^{14}\text{C}$, so long as we account for the mass difference. The larger mass of ^{14}C means that the diffusion fractionation factor is calculated to be 8.8 ‰ based on the atomic masses of $^{14}\text{CO}_2$, $^{12}\text{CO}_2$ and of bulk air (Southon, 2011).

We can show that $^{14}\text{CO}_2$ distribution in soils will be like that of $^{13}\text{CO}_2$, if we model its distribution through depth in the same synthetic soil gas environment. In Fig. 1(b) we present a modelled soil environment with defined atmospheric and production source CO_2 isotopic composition boundary conditions for $\delta^{14}\text{C}$, the ^{14}C equivalent to $\delta^{13}\text{C}$ (Stuiver and Polach, 1977):

$$\delta^{14}\text{C} = \left(\frac{A_s}{A_{abs}} - 1 \right) 1000, \quad (3)$$

where $\delta^{14}\text{C}$ is the isotopic composition in ‰, A_s is the measured activity of the sample, and A_{abs} is the activity of the oxalic acid standard (both unitless). As in Fig. 1(a), in panel (b) the profile with lower soil diffusivity, the downward penetration of atmospheric CO_2 into the soil profile is reduced, and as a consequence the isotopic depth profile more closely reflects (but does not equal) the composition of production (-50 ‰; dashed line). When the diffusion rate is high and transport is rapid, the atmospheric source is more readily able to penetrate the profile and mix with the production source. In both profiles, the measured value of soil CO_2 at a given depth will not equal the isotopic production value of -50 ‰, because of diffusion and diffusive mixing. Similar profiles of $\delta^{14}\text{C}$ of soil CO_2 with depth, highlighting the diffusive effects, have been presented by Wang et al. (1994).

Since $\delta^{14}\text{C}$ transport of soil CO_2 is like that of $\delta^{13}\text{C}$, it follows that we should apply corrections for $\delta^{14}\text{C}$ like those in Eq. (2) in order to calculate the isotopic composition of production. The $\delta^{14}\text{C}$ reformulation of Davidson's δ_f^{13} equation is as follows:

$$\delta_f^{14} = \frac{C_s(\delta_s^{14} - 8.8) - C_a(\delta_a^{14} - 8.8)}{1.0088(C_s - C_a)}, \quad (4)$$

where δ_f^{14} is the $\delta^{14}\text{C}$ composition of soil production, C_s and δ_s^{14} are the mole fraction and $\delta^{14}\text{C}$ composition of the soil CO_2 , and C_a and δ_a^{14} are the mole fraction and $\delta^{14}\text{C}$ composition of CO_2 in the air just above the soil. This Davidson reformulation for $\delta^{14}\text{C}$, δ_f^{14} , was applied to a model-generated profile of soil $\delta^{14}\text{C}$ at a 40 cm depth in Fig. 2(b), like in panel (a) for $\delta^{13}\text{C}$

($C_s = 14780$ ppm, $\delta_s^{14} = -39.3989$ ‰, $C_a = 380$ ppm and $\delta_a^{14} = 45.5276$ ‰; see inset box, Fig. 2(b)). As was the case for $\delta^{13}C$ in Fig. 2(a), the modelled values of C_s and δ_s^{14} at any depth will yield the true isotopic composition of production, -50 ‰ (dashed line), because this approach accounts for diffusion and diffusive mixing.

- 5 The typical approach that has been used for interpreting the ^{14}C composition of soil CO_2 and soil-respired CO_2 (e.g., Trumbore, 2000) differs from the $\delta^{14}C$ example above, because a $\delta^{13}C$ correction is applied to account for mass-dependent isotopic fractionation of biochemical origin, ultimately converting $\delta^{14}C$ to a variant called $\Delta^{14}C$ (Stuiver and Polach, 1977). The derivation of the mass-dependent correction is provided in Stuiver and Robinson (1974), where observations are normalized to an arbitrary baseline value of -25 ‰ for $\delta^{13}C$ (a value for terrestrial wood), and the ^{13}C fractionation factors
- 10 are squared to account for the $^{14}C/^{12}C$ fractionation factor as follows:

$$\begin{aligned} A_{SN} &= A_s \left[\frac{R_s(-25)}{R_s} \right]^2, \\ &= A_s \frac{\left[\left(1 - \frac{25}{1000} \right) * R_{VPDB} \right]^2}{\left[\left(1 + \frac{\delta^{13}C}{1000} \right) * R_{VPDB} \right]^2}, \\ &= A_s \frac{\left[\left(1 - \frac{25}{1000} \right) \right]^2}{\left[\left(1 + \frac{\delta^{13}C}{1000} \right) \right]^2}, \end{aligned} \quad (5)$$

- where A_{SN} is the normalized sample activity, A_s is the sample activity, and $\delta^{13}C$ is the isotopic composition of the sample (soil CO_2 in our case). As explained in Stuiver and Robinson (1974), the 0.975 term sometimes used in forms of A_{SN} is equivalent to $\left(1 - \frac{25}{1000} \right)$, which we will retain for clarity. The equation for $\Delta^{14}C$, the $\delta^{13}C$ corrected variant of $\delta^{14}C$, can then be created from Eq. (5) by substituting in delta notation for $\Delta^{14}C$ of $\Delta^{14}C = (A_{SN}/A_{abs} - 1) * 1000$ as follows Stuiver and Robinson (1974):
- 15

$$\Delta^{14}C = \left[\left(1 + \frac{\delta^{14}C}{1000} \right) \frac{\left(1 - \frac{25}{1000} \right)^2}{\left(1 + \frac{\delta^{13}C}{1000} \right)^2} - 1 \right] 1000. \quad (6)$$

- 20 Combining Eqs. (3) and (6):

$$\Delta^{14}C_{old} = \left[\left(\frac{A_s}{A_{abs}} \right) \frac{\left(1 - \frac{25}{1000} \right)^2}{\left(1 + \frac{\delta^{13}C}{1000} \right)^2} - 1 \right] 1000. \quad (7)$$

- For more information on the derivation of Eqs. (6) and (7) see Stuiver and Robinson (1974), page 88. In Eq. (7) we have added the subscript "old" to highlight that this is the common approach used to date for soil gas applications - we will introduce a "new" method with Eq. (15). The terms on the left-hand side of Eqs. (6) and (7) are identical. Note that A_{abs} in our notation is equivalent to A_O in Stuiver and Robinson (1974).
- 25

Eq. 7 uses $\delta^{13}\text{C}$ as an input parameter to make a mass-dependent correction to obtain $\Delta^{14}\text{C}$, but the profiles of $\delta^{13}\text{C}$ and $\delta^{14}\text{C}$ of soil CO_2 (Fig. 1) highlight that both vary within the soil because of diffusion and diffusive mixing. This makes it unclear what form of $\delta^{13}\text{C}$ should actually be used in the place of the mass-dependent correction in the soil gas environment ($\delta^{13}\text{C}$ of the soil CO_2 is measured, but $\delta^{13}\text{C}$ of biological production is not) as diffusive mixing is not a mass-dependent process.

- 5 When $\Delta^{14}\text{C}_{\text{old}}$ is modelled through depth like $\delta^{13}\text{C}$ and $\delta^{14}\text{C}$ in Figs. 1 and 2 it also varies with depth as shown in Fig. 2(c). However, using a $\Delta^{14}\text{C}$ variant of Davidson's δ_j (as for $\delta^{14}\text{C}$ in Fig. 2(b)) at the same 40 cm depth does not correctly reproduce the specified model value for the $\Delta^{14}\text{C}$ of production of -50 ‰, like it did for $\delta^{13}\text{C}$ and $\delta^{14}\text{C}$ ($C_s = 14780$ ppm, $\Delta_s = -48.4319$ ‰, $C_a = 380$ ppm and $\Delta_a = 10$ ‰; see inset box, 2(c)). We, therefore, adapted the mass-dependent correction in $\Delta^{14}\text{C}_{\text{old}}$ using Davidson's (1995) theory to demonstrate how and why it should be used for $\Delta^{14}\text{C}$ soil gas applications.

10 3 Methods – Model Description

We used an analytical gas transport model to simulate a range of natural soil profiles of $^{12}\text{CO}_2$, $^{13}\text{CO}_2$ and $^{14}\text{CO}_2$ in order to present soil gas transport theory. The model is based on Fick's second law of diffusion:

$$\theta \frac{\partial \text{Conc}}{\partial t} = \frac{\partial}{\partial z} \left(D(z, t) \frac{\partial \text{Conc}}{\partial z} \right) + P(z, t), \quad (8)$$

where θ is the soil air-filled pore space, *Conc* is the concentration, *t* is time, $D(z, t)$ is the soil gas diffusion function, and

- 15 $P(z, t)$ is the biological production function, with the latter two dependent on both depth *z* and time *t*.

The model was run in steady-state:

$$\frac{\partial \text{Conc}}{\partial t} = 0, \quad (9)$$

and both diffusion and production rates were constant with depth:

$$D(z) = D, \quad (10)$$

$$20 \quad P(z) = P. \quad (11)$$

The following boundary conditions were used:

$$C(z = 0) = \text{Conc}_{\text{atm}}, \quad (12)$$

$$\frac{\partial C}{\partial z} \Big|_{z=L} = 0, \quad (13)$$

where Conc_{atm} is the concentration of CO_2 in air just above the soil and *L* is the model lower spatial boundary, the point

- 25 below which no production or diffusion occurs. Eq. (8) is solved analytically to yield the following equation:

$$\text{Conc}(z) = \frac{P/L}{D} \left(L \times z - \frac{z^2}{2} \right) + \text{Conc}_{\text{atm}}. \quad (14)$$

In the model, isotopologues of CO₂ are treated as independent gases, with their own specific concentration gradients and diffusion rates (Cerling et al., 1991; Nickerson and Risk, 2009; Risk and Kellman, 2008). We assume total CO₂ to be ¹²CO₂ because of its high abundance. The error associated with this assumption is less than 0.01% (Amundson et al., 1998). Eq. (14) is thus applied for ¹³CO₂ and ¹⁴CO₂. For the full derivation see Nickerson et al. (2014) Section 2.3.

5

The analytical gas transport model was applied across a range of soil diffusivity (1×10^{-7} , 1×10^{-6} and 1×10^{-5} m² s⁻¹), and soil production rates (0.5, 1, 2, and 4 μmol CO₂ m⁻³ s⁻¹). The specific soil diffusivity and production rates used to generate each profile are stated in the figure caption of that profile. We used a δ¹³C of biological production (-25 ‰) and atmospheric CO₂ (δ_a; -8 ‰), and Δ¹⁴C of biological production (-50 ‰) and atmospheric CO₂ (Δ_a; 10 ‰) to represent realistic conditions found in nature. The other model boundary conditions were as follows: $L = 0.8$ m, $z = 0.025$ m, and $Conc_{atm} = 15833$ μmol m⁻³ (~380 ppm). The output of the model under these applied conditions were profiles of ¹²CO₂, ¹³CO₂, and ¹⁴CO₂ for each depth (z) down to the bottom boundary (L).

10

4 Results

4.1 Adapted Correction for Interpreting Radiocarbon Values of Soil and Soil-respired CO₂

Based on Davidson's (1995) theory and what we demonstrated with Fig. 2(c), rather than using the δ¹³C soil pore space as a mass-dependent correction, we suggest instead using the value δ_J¹³ (Eq. (2)), the biological production of δ¹³C, in its place in the denominator of Eq. (7) as follows:

15

$$\Delta^{14}C_{new} = \left[\left(\frac{A_S}{A_{abs}} \right) \frac{\left(1 - \frac{25}{1000} \right)^2}{\left(\frac{\delta_J^{13}}{1 + \frac{1}{1000}} \right)^2} - 1 \right] 1000. \quad (15)$$

Values of Δ¹⁴C_{new} through depth represent transport-fractionation-corrected soil CO₂ values of radiocarbon, and in comparison to Δ¹⁴C_{old}, they are corrected for mass-independent diffusive mixing.

20

A depth profile of Δ¹⁴C_{new} is presented in Fig. 3 (dashed line). To generate this soil depth profile we used the numbers from the simulated profiles in Fig. 2 and inserted them into Eq. 2 to determine δ_J¹³ at each depth. These values were then used in Eq. 15 to obtain Δ¹⁴C_{new} of soil CO₂ through depth. The Δ¹⁴C_{new} profile (dashed line) is more isotopically enriched compared to the Δ¹⁴C_{old} profile (solid line) in Fig. 3. Values “sampled” from the Δ¹⁴C_{old} profile (the same as the one presented in Fig. 2 (c)) were not able to reproduce the specified model value for the Δ¹⁴C of production of -50 ‰ using a Δ¹⁴C variant of Davidson's δ_J. To demonstrate that Δ¹⁴C_{new} does correct for gas-transport fractionations, it can be placed into Δ_J¹⁴, a Δ¹⁴C adaption of Davidson (1995) for ¹⁴C (Eq. (4)) as follows:

25

$$\Delta_J^{14} = \frac{C_S(\Delta^{14}C_{new} - 8.8) - C_A(\Delta_A^{14} - 8.8)}{1.0088(C_S - C_A)}, \quad (16)$$

where Δ_f^{14} is the $\Delta^{14}\text{C}$ composition of soil production, C_s and $\Delta^{14}\text{C}_{\text{new}}$ are the mole fraction and $\Delta^{14}\text{C}$ composition of the soil CO_2 , and C_a and Δ_a^{14} are the mole fraction and $\Delta^{14}\text{C}$ composition of CO_2 in the air just above the soil.

Unlike in the case of $\Delta^{14}\text{C}_{\text{old}}$ demonstrated in the inset box in Fig. 2 (c), using the same 40 cm depth from the $\Delta^{14}\text{C}_{\text{new}}$ profile, we were able to reproduce the specified model value for the $\Delta^{14}\text{C}$ of production of -50 ‰ ($C_s = 14780$ ppm, $\Delta_s = -39.3989$ ‰, $C_a = 380$ ppm and $\Delta_a = 45.5276$ ‰; see inset box, Fig. 3).

4.2 Workarounds and Establishing New Best Practice

In the soil gas environment, $\Delta^{14}\text{C}_{\text{new}}$ convention should be used to properly interpret soil-respired CO_2 from soil CO_2 , as it corrects for all related transport fractionations. For researchers who have soil CO_2 data previously interpreted using $\Delta^{14}\text{C}_{\text{old}}$, the following steps will help correct for transport fractionations: 1) use δ_s^{13} and $\Delta^{14}\text{C}_{\text{old}}$ to back out the activity of the sample (A_s); 2) calculate the isotopic composition of production for $\delta^{13}\text{C}$ using Eq. (2), δ_f^{13} ; 3) use δ_f^{13} and A_s in Eq. (7) to calculate $\Delta^{14}\text{C}_{\text{new}}$, and finally 4) determine the radiocarbon isotopic composition of production, using Eq. (16), Δ_f^{14} .

Going forward, several changes to best practice are recommended. On a lab level, for new soil CO_2 data, we propose that AMS laboratories report radiocarbon using Eq. (3), $\delta^{14}\text{C}$, the uncorrected radiocarbon variant, so that the first step above, i.e. use δ_s^{13} and $\Delta^{14}\text{C}_{\text{old}}$ to back out the activity of the sample (A_s), can be avoided, and researchers can proceed with steps 2-4. We also suggest that researchers measure δ^{13} alongside $\Delta^{14}\text{C}$, so that they are not dependent on the AMS measured δ^{13} for potential back-corrections

The Davidson (1995) δ_j method was the gradient approach we used in our study, but alternative gradient approaches such as those presented for $\delta^{13}\text{C}$ by Goffin et al. (2014) and Nickerson et al. (2014), and for $\Delta^{14}\text{C}$ by Phillips et al. (2013) would likely be similarly successful in producing depth-dependent compositions of production. They are, however, not quite as straightforward as the δ_j method.

6 Conclusions

This traditional $\Delta^{14}\text{C}$ solution, which uses $\delta^{13}\text{C}$ of soil CO_2 as a mass-dependent correction, is not appropriate for the soil gas environment, as it does not account for mass-independent mixing processes. We propose a new best practice for $\Delta^{14}\text{C}$ work in the soil gas environment that accounts for gas transport fractionations and produces true estimates of $\Delta^{14}\text{C}$ of production.

Author Contributions

JE, DB and DR conceptualized the theory and method for proving the new solution for radiocarbon applications in the soil gas environment. JE carried out the modeling, validation, visualization and writing of the original draft. DB, DR and JE reviewed and edited the draft.

5 Acknowledgements

Thanks to Thure Cerling for helpful discussions on the manuscript. JE is grateful for support from a Research-in-Residence Award from the Interuniversity Training in Continental-scale Ecology Project, US National Science Foundation Macrosystems Biology Program under award EF-1137336. JE was also funded by the Natural Sciences and Engineering Research Council of Canada (NSERC). Additional support was provided by the US Department of Energy, Office of Science, Office of Biological and Environmental Research, Terrestrial Ecosystem Science (TES) Program under award number DE-SC0010625.

References

- 15 Bowling, D. R., Egan, J. E., Hall, S. J. and Risk, D. A.: Environmental forcing does not induce diel or synoptic variation in the carbon isotope content of forest soil respiration, *Biogeosciences*, 12(16), 5143–5160, doi:10.5194/bg-12-5143-2015, 2015.
- Breecker, D. O., Payne, A. E., Quade, J., Banner, J. L., Ball, C. E., Meyer, K. W. and Cowan, B. D.: The sources and sinks of CO₂ in caves under mixed woodland and grassland vegetation, *Geochim. Cosmochim. Acta*, 96, 230–246, doi:10.1016/j.gca.2012.08.023, 2012.
- 20 Cerling, T. E.: The stable isotopic composition of modern soil carbonate and its relationship to climate, *Earth Planet. Sci. Lett.*, 71(2), 229–240, 1984.
- Cerling, T. E., Solomon, K. D., Quade, J. and Bowman, J. R.: On the isotopic composition of carbon in soil carbon dioxide, *Geochim. Cosmochim. Acta*, 55, 3403–3405, 1991.
- 25 Davidson, G. R.: The stable isotopic composition and measurement of carbon in soil CO₂, *Geochim. Cosmochim. Acta*, 59(12), 2485–2489, 1995.
- Goffin, S., Aubinet, M., Maier, M., Plain, C., Schack-Kirchner, H. and Longdoz, B.: Characterization of the soil CO₂ production and its carbon isotope composition in forest soil layers using the flux-gradient approach, *Agric. For. Meteorol.*, 188, 45–57, doi:10.1016/j.agrformet.2013.11.005, 2014.
- 30 Liang, L. L., Riveros-Iregui, D. A. and Risk, D. A.: Spatial and seasonal variabilities of the stable carbon isotope composition of soil CO₂ concentration and flux in complex terrain, *J. Geophys. Res. Biogeosciences*, 121(9), 2015JG003193, doi:10.1002/2015JG003193, 2016.
- Nickerson, N. and Risk, D.: Physical controls on the isotopic composition of soil-respired CO₂, *J. Geophys. Res.*, 114(G1), doi:10.1029/2008JG000766, 2009.

- Nickerson, N., Egan, J. and Risk, D.: Subsurface approaches for measuring soil CO₂ isotopologue flux: Theory and application, *J. Geophys. Res. Biogeosciences*, 119(4), 2013JG002508, doi:10.1002/2013JG002508, 2014.
- Phillips, C. L., McFarlane, K. J., Risk, D. and Desai, A. R.: Biological and physical influences on soil ¹⁴CO₂ seasonal dynamics in a temperate hardwood forest, *Biogeosciences*, 10(12), 7999–8012, doi:10.5194/bg-10-7999-2013, 2013.
- 5 Risk, D. and Kellman, L.: Isotopic fractionation in non-equilibrium diffusive environments, *Geophys. Res. Lett.*, 35(2), doi:10.1029/2007GL032374, 2008.
- Southon, J. R.: Are the fractionation corrections correct: Are the isotopic shifts for ¹⁴C/¹²C ratios in physical processes and chemical reactions really twice those for ¹³C/¹²C?, *Radiocarbon*, 53(4), 691–704, 2011.
- Stuiver, M. and Polach, H. A.: Discussion: Reporting of ¹⁴C Data, *Radiocarbon*, 19(3), 355–363, 1977.
- 10 Stuiver, M. and Robinson, S. W.: University of Washington Geosecs North Atlantic carbon-14 results, *Earth Planet. Sci. Lett.*, 23, 87–90, 1974.
- Trumbore, S.: Age of soil organic matter and soil respiration: radiocarbon constraints on belowground C dynamics, *Ecol. Appl.*, 10(2), 399–411, 2000.
- 15 Wang, Y., Amundson, R. and Trumbore, S.: A model for soil ¹⁴CO₂ and its implications for using ¹⁴C to date pedogenic carbonate, *Geochim. Cosmochim. Acta*, 58, 393–399, 1994.

Table 1. List of symbols used. Note the isotope composition ratios are also unitless but traditionally expressed using permil (‰) notation.

Symbol	Description	Unit
A_s	sample activity	unitless
5 A_{SN}	normalized sample activity, relative to $\delta^{13}\text{C}$ of terrestrial wood	unitless
A_{abs}	age-corrected absolute international standard for activity	unitless
$Conc$	CO_2 concentration	$\mu\text{mol m}^{-3}$
$Conc_{atm}$	CO_2 concentration in air just above the soil	$\mu\text{mol m}^{-3}$
C_a	CO_2 mole fraction in air just above the soil	$\mu\text{mol mol}^{-1}$
10 CO_2	CO_2 mole fraction relative to dry air	$\mu\text{mol mol}^{-1}$
C_s	CO_2 mole fraction in soil pore space	$\mu\text{mol mol}^{-1}$
D	soil gas diffusivity	$\text{m}^2 \text{s}^{-1}$
$D(z,t)$	soil gas diffusivity at depth z and time t	$\text{m}^2 \text{s}^{-1}$
$\delta^{13}\text{C}$	stable ($^{13}\text{C}/^{12}\text{C}$) isotope composition (relative to VPDB)	‰
15 $\delta^{14}\text{C}$	radiocarbon ($^{14}\text{C}/^{12}\text{C}$) isotope composition (relative to A_{abs})	‰
$\Delta^{14}\text{C}_{old}$	radiocarbon ($^{14}\text{C}/^{12}\text{C}$) isotope composition with $\delta^{13}\text{C}$ correction	‰
$\Delta^{14}\text{C}_{new}$	radiocarbon ($^{14}\text{C}/^{12}\text{C}$) isotope composition with $\delta_j^{13}\text{C}$ correction	‰
δ_a^{13}	$\delta^{13}\text{C}$ of CO_2 in air above the soil	‰
δ_a^{14}	$\delta^{14}\text{C}$ of CO_2 in air above the soil	‰
20 Δ_a	$\Delta^{14}\text{C}$ of CO_2 in air above the soil	‰
δ_j^{13}	$\delta^{13}\text{C}$ of CO_2 from soil production, calculated using Eq. (2)	‰
δ_j^{14}	$\delta^{14}\text{C}$ of CO_2 from soil production, calculated using Eq. (4)	‰
Δ_j^{14}	$\Delta^{14}\text{C}$ of CO_2 from soil production, calculated using Eq. (16)	‰
δ_s^{13}	$\delta^{13}\text{C}$ of CO_2 in soil pore space	‰
25 δ_s^{14}	$\delta^{14}\text{C}$ of CO_2 in soil pore space	‰
Δ_s	$\Delta^{14}\text{C}$ of CO_2 in soil pore space	‰
L	lower model depth boundary	m
$P(z,t)$	biological production rate at depth z and time t	$\mu\text{mol CO}_2 \text{m}^{-3} \text{s}^{-1}$
P	biological production rate	$\mu\text{mol CO}_2 \text{m}^{-3} \text{s}^{-1}$
30 R_s	isotopic ratio (heavy/light) of CO_2 sample	unitless
R_{VPDB}	isotopic ratio (heavy/light) of Vienna Pee Dee Belemnite standard	unitless
t	time	s
θ	air-filled porosity of soil	unitless
z	depth	m




Figure 1: Modelled steady-state diffusive vertical depth profiles for $\delta^{13}\text{C}$ and $\delta^{14}\text{C}$ of soil CO₂. In the top panel the $\delta^{13}\text{C}$ of atmospheric CO₂ (circle) is -8 ‰ and CO₂ from biological production (square with dashed line; δ_b) is -25 ‰. In the bottom panel the $\delta^{14}\text{C}$ of atmospheric CO₂ (circle) is 45.5 ‰ and CO₂ from biological production (square with dashed line) is -50 ‰. Both profiles in each of the panels have the same biological production rates and isotopic composition of biological production, but each profile has a different soil diffusivity as indicated.




Figure 2: Modelled steady-state diffusive vertical depth profiles for $\delta^{13}\text{C}$ (top panel; a), $\delta^{14}\text{C}$ (middle panel; b), and $\Delta^{14}\text{C}_{\text{old}}$ (bottom panel; c) of soil CO_2 . The three soil profiles were generated using the same soil production and diffusivity rates ($2 \mu\text{mol m}^{-3} \text{s}^{-1}$ and $1\text{e}^{-6} \text{m}^2 \text{s}^{-1}$, respectively). Panels 2(a) and 2(b) were prepared using $\delta^{13}\text{C}$ and $\delta^{14}\text{C}$ as noted. Panel 2(c) shows an approach consistent with present day, where the $\Delta^{14}\text{C}$ profile generated by the model incorporates the traditional Stuiver and Polach (1974) correction for biochemical fractionation. Inset “Calculated” panels show how, using input data read directly from each depth profile, a user would arrive at either the correct, or incorrect isotopic value of production using a Davidson approach to adjust for in-soil gas transport. The atmospheric source (Ca) composition is presented as a white circle, the soil CO_2 composition (Cs) is a black circle, and the isotopic composition of production is a black square. Note that values for the isotopic composition of soil in the three panels are rounded for ease of reading, but are actually -20.3832 ‰, -39.3989 ‰, and -48.4319 ‰ respectively for panels (a), (b), and (c). These values are drawn from the curve at a depth of 40 cm.




Figure 3: Modelled steady-state diffusive vertical depth profiles for $\Delta^{14}\text{C}_{\text{old}}$ (solid line; same profile as in Fig. 2 (c)) and $\Delta^{14}\text{C}_{\text{new}}$ (dashed line) of soil CO_2 . The $\Delta^{14}\text{C}_{\text{new}}$ soil profile was calculated from the profiles in Fig. 2 (soil production and diffusivity rates of $2 \mu\text{mol m}^{-3} \text{s}^{-1}$ and $1\text{e}^{-6} \text{m}^2 \text{s}^{-1}$, respectively). The inset “Calculated” panels show how, using input data read directly from the depth profile of $\Delta^{14}\text{C}_{\text{new}}$, a user would arrive at the correct value of production using a Davidson approach to adjust for in-soil gas transport. The atmospheric source (C_a) composition is presented as a white circle, the soil CO_2 composition (C_s) is a black circle, and the isotopic composition of production is a black square. Note that values for the isotopic composition of soil and atmosphere are rounded for ease of reading, but are actually -39.3989 ‰ and 45.5276 ‰, respectively. These values are drawn from the curve at a depth of 40 cm.