

Reply to referee comments

<https://www.biogeosciences-discuss.net/bg-2018-455/#discussion>

Paper: Review of key causes and sources for N₂O emissions and NO₃-leaching from organic arable crop rotations

In the manuscript attached below the numbers (6-98) given to the comments below are, when relevant, given in the remarks so referees easily can identify the new text.

General comments from referee 1

The subject matter "N₂O and NO₃ losses from organic agriculture" is an interesting subject, unfortunately the paper is poorly written. **A lack of organization (1)** results in repetition of certain points and completely missing other points. This lack of organization also creates a lack of focus. By the time I finished reading the paper, I still didn't have a good idea of what the authors were trying to achieve. Also, I had difficulty following the paper in places because the authors just attached databases as supplementary material, rather than actually including summary tables or figures based on the supplementary materials. The authors need to find some way to summarize the data and present the summary in a meaningful way (either **tables or figures (2)**). Once the data has been summarized, patterns may emerge, which can then be explored. But everything right now just seems haphazard. I also got the impression that the authors only used simple regression to look at drivers of N₂O/NO₃ losses. Why not some kind of **multivariate analysis (3)**. There are also many more comments in the attached document.

Responses to general comments from referee 1

The authors are grateful that the referee has taken the time to work thoroughly with the manuscript and for the many useful comments that have improved the paper.

1) **Lack of organization**

We understand that the referee found the manuscript difficult follow, and we have therefore simplified the structure of the paper. We have also reformulated and simplified the research questions and improved the reasoning for the new research questions in the introduction, thus providing a more focused review.

In section 3, we explain the nitrogen cycle in organic arable farming with focus on aspects that influence N₂O emissions and N-leaching. This provides the background for understanding and discussing how crop and soil management in organic farming can be adapted to reduce N losses. In comment no 28, the referee suggests that section 3 and 4 should be a part of the introduction. We think there is a risk that this will make the introduction too long and ill focused. Instead we have moved part of section 3 (crop rotation) to the introduction, and we have targeted section 3 towards supply of organic matter and soil N-dynamics in organic arable crop rotations. We have shortened and included the previous section 5 "Dynamics of SMN in organic arable crop production" in the new section 3. We have removed the section on soil acidity, and the section on soil structure. The text on crop yields has been moved to the new discussion section and shortened. The previous section 4.3 has been moved to the introduction.

As a start of the new sections 4 and 5 we focus on the processes responsible for N₂O emission and NO₃-leaching, respectively, and we show that the same mechanisms are responsible in organic and non-organic systems.

The new table of contents is as follows:

Abstract

Abbreviations

1. Introduction
2. Methodology
3. Drivers of SMN and degradable C in organic arable crop rotations
 - 3.1. Supply and quality of soil organic matter
 - 3.2. Soil biological activity
 - 3.3. Soil N-dynamics
4. Drivers of N₂O emissions in organic arable crop rotations
 - 4.1. Biogeochemical processes leading to N₂O emissions
 - 4.2. Legumes during plant growth
 - 4.3. Crop residues
 - 4.3.1. Freeze - thaw / dry - wet
 - 4.3.2. Soil and tillage effects
 - 4.4. Organic fertilizers
 - 4.5. Contribution of total N-input and high emission events to N₂O emissions
 - 4.6. Impact of earthworms
5. Drivers of NO₃ leaching in organic arable crop rotations
 - 5.1. Biogeochemical processes leading to NO₃-leaching
 - 5.2. Legumes
 - 5.2.1. Grain legumes
 - 5.2.2. Forage legumes
 - 5.3. Cover crops
 - 5.4. Tillage effects
6. Key drivers of N₂O emissions and NO₃ leaching and suggested mitigation strategies
7. Research and innovation needs
8. Conclusion

We address the following questions for organic arable crop rotations:

1. How does supply and quality of organic matter in above- and below ground residues and organic amendments influence availability of easily available N and degradable C?
2. How does supply of easily available N and degradable C drive N₂O emissions and how can these be mitigated?
3. Is there a lack of correlation between total N-input and N₂O emission in organic arable crop rotations, and are total N₂O emissions primarily driven by background emissions or by episodes with high N₂O fluxes following N additions?
4. What are the main drivers for NO₃ leaching in organic arable systems and how can the leaching be reduced?

2) Tables or figures of the data in the paper

The referees do miss tables and figures within the paper. We are providing a new figure illustrating N-dynamics in organic arable crop rotations, a new table with performance of selected indicators of SMN dynamic in organic and conventional arable systems (Extract from S1), and new figures that are extracting the essence of N₂O data presented in S2 and NO₃-leaching data presented in S3.

3) Want multivariate analyses instead of regression

Given the actual problems and data we find regression analysis to be the most relevant. As long as the referee do not suggest one or more other specific multivariate methods / models that can be used to solve our problems in a better way using our data, we keep regression analysis as a good method to use in our situation. We have added a better description on how the regression analyses were done and added the prints from the statistical analyses (S4, S5).

General comments from referee 2

While there is merit in the work, I think the paper needs a lot of work before it is ready to publish. Many of the points raised are **not comprehensively explored** (4) or the explanations are not sufficiently considered. Work in this area could enhance the paper. I think the tables also need further work. I think the **tables (5)** should be included in the main paper but in a revised format

[Responses to general comments from referee 2](#)

4) Not comprehensively explored or the explanations are not sufficiently considered

We hope that the reorganization of the MS will satisfy the requests from referee 2. We have found it difficult to respond directly to this comment, but we hope that the new clearer structure and better explanations help the manuscript. If there is something specific that we have still missed, we will be happy to attend to it

Tables should be included in the main paper

See response to comment 3).

[Comments given the paper from referee 1 or 2 \(the comments are numbered from 6 to 98 in the paper supplemented by the referees, for responses see below \(page 8\)](#)

Abstract

- 5) You include NO₃ leaching here, which is not an "emission". I think the term "losses" would be more appropriate. [Page 1, line 25](#)
- 6) This is a generalisation, you can get poorly managed organic fields too. Maybe use the term "in general"? [P1, l 28](#)
- 7) crop termination? senescence? [P1, l 31](#)
- 8) and the form of N, whether available or unavailable [P1, l 33](#)
- 9) `the risk "of" not "for", please replace throughout [P1, l 33](#)
- 10) remove "These" [P1, l 34](#)
- 11) for [P1, l 34](#)

Section 1 Introduction

- 12) combine these two sentences [P2, l 7](#)

- 13) the term "arable" refers to the land (i.e. land fit for agricultural production), so you don't produce "arable". the terms "crop production" or "agricultural production" would be better. P2, I 16
- 14) define these "challenges" P2, I 16
- 15) N use efficiency P2, I 16
- 16) ...efficient conversion of organic material to reactive N and reactive N into plant material.... P2, I 27
- 17) But it isn't just green manures that rely on BNF, you have intercropping, or the use of N fixing plants in rotation (e.g. Soybean). P2, I 30
- 18) why do you provide the symbol for nitrate here? you have already used the word "nitrate" a few times. Please indicate symbol at the first use (i.e. line 10). P2, I 35
- 19) I find that these 5 points follow a nice, logical order. However, the rest of the manuscript does not follow the same order. Why not? I would think that having everything consistent would be better, no? P3, I 2
- 20) I think the term "driver" is better word than "trigger". P3, I 2

Section2 M

- 21) I'm not sure what you mean by "characteristics". Soil types? Management? climate zone? please be more precise with your wording. P3, I 8
- 22) Not in the list of references. P3, I 13
- 23) The presence of NH4-N does not contribute directly to N2O emissions, NH4-N needs to be nitrified to NO3-N prior to denitrification to N2O P3, I 4
- 24) why "site"? Why not "mean annual precipitation" or "soil clay content" or something more meaningful? Site seems completely arbitrary. P3, I 4
- 25) I think this section should go in the section describing the stepwise regression. P3, 21-27
- 26) Cumulative P3, I 34

Section3

- 27) Shouldn't this follow the 5 points laid out in the introduction? The first point in the Intro was "what determines the dynamics of SMN concentrations and when do high SMN concentrations occur? I would actually move all of section 3 into the introduction. Probably all of section 4 should go into the introduction as well. P4, I 1

Section3.1

- 28) are longer and more diversified than non-organic rotations P4, I 5
- 29) Is this required? This sounds more like it should be in the introduction. While differences in rotations likely have an effect on emissions and leaching, I'm not sure that we need to know the area associated with the different plant species. And actually, these aren't really rotations, but only specific plant types within a rotation. P4, I 9-11
- 30) This seems redundant. Essentially you are saying that N fixing crops are more abundant because there are more N-fixing crops (i.e. legumes) as temporary fodder crops.
Also, what is a "temporary fodder crop"?
Also, I think the main reason why N fixing crops are more abundant in organic rotations is because organic farmers can not use synthetic fertilizers. P4, I 11-12

Section3.2

- 31) Isn't this still part of the "Crop Rotation"? P4, I 15-18

- 32) Do you mean just in Europe???? I think this is too much of a generalisation, many organic agriculture systems remove all of the organic material as the whole crop is used eg root stem and leaves - depleting soil fertility eg subsistence agriculture in Ethiopia P4, I 18
- 33) A single sentence is not a proper paragraph. Please revise. this could probably be part of the following paragraph that discusses the SOM quality. P4, I 20-25
- 34) Shouldn't this be part of section 3.3? P4, I 24
- 35) ...fertility as it is a short-term pool for nutrients (Marriott and Wander, 2006; Martyniuk et al., 2016). However, it can also enhance SMN... P4, I 29
- Section3.3**
- 36) how much higher? P4, I 37
- 37) You essentially say the same thing (i.e. earthworm biomass and population were higher in organic soils) over and over again. Please be more concise here. This can probably be trimmed by about 50-60%. P5, I 1-11
- Section3.4**
- 38) do you have a reference for this? P5, I 21
- 39) this is essentially a repeat of the first sentence in the previous paragraph P5, I 23-24.
- 40) How much? 50%? 80% P5, I 28
- 41) If you are talking about PMN pools, they are not the same as "concentrations". I would suggest writing "They showed that more diversified crop rotations resulted in larger PMN pools". P5, I 34 - 35
- 42) SOM.... stay consistent. P5, I 39
- 43) The DOK trial should be identified the first time it is referenced. This is already the second time. P6, I 5
- 44) This section is on the N supply. Keep the information on denitrification in the section on N2O emissions. P6, I 5
- 45) You can just use "DOK" here, now that you have already stated what the acronym refers to. P6, I 17
- 46) is this relevant? P6, I 21-24
- 47) can you be more specific what you mean by "soil life"? is it plants? macrofauna? fungi? P6, I 26
- 48) which properties? You mention "macropores" in the next sentence. Is this one of the properties? since this is the only one that you mention, why not just state that Marinari observed that organic fertilizers increased macropores and particularly elongated pores? P6, I 29
- 49) which are? P6, I 32
- 50) as long as the organic rotations include more perennial leys. P6, I 32-33
- 51) Is this realistic in the context of increased pressure for urbanisation? This suggests is the existing agricultural landbase is insufficient - that we need to cut down some trees? P6, I 38
- 52) this is a huge range.... Any idea why it ranges to much? are there differences between crop types? species? do different soils or climates respond differently? P6, I 39
- 53) There are many studies that look at how mouldboard ploughing effects N2O emissions. why have you not looked at any of these? P7, I 4
- 54) Do we really need a 2-sentence summary of each section? I think this is just repeating what was already said and is therefore unnecessary. P7, I 10-13
- Section4**
- 55) You should also include something on the paper by Balaine et al. 2013 (SSSAJ 77:1496-1505) on how gas diffusivity affects N2O fluxes. P7, I 14
- 56) you said earlier (lines 17 and 18 above) that nitrification is a biological process. So why is it stated that there are abiotic reactions during nitrification? P7, I 22
- 57) I understand what you are saying here, but it seems like an odd way of saying it. I also think you need to clarify that it is the native SOM, especially in organic farming when N applied as manure or slurry can be quickly considered part of the SOM. P8, I 1

- 58) this explanation of priming is inadequate. Please add a bit more information about what is meant by priming. [P8, I 3-5](#)
- 59) there should also be studies that use 15-N applications to determine whether the source is nitrification or denitrification. And this could also differ based on soil textures / structure / precipitation because of their effect on soil aeration and gas diffusivity [P8, I 10-11](#)
- 60) Please don't have paragraphs that consist solely of a single sentence. Also, this doesn't really explain why nitrifier denitrification would be a major N₂O source here. And also, I can't figure out if you mean that the soil has low content of easily degradable C, or if the OM that is applied has low content of easily degradable C. [P8, I 15-17](#)
- 61) Why nothing on nitrification inhibitors? if you can reduce the amount of nitrification, then you should also reduce NO₃ leaching. There are both synthetic and biological Nitrification inhibitors. [P8, I 18](#)
- 62) say the same thing in next sentence [P8, I 33](#)
- 63) lower than what? [P8, I 36](#)
- 64) Table S1 presents a lot of data (as do S2 and S3 actually). This makes it very difficult to come to any conclusions about the studies. Some tables and figures that summarize some of this data would be helpful. [P9, I 10](#)

Section5

- 65) You seem to use "organic grass-clover leys" as a synonym for "N-limited grass-clover ley". Why is that? are all organic grass-clover leys N limited? vice versa? This should be clarified. [P9, I 12-13](#)
- 66) why two different numbers? is one for the 1 yr and the other the 3-yr leys? I assume so, but it is not explicitly stated. [P9, I 19](#)
- 67) which conditions? why? [P9, I 22](#)
- 68) I feel like I am missing some important details from the Plaza study, because I can't the the obvious explanation of "priming". Why couldn't it just be from the decomposition/mineralization of the added catch crop? [P9, I 27-29](#)

Section6

- 69) this might be better in section 6.2. [P10, I 13-20](#)
- 70) incorporation of the CC? [P11, I 13](#)
- 71) There are few studies that actually show this directly, however Freeze/thaw has been consistently shown to increase mineral N availability (similar to the Birch effect). The release of N₂O during thawing could be related to a few things, one of which is the release of labile C and N due to cell lysis (the term "blasted" is the wrong term, by the way). But there are other theories as well. Disruption of soil aggregates due to F/T may also release labile C and N; frozen soils may have ice lenses that lead to N₂O accumulation under the ice, which can be released upon thawing, etc. There are many papers by Wagner-Riddle (in addition to the ones you mention here) that examine N₂O pulses during thawing of soils. I would suggest adding some of those citations here to aid in providing a more detailed description of how Freeze/thaw affects N₂O fluxes.
There is a good one in SBB (Congreves et al. 2018. vol. 117: 5-15) and another in Nature Geoscience: Wagner-Riddle et al. 2017. vol 10: 279-283. [P11, I 20-21](#)
- 72) sorry, but the way this supplementary material is set up makes it really difficult to find the information to which you are supposedly referring. The Tables must be organized in a way that they present summary rather than just raw data (although the raw data is good to have as supplementary materials). [P11, I 13](#)
- 74a) which study are you referring to? [P12, I 1](#)
- 73) This seems to be to be a very large leap in logic. [P12, I 2-4](#)
- 74) or it could be that deep incorporation of crop residues in soils with restricted air diffusivity will be mineralized much more slowly, reducing the amount of NO₃ available for denitrification. [P12, I 15](#)

75) There are many other studies that look at N₂O emissions from application of various organic N sources. And there is one paper that describes how the response in N₂O fluxes differs by soil type (Pelster et al. 2012. J. Environ. Qual. 41:427-435).

You need to expand these points more clearly. P12, I 36-37

76) if you want another example from a cool humid climate, there is also Chantigny et al. 2012. JEQ. 42:30-39 P13, I 2-3

77) provide numbers please. P13, I 23

78) what did these studies actually find? did they also find a relation between N₂O flux and SOM content? P14, I 5

79) Using the term "site" is completely meaningless. How was this entered into the model? Site is categorical, how was that worked into a multiple regression model? that makes no sense to me. Why not use site-specific variables (i.e. soil clay content; annual precipitation; soil sand content... or something like that). P14, I 16

80) how? You can make a case for the presence of labile N during reducing conditions, but where do you show the relation with labile C? P14, I 19-20

81) The inequality in length of study is really a problem here, because the numbers for Frick and Aberdeen are so different from the other two. I just am not sure how to normalize this in a way that makes sense. P14, I 23-25

82) how long was the measurement period for this study? P14, I 28

83) can you give the model (with co-efficients for the different variables). Also, are all of these just simple linear regression? did you try multiple regression? why not? P14, I 34

84) isn't this tautological? and therefore does not need to be stated. P14, I 34

85) what aspect? expression of the enzyme? enzyme activity? provide details. P15, I 9

86) if the N₂O/(N₂O+N₂) ratio is higher in organic agriculture that means that a larger proportion of N is emitted as N₂O, which seems to me to be the inverse of what you think you are saying here.

Maybe I am misunderstanding something. P15, I 11-13

87) in what way? P15, I 13-14

88) earthworms often have producing bacteria in their gut. This should also probably be explored a bit P15, I 20-21

Section 7

89) It seems to me that much of what is presented here is just a repetition of what was already provided in the section on N₂O. Which makes sense. Both N₂O emissions and NO₃ leaching are strongly related to available soil NO₃ and high water content (anaerobicity).

Here you talk about legumes, grain legumes, etc as possible triggers. But in my opinion, the trigger is the available NO₃ combined with high water (low O₂ content). All of this crop rotation information should be dealt with once in the section on soil mineral N. Not repeated in multiple sections. I would include section 7.2 as part of this as well as catch crops are ways to control the concentrations of NO₃. P15, I 20-21

90) were the weeds 50% taller? or was their coverage 50% greater? P18, I 30

91) how does tillage affect the soils? pore size distribution, continuity, etc? how about rainfall impact and splashing on tilled vs non tilled soils? what effects may that have on soil infiltration and percolation? Again, you are only discussing the soil mineral N aspect again (which has already been done in a slightly different context), but you are completely missing how tillage may affect NO₃ leaching in particular. P18, I 38-39

92) any idea why? P18, I 10-11

Section 8

93) You don't really provide much information on strategies in this section. At least not in any organized way. this whole section needs to be completely re-written. P20, I 5

94) This belongs in one of the previous sections as it is not a strategy to reduce N₂O and NO₃ losses P20, I 7-8

Section 9 conclusion which is numbered 10

- 95) why only rotary harrow? what is specific about this form of mixing that enhances the risk? P21, I 21
- 96) If you showed this somewhere, I completely missed the point. For me, this comes out of nowhere and with no justification. P21, I 22-23
- 97) could this be related to statistical power? Also, as you mention, site was a significant factor. was there a correlation after controlling (normalizing) for "site"? As I remember only reading about simple regression, I can only guess that this was not done. P21, I 25-26

Responses to comments given the paper from referee 1 or 2 (the comments are numbered from 6 to 98 in the paper supplemented by the referees)

During revision of the MS, there will be some adjustments in the sentences suggested below

- 6) We have replaced emission with loadings
- 7) Agree, this is included in the new version of the paper
- 8) Agree, senescence should also be included as a possible course. New text in paper ... after crop termination, harvest or senescence.
- 9) We do not understand what Referee mean. In the sentence it is already referred to "readily available nitrogen. We have added degradable carbon to make it clearer that we refer to C and N that that are rapidly available for microorganisms.
- 10) Agree, this is done in the new version of the paper
- 11) Do not agree in that.
- 12) "For" is changed to "of" throughout the paper
- 13) Agree, this is done in the new version of the paper
- 14) Agree. New sentence: We have chosen to focus on arable systems because crop and soil management vary more between organic and non-organic production of arable crops than for grassland....
- 15) Agree that writing challenges without explaining or defining what we mean are not OK. On the other and this is lengthy to explain and a bit of the scope of this paper, we therefore removed challenges and kept yield gap which is the main issue.
- 16) Agree, this is included in the new version of the paper
- 17) Agree. However, during rewriting of the Introduction the whole sentence was removed
- 18) Agree, However, during rewriting of the Introduction the whole sentence was removed
- 19) Agree, this is included in the new version of the paper
- 20) To make it easier to follow the papers structure, we are adding questions in the introduction that are better introducing the sections 3 and 4. To help the reader we have also introduced a table of contents. New text is given in the response to comment 1.
- 21) Agree, this is included in the new version of the paper
- 22) Based on the authors own field trials, literature databases and searches through Google Scholar, we compiled data on agronomic management, soil properties and yield level of organic arable crop rotations and measurements of SMN, N₂O emissions and NO₃ leaching from field trials relevant to organic crop rotations, climate and soil conditions in Europe.
- 23) This is now included: Molodovskaya, M., Singurindy, O., Richards, B.K., Warland, J., Johnson, M.S., Steenhuis, T.S.: Temporal Variability of Nitrous Oxide from Fertilized Croplands: Hot Moment Analysis. Soil Sci. Soc. Am. J. 76, 1728. <https://doi.org/10.2136/sssaj2012.0039>, 2012.
- 24) In this postulate the referee is not consistent with the main understanding. See for instance Firestone and Davidson, 1989; Butterbach-Bahl et al., 2013). No change is made.
- 25) Agree with referee, we have recalculated the regression and included %clay, pH and SOC in the regression. We have also added some more text to make it clearer what we did.

- 26) We do not understand this comment as this is in the section where we describe the stepwise regression. The expanded text will hopefully make this easier to follow.
- 27) Neither cumulated or cumulative is needed as N₂O emission is defined as the cumulative flux reported. The word is removed
- 28) See answer to comment 1 and 20. Section 3 and 4 are too long to be a part of the introduction as suggested by referee, and they are important part of this paper.
- 29) Agree,
- 30) Agree, the sentences are removed
- 31) The sentence is modified. New sentence: They further found that more legumes are included in fodder crops, and in catch crops, under sown cover crops and intercropping than in non-organic rotations
- 32) Agree. The sentence is moved to 3.1.
- 33) The referee is correct. We have now underlined that this is relevant for European cropping systems. New sentence: "... are returned to soil than in non-organic systems in cropping systems that are common in Europe"
- 34) Agree, the section is rewritten
- 35) Agree, the section is rewritten
- 36) ... soil fertility asset as it provides a short-term pool for plant nutrient supply (Marriott and Wander, 2006; Martyniuk et al., 2016). SMN is not included in this sentence because it comes in 3.3
- 37) Without being too comprehensive for this paper, where N₂O and NO₃ is the focus, it is difficult to give numbers on this. There are many ways to estimate/determine biological activity in soils. See for instance "Mäder, P., Fliessbach, A., Dubois, D., Gunst, L., Fried, P., Niggli, U., 2002. Soil fertility and biodiversity in organic farming. *Science* 296, 1694–7. <https://doi.org/10.1126/science.1071148>" where microbial biomass, dehydrogenase, protease, phosphatase, saccharase and mycorrhiza are used as indicators for microbial activity and earthworms, carabids, Staphylinids and spiders for faunal activity. The exact increase of biological activity in % will also vary with the sites climatic and soil conditions. The main thing is that it is a general trend with increased biological activity, which is shown in many investigations. No change is made.
- 38) Most of the examples are removed to make it less repeating. Earthworms do only have minor part of this subject so the referee has a good point.
- 39) Reference included: Pandey et al., 2018
- 40) Agree, the sentence is removed in the new version of the paper
- 41) This section is rewritten
- 42) Agree. The suggested sentence is included in the new version of the paper
- 43) Agree, this is done throughout the paper
- 44) The first time is removed, so this is the first time and adjusted accordingly.
- 45) We have removed this sentence.
- 46) To focus the MS more on the aspects where organic farming differ from most conventional farming, we have taken out the aspects of pH, and this text is therefore removed.
- 47) Same as for 46.
- 48) The text on soil structure is taken out of the MS
- 49) The text on soil structure is taken out of the MS
- 50) The text on soil structure is taken out of the MS
- 51) The text on soil structure is taken out of the MS
- 52) This article is not about organic farming or not, but about how different aspects that are typical for organic farming influence N₂O emission and NO₃ leaching. However, a much larger impact on

- feeding the world's population and the area needed for agricultural production, than if the yields are 80 or 100%, is the unequal distribution of resources in the world, the land that is taken out of use for roads, building ground or other purposes, all the food that is wasted, corn and wheat used for bio-energy and so on. However, the text on yields is shortened and moved to Section 6
- 53) yield level will be moved to the new discussion. Comment 53 relate to Meier et al. (2015) observed that 9 to 214% more land is needed to produce one arable crop unit by organic compared with non-organic production. The requested information is not easy to extract from the present article. We will rewrite this whole section when used in the introduction.
- 54) This section is about yields. The impact of ploughing on N₂O emission is discussed in the new section 4.3.3 Soil and tillage effects
- 55) We have removed the summaries
- 56) Thank you for this reference. A text is included in new section 4.1. Oxygen availability depends on soil microbial activity and gas diffusivity which is in dependent on soil moisture content, texture and density. Gas diffusivity is a promising predictor for N₂O fluxes from soils with varying bulk density as observed by Balaine et al., (2013) who found that the production of N₂O increased when the relative gas diffusivity was between 0.006 and 0.020 and the soil became anaerobic.
- 57) Yes the reviewer is right that « nitrification is a biological process” during this process NH₄ is oxides to NO₂- via hydroxylamine (NH₂OH) by ammonia oxidizing bacteria (AOB). During this process N₂O is produced as by product due to some partially understood abiotic reactions of hydroxylamine and NO₂-. Recent studies (i.e Liu et al 2017) have shown that there are certain abiotic reactions of hydroxylamine that can contribute to N₂O production in this process, also another hypothesis is that NO₂ (which is an intermediate in this process) can also be converted to N₂O through abiotic reactions. That's why we wrote partially understood biotic (for sure) and abiotic (unclear) reactions. We have made this clearer by adding “abiotic reactions of hydroxylamine”
- 58) We have changed the sentence slightly: “Emissions from soil organic matter (background emission) will vary between years because of variations in temperature and precipitation “ In addition we have defined background emission in the under terminology at the beginning of the paper.
- 59) We have decided to remove the text about priming as it does not have a major impact on N₂O emissions
- 60) This is the main picture. We have included some more on gas diffusivity (comment 56). It is to comprehensive for this review to cover all aspects of N₂O formation in this review.
- 61) The sentence is removed as the content is well enough covered above
- 62) Inhibitors are not allowed in organic farming. This is now specified in the introduction
- 63) The whole section is rewritten and moved to the introduction
- 64) The whole section is rewritten and moved to the introduction
- 65) See respons to comment 2).
- 66) The sentence is rewritten to emphasise that it is SMN content that matters:” SMN is normally very low under organic grass-clover leys (Table S1, Watson et al., 1993; Nadeem et al., 2012; Brozyna et al., 2013; Frøseth et al., 2014; Krauss et al., 2017b), thus grasses will quickly take up soil NO₃ in the root zone (Brophy et al., 1987).”
- 67) We have added the word “respectively” to make it clearer that this is Soil SMN after 1 and 3 years, respectively.
- 68) This sentence is removed, and the content is covered in the rest of the revised section 3.3

- 69) We have decided to remove the text about priming as it does not have a major impact on N₂O emissions
- 70) We were uncertain about what is the best place, because of the relation to growing clover. A part of this is now moved to new section 4.3.1, and a part is deleted.
- 71) Agree, this is included in the new version of the paper
- 72) Thanks for these references. Freezing/thawing is a large area and we think that for this article it is correct to emphasize what is relevant for organic systems: high content of SOM, soil biota, CC. We have included relevant information from Congreves et al. 2018. vol. 117: 5-15 and Wagner-Riddle et al. 2017. vol 10: 279-283. P11, l 20-21, which helps to complete the picture. New text: "The mechanisms behind freeze/thaw have been comprehensively reviewed by Congreves et al. (2018) showing that the causes for N₂O emissions are different for these two mechanisms and that freeze/thaw has a larger impact on N₂O emissions from temperate agroecosystems than drying/rewetting do. Wagner-Riddle et al. (2017) estimated that neglect of freeze/thaw emissions will underestimate global agricultural N₂O emissions by 17 to 28%. In the present article we do discuss freezing/thawing and drying/rewetting during conditions that are particularly relevant for organic crop rotations. Freezing/thawing of soil rich in organic matter and soil biota, or soil covered with plant residues may result in a N₂O boost as easily degradable C and N is released from cells lysis after frost."
- 73) See response to comment 2
- 74) It is understandable that the referee misunderstood this. The last sentence is removed as it is enough covered from earlier sentences.
- 75) The text is rewritten. We did not include "or it could be that deep incorporation of crop residues in soils with restricted air diffusivity will be mineralized much more slowly, reducing the amount of NO₃ available for denitrification" as we do not have any evidence for that. If referee does, we are grateful for suggested reference.
- 76) Thank you for this reference. Nice article. When revising the MS we have included these points and also expanded the discussion on the impact of N-content and the type of N-compound in the organic fertilizers on N₂O fluxes
- 77) I assume you mean this one: Chantigny, M.H., Angers, D.A., Rochette, P., Bélanger, G., Massé, D., Côté, D., 2007. Gaseous Nitrogen Emissions and Forage Nitrogen Uptake on Soils Fertilized with Raw and Treated Swine Manure. *J. Environ. Qual.* 36, 1864.
<https://doi.org/10.2134/jeq2007.0083>,
- 78) Numbers for N₂O-N and C are included. "Krauss et al. (2017b) found that fertilization with slurry and manure compost increased annual N₂O emissions during winter wheat after more than ten years of differentiated management compared to sole slurry fertilization (mean values in the period (369 days), were 2.2 and 2.9 kg N₂O-N ha⁻¹, respectively). They related this to higher microbial biomass and content of SOC. Mean values for the upper 10 cm in soil were 28 and 30 Mg C ha⁻¹ for fertilization with slurry and manure compost and sole slurry fertilization, respectively."
- 79) The text is rewritten to make it more clear that this refers to: Lack of correlation between N₂O emissions and N fertilization. Neither Pugesgaard et al (2017) nor Peyrard et al. (2016) have related their findings to the content of SOM.
- 80) See response to comment 25. We do not longer have a categorical variable. (That is however in principal no problem to include in a stepwise regression).
- 81) The sentence has been improved so it now should be easier to relate to the first sentence, which was the intention.: "The content of NH₄-N in soil did not affect peak N₂O fluxes. These findings indicate that denitrification is the main cause for high N₂O-flux rates in these studies"

- 82) Agree with referee. This is indeed problematic. We have tried to overcome this by estimating the impact of the highest emission on the mean daily N₂O emissions, but still this is tricky because longer periods with low emissions are not included in the shorter periods. The best would be to have yearly emissions, but we did the best we could do from the data available.
- 83) Agree that this is essential information. This is now included.
- 84) model (with co-efficients for the different variables) is now given in section 2 (Methodology). In the first model where we tested different factors we used multippel, stepwise regression. In the second model where we tested the impact of the highest peak on the total emission linear regression was the best option.
- 85) We do difference between flux and emission. Emissions relating to the total period. The sentence is rewritten to make this clearer." From this we can conclude that when the conditions for high N₂O fluxes are met at one or more days, there is a large chance for high total N₂O emissions in the period."
- 86) In order to focus the MS more on the aspects where organic farming differ from most conventional farming, we have taken out the aspects of pH, and this text is therefore removed.
- 87) The same as for comment 86
- 88) The same as for comment 86
- 89) We have moved the sentence that described this to an earlier position, and have made it clearer that body fluids is a part of earthworm gut. This, to make it easier to perceive for the reader
- 90) See comment 1. We have restructured the MS and have gone through the new section 5 to avoid repetition from section 3 and 4.
- 91) We have clarified by including that the weed incidence was higher.
- 92) We have focused the paper, and do think that general effects of tillage on nitrate leaching is out of the focus.
- 93) We have added an explanation and the new text is: "A field experiment over 13 years in the UK showed that N leaching in winter from fertilized grass (non-organic) was highly correlated with the preceding summer's soil moisture deficit, with the highest losses following dry summers (Tyson et al., 1997). In this case, poor grass growth due to drought lead to a buildup of NO₃ from unused fertilizer present in the autumn. Prolonged mineralization of organic fertilizers or crop residues due to drought may also lead a similar situation in organic farming systems."
- 94) See comment 1. We are restructuring the MS and have removed old section 8.
- 95) We have removed old section 8.
- 96) We have removed old section 8.
- 97) This is from section 6.4. It is understandable that this sentence was confusing.
- 98) This is from the literature review presented in section 6.4 and not from our statistical calculations as we were not able to run statistic on the impact of total N because of lack of data in the relevant experiments.
- 99) Miss the comment from referee.

Review of causes and sources for N₂O emissions and NO₃-leaching from organic arable crop rotations

Sissel Hansen¹, Randi Berland Frøseth², Maria Stenberg³, Jarosław Stalenga⁴, Jørgen E. Olesen⁵, Maike Krauss⁶, Paweł Radzikowski⁴, Jordi Doltra⁷, Shahid Nadeem⁸, Torfinn Torp⁹, Valentini Pappa¹⁰, Christine A. Watson^{11,12}

¹Norwegian Centre for Organic Agriculture, NO-6630 Tingvoll, Norway

²Department of Grain and Forage Seed Agronomy, Norwegian Institute of Bioeconomy (NIBIO), NO-1431 Ås, Norway

³Department of Soil and Environment, Swedish University of Agricultural Sciences (SLU), Skara, Sweden

⁴Department of Systems and Economics of Crop Production, Institute of Soil Science and Plant Cultivation - State Research Institute, 24-100 Puławy, Poland

⁵Department of Agroecology, Aarhus University, DK-8830 Tjele, Denmark

⁶Department of Soil Sciences, Research Institute of Organic Agriculture (FiBL), 5070 Frick, Switzerland

⁷Cantabrian Agricultural Research and Training Centre, CIFA, 39600 Muriedas, Cantabria, Spain

⁸Faculty of Environmental Sciences and Natural Resource Management (MINA), Norwegian University of Life Sciences (NMBU), 1430 Ås, Norway

⁹Department of Research, Norwegian Institute of Bioeconomy (NIBIO), NO-1431 Ås, Norway

¹⁰Texas A&M, Energy Institute, College Station, TX 77845-3372, USA

¹¹Department of Crop Production Ecology, Swedish University of Agricultural Sciences (SLU), Uppsala, Sweden

¹²SRUC, Craibstone Estate, Aberdeen, AB21 9YA, UK

Correspondence to: Sissel Hansen (sissel.hansen@norsok.no)

Abstract.

The emissions of nitrous oxide (N₂O) and leaching of nitrate (NO₃) from agricultural cropping systems have considerable negative impacts on climate and the environment. Although these environmental burdens are on average less per unit area in organic than in non-organic production, they are roughly similar per unit of product. If organic farming is to maintain its goal of being environmentally friendly, these loadings must be mitigated. We discuss the impact of possible drivers of N₂O emissions and NO₃ leaching within organic arable farming practice under European climatic conditions, and potential strategies to reduce these. Organic arable crop rotations are generally diverse with frequent use of legumes, intercropping and organic fertilizers. The soil organic matter content and share of active organic matter, soil structure, microbial and faunal activity are higher in such diverse rotations, and yields lower, than in non-organic arable cropping systems based on less diverse systems and inorganic fertilizers. Soil mineral nitrogen (SMN), N₂O emissions and NO₃ leaching are low under growing crops, but there is the potential for SMN accumulation and losses after crop termination, harvest or senescence. The risk of high N₂O fluxes increases when large amounts of herbage or organic fertilizers with readily available nitrogen (N) and degradable carbon are incorporated into the soil or left on the surface. Freezing / thawing, drying / rewetting, compacted and/or wet soil and mechanical mixing of crop residues into the soil further enhance the risk of high N₂O fluxes. N derived from soil organic matter (background emissions) do, however, seem to be the most important driver for N₂O emission from organic arable crop rotations and the correlation between yearly total N-input and N₂O emissions is weak. Incorporation of N rich plant residues or mechanical weeding followed by bare fallow increases the risk of NO₃ leaching. In contrast, strategic use of deep-rooted crops with long growing seasons or effective cover crops in the rotation reduces NO₃ leaching risk. Enhanced recycling of herbage from green manures, crop residues and cover crops through biogas or composting may increase N efficiency and reduce N₂O emissions and NO₃ leaching. Mixtures of legumes (e.g., clover or vetch) and non-legumes (e.g., grasses or Brassica species) are as efficient cover crops for reducing NO₃ leaching as monocultures of non-legume species. Continued regular use of cover crops has the potential to reduce NO₃ leaching and enhance soil organic matter but may enhance N₂O emissions. There is a need to optimise the use of crops and cover crops to enhance the synchrony of mineralisation with crop N uptake to enhance crop productivity, and this will at the same time reduce long-term risks of NO₃ leaching and N₂O emissions.

We use the following abbreviations: *BNF*: biological nitrogen fixation, *C*: Carbon, *CC*: cover crops, *CH₄*: methane, *EF*: Emission factor = % of N applied emitted as N₂O-N, *N*: nitrogen, *N₂O*: nitrous oxide, *NO₃*: nitrate, *PMN*: Potentially mineralizable N, *SMN*: Soil mineral nitrogen, *SOC*: Soil organic carbon, *SOM*: Soil organic matter.

and terminology: *Background emissions*: N₂O emissions that derive from N released from SOM as opposed to N added through fertilization, soil amendments or plant residue applied in the current year, *Degradable C*: Easily degradable carbon compounds, *Hot moments of N₂O flux*: Brief and disproportionally high short-term N₂O flux event due to the combination of multiple influencing factors (Molodovskaya et al., 2012), *N₂O emission*: The cumulative flux reported for one field treatment during the actual measurement period, *NO₃ leached*: NO₃

Kommentert [SH1]: 6

Kommentert [SH2]: 7

Kommentert [SH3]: 8

Kommentert [SH4]: 9

transported below the root zone, *PMN*: The amount of SMN released from organic matter under ideal soil environmental conditions, *SMN*: The content of mineral N in the form of ammonium (NH₄) and NO₃ in soil.

9. Introduction
10. Methodology
11. Drivers of SMN and degradable C in organic arable crop rotations
 - 11.1. Supply and quality of soil organic matter
 - 11.2. Soil biological activity
 - 11.3. Soil N-dynamics
12. Drivers of N₂O emissions in organic arable crop rotations
 - 12.1. Biogeochemical processes leading to N₂O emissions
 - 12.2. Legumes during plant growth
 - 12.3. Crop residues
 - 12.3.1. Freeze - thaw / dry - wet
 - 12.3.2. Soil and tillage effects
 - 12.4. Organic fertilizers
 - 12.5. Contribution of total N-input and high emission events to N₂O emissions
 - 12.6. Impact of earthworms
13. Drivers of NO₃ leaching in organic arable crop rotations
 - 13.1. Biogeochemical processes leading to NO₃-leaching
 - 13.2. Legumes
 - 13.2.1. Grain legumes
 - 13.2.2. Forage legumes
 - 13.3. Cover crops
 - 13.4. Tillage effects
14. Key drivers of N₂O emissions and NO₃ leaching and suggested mitigation strategies
15. Research and innovation needs
16. Conclusion

1 Introduction

Biologically available nitrogen (N) or reactive N is limited in most natural terrestrial ecosystems. In modern crop production, addition of N fertilizer has become crucial to achieve high crop yields. This has resulted in cropping systems where a substantial proportion of the N added is lost to the environment, and where the excess reactive N threatens the quality of air, water and ecosystems (Robertson and Vitousek, 2009). The emissions of N₂O have considerable environmental impacts through the contribution to global warming and ozone depletion (Ravishankara et al., 2009), and about 16 to 20 Tg N₂O-N is emitted annually to the atmosphere. Of this, close to 40% is anthropogenic, and agriculture accounts for 67–80% of the anthropogenic N₂O emissions (Ussiri and Lal, 2013). About half of the anthropogenic N₂O emissions originate from cultivated soils (Stehfest and Bouwman, 2006). In addition, agricultural soils are sources of indirect N₂O emissions resulting from downstream microbial turnover of N from NO₃ leaching or ammonia volatilization (IPCC, 2006). NO₃ lost by leaching may also contaminate drinking water and lead to eutrophication of freshwater and marine ecosystems (Dalgaard et al., 2014). The area under organic production increases worldwide (Willer and Lernoud, 2018). In Europe, 2.7% of the agricultural land is under organic farming, and in nine countries, 10% or more of the agricultural land is managed organically (Willer et al., 2018). In 2016, 43% (6 mill ha) of the organic farmed area in Europe was under arable crops.

Organic agriculture aims to be an environmentally friendly production system that sustains the health of soils, ecosystems and people. It should rely on ecological processes, biodiversity and nutrient cycles adapted to local conditions, rather than the use of inputs (IFOAM, 2019). Because of the serious consequences of N₂O emissions

Kommentert [SH5]: 13

and NO₃ leaching, these environmental burdens are important issues also for organic farming, and there is a continued debate whether an organic mode of crop production enhances or reduces greenhouse gas (GHG) emissions and NO₃ leaching from agriculture (Lorenz and Lal, 2016; McGee, 2015). We have chosen to focus on arable systems because crop and soil management vary more between organic and non-organic production of arable crops than for grassland (Barbieri et al., 2017), and the yield gap is larger (De Ponti et al., 2012). These conditions will affect N₂O emissions and NO₃ from the systems. Previous reviews have compared the net N₂O emissions and NO₃ leaching from these systems. Lower area-scaled, but roughly similar yield-scaled emissions (slightly higher, similar or slightly lower) are commonly observed for N₂O emissions (Skinner et al., 2019; Skinner et al., 2014; Tuomisto et al., 2012,) and NO₃ leaching (Aronsson et al., 2007; Benoit et al., 2014; Kirchmann and Bergström, 2001; Stopes et al., 2002; Tuomisto et al., 2012) from organic versus non-organic arable crop production.

In this review we focus on the drivers for N₂O emissions and NO₃-leaching under organic arable crop rotations under European climatic conditions. There is insufficient robust field data on N₂O emission and NO₃ leaching within organic arable crop rotations to allow for a meta-analysis to quantify the impact of key causes, so here we use the available data to identify sources and causes of N₂O emission and NO₃ leaching in these rotations, as a basis for suggesting targeted mitigation strategies. We define “organic arable crop rotations” as cropping systems with associated crop and soil management commonly used in European farms dominated by arable cropping and following the European Council Regulation (EC) No 834/2007s on organic farming (Council of the European Union, 2007). Among others, the use of synthetic N-fertilizers and N-inhibitors are prohibited, but manure and/or short-term leys may be used in these rotations. We designate “non-organic crop systems” as arable cropping systems generally based on inorganic fertilizers, use of pesticides and often using narrow crop rotations, commonly called conventional farming.

Both globally and in Europe organic rotations are longer and more diversified than non-organic rotations (Barbieri et al., 2017). This is essential for nitrogen supply, and for pest and weed control (Stockdale et al., 2001). Barbieri et al. (2017) found that catch crops and undersown cover crops are 2.4 and 8.7 times more frequent in organic compared to conventional systems, respectively. They further found that the share of pulses and temporary fodder crops (such as alfalfa, clover and ryegrass) were higher in organic than in non-organic crop rotations, and that the difference between organic and non-organic crop rotations was greater in this respect in Europe than in North America and globally. They further found that more legumes are included in fodder crops, and in catch crops, under sown cover crops and intercropping than in non-organic rotations. In addition to plant derived N, organic N are applied in manure or other organic fertilizers and amendments. The great diversity of N mineralisation patterns among the organic fertilizers and crop residues is a challenge for farm management to synchronize the N release with plant N uptake. If N is released during periods with poor plant uptake, then the content of soil mineral nitrogen (SMN) and other easily available N can accumulate, creating a large risk of N losses through gaseous emissions or through leaching. Because N is mainly applied through plant residues and a limited amount of organic fertilizers in arable organic systems, the N-turnover from biological activity is crucial for the content and type of SMN. Plants and organic fertilizers are also important sources of soil organic carbon (SOC). We address how supply and quality of organic matter in above- and belowground residues and organic amendments influence availability and type of SMN and degradable carbon.

Kommentert [SH6]: 14

Kommentert [SH7]: 15

Kommentert [SH8]: 62

Kommentert [SH9]: 29

Kommentert [SH10]: 31

Increasing the content of SOC enhances the risk of N₂O emissions (Li et al., 2005). This is true whether the soil has a high content of SOC or the content is increased by additions of organic matter to the soil and is caused by the tight link between SOC and microbial N₂O production (Sahrawat and Keeney, 1986). Because the impact of SOC on N₂O emission is dependent on NO₃ content in soil (e.g. Weier et al., 1992; Li et al., 2005), we address how supply of N and carbon through organic inputs drive N₂O emissions in organic arable rotations. Based on IPCC (2006) most inventories and farm models assume that 1% of total N-input by fertilizers, manure and plant residues are emitted as N₂O-N. Skinner et al. (2014) found no correlation between total N-input and N₂O emission in organic systems. If this is a general trend, the total N-input cannot be used to estimate N₂O emissions from organic crop rotations. In non-organic cropping systems, peak fluxes of N₂O are commonly observed shortly after fertilization with mineral fertilizers in moist soil (Smith et al., 2012), whereas in organic crop rotations the highest fluxes are often observed after incorporation of plant residues (e.g. Brozyna et al., 2013; Krauss et al., 2017b; Nadeem et al., 2012; Pappa et al., 2011; Skinner et al., 2019). Because of enhanced content of SOM and thus a larger impact of background emissions of N₂O in organic versus non-organic cropping systems (Skinner et al., 2014), increased background emissions are likely to have a major impact in organic-crop rotations. In order to design good mitigation strategies, it is useful to know the relative importance of these two sources of N₂O emissions.

The main sources for NO₃ leaching are NO₃ from nitrification of plant residues and added organic matter, as NO₃-fertilizers are prohibited in organic systems. The crop N requirement dependency on soil organic matter turnover may lead to an asynchrony between crop nutrient demand and the mineralization of soil organic N, which enhances the risk of NO₃ leaching (Crew et al., 2005; Di and Cameron, 2002). Furthermore, soil cultivation for weeding or incorporation of plant residues have been shown to influence NO₃ leaching (Askegaard et al., 2011). We discuss the main drivers for NO₃ leaching in organic arable cropping systems and the associated preventative measures.

We address the following questions for organic arable crop rotations:

5. How does supply and quality of organic matter in above- and below ground residues and organic amendments influence availability of easily available N and degradable C?
6. How does supply of easily available N and degradable C drive N₂O emissions and how can these be mitigated?
7. Is there a lack of correlation between total N-input and N₂O emission in organic arable crop rotations, and are total N₂O emissions primarily driven by background emissions or by episodes with high N₂O fluxes following N additions?
8. What are the main drivers for NO₃ leaching in organic arable systems and how can the leaching be reduced?

Kommentert [SH11]: 21

Kommentert [SH12]: 20

2. Methodology

Based on the authors own field trials, literature databases and searches through Google Scholar, we compiled data on agronomic management, soil properties and yield level of organic arable crop rotations and measurements of SMN, N₂O emissions and NO₃ leaching from field trials relevant to organic crop rotations, climate and soil conditions in Europe. For SMN and NO₃ leaching, we used the available literature and the data in Tables S1 and S3 to explore the importance of the determining factors for SMN and NO₃ leaching, and to identify factors of importance for NO₃ leaching in organic arable crop rotations. For SMN and NO₃ leaching the structure of the available data did not allow meaningful statistical analyses.

Kommentert [SH13]: 22

For N₂O we used data presented in Table S2 to explore the impact of background emissions and episodes with high N₂O fluxes. We aimed to analyse the impact of high emission events of N₂O fluxes on the total N₂O emission. However, we lacked daily measurements, and we lacked data for yearly periods. It was thus not possible to identify the full impact of hot moments as done by Molodovskaya et al. (2012). Because of the differences in measurement period, it was also not possible to make direct comparisons between the different field trials. To overcome this, we used a regression model based on N₂O emissions in the actual period and peak N₂O flux within this period, resulting in the following fitted model:

$$\ln(C/P) = -1.34 + 0.83 \ln(N+2), R^2 = 0.66, n = 97$$

(Eq. 1)

C is N₂O emission in the measurement period (-278 to 8566 g N₂O-N ha⁻¹); P is duration of the period (38 to 490 days); C/P express the average daily N₂O flux (-1.3 to 53.2 g N₂O-N ha⁻¹ day⁻¹) and N is the highest N₂O-flux rate (0.1 to 605 g N₂O-N ha⁻¹ day⁻¹) in the measurement period. The total analysis and data used is given in S4. One negative value for average daily N₂O flux (-1.3) in a barley/pea crop was removed from the analyses because it would have had a large impact on the results.

We also calculated the percentage contribution of the highest daily N₂O flux of the total N₂O emissions in the measurement periods for all trials presented in Table S2 (n=97), and correspondingly the sum of the fluxes for the days with the five highest flux rates as a percentage of the total N₂O emissions.

The impact of the following potential explanatory variables on the highest daily N₂O flux rates: clay (soil clay content %), pH (soil pH), SOC (soil organic C, g kg⁻¹ dry soil), WFPS (soil water filled pore space, %), NO₃ (soil content of NO₃, kg NO₃-N ha⁻¹) and NH₄ (soil content of NH₄, kg NH₄-N ha⁻¹) and temp (soil temperature, °C) was calculated by stepwise regression (α to enter = 0.15; α to remove = 0.15) by Minitab 18. The selection of these variables was based on expected impact on N₂O emission (Sect. 4.1) and the data we were able to obtain from these studies. We used data from the following studies: Ball et al. (2007a), Baral et al. (2017), Brozyna et al. (2013), Chirinda et al. (2010), Krauss et al. (2017b), Li et al. (2015a), Nadeem et al. (2012b) and Pugesgaard et al. (2017). The N₂O flux data were log-transformed to achieve near normality and variance homogeneity: $\ln(\text{daily N}_2\text{O flux} + 2)$, where daily N₂O flux = g N₂O-N ha⁻¹ day⁻¹ is the highest N₂O flux rate during the actual measurement period (Table S2, Highest daily flux rate). The total analysis and data used is given in S5.

Based on the stepwise regression, we achieved the following fitted regression model:

$$\ln(\text{daily N}_2\text{O flux} + 2) = -1.18 + 0.04 \text{ clay} + 0.06 \text{ SOC} + 0.03 \text{ WFPS} + 0.01 \text{ NO}_3 + 0.05 \text{ temp}, R^2=0.71, n=66$$

(Eq. 2)

We used a literature review to explore the impact of total N added and, N and C added through organic inputs from living plants, plant residues and organic fertilizers on N₂O emission as we did not have enough data on added N to be able to include this in the regression analyses.

3 Drivers of SMN and degradable C in organic arable crop rotations

3.1 Supply and quality of soil organic matter

Return of crop residues to the soil is standard practice in both organic and non-organic arable production; however, because of the more diverse crop rotations in organic production systems, larger and more diverse inputs of herbage from legume-based green manures, leys, cover crops (CC) and intercrops are returned to soil than in non-organic

Kommentert [SH14]: 84

Kommentert [SH15]: 25

Kommentert [SH16]: 84

Kommentert [SH17]: 28

cropping systems commonly used in Europe (Gattinger et al., 2012). The N content in the crop residues of legume-based systems is typically higher than from non-legume systems (Watson et al., 2002). Commonly used external sources of organic inputs in organic cropping systems are animal manures and slurries, composts or biogas residues, and organic fertilizers based on animal manure or municipal waste (Løes et al., 2017). The great diversity of N mineralisation patterns among the organic fertilizers and crop residues result in a large variation in how much N and C that are rapidly degradable in the organic inputs.

Kommentert [SH18]: 33

Through the application of organic amendments and various crop residues from arable and forage crops, carbon (C) and N is applied to soil, and the soil organic matter (SOM) content is often higher in organic than in non-organic arable crop rotations (Marriott and Wander, 2006; Marinari et al., 2007; Gomiero et al., 2011; Gattinger et al., 2012; Aguilera et al., 2013; Hu et al., 2018). The quality of SOM differs between non-organic and the more diversified organic, arable crop rotation, with a higher share of labile SOM (Lynch, 2015) and thus easily degradable organic matter (C and N) in soils in organic crop rotations (Marinari et al., 2007; Marriott and Wander, 2006; Martyniuk et al., 2016). The higher content of degradable SOM in organic crop rotations is a valuable soil fertility asset as it provides a short-term pool for plant nutrient supply (Marriott and Wander, 2006; Martyniuk et al., 2016). SOM turnover rates vary with soil texture and climate, being higher when organic carbon is less protected from decomposers (low clay content) and in warm climates with suitable moisture (Burke et al., 1989).

Kommentert [SH19]: 36

3.2 Soil biological activity

Soil microbes contribute directly to plant residue decomposition and to mineralization and turnover of SOM, as well as earthworms and other soil fauna (Lubbers et al., 2013; Kuiper et al., 2013). The inflow of degradable organic matter provides substrate for soil organisms, and application of organic matter increases the growth of microbial communities, their enzyme activities and the microbial diversity compared to an unfertilised control or soil fertilized with only mineral fertilizer (Anderson and Domsch, 1989; Marinari et al., 2007; Thangarajan et al., 2013), although such changes in the topsoil may be slow processes (Petersen et al., 2013). Accordingly, higher biological activity has commonly been found in arable soils managed organically compared with non-organically (Mäder et al., 2002; Gomiero, 2013; Hartmann et al., 2015; Lori et al., 2017). In their meta-analysis, Lori et al. (2017) found that organic systems had 32% to 82% greater microbial biomass C, microbial biomass N, total phospholipid fatty-acids, and dehydrogenase, urease and protease activities than conventional systems. They found that when both organic and non-organic systems included legumes, the organic system displayed a higher microbial N content than the non-organic counterpart. In cases where only the organic systems contained legumes, the difference in microbial N between the two systems was even more pronounced. The abundance of earthworms can be twice as high in organic compared with non-organic systems (Pfiffner and Mäder., 1997; Filser et al., 1999; Hansen and Engelstad, 1999; Riley et al., 2008). More abundant earthworm populations are found when large amounts of animal manure or green manure are applied to soil (Hansen and Engelstad, 1999; Frøseth et al., 2014), when autumn ploughing is avoided (Pfiffner and Luka, 2007) and in the absence of tractor traffic (Hansen and Engelstad, 1999).

Kommentert [SH20]: 36

Kommentert [SH21]: 37

Kommentert [SH22]: 38

3.3 Soil N-dynamics within organic arable cropping systems

Crop N supply and SMN in organic farming relies largely on mineralisation of N in soil organic matter, N in organic amendments and crop residues and BNF of legume-based crops (Gattinger et al., 2012; Lorenz and Lal,

2016). A principal sketch for N-dynamics in organic systems is given in Fig. 1 and an overview of performance of selected drivers of SMN in organic and non-organic crop rotations are given in Table 1.




Fig.1. The principle of N-dynamic in organic crop rotations

Table 1. Performance of selected drivers of SMN in organic and non-organic cropping systems

Indicator	Organic arable systems	Conventional arable systems	Source
Main source of SMN	Mineralisation of crop residues and soil organic matter	Synthetic fertilizers	Lorenz and Lal (2016)
BNF in SMN supply	High	Low	Kayser et al. (2010); Pandey et al. (2017)
Size of the potentially mineralizable N pool	Usually high	Usually medium	Poudel et al. (2002)
Microbial biomass nitrogen content	High	Low	Lori et al. (2017)
Release of SMN	Slow and more continuous	Rapid with clear peaks	Poudel et al. (2002)
Concentration of SMN during growing season	Usually low	Usually high	Brozyna et al. (2013) Frøseth et al. (2014) Krauss et al. (2017b)

Concentration of SMN off growing season	Usually low, high only after termination of legume-rich crops	Usually high	<i>Hansen et al. (2007)</i> <i>Jończyk and Martyniuk (2017)</i>
--	---	--------------	--

Despite substantial inputs of N from BNF (Kayser et al., 2010; Pandey et al., 2017) and from organic amendments, the N-supply is often below optimum for plant growth in arable organic farming (Berry et al., 2002; Tuomisto et al., 2012). BNF may be a more important N source, than N from organic manures and fertilizers in organic arable cropping systems (Pandey et al., 2018).

The major release of plant available N from added organic matter depends on the C:N ratio, the mineral N content and degradability of the C of added plant residues manure, compost and other decomposed amendments. Bhogal et al. (2016) showed that for pig slurry and poultry layer manure with C:N_{org} of 9-12:1, up to 70% of the organic N was mineralized after five growing seasons, whereas in cattle slurry and straw-based farmyard manure with C:N_{org} of 10-21:1, only 10-30% of N was net mineralised. For crop residues with a high C:N content there will be initial immobilisation and the net mineralisation may only start very late or after growing season of the main crop (Li et al., 2015b). The mineralization process of the more stable N can continue over years to decades. Simulation modelling has shown that even over a 20-year period, only 10-15% of organic N in applied manure may be taken up by crops, the rest being lost or retained in soil organic matter (Berntsen et al., 2007). The high microbial activity and high content of organic matter affect N-cycling. Hu et al. (2018b) found a higher net N mineralisation of added organic matter in soils having a prehistory of use of CC, indicating positive legacy effects of CC use, which was attributed to a greater microbial biomass N in systems with use of CC.

The design of the rotation, as well as its management, influences the PMN pool. Working in three different organic arable systems, Spargo et al. (2011) showed that the PMN pool amounted on average for 315 kg N ha⁻¹. They showed that more diversified crop rotations resulted in larger PMN pools. Poudel et al. (2002) reported 112 and 56% greater PMN pool in the organic system in comparison to the conventional and low-input systems, respectively. Moreover, they observed slower and more continuous release of mineral N in the organic systems compared to the more rapid release of mineral N from synthetic N fertilizers applied in non-organic systems. Moyo et al. (2016) reported higher PMN in soil under wheat following a cut and mulched red clover ley compared to after a ley where the residues had been removed, indicating the importance of total N input to the soil. The importance of total N input in plant residues for crop N uptake was also observed by Petersen et al. (2013).

SMN is found to be very low under grass-clover leys (Table S1, Watson et al., 1993; Nadeem et al., 2012; Brozyna et al., 2013; Frøseth et al., 2014; Krauss et al., 2017b), since grasses quickly take up soil NO₃ in the root zone (Brophy et al., 1987). Frøseth et al. (2014) observed low levels of SMN in a 1-year grass-clover ley, irrespective of whether the herbage was mulched or removed. After termination of a ley, the concentration of SMN usually increases (Table S1, Ball et al., 2007a; Brozyna et al., 2013; Krauss et al., 2017b). Even in the year following termination of a ley, the content of SMN can still be high (Hansen et al., 2007; Jończyk and Martyniuk, 2017). Kayser et al. (2010) pointed out that N provided by spring ploughing of both 1-year grass-clover ley and 3-year grassland ley resulted in high concentrations of SMN (0-90 cm, 61 kg N ha⁻¹ and 95 kg N ha⁻¹, respectively) in the following autumn after harvest of spring triticale. Much of this SMN may not have been available for the crops during spring and is likely to have been mineralised after the end of the growing season. In contrast, low concentrations of SMN were observed in topsoil and subsoil after spring ploughing of a 1-year grass-clover ley in

Kommentert [SH23]: 39

Kommentert [SH24]: 40

Kommentert [SH25]: 42

Kommentert [SH26]: 66

Kommentert [SH27]: 67

four field trials in Norway, in late spring and in autumn after harvest of a spring barley crop (Table S1) and in the following year (Frøseth, 2014, 2016).

4 Drivers of N₂O emissions in organic arable crop rotations

4.1 Mechanisms for N₂O emissions

Many processes contribute to N₂O production in soils, but the dominant mechanisms for N₂O emission in terrestrial agricultural soils are the microbial processes of nitrification, nitrifier denitrification (as a result of incomplete nitrification) and denitrification (Firestone and Davidson, 1989; Butterbach-Bahl et al., 2013). Nitrification and denitrification are both biological processes, thus the same mechanisms will cause N₂O emissions in organic as in non-organic farming systems. However, fertilisation, crop and soil management practices differ substantially between these two systems (Sect. 3), and the relative importance of the various triggers therefore differ (Sect. 6). Nitrification is the microbial oxidation of NH₃ to NO₂ and ultimately NO₃, where N₂O is produced as by-product through some partially understood biotic and abiotic reactions of hydroxylamine (Anderson, 1964, Liu et al., 2017). Nitrifier denitrification occurs when NO₂ produced during nitrification is reduced to N₂O (by denitrifying organisms), instead of being oxidized to NO₃, under fluctuating oxic-anoxic conditions (Firestone and Davidson, 1989). Denitrification is the microbial anaerobic reduction of NO₃ via NO₂ to gaseous NO, N₂O and N₂, which are ultimately transported from soil to atmosphere. Denitrification is the main source of N₂O production in soils, as N₂O yield potential of denitrification is much higher (1-100%) than nitrification (0.1-1%) (e.g., Andersson et al., 1993, Butterbach-Bahl et al., 2013).

Kommentert [SH28]: 57

The ratio between the gaseous products of denitrification depends on NO₃ availability, oxygen availability in the soil and/or microsites, amount of easily decomposable carbon as an energy source, soil pH and microbial community structure (Bakken et al., 2012). Oxygen availability depends on soil microbial activity and gas diffusivity, which depends on soil moisture content, texture and density. Gas diffusivity is a promising predictor for N₂O fluxes from soils with varying bulk density as observed by Balaine et al. (2013), who found that the production of N₂O increased when the relative gas diffusivity was between 0.006 and 0.020 and the soil became anaerobic.

Kommentert [SH29]: 56

As explained above, risk of N₂O emissions increases as soil carbon content increases (Li et al., 2005). N₂O and N₂ production correlates with total organic C, water soluble C and mineralizable C in soil, but increased availability of C also decreases the ratio of N₂O/N₂ (Sahrawat and Keeney, 1986). Emissions from soil organic matter (background emission) will vary between years because of variations in temperature and precipitation (Hansen et al., 2014; Brozyna et al., 2013) and between different categories of crops due to different time windows with high SMN (Dobbie and Smith, 2003).

Kommentert [SH30]: 58

Organic amendments and plant residues that provide carbon that is easily decomposable by microbes may enhance microbial activity and deplete soil oxygen through enhanced soil respiration. In addition, degradable carbon is an energy source for denitrifying bacteria. In accordance with this, Köster et al. (2011) concluded that bacterial denitrification was the main process for producing N₂O during the first three weeks after application of biogas residues, and high carbon availability was an important cause for this. Li et al. (2016) concluded that denitrification was the main cause for N₂O emission after addition of legume-based residues. Several studies have shown higher rates of N loss through denitrification from soils treated with organic amendments such as manure, composts, and

plant residues when compared to unamended or mineral N treated soils (Thangarajan et al., 2013). In line with this, incorporation of residues by tillage increases soil respiration and N₂O fluxes because of microbial stimulation (Krauss, 2017a).

Low soil pH inhibits the activity of N₂O reductase enzyme and thus N₂O:N₂ ratio increases (Liu et al., 2010). At higher soil pH, the denitrification rate is higher, but the N₂O:N₂ ratio is lower as a greater part is completely denitrified to N₂. At low temperatures, nitrous oxide reductase is hampered (Holtan-Hartwig et al., 2002), but on the other hand, denitrification rates are also reduced (Butterbach-Bahl et al., 2013).

4.2 Legumes during active plant growth

In general, unfertilized legumes have small N₂O emissions during their growing period, particularly when grown in mixtures with non-legumes. Low N₂O emission are found during growth of grain legumes (Dusenbury et al., 2008; Jensen et al., 2012; Jeuffroy et al., 2013; Pappa et al., 2011; Rochette and Janzen, 2005), green manure crops and CC (Baggs et al., 2000b; Brozyna et al., 2013; Peyrard et al., 2016; Li et al., 2015a; Shelton et al., 2018) as well as for grass-clover leys (Baggs et al., 2000b; Ball et al., 2002; Brozyna et al., 2013; Skinner et al., 2019; Krauss et al., 2017b; Nadeem et al., 2012). This is consistent with low SMN concentrations during growth (Sect. 3) and negligible N₂O emissions associated with BNF by the legume rhizobium symbioses (Rochette and Janzen, 2005; Carter and Ambus, 2006). However, following termination or senescence of the legume crops, reactive N released from dying roots and nodules may lead to enhanced N₂O emission (Rochette and Janzen, 2005).

Surface mulching of harvested herbage may theoretically enhance N₂O emissions due to mineral N released from the herbage. However, several studies show that mulching of grass-clover herbage on the growing ley only causes a slight increase in N₂O emissions (Brozyna et al., 2013; Möller and Stinner, 2009; Nadeem et al., 2012). None of these studies measured ammonia volatilization from mulched herbage, which could have been a major loss of mineralized N corresponding to the findings of Larsson et al. (1998). Volatilized NH₃ will be redeposited elsewhere and may result in increased N₂O formation downstream (IPCC, 2006).

4.3 Crop residues

As outlined in Sect. 4.1, there is an enhanced risk of N₂O emission from agricultural soils when easily degradable carbon and N are simultaneously available, and denitrification is probably the main source for this. Because legume-based crop residues also increase SMN (Sect. 3.3), increased N₂O emissions have been reported in field trials whether the residues are from grain legumes (Jeuffroy et al., 2013; Pappa et al., 2011), grass-clover (Baggs et al., 2000b; Ball et al., 2007a; Skinner et al., 2019; Nadeem et al., 2012; Brozyna et al., 2013), intercropped clover (Pappa et al., 2011) or CCs (Baggs et al., 2000b; Peyrard et al., 2016; Pugesgaard et al., 2017). The increase of N₂O fluxes after incorporation of crop residues and other plant material might however, be small and a negligible part of the total N₂O emissions (Peyrard et al., 2016; Pugesgaard et al., 2017; Shelton et al., 2018).

The C/N ratio of incorporated herbage do affect N₂O emissions, with higher emissions expected from herbage having low C/N-ratio (Chen et al., 2013). From this, one should expect higher N₂O emission from legume residues than from cereals or grasses (e.g. Rochette and Janzen, 2005). However, Larsson et al. (1998) observed the same N₂O-N EF (1% of applied-N) from mulched alfalfa (C/N-ratio 11) as from mulched grass with a C/N ratio of 21, but higher EF than from a mulched grass with a low N-content (C/N-ratio 36, EF = 0.1%). The N₂O fluxes might

be high despite a high C/N-ratio when the carbon source is easily degradable as observed by fodder radish by Li et al. (2015a) (Table S2).

N₂O emissions may also be associated with previous incorporation of plant residues. In accordance with this, Skinner et al. (2019) observed enhanced N₂O fluxes after a maize crop succeeding a grass-clover ley. Measurement of N₂O fluxes shortly after incorporation of plant material, or measurements in the following year, only tell part of the story. Enhanced content of various fractions of SOM derived from crop residues, ley and CC (Sect. 3.2) are likely to increase the long-term background emissions of N₂O (Sect. 4.1). In a ten-year-old field experiment with and without legume-rich CC in the crop rotation, Pugesgaard et al. (2017) concluded that crop residues were important source of N₂O, and that mineralizable C, rather than N input, was the main driver for N₂O emission. Contrary to this, Peyrard et al. (2016) observed in a three-year low-input field trial that although N₂O fluxes increased for a few days after incorporation of CC, the contribution of such events to cumulative N₂O emissions were negligible. In their study, however, the CC treatments started when the N₂O measurements started. More studies in long-term experiments with continuous use of CC are needed to verify the actual impact of crop residues in a long-term perspective in various field situations, because addition of plant material to soil also affect soil structure, soil biological activity and N turnover.

4.3.1 Freeze/thaw – dry/wet

The mechanisms behind freeze/thaw have been comprehensively reviewed by Congreves et al. (2018) showing that the causes for N₂O emissions are different for these two mechanisms and that freeze/thaw has a larger impact on N₂O emissions in temperate agroecosystems than drying/rewetting. Wagner-Riddle et al. (2017) estimated that neglect of freeze/thaw emissions underestimate global agricultural N₂O emissions by 17 to 28%. Freezing/thawing of soil rich in organic matter and soil biota, or soil covered with plant residues may result in a N₂O boost as easily degradable C and N is released from cells through lysis after frost. As summarized in the introduction and in Sect. 3 these conditions are particularly relevant for organic crop rotations. Flessa et al. (1995) observed that 46% of total annual N₂O emissions from a sunflower crop, solely fertilized with farmyard manure (12 Mg ha⁻¹) occurred during December and January, mainly due to high N₂O peak fluxes (650 g N₂O-N ha⁻¹day⁻¹) after thawing of the first freezing period during winter. Correspondingly, Westphal et al. (2018) did not observe any enhanced N₂O fluxes after late summer incorporation of a ley dominated by alfalfa (0-10 g N₂O-N ha⁻¹ day⁻¹), but fluxes were greatly enhanced during spring thaw in the following year (60 g N₂O-N ha⁻¹day⁻¹).

When CCs are killed by frost, N₂O fluxes will increase during thawing of the soil because of release of easily degradable C and N in the plant material. Li et al. (2015a) observed significantly higher emissions from the frost-sensitive fodder radish rich in readily degradable carbon than from other less frost sensitive CCs. Winter emissions were even greater when fodder radish was harvested in late autumn (30 October), leaving only roots and stubble (Table S2). This suggests that N and C in roots of frost sensitive CCs can be an important source for N₂O emissions after thawing. Also, in leys frost may enhance N₂O emissions. Sturite et al. (2014) observed that enhance N₂O emission during thawing of a frozen grass-clover ley correlated with clover content in the ley.

Under drought conditions, the nitrification process prevails and N₂O is produced at very low rates. However, with rewetting easily degradable N and C is mineralized, resulting in increased N₂O fluxes. Hansen et al. (2014) observed that the N₂O flux increased with increasing clover content during rewetting of a grass-clover ley after drought. Hence, both freezing/thawing and rewetting may have a large impact in organic systems.

Kommentert [SH31]: 71

Kommentert [SH32]: 72

Kommentert [SH33]: 70

4.3.2 Soil and tillage effects

N₂O emissions associated with crop residues are affected by tillage depth and soil type. Large N₂O emissions have been observed when crop residues are placed near the soil surface in heavy soil (Peyrard et al., 2016, Krauss et al., 2017b) whereas in lighter soil types the highest emission has been observed either after rotary harrow (Baggs et al., 2000a) or after ploughing (Petersen et al., 2011). When the crop residues are squeezed and mixed with a rotary harrow, easily available N and degradable C become available for denitrifying bacteria in soil and the potential for denitrification is large. In line with this, Krauss et al. (2017b) observed high N₂O fluxes few days after weeds and crop residues were superficially incorporated with a rotary harrow in a moist calcareous clay soil (WFPS 80 %) (Table S2, highest observed peak in single plot 800 g N₂O-N ha⁻¹day⁻¹). Similarly, Peyrard et al. (2016) observed enhanced N₂O fluxes (max rates 60 g N₂O-N ha⁻¹day⁻¹) up to several days after crop destruction when crop residues (sunflower, wheat, faba bean) were mulched or placed near the soil surface of a calcareous-clay, but not by ploughing or mechanical weeding. Baggs et al. (2000a) observed higher N₂O fluxes when lettuce residues were incorporated by rotary harrow than by ploughing (peak of 67 g N₂O-N ha⁻¹day⁻¹).

Restricted gas diffusivity is another possible explanation for the observed lower N₂O fluxes with deep incorporation of crop residues in dense soil. With reduced gas diffusivity more N₂O are likely reduced to N₂ in accordance with a general trend of a larger ratio of N₂O-N/(N₂O-N+N₂-N) close to the soil surface and smaller fluxes deeper in the soil profile (Sahrawat and Keeney, 1986). Kuntz et al. (2016) observed a decreased O₂ concentration at 8 cm soil depth and a corresponding reduction of N₂O to N₂ with surface application of carbon rich material. As another example, Petersen et al. (2011) found in their loamy sandy soil that the largest fluxes were observed when residues were incorporated by ploughing compared to reduced tillage. Possible explanations for this could be that residues came directly into contact with mineral N from the injected slurry after ploughing, thus fostering enhanced microbial turnover of C and N and that in this soil, the aeration with O₂ was still available at plough depth.

Kommenter [SH34]: 75

4.4 Organic fertilizers

Organic fertilizers vary widely in the content and types of N and C compounds causing large variations in N₂O emissions after application. Animal slurries have a higher content of NH₄-N and contain more easily degradable N and C than solid manures and composts and are thus stronger triggers for rapid N₂O emissions shortly after application (Charles et al., 2017, Sect. 3). In accordance with this, Krauss et al. (2017b) observed higher N₂O emissions shortly after application of cattle slurry than composted solid cattle manure. Correspondingly, in a field experiment with spring barley fertilized with various organic slurries, Baral et al. (2017) observed the highest N₂O EF in the treatment with highest application of organic matter, and thus highest content of easily degradable C. Mejjide et al. (2007) and Chantigny et al. (2007) found that the use of digested slurry, with lower content of degradable C compared to untreated pig slurry, reduced soil N₂O emissions by 25 and 50% respectively.

The effect of organic fertilizers does depend on soil type and content of SOC. Degradable C applied with organic fertilizers will to a greater extent trigger microbial respiration and denitrification in a soil with low content of SOC than in a soil with high content of SOC (Chantigny et al., 2010, Pelster et al., 2012), whereas in a soil with high content of SOC the impact of easily available N is higher (Petersen et al., 2008). Because of a higher content of SOC and a higher share of labile SOM in organic crop rotations compared with non-organic crop rotations (Sect.

Kommenter [SH35]: 76,77 incorporated in the text

3.1), the short-term effect of organic fertilizers with a high content of degradable C on N₂O emission are likely lower in organic than in non-organic crop rotations.

Also, the absence of synthetic fertilizers means that organic fertilizers are likely to have a smaller short-term impact on N₂O emission in organic than in non-organic crop production. Charles et al. (2017) found in a meta-analysis that the N₂O EF was higher when soils received organic amendments in combination with synthetic fertilizers. They found EFs for liquid manures + synthetic fertilizers: 2.14 % (± 0.53), composts + synthetic fertilizers: 0.37 % (± 0.24), and the corresponding EFs for manure: 1.12 % (± 0.18) and compost: 0.00% (± 0.17).

However, the long-term impact of manures is not included in these EFs. In contrast to the short-term fertilizer effect, a long-term fertilization with organic fertilizers may enhance N₂O emissions through enhanced background emissions. Chang et al. (1998) observed that annual N₂O emissions increased with manure rate when different rates of solid feedlot manure and thus N-application were applied for 21 years. Their manure rate was far greater than would be applied under organic farming conditions, but the possibility for enhanced background N₂O emissions after long-term input of organic matter through manuring and application of crop residues should be considered. Krauss et al. (2017b) found that fertilization with slurry and manure compost increased annual N₂O emissions during winter wheat after more than ten years of differentiated management compared to sole slurry fertilization (mean values in the period (369 days), were 2.2 and 2.9 kg N₂O-N ha⁻¹, respectively). They related this to higher microbial biomass and SOC. Mean values for the upper 10 cm in soil were 28 and 30 Mg C ha⁻¹ for fertilization with slurry and manure compost and sole slurry fertilization, respectively.

Kommentert [SH36]: 78

4.5 Contribution of total N-input and high emission events to N₂O emissions

Skinner et al. (2014) concluded in a review that soil characteristics (soil N content) had a greater impact on N₂O emissions from organic production than the total-N input by fertilization. Lack of correlation between N₂O emissions and N fertilization in organic production corresponds to the more recent findings of Krauss et al. (2017b) and Pugesgaard et al. (2017). Pugesgaard et al. (2017) observed no significant correlation between N₂O emissions and N input in fertilizer/manure, for either annual emissions or spring emissions, but N₂O emissions were correlated with N input in residues from the previous main crop and CC. This agrees with the findings of Bouwman et al. (2002), van Groenigen et al. (2010), Peyrard et al. (2016) and Shcherbak et al. (2014), who observed low EF's when N fertilization was below optimum as commonly found in organic production systems.

Kommentert [SH37]: 79

As discussed earlier N₂O emission in organic crop rotations are driven by enhanced background emissions from long term input of organic matter, and episodes with enhanced N₂O emission after application of crop residues or organic fertilizer, but are N₂O emissions primarily driven by background emissions or by episodes with high N₂O fluxes? We used data from mainly organic field trials (Table S2) and one non-organic trial fertilized with organic fertilizers (Baral et al., 2017) to calculate the impact of the highest N₂O fluxes on the total N₂O emissions. In Frick (CH), Edinburgh (UK), Aberdeen (UK) and Ås (NO), the highest daily flux rates were 605, 211, 297 and 94 g N₂O-N ha⁻¹day⁻¹, respectively (Table S2). Because of the high flux rates, we hypothesised that high emission were responsible for a major part of the N₂O emissions from these systems. The single days with the highest fluxes correspond to 18% (65 days measurement period) in Frick, 2% in Edinburgh (161 days), 17 % in Aberdeen (38 days), and 2% in Ås (218 days) of the cumulated N₂O emissions in the measurement periods. The five highest daily N₂O fluxes corresponded to 22, 7, 55 and 5% of the N₂O emission in the measurement periods in these investigations, respectively. In field trials conducted on well-structured sandy loams at either Foulum or

Kommentert [SH38]: 82

Flakkebjerg in Denmark (Table S2, 105 to 365 days), peak N₂O fluxes from one or five days, however, only constituted from <1-8% and 5-14% of the total emission in the periods, respectively. The highest daily flux rate in these trials was only 78 g N₂O-N ha⁻¹day⁻¹. This was in a non-organic treatment heavily fertilized with cattle slurry and digested sewage sludge (476 kg total N ha⁻¹, Table S2 (Baral et al., 2017)). From this we reject the hypothesis that high emission events were responsible for a major part of the N₂O emissions from these systems, rather background emissions seemed to be the major N₂O source. However, a simple regression model (Eq. 1) showed that the average daily N₂O flux correlated positively with episodes with high N₂O fluxes. From this we can conclude that when the conditions for high N₂O fluxes are met for one or more days, there is a large chance for high total N₂O emissions in the period. The small peaks in the Danish field trials reveals that well-drained sandy soils promote rapid water infiltration and good gas-diffusivity and in turn low N₂O emissions.

Kommentert [SH39]: 83

We wanted to explore the impact of soil conditions on the highest N₂O flux peak. In a stepwise regression (Eq. 2), the content of clay, SOC, NO₃-N in soil and soil temperature had significant positive impacts on peak N₂O fluxes in the selected investigations (Table S2). (P_{clay}, P_{SOC} < 0.001, P_{NO₃}, P_{temperature} < 0.01, P_{WFPS} < 0.05). The content of NH₄-N in soil did not affect peak N₂O fluxes. These findings indicate that denitrification is the main cause for high N₂O-flux rates in these studies. To visualize the impact of different factors on the mean and the highest N₂O flux, and the contribution of the 5 highest daily fluxes to the cumulated N₂O emissions in the measurement period we grouped the investigations according to % clay in soil, N added with organic fertilizers, SOC, soil pH, experimental period, type of crop and mean daily precipitation in the period (Fig. 2). Because of lack of information, we were neither able to include total-N input nor soil porosity in the regression analyses and the box-plots.

Kommentert [SH40]: 85

In line with the findings of Skinner et al. (2014) we did not find a clear impact of N added with organic fertilizers, but the soils with the lowest content of SOC, showed the lowest mean and lowest peak N₂O emission. This supports a hypothesis that the SOC content of the soil, and thus the content of SOM, are important drivers for N₂O emission. At high SOC contents other factors seem to be more important. The impact of type of crop on N₂O emissions do mainly follow the trend we have seen in other investigations (Sect.s 5.2, 5.3) with higher N₂O emissions when cereals succeed a short-term ley (grass-clover or clover) and that mixtures with monocotyledons and legumes do not have higher N₂O emissions than cultures with only monocotyledons. The high residual effects of previous leys supports the idea that background emissions are the main driving force for N₂O emission in organic crop rotations. The high N₂O emissions (up to 600 g N₂O-N ha⁻¹ day⁻¹) from one occasion in Switzerland with incorporation of weeds in wet soil have a large impact on all box-plots (Fig. 2). None of the other groupings in Fig. 2 showed any clear causal effect on N₂O emissions.

Kommentert [SH41]: 81

The low effect we observed of pH on N₂O emissions (Eq. 2, Fig. 2) is contradictory to enhanced N₂O emission often observed in acid soil caused by hampered N₂O reductase enzyme (Bakken et al., 2012). A reason for this could be that in these soils and with these managements other factors meant more for N₂O emissions than pH. Although denitrification was likely to have been the main cause of N₂O fluxes at the highest N₂O flux rates, the soil NO₃ concentration are often low between the episodes with high N₂O flux rates, as observed by Brozyna et al. (2013), Chirinda et al. (2010), Li et al. (2015a), Nadeem et al. (2012) and Pappa et al. (2008).



Fig.2. Box-plot (mean -, median x, upper and lower quartile) for (a) mean N₂O flux in the period (g N₂O-N ha⁻¹ day⁻¹) and the highest daily flux rate in the same period (g N₂O-N ha⁻¹ day⁻¹) and (b) % contribution of the 5 highest daily fluxes to cumulated N₂O emissions in the period. N=number of sites, n=number of observations

4.6 Impacts of earthworms

Abundant earthworm populations in organic crop rotations (Sect. 3.3) are likely to influence N₂O fluxes as they significantly affect mineralization and reduction of N compounds to N₂O and N₂ (Prieto, 2011). N₂O is emitted from intestinal microbes but is also released from nitrates emitted in body fluids in the earthworm gut as well as from casts, middens and burrows (Prieto, 2011). On the other hand, earthworms improve soil porosity and aggregate stability (Bronick and Lal, 2005) and thus gas-diffusivity and water infiltration in soils, which will reduce N₂O emission. Epigeic species (living near surface and feeding on surface litter) and anecic species (deep burrowing) are well known to enhance N₂O production, because they feed directly on decomposing herbage (Evers et al., 2010; Nebert et al., 2011; Lubbers et al., 2011). Endogeic earthworms that feed on SOM particles are most common in cultivated arable soils (Hansen and Engelstad, 1999), and they do not increase denitrification (Postma-Blaauw et al., 2006). There are too few published results to robustly predict the impact of earthworms in arable organic crop rotations on N₂O emissions as this will depend on local climatic and edaphic conditions.

Kommentert [SH42]: 89

5 Drivers of NO₃ leaching in organic arable crop rotations

5.1 Mechanisms for NO₃ leaching

NO₃ leaching is an abiotic process driven by diffusion and convection (e.g. Johnsson et al., 1987), where NO₃ is transported out of the root zone along with the downward water flow. In addition to soil water content, soil texture and structure are important in determining leaching rates. Fine textured soils have slower infiltration rates than coarse textured soils, and porous sandy soils are most vulnerable to leaching, also because these soils often have more shallow rooting depths than loamy soils (Askegaard et al., 2005). The impact of soil type was clearly demonstrated by Askegaard et al. (2011) who found that, depending on soil type (coarse sand>loamy sand>sandy loam) and precipitation, 20-100 kg N ha⁻¹ yr⁻¹ was leached on average for the crop rotations. In their study, the location on coarse sand had 200-300 more mm rainfall per year than the other locations. The leaching was considerably higher than for Swedish clay soils: 20 kg N ha⁻¹ yr⁻¹ (Stenberg et al., 2012).

Due to its high mobility in soil, NO₃ can easily be lost from the agroecosystem by leaching during periods of high drainage rates. A well-developed active root system enhances NO₃ uptake, while a poor root system will not utilize all the NO₃ within the soil profile (Dunbabin et al., 2003). NO₃ remaining in the soil after the growing season of crops, or mineralised subsequently, will greatly increase the risk of leaching loss. This could happen outside the growing season, but also when there is poor crop establishment caused by unfavourable seedbed structure or from crop failure caused by diseases or pests (Stenberg et al., 2012). If crop failure coincides with rainy weather, the risk of severe NO₃ leaching is large. This was observed by Torstensson et al. (2006) and De Notaris et al. (2018) in organic farming systems, where potato growth was restricted due to early crop termination following disease outbreaks. Torstensson et al. (2006) determined the annual leaching in the potato year to be 75 kg N ha⁻¹ after green manure and 98 kg N ha⁻¹ after pea/barley (Table S3), whereas De Notaris et al. (2018) observed substantially higher leaching rates. They measured 213 kg N ha⁻¹ yr⁻¹ leached when the potato followed green manure and 133

Kommentert [SH43]: 90

kg N ha⁻¹yr⁻¹ after grain legumes in a year with early occurring potato late blight. This was substantially higher than in previous years (140 and 78 kg N ha⁻¹, respectively).

Extreme rainfall events and/or periods with drought can significantly affect leaching for a variety of reasons. A field experiment over 13 years in the UK showed that N leaching in winter from fertilized grass (non-organic) was highly correlated with the preceding summer's soil moisture deficit, with the highest losses following dry summers (Tyson et al., 1997). In this case, poor grass growth due to drought led to a buildup of NO₃ from unused fertilizer present in the autumn. Prolonged mineralization of organic fertilizers or crop residues due to drought may also lead a similar situation in organic farming systems. Tosti et al. (2016) found, under Mediterranean rainfed conditions, that the risk of NO₃ leaching was mainly at the onset of drainage due to rainfall, i.e. at the initial stage of growth, and being typically variable among years depending on timing of heavy rains. Thus, amendments applied at pre-crop stage would be a risky practice for NO₃ leaching. Most N leaching studies in organic farming in Mediterranean environments focused on row and vegetables crops (e.g. Campanelli and Campali, 2012), because these systems are most demanding in N inputs and thus have higher N applications and potential leaching than in common arable crops.

Kommentert [SH44]: 93




Fig. 3. Lower and upper level of NO₃ leached (kg NO₃-N ha⁻¹yr⁻¹) in various field investigations: with or without cover crops (CC), different N amount applied, and with or without grass-clover ley. The data used are given in Table S3.

5.2 Legumes

In a crop rotation with a large contribution through BNF, some of the N inputs will be retained in crop residues and in particular in mulched green manures (Frøseth et al., 2014). In their review, Crews and Peoples (2005) found that when the N input was based on BNF, the proportion of the N retained in the soil was higher (58% of legume N) than in the fertilized systems (31% of fertilizer N). From this, it is likely to assume that the risk of N release outside the crop growing period are high in rotations without legumes. In their meta-analyses of crop yield and N dynamics as influenced by CCs, Tonitto et al. (2006) concluded, however, that on average, NO₃ leaching was reduced by 40% in legume-based systems relative to conventional fertilizer-based systems. The reason for this is

probably the large difference in N input between legume-based systems relative to conventional fertilizer-based systems. The response of NO₃ leaching to N input in fertilizer, manure and residues may also differ between sites due to soil type and precipitation (Pandey et al., 2018).

5.2.1 Grain legumes

Nitrate leaching reported from crop residues of grain legumes vary. Highest values are found when grain legumes are grown in monoculture rather than in mixtures with e.g. cereals, and when CCs are not used (Plaza-Bonilla et al., 2015). Stenberg et al. (2012) observed higher NO₃ leaching after faba bean compared to after non-leguminous crops. On a clay soil in Sweden, they observed an average leaching of 20 kg N ha⁻¹ yr⁻¹, which was twice that for spring cereals. On average over three years on loamy sand in Denmark, De Notaris et al. (2018) reported about twice as high NO₃ leaching following a barley/pea intercrop compared with spring wheat or spring barley. In a sandy soil in northwest Germany, Kayser et al. (2010) observed that 83 kg N ha⁻¹ leached in triticale following field bean. In a worst-case scenario, Askegaard et al. (2011) observed annual NO₃ leaching of 270 kg N ha⁻¹ during and after a lupin crop on a coarse sandy soil in a situation where the lupin crop did not ripen, leaving a large amount of N in crop residues (same experiment as De Notaris et al. (2018)). Pappa et al. (2008) observed very low N-leaching during and after a barley/pea intercrop, but they observed a significant effect of the pea cultivar on N leaching in the autumn and winter period.

5.2.2 Forage legumes

Many authors (Kayser et al., 2010; Neumann et al., 2011; Stalenga and Jończyk, 2008) emphasize that one of the most critical times for NO₃ leaching in organic crop rotations occur after soil incorporation of a grass-clover ley. N leaching is low during the growing period of grass-clover leys (Kayser et al., 2010), but because of the large amounts of mineralized N after termination of a grass-clover ley (Sect. 3.3), the risk of NO₃ leaching is large for one to two years after termination of these crops (Berntsen et al., 2005). The leaching may occur shortly after ley termination, during winter, or during the succeeding seasons, depending on time of incorporation, quality of the herbage, the weather and the crop sequence.

Stenberg et al. (2012) observed higher NO₃ leaching following termination of a grass-clover ley than following faba bean, but values were still low (4 kg N ha⁻¹ higher in average). They found the highest leaching when the grass-clover ley lasted for two years (up to 40 kg N leached ha⁻¹ yr⁻¹). This corresponds to the finding of Kayser et al. (2010), who observed greater NO₃ leaching during the winter after spring incorporation of a three-year ley than after a one-year ley (121 versus 83 kg N ha⁻¹, Fig. 3). However, the crop yield of triticale yielded much better after the three-year ley than after the one-year ley. The percentage share of clover (0-5, 30 and 50%) did not influence the amount leached after ley termination, neither the crop yield. Eriksen et al. (2008) measured NO₃ leaching after 1 to 8 year old grass/clover leys, but found that the length of the ley had no effect on NO₃ leaching. Stenberg et al. (2012) observed that cereals succeeding grass-clover ley had nearly double yearly N leaching compared to cereals with no legume pre-crop. The highest NO₃ leaching occurred after cultivation of a winter rye (48 kg N ha⁻¹ yr⁻¹). De Notaris et al. (2018) observed that NO₃ leaching during cultivation of spring wheat was about 50 kg N ha⁻¹ higher when the spring wheat succeeded a two-year green manure crop (alfalfa or grass-clover) than when it succeeded a grain legume (107 versus 50 kg N ha⁻¹). Similarly, Askegaard et al. (2011) observed peaks in NO₃ leaching in autumn and winter after ploughing-in grass-clover ley. At crop rotation level, inclusion of grass-clover

or alfalfa on 25% of the area increased NO_3 leaching rate by 6-12 kg N ha^{-1} (De Notaris et al., 2018). Forage legumes may also be undersown as intercrops to increase soil fertility in organic crop rotations. Pappa et al. (2008) found that clover intercropped in spring barley only increased annual NO_3 leaching by 1-2 $\text{kg NO}_3\text{-N ha}^{-1}$.

5.3 Cover crops

CCs are grown between main crops to minimize NO_3 leaching. Many field trials in non-organic systems have shown reduced leaching using CC (e.g., Rasse et al., 2000; Torstensson and Aronsson, 2000; Constantin et al., 2010; Valkama et al., 2015). This is also the case for organic crop rotations (Tonitto et al., 2006; Askegaard et al., 2011; Tosti et al., 2014; Tosti et al., 2016; De Notaris et al., 2018). The reduction in NO_3 leaching can be substantial. Studies in Nordic countries report reductions of 50-60% in N leaching (Askegaard et al., 2011; De Notaris et al., 2018; Torstensson and Aronsson, 2000, Fig. 3). If the cash crop fails, the effect of CCs on reduced leaching can be even higher. In a year where potato late blight caused crop failure in potato, the CCs reduced N-leaching by 95% when the potato succeeded a grain legume (from 133 to 6 kg N ha^{-1} leached), and by 92% when the potato succeeded a green manure ley (from 213 to 17 kg N ha^{-1} leached) (calculated from Table S3, De Notaris et al., 2018).

De Notaris et al. (2018) concluded that the use of CCs had a larger impact on leaching than a substantial variation in N surplus between alternative cropping systems. In three long-term field trials (13-17 years) in Northern France, Constantin et al. (2010) observed that CCs were the most efficient management option for reducing NO_3 leaching (from 36 to 62%). Good establishment and growth of the CC is essential to obtain sufficient uptake of SMN and thus reduce NO_3 leaching. Stenberg et al. (1999) found no significant reduction in NO_3 leaching during winter from a ryegrass CC that was undersown in spring. They explained this by poor CC establishment. De Notaris et al. (2018) also observed occasions with very small effect of CCs on NO_3 leaching. They related this to CC growth and identified threshold values in CC above-ground biomass determined in November, above which N leaching was reduced to a stable low level. NO_3 leaching from spring wheat averaged 15 $\text{kg N ha}^{-1} \text{ yr}^{-1}$ with CC biomass above 0.9 Mg ha^{-1} , and 41 $\text{kg N ha}^{-1} \text{ yr}^{-1}$ with CC biomass below 0.9 Mg ha^{-1} . In potatoes, the average N leaching was 11 and 80 $\text{kg N ha}^{-1} \text{ yr}^{-1}$ with CC biomass above and below 1.5 Mg ha^{-1} , respectively.

Including legumes in CC mixtures does not seem to reduce the ability of CCs to reduce NO_3 leaching (Tonitto et al., 2006; Tosti et al., 2014; De Notaris et al., 2018; Shelton et al., 2018). In a field trial with barley, hairy vetch and a 50:50 mixture of both species as CC, Tosti et al. (2014) found that, in all years, the barley/vetch mixture decreased N leaching to the same level of pure barley, both during its own growing cycle and after CC incorporation into the soil. De Notaris et al. (2018) concluded that the same degree of reduced NO_3 leaching was obtained with legume-based CCs as with non-legume CCs. The CC was either undersown in spring or after harvest of the main crop. The undersown legumes were white clover and red clover, and winter vetch was used in the mixture sown after harvest. Shelton et al. (2018) found greater NO_3 leaching from intercropped hairy vetch than from simultaneously grown wheat and wheat/hairy vetch mixture. When the CC is a pure stand of legumes, the CC does not necessarily reduce N leaching (Tosti et al., 2014; Valkama et al., 2015; Shelton et al., 2018). Tosti et al. (2014) concluded that hairy vetch sown as a pure crop in autumn showed high BNF, but no NO_3 leaching mitigation effect as compared to bare soil. Valkama et al. (2015) found in their meta-analysis of Nordic studies of undersown CCs that legumes (white and red clovers) in pure stand did not diminish the risk of NO_3 leaching.

5.4 Tillage

Kommentert [SH45]: 92

Tillage stimulates soil N mineralization, at least in the short term. Tillage is also often associated with soil incorporation of plant residues that may lead to net N immobilisation or mineralisation, depending on quality of the residues. Timing of tillage is therefore crucial for the fate of the mineralised N, whether SMN becomes available for crop uptake or subject to leaching. In general, incorporation should consider soil type, climate conditions and type of herbage (C/N ratio). Thorup-Kristensen and Dresbøll (2010) suggested late incorporation of CCs in high rainfall areas on sandy soils, and earlier in low rainfall areas on NO₃ retentive soils. Field studies have shown rapid N mineralization from N-rich plant material, even at low temperatures (Breland, 1994; Thorup-Kristensen and Dresbøll, 2010). Spring incorporation has therefore been recommended to increase N recovery by subsequent crops. However, under Scandinavian conditions, there may still be a deficit in crop-available N, even after spring incorporation of a green manure ley (Frøseth et al., 2014; Känkänen et al., 1998). Under such conditions, Torstensson and Aronsson (2000) suggest that late autumn incorporation of CCs, instead of spring incorporation, will be preferable with respect to N availability for the subsequent crop and will not substantially increase NO₃ leaching. Under Mediterranean climatic conditions, characterized by mild rainy winters and warm to hot dry summers, there is a risk of NO₃ leaching if residues are incorporated prior to the wet season. No studies were found that measured NO₃ leaching in relation to timing of tillage in organic arable farming under these conditions.

In organic crop production, the timing of cultivation for mechanical control of perennial weeds may conflict with the aims of high N use efficiency (Melander et al., 2016). Askegaard et al. (2011) found, on sandy soils in Denmark, that the management of crop and soil during autumn was the main determinant of N leaching. Stubble harrowing in autumn for controlling perennial weeds, followed by bare soil during winter, led to an average of 25 kg N ha⁻¹ more leached than for soils left untouched with a cover of weeds/volunteers. NO₃ leaching increased with increasing number of autumn soil cultivations.

Reducing tillage intensity may also enhance the need for weed management, and thereby the risk of NO₃ leaching. In their meta-analysis comparing different reduced tillage intensities in organic farming, Cooper et al. (2016) found that the weed incidence was consistently higher, by about 50 %, when tillage intensity was reduced, although this did not always lower the yields. Compared to conventional tillage, reduced tillage may reduce NO₃ leaching, but this depends on the establishment and growth of the succeeding crop (Känkänen et al., 1998).

Kommentert [SH46]: 91

Bare fallow has traditionally been a method to control perennial weeds by repeated tillage of superficial or deeper top soil layers. In practical organic farming, a bare fallow can sometimes be used before or after the main crop if conditions have promoted perennial weeds. However, if carried out in the growing season, soil temperature and moisture conditions favour soil microbial activity and therefore the build-up of SMN, which greatly increases the risk of NO₃ leaching (Borgen et al., 2012).

6. Key drivers of N₂O emissions and NO₃ leaching and suggested mitigation strategies

Easily available N and degradable C added through organic inputs enhances risks of high N₂O emissions in the short term because of enhanced biological activity (Sect. 3.3) and increased denitrification potential (Sect. 4.1), and in the longer term because of higher soil content of N and C in labile organic matter (Sect. 3.1) contributing to N₂O emissions from mineralized SOM (background emission, Sect. 4.1, 4.3, 4.4, 4.5). There is no strong correlation between total N-input and N₂O emissions in organic arable crop rotations (Sect. 4.5), which is another

indication of the strong impact of background emissions in these systems (Sect. 4.5). The same is also true of the relatively low impact of episodes with high N₂O fluxes on the total N₂O emission in various investigations (Sect. 4.5, Fig. 2).

From this, we postulate that background emissions in most organic crop rotations are a more important driver for N₂O emissions than episodes with high N₂O flux rate (hot moments). Nevertheless, reducing periods with hot moments of denitrification and thus high N₂O fluxes rates are important for reducing total emissions. One mitigation measure is to avoid large applications of residues from crops comprising easily available N and degradable C like clover or Brassica (Sect. 4.3). Mulching of grass-clover on top of a ley seems to only slightly increase N₂O-emission compared with no mulch (section 4.2). Incorporating organic material in the soil surface does not reduce the risk for high N₂O fluxes and N₂O emissions are often higher after surface application than after ploughing (Sect. 4.3.2). Mixing by rotary harrow can enhance emissions, particularly in moist soil. If the crop residues are removed, and composted, used in a biogas plant or treated by other methods before targeted soil application, such events can be avoided. To avoid that these measures result in that GHG emissions only are moved to another place, this requires measures that minimise GHG emissions during and after treatment of the plant residues (see sect 7). The largest stimulation of denitrification by application of plant residues or other organic material with a high content of degradable C seems to be in soils with a low content of SOC, particularly if the soil has a high NO₃ content of or easily available N is applied (Sect. 4.4). If crop residues consist of less degradable C and easily available N like straw from cereals and grain legumes, they do not stimulate rapid denitrification and enhanced N₂O flux rate (Sect. 4.3), and the N₂O flux is likely reduced (Xia et al., 2018).

Background emissions are highly influenced by content and mineralization patterns of SOM and release of SMN and degradable C (Sect.s 3 and 4.1). Thus, weather conditions will have a large impact on the background emissions, which are likely to be higher in warm and moist years, than in cold or dry years. These conditions cannot be influenced by farmers, but it is important to develop strategies to decrease the content of SMN to reduce the risk of N₂O emissions and NO₃ leaching in periods with risk of SMN accumulation. The choice and sequence of crops and the use of CC is a central strategy for simultaneously mitigating N₂O emissions and NO₃ leaching in organic systems. Poor timing of N released from crop residues from preceding crops and crop N uptake (Sect. 3.3 and 5.2) are major challenges. There is a need to reduce periods with bare soil, although this may conflict with the need for mechanical weed control (Melander et al., 2016).

CCs take up surplus SMN and protect soil from erosion, they reduce NO₃ leaching, and during growth CCs do not contribute to N₂O emissions (Sect. 4.2). However, in areas with frost, frost sensitive crops like fodder radish, will release easily available N and degradable C during thawing, and thus stimulate denitrification (Sect. 4.3.1). The choice of CC must be adapted to local conditions for ensuring good establishment and appropriate tolerance to frost and droughts. A mixture of legumes and non-legumes (for instance grasses or cereals) are just as efficient for reducing N leaching as sole non-legume CCs, whereas sole legumes are not as efficient (Sect. 5.3). CCs containing legumes have a lower C:N ratio than CC without legumes, which enhances the N fertilizer value for the following crop (Li et al., 2015b).

The knowledge about the rooting pattern of different crop species can be used as a tool for designing crop rotations that achieve higher N use efficiency and thereby reduces the risk of NO₃ leaching. Deep-rooted crops, especially tap rooted ones, can recover NO₃ from deeper soil layers before and after more shallow-rooted cash crops, such as

leak (Thorup-Kristensen, 2006). As shown by Fan et al. (2016), the root distribution and rooting depth may differ between varieties, although plant breeders do not normally select crops based on the root system.

For the choice of crop, species and varieties should be well-adapted to the climate conditions on the farm and the soil fertility level. Crops in good conditions also compete better against weeds. This decreases the need for soil management to achieve weed control, and thereby reduces the risk of NO₃ leaching. Timing of release of N from residues and amendments and crop uptake are crucial for minimizing the risk of NO₃ leaching. This can be achieved by timing of soil tillage and incorporation of residues. In any case, the effect of mitigation strategy is highly dependent on soil type and precipitation.

Because of the large impact of poorly aerated soil on N₂O emissions (Sect. 4.1) and plant growth, measures should be taken to improve and maintain a good soil structure. In organic production, soil fauna, microorganisms and the development and maintenance of soil structure, are supported by crop rotations that include legumes or grass-clover leys, use of CCs and application of organic fertilizers. Even so, traffic and tillage under wet soil conditions are damaging to soil structure and should therefore be avoided.

A way to reduce N₂O emissions and NO₃ leaching per unit produced is to increase yields in organic production as more land is commonly needed per unit product in organic than non-organic production (De Ponti et al., 2012, Meier et al., 2015). However, as discussed by Rööös et al. (2018) this is not straight forward as many of the available measures have negative side effects. More targeted and thus efficient use of N applied through crop residues and organic fertilizers seems to have co-benefits in terms for higher productivity as well as reduced N₂O emissions and NO₃ leaching.

7. Research and innovation needs

The knowledge of N cycling and losses in arable organic farming is constrained by few investigations that have adequately quantified the magnitude and timing of N₂O emissions and NO₃ leaching. Many of the reported results are from a few experiments, including a long-term experiment in Denmark (Olesen et al., 2000). Such long-term experiments are important for quantifying N transformation and loss processes in organic cropping systems, where the legacy effects of N in plant residues and other organic amendments are often considerably larger than in non-organic systems. Lessons from these experiments can help to develop methods and technologies to improve synchronisation of N released from crop residues and soil organic matter and the N demand of cash crops. However, there are few long-term experiments with organic arable crop rotations, and even fewer have been systematically used for quantifying the N cycling processes. Therefore, research on existing and new long-term experiments covering relevant soil and climatic conditions should be coordinated to make data and facilities available for research on N-dynamics, including N₂O emissions, NO₃ leaching, NH₃ volatilization and the fate of added N. In this way, useful lessons can be learned on the impact of long-term management on the N-dynamics. The availability of historical data on N-input and crop management is a prerequisite to achieve this, such data could be made available and structured through a coordination of long-term experiments.

CCs have been shown to efficiently reduce NO₃ leaching (Sect. 5.3), whereas they can reduce or increase N₂O emissions depending on the conditions (Sect. 4.2, 4.3). To optimise the use of CCs for crop productivity, environment and climate, improved knowledge is needed of locally adapted crop rotations and CC that maintain living plants that can take up available N and simultaneously reduce weed pressure.

Simulation models can be used to quantify N transformation and loss processes and how different management practices affect N storage in SOM, as well as short-term effects of crop residues and organic fertilizers on N₂O emissions and NO₃ leaching. However, process-based models fail to accurately simulate the impact of grass/clover leys on turnover of N and C (Doltra et al., 2019, Frøseth, 2016). To our knowledge, no model of field scale does include all C and N turnover processes, such as phyllodeposition and rhizodeposition. These processes may be of greater importance in organic than non-organic systems, because of the greater emphasis on fertility building measures. This may lead to underestimation of the amount of N and C returned to soil. Improving models on these aspects requires more comprehensive data on all C and N inputs and flows in the cropping systems. This can only be achieved through studies using isotopic labelling of C and N, and consideration should be given to conducting such studies also in the long-term experiments. There is further uncertainty on the release of N from legumes and thus the impact of legume residues on N₂O emission and NO₃ leaching (see Sect.s 4.2, 4.3 and 5.2). More research on the mineralization pattern of various legume residues would therefore be useful.

Heterogeneity is another aspect that makes it difficult to predict availability of N and C in organic systems and thus N₂O emission. In addition to the soil heterogeneity, variability is created from incorporated crop residues, CC, short term leys and organic fertilizer that are seldom evenly distributed. Uneven impact of frost thawing / drying rewetting will add to this variability and make it difficult to predict hot spots and hot moments of N₂O emissions. There is also a need to develop technologies to better measure this heterogeneity and to take account for the heterogeneity in N dynamic models adapted to organic crop rotations.

Composting of crop residues and manures is common on organic farms and is a way to make the organic matter more resistant and useful as a soil amendment. However, composting manure without emissions of N₂O and other greenhouse gases is challenging (Chadwick et al., 2011) and needs to be addressed to develop a more comprehensive picture of greenhouse gas emissions from organic arable production and how to mitigate them. Neither composting nor biogas fermentation are always feasible. Improved understanding of different pathways for microbial degradation of crop residues is needed as well, to better manage crop residues and organic waste for C and N retention properties.

8. Conclusions

Organic arable production is based on diverse crop rotations, N inputs through BNF in legumes, external inputs of manure and compost, and recovery of excess N in grassland and cover crops. This results in considerable inputs on N in organic matter from plant residues, green manures and application of organic fertilisers and soil amendments that enhance the content of total and labile SOC in organic arable production compared to non-organic production. When the conditions for mineralization of SOC are met this will lead to a high availability of easily available N and degradable C, and an enhanced content of SMN, if available N is not taken up by growing plants. Conditions with high content of SMN and degradable C provide hot moments for high N₂O fluxes. Degradable C will increase microbial growth and thus O₂ consumption in soil leading to anaerobic conditions and provide suitable conditions for denitrification of available NO₃. In organic arable crop rotations background emissions are more important in most situations than episodes with high N₂O fluxes for the total N₂O emission from organic arable crop rotations. SMN from mineralization of SOM and plant residues in combination with periods of bare soil or sparse plant growth and precipitation surplus provide drivers for NO₃ leaching.

Main mitigation strategies are targeted use of growing plants to take up surplus soil mineral N from root zone, and targeted use of crop residues and organic fertilizers to synchronise the availability of N with the crop demand. Continued use of CCs has a proven ability to reduce NO₃ leaching from organic arable crop rotations but does increase N₂O emissions in some situations. A CC mixture of legumes and non-legumes (for instance grasses or cereals) is just as efficient as sole non-legumes and has a better impact on soil fertility than non-legumes. More research is needed to develop locally adapted crop rotations and CC mixtures that at the same time reduce N₂O emissions and NO₃ leaching.

Author contributions: Sissel Hansen: Prepared the manuscript with contributions from all co-authors, Supplementary material Table S2, Fig. S2; Randi Berland Frøseth: Structure and content of the MS, mainly N leaching, complementary data from Norwegian field trials in Fig. 3, Table S1, Supplementary material Table S3; Maria Stenberg: Content of the MS, mainly N leaching, Supplementary material Table 3 and complementary data from Swedish field trials to this Table; Jaroslaw Stalenga: Content of the MS, mainly SMN, Supplementary material Table S1 and Table 1; Jørgen E. Olesen: Overall content and structure of the MS, Fig. 1, complementary data from Danish field trials in Table S2; Maike Krauss: Content of the MS, complementary data from Swiss field trial in Table S2; Paweł Radzikowski: Content of the MS, mainly earthworms, main editing of references and reference list; Jordi Doltra: Content of the MS, mainly from Mediterranean perspective; Shahid Nadeem: Content of the MS, mainly N₂O, complementary data from Norwegian field trial in Table S2; Torfinn Torp: Statistical analyses on cause of N₂O fluxes based on Table S2; Valentini Pappa: Complementary data from field trials in Edinburgh and Aberdeen in Table S2; Christine Watson: Overall development and content of the MS.

Acknowledgments. The FertilCrop project was financially supported by transnational funding bodies of the FP7 ERA-Net project, CORE Organic Plus, and from the European Commission. The authors also acknowledge support from the Scottish Government Strategic Research Programme and from Møre og Romsdal county council. The text in this report is the sole responsibility of the authors and does not necessarily reflect the views of the national funding bodies having financed this project. The authors acknowledge Bruce Ball for complementary data from a field trial in Aberdeen used in table S2, and an anonymous referee for valuable and comprehensive comments.

References

- Aguilera, E., Lassaletta, L., Gattinger, A., and Gimeno, B. S.: Managing soil carbon for climate change mitigation and adaptation in Mediterranean cropping systems: A meta-analysis, *Agr. Ecosyst. Environ.*, 168, 25–36, doi:10.1016/j.agee.2013.02.003, 2013.
- Anderson, J. H.: The metabolism of hydroxylamine to nitrite by *Nitrosomonas*, *Biochem. J.*, 91, 8–17, 1964.
- Anderson, T.-H., and Domsch, K. H.: Ratios of microbial biomass carbon to total organic carbon in arable soils, *Soil Biol. Biochem.*, 21(4), 471–479, doi:10.1016/0038-0717(89)90117-X, 1989.
- Anderson, I. C., Poth, M., Homstead, J., and Burdige, D.: A comparison of NO and N₂O production by the autotrophic nitrifier *Nitrosomonas europaea* and the heterotrophic nitrifier *Alcaligenes faecalis*, *Appl. Environ. Microbiol.*, 59, 3525–3533, 1993.

Aronsson, H., Torstensson, G., and Bergstrom, L.: Leaching and crop uptake of N, P and K from organic and conventional cropping systems on a clay soil, *Soil Use Manage.*, 23(1), 71–81, 2007.

Askegaard, M., Olesen, J. E., and Kristensen, K.: Nitrate leaching from organic arable crop rotations: effects of location, manure and catch crop, *Soil Use Manage.*, 21(2), 181–188 [online] Available from: [doi:10.1016/j.agee.2005.08.003](https://doi.org/10.1016/j.agee.2005.08.003), 2005.

Askegaard, M., Olesen, J. E., Rasmussen, I. A., and Kristensen, K.: Nitrate leaching from organic arable crop rotations is mostly determined by autumn field management, *Agr. Ecosyst. Environ.*, 142(3–4), 149–160, [doi:10.1016/j.agee.2011.04.014](https://doi.org/10.1016/j.agee.2011.04.014), 2011.

Baggs, E. M., Rees, R. M., Smith, K. A., and Vinten, A. J. A.: Nitrous oxide emission from soils after incorporating crop residues, *Soil Use Manage.*, 16, 82–87, 2000a.

Baggs, E. M., Watson, C. A., and Rees, R. M.: The fate of nitrogen from incorporated cover crop and green manure residues, *Nutr. Cycl. Agroecosys.*, 56(2), 153–163 [online] Available from: [doi:10.1016/j.agee.2000.08.008](https://doi.org/10.1016/j.agee.2000.08.008), 2000b.

Bakken, L. R., Bergaust, L., Liu, B., and Frostegård, A.: Regulation of denitrification at the cellular level: a clue to the understanding of N₂O emissions from soils., *Philos. T. R. Soc. B.*, 367(1593), 1226–34, [doi:10.1098/rstb.2011.0321](https://doi.org/10.1098/rstb.2011.0321), 2012.

Balaine, N., Clough, T.J., Beare, M.H., Thomas, S.M., Meenken, E.D., Ross, J.G.: Changes in Relative Gas Diffusivity Explain Soil Nitrous Oxide Flux Dynamics. *Soil Sci. Soc. Am. J.*, 77, 1496. <https://doi.org/10.2136/sssaj2013.04.0141>, 2013.

Ball, B. C., Mctaggart, I. P., and Watson, C. A.: Influence of organic ley-arable management and afforestation in sandy loam to clay loam soils on fluxes of N₂O and CH₄ in Scotland, *Agr. Ecosyst. Environ.*, 90(3), 305–317, [online] Available from: [doi:10.1016/j.agee.2002.03.008](https://doi.org/10.1016/j.agee.2002.03.008), 2002.

Ball, B. C., Watson, C. A., and Crichton, I.: Nitrous oxide emissions, cereal growth, N recovery and soil nitrogen status after ploughing organically managed grass/clover swards, *Soil Use Manage.*, 23(2), 145–155, [doi:10.1016/j.agee.2006.07.007](https://doi.org/10.1016/j.agee.2006.07.007), 2007a.

Baral, K. R., Labouriau, R., Olesen, J. E., and Petersen, S. O.: Nitrous oxide emissions and nitrogen use efficiency of manure and digestates applied to spring barley, *Agr. Ecosyst. Environ.*, 239, 188–198, [doi:10.1016/j.agee.2017.01.012](https://doi.org/10.1016/j.agee.2017.01.012), 2017.

Barbieri, P., Pellerin, S., Nesme, T.: Comparing crop rotations between organic and conventional farming. *Sci. Rep-UK.*, 7, 1–10, <https://doi.org/10.1038/s41598-017-14271-6>, 2017.

Benoit, M., Garnier, J., Anglade, J., and Billen, G.: Nitrate leaching from organic and conventional arable crop farms in the Seine Basin (France), *Nutr. Cycl. Agroecosys.*, 100(3), 285–299, 2014.

Berntsen, J., Petersen, B. M., Olesen, J. E., Eriksen, J. E., and Soegaard, K.: Simulation of residual effects and nitrate leaching after incorporation of different ley types, *Eur. J. Agron.*, 23(3), 290–304, [online] Available from: [doi:10.1016/j.agee.2005.08.007](https://doi.org/10.1016/j.agee.2005.08.007), 2005.

Berntsen, J., Petersen, B. M., Sørensen, P., and Olesen, J. E.: Simulating Residual Effects of Animal Manures Using N-15 Isotopes, *Plant Soil*, 290, 173–87, 2007.

Berry, B. M., Sylvester-Bradley, R., Philipp, L., Hatch, D. J., Cuttle, S. P., Rayns, F. W., and Gosling, P.: Is the productivity of organic farms restricted by the supply of available nitrogen? *Soil Use Manage.*, 18(3), 248–255, <https://doi.org/10.1079/SUM2002129>, 2002.

Bhogal, A., Williams, J.R., Nicholson, F.A., Chadwick, D.R., Chambers, K.H., and Chambers, B.J.: Mineralization of organic nitrogen from farm manure applications, *Soil Use Manage.*, 32, 32-43, 2016.

Borgen, S. K., Lunde, H. W., Bakken, L. R., Bleken, M. A., and Breland, T. A.: Nitrogen dynamics in stockless organic clover-grass and cereal rotations, *Nutr. Cycl. Agroecosys.*, 92(3), 363–378 [online] Available from: [wos:000302576100009](https://doi.org/10.1007/s10533-012-9300-9), 2012.

Bouwman, A. F., Boumans, L. J. M., and Batjes, N. H.: Modeling global annual N₂O and NO emissions from fertilized fields, *Global Biogeochem Cy.*, 16(4), [online] Available from: [isi:000180874100028](https://doi.org/10.1029/2001GB001428), 2002.

Breland, T. A.: Measured and predicted mineralization of clover green manure at low temperatures at different depths in two soils, *Plant Soil*, 166(1), 13–20, 1994.

Bronick, C. J., and Lal, R.: Soil structure and management: A review, *Geoderma*, 124(1–2), 3–22, [doi:10.1016/j.geoderma.2004.03.005](https://doi.org/10.1016/j.geoderma.2004.03.005), 2005.

Brophy, L.S., Heichel, G. H., and Russelle, M. P.: Nitrogen transfer from forage legumes to grass in a systematic planting design, *Crop Sci.*, 27, 753–758, 1987.

Brozyna, M. A., Petersen, S. O., Chirinda, N., and Olesen, J. E.: Effects of grass-clover management and cover crops on nitrogen cycling and nitrous oxide emissions in a stockless organic crop rotation, *Agr. Ecosyst. Environ.*, 181, 115–126, [doi:10.1016/j.agee.2013.09.013](https://doi.org/10.1016/j.agee.2013.09.013), 2013.

Burke, I. C., Yonker, C. M., Parton, W. J., Cole, C. V., Flach, K., and Schimel, D. S.: Texture, Climate, and Cultivation Effects on Soil Organic Matter Content in U.S. Grassland Soils, *Soil Sci. Soc. Am. J.*, 53(i), 800–805, 1989.

Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S.: Nitrous oxide emissions from soils: how well do we understand the processes and their controls?, *Philos. T. R. Soc. B.*, 368(1621), 20130122, [doi:10.1098/rstb.2013.0122](https://doi.org/10.1098/rstb.2013.0122), 2013.

Campanelli, G., and Canali, S.: Crop production and environmental effects in conventional and organic vegetable farming systems: The case of a long-term experiment in Mediterranean conditions (Central Italy), *J. Sustain. Agr.*, 36(6), 599-619, 2012.

Carter, M. S. and Ambus, P.: Biologically fixed N₂ as a source for N₂O production in a grass-clover mixture, measured by N⁻¹⁵, *Nutr. Cycl. Agroecosys.*, 74 (1), 13–26, [wos:000235949200002](https://doi.org/10.1007/s10533-006-9000-2), 2006.

Chadwick, D., Sommer, S., Thorman, R., Figueiro, D., Cardenas, L., Amon, B., and Misselbrook, T.: Manure management: Implications for greenhouse gas emissions, *Anim. Feed Sci. Tech.*, 166–167, 514–531, 2011.

Chang, C., Cho, C. M., and Janzen, H. H.: Nitrous oxide emission from long-term manured soils, *Soil Sci. Soc. Am. J.*, 62(3), 677–682, 1998.

Chantigny, M.H., Angers, D.A., Rochette, P., Bélanger, G., Massé, D., Côté, D.: Gaseous Nitrogen Emissions and Forage Nitrogen Uptake on Soils Fertilized with Raw and Treated Swine Manure. *J. Environ. Qual.*, 36, 1864. <https://doi.org/10.2134/jeq2007.0083.2007>.

Chantigny, M. H., Rochette, P., Angers, D.A., Bittman, S., Buckley, K., Massé, D., Bélanger, G., Eriksen-Hamel, N., Gasser, M.-O.: Soil Nitrous Oxide Emissions Following Band-Incorporation of Fertilizer Nitrogen and Swine Manure. *J. Environ. Qual.*, 39, 1545–1553. <https://doi.org/10.2134/jeq2009.0482>, 2010.

Charles, A., Rochette, P., Whalen, J. K., Angers, D. A., Chantigny, M. H., and Bertrand, N.: Global nitrous oxide emission factors from agricultural soils after addition of organic amendments: A meta-analysis, *Agr. Ecosyst. Environ.*, 236(3), 88–98, [doi:10.1016/j.agee.2016.11.021](https://doi.org/10.1016/j.agee.2016.11.021), 2017.

Chen, H. H., Li, X. C., Hu, F., and Shi, W.: Soil nitrous oxide emissions following crop residue addition: a meta-analysis, *Glob. Chang. Biol.*, 19(10), 2956–2964, doi:10.1111/gcb.12274, 2013.

Chirinda, N., Carter, M. S., Albert, K. R., Ambus, P., Olesen, J. E., Porter, J. R., and Petersen, S. O.: Emissions of nitrous oxide from arable organic and conventional cropping systems on two soil types, *Agr. Ecosyst. Environ.*, 136(3–4), 199–208, [online] Available from: isi:000276275400003, 2010.

Congreves, K.A., Wagner-Riddle, C., Si, B.C., Clough, T.J.: Nitrous oxide emissions and biogeochemical responses to soil freezing-thawing and drying-wetting, *Soil Biol. Biochem.*, 117, 5–15, doi:10.1016/j.soilbio.2017.10.040, 2018.

Constantin, J., Mary, B., Laurent, F., Aubrion, G., Fontaine, A., Kerveillant, P., and Beaudoin, N.: Effects of catch crops, no till and reduced nitrogen fertilization on nitrogen leaching and balance in three long-term experiments, *Agr. Ecosyst. Environ.*, 135(4), 268–278, doi:10.1016/j.agee.2009.10.005, 2010.

Cooper, J., Baranski, M., Stewart, G., Nobel-de Lange, M., Bärberi, P., Fließbach, A., Peigné, J., Berner, A., Brock, C., Casagrande, M., Crowley, O., David, C., De Vliegher, A., Döring, T. F., Dupont, A., Entz, M., Grosse, M., Haase, T., Halde, C., Hammerl, V., Huiting, H., Leithold, G., Messmer, M., Schloter, M., Sukkel, W., van der Heijden, M. G. A., Willekens, K., Wittwer, R., and Mäder, P.: Shallow non-inversion tillage in organic farming maintains crop yields and increases soil C stocks: a meta-analysis, *Agron. Sustain. Dev.*, 36(1), doi:10.1007/s13593-016-0354-1, 2016.

Council of the European Union.: Council Regulation (EC) No 834/2007 of 28 June 2007 on organic production and labelling of organic products and repealing Regulation (EEC) No 2092/91, Official Journal of the European Union, L 189/1, 2007. <http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32007R0834&from=EN>

Crews, T. E., and Peoples, M. B.: Can the synchrony of nitrogen supply and crop demand be improved in legume and fertilizer-based agroecosystems? A review, *Nutr. Cycl. Agroecosys.*, 72(2), 101–120, doi:10.1007/s10705-004-6480-1, 2005.

Dalgaard, T., Hansen, B., Hasler, B., Hertel, O., Hutchings, N.J., Jacobsen, B., Kronvang, B., Olesen, J.E., Schjørring, J., Termansen, M., and Vejre, H.: Policies for agricultural nitrogen management - trends, challenges and prospects for improved efficiency in Denmark, *Environ. Res. Lett.*, 9, 115002, 2014.

De Notaris, C., Rasmussen, J., Sørensen, P., and Olesen, J. E.: Nitrogen leaching: A crop rotation perspective on the effect of N surplus, field management and use of catch crops, *Agr. Ecosyst. Environ.*, 255, 1–11, doi:10.1016/J.AGEE.2017.12.009, 2018.

De Ponti, T., Rijk, B., and Van Ittersum, M. K.: The crop yield gap between organic and conventional agriculture, *Agr. Syst.*, 108, 1–9. <https://doi.org/10.1016/j.agsy.2011.12.004>, 2012.

Di, H. J., and Cameron, K. C.: Nitrate leaching in temperate agroecosystems: sources, factors and mitigating strategies, *Nutr. Cycl. Agroecosys.*, 64(3), 237–256, [online] Available from: isi:000179746500004, 2002.

Dobbie, K. E., Smith, K. A.: Nitrous oxide emission factors for agricultural soils in Great Britain: the impact of soil water filled pore space and other controlling variables. *Glob. Chang. Biol.*, 99, 204–218, <https://doi.org/10.1046/j.1365-2486.2003.00563.x>, 2003.

Doltra, J., Gallejones, P., Olesen, J.E., Hansen, S., Frøseth, R.B., Krauss, M., Stalenga, J., Jończyk, K., Martínez-Fernández, A., Pacini, G.C.: Simulating soil fertility management effects on crop yield and soil nitrogen dynamics

in field trials under organic farming in Europe. *F. Crop. Res.* 233, 1–11. <https://doi.org/10.1016/j.fcr.2018.12.008>, 2019.

Dunbabin, V., Diggle, A., and Rengel, Z.: Is there an optimal root architecture for nitrate capture in leaching environments?, *Plant. Cell. Environ.*, 26(6), 835–844, 2003.

Dusenbury, M.P., Engel, R.E., Miller, P.R., Lemke, R.L., Wallander, R.: Nitrous Oxide Emissions from a Northern Great Plains Soil as Influenced by Nitrogen Management and Cropping Systems, *J. Environ. Qual.*, 37, 542, <https://doi.org/10.2134/jeq2006.0395>, 2008.

Eriksen, J., Askegaard, M., and Sjøgaard, K.: Residual effect and nitrate leaching in grass-arable rotations: effect of grassland proportion, sward type and fertilizer history, *Soil Use Manage.*, 24, 373–382, 2008.

Evers A. K., Demers T. A., Gordon A. M., and Thevathasan, N. V.: The effect of earthworm (*Lumbricus terrestris* L.) population density and soil water content interactions on nitrous oxide emissions from agricultural soils, *Appl. Environ. Soil Sci.*, doi:10.1155/2010/737096, 2010.

Fan, J., McConkey, B., Wang, H., and Janzen, H.: Root distribution by depth for temperate agricultural crops, *Field. Crop. Res.*, 189, 68–74, doi:10.1016/j.fcr.2016.02.013, 2016.

Filser J., Dette A., Fromm H., Lang A., Munch J.C., Winter K., and Beese F.: Reactions of soil organisms in site-specific management. The first long-term study at the landscape scale, *Ecosyst. Suppl.*, 28, 139–147, 1999.

Firestone, M., and Davidson, E.: Microbiological basis of NO and N₂O production and consumption in soil, in *Exchange of trace gases between terrestrial ecosystems and the atmosphere.*, (Andreac, M. O. and Schimel, D. S. editors) pp. 7–21, John Wiley & Sons Ltd., 1989.

Flessa, H., Dörsch, P., and Beese, F.: Seasonal variation of N₂O and CH₄ fluxes in differently managed arable soils in southern Germany, *J. Geophys. Res.*, 100(23), 115–124, 1995.

Frøseth, R. B.: Nitrogen dynamics in an organic green manure – cereal rotation and mineralization of clover leaves at low temperature, *Philosophiae Doctor (PhD) Thesis 2016:25*, Norwegian University of Life Sciences, Ås, pp 58, 2016.

Frøseth, R. B., Bakken, A. K., Bleken, M. A., Riley, H., Pommeresche, R., Thorup-Kristensen, K., and Hansen, S.: Effects of green manure herbage management and its digestate from biogas production on barley yield, N recovery, soil structure and earthworm populations, *Eur. J. Agron.*, 52, doi:10.1016/j.eja.2013.10.006, 2014.

Gattinger, A., Muller, A., Haeni, M., Skinner, C., Fliessbach, A., Buchmann, N., Mäder, P., Stolze, M., Smith, P., Scialabba, N. E.-H., and Niggli, U.: Enhanced top soil carbon stocks under organic farming., *P. Natl. Acad. Sci. USA*, 109(44), 18226–18231, doi:10.1073/pnas.1209429109, 2012.

Gomiero, T.: Alternative Land Management Strategies and Their Impact on Soil Conservation, *Agriculture*, 3(3), 464–483, doi:10.3390/agriculture3030464, 2013.

Gomiero, T., Pimentel, D., and Paoletti, M.: Environmental Impact of Different Agricultural Management Practices: Conventional vs. Organic Agriculture, *CRC. Crit. Rev. Plant Sci.*, 30(1–2), 95–124, doi:10.1080/07352689.2011.554355, 2011.

Hansen, E. M., Eriksen, J., and Vinther, F. P.: Catch crop strategy and nitrate leaching following grazed grass-clover. *Soil Use Manage.*, 23(4), pp.348–358., 2007.

Hansen, S., and Engelstad, F.: Earthworm populations in a cool and wet district as affected by tractor traffic and fertilisation, *Appl. Soil Ecol.*, 13(3), doi:10.1016/S0929-1393(99)00037-2, 1999.

Hansen, S., Bernard, M.-E., Rochette, P., Whalen, J. K., and Dörsch, P.: Nitrous oxide emissions from a fertile grassland in Western Norway following the application of inorganic and organic fertilizers, *Nutr. Cycl. Agroecosys.*, 98(1), 71–85, doi:10.1007/s10705-014-9597-x, 2014.

Hartmann, M., Frey, B., Mayer, J., Mäder, P., and Widmer, F.: Distinct soil microbial diversity under long-term organic and conventional farming, *ISME J.*, 9(5), 1177–1194, doi:10.1038/ismej.2014.210, 2015.

Holtan-Hartwig, L., Dorsch, P., and Bakken, L. R.: Low temperature control of soil denitrifying communities: kinetics of N₂O production and reduction, *Soil Biol. Biochem.*, 34(11), 1797–1806, [online] Available from: isi:000179996900029, 2002.

Hu, T., Sørensen, P., and Olesen, J. E.: Soil carbon varies between different organic and conventional management schemes in arable agriculture, *Eur. J. Agron.*, 94, 79-88, 2018a.

Hu, T., Olesen, J.E., Christensen, B.T. & Sørensen, P.: Release of C and N from fodder radish (*Raphanus sativus*) shoots and roots incubated in soils of different management history, *Acta Agr. Scand. B.-S. P.* (in press), 2018b.

IFOAM: The Principles of Organic Agriculture. https://www.ifoam.bio/sites/default/files/poa_english_web.pdf, downloaded April 2019.

IPCC: Guidelines for National Greenhouse Gas Inventories. Agriculture, Forestry and Other Land Use., International Panel of Climate Change, <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index>. [online] Available from: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html>, 2006.

Jensen, L.S., Salo, T., Palmason, F., Breland, T.A., Henriksen, T.M., Stenberg, B., Pedersen, A., Lundström, C., and Esala, M.: Influence of biochemical quality on C and N mineralisation from a broad variety of plant materials in soil, *Plant Soil* 273, 307-326, 2005.

Jeuffroy, M.H., Baranger, E., Carrouée, B., de Chezelles, E., Gosme, M., Hénault, C., Schneider, A., Cellier, P.: Nitrous oxide emissions from crop rotations including wheat, oilseed rape and dry peas, *Biogeosciences* 10, 1787–1797, <https://doi.org/10.5194/bg-10-1787-2013>, 2013.

Johnsson, H., Bergström, L., Jansson, P.-E., and Paustian, K.: Simulated nitrogen dynamics and losses in a layered agricultural soil, *Agr. Ecosyst. Environ.*, 18, 333-356, 1987.

Jończyk, K., and Martyniuk, S.: Productivity of crop rotations and selected indicators of soil fertility in different types of ecological farms, *J. Res. Appl. Agric. Eng.*, 62(3), 153-157, 2017.

Kayser, M., Müller, J., and Isselstein, J.: Nitrogen management in organic farming: Comparison of crop rotation residual effects on yields, N leaching and soil conditions, *Nutr. Cycl. Agroecosys.*, 87(1), 21–31, doi:10.1007/s10705-009-9309-0, 2010.

Kirchmann, H. and Bergström, L.: Do organic farming practices reduce nitrate leaching?, *Commun. Soil Sci. Plan.*, 32(7–8), 997–1028, [online] Available from: isi:000170101800005, 2001.

Krauss, M., Krause, H. M., Spangler, S., Kandeler, E., Behrens, S., Kappler, A., Mäder, P., and Gattinger, A.: Tillage system affects fertilizer-induced nitrous oxide emissions, *Biol. Fert. Soils*, 53(1), 49–59., doi.org/10.1007/s00374-016-1152-2, 2017a.

Krauss, M., Ruser, R., Müller, T., Hansen, S., Mäder, P., and Gattinger, A.: Impact of reduced tillage on greenhouse gas emissions and soil carbon stocks in an organic grass-clover ley - winter wheat cropping sequence, *Agr. Ecosyst. Environ.*, 239, 324–333, doi:10.1016/j.agee.2017.01.029, 2017b.

Kuiper, I., Deyn G. B., Thakur M. P., and Groenigen J. W.: Soil invertebrate fauna affect N₂O emissions from soil, *Glob. Chang. Biol.*, 19(9): 2814-2825, 2013.

Kuntz, M., Morely, N. J., Hallett, P. D., Watson, C., and Baggs, E. M.: Residue-C effects on dinitrification vary with soil depth, *Soil Biol. Biochem.*, 103, 365–375, 2016.

Köster, J. R., Cárdenas, L., Senbayram, M., Bol, R., Well, R., Butler, M., Mühling, K. H., and Dittert, K.: Rapid shift from denitrification to nitrification in soil after biogas residue application as indicated by nitrous oxide isotopomers, *Soil Biol. Biochem.*, 43(8), 1671–1677, doi:10.1016/j.soilbio.2011.04.004, 2011.

Känkänen, H., Kangas, A., Mela, T., Nikunen, U., Tuuri, H., and Vuorinen, M.: Timing incorporation of different green manure crops to minimize the risk of nitrogen leaching, *Agric. Food Sci. Finland.*, 7(5–6), 553–567, 1998.

Larsson, L., Ferm, M., Kasimir-Klemetsson, A., and Klemetsson, L.: Ammonia and nitrous oxide emissions from grass and alfalfa mulches, *Nutr. Cycl. Agroecosys.*, 51(1), 41–46, 1998.

Li, C. S., Frolking, S., and Butterbach-Bahl, K.: Carbon Sequestration in Arable Soils Is Likely to Increase Nitrous Oxide Emissions, Offsetting Reductions in Climate Radiative Forcing, *Climatic Change*, 72, 321–38, 2005.

Li, X., Petersen, S. O., Sørensen, P., and Olesen, J. E.: Effects of contrasting catch crops on nitrogen availability and nitrous oxide emissions in an organic cropping system, *Agr. Ecosyst. Environ.*, 199, 382–393, doi:10.1016/j.agee.2014.10.016, 2015a.

Li, X., Sørensen, P., Li, F., Petersen, S.O. & Olesen, J.E.: Quantifying biological nitrogen fixation of different catch crops, and the residual effects of roots and tops on nitrogen uptake in barley using in situ ¹⁵N labelling. *Plant Soil*, 395, 273–287, 2015b.

Li, X., Sørensen, P., Olesen, J. E., and Petersen, S. O.: Evidence for denitrification as main source of N₂O emission from residue-amended soil, *Soil Biol. Biochem.*, 92, 153–160, doi:10.1016/j.soilbio.2015.10.008, 2016.

Liu, B., Mørkved, P. T., Frostegård, Å., and Bakken, L. R.: Denitrification gene pools, transcription and kinetics of NO, N₂O and N₂ production as affected by soil pH, *FEMS Microbiol. Ecol.*, 72(3), 407–417, doi:10.1111/j.1574-6941.2010.00856.x, 2010.

Liu, S., Han, P., Hink, L., Prosser, J., Wagner M., and Brüggemann N.: Abiotic conversion of extracellular NH₂OH contributes to N₂O emission during ammonia oxidation. *Environ. Sci. Technol.*, 51, 13122–13132. DOI:10.1021/acs.est.7b02360, 2017.

Lorenz, K., and Lal, R.: Environmental Impact of Organic Agriculture, In *Adv. Agron.* 139, 99–152, Academic Press, 2016.

Lori, M., Symnaczik, S., Mäder, P., De Deyn, G., Gattinger, A., and Dentener, F.: Organic farming enhances soil microbial abundance and activity—A meta-analysis and meta-regression, *PLoS One*, 12(7), 1–25, doi:10.1371/journal.pone.0180442, 2017.

Lubbers, I. M., Brussaard, L., Otten, W., and Van Groenigen, J. W.: Earthworm induced N mineralization in fertilized grassland increases both N₂O emission and crop N uptake, *Eur. J. Soil Sci.*, 62(1), 152–161, 2011.

Lubbers I. M., Van Groenigen K. J., Fonte S. J., Six J., Brussaard L., and Van Groenigen J. W.: Greenhouse-gas emissions from soils increased by earthworms, *Nat. Clim. Chang.*, 3(3), 187–194, 2013.

Lynch, D. H.: Nutrient Cycling and Soil Health in Organic Cropping Systems - Importance of Management Strategies and Soil Resilience, *Sustain. Agric. Res.*, 4(3), 80, doi:10.5539/sar.v4n3p80, 2015.

Løes, A.-K., Bünemann, E. K., Cooper, J., Hörtenhuber, S., Magid, J., Oberson, A., Möller, K.: Nutrient supply to organic agriculture as governed by EU regulations and standards in six European countries, *Org. Agric.*, 7, 395–418. <https://doi.org/10.1007/s13165-016-0165-3>, 2017.

Marinari, S., Masciandaro, G., Ceccanti, B., and Grego, S.: Evolution of soil organic matter changes using pyrolysis and metabolic indices: A comparison between organic and mineral fertilization, *Bioresource Technol.*, 98(13), 2495–2502, doi:10.1016/j.biortech.2006.09.001, 2007.

Marriott, E. E., and Wander, M. M.: Total and Labile Soil Organic Matter in Organic and Conventional Farming Systems, *Soil Sci. Soc. Am. J.*, 70(3), 950, doi:10.2136/sssaj2005.0241, 2006.

Martyniuk, S., Koziel, M., Stalenga, J., and Jonczyk, K.: Labile fractions of soil organic matter and microbial characteristics of soil under organic and conventional crop management systems, *Biol. Agric. Hortic.*, 32, 1–6, doi:10.1080/01448765.2015.1015054, 2016.

McGee, J. A.: Does certified organic farming reduce greenhouse gas emissions from agricultural production?, *Agric. Hum. Values*, 32(2), 255–263, doi:10.1007/s10460-014-9543-1, 2015.

Meier, M. S., Stoessel, F., Jungbluth, N., Juraske, R., Schader, C., and Stolze, M.: Environmental impacts of organic and conventional agricultural products - Are the differences captured by life cycle assessment?, *J. Environ. Manage.*, 149, 193–208, doi:10.1016/j.jenvman.2014.10.006, 2015.

Mejjide, A., Díez, J. A., Sánchez-Martín, L., López-Fernández, S., and Vallejo, A.: Nitrogen oxide emissions from an irrigated maize crop amended with treated pig slurries and composts in a Mediterranean climate, *Agr. Ecosyst. Environ.*, 121(4), 383–394, doi:10.1016/j.agee.2006.11.020, 2007.

Melander, B., Rasmussen, I.A., and Olesen, J.E.: Incompatibility between fertility building methods and the management of perennial weeds in organic cropping systems, *Agr. Ecosyst. Environ.*, 220, 184–192, 2016.

Molodovskaya, M., Singurindy, O., Richards, B. K., Warland, J., Johnson, M. S., Steenhuis, T. S.: Temporal Variability of Nitrous Oxide from Fertilized Croplands: Hot Moment Analysis, *Soil Sci. Soc. Am. J.*, 76, 1728. <https://doi.org/10.2136/sssaj2012.0039>, 2012.

Mäder, P., Fliessbach, A., Dubois, D., Gunst, L., Fried, P., and Niggli, U.: Soil fertility and biodiversity in organic farming., *Science*, 296(5573), 1694–7, doi:10.1126/science.1071148, 2002.

Moyo, H., Davies, W. P., Cannon, N. D., and Conway, J. S.: Influences of two year red clover-grass ley management on nitrogen economy and following wheat performance, *Biol. Agric. Hort.*, 32:2, 98-109, DOI: 10.1080/01448765.2015.1057865, 2016.

Möller, K., and Stinner, W.: Effects of different manuring systems with and without biogas digestion on soil mineral nitrogen content and on gaseous nitrogen losses (ammonia, nitrous oxides), *Eur. J. Agron.*, 30(1), 1–16, doi:10.1016/j.eja.2008.06.003, 2009.

Nadeem, S., Hansen, S., Bleken, M. A., and Dörsch, P.: N₂O emission from organic barley cultivation as affected by green manure management, *Biogeosciences*, 9(7), doi:10.5194/bg-9-2747-2012, 2012.

Nebert L. D., Bloem J., Lubbers I. M., and van Groenigen J. W.: Association of earthworm-denitrifier interactions with increased emission of nitrous oxide from soil mesocosms amended with crop residue, *Appl. Environ. Microbiol.*, 77(12), 4097–4104, 2011.

Neumann, A., Torstensson, G., and Aronsson, H.: Losses of nitrogen and phosphorus via the drainage system from organic crop rotations with and without livestock on a clay soil in southwest Sweden, *Org. Agric.*, 1(4), 217–229, doi:10.1007/s13165-011-0017-0, 2011.

Olesen, J.E., Askegaard, M., and Rasmussen, I.A.: Design of an organic farming crop rotation experiment. *Acta Agriculturae Scandinavica, Section B, Soil and Plant Science* 50, 13-21, 2000.

Kommentert [SH47]: 23

Pandey, A., Li, F., Askegaard, M., and Olesen, J. E.: Biological nitrogen fixation in organic and conventional arable crop rotations in Denmark, *Eur. J. Agron.*, 90, 87-95, 2017.

Pandey, A., Li, F., Askegaard, M., Rasmussen, I.A., and Olesen, J.E.: Nitrogen balances in organic and conventional arable crop rotations and their relations to nitrogen yield and leaching losses, *Agr. Ecosyst. Environ.*, 265, 350-362, 2018.

Pappa, V. A., Rees, R. M., Walker, W. A., and Baddely, J. A.: Intercropping can reduce environmental impacts, *Agric. Environ. VII L. Manag. a Chang. Environ. Proc. SAC SEPA Bienn. Conf.*, 287-290, 2008.

Pappa, V. A., Rees, R. M., Walker, R. L., Baddeley, J. A., and Watson, C. A.: Nitrous oxide emissions and nitrate leaching in an arable rotation resulting from the presence of an intercrop, *Agr. Ecosyst. Environ.*, 141(1-2), 153-161, doi:10.1016/j.agee.2011.02.025, 2011.

Pelster, D.E., Chantigny, M.H., Rochette, P., Angers, D.A., Rieux, C., Vanasse, A.: Nitrous Oxide Emissions Respond Differently to Mineral and Organic Nitrogen Sources in Contrasting Soil Types. *J. Environ. Qual.*, 41, 427, <https://doi.org/10.2134/jeq2011.0261>, 2012

Petersen, S.O., Schjonning, P., Thomsen, I.K., Christensen, B.T.: Nitrous oxide evolution from structurally intact soil as influenced by tillage and soil water content, *Soil Biol. Biochem.*, 40, 967-977, <https://doi.org/10.1016/j.soilbio.2007.11.017>, 2008.

Petersen, S. O., Mutegi, J. K., Hansen, E. M., and Munkholm, L. J.: Tillage effects on N₂O emissions as influenced by a winter cover crop, *Soil Biol. Biochem.*, 43(7), 1509-1517, doi:10.1016/j.soilbio.2011.03.028, 2011.

Petersen, S.O., Schjonning, P., Olesen, J.E., Christensen, S., and Christensen, B.T.: Sources of Nitrogen for Winter Wheat in Organic Cropping Systems, *Soil Sci. Soc. Am. J.*, 77(1), 155-165, 2013.

Peyraud, C., Mary, B., Perrin, P., Vericel, G., Grehan, E., Justes, E., and Leonard, J.: N₂O emissions of low input cropping systems as affected by legume and cover crops use, *Agr. Ecosyst. Environ.*, 224, 145-156, doi:10.1016/j.agee.2016.03.028, 2016.

Pfiffner L., Mäder P.: Effects of biodynamic, organic and conventional production systems on earthworm populations, *Biol. Agric. Hortic.*, 15:1-4, 2-10, DOI: 10.1080/01448765.1997.9755177, 1997.

Pfiffner, L., Luka, H.: Earthworm Populations in Two Low-Input Cereal Farming Systems, *Appl. Soil Ecol.*, 37, 184-91, doi:10.1016/j.apsoil.2007.06.005, 2007.

Plaza-Bonilla, D., Nolot, J.M., Raffailac, D., and Justes, E.: Cover crops mitigate nitrate leaching in cropping systems including grain legumes: field evidence and model simulations, *Agr. Ecosyst. Environ.*, 212,1-12, 2015.

Postma-Blaauw, M. B., Bloem, J., Faber, J. H., Van Groenigen, J. W., De Goede, R. G., and Brussaard, L.: Earthworm species composition affects the soil bacterial community and net nitrogen mineralization, *Pedobiologia*, 50(3), 243-256, 2006.

Poudel D. D., Horwath W. R., Lanini W. T., Temple S. R., Van Bruggen A. H. C.: Comparison of soil N availability and leaching potential, crop yields and weeds in organic, low-input and conventional farming systems in northern California, *Agr. Ecosyst. Environ.*, 90,125-137, 2002.

Prieto, R. A.: The role of earthworms in nitrous oxide emissions from forage agroecosystems, In *Masters Abstracts International*, (Vol. 50, No. 02), 2011.

Pugesgaard, S., Petersen, S. O., Chirinda, N., and Olesen, J. E.: Crop residues as driver for N₂O emissions from a sandy loam soil, *Agric. Forest. Meteorol.*, 233, 45-54, doi:10.1016/j.agrformet.2016.11.007, 2017.

Rasse, D., Ritchie, J., Peterson, W., Wei, J., and Smucker, A.: Rye cover crop and nitrogen fertilization effects on nitrate leaching in inbred maize fields, *J. Environ. Qual.*, 29, 298–304, doi:10.2134/jeq2000.00472425002900010037x, 2000.

Ravishankara, A. R., Daniel, J. S., and Portmann, R. W.: Nitrous Oxide (N₂O): The Dominant Ozone-Depleting Substance Emitted in the 21st Century, *Science* 326, 123–125, doi:10.1126/science.1176985, 2009.

Riley, H., Pommeresche, R., Eltun, R., Hansen, S., and Korsae, A.: Soil structure, organic matter and earthworm activity in a comparison of cropping systems with contrasting tillage, rotations, fertilizer levels and manure use, *Agr. Ecosyst. Environ.*, 124(3–4), 275–284, [online] Available from: isi:000254944900016, 2008.

Robertson, G. P., and Vitousek, P. M.: Nitrogen in Agriculture: Balancing the Cost of an Essential Resource, *Annu. Rev. Envi. Resour.*, 34(1), 97–125, doi:10.1146/annurev.enviro.032108.105046, 2009.

Rochette, P., and Janzen, H. H.: Towards a revised coefficient for estimating N₂O emissions from legumes, *Nutr. Cycl. Agroecosys.*, 73(2–3), 171–179, [online] Available from: wos:000233363200006, 2005.

Röös, E., Mie, A., Wivstad, M., Salomon, E., Johansson, B., Gunnarsson, S., Wallenbeck, A., Hoffmann, R., Nilsson, U., Sundberg, S., and Watson, C. A.: Risks and opportunities of increasing yields in organic farming. A review, *Agron. Sustain. Dev.*, 38, 14, <https://doi.org/10.1007/s13593-018-0489-3>, 2018.

Sahrawat, K. L., and Keeney, D. R.: Nitrous Oxide Emissions from Soils, *Adv. Soil Sci.*, 4, 103–47, 1986.

Shcherbak, I., Millar, N., and Robertson, G. P.: Global metaanalysis of the nonlinear response of soil nitrous oxide (N₂O) emissions to fertilizer nitrogen, *P. Natl. Acad. Sci. USA*, 111(25), 9199–9204, doi:10.1073/pnas.1322434111, 2014.

Shelton, R. E., Jacobsen, K. L., and McCulley, R. L.: Cover Crops and Fertilization Alter Nitrogen Loss in Organic and Conventional Conservation Agriculture Systems, *Front. Plant Sci.*, 8, 1–14, doi:10.3389/fpls.2017.02260, 2018.

Skinner, C., Gattinger, A., Müller, A., Mäder, P., Fließbach, A., Stolze, M., Ruser, R., and Niggli, U.: Greenhouse gas fluxes from agricultural soils under organic and non-organic management — A global meta-analysis, *Sci. Total Environ.*, 468–469, 553–563, doi:http://dx.doi.org/10.1016/j.scitotenv.2013.08.098, 2014.

Skinner, C., Gattinger, A., Krauss, M., Krause, H.-M., Mayer, J., van der Heijden, M. G. A., Mäder, P.: The impact of long-term organic farming on soil-derived greenhouse gas emissions, *Sci. Rep.* 9, 1702. <https://doi.org/10.1038/s41598-018-38207-w>, 2019.

Smith, K. A., Dobbie, K. E., Thorman, R., Watson, C. J., Chadwick, D. R., Yamulki, S., Ball, B. C.: The effect of N fertilizer forms on nitrous oxide emissions from UK arable land and grassland, *Nutr. Cycl. Agroecosys.*, 93, 127–149, 2012.

Spargo, J. T., Cavigelli, M. A., Mirsky, S. B., Maul, J. E., and Meisinger, J. J.: Mineralizable soil nitrogen and labile soil organic matter in diverse long-term cropping systems, *Nutr. Cycl. Agroecosys.*, 90, 253–266, 2011.

Stalenga, J., and Jończyk, K.: Nutrient management and soil organic matter balance in the organic crop production system evaluated by NDICEA model, *J. Res. Appl. Agric. Eng.*, 53(4), 78–84, 2008.

Stehfest, E., and Bouwman, L.: N₂O and NO emission from agricultural fields and soils under natural vegetation: summarizing available measurement data and modeling of global annual emissions, *Nutr. Cycl. Agroecosys.*, 74(3), 207–228, [online] Available from: isi:000240619700001, 2006.

Stenberg, M., Aronsson, H., Linden, B., Rydberg, T., and Gustafson, A.: Soil mineral nitrogen and nitrate leaching losses in soil tillage systems combined with a catch crop, *Soil Till. Res.*, 50(2), 115–125, doi:10.1016/S0167-1987(98)00197-4, 1999.

Stenberg, M., Ulén, B., Söderström, M., Roland, B., Delin, K., and Helander, C. A.: Tile drain losses of nitrogen and phosphorus from fields under integrated and organic crop rotations. A four-year study on a clay soil in southwest Sweden, *Sci. Total Environ.*, 434, 79–89, doi:10.1016/j.scitotenv.2011.12.039, 2012.

Stockdale, E. A., Lampkin, N. H., Hovi, M., Keatinge, R., Lennartsson, E. K. M., MacDonald, D. W., Padel, S., Tattersall, F. H., Wolfe, M. S., and Watson, C. A.: Agronomic and Environmental Implications of Organic Farming Systems, *Adv. Agron.* 70, 261-327, 2001.

Stopes, C., Lord, E. I., Philipps, L., and Woodward, L.: Nitrate leaching from organic farms and conventional farms following best practice, *Soil Use Manage.*, 18, 256–263, [online] Available from: isi:000178455700004, 2002.

Sturite, I., Rivedal, S., and Dörsch, P.: Nitrous oxide emissions from clover rich leys during the long Northern winter., in Cordovil C.M.d.S.(Ed.), *Proceedings of the 18th Nitrogen Workshop – The nitrogen challenge: building a blueprint for nitrogen use efficiency and food security*, 30th June – 3rd July 2014, Lisboa, Portugal, pp. 517–518., 2014.

Thangarajan, R., Bolan, N. S., Tian, G. L., Naidu, R., and Kunhikrishnan, A.: Role of organic amendment application on greenhouse gas emission from soil, *Sci. Total Environ.*, 465, 72–96, doi:10.1016/j.scitotenv.2013.01.031, 2013.

Thorup-Kristensen, K.: Effect of deep and shallow root systems on the dynamics of soil inorganic N during 3-year crop rotations, *Plant Soil*, 288(1–2), 233–248, [online] Available from: isi:000243001400018, 2006.

Thorup-Kristensen, K., and Dresbøll, D. B.: Incorporation time of nitrogen catch crops influences the N effect for the succeeding crop, *Soil Use Manage.*, 26(1), 27–35, [online] Available from: wos:000274666500005, 2010.

Tonitto, C., David, M. B., and Drinkwater, L.: Replacing bare fallows with cover crops in fertilizer-intensive cropping systems: A meta-analysis of crop yield and N dynamics, *Agr. Ecosyst. Environ.*, 112, 58–72, doi:10.1007/s10705-010-9354-8, 2006.

Torstensson, G., and Aronsson, H.: Nitrogen leaching and crop availability in manured catch crop systems in Sweden, *Nutr. Cycl. Agroecosys.*, 56(2), 139–152, doi:10.1023/A:1009821519042, 2000.

Torstensson, G., Aronsson, H., and Bergstrom, L.: Nutrient use efficiencies and leaching of organic and conventional cropping systems in Sweden, *Agron. J.*, 98(3), 603–615 [online] Available from: isi:000237867600023, 2006.

Tosti, G., Benincasa, P., Farneselli, M., Tei, F., and Guiducci, M.: Barley-hairy vetch mixture as cover crop for green manuring and the mitigation of N leaching risk, *Eur. J. Agron.*, 54, 34–39, doi:10.1016/J.EJA.2013.11.012, 2014.

Tosti, G., Farneselli, M., Benincasa, P., and Guiducci, M.: Nitrogen fertilization strategies for organic wheat production: Crop yield and nitrate leaching, *Agron. J.*, 108(2), 770–781, doi:10.2134/agronj2015.0464, 2016.

Tuomisto, H. L., Hodge, I. D., Riordan, P., and Macdonald, D. W.: Does organic farming reduce environmental impacts? - A meta-analysis of European research, *J. Environ. Manage.*, 112(834), 309–320, doi:10.1016/j.jenvman.2012.08.018, 2012.

Tyson, K. C., Scholefield, D., Jarvis, S. C., and Stone, A. C.: A comparison of animal output and nitrogen leaching losses recorded from drained fertilized grass and grass/clover pasture, *J. Agric. Sci.*, 129(3), 315-323, 1997.

Ussiri, D., and Lal, R.: Global Sources of Nitrous Oxide, in *Soil Emission of Nitrous Oxide and its Mitigation*, pp. 131–175, Springer Netherlands, Dordrecht., 2013.

Valkama, E., Lemola, R., Känkänen, H., and Turtola, E.: Meta-analysis of the effects of undersown catch crops on nitrogen leaching loss and grain yields in the Nordic countries, *Agr. Ecosyst. Environ.*, 203, 93–101, doi:10.1016/j.agee.2015.01.023, 2015.

Van Groenigen, J. W., Velthof, G. L., Oenema, O., Van Groenigen, K. J., and Van Kessel, C.: Towards an agronomic assessment of N₂O emissions: A case study for arable crops, *Eur. J. Soil Sci.*, 61(6), 903–913, doi:10.1111/j.1365-2389.2009.01217.x, 2010.

Wagner-Riddle, C., Congreves, K.A., Abalos, D., Berg, A.A., Brown, S.E., Ambadan, J.T., Gao, X., Tenuta, M.: Globally important nitrous oxide emissions from croplands induced by freeze-thaw cycles, *Nat. Geosci.* 10, 279–283. <https://doi.org/10.1038/ngeo2907>, 2017.

Watson, C.A., Fowler, S.M., and Wilman, D.: Soil inorganic-N and nitrate leaching on organic farms, *J. Agric. Sci.*, 120(3), 361-369, 1993.

Watson, C.A., Atkinson, D., Gosling, P., Jackson, L.R., and Rayns, F.W.: Managing soil fertility in organic farming systems, *Soil Use Manage.*, 18, 239-247, 2002.

Weier, K.L., Doran, W.L., Power, J.F., and Walters, D.T: Denitrification and the dinitrogen/nitrous oxide ratio as affected by soil water, available carbon, and nitrate. *Soil Sci. Soc. Am. J.* 57, 66-72, 1992.

Westphal, M., Tenuta, M., and Entz, M. H.: Nitrous oxide emissions with organic crop production depends on fall soil moisture, *Agr. Ecosyst. Environ.*, 254, 41–49, doi:10.1016/j.agee.2017.11.005, 2018.

Willer, H., Schaack, D., and Lernoud, J.: Organic farming and market development in Europe and the European Union. In: Willer, H., and Lernoud, J. (Eds.): *The World of Organic Agriculture. Statistics and Emerging Trends 2018*. Research Institute of Organic Agriculture (FiBL), Frick, and IFOAM – Organics International, Bonn, p 218 – 255, 2018.

Willer, H. and Lernoud, J. (Eds.): *The World of Organic Agriculture. Statistics and Emerging Trends 2018*. Research Institute of Organic Agriculture (FiBL), Frick, and IFOAM – Organics International, Bonn, 2018.

Xia, L., Lam, S.K., Wolf, B., Kiese, R., Chen, D., Butterbach-Bahl, K.: Trade-offs between soil carbon sequestration and reactive nitrogen losses under straw return in global agroecosystems, *Glob. Chang. Biol.*, 24, 5919–5932. <https://doi.org/10.1111/gcb.14466>, 2018.

Captions supplementary material

S1. Soil mineral nitrogen (SMN) contents in soil profiles in organic field trials in Norway and Poland as influenced by crop rotation, soil tillage and N fertilization.

S2. N₂O emission and the five highest daily N₂O flux rates in the given measurement periods for organic field trials in Switzerland, Denmark, Scotland and Norway. WFPS, soil temperature and soil mineral-N at 0-20 cm depth are given for the day with highest flux rate. Abbreviations: CS = Cattle Slurry, CCM = composted cattle

manure, PS = Pig slurry, P=ploughing, H = Harrowing, CCinc= Cover crop incorporated, CCs= Cover Crop under sown, CCh = cover crop harvested

S3. Annual N leaching (total N or nitrate N) reported from organic field trials in Europe, as influenced by crops, soil type and N applied as organic fertilizers.

S4. Statistical analyses Eq. (1).

S5. Statistical analyses Eq. (2).