

Multi-year effect of wetting on CH₄ flux at taiga-tundra boundary in northeastern Siberia deduced from stable isotope ratios of CH₄

Ryo Shingubara^{1,a}, Atsuko Sugimoto^{2,3,4}, Jun Murase⁵, Go Iwahana^{2,6}, Shunsuke Tei^{2,3}, Maochang Liang^{1,b}, Shinya Takano¹, Tomoki Morozumi¹, Trofim C. Maximov^{7,8}

5

¹Graduate School of Environmental Science, Hokkaido University, Sapporo, 060-0810, Japan

²Arctic Research Center, Hokkaido University, Sapporo, 001-0021, Japan.

³Faculty of Environmental Earth Science, Hokkaido University, Sapporo, 060-0810, Japan.

10 ⁴Global Station for Arctic Research, Global Institution for Collaborative Research and Education, Hokkaido University, Sapporo, 060-0808, Japan.

⁵Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya, 464-8601, Japan.

⁶International Arctic Research Center, University of Alaska Fairbanks, Fairbanks, 99775-7340, USA.

⁷Institute for Biological Problems of Cryolithozone, Siberian Branch, Russian Academy of Sciences, Yakutsk, 677890, Russia.

15 ⁸Institute of Natural Sciences, North-Eastern Federal University, Yakutsk, 677000, Russia.

^aCurrently at: Graduate School of Environmental Studies, Nagoya University, Nagoya, 464-8601, Japan.

^bCurrently at: College of Horticulture and Gardening, Yangtze University, Jingzhou, 434023, China.

Correspondence to: Atsuko Sugimoto (atsukos@ees.hokudai.ac.jp)

Abstract. The response of CH₄ emission from natural wetlands due to meteorological conditions is important because of its strong greenhouse effect. To understand the relationship between CH₄ flux and wetting, we observed interannual variations in chamber CH₄ flux, as well as the concentration, δ¹³C, and δD of dissolved CH₄ during the summer from 2009 to 2013 at the taiga-tundra boundary in the vicinity of Chokurdakh (70° 37' N, 147° 55' E), located on the lowlands of the Indigirka River in northeastern Siberia. We also conducted soil incubation experiments to interpret δ¹³C and δD of dissolved CH₄ and to investigate variations in CH₄ production and oxidation processes. Methane flux showed large interannual variations in wet areas of sphagnum mosses and sedges (36–140 mg CH₄ m⁻² day⁻¹ emitted). Increased CH₄ emission was recorded in the summer of 2011 when a wetting event with extreme precipitation occurred. Although water level decreased from 2011 to 2013, CH₄ emission remained relatively high in 2012, and increased further in 2013. Thaw depth became deeper from 2011 to 2013, which may partly explain the increase in CH₄ emission. Moreover, dissolved CH₄ concentration rose sharply by one order of magnitude from 2011 to 2012, and increased further from 2012 to 2013. Large variations in δ¹³C and δD of dissolved CH₄ were observed in 2011, and smaller variations were seen in 2012 and 2013, suggesting both enhancement of CH₄ production and less significance of CH₄ oxidation relative to the larger pool of dissolved CH₄. These multi-year effects of wetting on CH₄ dynamics may have been caused by continued soil reduction across multiple years following the wetting. Delayed activation of acetoclastic methanogenesis following soil reduction could also have contributed to the enhancement of CH₄ production. These processes suggest that duration of water saturation in the active layer can be important for predicting CH₄ emission following a wetting event in permafrost ecosystem.

1 Introduction

Atmospheric CH₄ has an important greenhouse effect (Myhre et al., 2013). The largest source of atmospheric CH₄ is the emission from natural wetlands, which is considered to be the main driver of interannual variations in the global CH₄ emission, depending on meteorological conditions such as air temperature and precipitation (Ciais et al., 2013). For instance, Dlugokencky et al. (2009) reported that high temperatures in the Arctic and high precipitation in the tropics led to high CH₄ emissions from natural wetlands, which caused the observed large growth rates in atmospheric CH₄ concentration during 2007 and 2008. Atmospheric CH₄ has been increasing from 2007 through the present (Nisbet et al., 2014).

Methane flux from wetland soil to the atmosphere (we define a positive flux value as CH₄ emission) is determined by three processes: CH₄ production, oxidation, and transport (Lai, 2009). Methane is produced by strictly anaerobic *Archaea* (methanogens) mainly via hydrogenotrophic methanogenesis ($4\text{H}_2 + \text{CO}_2 \rightarrow \text{CH}_4 + 2\text{H}_2\text{O}$) or acetoclastic methanogenesis ($\text{CH}_3\text{COOH} \rightarrow \text{CH}_4 + \text{CO}_2$) as an end product of organic matter decomposition (Lai, 2009). In the soil's aerobic zone, CH₄ is oxidized to CO₂ by methanotrophic bacteria utilizing O₂, which reduces CH₄ emission to the atmosphere (Lai, 2009). Underground CH₄ is transported to the atmosphere via bubble ebullition, diffusion through soil layers and surface water, and via aerenchyma of vascular plants (Lai, 2009).

High water levels can lead to development of reducing conditions in soil, which can promote CH₄ production or depress CH₄ oxidation, both leading to increases in CH₄ flux (Lai, 2009). This is reflected in the widely observed positive relationship between water level and CH₄ flux, found in a meta-analysis across the circum-Arctic permafrost zone (Olefeldt et al., 2013). Meanwhile, Desyatkin et al. (2014) observed increases in CH₄ flux during the second consecutive year of flooding at a thermokarst depression in boreal eastern Siberia. Treat et al. (2007) reported observations at a temperate fen in the northeastern USA showing that high water level coincided with high CH₄ flux in interannual variations. However, water level correlated negatively with CH₄ flux over shorter timescales, namely as monthly means or individual measurements. These observational results imply that wetting is not directly related to CH₄ flux in wetlands. To understand the relationship between wetting and CH₄ flux, it is necessary to assess the underlying processes.

Stable isotopes of CH₄ have been used to estimate production pathways of CH₄ (Sugimoto and Wada, 1993; Sugimoto and Wada, 1995; McCalley et al., 2014; Itoh et al., 2015), determine the fraction of oxidized CH₄ versus produced CH₄ (Marik et al., 2002; Preuss et al., 2013) and to study mechanisms of CH₄ transport by plants (Chanton, 2005). When CH₄ in soil is lost by oxidation or diffusion, both $\delta^{13}\text{C}$ and δD of the remaining CH₄ increase. While the hydrogen isotope ratio increases more than that of carbon during oxidation, both ratios are considered to change to the same extent during diffusion. Thus it is useful to analyze both carbon and hydrogen isotopes of CH₄ to distinguish the effects of both of these processes (Chanton, 2005).

The taiga-tundra boundary ecosystem (or transition zone) contains vegetation types of both taiga and tundra ecosystems. Liang et al. (2014) reported that the distribution of vegetation types at the taiga-tundra boundary on the lowland of the Indigirka River in northeastern Siberia is controlled by soil moisture, which corresponds to microtopography. Larches, the dominant tree species in the taiga forests of eastern Siberia, grow on micro-reliefs with higher ground level and drier soil, while wetland vegetation such as sphagnum mosses and sedges, typically seen in wet tundra (van Huissteden et al., 2005; van der Molen et al., 2007), dominates lower and wetter micro-reliefs. Thus, it is reasonable to assume that the taiga-tundra boundary ecosystem has various micro-reliefs in terms of interannual variation in soil wetness conditions: always wet micro-reliefs, always dry micro-reliefs, and micro-reliefs with large interannual wetness variations. Hence, this ecosystem is a suitable area to evaluate the processes controlling CH₄ flux in relation to soil wetting and/or drying on an interannual timescale.

In this study, to understand relationships between CH₄ flux and environmental factors, we observed interannual variations in chamber CH₄ flux, along with the concentration, $\delta^{13}\text{C}$, and δD of dissolved CH₄ during the summer, from 2009 to 2013, at the taiga-tundra boundary located on Indigirka River lowlands in northeastern Siberia. We also conducted soil incubation experiments to investigate how δ values of CH₄ reflect CH₄ production and oxidation processes in this ecosystem. In 2011, a wetting event with a significant amount of precipitation occurred. We focused in particular on the responses of CH₄ flux and other underlying processes to this unusual wetting event.

2 Methods

2.1 Study sites

The taiga-tundra boundary on the lowlands of the Indigirka River was selected as our study area. Observations and sampling were conducted at three sites (V: Verkhny Khatistakha, K: Kodac, and B: Boydom) in the vicinity of Chokurdakh (70° 37' N, 147° 55' E), Republic of Sakha (Yakutia), Russia (Fig. 1 and Table 1). The sites are located in the Russian Arctic with an annual mean air temperature of -13.9 °C and an annual mean precipitation of 208 mm for the period of 1950–2008, according to the Baseline Meteorological Data in Siberia Database (Yabuki et al., 2011). Sites V, K, and B are alongside the Indigirka River or its tributary, and tree density decreases from site V to site B.

These study sites are underlain by continuous permafrost (Iwahana et al., 2014). Normally, snowmelt and the start of active layer thawing begin in the latter half of May through the first half of June, and the growing season occurs from the end of June through the beginning of August. Air temperature and surface soil temperature (10 cm depth) peak in July, whereas the maximum thaw depth occurs from the latter half of August to the first half of September. The freezing of the active layer starts in the latter half of September to October and the whole active layer freezes from November to December.

Observations of CH₄ flux were conducted at seven points with three typical vegetation types, as summarized in Table 1. These vegetation types are distributed in patches, corresponding to microtopography and soil moisture (Liang et al., 2014). Micro-relief with a higher ground level is covered by green moss, larch trees, and shrubs of willows or dwarf birches. On the other hand, lower micro-relief is covered by wetland vegetation of sphagnum moss or sedges. In this study, the former vegetation type was termed 'tree mound', and the latter type was termed 'wet area'. Observation points in tree mounds were selected at each of the sites V, K, and B, and termed 'tree mound_V', 'tree mound_K', 'tree mound_B' (Table 1). For observation points of wet areas, a micro-relief covered by sphagnum moss in site K was termed 'sphagnum_K', and points covered by sedges including especially cotton-sedges (*Eriophorum angustifolium*) in sites V, K, and B were termed 'sedge_V', 'sedge_K', and 'sedge_B', respectively. Measurements of volumetric water content in the surface soil layer (0–20 cm) by TDR (time domain reflectometry; TDR-341F, Fujiwara Scientific Company, Japan) showed that tree mounds were drier than wet areas; this will be described in Sect. 3.1 (Table 1).

2.2 Field observations and samplings

Methane flux was observed using the chamber method in each of the typical vegetation types described in Sect. 2.1 during the summer from 2009 to 2013. A transparent cylindrical flux chamber (acrylic resin, base area 4.7×10^2 cm², height 25 cm) was installed on the ground. The headspace gas of the chamber (ca. 12 L) was circulated with a pump (ca. 1 L min⁻¹). The chamber was closed for 15–30 min and headspace gas was sampled two to three times after chamber closure. In most cases, the chamber was closed for 30 min and headspace gas was collected at 0 min, 15 min, and 30 min after closure. Samples were kept in pre-evacuated glass vials with butyl rubber septa. To minimize soil disturbance, we stepped on wooden boards at observation points. In 2009 and 2010, CH₄ flux measurements were conducted in the latter half of July, and from 2011 to

2013, observations were conducted continuously from early July to the end of July or early August. For all of these years, the observation period included the warmest season when CH₄ emission was expected to be the most active (Table S2).

For measurements of dissolved CH₄, surface water and soil pore water were sampled in wet areas from 2011 to 2013. Surface water was directly taken up by a 50 mL plastic syringe with a three-way cock attached to its tip, whereas soil pore water was sampled by a 50 mL syringe (with a three-way cock attached) through a plastic tube inserted in the soil. Soon after collecting water samples, dissolved CH₄ was extracted inside the syringes by the headspace method, after adding 15–35 mL of the atmosphere prepared in a 10 L aluminum bag. This atmosphere was collected beforehand at Chokurdakh village or our observation sites, and filtered by Molecular Sieves 5A (1/16 pellets, FUJIFILM Wako Pure Chemical Corporation, Japan). The atmosphere was analyzed later for CH₄ concentration and isotopic compositions as a background sample (2.0–4.3 ppm for CH₄ concentration, –53‰ to –45‰ for δ¹³C of CH₄, and –168‰ to –78‰ for δD of CH₄). The syringes were vigorously shaken for one minute and left standing for five minutes to ensure equilibration. Finally, headspace gas in the syringes was preserved in 10–20 mL pre-evacuated glass vials with rubber septa.

Concurrently with each flux measurement, soil temperature around the flux chamber was measured with a temperature sensor in an ORP electrode (PST-2739C, DKK-TOA Corporation, Japan) with an ORP meter (RM-30P or RM-20P). After flux measurement samples were collected, thaw depth was observed on the same day around each chamber by inserting a steel rod into the ground. From 2011 on, water level was also measured after flux measurements around each chamber in wet areas using a scale. The water level was expressed as height relative to the ground surface or the moss surface. Observation dates of these environmental factors are shown in Table S2.

2.3 Soil incubation experiments and microbial community analysis

Soil incubation experiments were conducted to estimate δ¹³C and δD of produced CH₄ and fractionation factors of CH₄ oxidation for carbon and hydrogen isotopes. For CH₄ production experiments, surface soil was sampled in all the wet areas in Table 1 (sedge_V, sphagnum_K, sedge_K, and sedge_B) during summer 2013. Samples were taken at 10 cm depth at each sampling location. To observe vertical variations in δ values of produced CH₄ within the thaw layer, we also collected samples from two additional depths (20 cm and 30 cm) at sedge_K, which is a location typical of the taiga-tundra boundary region. These samples were from organic layers, except for the samples from 30 cm, which were from the top of the mineral layer.

Approximately 10 mL of soil was directly transferred into each plastic syringe (60 mL maximum capacity) along with in situ water (approximately 50 mL) to prevent the sample from being oxidized by the atmosphere. Syringes were preserved in water to ensure no leakage and were immediately pre-incubated for 4–8 days, then incubated in triplicate for 8 days. Pre-incubation and incubation temperatures were set at 5 °C. We also incubated syringes at 10 °C for samples from 10 cm depth at sedge_K to investigate temperature dependence of δ values of produced CH₄. For each of these seven incubation treatments (sphagnum_K, sedge_V, sedge_K, and sphagnum_K, 10 cm depth, 5 °C; sedge_K, 20 cm and 30 cm depths, 5 °C; sedge_K, 10 cm depth, 10 °C), three replicate soil samples were prepared. Water in each incubation syringe was

sampled twice at the start and the end of incubation, and dissolved CH₄ was extracted using the headspace method described in Sect. 2.2. As a consequence, dissolved CH₄ samples were collected in triplicate for each of the initial and final conditions of one incubation treatment.

To interpret CH₄ production in the incubation experiments (Sect. 2.3), phylogenetic composition of methanogens in the surface soil was additionally analyzed in 2016 using 16S rRNA gene sequencing. In July 2016, soil samples from 10 cm depth were collected in 10 mL plastic tubes in triplicate in the same four wet areas as the anaerobic incubation experiments, and kept frozen until analysis. DNA was extracted from 3 g of the soil samples as described by Ikeda et al. (2004). Extracted DNA was purified using the OneStep™ PCR Inhibitor Removal Kit (Zymo Research, Calif.) and quantified using the QuantiT PicoGreen dsDNA assay Kit (Invitrogen, Carlsbad, Calif.). Amplicon sequencing was conducted targeting the V3/V4 regions of 16S rRNA genes (Caporaso et al. 2011). Sequences obtained were processed through the QIIME pipeline (Caporaso et al. 2010). A representative sequence was picked from each operational taxonomic unit (OTU), and the Greengenes reference database (version 13.8) was used to assign taxonomic information and calculate the relative abundance of methanogenic archaea present.

For CH₄ oxidation, surface organic layers (0–13 cm depth) were sampled at sphagnum_K and sedge_K in July 2012, and then kept in a refrigerator until the experiment (6 days). These soil samples were cut into small pieces and mixed well with air. Ten grams (about 40 mL) of soil sample were transferred into plastic syringes (maximum 120 mL) in quadruplicate for each sampling location. Approximately 80 mL of air and 0.2–2 mL of 25% CH₄ gas were added to each syringe so the total volume in each syringe was 120 mL and the headspace CH₄ concentration was 5.0×10^2 – 4.8×10^3 ppm. Syringes were preserved in water and incubated at 8 °C for 8 days. Headspace gas was sampled on day 0, day 4, and day 8 from each syringe into 20 mL pre-evacuated glass vials with rubber septa. Consequently, quadruplicate gas samples were collected for each location and each sampling day.

2.4 Sample analysis and data processing

Methane concentrations in air samples were analyzed using a gas chromatograph (HP6890 series G1530A, Hewlett Packard, USA) equipped with a flame ionization detector and a CP-carboplot capillary column (Varian, USA). Methane flux was calculated from CH₄ concentration in chamber headspace by a linear regression of two to three concentration values against the time elapsed since chamber closure. The detection limit of CH₄ flux for each observation was calculated as 0.8–2.4 mg CH₄ m⁻² day⁻¹, based on whether the change of chamber CH₄ concentration during the observation was significant relative to the precision of CH₄ concentration analysis. Regression r^2 was calculated (formally) as ≥ 0.87 , when the flux value was larger than 2 mg CH₄ m⁻² day⁻¹. Dissolved CH₄ concentrations were obtained from calculation of the headspace method where equilibrations of CH₄ between gas and water phases are described by the Bunsen absorption coefficient of CH₄ (Yamamoto et al., 1976).

Carbon and hydrogen isotope ratios of in situ dissolved CH₄ and CH₄ samples from both incubation experiments were analyzed on a GC/GC/C/IRMS (modified after Sugimoto, 1996)—which is a continuous flow system consisting of two

gas chromatographs, a combustion reactor, and an isotope ratio mass spectrometer (MAT253, Thermo Fisher Scientific, USA) —and on a GC/GC/P/IRMS (P: pyrolysis in a HTC reactor of GC IsoLink, Thermo Fisher Scientific), respectively. Carbon and hydrogen isotope ratios obtained were represented relative to VPDB and VSMOW, respectively. Precisions of the analyses were $\pm 0.2\%$ and $\pm 2\%$ for $\delta^{13}\text{C}$ and δD , respectively. When calculating $\delta^{13}\text{C}$ and δD of dissolved CH_4 , the effect of CH_4 in background air was removed based on the mass balance. In the aerobic incubation experiments, the fractionation factors of CH_4 oxidation for carbon and hydrogen were calculated using the following Rayleigh distillation equation:

$$\ln \frac{R_t}{R_0} = \left(\frac{1}{\alpha_{\text{ox}}} - 1 \right) \ln \frac{[\text{CH}_4]_t}{[\text{CH}_4]_0}, \quad (1)$$

where R_0 and R_t represent isotope ratios under initial conditions and at time t , respectively; α_{ox} is the fractionation factor for CH_4 oxidation (defined so that $\alpha_{\text{ox}} > 1$); and $[\text{CH}_4]_0$ and $[\text{CH}_4]_t$ are CH_4 concentrations under initial conditions and at time t , respectively.

All statistical tests for detecting differences in CH_4 fluxes or dissolved CH_4 concentrations were conducted using R software (version 3.3.3). Mann-Whitney's U test was applied to compare magnitudes between two years of data and Steel-Dwass's multiple comparison test was used to compare magnitudes among three years or more of data.

15 2.5 Meteorological data

Air temperature and precipitation observed at a weather station in Chokurdakh (WMO station 21946) were used to investigate interannual variations in meteorological conditions during our observation period of CH_4 flux (2009–2013) and during the preceding two years (2007–2008). The distance between the weather station and our farthest observation site (site V) is approximately 45 km (Fig. 1). These data were obtained from GHCN-Daily, a NOAA database (Menne et al., 2012a, 2012b).

3 Results

3.1 Environmental factors

Soil wetness conditions and thaw depth differed among vegetation types (Table 1). Tree mounds had lower surface water content (2.1–17%) than wet areas (42–48%). Among the two types of wet areas, the water level was lower in wet areas of sphagnum mosses than those of sedges (Fig. 2). Wet areas of sedges experienced water levels higher than the ground surface (defined as 0 cm), reaching more than 10 cm above the ground surface. Corresponding with soil water content, the thaw depth was shallower at dry tree mounds (20–23 cm), and deeper in wet areas (31–56 cm). In wet areas, thaw depth became deeper from 2011 (22 ± 4 cm) to 2012 (25 ± 8 cm) and 2013 (35 ± 7 cm) in observations made during mid-July (Table S1). The overall average thaw depth observed on days when flux measurements were taken was 31 ± 12 cm ($n = 77$, 9–58 cm between Jul 3 and Aug 9).

Figure 2 shows persistently low annual precipitation (162–173 mm) from 2007 to 2009. In 2010, July air temperature was characteristically high (15.5 °C) accompanying low monthly precipitation (8 mm). These show dry conditions during our flux observations in 2009 and 2010. Parmentier et al. (2011) reported that water level was lower in summer 2009 than the previous two summers at a tundra research station (Kytalyk) in the vicinity, approximately 30 km to northwest of Chokurdakh. In contrast, precipitation in July 2011 was extremely high (94 mm) with a relatively mild temperature (13.0 °C), which caused an unusual wetting. High precipitation continued in August (74 mm) and September (67 mm) of the same year. Corresponding with this heavy rainfall, water levels were also high in 2011, and subsequent observations show a clear decrease from 2011 to 2013 in wet areas of sedges ($p < 0.005$). Water levels also declined in wet area of sphagnum mosses, with values of –9 cm, –10 cm, and less than –12 cm in 2011, 2012, and 2013, respectively.

10 3.2 CH₄ flux and dissolved CH₄ concentration

Obtained CH₄ flux shows clear spatial and interannual variations (Fig. 3). Individual flux measurements ($n = 143$ in total) are summarized as mean values for the main summer seasons. From 2011 to 2013, continuous flux observations (Table S2) were conducted in concert with dissolved CH₄ analysis, and the interannual variation during this period will be discussed in detail.

With regards to the spatial variation of CH₄ flux, tree mounds had consistently small values around the detection limit for all measurements (–4.9 to 1.9 mg CH₄ m^{–2} day^{–1}), while wet areas showed CH₄ emissions. From 2009 to 2013, the CH₄ flux in wet areas showed large interannual variations ranging from 36 to 140 mg CH₄ m^{–2} day^{–1}. The flux increased in 2011 when the wetting event occurred, then remained relatively large in 2012 (compared to 2009 and 2010). Moreover, the flux increased again from 2011/2012 to 2013 ($p < 0.05$). No statistically significant correlation was found when CH₄ flux was plotted against soil temperature (10 cm depth), thaw depth, or water level using all the data from wet areas (Fig. S1).

In addition to CH₄ flux, dissolved CH₄ concentration increased after the wetting event in 2011 (Fig. 4). From 2011 to 2012, CH₄ concentration in soil pore water at 10 cm depth (Fig. 4b) exhibited a sharp increase of one order of magnitude ($p < 0.005$). It remained high from 2012 to 2013, and the concentrations in surface water and that at 20 cm depth (Fig. 4a and c) also increased significantly over the same period ($p < 0.05$). No significant difference in concentration was observed at 30 cm depth between 2012 and 2013. In terms of vertical variation, the concentration in surface water was lower than that in soil pore water (10, 20, and 30 cm depth).

3.3 δ¹³C and δD of in situ dissolved CH₄

Variability of both δ¹³C and δD of dissolved CH₄ was smaller in deeper layers, showing different patterns between δ¹³C and δD, and across years (Fig. 5). The δ¹³C of dissolved CH₄ had similarly large ranges (–68‰ to –40‰) in surface water and at 10 cm and 20 cm depths, compared to a small range (–53‰ to –46‰) at 30 cm depth. The δD of dissolved CH₄ was variable only in surface water (–415‰ to –308‰) and at 10 cm depth (–417‰ to –341‰), whereas it had a constant value of around –408‰ at 20 cm and 30 cm depths. Additionally, δ¹³C values approached a relatively high value (approximately –50‰) at depth, while δD values converged to almost their lowest value. In terms of interannual variations in δ¹³C and δD of dissolved

CH₄ from 2011 to 2013, both δ¹³C and δD values in surface soil pore water (10 cm depth) were scattered more widely in 2011, showing standard deviations (SD) of 6.6‰ and 24‰, respectively, whereas their ranges were smaller in 2012 and 2013 (SD: 3.3‰ and 17‰ at maxima, respectively).

As shown in Fig. 6, convergence of δ¹³C and δD of dissolved CH₄ is associated with dissolved CH₄ concentrations.

- 5 The δ¹³C and δD values of dissolved CH₄, including surface water and 10 cm depth, converged at high CH₄ concentrations to the values seen in deeper soil layers (δ¹³C = -50 ± 5‰ and δD = -408 ± 5‰ at > 200 μmol CH₄ L⁻¹).

3.4 Soil incubation experiments and microbial community analysis

In the anaerobic incubation experiment, the CH₄ production rate was different among sampling locations (Fig. 7); the rate was higher for sedge_K and sedge_B (0.66 ± 0.15 μmol day⁻¹ and 0.43 ± 0.09 μmol day⁻¹, respectively) than sedge_V and sphagnum_K (0.24 ± 0.02 μmol day⁻¹ and 0.08 ± 0.01 μmol day⁻¹, respectively). In sedge_K, the sampling location tested in detail, production was more rapid for shallower soil layers among the 10 cm, 20 cm and 30 cm depths (0.66 ± 0.15 μmol day⁻¹, 0.33 ± 0.06 μmol day⁻¹, 0.003 ± 0.004 μmol day⁻¹, respectively; *p* < 0.01 in Welch's ANOVA test), while no difference in the rate was found between incubation temperatures (0.66 ± 0.15 μmol day⁻¹ gdw⁻¹ at 5 °C and 0.74 ± 0.14 μmol day⁻¹ gdw⁻¹ at 10 °C, *p* > 0.5 in *t*-test). When the CH₄ production rate was high, the δ¹³C and δD values of produced CH₄ were less variable irrespective of sampling location, sampling depth, or incubation temperature. The δ¹³C value of produced CH₄ at a high production rate (> 0.26 μmol day⁻¹ gdw⁻¹) was -55 ± 4‰ (n = 12). Similarly, δD under rapid CH₄ production was -410 ± 9‰ (n = 12). These δ¹³C and δD values of CH₄ obtained under rapid production were mostly comparable with the δ values of in situ dissolved CH₄ that converged in deep soil layers (δ¹³C = -50 ± 2‰ at 30 cm depth and δD = -408 ± 5‰ at 20–30 cm depth; Fig. 5c and d), although δ¹³C values in the incubation experiment were slightly lower than those in situ.

In the microbial community analysis using 16S rRNA gene sequencing (Fig. 8), soil with high rates of CH₄ production shown in the incubation experiment (sedge_K and sedge_B as in Fig. 7) had more abundant total methanogens within the detected archaea than that with slow CH₄ production rates (sphagnum_K and sedge_V). Acetoclastic methanogens in the order Methanosarcinales were higher in proportion among methanogens in sedge_K and sedge_B, where δ¹³C values of produced CH₄ in the incubation were higher. In contrast, Methanosarcinales were fewer in proportion in sphagnum_K, where δ¹³C of the produced CH₄ was lower.

In the CH₄ oxidation experiment, CH₄ concentration in headspace declined continuously in every sample (Fig. S2). As CH₄ oxidation proceeded, both δD and δ¹³C of the remaining CH₄ increased with a linear relationship between them (Fig. 9, S2). Observed slope Δ(δD)/Δ(δ¹³C) was 11, indicating a much larger fractionation of hydrogen than carbon, regardless of vegetation types in wet areas (sphagnum or sedge). The hydrogen isotope fractionation factors of CH₄ oxidation calculated from the data shown in Fig. 9 were 1.25 and 1.16 for wet areas of sphagnum and sedge, respectively, while carbon isotope fractionations were 1.021 and 1.015, respectively.

4 Discussion

4.1 CH₄ flux at tree mounds and wet areas at the taiga-tundra boundary on the Indigirka River lowlands

Methane flux observed in our study was clearly larger at wet areas than at dry tree mounds (Table 1, Fig. 3). Such differences in CH₄ flux between wetland vegetation and dry areas with trees or shrubs is generally observed (van Huissteden et al., 2005; van der Molen et al., 2007; Flessa et al., 2008) and is consistent with the fact that CH₄ production requires reducing conditions in soil (Conrad, 2007). Our CH₄ flux in wet areas (36–140 mg CH₄ m⁻² day⁻¹) was comparable to that reported for wet tundras (32–101 mg CH₄ m⁻² day⁻¹) or permafrost fens (42–147 mg CH₄ m⁻² day⁻¹) in a database across permafrost zones compiled by Olefeldt et al. (2013). In forests, many studies have observed CH₄ absorption instead of emission (King et al., 1997; Dutaur and Verchot, 2007; Flessa et al., 2008; Morishita et al., 2014). However, our observations at tree mounds rarely found CH₄ absorption or emission. In addition, CH₄ was not consumed even under O₂- and CH₄-rich conditions in incubation experiments of tree mound soil from site K (Murase et al., 2014), indicating that a lack of methanotrophic bacterial activity limited CH₄ absorption at this vegetation type. Our results show that CH₄ emission from wet areas is expected to make a greater contribution to ecosystem-scale CH₄ exchange at the taiga-tundra boundary on the Indigirka River lowlands.

4.2 Methane flux, production, and oxidation responses to the wetting event

In 2009 and 2010 the CH₄ emission in wet areas was low (Fig. 3), even at relatively high soil temperatures in 2010 (Fig. S1), under dry conditions that were not directly observed in this study. The wetting event in 2011 initiated the high CH₄ emission that continued through 2013 despite decreasing water levels (Fig. 2). Moreover, a further increase in flux was observed in 2013, accompanying a build-up of dissolved CH₄ (2011–2013) as shown in Fig. 4.

These interannual variations from 2011 to 2013 could be caused by the development of reducing soil conditions over multiple years after the wetting event. Reducing soil conditions may have developed, to some extent, as a result of the extreme precipitation in the summer of 2011 (Fig. 2). The surface soil layer, particularly under high water levels, could eliminate O₂ from soil pore spaces due to water saturation. These reducing conditions may have been preserved by freezing of the soil throughout the following winter. Additionally, a surface soil layer saturated with ice could have prevented snowmelt water (rich in O₂) from infiltrating the soil during the spring thaw season of 2012 (Woo, 2012). These processes would have led to the continuation of reducing conditions in the soil, which were created in summer 2011, into 2012. Through further decomposition of soil organic matter with the consumption of O₂, reducing soil conditions may have been exacerbated in the water-saturated soil layer to a greater extent in summer 2012. Water in the saturated soil layer may be retained without exchange for a relatively long time in our study sites, because lateral runoff is assumed to be small in the flat floodplain that comprises the Indigirka River lowlands (Nassif and Wilson, 1975). In addition, deep percolation loss is prevented by the impermeable permafrost layer (Woo, 2012). From summer 2012 to summer 2013, reducing conditions in the soil may have been similarly prolonged, especially in the deep soil layer, despite the decrease in water level from

summer 2011 to summer 2013. This continuous soil reduction from 2011 to 2013 could have promoted CH₄ production and/or decreased CH₄ oxidation, which may explain the increase in dissolved CH₄ concentration and CH₄ flux in wet areas following the wetting event and continuing until 2013 (Fig. 3 and 4).

In 2011, $\delta^{13}\text{C}$ and δD of dissolved CH₄ (10 cm depth) were scattered broadly across a wide range, whereas in 2012 and 2013 the ranges were narrower and they clustered around a high $\delta^{13}\text{C}$ value (-50‰) and a low δD value (-408‰ ; Fig. 5b). Considering that δD increased much more rapidly than $\delta^{13}\text{C}$ in our oxidation experiment (Fig. 9), δD can be considered as a sensitive indicator of CH₄ oxidation. In contrast, $\delta^{13}\text{C}$ is not a good indicator because its fractionation factor of CH₄ oxidation (1.015-1.021) was similar to that of CH₄ diffusion (1.019; Chanton, 2005), thus the effects of CH₄ oxidation and diffusion cannot be discerned by $\delta^{13}\text{C}$. Additionally, δD of dissolved CH₄ (Fig. 5) was clearly lower in deeper layers (20 cm and 30 cm depths) than in shallower layers (surface water and 10 cm depth), which indicates δD showed CH₄ oxidation in situ as well, because shallower layers are provided with O₂ from the atmosphere and precipitation. The effect of CH₄ exchange between surface dissolved CH₄ and atmospheric CH₄ can be excluded, because all the dissolved CH₄ observed in this study was highly oversaturated ($> 0.3 \mu\text{mol L}^{-1}$, Fig. 4) compared to the equilibrium concentration of atmospheric CH₄ (4–5 nmol L⁻¹, assuming 1–10 °C water temperature and 2 ppm atmospheric CH₄ concentration; Yamamoto et al., 1976). Thus, δD values at 10 cm in 2011 were scattered broadly compared with those in 2012 and 2013 that clustered around a low value, suggesting that CH₄ oxidation was significant in the surface soil layer during the year of the wetting event (2011). In 2012 and 2013, CH₄ oxidation became insignificant, relative to the larger pool of dissolved CH₄ (Fig. 4). In the CH₄ production incubation experiment, $\delta^{13}\text{C}$ and δD of produced CH₄ were less variable at higher production rates ($\delta^{13}\text{C} = -55 \pm 4\text{‰}$ and $\delta\text{D} = -410 \pm 9\text{‰}$ as in Fig. 7). Analogously, those of in situ dissolved CH₄ converged at a high CH₄ concentration around similar values ($\delta^{13}\text{C} = -50 \pm 5\text{‰}$ and $\delta\text{D} = -408 \pm 5\text{‰}$ in Fig. 6). This suggests that δ values of produced CH₄ became almost constant under rapid CH₄ production in situ and that the convergence of δ values of dissolved CH₄ observed in situ reflect rapid CH₄ production. Hence, the narrow ranges of $\delta^{13}\text{C}$ and δD values of dissolved CH₄ at 10 cm depth observed in 2012 and 2013 ($\delta^{13}\text{C}$: around -50‰ and δD : around -408‰ , Fig. 5b) suggest enhanced CH₄ production relative to the wetting year (2011).

Multi-year effects of wetting on CH₄ flux through soil reduction processes have been previously proposed by Kumagai and Konno (1998) and Desyatkin et al. (2014) as one possible factor for explaining the increase in CH₄ flux after wetting. Kumagai and Konno (1998) reported a CH₄ flux increase at a temperate rice field in Japan one year after the rice field was irrigated and restored from farmland that had been drained for eight years. Desyatkin et al. (2014) observed flux increases at a thermokarst depression in boreal eastern Siberia during the second consecutive year of flooding following large volumes of precipitation. On the other hand, studies at natural wetlands in the northeastern USA (Smemo and Yavitt, 2006; Treat et al., 2007; Olson et al., 2013) and southern Canada (Moore et al., 2011) reported that interannual variations of CH₄ flux correspond with those of water level and/or precipitation in the current year. In our study area, multi-year soil reduction may be important because soil temperature is generally lower than 11 °C (10 cm to 30 cm depth; Fig. S1a and Iwahana et al., 2014) due to a shallow active layer underlain by permafrost. Therefore, decomposition of organic matter can

be slow (Treat et al., 2015), which would slowly decrease soil redox potential, allowing it to remain relatively high in the first year of wetting.

In the following two years (2012–2013), we observed redox potential values lower than -100 mV in wet areas (Table S6), which are well below the upper limit for CH_4 production in soil (Conrad, 2007; Street et al., 2016). Methane
5 production at a potential higher than -100 mV can also occur, because soil is heterogeneous and can have more reducing microsites than the rest of the bulk soil, where redox potential can be measured (Teh et al., 2005; Teh and Silver, 2006).

In addition to the multi-year soil reduction, it appears that the wetting event led to the thaw depth increase in wet areas from 2011 to 2013 (Table S1). Although thaw depth increased, summer air temperatures decreased from 2011 (7.7, 13.0 °C as June and July mean temperatures, respectively) to 2012 (7.4, 9.2 °C) and 2013 (6.6, 10.5 °C) as shown in Fig. 2.
10 The wetting event may have led to the CH_4 flux increase from 2011 to 2013 (Fig. 3) partly through the thaw depth increase, by thickening the soil layer where CH_4 production occurs (Nakano et al., 2000; van Huissteden et al., 2005). However, the clear increase in dissolved CH_4 concentration (Fig. 4) and the enhanced CH_4 production and less significant CH_4 oxidation reflected in our isotopic data (Fig. 5b) cannot be explained by the thaw depth increase. Additionally, in the incubation experiment of CH_4 production (Fig. 7), the CH_4 production rate under anaerobic conditions was slower in the deeper layer,
15 especially at 30 cm depth (mineral soil) compared to 10 cm and 20 cm depths (organic soil) in sedge_K. Treat et al. (2015) also reported, from a pan-Arctic synthesis of anaerobic incubations, that difference in soil types (organic/mineral) and that in substrate quality along depth are important controls on CH_4 production rate. Our results from the incubations suggests that the deep layer comprised of mineral soil, where CH_4 production becomes active when thaw depth increase, is not the main region for CH_4 production.

20 This study did not evaluate vegetation cover quantitatively, and the wetting event might have also led to some vegetation change (such as increase of sedges), although no drastic changes were found visually in the observed wet areas. Increase in cover by sedges might have raised CH_4 flux partly by providing labile organic substrate for CH_4 production or conduits for the CH_4 transport from the soil to the atmosphere (Chanton, 2005; Lai, 2009; Ström et al., 2015).

4.3 Process behind CH_4 production response

25 When CH_4 production is initiated after the onset of anoxia in rice paddy soil, it first occurs via hydrogenotrophic methanogenesis, and then by both hydrogenotrophic and acetoclastic methanogenesis, which increases CH_4 production rate (Conrad, 2007). Afterwards, the ratio of acetoclastic to hydrogenotrophic methanogenesis can stabilize (Roy et al., 1997). Considering that this ratio is an important control on isotopic compositions of produced CH_4 , stabilization of production pathways might explain the convergence in δ values of dissolved CH_4 at our study sites under high CH_4 concentration (Fig.
30 6), and the reduced variability of δ values of produced CH_4 in our experiment under rapid production conditions (Fig. 7). As acetoclastic methanogenesis leads to higher $\delta^{13}\text{C}$ in produced CH_4 than hydrogenotrophic methanogenesis (Sugimoto and Wada, 1993), acetoclastic methanogenesis may have been activated when dissolved CH_4 concentration or CH_4 production rate were high during our study. This interpretation is supported by the microbial community analysis (Fig. 8), where

acetoclastic methanogens of Methanosarcinales were more abundant in wet areas, with a higher $\delta^{13}\text{C}$ of produced CH_4 in the incubation. Therefore, the high and less-variable $\delta^{13}\text{C}$ values observed at 10 cm depth in 2012 and 2013 (Fig. 5b) suggest a greater contribution from acetoclastic methanogenesis compared to the wetting year (2011). Similar to findings from rice paddy soil (Conrad, 2007), acetoclastic methanogenesis may have experienced delayed activation after anoxic conditions began in 2011, which could also have promoted CH_4 production in 2012 and 2013.

5 Concluding remarks

At the taiga-tundra boundary on the Indigirka River lowlands, we observed an increase in CH_4 flux in wet areas following the wetting event in 2011, and a further increase in flux in 2013. Our results show interannual variations in $\delta^{13}\text{C}$ and δD of dissolved CH_4 , and when compared with our incubation experiments, suggest both enhancement of CH_4 production and less significance of CH_4 oxidation in 2012 and 2013 compared to 2011. This enhancement of production could be partly caused by activation of acetoclastic methanogenesis following the development of reducing soil conditions after the wetting event. Analyses of isotopic compositions of CH_4 both in situ and in incubation experiments can be combined to investigate the effects of CH_4 production and oxidation on these isotopic compositions, and to clarify the relationship between CH_4 flux and wetting. In the future, measuring the $\delta^{13}\text{C}$ of dissolved CO_2 would be useful to further validate activation of acetoclastic methanogenesis (Sugimoto and Wada, 1993; McCalley et al., 2014; Itoh et al., 2015). Outside of these processes, the wetting event might have affected CH_4 flux partly via the thaw depth increase or some amount of vegetation change. It would be useful to analyze $\delta^{13}\text{C}$ and δD values of emitted CH_4 in order to assess changes in CH_4 transport (such as by increase of sedge cover) and to investigate the relationship between dissolved CH_4 concentration and CH_4 flux in detail (Chanton, 2005).

In recent years, strong storm activity and wetting events in terrestrial ecosystems have been observed in northern regions (Iijima et al., 2016). A wetting event at the taiga-tundra boundary can switch micro-reliefs with large interannual variations in soil wetness conditions to significant CH_4 sources; we observed clear increases in CH_4 flux at wet areas after the wetting event. In order to predict CH_4 flux following a wetting event in a permafrost ecosystem, our results show the multi-year process of soil reduction affected by the duration of water saturation in the active layer.

25 Author contribution

Ryo Shingubara and Atsuko Sugimoto designed the experiments and Ryo Shingubara carried them out. Go Iwahana, Shunsuke Tei, Liang Maochang, Shinya Takano, Tomoki Morozumi, and Trofim C. Maximov helped with sampling, in situ data collection, and preparing resources for the fieldwork. Jun Murase contributed to the laboratory analysis. Ryo Shingubara prepared the manuscript with contributions from all co-authors.

30

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

5 This research was supported by JSPS Grant-in-Aid for Scientific Research (KAKENHI) numbers 21403011 and 16H05618, a Grant-in-Aid from the Global COE Program ‘Establishment of Center for Integrated Field Environmental Science’ (IFES-GCOE) funded by the Ministry of Education, Culture, Sports, Science and Technology-Japan (MEXT), JST (Strategic International Collaborative Research Program: SICORP)-EU cooperative research project ‘Dynamics of permafrost and methane emission in Arctic terrestrial ecosystem in Eastern Siberia’, the Green Network of Excellence (GRENE) Program
10 funded by MEXT, and the COPERA (C budget of ecosystems and cities and villages on permafrost in eastern Russian Arctic) project funded by the Belmont Forum through JST. We sincerely thank A. Kononov, R. Petrov, E. Starostin, A. Alexeeva and other members of the Institute for Biological Problems of Cryolithozone SB RAS, and T. Stryukova, S. Ianygin and other staff at the Allikhovsky Ulus Inspectorate of Nature Protection for supporting our fieldwork in the vicinity of Chokurdakh. We also wish to acknowledge help from Y. Hoshino, S. Nunohashi, K. Tanaka, K. Saito, H. Kudo, and R.
15 Shakhmatov in our research group at Hokkaido University.

References

- Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N., Peña, A. G.,
Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T., Knights, D., Koenig, J. E., Ley, R. E., Lozupone, C. A.,
20 McDonald, D., Muegge, B. D., Pirrung, M., Reeder, J., Sevinsky, J. R., Turnbaugh, P. J., Walters, W. A., Widmann, J., Yatsunenko, T., Zaneveld, J. and Knight, R.: QIIME allows analysis of high-throughput community sequencing data, *Nature Methods*, 7(5), 335–336, doi:10.1038/nmeth.f.303, 2010.
- Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh, P. J., Fierer, N. and Knight, R.: Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample, *PNAS*, 108(Supplement 1),
25 4516–4522, doi:10.1073/pnas.1000080107, 2011.
- Chanton, J. P.: The effect of gas transport on the isotope signature of methane in wetlands, *Organic Geochemistry*, 36(5), 753–768, doi:10.1016/j.orggeochem.2004.10.007, 2005.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S. and Thornton, P.: Carbon and other biogeochemical cycles, in
30 *Climate change 2013: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of*

- the Intergovernmental Panel on Climate Change, pp. 465–570, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2014.
- Conrad, R.: Microbial ecology of methanogens and methanotrophs, in *Advances in Agronomy*, vol. 96, pp. 1–63, Academic Press., 2007.
- 5 Desyatkin, A. R., Takakai, F. and Hatano, R.: Flood effect on CH₄ emission from the alas in Central Yakutia, East Siberia, *Soil Science and Plant Nutrition*, 60(2), 242–253, doi:10.1080/00380768.2014.883486, 2014.
- Dlugokencky, E. J., Bruhwiler, L., White, J. W. C., Emmons, L. K., Novelli, P. C., Montzka, S. A., Masarie, K. A., Lang, P. M., Crotwell, A. M., Miller, J. B. and Gatti, L. V.: Observational constraints on recent increases in the atmospheric CH₄ burden, *Geophysical Research Letters*, 36(18), L18803, doi:10.1029/2009GL039780, 2009.
- 10 Dutaur, L. and Verchot, L. V.: A global inventory of the soil CH₄ sink, *Global Biogeochemical Cycles*, 21(4), GB4013, doi:10.1029/2006GB002734, 2007.
- Flessa, H., Rodionov, A., Guggenberger, G., Fuchs, H., Magdon, P., Shibistova, O., Zrazhevskaya, G., Mikheyeva, N., Kasansky, O. A. and Blodau, C.: Landscape controls of CH₄ fluxes in a catchment of the forest tundra ecotone in northern Siberia, *Global Change Biology*, 14(9), 2040–2056, doi:10.1111/j.1365-2486.2008.01633.x, 2008.
- 15 van Huissteden, J., Maximov, T. C. and Dolman, A. J.: High methane flux from an arctic floodplain (Indigirka lowlands, eastern Siberia): METHANE FLUX ARCTIC FLOODPLAIN SIBERIA, *Journal of Geophysical Research: Biogeosciences*, 110(G2), G02002, doi:10.1029/2005JG000010, 2005.
- Iijima, Y., Nakamura, T., Park, H., Tachibana, Y. and Fedorov, A. N.: Enhancement of Arctic storm activity in relation to permafrost degradation in eastern Siberia, *International Journal of Climatology*, 36(13), 4265–4275, doi:10.1002/joc.4629, 2016.
- 20 Ikeda, S., Watanabe, K. N., Minamisawa, K. and Ytow, N.: Evaluation of Soil DNA from Arable Land in Japan Using a Modified Direct-extraction Method, *Microbes Environ.*, 19(4), 301–309, doi:10.1264/jsme2.19.301, 2004.
- Itoh, M., Shimamura, T., Ohte, N. and Takemon, Y.: Differences in hydrophyte life forms induce spatial heterogeneity of CH₄ production and its carbon isotopic signature in a temperate bog peatland: $\delta^{13}\text{C}_{\text{CH}_4}$ variations in hydrophyte species, *Journal of Geophysical Research: Biogeosciences*, 120(7), 1177–1195, doi:10.1002/2014JG002881, 2015.
- 25 Iwahana, G., Takano, S., Petrov, R. E., Tei, S., Shingubara, R., Maximov, T. C., Fedorov, A. N., Desyatkin, A. R., Nikolaev, A. N., Desyatkin, R. V. and Sugimoto, A.: Geocryological characteristics of the upper permafrost in a tundra-forest transition of the Indigirka River Valley, Russia, *Polar Science*, 8(2), 96–113, doi:10.1016/j.polar.2014.01.005, 2014.
- King, G.: Responses of atmospheric methane consumption by soils to global climate change, *Global Change Biology*, 3(4), 351–362, doi:10.1046/j.1365-2486.1997.00090.x, 1997.
- 30 Kumagai K. and Konno Y.: Methane emissions from rice paddy fields after upland farming, *Japanese Journal of Soil Science and Plant Nutrition*, 69(4), 333–339, doi:10.20710/dojo.69.4_333, 1998 (in Japanese, with English abstract).
- Lai, D. Y. F.: Methane dynamics in northern peatlands: A review, *Pedosphere*, 19(4), 409–421, doi:10.1016/S1002-0160(09)00003-4, 2009.

- Liang, M., Sugimoto, A., Tei, S., Bragin, I. V., Takano, S., Morozumi, T., Shingubara, R., Maximov, T. C., Kiyashko, S. I., Velivetskaya, T. A. and Ignatiev, A. V.: Importance of soil moisture and N availability to larch growth and distribution in the Arctic taiga-tundra boundary ecosystem, northeastern Siberia, *Polar Science*, 8(4), 327–341, doi:10.1016/j.polar.2014.07.008, 2014.
- 5 Marik, T., Fischer, H., Conen, F. and Smith, K.: Seasonal variations in stable carbon and hydrogen isotope ratios in methane from rice fields: Seasonal variations in CH₄ isotope ratios from rice, *Global Biogeochemical Cycles*, 16(4), 1094, doi:10.1029/2001GB001428, 2002.
- McCalley, C. K., Woodcroft, B. J., Hodgkins, S. B., Wehr, R. A., Kim, E.-H., Mondav, R., Crill, P. M., Chanton, J. P., Rich, V. I., Tyson, G. W. and Saleska, S. R.: Methane dynamics regulated by microbial community response to permafrost thaw, *Nature*, 514(7523), 478–481, doi:10.1038/nature13798, 2014.
- 10 Menne, M. J., Durre, I., Vose, R. S., Gleason, B. E. and Houston, T. G.: An overview of the Global Historical Climatology Network-Daily database, *J. Atmos. Oceanic Technol.*, 29(7), 897–910, doi:10.1175/JTECH-D-11-00103.1, 2012a.
- Menne, M. J., Durre, I., Korzeniewski, B., McNeal, S., Thomas, K., Yin, X., Anthony, S., Ray, R., Vose, R. S., Gleason, B. E. and Houston, T. G.: Global Historical Climatology Network - Daily (GHCN-Daily), Version 3.22, NOAA National Data Climatic Data Center, doi:10.7289/V5D21VHZ, 2012b.
- 15 van der Molen, M. K., van Huissteden, J., Parmentier, F. J. W., Petrescu, A. M. R., Dolman, A. J., Maximov, T. C., Kononov, A. V., Karsanaev, S. V. and Suzdalov, D. A.: The growing season greenhouse gas balance of a continental tundra site in the Indigirka lowlands, NE Siberia, *Biogeosciences*, 4(6), 985–1003, doi:10.5194/bg-4-985-2007, 2007.
- Moore, T. R., De Young, A., Bubier, J. L., Humphreys, E. R., Lafleur, P. M. and Roulet, N. T.: A multi-year record of methane flux at the Mer Bleue Bog, southern Canada, *Ecosystems*, 14(4), 646, doi:10.1007/s10021-011-9435-9, 2011.
- 20 Morishita, T., Matsuura, Y., Kajimoto, T., Osawa, A., Zyryanova, O. A. and Prokushkin, A. S.: CH₄ and N₂O dynamics of a *Larix gmelinii* forest in a continuous permafrost region of central Siberia during the growing season, *Polar Science*, 8(2), 156–165, doi:10.1016/j.polar.2014.01.004, 2014.
- Murase, J., Sugimoto, A., Shingubara, R. and Maximov, T. C.: Methane oxidation potential of Arctic wetland soil of a taiga-tundra ecotone in Northeastern Siberia, Yokohama, Japan., 2014.
- 25 Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T. and Zhang, H.: Anthropogenic and natural radiative forcing, in *Climate change 2013: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 658–740, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.
- 30 Nakano, T., Kuniyoshi, S. and Fukuda, M.: Temporal variation in methane emission from tundra wetlands in a permafrost area, northeastern Siberia, *Atmospheric Environment*, 34(8), 1205–1213, doi:10.1016/S1352-2310(99)00373-8, 2000.
- Nassif, S. H. and Wilson, E. M.: The Influence of slope and rain intensity on runoff and infiltration, *Hydrological Sciences Bulletin*, 20(4), 539–553, doi:10.1080/02626667509491586, 1975.

- Nisbet, E. G., Dlugokencky, E. J. and Bousquet, P.: Methane on the Rise—Again, *Science*, 343(6170), 493–495, doi:10.1126/science.1247828, 2014.
- Olefeldt, D., Turetsky, M. R., Crill, P. M. and McGuire, A. D.: Environmental and physical controls on northern terrestrial methane emissions across permafrost zones, *Global Change Biology*, 19(2), 589–603, doi:10.1111/gcb.12071, 2013.
- 5 Olson, D. M., Griffis, T. J., Noormets, A., Kolka, R. and Chen, J.: Interannual, seasonal, and retrospective analysis of the methane and carbon dioxide budgets of a temperate peatland, *Journal of Geophysical Research: Biogeosciences*, 118(1), 226–238, doi:10.1002/jgrg.20031, 2013.
- Parmentier, F. J. W., Huissteden, J. van, Molen, M. K. van der, Schaepman-Strub, G., Karsanaev, S. A., Maximov, T. C. and Dolman, A. J.: Spatial and temporal dynamics in eddy covariance observations of methane fluxes at a tundra site in northeastern Siberia, *Journal of Geophysical Research: Biogeosciences*, 116(G3), doi:10.1029/2010JG001637, 2011.
- 10 Preuss, I., Knoblauch, C., Gebert, J. and Pfeiffer, E.-M.: Improved quantification of microbial CH₄ oxidation efficiency in arctic wetland soils using carbon isotope fractionation, *Biogeosciences*, 10(4), 2539–2552, doi:https://doi.org/10.5194/bg-10-2539-2013, 2013.
- Roy, R., Klüber, H. D. and Conrad, R.: Early initiation of methane production in anoxic rice soil despite the presence of oxidants, *FEMS Microbiol Ecol*, 24(4), 311–320, doi:10.1111/j.1574-6941.1997.tb00448.x, 1997.
- 15 Smemo, K. A. and Yavitt, J. B.: A multi-year perspective on methane cycling in a shallow peat fen in central New York State, USA, *Wetlands*, 26(1), 20–29, doi:10.1672/0277-5212(2006)26[20:AMPOMC]2.0.CO;2, 2006.
- Street, L. E., Dean, J. F., Billett, M. F., Baxter, R., Dinsmore, K. J., Lessels, J. S., Subke, J.-A., Tetzlaff, D. and Wookey, P. A.: Redox dynamics in the active layer of an Arctic headwater catchment; examining the potential for transfer of dissolved methane from soils to stream water, *Journal of Geophysical Research: Biogeosciences*, 121(11), 2776–2792, doi:10.1002/2016JG003387, 2016.
- 20 Ström, L., Falk, J. M., Skov, K., Jackowicz-Korczynski, M., Mastepanov, M., Christensen, T. R., Lund, M. and Schmidt, N. M.: Controls of spatial and temporal variability in CH₄ flux in a high arctic fen over three years, *Biogeochemistry*, 125(1), 21–35, doi:10.1007/s10533-015-0109-0, 2015.
- 25 Sugimoto, A.: GC/GC/C/IRMS system for carbo isotope measurement of low level methane concentration, *Geochemical Journal*, 30(3), 195–200, doi:10.2343/geochemj.30.195, 1996.
- Sugimoto, A. and Wada, E.: Carbon isotopic composition of bacterial methane in a soil incubation experiment: Contributions of acetate and CO₂/H₂, *Geochimica et Cosmochimica Acta*, 57(16), 4015–4027, doi:10.1016/0016-7037(93)90350-6, 1993.
- 30 Sugimoto, A. and Wada, E.: Hydrogen isotopic composition of bacterial methane: CO₂/H₂ reduction and acetate fermentation, *Geochimica et Cosmochimica Acta*, 59(7), 1329–1337, doi:10.1016/0016-7037(95)00047-4, 1995.
- Teh, Y. A. and Silver, W. L.: Effects of soil structure destruction on methane production and carbon partitioning between methanogenic pathways in tropical rain forest soils, *Journal of Geophysical Research: Biogeosciences*, 111(G1), G01003, doi:10.1029/2005JG000020, 2006.

- Teh, Y. A., Silver, W. L. and Conrad, M. E.: Oxygen effects on methane production and oxidation in humid tropical forest soils, *Global Change Biology*, 11(8), 1283–1297, doi:10.1111/j.1365-2486.2005.00983.x, 2005.
- Treat, C. C., Bubier, J. L., Varner, R. K. and Crill, P. M.: Timescale dependence of environmental and plant-mediated controls on CH₄ flux in a temperate fen, *Journal of Geophysical Research*, 112(G1), doi:10.1029/2006JG000210, 2007.
- Treat, C. C., Natali, S. M., Ernakovich, J., Iversen, C. M., Lupascu, M., McGuire, A. D., Norby, R. J., Chowdhury, T. R., Richter, A., Šantrůčková, H., Schädel, C., Schuur, E. A. G., Sloan, V. L., Turetsky, M. R. and Waldrop, M. P.: A pan-Arctic synthesis of CH₄ and CO₂ production from anoxic soil incubations, *Global Change Biology*, 21(7), 2787–2803, doi:10.1111/gcb.12875, 2015.
- 10 Woo, M.-K.: *Permafrost hydrology*, Springer, Berlin, Heidelberg, Germany, <https://doi.org/10.1007/978-3-642-23462-0>, 2012.
- Yabuki, H., Park, H., Suzuki, R. and Ohata, T.: *Baseline Meteorological Data in Siberia (BMDS) Version 5.0*, 5.00, Arctic Data archive System (ADS), Japan, [online] Available from: <https://ads.nipr.ac.jp/dataset/A20131107-002>, 2011.
- 15 Yamamoto, S., Alcauskas, J. B. and Crozler, T. E.: Solubility of methane in distilled water and seawater, *Journal of Chemical and Engineering Data*, 21(1), 78–80, doi:10.1021/je60068a029, 1976.


Table 1. Observation points of chamber CH₄ flux. Concentration and isotopic compositions of dissolved CH₄ were also observed in the following wet areas.

Site	Landscape	Observation points and surface conditions	Dominant vegetation	Volumetric water content (%) ^b	Thaw depth (cm) ^c
V (Verkhny Khatistakha) 70° 15' N 147° 28' E	Larch forest and wetland	tree mound_V	Green moss, <i>Larix gmelinii</i>	17 ± 5 (n = 3)	23 ± 3 (n = 5)
		sedge_V (wet area)	<i>Carex spp.</i> , <i>Eriophorum angustifolium</i>	48 ± 4 (n = 3)	56 ± 3 (n = 4)
K (Kodac) ^a 70° 34' N 148° 16' E	Typical taiga-tundra boundary	tree mound_K	Green moss, <i>Larix gmelinii</i>	2.1 ± 0.6 (n = 4)	23 ± 4 (n = 9)
		sphagnum_K (wet area)	<i>Sphagnum squarrosum</i>	42 ± 5 (n = 6)	31 ± 8 (n = 15)
		sedge_K (wet area)	<i>Eriophorum angustifolium</i>	44 ± 4 (n = 6)	32 ± 13 (n = 28)
B (Boydom) 70° 38' N 148° 09' E	Low-centered polygon	tree mound_B	Green moss, <i>Larix gmelinii</i>	6 ± 2 (n = 5)	20 ± 4 (n = 8)
		sedge_B (wet area)	<i>Eriophorum angustifolium</i>	46 ± 2 (n = 5)	36 ± 9 (n = 8)


^a Site K was previously named as Kryvaya (Iwahana et al., 2014) or Kodak (Liang et al., 2014).

5 ^b Observed for the surface soil layer down to 20 cm on 1 to 3 days in July 2011 at each observation point (see Table S2 for detailed observation dates). Standard deviations are shown.


^c Observed from early July to early August during 2010–2013 (see Table S1 for the interannual variation and Table S2 for detailed observation dates). Standard deviations are shown.



5 **Figure 1: Locations of study sites. (a) Location of study region in northeastern Siberia (the Generic Mapping Tools 5.0.0). (b) Satellite image of Indigirka River lowland around Chokurdakh village (Ckh: 70° 37' N, 147° 55' E) from Landsat 8. Observation sites (V, K, B) were selected in this region alongside the main stem and a tributary of the Indigirka River.**



5 **Figure 2: Interannual variations in precipitation (bars) and air temperature (solid lines) observed at a weather station at Chokurdakh for the cold season with snow cover (C: total from October in the previous year to May in the current year) and the warm season (JJAS), and water level (dotted lines) measured in wet areas of sedges (triangle) and sphagnum mosses (square). Water level was very low (< -12 cm) in the wet area of sphagnum in 2013, and could not be measured. Error bars represent standard deviations. Methane flux was observed during the main summers (early July to early August) from 2009 to 2013.**



5 **Figure 3: Interannual variations in averaged CH₄ flux in tree mounds (denoted as “Tr”) and wet areas (“Wt”) for main summer seasons from 2009 to 2013. Replication numbers (“n”) are shown for each averaged flux value, and standard deviations are represented by error bars. Different letters show statistical interannual differences in the flux values for wet areas. See Table S2 for flux values at respective observation points.**




Figure 4: Interannual variation in dissolved CH₄ concentration in (a) surface water and soil pore water at (b) 10 cm, (c) 20 cm, and (d) 30 cm depths in wet areas from 2011 (wetting event) to 2013. Different letters in each panel denote statistical differences among years in averaged concentration across the four wet areas ($p < 0.05$). Error bars represent standard deviations. See Table S3 for numerical values of dissolved CH₄ concentrations.




Figure 5: In situ $\delta^{13}\text{C}$ versus δD of dissolved CH_4 in (a) surface water and soil pore water at (b) 10 cm depth, (c) 20 cm depth, and (d) 30 cm depth from the wet event in 2011 to 2013. Individual delta values are shown here. See Table S3 for averaged delta values for each observation point and each year.

5





Figure 6: In situ (a) $\delta^{13}\text{C}$ and (b) δD versus concentration of dissolved CH_4 at four depths (surface water, 10 cm, 20 cm, and 30 cm) in wet areas from 2011 to 2013.



5 Figure 7: (a) $\delta^{13}\text{C}$ and (b) δD of produced CH₄ versus CH₄ production rate in the anaerobic soil incubation experiment. Production rates are shown in moles of produced CH₄ per day and per weight of dry soil in gram. Soil samples were collected at four observation points (sphagnum_K, sedge_V, sedge_K, and sedge_B) at three depths (10 cm, 20 cm, and 30 cm) and incubated at two temperatures (5 °C and 10 °C). These samples contain organic layers except for those collected at 30 cm. Error bars represent standard deviations.





Figure 8: Phylogenetic compositions of methanogenic archaea by order in wet areas. Soil samples (organic layers) were taken in triplicate from 10 cm depth in each wet area in July 2016. See Table S5 for detailed results.



5 **Figure 9: Enrichment of D/H (CH₄) and ¹³C/¹²C (CH₄) through CH₄ oxidation during the aerobic incubation experiment of surface organic layers from wet areas of sphagnum mosses and sedges in site K. Individual delta values of the headspace CH₄ from each incubated syringe and each day are shown. Initial isotopic compositions of the headspace CH₄ were -66‰ to -65‰ for δ¹³C and -167‰ to -162‰ for δD.**