

Dear editor,

We have now made all the technical corrections suggested by the referees except the one (#1) below which we disagree with. On behalf of the authors, I would like to thank you for taking the time to review this manuscript.

5 Referee #2:

1. I find 'forest floor respiration' somewhat misleading as respired CO₂ originates also from deeper soil layer - consider to rename it to soil respiration or soil CO₂ efflux.

We disagree because the respiration we measured did not only include soil respiration but also plant respiration. We can't use soil CO₂ efflux either as it by definition does not include the above-ground parts of the plants, only the roots. We think that
10 'forest floor respiration', which we defined as the sum of autotrophic and heterotrophic respiration (Pg. 7 L. 21-22), is the best definition to use in our case.

5. Table 1: Soil properties. Is the classification as 'humus layer' correct? The extremely low density would rather suggest that it is a 'litter layer' that is also more clearly distinguishable from peat. What is the thickness?

15 Yes, it should be litter layer and it's now corrected. Unfortunately, the litter layer thickness was not recorded.

Best regards,

Mika Korkiakoski

Greenhouse gas and energy fluxes in a boreal peatland forest after clearcutting

Mika Korhikoski¹, Juha-Pekka Tuovinen¹, Timo Penttilä², Sakari Sarkkola², Paavo Ojanen³, Kari Minkkinen³, Juuso Rainne¹, Tuomas Laurila¹, Annalea Lohila¹

5 ¹Finnish Meteorological Institute, P.O. Box 503, FI-00101 Helsinki, Finland

²Natural Resources Institute Finland, Viikinkaari 4, 00790 Helsinki, Finland

³University of Helsinki, Department of Forest Sciences, P.O. Box 27, 00014 University of Helsinki, Finland

Correspondence to: Mika Korhikoski (mika.korhikoski@fmi.fi)

Abstract.

10 The most common forest management method in Fennoscandia is rotation forestry including clearcutting and forest regeneration. In clearcutting, stem wood is removed and the logging residues are either removed or left on site. Clearcutting changes the microclimate and vegetation structure at the site, both affecting the site's carbon balance. Peat soils with poor aeration and high carbon densities are especially prone to such changes, and significant changes in greenhouse gas exchange can be expected. We measured carbon dioxide (CO₂) and energy fluxes with the eddy covariance method for two years (April
15 2016 – March 2018) after clearcutting a drained peatland forest. We observed a significant rise (23 cm) in the water table level and a large CO₂ source (first year: 3086 ± 148 g CO₂ m⁻² yr⁻¹, second year: 2072 ± 124 g CO₂ m⁻² yr⁻¹). These large CO₂ emissions resulted from the very low gross primary production (GPP) following the removal of photosynthesizing trees and the decline of ground vegetation, unable to compensate for the decomposition of logging residues and peat. During the second summer (June–August) after the clearcutting, GPP had already increased by 96% and total ecosystem respiration decreased by
20 14% from the previous summer. The mean daytime ratio of sensible to latent heat flux decreased after harvesting from 2.6 in May 2016 to 1.0 in August 2016 and in 2017 it varied mostly within 0.6–1.0. In April–September, the mean daytime sensible heat flux was 33% lower and latent heat flux 40% higher in 2017, probably due to the recovery of ground vegetation that increased evapotranspiration and albedo of the site. In addition to CO₂ and energy fluxes, we measured methane (CH₄) and nitrous oxide (N₂O) fluxes with manual chambers. After the clearcutting, the site turned from a small CH₄ sink into a small
25 source and from N₂O-neutral to a significant N₂O source. Compared to the large CO₂ emissions, the 100 yr global warming potential (GWP₁₀₀) of the CH₄ emissions was negligible. Also, the GWP₁₀₀ due to increased N₂O emissions was less than 10% of that of the CO₂ emission change.

1 Introduction

~~The n~~Northern peatlands cover approximately 3% of the Earth's land surface (Clarke and Rieley, 2010), most of which is located in the boreal region (Fischlin et al., 2007). ~~The b~~Boreal and subarctic peatlands are substantial reservoirs of carbon (C), storing 270–550 Pg C in total (Turunen et al., 2002; Yu, 2011). These reservoirs are affected by peatland drainage, which has been a common practice in northern countries. In Finland, more than half of the original peatland area of 100 000 km² has been drained, mostly for forestry (Päivänen and Hånell, 2012). Drainage lowers water table level (WTL), which accelerates the peat decomposition rate due to the increased availability of oxygen (Drzymulska, 2016). This often leads to higher carbon dioxide (CO₂) emissions from the soil (Maljanen et al., 2010; Ojanen et al., 2013). On the other hand, a well-performed drainage increases root aeration and nutrient availability, which enhances tree growth and CO₂ uptake by trees (Minkkinen et al., 2001). In addition, methane (CH₄) emissions decrease, and the soil may even turn into a net CH₄ sink if the site is well-drained (Maljanen et al., 2010; Ojanen et al., 2010). This results from the thickening of the oxic peat layer, which enhances CH₄ oxidation in the soil and from the disappearance of deep-rooted vascular plants, which in natural mires feed anaerobic microbes with fresh carbon and transport methane from the anoxic peat layer to atmosphere through their aerenchyma. In contrast to CH₄, nitrous oxide (N₂O) emissions often increase after drainage at nitrogen-rich minerotrophic peatlands (Maljanen et al., 2010; Ojanen et al., 2010, 2018) because nitrification (the process that produces nitrate and N₂O as a by-product) needs nitrogen and oxic conditions.

The most common method of forest management in Finland is rotation forestry including clearcutting and forest regeneration. In clearcutting, stem wood is removed, while the logging residues, i.e. foliage biomass, branches, stumps and roots, are either removed or left on site. After clearcutting, the site is prepared, e.g., by mounding or scalping, and a new tree generation is established by planting or sowing, or naturally from surrounding seed-trees. Removing the trees changes the local microclimate as more solar radiation reaches the soil surface, which increases soil temperature (Edwards and Ross-Todd, 1983; Londo et al., 1999) and its diel variation. This affects the carbon cycle, as higher soil temperature potentially increases soil respiration. In peatland forests, however, removing tree biomass raises the WTL by diminishing transpiration (Sarkkola et al., 2010), which in turn may decrease soil respiration and slow down the peat decomposition rate due to the reduced volume of aerated peat. On the other hand, photosynthesis is diminished due to the removal of trees and decline of ground vegetation. Mäkiranta et al. (2010) found that ground vegetation recovers rather fast in a peatland forest after clearcutting; however, after three years the recovery was still insufficient to compensate for the high ecosystem respiration produced by the large amount of fresh organic matter. The increase in ecosystem respiration is accounted for by the decay of logging residues and possibly also by increased soil organic matter decomposition under these residues (Mäkiranta et al., 2012; Ojanen et al., 2017). As boreal forest grows slowly, it may take from 8 up to 20 years to turn the forest back to a net carbon sink (Fredeen et al., 2007; Kolari et al., 2004; Mäkiranta et al., 2010; Pypker and Fredeen, 2002; Rannik et al., 2002; Schulze et al., 1999).

The CO₂ and energy fluxes between ecosystems and the atmosphere are commonly measured with the eddy covariance (EC) method (Aubinet et al., 2012). Previous work concerning clearcutting in forests with mineral soil has involved both EC (Clark

et al., 2004; Humphreys et al., 2005; Kolari et al., 2004; Kowalski et al., 2004, 2003; Machimura et al., 2005; Mamkin et al., 2016; Takagi et al., 2009; Williams et al., 2013) and soil chambers (Howard et al., 2004; Londo et al., 1999; Zerva and Mencuccini, 2005). Also, soil chambers have been used on clear-cut peatland forests (Mäkiranta et al., 2010; Pearson et al., 2012), whereas EC measurements have been lacking. In addition, most of this peatland forest gas exchange research has concentrated on carbon dynamics and overlooked the possibly important variations in energy and water fluxes.

Exchange of sensible and latent heat constitutes an important part of the surface energy balance, driving many local, regional and global scale climatological processes. The available energy for these turbulent fluxes is determined predominantly by net radiation and the ground heat flux. The partitioning of available energy between the sensible and latent heat fluxes, which can be described in terms of the Bowen ratio, is strongly influenced by vegetation and soil properties (Betts et al., 2007). Also, release of latent heat to the atmosphere, i.e., evapotranspiration, is a key component of the water cycle in peatlands and forests, also affecting soil moisture and forest productivity, which further affect the GHG fluxes in a forest. In a temperate deciduous broadleaf forest, evapotranspiration has been found to recover rapidly after clearcutting: the latent heat flux increased over the first three years, while the sensible heat flux declined correspondingly (Williams et al., 2013). It is unknown if this also happens in peatland forests.

In this study, we investigated the GHG and energy fluxes between the forest floor and the atmosphere and their environmental drivers after clearcutting in a boreal peatland forest in southern Finland. The study was based on EC measurements made for two years (April 2016 – March 2018) after clearcutting and on soil chamber measurements performed during the period of June 2015 – August 2017. Our specific aims were:

1. To estimate the magnitude of CO₂ fluxes and their environmental drivers after clearcutting.
2. To quantify the development of surface heat fluxes after clearcutting.
3. To investigate how soil CH₄ and N₂O fluxes change due to clearcutting.

2 Materials and methods

2.1 Measurement site

The measurements were set up in a nutrient-rich peatland forest called Lettosuo, which is located in southern Finland (60°38' N, 23°57' E). The site was drained with widely-spaced, manually dug ditches probably during the 1930s, then drained more effectively in 1969 and fertilized with phosphorus and potassium. The distance between the ditches is on average 45 m, and they were dug ca. 1 m deep but have since been partially filled with vegetation. After drainage and before clearcutting in 2016, the tree stand was dominated by Scots pine (*Pinus sylvestris*) with some pubescent birch (*Betula pubescens*). The understorey included mostly Norway spruce (*Picea abies*) and some small-sized pubescent birch. The tree stand was quite dense, which made the ground vegetation and moss layer patchy and variable due to irregular shading. Ground vegetation included herbs, such as *Trientalis europaea* and *Dryopteris carthusiana*, and dwarf shrubs, such as *Vaccinium myrtillus* (Bhuiyan et al., 2017).

The moss layer was dominated by *Pleurozium schreberi* and *Dicranum polysetum* with some *Sphagnum* mosses, such as *Sphagnum capillifolium*, *S. angustifolium* and *S. russowii*, in moist patches.

Before clearcutting, the average soil organic carbon content at the site was 58% and carbon distribution was rather uniform across different soil layers (Table 1). The total soil nitrogen content was lowest in ~~humus-litter layer~~ (1.7%) and varied from 2.2% up to 2.5% among the other soil layers. The CN ratio averaged at 27 which is typical for peatland forests with a fen history. The mean soil bulk density of the 0–40 cm layer was 0.14 g cm⁻³. The present-day peat thickness at Lettosuo varies mostly between 1.5 and 2.5 m. Assuming a thickness of 2 m results in soil organic carbon and nitrogen stock estimates of 156±72 kg C m⁻² and 6.4±2.9 kg N m⁻² (±SD).

Clearcutting was performed within a trapezoidal area of ~~2.35~~23500 hectares-m² (Fig. 1) between 29 February and 16 March 2016. After clearcutting, the logging residues were left at the site. The previous ground vegetation was almost totally destroyed in the harvesting operation and the following drastic increase in solar radiation. In the following summer, some species adapted to the open, well-lighted conditions; for example, *Rubus idaeus*, *Carex canescens* and *Dryopteris carthusiana* were observed here and there within the clear-cut site. Mounding was performed on 1–2 August 2016, and spruce (*Picea abies*) seedlings were planted in 2017. In addition to the clear-cut site, we conducted GHG measurements on a similarly sized control site located south-west of the clear-cut site, where the forest was left in its original state with similar tree stand and vegetation composition as at the pre-harvest clear-cut site.

2.2 Measurement system

An EC system to measure turbulent CO₂ and energy fluxes was set up in the north-eastern part of the clearcut (Fig. 1), and the measurements started on 8 April 2016, approximately three weeks after the clearcutting had ended. The measurements continued until 7 April 2018. From this point on, the time periods of 8 April 2016 – 7 April 2017 and 8 April 2017 – 7 April 2018 are referred to as the first and second EC measurement year, respectively. The EC system included a three-axis sonic anemometer (uSonic-3 Scientific, METEK, Elmshorn, Germany) for wind speed and air temperature and a closed-path infrared gas analyser (LI-7000, Licor Biosciences, Lincoln, NE, USA) for CO₂ and H₂O mixing ratios. A sampling rate of 10 Hz was used for the EC system. The measurement height was 2.75 m, the flow rate was about 6 l min⁻¹, and the length of the inlet tube (inner diameter 3.1 mm, Bevaline IV) was 8 m. The mouth of the inlet tube was positioned 15 cm below the sonic anemometer. Air with a zero CO₂ concentration was used as the reference gas when calibrating the gas analyser. The micrometeorological sign convention is used throughout the paper: a positive flux indicates a flux from the ecosystem to the atmosphere (net emission) and a negative flux indicates a flux from the atmosphere into the ecosystem (net uptake).

Auxiliary meteorological measurements were installed at the centre of the clearcut (80 m south of the EC mast) on 24 July 2015, i.e., before the clearcutting. A similar system was installed two weeks earlier for the control site (180 m south-east of the EC mast). Air temperature and relative humidity were measured at 2 m height (HMP155, Vaisala Corporation, Vantaa, Finland) and soil temperature profile at 5, 10, 20 and 30 cm depths (Pt100, Vaisala Corporation, Vantaa, Finland); the measurements also included net radiation (NR Lite2 Net Radiometer, Kipp & Zonen, Delft, The Netherlands),

photosynthetically active photon flux density (PPFD) (PQS1 PAR Quantum Sensor, Kipp & Zonen, Delft, The Netherlands) and ground heat flux (HFP01, Hukseflux Thermal Sensors B.V., Delft, The Netherlands). In addition to these, global radiation (Pyranometer CMP3, Kipp & Zonen, Delft, The Netherlands) was measured from another EC mast above the canopy of the surrounding forest (250 m south from the clear-cut EC mast). The data were collected by data loggers (QML201C, Vaisala Corporation, Vantaa, Finland) as 30 min averages.

2.3 EC data processing

Half-hourly turbulent fluxes were calculated using standard EC methods (Aubinet et al., 2012). The 10 Hz raw data were block-averaged, and a double rotation of the coordinate system was applied (McMillen, 1988). The time lag between the gas analyser and sonic anemometer signals was determined by cross-correlation analysis for each flux variable and 30 min period. Water vapour fluctuations affecting flux measurements with LI-7000 were compensated for (Webb et al., 1980); a corresponding compensation is not necessary for temperature (Rannik et al., 1997). The fluxes were corrected for systematic losses due to block averaging and attenuation of the highest frequencies in the cospectra between vertical wind speed and mixing ratio; the transfer function method of Moore (1986) was used for this. For the high-frequency losses, the transfer functions describing the flux attenuation were determined separately for CO₂ and water vapour fluxes (with half-power frequencies of 1.3 and 0.6 Hz, respectively) using temperature cospectra as the reference. These functions were convoluted with generic cospectral distributions (Kaimal and Finnigan, 1994) to calculate the flux correction as a function of wind speed and atmospheric stability.

The 30 min averaged data were screened according to the following acceptance criteria: relative stationarity (Foken and Wichura, 1996) < 100%, internal LI-7000 pressure > 60 kPa, CO₂ mixing ratio > 350 ppm, wind direction within 80–315°, number of spikes in the vertical wind speed and CO₂ concentration data < 150 of 18000. Also, the periods of weak turbulence were discarded by applying a friction velocity (u^*) limit of 0.125 m s⁻¹ (Fig. S1). In addition, the footprint accumulated within the target area shown in Fig. 1 was required to exceed a limit of 0.75 to ensure that the measured flux originated predominantly from the clear-cut site. The footprints were calculated using the model developed by Kormann and Meixner (2001) and input data measured with the sonic anemometer.

In addition to the vertical turbulent fluxes, the CO₂ fluxes associated with the storage of CO₂ below the measurement height were calculated from the change in the mean CO₂ concentration profile between consecutive half-hour periods. It was assumed that the CO₂ concentration in the air column from ground level to the measurement height of 2.75 m was constant.

There were only two periods when a measurement gap in the data was longer than five days: 30 September – 5 October 2016 and 28 April – 4 May 2017. After applying all the data filters described above, 30% of the 32635 half-hour periods recorded within the measurement period were accepted for further analysis (Table S1, Fig. S2). The gap-filling procedure and uncertainty analysis of the net ecosystem exchange of CO₂ (NEE) are described in Appendix.

2.4 Surface energy balance and Bowen ratio

The surface energy balance can be expressed as

$$Q_H + Q_E = Q_N - Q_G - Q_S, \quad (1)$$

where Q_H is sensible heat flux, Q_E is latent heat flux, Q_N is net radiation, Q_G is the ground heat flux, and Q_S is the sum of storage fluxes from other energy sinks and sources. In this study, we assumed that $Q_S = 0$.

The Bowen ratio, defined as

$$\beta = \frac{Q_H}{Q_E}, \quad (2)$$

is used to describe the partitioning of $Q_N - Q_G$ to Q_H and Q_E at the surface. When $\beta < 1$, more available energy at the surface is released to the atmosphere as latent heat than as sensible heat, and vice versa when $\beta > 1$. In this study, we present monthly Bowen ratios that were calculated from the monthly mean daytime (10:00–16:00, UTC+2) fluxes of Q_H and Q_E . Gap-filling of the energy fluxes is described in Appendix.

2.5 Chamber measurements of GHG fluxes

For manual chamber measurements, sampling transects were set up within the clear-cut and control sites, where the GHG fluxes were measured between 29 June 2015 and 29 August 2017, mostly during the snow-free periods. The measurement interval varied between one week and one month, but there were longer gaps in autumn 2015 and spring 2016. The transect had two flux measurement plots at a distance of 4, 8, 12 and 22.5 m from the ditch, and at each distance there was an automatic WTL logger close to the flux measurement plots (see Sect. 2.7 for details). In addition, all the flux plots included a soil temperature data logger (iButton DS1921G, Maxim Integrated Products) at 5 cm depth and two of them (8 and 22.5 m from the ditch) also had a similar logger at 30 cm depth. Before starting the measurements, 2 cm deep grooves were carved into the soil surface for the chambers, and the grooves were occasionally renewed when necessary to keep the chamber sealing adequate. It should be noted that, even though logging residues were left at the site, the measurement plots did not have any above-ground residues. The fluxes were measured using a closed-chamber system with an opaque cylindrical chamber (height 30.5 cm, diameter 31.5 cm) including a mixing fan. The measurements were made in two different ways: using (1) a portable analyser and (2) a stationary analyser. As a portable analyser, we employed a Gaset DX4015 (Gaset Technologies Oy, Helsinki, Finland), based on Fourier transform infrared spectroscopy, to measure CO_2 , CH_4 and N_2O mixing ratios every 5 s. At the clear-cut site, only the portable gas analyser was used. In this setup, the sampled air was circulated in a loop between the gas analyser and the chamber, and the closure time was 10–11 min. At the control site, the portable analyser was used in 2015 for all the gases, but from 2016 onwards the analyser was used only for measuring the N_2O mixing ratios. Starting from 2016 at the control site, we connected the portable gas analyser in series with a Picarro G1130 cavity ring-down spectroscopy gas analyser (Picarro Inc., Santa Clara, CA, USA) to acquire more precise CO_2 and CH_4 mixing ratios. In this system, the gas analysers were stored in a cabin and connected to the chamber with a 50 m long tubing. The instruments were compared against

each other and observed to result in closely similar flux estimates with the 10–11 min closure time. The error caused by not returning the sampled air back to the chamber was corrected for in the data analysis.

The fluxes were calculated the same way for all the gases (Korkiakoski et al., 2017). In short, both linear and exponential regression models were first fitted to the mixing ratio time series using the least-squares approach. The start and end points of the chamber closure were visually identified from the data for each closure. The first minute of each measurement was discarded to ensure that the sample air was properly mixed inside the chamber. After fitting, the mass flux (F) was calculated as

$$F = \left(\frac{dC(t)}{dt} \right)_{t=0} \frac{MPV}{RTA}, \quad (3)$$

where $\left(\frac{dC(t)}{dt} \right)_{t=0}$ is the mixing ratio change in time determined from a linear or exponential model at the beginning of the closure, M is the molecular mass of CO₂, CH₄ or N₂O (44.01, 16.04 and 44.01 g mol⁻¹, respectively), P is air pressure, R is the universal gas constant (8.314 J mol⁻¹ K⁻¹), T is the mean chamber headspace temperature during closure, and V and A are the volume and the base area of the chamber headspace, respectively. The snow depth and the height of mosses and other vegetation in the chamber headspace volume were taken into account, ignoring the pore space in the soil and snow. However, if the soil surface was frozen, the measurements were not made as it was not possible to properly seal the chamber. Finally, analyser-specific flux limits were determined to choose between the linear and exponential models (Korkiakoski et al., 2017). If the flux calculated with the linear model was smaller than the limit, then this estimate was considered more robust for the noisy data and adopted for the later analysis. These limits for the portable system were 5.6 μg CO₂ m⁻² s⁻¹, 9.7 ng CH₄ m⁻² s⁻¹ and 12.5 ng N₂O m⁻² s⁻¹. We did not define a corresponding limit for the CO₂ flux measured with the Picarro gas analyser at the control site as the fluxes were always sufficiently large for using exponential fitting, but for CH₄ the limit was set to 0.7 ng CH₄ m⁻² s⁻¹.

The CO₂ flux measured with chambers represents forest floor respiration (R_{ff}), which is defined as the sum of heterotrophic and autotrophic respiration. As clearcutting affects soil processes, it should be noted that the R_{ff} before and after clearcutting consist of different components. Before the clearcutting (the 2015 data), R_{ff} includes ground vegetation and living roots of trees and ground vegetation, but after clearcutting (2016–2017) it instead includes the survived and regrown ground vegetation and living and dead roots.

The summertime sum of R_{ff} was calculated for each summer separately by using Eq. A3 with hourly soil temperature at 5 cm depth. The uncertainty of this sum was estimated from the minimum and maximum sums based on the standard errors of model parameters (R_0 and E). The minimum and maximum sums were slightly asymmetrical around the mean sum, so the largest deviation from the mean sum was adopted as a conservative uncertainty estimate.

30 2.76 Water table level measurements and analysis

The water table levels within both the clear-cut and non-managed control site were measured for one year before (2015) and two years after the clearcutting (2016–2017). Four automatic monitoring plots consisting of dipwells (perforated plastic tubes

120 cm long, 3.5 cm diameter) were set up at the centre of each site (Fig. 1), located at a distance of 4, 8, 12 and 22.5 m in a transect perpendicular to the ditch (ditch spacing was 45 m). In addition, in order to calibrate the automatic water table measurement data, manually monitored dipwells were installed close to the automatically monitored dipwells within the clear-cut and control sites. WTL was measured manually at weekly or fortnightly intervals during March–November. From the automated dipwells, WTL was recorded with automatic probes (TruTrack WT-HR -logger, Intech Instruments Ltd, Auckland, New Zealand; Odyssey Capacitance Water Level Logger, Dataflow Systems Limited, Christchurch, New Zealand) at hourly intervals. The recorded values were then calibrated with linear regression using the manually measured WTL data from both the control and clear-cut site.

2.6.7 Statistical analysis

Linear mixed effect model was used for testing for differences in the daily mean CO₂, CH₄ and N₂O fluxes between the clear-cut and control sites and between the years in the chamber measurement data. In both cases, the chamber plots were treated as a random effect. The fixed effects of the model were the site type (clear-cut or control) and the measurement year when comparing the measurements made at different sites and in different years, respectively. The linear mixed effect model was carried out with the R programming language (R Core Team, 2018, version 3.5.0) using the ‘lme4’ package. The normality of model residuals was visually checked using quantile-quantile plots. The differences were tested with Tukey’s HSD post hoc test.

The analysis of the clearcutting effects on WTL was based on the paired treatment approach (also called calibration period – control area method) (e.g., Kaila et al., 2014; Laurén et al., 2009). We first calculated linear regressions between the WTL within the control and clear-cut sites for the pre-treatment period using the WTL logger data from 2015. Then we used this regression model and the post-treatment WTL data from the control site to predict WTL for the clear-cut site as if it had not been harvested. The clear-cut effect was calculated as the difference between the calibrated post-clear-cut WTL measurements and the predicted background WTL values in the clear-cut site after the harvest.

2.7 Water table level measurements and analysis

~~The water table levels within both the clear-cut and non-managed control site were measured for one year before (2015) and two years after the clearcutting (2016–2017). Four automatic monitoring plots consisting of dipwells (perforated plastic tubes 120 cm long, 3.5 cm diameter) were set up at the centre of each site (Fig. 1), located at a distance of 4, 8, 12 and 22.5 m in a transect perpendicular to the ditch (ditch spacing was 45 m). In addition, in order to calibrate the automatic water table measurement data, manually monitored dipwells were installed close to the automatically monitored dipwells within the clear-cut and control sites. WTL was measured manually at weekly or fortnightly intervals during March–November. From the automated dipwells, WTL was recorded with automatic probes (TruTrack WT-HR -logger, Intech Instruments Ltd, Auckland, New Zealand; Odyssey Capacitance Water Level Logger, Dataflow Systems Limited, Christchurch, New Zealand) at hourly~~

~~intervals. The recorded values were then calibrated with linear regression using the manually measured WTL data from both the control and clear cut site.~~

~~The analysis of the clearcutting effects on WTL was based on the paired treatment approach (also called calibration period—control area method) (e.g., Kaila et al., 2014; Laurén et al., 2009). We first calculated linear regressions between the WTL within the control and clear cut sites for the pre-treatment period using the WTL logger data from 2015. Then we used this regression model and the post-treatment WTL data from the control site to predict WTL for the clear cut site as if it had not been harvested. The clear cut effect was calculated as the difference between the calibrated post clear cut WTL measurements and the predicted background WTL values in the clear cut site after the harvest.~~

3 Results

10 3.1 Meteorological and hydrological conditions

The long-term (1981–2010) mean annual, winter (DJF), and summer (JJA) air temperatures at the nearby (Jokioinen, 35 km northwest of Lettosuo) weather station were 4.6, –5.3, and 15.2 °C, respectively (Pirinen et al., 2012). The mean annual temperature before the clearcutting in 2015 at Jokioinen was 6.2 °C. Also, the winter (2015–2016) before the clearcutting was warmer (–3.4 °C) while the summer 2015 was colder (14.4 °C) than the long-term mean. The mean post-clearcut annual air temperatures at the EC site during the first and second EC measurement year were 5.6 °C and 4.4 °C, respectively (Fig. 2a); which this difference was reflected in soil temperatures (Fig. 2b). These air temperatures were higher and about the same, respectively, than the long-term (1981–2010) average at Jokioinen (4.6 °C; Pirinen et al., 2012). All Both post-clearcut winters (DJF) were warmer (2015–2016: –3.4 °C (Jokioinen), 2016–2017: –3.0 °C, 2017–2018: –3.4 °C) than the long-term average (–5.3 °C). However, the mean summer (JJA) temperature at Jokioinen in 2015 (14.4 °C) was 0.8 °C cooler than the long-term average (15.2 °C), while the JJA temperature at the EC site in 2016 was similar to (15.3 °C) and in 2017 cooler (by 1.6 °C) than in 1981–2010. In addition, compared to the precipitation at Jokioinen (long-term average 627 mm), the first EC measurement year was drier (502 mm) and the second year similar (656 mm) (Fig. 2c) while the pre-clearcut year of 2015 was wetter (680 mm) than both post-clearcut years. Especially the autumn (SON) 2016 and winter 2016–2017 were much drier, while the springs and summers were quite similar to the long-term average conditions.

Both winters had a shallower snow cover than the long-term average annual maximum (28 cm; Pirinen et al., 2012). The maximum snow depths at Jokioinen were 12 cm and 22 cm in 2016–2017 and 2017–2018, respectively. Due to the especially shallow snow cover in the first winter and the high temperatures in March, all the snow had melted by 23 March 2017 (Fig. 2d). However, in 2018, the snow cover was still intact when the measurement period ended in 7 April.

The pre-harvest soil temperatures at 30 cm depth were similar within the clearcutting and control sites, but clearcutting increased their temporal variation (Fig. S3). After clearcutting in late spring, the daily mean 30 cm soil temperature rose faster and was higher throughout the summer within the clearcut than at the control site. Also, the 30 cm temperature within the clearcut started to decline earlier in autumn and was lower during winter within the clearcut than the control site. The monthly

mean diel variation of soil temperature at 5 cm depth ranged from 0.8 to 1.3 °C in July–September 2015 before clearcutting (Fig. S4). After clearcutting, this variation increased to 2.3–2.9 °C and 2.1–3.7 °C in 2016 and 2017, respectively. On average the diel variation was 1.8 °C larger in July–September after the clearcutting than before it (Figs. 2b and S4). Increases in the diel temperature range were also observed at the 10 cm (1.2 °C) and 20 cm (0.33 °C) depths, but not anymore at 30 cm.

5 The mean WTL in July–August 2015 was –33 cm within the clear-cut area, while at the control site it was –50 cm (Fig. 3). After clearcutting, the WTL rose and remained continuously at a markedly higher level than the predicted background WTL and the measured WTL at the control site. At the clear-cut site, the WTL rose to –22 cm and –27 cm in July–August 2016 and 2017, respectively. On the other hand, the WTL at the control site sank to –55 cm during these months in both post-harvest years. The largest post-treatment rise occurred in July–October when the average WTL was –23 cm and –24 cm in the clear-cut and
10 –46 cm and –49 cm in the background model, corresponding to a 23 cm and 25 cm rise in WTL in 2016 and 2017, respectively (Fig. 3). As a consequence of heavy rain episodes in summer and autumn in 2017, the amplitude of WTL variations was larger in this than the previous year.

3.2 CO₂ exchange on an ecosystem level

3.2.1 Seasonal and diel variations

15 When the measurements started at the beginning of April 2016, the daily mean NEE was mainly < 0.08 mg CO₂ m⁻² s⁻¹ but started to increase with temperature at the end of April, after the daily mean temperature exceeded 5 °C (Figs. 2a and 4). There was a substantial temporary drop in NEE in the first half of June after a cold spell when the daily mean temperature decreased from 20 to 10 °C. By the end of June, NEE stabilized at around 0.21 mg CO₂ m⁻² s⁻¹. From the end of August, NEE decreased until reaching a stable level of 0.03 mg CO₂ m⁻² s⁻¹ in January 2017. At the end of March 2017, NEE started to gradually
20 increase, and in mid-May it quickly rose to 0.08 mg CO₂ m⁻² s⁻¹. After this, NEE continued to increase, stabilizing at around 0.13 mg CO₂ m⁻² s⁻¹ in August and then decreasing from the end of September onwards.

Even though all the big trees were removed in clearcutting and most of the ground vegetation was destroyed, some small understorey trees were left at the site and weak photosynthesis could be observed simultaneously with the increased respiration in the end of April 2016 (Fig. 4). In May, the estimated magnitude of gross primary production (|GPP|, Appendix A) increased
25 to 0.05 mg CO₂ m⁻² s⁻¹, which was about 25% of the respiration rate at that time. However, |GPP| decreased to 0.02 mg CO₂ m⁻² s⁻¹ in June, after which it started to increase again, reaching its maximum at 0.08 g CO₂ m⁻² s⁻¹ at the end of July and the beginning of August. From that point on, photosynthesis started to weaken, marking the ending of the growing season, and ceased in mid-October, about a month before the first snow. In 2017, |GPP| increased from mid-May until July, when it stabilized at around 0.11 mg CO₂ m⁻² s⁻¹. Similarly to 2016, |GPP| decreased from August onwards.

30 The correlation between the night-time NEE (i.e., R_{eco} in Eq. A3) and the 5 cm soil temperature was stronger in summer 2017 than 2016 (Fig. 5a). The model fits (Eq. A3) indicate that the temperature response of ecosystem respiration (parameter E) was also stronger in 2017. However, the base respiration rate R_0 (R_{eco} at 10 °C) was larger in the first than the second summer

(0.199 vs. 0.173 mg CO₂ m⁻² s⁻¹). In addition, no correlation was found between the model residuals and WTL in either summer (Fig. S5). Like the temperature response of R_{eco} , the light response of GPP was stronger in summer 2017 than 2016 (Fig. 5b). Also, the corresponding maximum gross photosynthesis rate (GP_{max} , Eq. A2) more than doubled (-0.226 vs. -0.100 mg CO₂ m⁻² s⁻¹).

5 The site was on average a CO₂ source throughout the day, also during the summer. It should be noted, however, that there were still multiple half-hour periods, especially in August 2017, when the site acted as a CO₂ sink. A noticeable diel variation in NEE was observed mainly from April to October (Fig. 6). Between April and August, the mean diel NEE cycle had a minimum during the morning hours (5:00–10:00, UTC+2) and the highest emissions during the evening/night (21:00–1:00). After August, the lowest emissions took place later, at around noon. On the other hand, the highest emissions were observed earlier, at 20:00 in September and at 18:00 in October. In October, the diel cycle of NEE was still noticeable; it vanished in November and reappeared in March 2017. In spite of the different mean fluxes, the amplitudes of their diel cycle were rather similar in 2016 and 2017, except from July to September when the amplitudes were much larger in 2017. Also, unlike in 2016, systematic diel variation was still obvious in November 2017. Even though such variation indicates significant photosynthesis during the midday hours, the monthly mean diel NEE cycle consistently showed positive fluxes.

15 3.2.2 CO₂ balances

The annual CO₂ balances of the first and the second year after clearcutting were 3086 ± 148 g CO₂ m⁻² (\pm uncertainty; see Appendix) and 2072 ± 124 g CO₂ m⁻², respectively. About half of the annual CO₂ emissions during both EC measurement years took place during the summer months (JJA), totalling 1558 ± 99 g CO₂ m⁻² (Table 2, Fig. 7), and 73% (2256 ± 146 g CO₂ m⁻²) of the annual total was accumulated during the May–September period. In 2017, however, these emissions decreased to 915 ± 79 g CO₂ m⁻² (June–August) and 1401 ± 109 g CO₂ m⁻² (May–September), i.e., by 41% and 38%, respectively. The effect of storage fluxes on the CO₂ balance was negligible as the daily mean storage flux typically varied within ± 0.009 mg CO₂ m⁻² s⁻¹ during summer and within ± 0.004 mg CO₂ m⁻² s⁻¹ during the other seasons.

The most significant period of CO₂ uptake was from June to September (Fig. 7). Considering only the summer months (JJA), the integrated GPP was 389 and 761 g CO₂ m⁻² in 2016 and 2017, respectively (Table 2), i.e., the mean CO₂ uptake in the second summer after clearcutting increased by 96% from the first summer. On the other hand, the total summertime R_{eco} decreased by 14%, from 1928 g CO₂ m⁻² in 2016 to 1652 g CO₂ m⁻² in 2017 (Table 2).

3.3 Soil GHG fluxes

The summertime (JJA) R_{ff} measured at the soon-to-be clear-cut site in 2015 was 1611 ± 191 g CO₂ m⁻² (\pm uncertainty; see Sect. 2.5) and the measured CO₂ fluxes varied from 0.04 to 0.36 g CO₂ m⁻² s⁻¹ (Fig. 8a). The CH₄ flux averaged over all the measurement plots was -26 ± 5 ng CH₄ m⁻² s⁻¹ before the clearcutting. All the measured CH₄ fluxes were negative (Fig. 8b) ranging from -103 to -2 ng CH₄ m⁻² s⁻¹. N₂O fluxes varied mostly from -17 to 33 ng N₂O m⁻² s⁻¹ and averaged at 1 ± 5 ng N₂O m⁻² s⁻¹ (Fig. 8c). At the control site, the summertime fluxes of CO₂ (from 0.03 to 0.39 g CO₂ m⁻² s⁻¹) and N₂O (mean: 11

$\pm 9 \text{ ng N}_2\text{O m}^{-2} \text{ s}^{-1}$) in 2015 were not significantly different from those at the soon-to-be clear-cut site. However, the CH_4 fluxes (mean: $-10 \pm 3 \text{ ng CH}_4 \text{ m}^{-2} \text{ s}^{-1}$) at the control site were significantly ($p < 0.02$) larger than at the soon-to-be clear-cut site. After the clearcutting, the total summertime R_{ff} decreased to $980 \pm 299 \text{ g CO}_2 \text{ m}^{-2}$ in summer 2016 and remained at the same level ($1047 \pm 113 \text{ g CO}_2 \text{ m}^{-2}$) in 2017. These numbers were 51% and 63% of the R_{eco} for summers 2016 and 2017, respectively (Fig. 9). Also, the temperature response of R_{ff} got weaker and the R_0 parameter (R_{ff} at $10 \text{ }^\circ\text{C}$) decreased while at the control site the responses remained quite similar over the years (Fig. 10). The strength of temperature response and R_0 partly recovered in 2017 but were still weaker than before the clearcutting. In contrast to the R_{ff} at the clear-cut, the summertime R_{ff} at the control site increased from $1852 \pm 1353 \text{ g CO}_2 \text{ m}^{-2}$ in 2015 to $2280 \pm 346 \text{ g CO}_2 \text{ m}^{-2}$ and $2438 \pm 225 \text{ g CO}_2 \text{ m}^{-2}$ in 2016 and 2017, respectively. However, the increase was significant only when comparing the summers 2015 and 2017 ($p < 0.05$). The clear-cut and control sites were significantly different from each other in both summers ($p < 0.001$) after clearcutting.

CH_4 fluxes changed markedly after the clearcutting as the previously small mean CH_4 sink turned into a small CH_4 source in the first ($4 \pm 3 \text{ ng CH}_4 \text{ m}^{-2} \text{ s}^{-1}$) and second ($6 \pm 2 \text{ ng CH}_4 \text{ m}^{-2} \text{ s}^{-1}$) year after the clearcutting (Fig. 8b); the changes were significant for both years ($p < 0.001$). However, the post-clearcut years were not significantly different from each other ($p = 0.85$). The post-clearcut fluxes varied mostly from -14 to $28 \text{ ng CH}_4 \text{ m}^{-2} \text{ s}^{-1}$ with some occasional emission peaks reaching up to $140 \text{ ng CH}_4 \text{ m}^{-2} \text{ s}^{-1}$. Most of the measured fluxes were positive (emission), even though a single measurement plot could act both as a source and a sink on different measurement days. The control site remained as a CH_4 sink in 2016 and 2017 and neither of the mean summer fluxes was significantly different from the pre-harvest summer mean. However, the mean CH_4 fluxes were significantly different ($p < 0.001$) between the sites in both post-harvest summers.

Similarly to CH_4 , the post-clearcut N_2O fluxes were significantly higher than the pre-clearcut ones ($p < 0.001$), and the post-clearcut years did not differ significantly from each other. After the clearcutting, the mean annual fluxes were 228 ± 26 and $212 \pm 21 \text{ ng N}_2\text{O m}^{-2} \text{ s}^{-1}$ in 2016 and 2017, respectively (Fig. 8c). All the measurement plots turned to large N_2O sources, and the largest measured emission was $1339 \text{ ng N}_2\text{O m}^{-2} \text{ s}^{-1}$. All the measured fluxes were positive, except for a plot furthest from the ditch that acted as a temporary N_2O sink in 2017, especially in the June–July period. The N_2O fluxes at the control site remained low in summer 2016 (mean: $16 \pm 11 \text{ ng N}_2\text{O m}^{-2} \text{ s}^{-1}$), but increased in summer 2017 (mean: $70 \pm 10 \text{ ng N}_2\text{O m}^{-2} \text{ s}^{-1}$). However, the post-harvest emissions at the control site were significantly lower than those at the clear-cut site ($p < 0.001$). Both the spatial and temporal variation were large: the largest emissions occurred in June while from July onwards the emissions decreased (Fig. 8c).

3.4 Energy fluxes

The daily sums of sensible heat (Q_{H}) were about $1 \text{ MJ m}^{-2} \text{ day}^{-1}$ on average at the start of the EC measurement period in April 2016 and increased to a maximum of $4.7 \text{ MJ m}^{-2} \text{ day}^{-1}$ in June (Fig. 11). In July, the daily Q_{H} sum was already decreasing with the decreasing net radiation (Q_{N}), and it turned negative in October. On the other hand, the daily sums of latent heat flux (Q_{E}) were quite stable from April until August averaging at $1.0 \pm 0.04 \text{ MJ m}^{-2} \text{ day}^{-1}$ (\pm standard error of mean). In September, Q_{E}

started to decrease and reached zero at the end of October. The ground heat flux (Q_G) varied within 0–0.5 MJ m⁻² day⁻¹ from April to August, after which it varied mostly between –0.5 and 0 MJ m⁻² day⁻¹ until the end of April 2017.

In 2017, the seasonal dynamics of the energy fluxes were similar to 2016, but the magnitude of fluxes changed considerably. The summertime average daily sum of Q_H decreased from 1.8 to 1.2 MJ m⁻² day⁻¹, while that of Q_E increased from 1.3 to 2.3 MJ m⁻² day⁻¹. Also, the monthly sums of these fluxes were markedly different in the second year after the clearcutting. During the period of the highest fluxes (April–September), the Q_H sum was 33% smaller in 2017 than 2016, while the Q_E and Q_N sums were higher by 40% and 13%, respectively. The changes in Q_G were minor.

The monthly mean midday Q_H in May and June 2016 were 206 W m⁻² and 165 W m⁻², respectively (Fig. 12a). After June, the midday Q_H started to decrease with decreasing Q_N , and the night-time Q_H turned from zero to slightly negative. The midday Q_E also increased from April to June, reaching 88 W m⁻², and remained quite stable until September. The resulting monthly mean daytime Bowen ratio (β), increased from 1.7 in April to 2.6 in May 2016 and then declined to 0.8 in August. In 2017, the mean midday Q_H was similar to that in 2016, except in July and September when the fluxes were 34% and 23% lower, respectively, than in 2016 (Fig. 12b). However, the daily maximum Q_E was either similar to or higher (up to 48%) than Q_H in all months except May 2017. Correspondingly, β was lower than one (0.6–1.0) during the period of high fluxes in 2017, except in May when it was 1.7.

4 Discussion

Dynamics of the CO₂ fluxes and flux components

The study site was a large source of CO₂ during the two EC measurement years after clearcutting, but the emissions were 33% smaller in the second than the first year. |GPP| remained low and stable during the first summer after clearcutting, reflecting the fact that neither vascular ground vegetation nor moss cover had developed substantially by that time. However, during the second summer, the total |GPP| increased by 96% from the first post-clearcut summer, and the increase in the coverage of ground vegetation was already noticeable (although not directly measured). This relatively fast recovery of ground vegetation is similar to the leaf area index changes observed by Mäkiranta et al. (2010) after clearcutting in a nutrient-poor peatland forest. R_{eco} was larger in the first than in the second post-harvest year which could be due to the first year being warmer or due to changes in the respiration of either residues or forest floor in the second year. The chamber measurements, which were not affected by aboveground logging residues, showed a 39% and 35% decrease in R_{ff} after clearcutting in the first and second summer after clearcutting, respectively. Possible reasons for this decrease include reduced peat decomposition due to raised WTL, ceased autotrophic respiration of (the harvested) trees roots and decreased autotrophic respiration of ground vegetation, even though the latter slightly recovered during the summer as the ground vegetation started to regrow. The CO₂ flux from the decomposing belowground logging residues (roots) was not sufficiently large to compensate for this decrease in R_{ff} . In summer 2016, R_{ff} was 51% of the R_{eco} suggesting that half of the R_{eco} during that time originated from the aboveground logging residues, which were not included in the chamber measurements of R_{ff} . In summer 2017, however, R_{eco} decreased by 14%, but

R_{ff} remained about the same, indicating that a larger proportion (63%) of R_{eco} originated from the forest floor respiration than in the previous summer. This could be due to the decrease in decomposition rate of the aboveground logging residues due to lower temperatures and because the residues had already partly decomposed during the previous summer. However, the lower temperatures would also decrease R_{ff} , but this did not happen because in 2017 WTL was 5 cm lower, which enhanced peat decomposition. In addition, the NEE in summer 2017 was approximately equal to R_{ff} , which means that the increase in |GPP| was sufficient to balance the CO₂ emissions from the logging residues, but not from soil and plants.

After the removal of the canopy, more solar radiation reached the ground surface, heating it more during the daytime, while the heat transfer from the soil to the atmosphere was enhanced during the night; together these resulted in a higher diel variation in soil temperature. On the other hand, the insulating impact of logging residues may have partly dampened the amplitude of the diel temperature cycle, as observed by (Ojanen et al., 2017). The higher soil temperatures during the daytime may have enhanced peat respiration in the layer closest to the surface. However, it has been shown in previous studies that the temperature response of respiration weakens when the soil gets dry, often resulting in lower respiration rates (Mäkiranta et al., 2009, 2010). Such weakening of the temperature response was also observed at Lettosuo especially in the first summer after the harvest when the almost bare soil surface was exposed to more solar radiation after clearcutting. Also, the summer 2016 was warmer and dryer than on average, which probably enhanced this effect. Moreover, the addition of logging residues may have increased R_{eco} substantially, as demonstrated by Mäkiranta et al. (2012) who observed that the plots with logging residues had a CO₂ efflux twice as high as the plots without them. However, a later study by Ojanen et al. (2017) did not show a consistent increase in soil organic matter decomposition rates at similar sites. In previous studies, a decrease has been observed in both R_{eco} (Kowalski et al., 2004, 2003; Takagi et al., 2009) and R_{ff} (Mäkiranta et al., 2010; Takagi et al., 2009; Zerva and Mencuccini, 2005) after clearcutting in peatland and upland forests. However, also increasing R_{eco} and R_{ff} have been reported (Kowalski et al., 2004; Londo et al., 1999). The differences between these results are most likely caused by a different effect of harvest on soil microbial activity, which is mostly controlled by soil temperature, moisture and nutrient availability (Davidson et al., 1998; Fontaine et al., 2004, 2007; Lloyd and Taylor, 1994). Other possible reasons for this variability include differences in, for example, root respiration, the type and amount of logging residue, and WTL, especially in peatland forests.

In peat soils, the decomposition of organic matter is mainly controlled by soil temperature and WTL (Blodau et al., 2004; Mäkiranta et al., 2009; Silvola et al., 1996), but soil temperature is typically the most influential factor explaining the temporal dynamics in R_{ff} (e.g. Mäkiranta et al., 2008; Ojanen et al., 2010). At Lettosuo, the correlation between the half-hourly R_{eco} and soil temperature was mostly weak. This could be due to logging residues as their decomposition rate probably does not depend as much on soil temperature as on the temperature and moisture of the logging residues themselves. At times, especially when the temperature range was larger (in May and June), a significant relationship (correlation coefficient peaking at 0.57) was observed, and there was also an obvious temperature response that followed the annual cycle. Also, when WTL is closer (> – 15 cm) to the surface, it is usually a significant predictor of R_{ff} in peat soils (Chimner and Cooper, 2003; Riutta et al., 2007). At Lettosuo after clearcutting, WTL varied mostly within 20–30 cm below the soil surface, and there was no correlation between R_{eco} and WTL after removing the effect of soil temperature on R_{eco} (Fig. S5). Because R_{eco} combines many respiration

components in addition to R_{ff} , it is likely that the high CO_2 efflux from logging residues, for example, masks the possible WTL effect on R_{ff} .

Our results show that after clearcutting a peatland forest is a large source of CO_2 to the atmosphere. Nutrient-rich drained peatland forests with a growing tree stand have generally been found to be CO_2 neutral or small sinks of CO_2 (Meyer et al., 2013; Ojanen et al., 2013; Uri et al., 2017) as the carbon accumulated by trees typically balances the carbon emissions caused by the decomposing peat layer. This suggests that also Lettosuo probably was close to CO_2 neutral state before the clearcutting. Thus the impact of clearcutting from CO_2 neutral to a source of $3086 \pm 148 \text{ g CO}_2 \text{ m}^{-2} \text{ yr}^{-1}$ in the first year after clearcutting must be considered substantial. This also means that the site may need to be considered a net CO_2 source in terms of the long-term balance even if the forest had been a CO_2 sink at a particular growth stage (Hommeltenberg et al., 2014), depending on how long the post-clearcut emissions continue, even at the forest had been a CO_2 sink at a certain growth stage, due to these emissions it may still need to be considered a net CO_2 source in terms of the long term balance (Hommeltenberg et al., 2014). In this study, however, we only focus on the immediate impact of clearcutting.

Mäkiranta et al. (2010) measured CO_2 exchange in a clear-cut nutrient-poor peatland forest with soil chambers during a growing season (May–October) and reported slightly smaller NEE ($1990 \text{ g CO}_2 \text{ m}^{-2} \text{ season}^{-1}$) than the NEE we measured at Lettosuo during a similar timeframe ($2340 \text{ g CO}_2 \text{ m}^{-2} \text{ season}^{-1}$). Among mineral soil forests, on the other hand, the annual CO_2 exchange measured with the EC method right after the clearcutting varies a lot; the annual balances ranging from ca. 400 to $4700 \text{ g CO}_2 \text{ m}^{-2} \text{ yr}^{-1}$ (Amiro et al., 2010; Clark et al., 2004; Humphreys et al., 2005; Kowalski et al., 2004, 2003; Takagi et al., 2009; Williams et al., 2013). If considering only mineral soil forests, the forests in warmer climates typically emit more carbon after the harvest, but they also recover faster than the boreal forests (Amiro et al., 2010). Compared to the studies cited above, the clear-cut site at Lettosuo was the second largest source of CO_2 immediately after clearcutting, being second only to the Slash pine plantation in Florida (Clark et al., 2004). The high CO_2 emission at Lettosuo as compared to other clearcut sites on mineral soil is apparently due to the decomposition of the oxic peat layer in this nutrient-rich forest, which are known to act as CO_2 sources up to $1000 \text{ g CO}_2 \text{ m}^{-2} \text{ yr}^{-1}$ (Ojanen et al., 2013; Uri et al., 2017). Mineral soil forests do not have a peat layer or any other carbon storage that could cause CO_2 emissions of such magnitude.

25 Soil CH_4 and N_2O fluxes

The CH_4 flux in peat soils is mostly controlled by WTL (Martikainen et al., 1995; Roulet et al., 1993), which divides the peat column into anoxic and oxic layers in which CH_4 production and oxidation occur, respectively. After clearcutting, the WTL rose from -33 cm in 2015 to -22 cm in 2016 during July-August, and the highest estimated rise in WTL occurred in July-October when clearcutting was estimated to raise WTL by 23 cm on average when compared to the background model. This made the top-soil oxic layer much thinner and the conditions less favourable for CH_4 oxidation, allowing a smaller amount of the CH_4 produced in the anoxic layer to be oxidized before reaching the atmosphere. The WTL rise was enough to turn the clear-cut site from a CH_4 sink into a small CH_4 source while the control site remained a consistent CH_4 sink. A similar turn of a clear-cut site from a CH_4 sink to a source has previously been found for both peat (Zerva and Mencuccini, 2005) and mineral

soils (Sundqvist et al., 2014). However, Huttunen et al. (2003) measured mostly net CH₄ uptake in two drained peatland forest sites in southern Finland also after clearcutting, and the difference between the control and clear-cut sites was not statistically significant in either case.

The N₂O fluxes were highly variable both before and after clearcutting. However, there was a strong increase from the mean pre-clearcut average flux of 1 ± 5 ng N₂O m⁻² s⁻¹ to the post-clearcut flux of 228 ± 26 ng N₂O m⁻² s⁻¹. On the other hand, the emissions were small at the control site in 2016, but increased in 2017, especially in June, even though the increase was markedly smaller than at the clear-cut site. This is probably due to higher precipitation in 2017, which increased soil moisture and enhanced denitrification. Increases of N₂O emissions after clearcutting have also been observed in peat and mineral soil forests by Huttunen et al. (2003) and Saari et al. (2009), but the fluxes at these sites were an order of magnitude smaller than at Lettosuo. It should be noted that the N₂O measurements in this study do not include any above-ground logging residues, which are expected to raise N₂O emissions further; Mäkiranta et al. (2012) found about three times as large N₂O emissions from the plots with logging residues as from the plots without them.

Compared to the change in NEE (i.e. CO₂ fluxes) at the site, the change in CH₄ emissions is negligible when considering its climatic impact in terms of the global warming potential over 100 yr (GWP₁₀₀ = 34; IPCC, 2013). Even though the change in N₂O fluxes to the present level at our site was large, the climatic effect of N₂O was only about 10% of that due to the even larger change in NEE, when considering GWP over a period of up to 100 years (GWP₁₀₀ = 298; IPCC, 2013).

Energy fluxes

The mean daily Q_E at Lettosuo during the spring and summer right after the clearcutting varied mostly within 0.5–2.5 MJ m⁻² day⁻¹ and within 1.0–4.0 MJ m⁻² day⁻¹ during the second spring and summer. Compared to the values of 2–10 MJ m⁻² day⁻¹ measured in an upland forest in Western Russia (Mamkin et al., 2016), our numbers are low. In the case of Q_H , the fluxes at Lettosuo were systematically lower than those reported by Mamkin et al. (2016), but the difference was not as large as for Q_E . At Lettosuo, β was lower in the second summer after the clearcutting as compared to the first summer; in summer 2016 β was >1 most of the time, whereas in summer 2017 it was <1 for the whole summer. Thus the pattern was somewhat different from the results of Mamkin et al. (2016) who reported that $\beta < 1$ already throughout the first summer after the clearcutting. One reason for the smaller Q_E at Lettosuo could be the drier-than-normal summers in 2016 and 2017 and the drier surface layer of the peat.

In summer 2017 at Lettosuo, Q_H decreased slightly and Q_E almost doubled compared to 2016, suggesting a recovery of transpiration due to the gradual recovery of ground vegetation. This is in agreement with the results obtained by Williams et al. (2013) for a clear-cut Norway spruce plantation in the north-east USA, where a recovery of evapotranspiration and a decrease in Q_H were observed across the years after the clearcutting. Moreover, the recovered ground vegetation at Lettosuo most likely increased albedo and thus prevented the soil from heating as much as in 2016, resulting in a smaller Q_H over the whole summer. The clear-cut site at Lettosuo is small, but a larger scale clearcutting would probably affect local and regional

climate due to changes in surface energy balance fluxes. This is because Q_H and Q_E affect the properties and growth rate of the planetary boundary layer, influencing convection and the long-range transport of heat and humidity.

5 Conclusions

Based on our measurements, we conclude that after clearcutting a nutrient-rich peatland forest is a large CO₂ source. This is due both to the decomposition of peat and logging residues and to the reduction of gross primary production as a result of the removal of photosynthesizing trees and the decline of ground vegetation and understorey. Removing the trees decreased transpiration, which caused the WTL to rise by 23 cm compared to the background model during July-October, which in turn likely decreased the peat decomposition rate due to the decreased volume of aerobic peat. Plant respiration also decreased as the plants were removed or destroyed. On the other hand, decomposition of logging residues increased CO₂ emissions from the site; the emissions from the residues were estimated to be 49% of the total ecosystem respiration in the first summer after clearcutting. In the second summer, ground vegetation and its primary production recovered noticeably. On the other hand, the CO₂ emissions from the logging residues decreased as part of the residues had decomposed during the previous summer. In total, these changes reduced the net CO₂ emissions of the site by 41% compared to the first summer.

The soil turned from a small CH₄ sink into a small source after clearcutting due to the WTL rise. However, the radiative forcing related to this change was insignificant compared to that due to the change in NEE. Clearcutting turned the soil into a substantial source of N₂O. This change produced a 100 yr GWP of about 10% of that due to the increased NEE at the site.

The mean daytime latent heat flux almost doubled from the first to the second year after clear-cutting, suggesting that the transpiration of ground vegetation had recovered. The recovered ground vegetation most likely also increased albedo and thus prevented the soil from heating as much as in 2016, which resulted in a smaller sensible heat flux.

Overall, the results of this study show that clearcutting peatland forests exerts a strong climatic warming effect through accelerated emissions of greenhouse gases. However, this study only demonstrates a short-term impact of two years, and more extensive measurements are required to gain knowledge of the long-term effects of clearcutting in peatland forests.

Appendix A: Gap-filling CO₂ data

To calculate seasonal and annual CO₂ balances, full time series are needed and thus any gaps in the CO₂ flux data need to be filled. For this, and to analyse the components of the carbon balance, we partitioned the measured CO₂ flux, i.e., net ecosystem exchange (NEE), into gross primary production (GPP) and total ecosystem respiration (R_{eco}):

$$5 \quad \text{NEE} = \text{GPP} + R_{\text{eco}}. \quad (\text{A1})$$

For gap-filling, we expressed GPP as a function of irradiance (e.g., Aurela et al., 2015):

$$\text{GPP} = \frac{\alpha \times \text{PPFD} \times \text{GP}_{\text{max}}}{\alpha \times \text{PPFD} + \text{GP}_{\text{max}}}, \quad (\text{A2})$$

where PPFD is the photosynthetic photon flux density, α is the initial slope of the NEE response to PPFD, and GP_{max} is the asymptotic gross photosynthesis rate in optimal light conditions. R_{eco} was assumed to follow the Arrhenius-type model described by Lloyd and Taylor (1994):

$$10 \quad R_{\text{eco}} = R_0 \times \exp \left[E \left(\frac{1}{T_0} - \frac{1}{T_{\text{soil}} - T_1} \right) \right], \quad (\text{A3})$$

where R_0 is the ecosystem respiration at 10 °C, E is the temperature sensitivity of the respiration, T_{soil} is the measured soil temperature at 5 cm depth, $T_0 = 56.02$ K and $T_1 = 227.13$ K.

The parameters E and R_0 were defined from the night-time data ($\text{PPFD} < 20 \mu\text{mol m}^{-2} \text{s}^{-1}$) in two parts. First, the parameter E , which was allowed to vary within 200–500 K, was determined with a 15 day moving window for each day, with the minimum number of observations set to 12. If there were not enough observations within a time window, then the window size was increased by one day both in the beginning and the end until enough data were found. The resulting window size varied between 15 and 27 days, but only 22 gap-filled days had a longer window than 15 days. Next, all the E values that hit the allowed boundary values (200 and 500 K) were discarded and filled with 14 day moving medians. Finally, a 15 day moving window, similar to the one in the first part, was used to determine R_0 by using the fixed E values. Also, using the same moving window, GP_{max} and α were determined from the daytime data. However, from 1 November 2016 until 8 March 2017 with no significant CO₂ uptake, a 5 day moving average was used to fill the gaps in the measured NEE.

All the calculations and analyses were made with the Python programming language (Python Software Foundation, version 2.7, <https://www.python.org>). For the fits, the least-squares method was used through the “polyfit” function of NumPy (<http://www.numpy.org/>) library for the linear regression and the “curve_fit” function of SciPy (<http://www.scipy.org/>) library for the nonlinear fits.

Appendix B: Uncertainty analysis of NEE

The CO₂ balance obtained from EC measurements has multiple potential error sources due to instrumental, statistical and methodological uncertainties. We included the most significant, although not all, random error sources. The random error

including the statistical measurement error (E_{meas}) inherent in EC measurements and the error caused by gap filling of missing data (E_{gap}) was estimated as follows (Räsänen et al., 2017):

$$E_{\text{meas/gap}} = \sqrt{\sum_i \frac{(\text{NEE}_{i,\text{obs}} - \text{NEE}_{i,\text{mod}})^2}{n_{\text{obs/mod}}} \sqrt{n_{\text{meas/gap}}}}, \quad (\text{B1})$$

where NEE_{obs} is the 30 min flux that passed all the filtering procedures and NEE_{mod} is the corresponding fitted NEE (Eqs. A1–A3), and $n_{\text{meas/gap}}$ the number of the measured or gap-filled data. This method provides a conservative estimate for E_{meas} (Aurela et al., 2002) and for E_{gap} includes the effect of random variability on the model fits.

The annual systematic error caused by the friction velocity filtering (E_{ustar}) was estimated by recalculating the annual balance with modified data sets that were screened with two different u_* limits (0.075 and 0.175 m s⁻¹). E_{ustar} was calculated as the average difference between the annual balance calculated with the optimal u_* threshold (0.125 m s⁻¹) and with the annual balances calculated with the modified u_* limits. Similarly, two additional different footprint limits (0.65 and 0.85) were adopted to estimate the annual error caused by the footprint filtering (E_{fp}).

The total uncertainty of the annual balance (E_{tot}) was calculated with the standard error propagation principle:

$$E_{\text{tot}} = \sqrt{E_{\text{meas}}^2 + E_{\text{gap}}^2 + E_{\text{ustar}}^2 + E_{\text{fp}}^2}. \quad (\text{B2})$$

Appendix C: Gap-filling energy fluxes

Energy fluxes were gap-filled in several steps following the procedure described by Kowalski et al. (2003). First, the gaps in daytime Q_H ($Q_N > 0$) were filled with monthly linear regressions with net radiation. Next, the night-time ($Q_N < 0$) gaps in Q_H were replaced by the corresponding Q_N . Finally, the daytime gaps in Q_E were filled in such a way that the monthly mean energy balance closure was achieved, while during the night the missing Q_E data were set to 0.

Data availability. The measured flux and meteorological data are available in Zenodo (<https://zenodo.org/>) research data depository.

5 *Author contributions.* TP, PO, KM, TL and AL designed the study. Field measurements and maintenance were carried out by MK, JR, TL, and AL. J-PT made spectral corrections and footprint analysis for the EC data. SS and PO corrected the WTL data and did the WTL background modelling. Rest of the data analysis was carried out by MK. MK wrote the paper with contributions from all coauthors.

Competing interests. The authors declare that they have no conflict of interest.

10

Acknowledgements. We are grateful for the financial support from the Maj and Tor Nessling foundation and from the Ministry of Transport and Communications through the Integrated Carbon Observing System (ICOS) research.

References

- Amiro, B. D., Barr, A. G., Barr, J. G., Black, T. A., Bracho, R., Brown, M., Chen, J., Clark, K. L., Davis, K. J., Desai, A. R., Dore, S., Engel, V., Fuentes, J. D., Goldstein, A. H., Goulden, M. L., Kolb, T. E., Lavigne, M. B., Law, B. E., Margolis, H. A., Martin, T., McCaughey, J. H., Misson, L., Montes-Helu, M., Noormets, A., Randerson, J. T., Starr, G. and Xiao, J.: Ecosystem carbon dioxide fluxes after disturbance in forests of North America, *J. Geophys. Res. Biogeosciences*, 115(4), doi:10.1029/2010JG001390, 2010.
- Aubinet, M., Vesala, T. and Papale, D., Eds.: *Eddy Covariance: a practical guide to measurement and data analysis*, Springer Science & Business Media., 2012.
- 10 Aurela, M., Lohila, A., Tuovinen, J.-P., Hatakka, J., Penttilä, T. and Laurila, T.: Carbon dioxide and energy flux measurements in four northern-boreal ecosystems at Pallas, *Boreal Environ. Res.*, 20(4), 455–473, 2015.
- Betts, A. K., Desjardins, R. L. and Worth, D.: Impact of agriculture, forest and cloud feedback on the surface energy budget in BOREAS, *Agric. For. Meteorol.*, 142(2–4), 156–169, doi:10.1016/j.agrformet.2006.08.020, 2007.
- Bhuiyan, R., Minkkinen, K., Helmisaari, H.-S., Ojanen, P., Penttilä, T. and Laiho, R.: Estimating fine-root production by tree species and understorey functional groups in two contrasting peatland forests, *Plant Soil*, 412(1–2), 299–316, doi:10.1007/s11104-016-3070-3, 2017.
- 15 Blodau, C., Basiliko, N. and Moore, T.: Carbon turnover in peatland mesocosms exposed to different water table levels, *Biogeochemistry*, 67(3), 331–351, doi:10.1023/B:BIOG.0000015788.30164.e2, 2004.
- Chimner, R. A. and Cooper, D. J.: Influence of water table levels on CO₂ emissions in a Colorado subalpine fen: An in situ microcosm study, *Soil Biol. Biochem.*, 35(3), 345–351, doi:10.1016/S0038-0717(02)00284-5, 2003.
- 20 Clark, K. L., Gholz, H. L. and Castro, M. S.: Carbon dynamics along a chronosequence of slash pine plantations in north Florida, *Ecol. Appl.*, 14(4), 1154–1171 [online] Available from: <http://www.esajournals.org/doi/pdf/10.1890/02-5391%5Cnhttp://www.esajournals.org/doi/abs/10.1890/02-5391>, 2004.
- Clarke, D. and Rieley, J.: *Strategy for Responsible Peatland Management*, edited by D. Clarke and J. Rieley, International Peat Society, Saarijärvi, Finland., 2010.
- 25 Davidson, E., Belk, E. and Boone, R. D.: Soil water content and temperature as independent or confounded factors controlling soil respiration in a temperate mixed hardwood forest, *Glob. Chang. Biol.*, 4(2), 217–227, doi:10.1046/j.1365-2486.1998.00128.x, 1998.
- Drzymulska, D.: Peat decomposition - Shaping factors, significance in environmental studies and methods of determination; A literature review, *Geologos*, 22(1), 61–69, doi:10.1515/logos-2016-0005, 2016.
- 30 Edwards, N. T. and Ross-Todd, B. M.: Soil Carbon Dynamics in a Mixed Deciduous Forest Following Clear-Cutting with and without Residue Removal, *Soil Sci. Soc. Am. J.*, 47(5), 1014–1021, doi:10.2136/sssaj1983.03615995004700050035x, 1983.

- Fischlin, A., Midgley, G. F., Price, J., Leemans, R., Gopal, B., Turley, C., Rounsevell, M., Dube, P., Tarazona, J. and Velichko, A.: Ecosystems, their properties, goods, and services, in: *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of working group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by M. L. Parry, O. F. Canziani, J. P. Palutikof, P. J. van der Linden, and C. E. Hanson, Cambridge University Press, Cambridge.
- 5 [online] Available from: <http://www.treesearch.fs.fed.us/pubs/33102>, 2007.
- Foken, T. and Wichura, B.: Tools for quality assessment of surface-based flux measurements, *Agric. For. Meteorol.*, 78(1–2), 83–105, doi:10.1016/0168-1923(95)02248-1, 1996.
- Fontaine, S., Bardoux, G., Abbadie, L. and Mariotti, A.: Carbon input to soil may decrease soil carbon content, *Ecol. Lett.*, 7(4), 314–320, doi:10.1111/j.1461-0248.2004.00579.x, 2004.
- 10 Fontaine, S., Barot, S., Barré, P., Bdioui, N., Mary, B. and Rumpel, C.: Stability of organic carbon in deep soil layers controlled by fresh carbon supply, *Nature*, 450(7167), 277–280, doi:10.1038/nature06275, 2007.
- Fredeen, A. L., Waughtal, J. D. and Pypker, T. G.: When do replanted sub-boreal clearcuts become net sinks for CO₂?, *For. Ecol. Manage.*, 239(1–3), 210–216, doi:10.1016/j.foreco.2006.12.011, 2007.
- Hommeltenberg, J., Schmid, H. P., Drösler, M. and Werle, P.: Can a bog drained for forestry be a stronger carbon sink than a
 15 natural bog forest?, *Biogeosciences*, 11(13), 3477–3493, doi:10.5194/bg-11-3477-2014, 2014.
- Howard, E. A., Gower, S. T., Foley, J. A. and Kucharik, C. J.: Effects of logging on carbon dynamics of a jack pine forest in Saskatchewan, Canada, *Glob. Chang. Biol.*, 10(8), 1267–1284, doi:10.1111/j.1365-2486.2004.00804.x, 2004.
- Humphreys, E. R., Black, T. A., Morgenstern, K., Li, Z. and Nesic, Z.: Net ecosystem production of a Douglas-fir stand for 3 years following clearcut harvesting, *Glob. Chang. Biol.*, 11(3), 450–464, doi:10.1111/j.1365-2486.2005.00914.x, 2005.
- 20 Huttunen, J. T., Nykänen, H., Martikainen, P. J. and Nieminen, M.: Fluxes of nitrous oxide and methane from drained peatlands following forest clear-felling in southern Finland, *Plant Soil*, 255(2), 457–462, doi:10.1023/A:1026035427891, 2003.
- IPCC: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.
- 25 Kaila, A., Sarkkola, S., Laurén, A., Ukonmaanaho, L., Koivusalo, H., Xiao, L., O’Driscoll, C., Asam, Z. U. Z., Tervahauta, A. and Nieminen, M.: Phosphorus export from drained Scots pine mires after clear-felling and bioenergy harvesting, *For. Ecol. Manage.*, 325, 99–107, doi:10.1016/j.foreco.2014.03.025, 2014.
- Kaimal, J. C. and Finnigan, J. J.: *Atmospheric boundary layer flows: their structure and measurement*, Oxford University Press, Oxford., 1994.
- 30 Kolari, P., Pumpanen, J., Rannik, Ü., Ilvesniemi, H., Hari, P. and Berninger, F.: Carbon balance of different aged Scots pine forests in Southern Finland, *Glob. Chang. Biol.*, 10(7), 1106–1119, doi:10.1111/j.1365-2486.2004.00797.x, 2004.
- Korkiakoski, M., Tuovinen, J.-P., Aurela, M., Koskinen, M., Minkkinen, K., Ojanen, P., Penttilä, T., Rainne, J., Laurila, T. and Lohila, A.: Methane exchange at the peatland forest floor - Automatic chamber system exposes the dynamics of small fluxes, *Biogeosciences*, 14(7), 1947–1967, doi:10.5194/bg-14-1947-2017, 2017.

- Kormann, R. and Meixner, F. X.: An analytical footprint model for non-neutral stratification, *Boundary-Layer Meteorol.*, 99(2), 207–224, doi:10.1023/A:1018991015119, 2001.
- Kowalski, A. S., Loustau, D., Berbigier, P., Manca, G., Tedeschi, V., Borghetti, M., Valentini, R., Kolari, P., Berninger, F., Rannik, Ü., Hari, P., Rayment, M., Mencuccini, M., Moncrieff, J. and Grace, J.: Paired comparisons of carbon exchange
5 between undisturbed and regenerating stands in four managed forest in Europe, *Glob. Chang. Biol.*, 10(10), 1707–1723, doi:10.1111/j.1365-2486.2004.00846.x, 2004.
- Kowalski, S., Sartore, M., Burrell, R., Berbigier, P. and Loustau, D.: The annual carbon budget of a French pine forest (*Pinus pinaster*) following harvest, *Glob. Chang. Biol.*, 9(7), 1051–1065, doi:10.1046/j.1365-2486.2003.00627.x, 2003.
- Laurén, A., Heinonen, J., Koivusalo, H., Sarkkola, S., Tattari, S., Mattsson, T., Ahtiainen, M., Joensuu, S., Kokkonen, T. and
10 Finér, L.: Implications of uncertainty in a pre-treatment dataset when estimating treatment effects in paired catchment studies: Phosphorus loads from forest clear-cuts, *Water. Air. Soil Pollut.*, 196(1–4), 251–261, doi:10.1007/s11270-008-9773-1, 2009.
- Lloyd, J. and Taylor, J. A.: On the Temperature Dependence of Soil Respiration, *Funct. Ecol.*, 8, 315–323, 1994.
- Londo, A. J., Messina, M. G. and Schoenholtz, S. H.: Forest Harvesting Effects on Soil Temperature, Moisture, and Respiration in a Bottomland Hardwood Forest, *Soil Sci. Soc. Am. J.*, 63(3), 637–644, doi:10.2136/sssaj1999.03615995006300030029x,
15 1999.
- Machimura, T., Kobayashi, Y., Iwahana, G., Hirano, T., Lopez, L., Fukuda, M. and Fedorov, A. N.: Change of carbon dioxide budget during three years after deforestation in eastern siberian larch forest, *J. Agric. Meteorol.*, 60(5), 653–656, 2005.
- Mäkiranta, P., Minkkinen, K., Hytönen, J. and Laine, J.: Factors causing temporal and spatial variation in heterotrophic and rhizospheric components of soil respiration in afforested organic soil croplands in Finland, *Soil Biol. Biochem.*, 40(7), 1592–
20 1600, doi:10.1016/j.soilbio.2008.01.009, 2008.
- Mäkiranta, P., Laiho, R., Fritze, H., Hytönen, J., Laine, J. and Minkkinen, K.: Indirect regulation of heterotrophic peat soil respiration by water level via microbial community structure and temperature sensitivity, *Soil Biol. Biochem.*, 41(4), 695–703, doi:10.1016/j.soilbio.2009.01.004, 2009.
- Mäkiranta, P., Riutta, T., Penttilä, T. and Minkkinen, K.: Dynamics of net ecosystem CO₂ exchange and heterotrophic soil
25 respiration following clearfelling in a drained peatland forest, *Agric. For. Meteorol.*, 150(12), 1585–1596, doi:10.1016/j.agrformet.2010.08.010, 2010.
- Mäkiranta, P., Laiho, R., Penttilä, T. and Minkkinen, K.: The impact of logging residue on soil GHG fluxes in a drained peatland forest, *Soil Biol. Biochem.*, 48, 1–9, doi:10.1016/j.soilbio.2012.01.005, 2012.
- Maljanen, M., Sigurdsson, B. D., Guðmundsson, J., Óskarsson, H., Huttunen, J. T. and Martikainen, P. J.: Greenhouse gas
30 balances of managed peatlands in the Nordic countries present knowledge and gaps, *Biogeosciences*, 7(9), 2711–2738, doi:10.5194/bg-7-2711-2010, 2010.
- Mamkin, V., Kurbatova, J., Avilov, V., Mukhartova, Y., Krupenko, A., Ivanov, D., Levashova, N. and Olchev, A.: Changes in net ecosystem exchange of CO₂, latent and sensible heat fluxes in a recently clear-cut spruce forest in western Russia: results from an experimental and modeling analysis, *Environ. Res. Lett.*, 11(12), 125012, doi:10.1088/1748-9326/aa5189,

2016.

Martikainen, P. J., Nykänen, H., Alm, J. and Silvola, J.: Change in Fluxes of Carbon-Dioxide, Methane and Nitrous-Oxide Due To Forest Drainage of Mire Sites of Different Trophic, *Plant Soil*, 168, 571–577, doi:10.1007/bf00029370, 1995.

5 McMillen, R. T.: An eddy correlation technique with extended applicability to non-simple terrain, *Boundary-Layer Meteorol.*, 43(3), 231–245, doi:10.1007/BF00128405, 1988.

Meyer, A., Tarvainen, L., Noursatpour, A., Björk, R. G., Ernfors, M., Grelle, A., Kasimir-Klemetsson, Å., Lindroth, A., Rantfors, M., Rütting, T., Wallin, G., Weslien, P. and Klemetsson, L.: A fertile peatland forest does not constitute a major greenhouse gas sink, *Biogeosciences*, 10(11), 7739–7758, doi:10.5194/bg-10-7739-2013, 2013.

10 Minkkinen, K., Laine, J. and Hökkä, H.: Tree stand development and carbon sequestration in drained peatland stands in Finland—a simulation study, *Silva Fenn.*, 35(January), 55–69 [online] Available from: http://www.researchgate.net/publication/228505488_Tree_stand_development_and_carbon_sequestration_in_drained_peatland_stands_in_Finland-a_simulation_study/file/79e4150a33d5148608.pdf, 2001.

Moore, C. J.: Frequency response corrections for eddy correlation systems, *Boundary-Layer Meteorol.*, 37, 17–35, doi:10.1007/BF00122754, 1986.

15 Ojanen, P., Minkkinen, K., Alm, J. and Penttilä, T.: Soil-atmosphere CO₂, CH₄ and N₂O fluxes in boreal forestry-drained peatlands, *For. Ecol. Manage.*, 260(3), 411–421, doi:10.1016/j.foreco.2010.04.036, 2010.

Ojanen, P., Minkkinen, K. and Penttilä, T.: The current greenhouse gas impact of forestry-drained boreal peatlands, *For. Ecol. Manage.*, 289, 201–208, doi:10.1016/j.foreco.2012.10.008, 2013.

20 Ojanen, P., Mäkiranta, P., Penttilä, T. and Minkkinen, K.: Do logging residue piles trigger extra decomposition of soil organic matter?, *For. Ecol. Manage.*, 405(June), 367–380, doi:10.1016/j.foreco.2017.09.055, 2017.

Ojanen, P., Minkkinen, K., Alm, J. and Penttilä, T.: Corrigendum to “Soil–atmosphere CO₂, CH₄ and N₂O fluxes in boreal forestry-drained peatlands” [*For. Ecol. Manage.* 260 (2010) 411–421](S0378112710002409)(10.1016/j.foreco.2010.04.036), *For. Ecol. Manage.*, 412(February), 95–96, doi:10.1016/j.foreco.2018.01.020, 2018.

25 Päivänen, J. and Hånell, B.: Peatland ecology and forestry: a sound approach, University of Helsinki Department of Forest Sciences Publications 3, Helsinki, Finland., 2012.

Pearson, M., Saarinen, M., Minkkinen, K., Silvan, N. and Laine, J.: Short-term impacts of soil preparation on greenhouse gas fluxes: A case study in nutrient-poor, clearcut peatland forest, *For. Ecol. Manage.*, 283, 10–26, doi:10.1016/j.foreco.2012.07.011, 2012.

30 Pirinen, P., Simola, H., Aalto, J., Kaukoranta, J.-P., Karlsson, P. and Ruuhela, R.: Climatological statistics of Finland 1981–2010, Reports, 2., Finnish Meteorological Institute, Helsinki, Finland., 2012.

Pypker, T. G. and Fredeen, A. L.: Ecosystem CO₂ flux over two growing seasons for a sub-Boreal clearcut 5 and 6 years after harvest, *Agric. For. Meteorol.*, 114(1–2), 15–30, doi:10.1016/S0168-1923(02)00139-9, 2002.

R Core Team: R: A language and Environment for Statistical Computing., [online] Available from: <https://www.r-project.org>, 2018.

- Rannik, Ü., Vesala, T. and Keskinen, R.: On the damping of temperature fluctuations in a circular tube relevant to the eddy covariance measurement technique, *J. Geophys. Res. Atmos.*, 102(D11), 12789–12794, 1997.
- Rannik, Ü., Altimir, N., Raittila, J., Suni, T., Gaman, A., Hussein, T., Hölttä, T., Lassila, H., Latokartano, M., Lauri, A., Natsheh, A., Petäjä, T., Sorjamaa, R., Ylä-Mella, H., Keronen, P., Berninger, F., Vesala, T., Hari, P. and Kulmala, M.: Fluxes of carbon dioxide and water vapour over Scots pine forest and clearing, *Agric. For. Meteorol.*, 111(3), 187–202, doi:10.1016/S0168-1923(02)00022-9, 2002.
- Räsänen, M., Aurela, M., Vakkari, V., Beukes, J. P., Tuovinen, J.-P., Van Zyl, P. G., Josipovic, M., Venter, A. D., Jaars, K., Siebert, S. J., Laurila, T., Rinne, J. and Laakso, L.: Carbon balance of a grazed savanna grassland ecosystem in South Africa, *Biogeosciences*, 14(5), 1039–1054, doi:10.5194/bg-14-1039-2017, 2017.
- 10 Riutta, T., Laine, J. and Tuittila, E.-S.: Sensitivity of CO₂ exchange of fen ecosystem components to water level variation, *Ecosystems*, 10(5), 718–733, doi:10.1007/s10021-007-9046-7, 2007.
- Roulet, N., Ash, R., Quinton, W. and Moore, T.: Methane flux from drained northern peatlands: Effect of a persistent water table lowering on flux, *Global Biogeochem. Cycles*, 7(4), 749–769, 1993.
- Saari, P., Saarnio, S., Kukkonen, J. V. K., Akkanen, J., Heinonen, J., Saari, V. and Alm, J.: DOC and N₂O dynamics in upland and peatland forest soils after clear-cutting and soil preparation, *Biogeochemistry*, 94(3), 217–231, doi:10.1007/s10533-009-9320-1, 2009.
- Sarkkola, S., Hökkä, H., Koivusalo, H., Nieminen, M., Ahti, E., Päivänen, J. and Laine, J.: Role of tree stand evapotranspiration in maintaining satisfactory drainage conditions in drained peatlands, *Can. J. For. Res.*, 40(8), 1485–1496, doi:10.1139/X10-084, 2010.
- 20 Schulze, E.-D., Lloyd, J., Kelliher, F. M., Wirth, C., Rebmann, C., Luhker, B., Mund, M., Knohl, A., Milyukova, I. M., Schulze, W., Ziegler, W., Varlagin, A. B., Sogachev, A. F., Valentini, R., Dore, S., Grigoriev, S., Kolle, O., Panfyorov, M. I., Tchebakova, N. and Vygodskaya, N. N.: Productivity of forests in the eurosiberian boreal region and their potential to act as a carbon sink - a synthesis, *Glob. Chang. Biol.*, 5(6), 703–722, doi:10.1046/j.1365-2486.1999.00266.x, 1999.
- Silvola, J., Alm, J., Ahlholm, U., Nykänen, H. and Martikainen, P. J.: CO₂ Fluxes from Peat in Boreal Mires under Varying Temperature and Moisture Conditions, *J. Ecol.*, 84(2), 219–228, 1996.
- 25 Sundqvist, E., Vestin, P., Crill, P. M., Persson, T. and Lindroth, A.: Short-term effects of thinning, clear-cutting and stump harvesting on methane exchange in a boreal forest, *Biogeosciences*, 11(21), 6095–6105, doi:10.5194/bg-11-6095-2014, 2014.
- Takagi, K., Fukuzawa, K., Liang, N., Kayama, M., Nomura, M., Hojyo, H., Sugata, S., Shibata, H., Fukazawa, T., Takahashi, Y., Nakaji, T., Oguma, H., Mano, M., Akibayashi, Y., Murayama, T., Koike, T., Sasa, K. and Fujinuma, Y.: Change in CO₂ balance under a series of forestry activities in a cool-temperate mixed forest with dense undergrowth, *Glob. Chang. Biol.*, 15(5), 1275–1288, doi:10.1111/j.1365-2486.2008.01795.x, 2009.
- 30 Turunen, J., Tomppo, E., Tolonen, K. and Reinikainen, A.: Estimating carbon accumulation rates of undrained mires in Finland—application to boreal and subarctic regions, *The Holocene*, 12(1), 69–80, doi:10.1191/0959683602h1522rp, 2002.
- Uri, V., Kukumägi, M., Aosaar, J., Varik, M., Becker, H., Morozov, G. and Karoles, K.: Ecosystems carbon budgets of

differently aged downy birch stands growing on well-drained peatlands, *For. Ecol. Manage.*, 399, 82–93, doi:10.1016/j.foreco.2017.05.023, 2017.

Webb, E. K., Pearman, G. I. and Leuning, R.: Correction of flux measurements for density effects due to heat and water vapour transfer, *Q. J. Roy. Meteorol. Soc.*, 106(447), 85–100, doi:http://dx.doi.org/10.1002/qj.49710644707, 1980.

- 5 Williams, C., Vanderhoof, M., Khomik, M. and Ghimire, B.: Post-clearcut dynamics of carbon, water and energy exchanges in a midlatitude temperate, deciduous broadleaf forest environment, *Glob. Chang. Biol.*, 20(3), 992–1007, doi:10.1111/gcb.12388, 2013.

Yu, Z.: Holocene carbon flux histories of the world's peatlands: Global carbon-cycle implications, *Holocene*, 21(5), 761–774, doi:10.1177/0959683610386982, 2011.

- 10 Zerva, A. and Mencuccini, M.: Short-term effects of clearfelling on soil CO₂, CH₄, and N₂O fluxes in a Sitka spruce plantation, *Soil Biol. Biochem.*, 37(11), 2025–2036, doi:10.1016/j.soilbio.2005.03.004, 2005.

Table 1. Mean soil organic carbon, nitrogen, CN ratio and bulk density and their standard deviation in different soil layers determined before the harvest (n=3–33).

Layer	Soil organic carbon (%)	Soil nitrogen (%)	CN ratio	Bulk density (g cm ⁻³)
Litter	56.2 ± 2.3	1.7 ± 0.4	33.2 ± 2.3	0.005 ± 0.003
0-10 cm	55.2 ± 2.1	2.2 ± 0.2	24.9 ± 2.1	0.12 ± 0.03
10-20 cm	58.9 ± 1.6	2.5 ± 0.2	23.8 ± 1.6	0.18 ± 0.02
20-30 cm	59.1 ± 0.8	2.4 ± 0.2	24.6 ± 0.8	0.13 ± 0.03
30-40 cm	59.2 ± 0.3	2.2 ± 0.1	26.4 ± 0.3	0.13 ± 0.04

Layer	Soil organic carbon (%)	Soil nitrogen (%)	CN ratio	Bulk density (g cm ⁻³)
Humus	56.2 ± 2.3	1.7 ± 0.4	33.2 ± 2.3	0.005 ± 0.003
0-10 cm	55.2 ± 2.1	2.2 ± 0.2	24.9 ± 2.1	0.12 ± 0.03
10-20 cm	58.9 ± 1.6	2.5 ± 0.2	23.8 ± 1.6	0.18 ± 0.02
20-30 cm	59.1 ± 0.8	2.4 ± 0.2	24.6 ± 0.8	0.13 ± 0.03
30-40 cm	59.2 ± 0.3	2.2 ± 0.1	26.4 ± 0.3	0.13 ± 0.04

5

Table 2. Annual and summertime (JJA) balances of net ecosystem exchange (NEE), modelled gross primary production (GPP) and modelled total ecosystem respiration (R_{eco}), and the modelled summertime sums of forest floor respiration (R_{ff}).

	Annual		Summer (JJA)	
	Apr 2016 - Mar 2017	Apr 2017 - Mar 2018	2016	2017
NEE [g CO ₂ m ⁻²]	3086±148	2072±124	1558±99	915±79
GPP [g CO ₂ m ⁻²]	659	1106	389	761
R _{eco} [g CO ₂ m ⁻²]	3727	3135	1928	1652
R _{ff} [g CO ₂ m ⁻²]	-	-	980±299	1047±113

5

	Annual		Summer (JJA)	
	Apr 2016 - Mar 2017	Apr 2017 - Mar 2018	2016	2017
NEE [g CO ₂ m ⁻²]	3086±148	2072±124	1558±99	915±79
GPP [g CO ₂ m ⁻²]	659	1106	389	761
R _{eco} [g CO ₂ m ⁻²]	3727	3135	1928	1652
R _{ff} [g CO ₂ m ⁻²]	-	-	980±299	1047±113



Figure 1. Aerial view of the clear-cut site and the surrounding forest at Lettosuo. The red dot shows the location of the eddy covariance mast, and the red lines surround the target area ($80^{\circ} - 315^{\circ}$). The meteorological measurements were conducted at the location indicated by the orange dot, while the orange line marks the water table, soil temperature and chamber measurements (4, 8, 12 and 22.5 m from the ditch). Similar measurements of greenhouse gas fluxes, water table and soil temperature within the control site, surrounded by the green lines, are shown with a light blue line. The windroses on the left show the cumulative footprint contributions (in %) of the accepted flux data.




Figure 2. Time series of (a) daily mean air temperature (T_{air}) and (b) hourly mean soil temperatures at 5, 10, 20 and 30 cm depths (T_{soil}) measured at Lettosuo, and (c) daily precipitation sum and (d) daily mean snow level recorded at the Jokioinen observatory (35 km northwest of Lettosuo). The clearcutting (red vertical line) was carried out in February–

5 March 2016.




Figure 3. Mean daily water table level at the control site (measured; green) and the clear-cut site with (measured; black line) and without (estimated; grey line) the harvesting effect averaged over all the measurement points (4, 8, 12 and 22.5 m from the ditch) from June 2015 to October 2017. The clearcutting (red vertical line) was carried out in
 5 February–March 2016.




Figure 4. The gap-filled time series of daily mean net ecosystem exchange (NEE, top panel) and the corresponding modelled gross primary production (GPP, middle panel) and ecosystem respiration (R_{eco} , bottom panel) based on Eq. (A2) and Eq. (A3), respectively.




Figure 5. Temperature (a; Eq. A3) and light responses (b; Eq. A2) of 30 min CO₂ fluxes for summers (JJA) 2016 (black) and 2017 (blue). The temperature response equation was fitted to the night-time data (PPFD < 20 μmol m⁻² s⁻¹) and the light response equation to the daytime (PPFD > 20 μmol m⁻² s⁻¹) data.




Figure 6. Monthly mean diel cycles of gap-filled net ecosystem exchange (NEE) measurements in the first (2016, grey), second (2017, black) and third (2018, blue) calendar year after the clearcutting.




Figure 7. The monthly sums of net ecosystem exchange (NEE), gross primary production (GPP) and ecosystem respiration (R_{eco}) based on EC data in April 2016 – March 2018.




Figure 8. Hourly mean fluxes measured with manual chambers of (a) CO₂, (b) CH₄ and (c) N₂O averaged over all measurement points (4, 8, 12 and 22.5 m from the ditch) at the clear-cut (black) and control (green) sites from June 2015 to August 2017. The error bars show the standard error of the mean. The clearcutting (red vertical line) was carried out in February–March 2016.




Figure 9. The division of ecosystem respiration (R_{eco} , based on EC measurements) to forest floor respiration R_{ff} (chamber measurements) and respiration of logging residues ($R_{residue} = R_{eco} - R_{ff}$) for summers 2016 and 2017.




Figure 10. Temperature response (Eq. A3) of the hourly mean CO₂ fluxes measured with manual chambers at the clear-cut (a) and control (b) site in 2015 (black), 2016 (blue) and 2017 (green).





Figure 11. Daily sums of net radiation (Q_N), sensible heat flux (Q_H), latent heat flux (Q_E) and ground heat flux (Q_G).

(a)



(b)




Figure 12. Monthly mean diel cycles of net radiation (Q_N), air temperature (T_{air}), sensible heat flux (Q_H), latent heat flux (Q_E) and ground heat flux (Q_G) and the monthly mean daytime (10:00–16:00, UTC+2) Bowen ratios (β) from April to September in 2016 (a) and 2017 (b).