

May 9, 2019

Dear Dr Ciavatta,
Associate Editor, Biogeosciences

We are re-submitting our manuscript “*Rates and drivers of Red Sea plankton community metabolism*” to BGD. In this new version of the paper, we have addressed the issues raised by the reviewers from our last submission. Regarding the main concerns from the reviewers, the principal changes to our manuscript are detailed in the following section:

1. About the different variables used to explain the changes of planktonic metabolism in the Red Sea (i.e., latitude, water depth and seasonality) and reviewers’ suggestion to rewrite the narrative of the manuscript by focussing on one or two variables.

Reply

We had modified the text and had mainly focused on the latitudinal and seasonal variability of planktonic metabolism and their relationship with temperature and nutrient availability (both tightly related to the latitudinal gradient that characterised the Red Sea). Although changes throughout the water column are relevant, we agreed with the reviewer that it added an unnecessary complexity to the text.

Action

We re-wrote most of the results section (now between lines 322–448), and modified the figures so that Figure 2 describes the latitudinal gradient of variables only in surface waters. Figure 3: we modify the format and the data presented, so we only included data within the photic layer. We removed Figures 4–5, and Figures 1A and 2A. We also removed Table 1, and Table 2. The information from Table 2 is now presented as a figure (now Figure 5) with the overall information and with all the data plotted. The changes in the results are in agreement with changes in the narrative of the discussion section. The main changes can be tracked between lines 574–580 and between lines 583–588 of the new manuscript.

2. About the methodology chosen and length of the incubations.

Reply

The Winkler titration method used in this study, is a methodology extensively used in studies that aim to measure autotrophic planktonic photosynthetic rates (i.e. gross primary production rates) and respiration rates derived from both the autotrophic and heterotrophic plankton community (see e.g., Williams et al., 1979; Duarte and Agustí, 1998; Bender et al., 1999; Robinson and Williams, 1999; Ducklow et al., 2000; Serret et al., 2001; Robinson et al., 2002; Serret et al., 2009; García-Martín et al., 2017). The selected incubation time allows to fully take into account diel changes of both metabolic rates, as community respiration rates occur during day and night.

Action

We have clarified this point in the methods section 3.3 (between lines 200–208). In addition, we have ensured that the sampling, filling and fixing of the samples is

described in sufficient detail. We also clarified how we calculated gross primary production, community respiration and net community production (which is the balance between the autotrophic and heterotrophic metabolism, and not only the result of photosynthetic activity). These calculations can be seen between lines 256 and 277.

3. About the lack of statistical analyses.

Reply

We agreed with the reviewers that a section with a detailed description of all the statistical analyses was missing, and that the statistical analyses mentioned in the results were not properly clarified, hence the reviewers were under the impression that we did not perform any statistics to support our conclusions.

Action

We have included a detailed description of all the statistical analyses done (see section 2.4 between lines 288–321). In addition, the results of each analysis are detailed whenever they are presented. See e.g. line 333 or line 438.

4. We also detected some typos and errors in three figures and one table that have been rectified, yet neither of those changed our main conclusions in any way.

Figures:

- Figure 8C (now Figure 4G) R^2 is 0.38 not 0.39 and, R^2 in Figure 8I is 0.16 not 0.17
- Figure 7A (now Figure 6A), the slope is 0.73 not 0.74
- Figure 9B (now Figure 7B), the slope is -0.72 not -0.73

The detailed answer to the reviewers with the updated line numbers is presented in the following section.

I would be grateful if this newly revised version of the manuscript was considered for publication in Biogeosciences.

Sincerely,

Daffne C. López-Sandoval

Red Sea Research Center

King Abdullah University of Science and Technology (KAUST)

Thuwal, Jeddah, 23955-6900, Kingdom of Saudi Arabia

Phone (+966) 12 808 2659

Reply to reviewers

Reviewer 1

2.1.1 General comments

The authors describe a dataset of environmental variables related to the metabolism of planktonic communities along a depth and latitudinal gradient in a seasonal resolution in the Red Sea. The authors conclude that gross primary production relates positively to sea surface temperature and nutrient availability. The dataset is extensive, and the research questions (for this part of the Red Sea), to my knowledge, are novel and worthy of publication. The abstract is clear and reads well, but shows a different narrative than the rest of the manuscript. Thus, I suggest for the authors to consider rewriting the manuscript. As mentioned in the author contributions, the manuscript is written by several people and this is noticeable (see specific comments). The abstract mentions the latitudinal gradient but the ms introduces two more variables, i.e. depth and seasonality. While interesting variables, they make the story confusing at times and harder to disentangle the story the authors want to tell (according to the abstract). Concerns about the methods used are mentioned in the specific comments and need to be addressed first. Proper description of statistical analyses is lacking.

My recommendation is that the ms needs major revisions, but only if methodological concerns can be addressed adequately. Then, I suggest a complete overhaul of the manuscripts narrative by focusing on 1 or 2 of the 3 major variables (latitude, water depth and seasonality) and stick with these in the entire narrative. Also, there needs to be a clear description of used statistics in the M&M section and figures and tables should be cut back and/or improved. Consistency in the presentation of the results (including the statistics) and the use of abbreviations (as well as changing them) is recommended

Reply to general comments

We sincerely appreciate the thorough revision, comments and the time devoted to review our manuscript in such a rigorous manner, as it helped us to greatly improve our manuscript. We have addressed all the reviewer's comments, and significant changes were done to the manuscript, figures and tables. Also, we have included a detailed description of the statistical analyses.

Action

The changes can be tracked in the new version of the manuscript in the Results section between lines 322–448. We modified the figures according to the new narrative of the results as such that Figure 2 now only describes the latitudinal gradient of variables in surface waters. In Figure 3, we modified the format and the data presented, so now we only focus on data within the photic layer. We removed Figures 4–5, Figure 1A and 2A, as well as Table 1 and Table 2. The information from Table 2 is now presented as a figure (now Figure 5) with the overall information and with all the data plotted. The changes in the results are in agreement with changes in the narrative of the discussion section. The main changes can be tracked between lines 574–580 and between lines 584–588 of the new manuscript.

2.1.2 Reply to specific comments

Reviewer comment 1

Title: Says Red Sea but Gulfs are not included.

Reply

We agree with the reviewer that our study does not include data from the Gulf of Aqaba nor does it include the western half of the Red Sea. However, this detail is carefully described in the Introduction (lines 153) and Methods sections (between lines 168–170) and in Figure 1, so the reader will not be misled.

Action

We preferred to leave the title as it is, since we provided sufficient detail in the manuscript on which areas were sampled.

Reviewer comment 2

Abstract: Line 10: Mentioning “Low productive waters” immediately brings down the importance of the story.

Action

We modify the text accordingly and deleted “low productive waters” (see line 10)

Reviewer comment 3

The first paragraph is loaded with self-referencing while many others are not or less.

Reply

There is no intent to load the paper unnecessarily with self-references, but it turns out that the co-authors of this paper have published a large number of relevant articles on the topic. We have now, however, added additional works, by other researchers.

Action

These new references can be tracked between lines 39–45 and between 129–130.

Reviewer comment 4

Page 2, line 4-5 and 11: Introduce abbreviations once (see technical corrections) and use them consistently throughout the ms.

Reply

Thank you for your comment. We have modified the text accordingly.

Action

Abbreviations were introduced between lines 13 and 14 and subsequently used consistently.

Reviewer comment 5

Page 2: The abbreviations of GPP, CR and NCP are presented with units of daily oxygen produced or used. However, these abbreviations are normally used for daily production and use of carbon. I suggest the authors change the abbreviations for these processes and/or use a conversion factor to present daily carbon production and use.

Reply

This and other comments from the reviewer, led us to believe that there is a misunderstanding as they seem to suggest that our work was focused on primary production, which is performed by the photosynthetic components of the plankton community during the daytime and that has gross and net components (as phytoplankton excrete and respire carbon). However, our paper focuses on the entire plankton community, both photosynthetic and heterotrophic (e.g. bacteria), where the net community production (NCP) represents the organic matter remaining after consumption of the GPP through respiration by plants (autotrophs), microbes (either autotrophs or heterotrophs), and animals (heterotrophs) (Ducklow and Doney 2013). Hence, it is the balance of the photosynthetic processes (GPP) and respiratory activity of the entire plankton community (not just phytoplankton) measured during 24 h periods. Studies that focus on the photosynthetic component of the plankton community (e.g. Net Primary Production, NPP) report values for the daylight period only whereas studies such as this, report (24 h) rates. For instance, published syntheses of community metabolism rates report values per day (24 h), e.g., Robinson and Williams 2005, Regaudie-de-Gioux 2012, 2013).

Reports of daily GPP, NCP and CR rates in oxygen units has remained the case since Haaken H. Gran used the light and dark bottle method in combination with the Winkler method to report community metabolic rates (Gaarder and Gran 1927; Gran 1927). The reviewer can find other examples in McIntire et al. 1964; Williams et al. 1979; Williams et al. 1983; Owens and Crumpton 1995; Robinson and Williams 1999; López-Urrutia et al. 2006; Stanley et al. 2010; García-Martín et al. 2017. The use of 24 h to report rates is indeed not just a tradition but is justified as the metabolic budget needs be resolved over 24 h to be completed. For photosynthesis this does not apply as it is a light-dependent process, but respiration, which is necessary to define net community production, occurs at night as well. Moreover, all of the above papers report rates based on oxygen units since this is the property measured, and converting these to C requires some assumptions, such as a theoretical PQ value. However, to allow comparisons with another component of carbon cycling, we have provided an estimate of what, for example, the mean GPP reported here represents in terms of carbon production, assuming a PQ=1.

Action

We have clarified this point in the Methods section 3.3 (between lines 200–208). In addition, we clarified how we calculated gross primary production, community respiration and net community production (which is the balance between the autotrophic and heterotrophic metabolism, and not only the result of photosynthetic activity). These calculations can be seen between lines 256 and 277. The average GPP rates in carbon units can be tracked between lines 401–402. We have also included the equivalent of CR in carbon units (assuming a RQ=1) in line 403.

Reviewer comment 6

Page 4, line 7-8: There are plenty of references that describe metabolism in the northern part of the Red Sea (e.g. Rahav et al. 2015 MEPS, Tilstra et al 2018 Frontiers, Levanon-Spanier et al 1979 Deep Sea Res.)

Reply

Thank you for the references, we have included them in our manuscript.

Action

Please find the suggested references and others more between lines 129–130.

Reviewer comment 7

Page 5-6: Silicate is measured, mentioned in the results and in many figures/tables (with significant interactions) but nowhere mentioned in the Discussion. If not important, mention briefly in Discussion. Page 6-7, line 20 and 1 (resp.): Was NH₄ determined? If not, then you have NO_x values, not DIN.

Reply

Thank you for pointing this out. Regarding the lack of discussion about the specific relationship between metabolic rates and silicates, we would like to mention that we did not discuss in detail the relationship between metabolic rates and any of the inorganic nutrients measured, as we aimed to discuss the overall patterns found among all variables (i.e., nutrients, temperature, autotrophic biomass).

Action

We have replaced DIN with NO_x throughout the manuscript. Please see lines 329 and 389, as well as Figure 3, and Table 1.

Reviewer comment 8

Page 7, line 10: provide actual depths of PAR measurements. Also, I am confused about the use of 100%, 60-20 and 8-1 as table 1 and 2 give different ranges. Is the data comparable if different depths of sampling were used?

Reply

The reviewer is absolutely right, and we understand the reviewer's confusion. During our surveys, the samples to quantify planktonic metabolic rates were consistently taken within the first optical depth (ζ), towards the bottom of the photic layer, and one intermediate sample was either taken where we found the max. Chl-a fluorescence, or in case the Chl-a max was at the surface or the bottom layers, the intermediate sample was collected towards the middle of the euphotic layer (approx. 2.3 ζ). Therefore, our measurements are comparable. We chose the sampling based on the optical depths instead of physical depths (i.e., depth in m) as they are biologically relevant to describe metabolic processes such as GPP.

Action

This has been clarified in the manuscript between lines 209–240

Reviewer comment 9

Page 7: Were samples for metabolic rates filtered? Does the planktonic community include both single and multicellular organisms? Were the optodes adjusted for salinity? A major, potential, flaw in the methods used for metabolic rates is that it appears as though net photosynthesis was measured for 24h. If correct, this includes an approx. 12-hour period of darkness and thus results in data that cannot be used for calculations for gross photosynthesis, i.e. O₂ measurements will be severely lower due to dark respiration. net photosynthesis should have been measured only during daylight and respiration rates should have been measured in 2 phases; during the daytime and during nighttime, so approximately 12:12 h as respiration rates can have a diurnal rhythm. So

extrapolating these data to daily rates could result in a wrong estimation of gross photosynthesis. Also, how was O₂ production data extrapolated to per day? I suggest authors stick to hourly values for oxygen rates. If methods are used correctly, carbon budgets can be calculated using a conversion factor. If net photosynthesis was measured during daylight and respiration for 24 hours, the authors need to state assumptions of the values to their manuscript (potential over- or underestimation of rates)

Reply

We agree with the reviewer that to measure net photosynthesis (i.e., the net organic carbon production in the light), a 24-h incubation can carry potential bias. However, in our study, we are neither measuring nor reporting net photosynthetic rates. As clarified in our reply to specific comments (5) and also stated in our introduction and methodology, our goal was to quantify the daily net community production. Therefore, we aimed to estimate the metabolic contribution not only from the autotrophic community but the entire planktonic community (i.e., the balance between the production and respiration of organic material). The selected methodology to quantify planktonic metabolism is based on the extensively used dark and light method in combination with the Winkler titration method (as mentioned in our reply AC5) and not optodes (as mentioned by the reviewer), in a 24 h incubation period, hence we report our results as daily rates.

Action

We have clarified why we used this methodology, and supported with references that the methodology chosen is largely used for the goals of this manuscript. These points can be seen in the methods section 3.3 (between lines 200–208)

Reviewer comment 10

Page 9, statistics: Need to be expanded with actual models used.

Reply

We agree with the reviewer that a section with a detailed description of all the statistical analyses was missing.

Action:

We have included a detailed description of all the statistical analyses done (see section 2.4 between lines 288–321) and in addition the results of each analysis are detailed whenever they are presented. See e.g. line 333 or line 438.

Reviewer comment 11

Page 9, line 11: NO_x, not DIN.

Action

We modified the text accordingly, as seen now in line 329.

Reviewer comment 12

Page 10, line 17: 56% of heterotrophs suggests dominance of this trophic strategy

Reply

We agreed with the reviewer

Action

We have modified the results section and we no longer describe by depths the autotrophic or heterotrophic status of the communities

Reviewer comment 13

Page 11, line 4: What models were used to test this?

Reply

We determined if plankton metabolism and nitrite and nitrate were correlated by using a Pearson's correlation

Action

We re-analysed the data and the results are shown between lines 389–390

Reviewer comment 14

Page 11, line 8: Which analysis?

Reply

We evaluated the relationship between GPP with CR and NCP by performing an OLS linear regression.

Action

This information has now been clarified in the statistical analyses section (between lines 296–317) and in the associated figure (Figure 6).

Reviewer comment 15

Page 11, line 11: Introduction of this statistical method should be in the appropriate section

Reply

We agree with the reviewer and have included it in the statistical analyses.

Action

Please refer to line 317, the rationale behind the Arrhenius plots was already described in section 2.3. This information can now also be found between lines 278–286.

Reviewer comment 16

Page 11: How were AE values calculated? - Page 11: AE are presented as negatives, are they? Next page the authors mention a positive value.

Reply

The procedure to obtain the activation energies were explained in the methods section 2.3 (page 8 between lines 16–21, now between lines 278–286). That being said, we determined the activation energies by fitting an OLS linear regression to the relationship between the natural logarithm of Chl-a specific metabolic rates and the inverse of the absolute temperature.

The slopes of these so-called Arrhenius plots represent the average activation energy. The negative values seen in Figure 10 (now figures 7 and 8), resulted from the way we

plotted the normalised metabolic rates against the inverse of the absolute temperature multiplied by the Boltzmann's constant (from 38 to 39.5 eV⁻¹, lower x-axis), the slope of the resulting relationship is negative. However, please note that the relationship with temperature (upper x-axis) would yield a positive slope if plotted from 20.7 to 32.3 °C.

Reviewer comment 17

Page 13, line 14: GPP is said to be low, compared to what?

Action

We clarified this point in the new version of the manuscript, please see in line 622.

Reviewer comment 18

Page 15, line 18: How was AE standardized to Chl-a?

Reply

We understand the confusion; the sentence was poorly written. We aimed to describe the activation energies obtained from the relationship between the Chl-a specific GPP data and temperature.

Action

We completely changed this section of the discussion, please refer to lines 728–732.

Reviewer comment 19

Page 16, line 2: What is “the ocean”?

Action

We modified this noun, as we meant open oceanic waters. Please see line 730.

Reviewer comment 20

Page 16: Opens with “Surprisingly” and a discussion, then the next paragraph mentions a contradiction that is not surprising. What is the contradiction exactly the authors mean?

Reply

AC22: Thank you for highlighting this point

Action

We deleted this adverb as it is misleading

Reviewer comment 21

RC23 Page 16, line 6-7: Authors compare results with other references but need to mention actual values.

Reply

We added the requested information

Action

The changes can now be tracked between lines 734–737 in the new version of the manuscript

Reviewer comment 22

Figure 3: Thickness of the pink or green seems to say something about how significant it is but this is said nowhere. In line with this, the diagonal dark green lines seem to signify extreme significance instead of same variable and thus not tested. DIN is NO_x. Are variables tested at different depths than metabolic rates of plankton? If so, how can you relate the 2?

Reply

We agree with the reviewer's comments.

Action

We modified this graph to present information relevant to the depths where we analysed metabolic rates (i.e., only photic layer). In addition, we modified the way the results are presented, and we explained the color code.

Reviewer comment 23

Figure 4-6: Lots of white space and hard to see with tiny colored dots anyway. Revise these figures. I suggest to distill from them the most important results you want to show and add the rest to the supplementary section.

Action

We decided to remove the figures in this new version of the manuscript

Reviewer comment 24

Figure 7: could be mentioned with text in the results section. Suggest moving figure to supplements.

Reply

The information regarding this figure is mentioned in the results, but as it also provides information to derive the GPP threshold, we prefer to keep it as a main result.

Reviewer comment 25

Figure 9: Same as Figure 7, B is missing a parenthesis on the y-axis - Figure 10: Same as Figure 7.

Reply

Since figure 9 and 10 are both explaining one of the main results (i.e. the metabolic response of metabolic rates to temperature) we prefer to keep them as main figures.

Action

We added the missing parenthesis to the y-axis of the figures.

Reviewer comment 26

Please use continues line numbers for the manuscript

Reply

We followed the format and template designated by the journal.

Action

However, in the new version of the manuscript we changed to continuous lines as suggested by the reviewer.

Reviewer comment 27

Page 1, line 8-9: Please rewrite, it reads as if you want to understand their variability and their present and their future but you want to understand their variability in the present and the future

Action

This has been modified as suggested, please refer to line 9

Reviewer comment 28

Page 2, line 4-5: Add community

Action

Done as suggested, please see in line 38

Reviewer comment 29

Page 2, line 11: First mention of NCP, introduce abbreviation.

Reply

NCP was mentioned for the first time on Page 1 line 14

Reviewer comment 30

Page 3, line 1: "The Red Sea is a semi-enclosed"

Action

We modified the text, please see line 77

Reviewer comment 31

Page 3, line 3-5: Consider merging this sentence with the previous one

Action

Please see the changes between lines 79–80

Reviewer comment 32

Page 3, line 9: "throughout the year"

Action

Changed as suggested, see line 85

Reviewer comment 33

Page 3, line 10: Delete the dot before the references

Action

Noted, see line 86

Reviewer comment 33

Page 4, line 12: Add “relatively” to “unproductive waters”

Action

Done as suggested, see line 133

Reviewer comment 34

Page 4, line 18: Add “latitudinal gradient” to the sentence

Action

Done as suggested, see line 150

Reviewer comment 35

Page 10, line 8-10: I suggest to start the Results section with this sentence

Reply

Thank you for the suggestion but we prefer to keep the sentence as a closing sentence

Reviewer comment 36

Page 10, line 16: net autotrophic?

Reply

This entire section of the discussion was modified

Reviewer comment 37

Page 12, line 8: Please stay consistent, use R2.

Reply

This section was modified

Reviewer comment 38

Page 13, line 9: Heterotrophic suggest no autotrophs, add “net”

Reply

This section was modified

Reviewer comment 39

Page 15, line 6-9: Please rewrite.

Reply

Modified as indicated, please see lines 719–722

Reviewer comment 40

Page 16, line 4: Add i.e. or parentheses after 2.5 _C

Action

We modified the text in the section but indicated the range inside a parenthesis (lines 732–733)

Reviewer comment 41

Page 19, line 6: Heterotrophic

Action

Typographical error corrected, see line 939

Reviewer comment 42

Figure A1: Add axis titles to every part of the figure, having double axes without titles is confusing, especially since the 27 _N axis title (Temperature) is not on any axis.

Action

We are not including this figure in the new version of the manuscript

Reviewer comment 43

Table 1: Add Silicate to the table description. Also, it is unclear which header belongs to which environmental variable. Also, I fail to see the benefit of the min and max values

Action

We are not including this table in the new version of the manuscript

Reviewer comment 44

Present data as mean +/- SE

Action

We clarified this point in line 322. Whenever we presented mean values, we added the standard error of the mean

Reviewer comment 45

Table 2: N does not need decimals

Action

We are not including this table in the new version of the manuscript

Reviewer comment 46

What does “rank” mean? % PAR differs from Table 1

Action

We are not including this table in the new version of the manuscript. Data from table 2 are now being presented in a figure (new figure 5) with the overall information by seasons, but with all the data points included in the study.

Reviewer comment 46

Table 3: Upper part are, what seems to be, Pearson rank coefficients, not the units given in the description. The lower part seems to be p-values, mention this in the description. A hyphen is not the same as a blanc.

Reply

Thank you for pointing this out. It is indeed a correlation matrix.

Action

We modified the table (now Table 1) and only presented the correlation of metabolic rates with main explanatory variables.

Reviewer 2

2.2.1 General comments

The authors quantified plankton metabolic rates along the Red Sea. They have shown that Chla and plankton community metabolism (GPP and CR) increase with temperature. Contrary to previous results they have observed a higher Activation Energy for GPP than for CR showing a positive relationship between NCP and Temperature. These results have been explained by the authors as a consequence of the high nutrient availability in warmer waters and the lack of external organic carbon sources to sustain a heterotrophic metabolism constraining the CR.

The dataset are very interesting and merit been published, however, the way how the results have been presented, the lack of statistical analyses and the methodology proposed

are not the most suitable to achieve the main goal proposed in the manuscript.

Therefore, I consider the ms still needs major revision in order to be published and providing the authors follow the reviewers recommendations.

2.2.2 Reply to general comments

Reply

Thank you for your comments and suggestions. We have addressed the changes and recommendations of the reviewer and provide a detailed answer to each of the points made in the following section. First, we completely agree with the reviewer that a detailed

description of the statistical analyses performed was missing in the methodology section. We have included a detailed description of the statistical analyses in a new section (2.4), and this change can be tracked now between lines 288–323 in the latest version of the manuscript. Regarding the primary concern of Reviewer 2, which was also pointed out by the first referee, related to the methodological approach we used to quantify planktonic metabolic rates, we think, there is a misunderstanding. The methodology used to quantify planktonic metabolism is based on the extensively-used dark and light method (in combination with the Winkler titration method). The reviewer indicated that the methodology used was not suitable, and suggested that a shorter incubation period (6–12 h) was more appropriate to quantify NCP. We want to point out that NCP represents the organic matter remaining after consumption of the GPP through respiration by plants (autotrophs), microbes (either autotrophs or heterotrophs), and animals (heterotrophs) (Ducklow and Doney 2013), and to account for those process, the standard incubation time for *in vitro* incubations is 24 h. This incubation length is needed because contrary to photosynthesis, which can be resolved during daylight, the losses due to respiration (which are necessary to define NCP) also occurs at night.

Action

We have clarified this point in the methods section 3.3 (between lines 200–208). In addition, we made sure that a detailed description of the sampling, filling and fixing of the samples was provided. We also clarified how we calculated gross primary production, the community respiration and the net community production (which is the balance between the autotrophic and heterotrophic metabolism, and not only the result of photosynthetic activity). These calculations can be seen between lines 256 and 277.

2.2.2 Specific comments

Reviewer comment 1

First, according to the title and the abstract the authors consider as drivers of the plankton community metabolism in the Red Sea, the Chla and temperature. However, other important parameters such as, temporal and spatial variability, salinity and nutrients seem to govern the plankton community metabolism within this particular ecosystem and are not included in the abstract. Therefore, this lack of agreement between the ms, the conclusion and the abstract. is confusing. In my opinion, there is a large floor in the experimental design proposed and it is difficult to resolve.

Reply

We appreciate the reviewer's comment, and agree that the abstract highlighted our main findings and did not detail all the results. The abstract was indeed mostly orientated towards the effect of temperature and nutrient availability on metabolic rates as we found that those were the main controlling drivers. That was consistently explained in our results, discussion and conclusion; therefore, we do not find disagreement in our statements. However, following the suggestion of referee #1, we changed the narrative of the text to focus only on the variables highlighted in the abstract (i.e., temperature and nutrient availability; which are tightly related to the latitudinal gradient that characterises the basin) leaving aside the changes of planktonic metabolism throughout the water column.

Action

The changes can be tracked in the new version of the manuscript in the result section between lines 322–448. We have modified the figures accordingly so that Figure 2 now describes the latitudinal gradient of variables only in surface waters. In Figure 3 we have modified the format and the data presented, so now we only centre on data within the photic layer. We removed Figures 4–5, Figure 1A and 2A. We also removed Table 1, and Table 2. The information from Table 2 is now presented as a figure (now Figure 5) with the overall information and with all the data plotted. The changes in the results are in agreement with changes in the narrative of the discussion section. The main changes can be tracked between lines 574–580 and between lines 584–588 of the new manuscript.

Reviewer comment 2

All samples included the deepest ones have been incubated on deck with surface water. During some of the surveys there is an important thermal variability. The authors have attempted to mitigate the issue by including just those samples above the thermocline. However, Material and Methods mention that changes in temperature and PAR in the incubation tanks were recorded with HOBO data loggers. Therefore, those data should be shown in a table in order to select objectively the samples for the analyses. Hence, eliminating those samples that register thermal differences above 2 °C with the in situ temperatures. In addition, samples adapted to cool temperatures such as those at the bottom will respond more drastically to artificial increments of temperature than surface ones (for example. Apple et al. 2006. AME. 43: 243–254) resulting in erroneous conclusions. Therefore, Figure A1 is important and should be included in the Ms.

Reply

As mentioned in our previous answer, after careful consideration, we decided to focus on the overall patterns observed in planktonic metabolism leaving aside the variability that takes place through the water column.

Action

We removed Figure A1 from the current version of the manuscript; however, all metabolic rates are now shown in the newly generated Figure 5.

Reviewer comment 3

Other figures such as 4-6 do not show crucial information in the current format. Figure 3 and Table 3 to me are redundant

Action

We removed Figures 4–6 from the current version of the manuscript, and the overall results are now presented as a new figure (Figure 5). We modified Table 3 (now Table 1) to only show complimentary information to Figure 3.

Reviewer comment 4

The paragraph 10-15 page 6 the authors should indicate if samples were collected before sunrise (to avoid any light on the samples) and if the incubation started at the sunrise to estimate the full light period. The authors say, the samples were collected between 7 to 9 and to me this sounds very late to incubate and obtain the full light period nor precisely.

Reply

The samples were incubated for 24-h, covering an entire dark-light period, thus there is no need to estimate the light period.

Reviewer comment 5

In The net community metabolism..... page 7, NCP should be estimated during the light period (NCP 6 to 12 hours).

Reply

We believe that there is a misunderstanding regarding the process we measured. The reviewer's comment seemed to suggest that our work was focused on primary production, which is performed by the photosynthetic components of the plankton community during the daytime and that has gross and net components (as phytoplankton excrete and respire carbon). However, our paper focuses on the entire plankton community, both photosynthetic and heterotrophic (e.g. bacteria), where the net community production (NCP) represents the organic matter remaining after consumption of the GPP through respiration by plants (autotrophs), microbes (either autotrophs or heterotrophs), and animals (heterotrophs) (Ducklow and Doney 2013). Studies that focus on the photosynthetic component of the plankton community (e.g. Net Primary Production, NPP) report values for the daylight period only. Whereas studies, such as this, report (24 h) rates. For instance, published syntheses of community metabolism rates report values per day (24 h), e.g., Robinson and Williams 2005, Regaudie-de-Gioux 2012, 2013). The use of 24 h to report rates is justified as the metabolic budget needs to be resolved over 24 h to be completed, not required for

photosynthesis as it is light-dependent, but respiration, which is necessary to define net community production, which occurs both during day and at night.

Reviewer comment 6

The authors should show, the variation coefficient of the pool data and also the original CR, NCP and GPP data including their SE.

Action

The data are now shown in Figure 5, with a detailed description of the statistical analyses. In addition, the relevant information regarding the mean and the standard error of the mean are shown. See e.g., lines 386, 387, 435.

Reviewer comment 7

Because, In these oligotrophic areas the metabolic rates are very low and can be difficult to detect. Therefore, the methodology needs to be very precise in the processes of filling, incubating and fixing the bottles.

Reply

We agree with the reviewer, however, the information about filling the bottles and all special cares during the sampling is detailed in methods section 2.3 (Page 7: lines 11–15). Now between lines 242–250.

Reviewer comment 8

The paragraph 20 in page 8 It should be indicated the Arrhenius plots the authors Mention

Reply

The Arrhenius plots were described in the methods section 2.3 (P8, between lines 13 and 20) and additionally explained in the results section 3.3 (P11: lines 16 and P12 lines 1–3) and Figures 9 and 10.

Action

The temperature-dependence of planktonic metabolism explained with the Arrhenius plots can now be seen between lines 463–471 and in Figures 7 and 8.

Reviewer comment 9

The paragraph 10 in page 10 should be transferred from the Results to the Discussion.

Reply

The sentence on P10, between lines 8–10 is a closing statement with the main results shown in previous paragraphs, we are not discussing any results. Therefore, we prefer to keep it as it is.

Reviewers comment 10

And also the first paragraph of the 3.2 Variability of plankton : : . Is already mentioned in M and M.

Reply

We agree with the reviewer's comment and have modified the text accordingly

Action

We have deleted this sentence

Reviewers comment 11

The name of the KAUST is excessive. I would use just one larger map with different colours or shapes to show the stations at each survey or season.

Reply

We decided to not modify the figure as some of the stations were sampled in the same location more than twice, and different shapes or points overlap

Action

We did remove the name of the university to leave it only in one panel

Reviewers comment 12

Figure 2. I consider in this figure is difficult to detect the thermocline and the vertical profiles of Chla and salinity. I consider that nutrient profiles should also included

Reply

It was not possible to determine the depth of the thermocline in Figure 2, and it was not intended to do so. The figure summarises the main characteristics along the latitudinal axis that we sampled (i.e., the increasing temperature and phytoplankton chlorophyll-a toward the southern region with an increasing salinity towards the north).

Action

We modified Figure 2, so that now it only shows the variability of temperature, salinity and chlorophyll-a in surface waters.

Reply to reviewer 13

Figure 8, 9 and 10. To test one of the main conclusions, if AE is higher for GPP than for CR, authors should test statistically if the slopes are different. I would test also the slopes for the figures 9 and 10 explaining the consequences of the statistical differences in the cases observed

Reply

We performed an analysis of covariance (ANCOVA)

Action

We have mentioned this in section 2.4 between lines 318–320 and described the results between lines 467–471.

Reply to reviewer 14

In the figure 9, the RMA analyses have been included but it is not necessary in this case because temperature is not a rate. In addition, the authors have not explained when the RMA or OLS should be used in M and M.

Reply

We agree with the reviewer that it is not necessary to provide the results of the RMA analyses

Action

We removed these analyses and their results from our manuscript

1 **Rates and drivers of Red Sea plankton community metabolism**

2 Daffne C. López-Sandoval¹, Katherine Rowe¹, Paloma Carillo-de-Albonoz¹, Carlos M. Duarte¹ and

3 Susana Agusti¹,

4 ¹ Red Sea Research Center, King Abdullah University of Science and Technology (KAUST), Thuwal-Jeddah, 23955-6900,

5 Saudi Arabia

6 *Correspondence to:* Daffne C. López-Sandoval (daffne.lopezsandoval@kaust.edu.sa)

7 **Abstract**

8 Resolving the environmental drivers shaping planktonic communities is fundamental to understanding
9 their variability, in the present and the future, across the ocean. More specifically, resolving the
10 temperature-dependence of planktonic communities is essential to predict the response of marine
11 ecosystems to warming scenarios, as ocean warming leads to oligotrophication of the subtropical ocean.

12 Here we quantified plankton metabolic rates along the Red Sea, a uniquely oligotrophic and warm
13 environment, and analysed the drivers that regulate gross primary production (GPP), community
14 respiration (CR) and the net community production (NCP). The study was conducted on six

15 oceanographic surveys following a north-south transect along the Saudi Arabian coast. Our findings
16 revealed that GPP and CR rates increased with increasing temperature ($R^2= 0.41$ and 0.19 , respectively,
17 $p < 0.001$ in both cases), with a higher activation energy (E_a) for GPP (1.2 ± 0.17 eV) than for CR (0.73
18 ± 0.17 eV). The higher E_a for GPP than for CR resulted in a positive relationship between NCP and
19 temperature. This unusual relationship is likely driven by 1) the relatively higher nutrient availability
20 found towards the warmer region (*i.e.*, the South of the Red Sea), which favours GPP rates above the
21 threshold that separates autotrophic from heterotrophic communities ($1.7 \text{ mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$) in this
22 region. 2) Due to the arid nature, the basin lacks riverine and terrestrial inputs of organic carbon to
23 subsidise a higher metabolic response of heterotrophic communities, thus constraining CR rates. Our

Style Definition: Heading 4: Font: 12 pt, Not Bold

Style Definition: MS title: Font: 12 pt, Not Bold

Formatted: Section start: Continuous, Numbering: Continuous

Formatted: Font: 17 pt, Bold, Font color: Text 1

Formatted: Font color: Text 1

Deleted: Agusti¹

Formatted: Font color: Text 1

Deleted: in low productive waters

Deleted: unique

Deleted: coasts

Deleted: Chl-*a* specific

Formatted: Superscript

Deleted: P

Deleted: AE

Deleted: ±

Formatted: Font: norma

Deleted: AE

Deleted: and

Deleted:).

35 study demonstrates that GPP increases steeply with increasing temperature in the warm ocean when
36 relatively high nutrient inputs are present.

Formatted: Font color: Text 1

37 1 Introduction

Formatted: Font color: Text 1

38 The balance between gross primary production and community respiration, which involves both
39 autotrophic and heterotrophic metabolic activity (Williams, 1993; Cullen, 2001; Ducklow and Doney,
40 2013), sets the metabolic status of an ecosystem by defining the carbon available to fuel pelagic food
41 webs and determining whether plankton communities act as a source or sink of CO₂ (Del Giorgio et al.,
42 1997; Williams, 1998). Whereas GPP typically satisfies the respiratory demands within the food web
43 across productive waters, the oligotrophic ocean often requires allochthonous inputs of organic carbon
44 to meet the metabolic requirements of heterotrophic organisms (Smith and Mackenzie, 1987). Due to
45 comparatively higher carbon consumption, relative to the production, planktonic communities in low
46 productive systems are in close metabolic balance (i.e., the net community production (NCP = 0, or
47 GPP = CR) or experience a net metabolic imbalance (i.e. NCP < 0, GPP < CR) (Smith and Hollibaugh,
48 1993; Duarte and Agustí, 1998; Duarte et al., 2013).

Deleted: Plankton community metabolism, the

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted:), defi

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: determines

Formatted: Font color: Text 1

Deleted: (Duarte and Agustí, 1998; Williams, 1998; Duarte et al., 2011).

Formatted: Font color: Text 1

Deleted: (Duarte et al., 2013).

Formatted: Font color: Text 1

Deleted: (GPP),

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: metabolism,

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: , which are a characteristic of these systems,

Formatted: Font color: Text 1

Deleted: with increasing temperature

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: subsidize

Formatted: Font color: Text 1

49 In tropical and subtropical oligotrophic regions, the high temperatures may amplify the
50 metabolic imbalances in plankton communities, as CR tends to increase faster than GPP (Harris et al.,
51 2006; Regaudie-de-Gioux and Duarte, 2012) if the allochthonous sources of organic carbon are enough
52 to subsidize their carbon demand. These allochthonous inputs may be delivered from land through
53 riverine discharge, from the atmosphere through atmospheric deposition of dust and volatile organic

65 carbon (Jurado et al., 2008), or are exported from productive coastal habitats (Duarte et al., 2013;
66 Barrón and Duarte, 2015).

67 The Red Sea is a semi-enclosed highly oligotrophic basin (Acker et al., 2008; Raitzos et al.,
68 2013). It is known as one of the warmest tropical seas, with maximum sea surface temperatures ranging
69 from 33.0 to 33.9 °C during the summer period (Chaidez et al., 2017; Osman et al., 2018), and between
70 34–35 °C in certain regions (Rasul and Stewart, 2015; Garcias-Bonet and Duarte, 2017; Almahasheer et
71 al., 2018). Due to the prevailing arid conditions, the Red Sea experiences large evaporation rates (nearly
72 2 cm yr⁻¹ of freshwater from the surface layers), while the lack of river runoff and low precipitation
73 rates make this system one of the saltiest seas on the planet (Sofianos, 2002; Sofianos and Johns, 2015;
74 Zarokanellos et al., 2017). Two wind patterns govern the region: in the northern part, the wind coming
75 from the northwest remains relatively constant throughout the year, while in the southern area, the
76 Indian Monsoon system regulates the wind dynamics (Sofianos, 2002; Sofianos and Johns, 2015).
77 During the winter monsoon, the wind changes direction, and this wind reversal along with the
78 thermohaline forces drives the overall circulation and favours the exchange of water with the Indian
79 Ocean (Sofianos, 2002; Zarokanellos et al., 2017).

80 Due to the almost negligible terrestrial inputs, the intrusion of nutrient-rich waters from the
81 Indian Ocean through the Bab-el-Mandeb Strait (Sofianos and Johns, 2007; Raitzos et al., 2015; Kürten
82 et al., 2016), together with aeolian dust and aerosol deposition (Chen et al., 2007; Engelbrecht et al.,
83 2017), represent the primary sources of nutrients into the basin. Thus, nutrient availability in the Red
84 Sea follows a latitudinal pattern that is opposite to the one of salinity, but parallel to the thermal

- Formatted: Font color: Text 1
- Deleted: ,
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Deleted: -
- Formatted: Font color: Text 1
- Deleted: As a result
- Formatted: Font color: Text 1
- Deleted: its desert
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Deleted: In
- Formatted: Font color: Text 1
- Deleted: through
- Formatted: Font color: Text 1
- Deleted: .
- Formatted: Font color: Text 1
- Deleted: some of
- Formatted: Font color: Text 1

93 gradient, with nutrient-rich and warmer waters towards the Southern Red Sea compared to the cooler
94 and more oligotrophic Northern Red Sea (Sofianos, 2002; Raitso et al., 2015).

95 Studies based on ocean color data revealed that chlorophyll-a (Chl-a) concentrations decline
96 from the Southern Red Sea to the Northern Red Sea (Raitso et al., 2013; Kheireddine et al., 2017;
97 Qurban et al., 2017) and depict a clear seasonality. During winter time, when the maximum exchange of
98 water with the Indian Ocean takes place, Chl-a concentration peaks, decreasing towards the summer
99 period when the water column is mostly stratified (Sofianos, 2002). Measurements of primary
100 production also revealed that phytoplankton photosynthetic rates follow the same south to north
101 gradient as Chl-a and nutrient concentration (Qurban et al., 2017). However, reports regarding the
102 metabolic balance of the plankton communities are scarce, mostly focus on the contribution of the
103 autotrophic community via photosynthetic processes (Levanon-Spanier et al., 1979; Qurban et al., 2014;
104 Rahav et al., 2015), or are restricted to specific regions (Tilstra et al., 2018).

105 Based on available evidence, we hypothesise that the high gross primary production expected in
106 the Southern Red Sea may be counterbalanced by a higher respiratory demand in these warm waters
107 and that NCP might decline towards the relatively unproductive waters of the Northern Red Sea. With
108 the expected decrease in GPP towards the northern region, planktonic metabolism might be driven
109 mainly by heterotrophic communities (Duarte and Agustí, 1998; Duarte et al., 2013). However, the
110 absence of significant allochthonous subsidies in the basin may hamper the metabolic response of the
111 heterotrophic plankton communities. Hence, it remains unclear what the metabolic balance of plankton
112 communities is and whether a south to north latitudinal gradient in NCP exists in the Red Sea.

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: ¶

Formatted: Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Deleted: However, there is no information regarding the

metabolism of the plankton communities.

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Deleted: : (1)

Formatted: Font color: Text 1

Deleted: (Harris et al., 2006; Regaudie-de-Gioux and Duarte,

2012), and that (2) NCP might decline towards the

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

119 Here we report the variability of plankton community metabolism (GPP, CR and NCP) along a
120 latitudinal gradient in the Red Sea, and examine if the temperature-dependence of planktonic metabolic
121 rates in this basin are consistent with those reported for the global ocean (López-Urrutia et al., 2006;
122 Regaudie-de-Gioux and Duarte, 2013; Garcia-Corral et al., 2017). We did so by measurements
123 conducted as part of six surveys along the south-north latitudinal gradient in the Saudi Economic
124 Exclusive Zone in Red Sea waters. We determined plankton metabolic rates between winter 2016 and
125 spring 2018, thus allowing us to 1) delineate the seasonal variability of the gross primary production
126 and community respiration along the Red Sea, 2) quantify changes in the metabolic balance (net
127 community production) and 3) test the hypothesized roles of productivity gradients and temperature in
128 driving NCP.

Formatted: Font color: Text 1

Deleted: in

Formatted: Font color: Text 1

Deleted: at three different optical depths

Formatted: Font color: Text 1

Deleted: (GPP)

Formatted: Font color: Text 1

Deleted: (CR)

Formatted: Font color: Text 1

Deleted: , NCP

Formatted: Font color: Text 1

134 2. Material and Methods

135 2.1 Field Sampling

136 We conducted six oceanographic surveys; two during autumn (October and November 2016),
137 two during winter (February 2016 and January 2017), one in summer (August 2017), and one in spring
138 (March 2018) on board the R/V *Thuwal* and R/V *Al Azizi*. Sampling was conducted following a
139 latitudinal transect along the Red Sea within a region limited by coordinates 17.25 °N to 27.82 °N and
140 34.83 °E to 41.39 °E (Figure 1). At each station, vertical profiles of temperature and salinity were
141 obtained with a Sea-Bird SBE 911 plus CTD profiler (Sea-Bird Electronics, Bellvue, WA, USA),
142 equipped with additional sensors to measure the attenuation of photosynthetically active radiation
143 (PAR) (Biospherical/Licor PAR/Irradiance Sensor), *in vivo* fluorescence (WetLabs ECO FL
144 fluorometer), and dissolved oxygen concentration (Seabird SBE 43 Dissolved Oxygen Sensor). Water
145 samples for chemical and biological measurements were collected between 7:00 and 9:00 am local time,
146 using a rosette sampler equipped with 12 Teflon Niskin bottles (12 L) that were provided with silicone
147 O-rings and seals.

148 2.2 Inorganic nutrients and chlorophyll-a concentration

149 Water samples for nutrient analyses were collected in 50 mL polyethene bottles and kept frozen
150 (-20 °C) until determination. Inorganic nutrient concentration was determined with a SEAL AA3
151 Segmented Flow Analyzer (SEAL Analytical Inc., WI, USA) using standard methods (Hansen and
152 Koroleff, 1999). The detection limits were 0.05 μM for nitrate, 0.01 μM for nitrite, 0.01 μM for

Deleted: [

Formatted: Font color: Text 1

Deleted:]

Formatted: Font color: Text 1

Deleted: Fig. 1), sampling between five and seven stations per survey....

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Deleted: nutrients analysis

Formatted: Font color: Text 1

Deleted: (Hansen and Koroleff, 1999).

Formatted: Font color: Text 1

159 phosphate and 0.08 μM for silicate. For the chlorophyll-*a* analysis, 200 mL samples were taken at ten
160 discrete depths (between 5 and 200 m) and filtered through Whatman GF/F filters. The filters were kept
161 frozen (-20 °C) until further analysis. Pigments were extracted for 24 h using 90 % acetone and left
162 overnight in the dark at 4 °C. Chl-*a* concentration was estimated with the non-acidification technique
163 using a Trilogy Fluorometer equipped with CHL-NA module (Turner Designs, San Jose, USA),
164 previously calibrated with pure Chl-*a*.

165 2.3 Net community metabolism, community respiration and gross primary production

166 Plankton metabolic rates were determined *in vitro* by measuring the changes in dissolved
167 oxygen concentration after 24 h light-dark bottle (Winkler) incubations (Carpenter, 1965). This
168 methodology, commonly used to determine plankton metabolic rates (Williams et al., 1979; Duarte and
169 Agustí, 1998; Bender et al., 1999; Robinson and Williams, 1999; Ducklow et al., 2000; Serret et al.,
170 2001; Robinson et al., 2002; Serret et al., 2009; García-Martín et al., 2017), allows to account for the
171 diel cycle of oxygen and carbon fluxes derived from photosynthetic mechanisms (light-dependent
172 reactions) and also those linked to the acquisition of energy by both autotrophic and heterotrophic
173 microorganisms (light and dark-dependent reactions) (Robinson and Williams, 2005; Williams and del
174 Giorgio, 2005).

175 Water samples were collected at three different optical depths (ζ) through the water column.
176 One at the surface (100–80 % of incident PAR), another towards the bottom of the photic layer (8–1 %
177 of incident PAR), and one intermediate sample, at a depth of the chlorophyll maximum (Chl-*a* max). In
178 case the Chl-*a* max was sampled at the surface or bottom layers, the intermediate sample was taken

Formatted: Font color: Text 1

Deleted: (Chl-*a*),

Formatted: Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: (-20 °C).

Deleted: for 24 h

Formatted: Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: Planktonic

Formatted: Font color: Text 1

Deleted: by

Formatted: Font color: Text 1

Deleted: during

Formatted: Font color: Text 1

Formatted: Font color: Text 1, English (UK)

Deleted: We selected

Formatted: Font color: Text 1

Deleted: ,

Deleted: of them

Formatted: Font color: Text 1

Formatted: Font color: Text 1

188 between 1.5–2.3 ζ (i.e., 22–10% of incident PAR). Seawater was collected directly from the Niskin
189 bottles to fill a total of 21 (100 mL) Winkler bottles. The bottles were carefully filled using silicone
190 tubing and allowing the water to overflow during the filling, taking special care to avoid the formation
191 of air bubbles. Surface samples were collected in 100 mL quartz bottles. From each depth, seven of the
192 bottles were immediately fixed with Manganese sulfate (MnSO_4) and Potassium hydroxide/Potassium
193 iodide solution (KI/KOH) to determine the initial oxygen concentration while the other 14, seven light
194 and seven black bottles, were incubated on deck in surface water flow through tanks. Due to the
195 difference in temperature between the surface and deep waters, particularly during the summer and
196 autumn surveys, we decided to include in our analyses only those samples collected above the
197 thermocline. Changes in temperature and PAR in the incubation tanks were recorded with HOBO
198 Pendant data loggers (Onset, Massachusetts, USA).

199 Before the incubation, the bottles were covered with neutral mesh to reduce the incident PAR
200 radiation according to the sampled depth. At the end of the incubation period, light and dark bottles
201 from each depth were fixed to determine final O_2 concentrations. Oxygen concentration was measured
202 by automated high-precision Winkler titration with a potentiometric end-point detection (Oudot et al.,
203 1988) using a Mettler Toledo T50 Titration Excellence auto-titrator attached to an Inmotion
204 autosampler. NCP was calculated as the difference in the oxygen concentration between the light bottles
205 after the 24 h incubation period ($[\text{O}_2]_{\text{L-24h}}$) and the oxygen concentration measured before the
206 incubation ($[\text{O}_2]_{\text{Tzero}}$) (i.e., $\text{NCP} = ([\text{O}_2]_{\text{L-24h}} - [\text{O}_2]_{\text{Tzero}})$). CR rates ($\text{mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$) were calculated as
207 the difference of the oxygen concentration after the 24 h incubation period in the dark bottles ($[\text{O}_2]_{\text{D}}$).

Deleted: . 100 %, 60–20 % and 8–1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: (100 % PAR)

Formatted: Font color: Text 1

Deleted:

Formatted: Font color: Text 1

Deleted: was

Formatted: Font color: Text 1

Deleted: ¶

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: Net community metabolism (NCP, $\text{mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$)

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: . Community respiration

Formatted: Font color: Text 1

Deleted: CR,

Formatted: Font color: Text 1

216 $_{24h}$) and the initial oxygen concentration ($[O_2]_{Tzero}$) (i.e., $CR = [O_2]_{Tzero} - ([O_2]_{D-24h})$). GPP ($mmol O_2 m^{-3} d^{-1}$) was calculated as the sum of NCP and CR.

218 Due to the consistent relationship existing between plankton metabolism and temperature across
219 diverse marine regions (Regaudie-de-Gioux and Duarte, 2012; García-Corral et al., 2014), we examined
220 how plankton metabolic rates covariate with temperature in the Red Sea, a system whose temperature
221 range is higher than previously encountered in marine planktonic metabolism research. We determined
222 the relationship between metabolic rates and temperature by fitting an ordinary least squares linear
223 regression equation to the relationship between the natural logarithm of the Chl- a specific metabolic
224 rates and the inverse of the absolute temperature * k , which is the Boltzmann's constant ($8.617734 * 10^{-5}$
225 eV K $^{-1}$). In these Arrhenius plots, the slope represents the average activation energy (E_a), characterising
226 the extent of thermal-dependence of metabolic processes.

227 2.4 Statistical Analyses

228 Statistical analyses and figures were done using the statistical and machine learning toolbox in
229 Matlab version R2018b (Mathworks Inc, Natick, MA, USA) and with the R statistical computing
230 package using RStudio 1.1419. Pearson correlation tests were used (corrplot function in R) to determine
231 the relationship between environmental variables (temperature, nitrate + nitrite (NO $_x$), phosphate and
232 silicate concentration) and their latitudinal distribution, and to determine the relationship between
233 volumetric measurements of GPP, CR, NCP, and environmental variables (Temperature, NO $_x$
234 concentration, Chl- a , and latitude). We used ordinary least squares (OLS) simple regression models
235 (fitlm function in Matlab) to describe the potential relationships between different planktonic metabolic

Deleted: . Gross primary production (

Formatted: Font color: Text 1

Deleted: ,

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted:

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: OLS (Ordinary

Formatted: Font color: Text 1

Deleted:)

Formatted: Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Deleted: E_a

Formatted: Font color: Text 1

Formatted: Font: 12 pt, Font color: Text 1

Formatted: Heading 2, Indent: First line: 0 cm

Deleted: Statistics

Formatted: Font: 12 pt, Font color: Text 1

Deleted: analysis was done using GraphPad Prism 7 (GraphPad Software, La Jolla, CA, USA) and MVAapp_v2.0 (Julkowska et al., MVApp.pre-release_v2.0 mmjulkowska/MVApp: MVApp.pre-release_v2.0. DOI: 10.5281/zenodo.1067974).

Formatted: Font: 12 pt, Font color: Text 1

248 rates, between metabolic rates and environmental variables, and to predict the response of the Chl-a
249 normalised GPP (and CR) to temperature (Arrhenius plots described in section 2.3). To test if the
250 activation energies (obtained from the Arrhenius plots) were significantly different, we performed an
251 analysis of covariance (ANCOVA) by using the aocool in Matlab. The variability of planktonic
252 metabolic rates between cruises was statistically analysed using non-parametric Kruskal-Wallis tests.
253 Mean values and their standard error of the mean (SE) are reported throughout the text.

254 3. Results

255 3.1 Latitudinal variability of physico-chemical properties and Chl-a concentration

256 Hydrographic (temperature and salinity) and chemical variables (nutrient concentrations)
257 depicted a marked latitudinal gradient typical of the Red Sea. At the southern-most area, sea surface
258 temperature (SST) fluctuated between 28 °C (winter-spring) and 32 °C (summer), while at the far-
259 northern sampling site SST ranged between 23 °C (winter) and 27–28 °N (summer-autumn) (Figure 2).
260 Overall, all macronutrients observed a significant inverse correlation with latitude (Pearson correlation
261 coefficients $r < -0.4$, $p < 0.05$) (Figure 3). Nitrite+nitrate (NO_x) decreased from $6.1 \pm 0.9 \mu\text{M}$ in the
262 southern region to $2.9 \pm 0.3 \mu\text{M}$ towards the northern Red Sea, while on average, phosphate
263 concentration ranged from $0.5 \pm 0.01 \mu\text{M}$ in the south of the Red Sea to $0.1 \pm 0.01 \mu\text{M}$ towards the
264 northern stations (data not shown). Phytoplankton biomass (measured as Chl-a concentration) also
265 decreased significantly towards the north of the Red Sea (Pearson's correlation, $r = -0.41$, $p < 0.001$)
266 (Table 1). We found the highest autotrophic biomass during the autumn and winter cruises. During this

Formatted: Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Deleted: The

Deleted: profiles revealed strong

Deleted: and seasonal variation in the structure

Deleted: water column along the basin (Figures 2 and A1). During the summer

Deleted:), ranged from 32.5 °C, at the southern-most station (17 °N), to 29.3 °C

Deleted: (27 °N

Deleted: During this survey, the

Deleted: of dissolved inorganic nitrogen (DIN, NO₂ + NO₃) decreased...

Deleted: 61 μM (±

Deleted: 11) (at

Deleted: first optical depth)

Deleted: 12 μM (± 0.04) between 15–5 % of PAR, while phytoplankton chlorophyll-a concentration remained ~ 0.2 μg Chl-a l⁻¹ (Table 1). In Autumn, SST ranged from 31.2 °C (southern-most station) to 27.4 °C (northern-most station) (Figure 2), while nutrient availability and

Formatted: Font: Not Italic

Deleted: increased, particularly at the southern stations where phytoplankton Chl-a concentration was above 0.8 μg l⁻¹ in the surface layers (Figure 2, Table 1). During winter,

Deleted: water column generally remained well mixed, with temperatures fairly similar along

Deleted: upper 100 m (Figure A1).

292 period, surface Chl-a ranged from 0.6–0.8 mg m⁻³ in the southern region to 0.2–0.3 mg m⁻³ in the north
293 (Figure 2). In general, our results confirm that all variables correlated significantly with latitude,
294 highlighting the prevalence of the south-north gradient in temperature, salinity, nutrient availability and
295 chlorophyll-a concentration across the Red Sea.

296 3.2 Variability of plankton metabolism measured along the Red Sea

297 Analogous to the environmental variability, planktonic metabolism followed the same
298 significant north-south decreasing pattern with latitude (Figure 4). The inverse correlation of GPP rates
299 with latitude was highly significant (Pearson correlation coefficient $r = -0.60$, $p < 0.001$) (Table 1), as
300 found for autotrophic biomass, thus, explaining the strong correlation observed between GPP and Chl-a
301 concentration (Pearson correlation coefficient $r = 0.69$) (Table 1). GPP rates decreased on average by
302 79%, from 4.1 ± 0.5 mmol O₂ m⁻³ d⁻¹ (≈ 49.2 mgC m⁻³ d⁻¹; assuming a photosynthetic quotient, PQ = 1)
303 at the southernmost station of the Red Sea to 0.9 ± 0.1 (≈ 10 mgC m⁻³ d⁻¹; PQ = 1) at the northern site,
304 while CR decreased on average by 73 %, from 3 ± 0.4 mmol O₂ m⁻³ d⁻¹ (≈ 36 mgC m⁻³ d⁻¹; assuming a
305 respiratory quotient, RQ = 1) in the south to 0.8 ± 0.1 in the north (≈ 9.6 mgC m⁻³ d⁻¹; RQ = 1) (Figure
306 4). We did not find any significant correlation between NO_x availability and GPP (Pearson correlation
307 coefficient, $r = 0.01$, $p > 0.05$). CR (Pearson correlation coefficient, $r = 0.19$, $p > 0.05$) nor NCP rates
308 (Pearson correlation coefficient, $r = -0.19$, $p > 0.05$) (Table 1); however, all metabolic rates were
309 positively correlated with temperature (Table 1).

310 The highest GPP and CR rates measured along the Red Sea came from data collected during the
311 autumn and winter cruises, when GPP and CR rates reached values above 6 and 4 mmol O₂ m⁻³ d⁻¹,

Deleted: Chl-a concentration along the basin peaked (Figure 2) while DIN, phosphate and silicate concentration remained similar within the first two optical depths (Table 1).
We identified two distinctive patterns portraying nutrient distribution along the Red Sea. Nutrient concentration throughout the water column decreased with latitude (Figure 3), while DIN concentration increased with temperature in the first optical depth (37 % PAR) and at the base of the euphotic zone (1–0.1 % PAR) (Figure 3A and 3B). Phosphate and silicate concentrations were also positive correlated with temperature at the base of the photic layer (Pearson correlation, DIN, $r = 0.39$, $P = 0.01$; phosphate, $r = 0.48$, $P < 0.001$; silicate, $r = 0.42$, $P < 0.001$) (Figure 3B). Below the first optical depth and above the base of the euphotic zone, nutrient availability and temperature were not correlated (data not shown).

Deleted: to

Formatted: Font: Not Italic

Formatted: Font color: Auto

Deleted: ¶

Formatted: Font color: Text 1

Deleted: The measurements of plankton metabolic rates taken from the six oceanographic surveys (between winter 2016 and spring 2018) allowed us to define the general variability patterns of gross primary production (GPP) and community respiration (CR) along the Red Sea (Figures 4 and 5, Table 2). Plankton communities were autotrophic when all surveys were taken in concert, with heterotrophic communities representing 38 %, 32 % and 56 % of the communities assessed between 100–37, 36–6, and 5–1 % PAR, respectively (Table 2). Part of the variability in community metabolism was explained by seasonal differences, as plankton communities tended to be mostly heterotrophic (80 % of NCP < 0) along the basin during spring, while between summer and winter, NCP rates < 0 were mostly restricted to the northern part of the Red Sea (above 21 °N) (Figure 6), GPP (and CR) rates peaked in autumn and winter (3–8 mmol O₂ m⁻³ d⁻¹) in the southern stations, while in the stations sampled towards the north GPP rates remained ~ 1.5 mmol O₂ m⁻³ d⁻¹ throughout the year (Table 2, Figure 4). Plankton metabolic rates were independent of dissolved inorganic nitrogen concentration ($P = 0.99$, 0.47, and 0.43 for GPP, CR, and NCP, respectively), but increased significantly with increasing Chl-a concentration ($R^2_{(GPP)} = 0.50$, $R^2_{(CR)} = 0.45$ and $R^2_{(NCP)} = 0.14$, $P < 0.001$ in all cases) and temperature ($R^2_{(GPP)} = 0.23$, $R^2_{(CR)} = 0.15$, $R^2_{(NCP)} = 0.11$, $P < 0.001$ in all cases) (Table 3, Figure 8).

351 respectively (Figure 5), and when the mean values were the highest ($GPP_{\text{autumn-winter}} = 2.9 \pm 0.3 - 2.3 \pm$
352 $0.3 \text{ mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$; $CR_{\text{autumn-winter}} = 2.5 \pm 0.3 - 2 \pm 0.2$) (Figure 5). However, despite the overall
353 variability between autumn-winter and spring-summer, when all data are taken in concert, planktonic
354 GPP and CR rates were not significantly different between seasons (Kruskal-Wallis H test, $\chi^2 = 6.83$, p
355 $= 0.08$; $\chi^2 = 4.14$, $p = 0.25$, respectively). Furthermore, the balance between planktonic autotrophic
356 production (GPP) and the respiratory losses (due to the heterotrophic and autotrophic metabolism, CR)
357 (i.e., NCP rates), revealed that NCP rates also decreased towards the northern region (by 94%). From
358 $1.1 \pm 0.3 \text{ mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$ at the southern stations to $0.1 \pm 0.1 \text{ mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$ above 26°N (Figure 4).
359 The average NCP from our cruises was $0.3 \pm 0.1 \text{ mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$ (Figure 5), which indicates an overall
360 prevalence of autotrophic communities (Figure 5). However, a closer look to our data revealed that
361 during spring, the mean NCP rate was $-0.31 \pm 0.24 \text{ mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$ (Figure 5), while during the
362 summer, NCP rates in the northern region ranged from -0.64 to $-0.09 \text{ mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$, which evidenced
363 that planktonic metabolism was governed by heterotrophic communities during the spring and also
364 during the summer at the northern region.

365 When we evaluated the relationship of GPP with CR and NCP, the analysis showed that both
366 CR and NCP increased significantly with GPP ($R^2 = 0.62$ and 0.49 , respectively; $p < 0.001$) (Figure 6).
367 From the functional relationships between GPP with CR and NCP, we calculated the threshold of GPP
368 for metabolic equilibrium for the region. By solving for $GPP = CR$ and for $NCP = 0$ (from the
369 relationship describing NCP as a function of GPP), and by using the slope and intercept shown in
370 figures 6A and 6B, we determined that the GPP threshold that separates autotrophic from heterotrophic
371 planktonic communities in the Red Sea is $1.7 \text{ mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$ (range $1.2\text{--}1.9 \text{ mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$).

Deleted: (Figure 7A and B),

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: 50

Formatted: Font color: Text 1

Deleted: P

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: the fitted linear regressions

Formatted: Font color: Text 1

Deleted: obtained from the OLS analysis (Figures 7A and B) for GPP, where $GPP = CR$ and GPP where $NCP = 0$

Formatted: Font color: Text 1

378 **3.3 Metabolic rates and temperature**

379 Due to the pervasive influence of temperature in regulating metabolic rates, we further explored
380 the temperature-dependence of GPP and CR by analysing the relationship between chlorophyll-a
381 specific metabolic rates and temperature. Our analysis revealed that both GPP and CR tended to
382 increase with temperature albeit with different activation energies (i.e., E_a was significantly higher for
383 GPP (-1.2 ± 0.2 eV) than for CR rates (-0.73 ± 0.2 eV). ANCOVA, $F = 3.94$, $p = 0.04$) (Figure 7). We
384 also tested whether the temperature-dependence response was consistent between cruises (Figure 8).
385 Our results indicated a relatively higher activation energy for GPP during the summer cruise (-2.3 ± 0.8
386 eV) and in spring for CR (-2.6 ± 0.9 eV). However, the observed differences in the activation energies
387 for GPP were not significantly different between seasons (ANCOVA, $F = 0.38$, $p = 0.8$).
388

- Deleted: We ... [1]
- Formatted ... [1]
- Deleted: , ... [2]
- Formatted ... [2]
- Deleted: the normalised metabolic rates (Chlorophyll ... [3]
- Formatted ... [3]
- Deleted:) ... [4]
- Formatted ... [4]
- Deleted: This ... [5]
- Formatted ... [5]
- Deleted: showed ... [6]
- Formatted ... [6]
- Deleted: AE) (Figure 9). The AE ... [7]
- Formatted ... [7]
- Deleted: (- ... [8]
- Formatted ... [8]
- Deleted:). ... [9]
- Formatted ... [9]
- Deleted: observed in GPP and CR varied ... [10]
- Formatted ... [10]
- Deleted: seasons ... [11]
- Formatted ... [11]
- Deleted: 10). The ... [12]
- Formatted ... [12]
- Deleted: revealed that during summer, the AE for GPP rates was ... [13]
- Formatted ... [13]
- Deleted: 32 ... [14]
- Formatted ... [14]
- Deleted: 76 ... [15]
- Formatted ... [15]
- Deleted: than for the rest of ... [16]
- Formatted ... [16]
- Deleted: seasons (values ranged between 1.5–1.8 eV); although ... [17]
- Formatted ... [17]
- Deleted: seasonal ... [18]
- Formatted ... [18]
- Deleted: ($F = 0.39$, $dF = 3$, $P = 0.76$). The metabolic energy relation ... [19]
- Formatted ... [19]
- Deleted: 5.25, $dF = 3$, $P =$... [20]
- Formatted ... [20]
- Deleted: 002), with the highest values found in spring ($-2.63 \pm$... [21]
- Formatted ... [21]
- Deleted: 9 eV), and the AE for CR decreasing to -1.4 eV during ... [22]
- Formatted ... [22]
- Deleted: ... [23]
- Formatted ... [23]
- Formatted ... [24]

445 **4. Discussion**

446 **4.1 Variability of plankton community metabolic rates along the Red Sea**

447 Our results demonstrate that planktonic metabolic rates are markedly different between the
448 southern and northern regimes of the Red Sea, with an increase from the southern to the northern
449 regions in the overall mean GPP and CR by a factor of 5 and 4, respectively (i.e., an absolute increase in
450 GPP rates of $3.2 \text{ mmol O}_2 \text{ m}^{-3} \text{ d}^{-1} \approx 38.4 \text{ mgC m}^{-3} \text{ d}^{-1}$, and an absolute increase in CR rates of 2.2 mmol
451 $\text{O}_2 \text{ m}^{-3} \text{ d}^{-1} \approx 26.4 \text{ mgC m}^{-3} \text{ d}^{-1}$). Although, *sensu stricto*, the overall balance between autotrophic
452 metabolism and planktonic community respiration (i.e. NCP) indicated a prevalence of autotrophic
453 communities during our samplings along the Red Sea, heterotrophic communities prevailed during the
454 spring, and in the northern stations during the summer, which highlights the shift in the trophic
455 conditions in the basin. Consistent with these findings, our data revealed that the GPP threshold that
456 separated autotrophic from heterotrophic communities in the Red Sea ($1.7 \text{ mmol O}_2 \text{ m}^{-3} \text{ d}^{-1}$) is similar to
457 that reported across oceanic communities elsewhere (Duarte and Agustí, 1998; Duarte and Regaudie-de-
458 Gioux, 2009), agreeing with the oligotrophic characteristics that govern at certain periods or locations
459 the basin. The latitudinal differences depicted in our results mirror the increasing north-south pattern in
460 Chl-*a* concentration and photosynthetic carbon fixation rates previously reported for the Red Sea (Acker
461 et al., 2008; Raitso et al., 2013; Qurban et al., 2014; Kheireddine et al., 2017), and which are supported
462 by the presence of different planktonic communities (Al-aidaroos et al., 2016; Pearman et al., 2016;
463 Robitzsch et al., 2016; Kheireddine et al., 2017; Kottuparambil and Agusti, 2018).

464

- Deleted: (below 21–22 °N)
- Formatted: Font color: Text 1
- Deleted: the
- Formatted: Font color: Text 1
- Deleted: half
- Formatted: Font color: Text 1
- Deleted: basin (above 21–22 °N). Gross primary production and community respiration rates varied, on average, by a factor of 3–5 between the northern and
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Deleted: (during
- Formatted: Font color: Text 1
- Deleted: autumn and winter period). The metabolic
- Formatted: Font color: Text 1
- Deleted: (NCP) showed that
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Deleted: tended towards heterotrophy in the northern region (particularly
- Formatted: Font color: Text 1
- Deleted: summer and winter period), whereas the southern Red Sea tended to be mostly autotrophic, except in
- Formatted: Font color: Text 1
- Deleted: when all
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Deleted: < 22 °N were heterotrophic.
- Formatted: Font color: Text 1
- Deleted: was
- Formatted: Font color: Text 1
- Deleted: ,
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Deleted: . ¶

483 The lower productivity of the northern section of the Red Sea, explains the dominance of
 484 heterotrophic communities therein. Still, sustaining heterotrophy in oligotrophic regions requires an
 485 allochthonous source of organic matter (Duarte et al. 2011, 2013). The arid nature of the northern Red
 486 Sea, with the watershed consisting mostly of deserts, leads to the absence of rivers and significant
 487 organic carbon inputs to the sea. Dust inputs are important, however, and whereas they have shown no
 488 effect on primary production (Torfstein and Kienast, 2018), they are a source of organic carbon (Jurado
 489 et al. 2009) that can partially supply the organic matter required to sustain heterotrophic communities.
 490 Moreover, the Red Sea supports highly productive coral reef, mangrove, seagrass and algal
 491 communities in the extensive shallow coastal areas (Rasul et al., 2015; Almahasheer et al., 2016), which
 492 may export significant organic carbon to the pelagic compartment, thereby helping to sustain
 493 heterotrophic plankton communities in the northern Red Sea.

- Deleted:** low
- Formatted:** Font color: Text 1
- Deleted:** Northern
- Deleted:** (above 21–22 °N), with average GPP of 1.57 ± 0.16 mmol O₂ m⁻³ d⁻¹
- Deleted:** prevalence
- Formatted:** Font color: Text 1
- Deleted:** Sustaining
- Formatted:** Font color: Text 1
- Formatted:** Font color: Text 1
- Formatted:** Font color: Text 1
- Deleted:** carbon
- Formatted:** Font color: Text 1
- Deleted:** contains
- Formatted:** Font color: Text 1
- Formatted:** Font color: Text 1
- Formatted:** Font color: Text 1
- Deleted:** Northern
- Formatted:** Font color: Text 1
- Deleted:** The significant correlations observed between metabolic rates with Chl-*a* concentrations (positive relationship), and with latitude (inverse correlation) further supports the idea that different ecological domains govern planktonic metabolism in the Red Sea. In the southern half (below 21–22 °N), the balance between GPP and CR generally resulted in positive NCP values, i.e. autotrophic communities, while in the northern half (above 21–22 °N) the metabolic balance resulted in NCP values <= 0, i.e. heterotrophic communities. The latitudinal differences depicted in our results mirror the pattern of increasing Chl-*a* concentration and primary [25]
- Formatted:** Font color: Text 1
- Deleted:** the
- Formatted:** Font color: Text 1
- Formatted:** Font color: Text 1
- Formatted:** Font color: Text 1
- Deleted:** the chlorophyll-*a* specific
- Formatted:** Font color: Text 1

494 4.2 Temperature and metabolic balance in the Red Sea

495 Temperature is a master variable that regulates many components of ocean dynamics, such as
 496 vertical stratification, and most aspects of organismal biology, from setting boundaries in the
 497 distribution of organisms (Clarke, 1996) to controlling biochemical reactions that constrain the energy
 498 for metabolic processes (Gillooly et al., 2001). Hence, temperature is likely a significant driver of
 499 metabolic processes in the Red Sea, one of the warmest tropical marine ecosystems (Raitsos et al.,
 500 2011; Chaidez et al., 2017). Indeed, our results showed a positive response of planktonic metabolism to
 501 temperature. Moreover, the functional relationships between metabolic rates with temperature suggested
 502 that both GPP and CR were positively enhanced with increasing temperature; but at a different pace.

540 The metabolic theory of ecology (MTE) relates the metabolic rate of an organism with its mass
 541 and temperature. This theory hypothesizes that individual metabolic rates relate to temperature with a
 542 relatively constant activation energy ($E_a \sim 0.63$ eV) for a wide range of taxa, from unicellular organisms
 543 to plants and animals (Gillooly et al., 2001; Brown et al., 2004). For aerobic respiration, E_a values vary
 544 between 0.41 and 0.74 eV at temperatures between 0–40 °C (Gillooly et al., 2005), while for
 545 photosynthetic processes, the predicted E_a is lower, ~ 0.32 eV (Allen et al., 2005). From a thorough
 546 compilation of data obtained for a wide range of marine systems (from polar to subtropical and tropical
 547 oceanic regions), Regaudie-de-Gioux and Duarte (2012) found that overall, the activation energies for
 548 photosynthetic production (GPP) varied between 0.29–0.32 eV, and for respiratory processes (CR)
 549 between 0.65 and 0.66 eV.

550 The E_a for GPP (-1.2 ± 0.17 eV) obtained for the Red Sea was higher than the overall value
 551 predicted by the MTE, while the E_a values for CR were below those for GPP (0.72 ± 0.17 eV) unlike
 552 observed elsewhere in open oceanic waters (Regaudie-de-Gioux and Duarte 2011, Garcia-Corral et al.
 553 2017). Furthermore, these E_a values imply that GPP rates increased faster (5.1-fold) than CR rates (2.7-
 554 fold), in the Red Sea's thermal range (22–32.5 °C). These findings differ with the expected double
 555 increase of heterotrophic respiration (regarding photosynthetic processes) with temperature (Harris et
 556 al., 2006), but are closer to results obtained by Garcia-Corral et al. (2017), who recently reported
 557 activation energies for GPP of -0.86, -1.48 and -1.07 for the Atlantic, Indian, and Pacific oceans,
 558 respectively, while E_a for CR found in the Atlantic, Indian and the Pacific oceans were -0.77, -0.57 and
 559 -0.82, respectively.

- Deleted:), that
- Formatted ... [26]
- Deleted: .
- Formatted ... [27]
- Deleted: vary with
- Formatted ... [28]
- Deleted: an activation energy (minimum kinetic energy for the
- Formatted ... [29]
- Formatted ... [30]
- Deleted: (AE
- Formatted ... [31]
- Formatted ... [32]
- Formatted ... [33]
- Formatted ... [34]
- Deleted: AE
- Formatted ... [35]
- Formatted ... [36]
- Formatted ... [37]
- Deleted: in
- Formatted ... [38]
- Formatted ... [39]
- Formatted ... [40]
- Deleted: AE
- Formatted ... [41]
- Formatted ... [42]
- Deleted: AE
- Formatted ... [43]
- Formatted ... [44]
- Formatted ... [45]
- Formatted ... [46]
- Formatted ... [47]
- Formatted ... [49]
- Deleted: Chl-standardized activation energies for photosynthes
- Formatted ... [48]
- Formatted ... [51]
- Deleted: We found that AE for Chl-*a* standardized GPP and CR
- Formatted ... [52]
- Formatted ... [53]
- Formatted ... [54]
- Formatted ... [55]
- Deleted: from oligotrophic subtropical and tropical regions
- Formatted ... [56]
- Formatted ... [57]

595 The apparent contradiction between our findings and the general patterns predicted by the MTE
596 is, however, not surprising. In their model, Allen et al. (2005) predict the activation energy of
597 photosynthesis per chloroplast (for temperatures between 0–30 °C) using the temperature dependence
598 parameters obtained by Bernacchi et al. (2001) for RuBisCO carboxylation rates in one species (tobacco
599 leaves). Although the temperature range selected by Allen et al. (2005) comprises the optimum
600 temperatures of growth rates for a wide range of functional groups of marine primary producers (Chen,
601 2015; Thomas et al., 2016), the temperature observed in the Red Sea exceeded this range. Due to the
602 fast generation times of microbes (Collins, 2010), we can expect that photosynthetic planktonic
603 communities are acclimated or even locally adapted to the thermal conditions they experience. So by
604 favouring certain photosynthetic or thermal traits, they can enhance their metabolism and growth to the
605 temperatures they experience, up to their thermal optimum (Galmes et al., 2015; Thomas et al., 2016).
606 ~~Therefore, it is likely that the acclimation or local adaptation (in the long term) of photosynthetic traits~~
607 ~~in Red Sea plankton optimises the metabolic response at the high-temperatures reached, resulting in a~~
608 ~~steeper response to temperature than predicted by the MTE. Moreover, as the trait responses to~~
609 ~~temperature vary among phylogenetic groups (Galmes et al., 2015; Galmés et al., 2016; Thomas et al.,~~
610 ~~2016), we anticipated a certain degree of discrepancy if we characterise the photosynthetic response~~
611 ~~(GPP) of planktonic communities forming an ecosystem, by considering only one trait (i.e., RuBisCO~~
612 ~~carboxylation) of one species.~~

613 However, we must bear in mind that the metabolic response of individuals is not only
614 temperature-dependent, and that resource supply also plays an essential role (Brown et al., 2004; Allen
615 and Gillooly, 2009). Our results evidenced that the increased response of planktonic metabolism

Formatted: Font color: Text 1

Deleted: It

Formatted: Font color: Text 1

Deleted: therefore

Formatted: Font color: Text 1

Deleted: of

Formatted: Font color: Text 1

Deleted: there

Formatted: Font color: Text 1

Deleted: consider

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: ; thus their effect on planktonic communities can be intertwined (Raven and Geider, 1988; Marañón et al., 2014). A closer look at our

Formatted: Font color: Text 1

624 towards warmer temperatures was mostly confined to the southern half of the Red Sea, a region that
 625 receives the direct inflow of the enriched Intermediate Water coming from the Gulf of Aden during the
 626 winter monsoon (Raitos et al., 2015; Wafar et al., 2016). Recent findings have demonstrated that mass-
 627 specific carbon fixation rates of phytoplankton communities can be enhanced with temperature when
 628 nutrients are not limiting their growth (Marañón et al., 2014; Marañón et al., 2018). Therefore, it is
 629 likely that the intertwined effect of both the warmer temperatures and the larger nutrient availability
 630 towards the south of the Red Sea are key drivers regulating the metabolic response of planktonic
 631 communities. Thus, unlike the global ocean, where nutrient concentration is inversely correlated with
 632 temperature (e.g., Agawin et al. 2000), in the Red Sea nutrient concentration and temperature are
 633 positively correlated. This anomaly may explain the steep E_a for GPP, as primary producers in the
 634 warmer region are being supported by the inflow of the nutrient-enriched waters from the Indian Ocean.

635 The elevated E_a for GPP compared to CR in Red Sea plankton is also an anomaly, likely
 636 associated with the lack of allochthonous nutrient supply due to the absence of rivers and vegetation in
 637 the arid watershed of the Red Sea. The warm oligotrophic ocean is characterised by plankton
 638 communities that are in metabolic balance or net metabolic imbalanced (Duarte and Agusti 2008,
 639 Duarte et al. 2013). In contrast, the warm Southern Red Sea tends to support autotrophic metabolism,
 640 sustained by the input of nutrient-enriched waters while low allochthonous carbon inputs may constrain
 641 CR. As a result, NCP tends to increase, rather than decrease with increasing temperature (Regaudie-de-
 642 Gioux and Duarte 2011, Garcia-Corral et al. 2017). These patterns in plankton metabolism in the
 643 oligotrophic and warm Red Sea deviate from those characterising the subtropical and tropical gyres of
 644 the open ocean, but it provides an opportunity to explore the mechanistic basis for the global patterns in

Deleted: also linked to higher autotrophic biomass, and

Deleted: (below 21–22 °N). This

Formatted: Font color: Text 1

Deleted: These

Formatted: Font color: Text 1

Deleted: are in line with work demonstrating an increased metabolic response (i.e.,

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted:)

Formatted: Font color: Text 1

Deleted: phytoplanktonic

Formatted: Font color: Text 1

Deleted: ¶
Therefore

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: AE

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: AE

Formatted: Font color: Text 1

Deleted: rich

Formatted: Font color: Text 1

Deleted: Hence, the

Formatted: Font color: Text 1

Deleted: . This anomaly

Formatted: Font color: Text 1

559 plankton metabolism with temperature, which would otherwise remain obscured by the underlying
560 prevalent negative relationship with nutrient concentrations.

Deleted: with

Formatted: Font color: Text 1

561 **5. Conclusions**

562 Our results show that plankton metabolism in the Red Sea presents a remarkably different
563 pattern compared to other warm and oligotrophic marine systems (e.g., the subtropical and tropical
564 gyres). In this region, autotrophic plankton communities prevailed and are supported by relatively high
565 GPP rates; above the threshold separating heterotrophic low-productivity communities from autotrophic
566 ones. Metabolically-balanced or net heterotrophic plankton communities dominated in the Northern Red
567 Sea, whereas autotrophic communities, supported by nutrient inputs from the Gulf of Aden, were
568 predominant in the south. Elevated temperatures contributed to an enhanced metabolic activity of
569 planktonic organisms due to the increase in kinetic energy (favouring enzymatic reactions) with
570 temperature. Plankton communities in the Red Sea, however, displayed activation energies for GPP that
571 were higher than those for CR, resulting in a positive relationship between NCP and temperature. Those
572 findings represent anomalies in the relationship between metabolic rates and temperature compared to
573 the warm, oligotrophic open ocean. These anomalies are likely related to the higher nutrient supply
574 from nutrient-rich Indian Ocean waters in the warm Southern Red Sea, suggesting that GPP can respond
575 strongly to the temperature in the warm ocean when supported by high nutrient inputs, relative to those
576 in the subtropical gyres.

Deleted: Despite the lack of significant relationship between nutrient availability with metabolic rates, the close relationship found between planktonic metabolic rates with autotrophic biomass (Chl-*a*), and the significant relationship between Chl-*a* concentration and nutrient availability, suggest that variables regulating phytoplanktonic metabolism are also defining the metabolic response of the rest of the planktonic organisms in the region. The overall low nutrient concentration in the mixed layer along the basin (except a few stations in the south during autumn), and the relatively high Chl-*a* concentration during autumn and winter in the Southern Red Sea suggest fast turnover rates of the nutrient pools, a common feature in oligotrophic environments (Capblancq, 1990). Hence, Red Sea plankton communities are likely to be efficient using nutrients as they seemed to be rapidly consumed, with an associated response of autotrophic and heterotrophic communities (i.e., increase in GPP and CR rates).¶

Formatted: Font color: Text 1

Deleted: heterotrophic

Formatted: Font color: Text 1

Deleted: AEs

Formatted: Font color: Text 1

696 **Author Contributions**

697 DCL-S, CMD, and SA designed the study; KR and PCdA obtained the data and provided technical
698 support; DCL-S analysed the data; DCL-S wrote the article with a substantial contribution of CMD, and
699 SA; all authors discussed the results and commented on the manuscript.

700

701 **Acknowledgements**

702 The research reported in this publication was supported by funding from King Abdullah University of
703 Science and Technology (KAUST), under award number BAS/1/1071-01-10 assigned to CMD,
704 BAS/1/1072-01-01 assigned to SA, and CCF/1/1973-21-01 assigned to the Red Sea Research Center.

Formatted: Font: 12 pt, Font color: Text 1

Deleted: provide

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: 12 pt, Font color: Text 1

Formatted: Font color: Text 1

Deleted: FCC

Formatted: Font color: Text 1

707 References

- 708 Acker, J., Leptoukh, G., Shen, S., Zhu, T., and Kempler, S.: Remotely-sensed chlorophyll a observations of the northern Red Sea indicate
709 seasonal variability and influence of coastal reefs, *Journal of Marine Systems*, 69, 191-204, 10.1016/j.jmarsys.2005.12.006, 2008.
- 710 Agawin, N. S., Duarte, C. M., and Agustí, S.: Nutrient and temperature control of the contribution of picoplankton to phytoplankton biomass
711 and production, *Limnology and Oceanography*, 45, 591-600, 2000.
- 712 Al-aidaroos, A. M., Karati, K. K., El-sherbiny, M. M., Devassy, R. P., and Kürten, B.: Latitudinal environmental gradients and diel variability
713 influence abundance and community structure of Chaetognatha in Red Sea coral reefs, *Systematics and Biodiversity*, 15, 35-48,
714 10.1080/14772000.2016.1211200, 2016.
- 715 Almahasheer, H., Abdulaziz, A., and Duarte, C. M.: Decadal stability of Red Sea mangroves, *Estuarine Coastal and Shelf Science*, 169, 164-
716 172, 2016.
- 717 Almahasheer, H., Duarte, C. M., and Irigoien, X.: Leaf Nutrient Resorption and Export Fluxes of *Avicennia marina* in the Central Red Sea
718 Area, *Frontiers in Marine Science*, 5, 10.3389/fmars.2018.00204, 2018.
- 719 Allen, A., Gillooly, J., and Brown, J.: Linking the global carbon cycle to individual metabolism, *Functional Ecology*, 19, 202-213, 2005.
- 720 Allen, A. P., and Gillooly, J. F.: Towards an integration of ecological stoichiometry and the metabolic theory of ecology to better understand
721 nutrient cycling, *Ecol Lett*, 12, 369-384, 10.1111/j.1461-0248.2009.01302.x, 2009.
- 722 Barrón, C., and Duarte, C. M.: Dissolved organic carbon pools and export from the coastal ocean, *Global Biogeochemical Cycles*, 29, 1725-
723 1738, 2015.
- 724 Bender, M., Orchardo, J., Dickson, M.-L., Barber, R., and Lindley, S.: In vitro O₂ fluxes compared with 14 C production and other rate
725 terms during the JGOFS Equatorial Pacific experiment, *Deep Sea Research Part I: Oceanographic Research Papers*, 46, 637-654, 1999.
- 726 Bernacchi, C., Singasas, E., Pimentel, C., Portis Jr, A., and Long, S.: Improved temperature response functions for models of Rubisco-limited
727 photosynthesis, *Plant, Cell & Environment*, 24, 253-259, 2001.
- 728 Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., and West, G. B.: Toward a metabolic theory of ecology, *Ecology*, 85, 1771-1789,
729 2004.
- 730 Capblancq, J.: Nutrient dynamics and pelagic food web interactions in oligotrophic and eutrophic environments: an overview, *Hydrobiologia*,
731 207, 1-14, 1990.
- 732 Carpenter, J. H.: The accuracy of the Winkler method for dissolved oxygen analysis, *Limnology and Oceanography*, 10, 135-140, 1965.
- 733 Clarke, A.: The influence of climate change on the distribution and evolution of organisms, *Animals and Temperature. Phenotypic and
734 Evolutionary Adaptation*, 377-407, 1996.
- 735 Collins, S.: Many Possible Worlds: Expanding the Ecological Scenarios in Experimental Evolution, *Evolutionary Biology*, 38, 3-14,
736 10.1007/s11692-010-9106-3, 2010.
- 737 Cullen, J.: Primary production methods, *Marine Ecology Progress Series*, 52, 88, 2001.
- 738 Chaidez, V., Dreano, D., Agustí, S., Duarte, C. M., and Hoteit, I.: Decadal trends in Red Sea maximum surface temperature, *Scientific
739 Reports*, 7, 8144, 10.1038/s41598-017-08146-z, 2017.
- 740 Chen, B.: Patterns of thermal limits of phytoplankton, *Journal of Plankton Research*, 37, 285-292, 10.1093/plankt/fbv009, 2015.
- 741 Chen, Y., Mills, S., Street, J., Golan, D., Post, A., Jacobson, M., and Paytan, A.: Estimates of atmospheric dry deposition and associated
742 input of nutrients to Gulf of Aqaba seawater, *Journal of Geophysical Research: Atmospheres*, 112, 2007.
- 743 Del Giorgio, P. A., Cole, J. J., and Cimleris, A.: Respiration rates in bacteria exceed phytoplankton production in unproductive aquatic
744 systems, *Nature*, 385, 148, 1997.
- 745 Dell, A. I., Pawar, S., and Savage, V. M.: Systematic variation in the temperature dependence of physiological and ecological traits,
746 *Proceedings of the National Academy of Sciences*, 108, 10591-10596, 2011.
- 747 Duarte, C. M., and Agustí, S.: The CO₂ balance of unproductive aquatic ecosystems, *Science*, 281, 234-236, 1998.
- 748 Duarte, C. M., and Regaudie-de-Gioux, A.: Thresholds of gross primary production for the metabolic balance of marine planktonic
749 communities, *Limnology and Oceanography*, 54, 1015-1022, 2009.
- 750 Duarte, C. M., Regaudie-de-Gioux, A., Arrieta, J. M., Delgado-Huertas, A., and Agustí, S.: The Oligotrophic Ocean Is Heterotrophic, *Annual
751 Review of Marine Science*, 5, 551-569, 10.1146/annurev-marine-121211-172337, 2013.
- 752 Ducklow, H. W., Dickson, M.-L., Kirchman, D. L., Steward, G., Orchardo, J., Marra, J., and Azam, F.: Constraining bacterial production,
753 conversion efficiency and respiration in the Ross Sea, Antarctica, January–February, 1997, *Deep Sea Research Part II: Topical Studies in
754 Oceanography*, 47, 3227-3247, 2000.
- 755 Ducklow, H. W., and Doney, S. C.: What is the metabolic state of the oligotrophic ocean? A debate, *Ann Rev Mar Sci*, 5, 525-533,
756 10.1146/annurev-marine-121211-172331, 2013.
- 757 Engelbrecht, J. P., Stenichkov, G., Prakash, P. J., Lersch, T., Anisimov, A., and Shevchenko, I.: Physical and chemical properties of deposited
758 airborne particulates over the Arabian Red Sea coastal plain, *Atmospheric Chemistry and Physics*, 17, 11467-11490, 2017.

Formatted: Font: 12 pt, Font color: Text 1

Formatted: Font color: Text 1

Deleted: Agustí, S., and Regaudie-de-Gioux, A.: The Role of Marine Biota in the Metabolism of the Biosphere, in: *The Role of Marine Biota in the Functioning of the Biosphere*, edited by: Duarte, C. M., Fundacion BBVA, Bilbao, Spain, 39-53, 2011.¶ Duarte, C. M.,

764 Galmes, J., Kapralov, M., Copolovici, L., Hermida-Carrera, C., and Niinemets, Ü.: Temperature responses of the Rubisco maximum
765 carboxylase activity across domains of life: phylogenetic signals, trade-offs, and importance for carbon gain, *Photosynthesis research*, 123,
766 183-201, 2015.

767 Galmés, J., Hermida-Carrera, C., Laanisto, L., and Niinemets, Ü.: A compendium of temperature responses of Rubisco kinetic traits:
768 variability among and within photosynthetic groups and impacts on photosynthesis modeling, *Journal of experimental botany*, 67, 5067-
769 5091, 2016.

770 García-Corral, L., Barber, E., Gioux, A. R. d., Sal, S., Holding, J., Agustí, S., Navarro, N., Serret, P., Mozeti, P., and Duarte, C.: Temperature
771 dependence of planktonic metabolism in the subtropical North Atlantic Ocean, *Biogeosciences*, 11, 4529-4540, 2014.

772 García-Corral, L. S., Holding, J. M., Carrillo-de-Albornoz, P., Steckbauer, A., Pérez-Lorenzo, M., Navarro, N., Serret, P., Gasol, J. M.,
773 Morán, X. A. G., Estrada, M., Fraile-Nuez, E., Benítez-Barrios, V., Agustí, S., and Duarte, C. M.: Temperature dependence of plankton
774 community metabolism in the subtropical and tropical oceans, *Global Biogeochemical Cycles*, 31, 1141-1154, 10.1002/2017gb005629,
775 2017.

776 [García-Martín, E. E., Daniels, C. J., Davidson, K., Davis, C. E., Mahaffey, C., Mayers, K. M. J., McNeill, S., Poulton, A. J., Purdie, D. A.,
777 Tarran, G. A., and Robinson, C.: Seasonal changes in plankton respiration and bacterial metabolism in a temperate shelf sea, *Progress in
778 Oceanography*, 10.1016/j.pocan.2017.12.002, 2017.](#)

779 Garcias-Bonet, N., and Duarte, C. M.: Methane Production by Seagrass Ecosystems in the Red Sea, *Frontiers in Marine Science*, 4,
780 10.3389/fmars.2017.00340, 2017.

781 Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M., and Charnov, E. L.: Effects of size and temperature on metabolic rate, *science*,
782 293, 2248-2251, 2001.

783 Gillooly, J. F., Allen, A. P., West, G. B., and Brown, J. H.: The rate of DNA evolution: effects of body size and temperature on the molecular
784 clock, *Proceedings of the National Academy of Sciences of the United States of America*, 102, 140-145, 2005.

785 Hansen, H. P., and Koroleff, F.: Determination of nutrients, in: *Methods of seawater analysis*, edited by: K. Grasshoff, K. Kremling, and
786 Ehrhardt, M., Wiley-VCH Verlag, Weinheim, Germany, 159-228, 1999.

787 Harris, L. A., Duarte, C. M., and Nixon, S. W.: Allometric laws and prediction in estuarine and coastal ecology, *Estuaries and Coasts*, 29,
788 340-344, 2006.

789 Jurado, E., Dachs, J., Duarte, C. M., and Simo, R.: Atmospheric deposition of organic and black carbon to the global oceans, *Atmospheric
790 Environment*, 42, 7931-7939, 2008.

791 Kheireddine, M., Ouhssain, M., Claustre, H., Uitz, J., Gentili, B., and Jones, B.: Assessing pigment-based phytoplankton community
792 distributions in the Red Sea, *Frontiers in Marine Science*, 2017.

793 Kottaparambil, S., and Agustí, S.: PAHs sensitivity of picophytoplankton populations in the Red Sea, *Environmental Pollution*, 239, 607-
794 616, 2018.

795 Kürten, B., Al-Aidaros, A. M., Kürten, S., El-Sherbiny, M. M., Devassy, R. P., Struck, U., Zarokanellos, N., Jones, B. H., Hansen, T.,
796 Bruss, G., and Sommer, U.: Carbon and nitrogen stable isotope ratios of pelagic zooplankton elucidate ecohydrographic features in the
797 oligotrophic Red Sea, *Progress in Oceanography*, 140, 69-90, 10.1016/j.pocan.2015.11.003, 2016.

798 [Levanon-Spanier, I., Padan, E., and Reiss, Z.: Primary production in a desert-enclosed sea—the Gulf of Elat \(Aqaba\), *Red Sea, Deep Sea
799 Research Part A, Oceanographic Research Papers*, 26, 673-685, 1979.](#)

800 López-Urrutia, A., San Martín, E., Harris, R. P., and Irigoien, X.: Scaling the metabolic balance of the oceans, *Proceedings of the National
801 Academy of Sciences*, 103, 8739-8744, 2006.

802 Maraño, E., Cermeño, P., Huete-Ortega, M., López-Sandoval, D. C., Mouriño-Carballido, B., and Rodríguez-Ramos, T.: Resource supply
803 overrides temperature as a controlling factor of marine phytoplankton growth, *PLoS one*, 9, e99312, 2014.

804 Maraño, E., Lorenzo, M. P., Cermeño, P., and Mouriño-Carballido, B.: Nutrient limitation suppresses the temperature dependence of
805 phytoplankton metabolic rates, *The ISME journal*, 2018.

806 Osman, E. O., Smith, D. J., Ziegler, M., Kürten, B., Conrad, C., El-Haddad, K. M., Voolstra, C. R., and Suggett, D. J.: Thermal refugia
807 against coral bleaching throughout the northern Red Sea, *Global change biology*, 24, e474-e484, 2018.

808 Oudot, C., Gerard, R., Morin, P., and Gningue, I.: Precise shipboard determination of dissolved oxygen (Winkler procedure) for productivity
809 studies with a commercial system1, *Limnology and Oceanography*, 33, 146-150, 1988.

810 Padfield, D., Lowe, C., Buckling, A., Ffrench-Constant, R., Student Research, T., Jennings, S., Shelley, F., Olafsson, J. S., and Yvon-
811 Durocher, G.: Metabolic compensation constrains the temperature dependence of gross primary production, *Ecol Lett*, 20, 1250-1260,
812 10.1111/ele.12820, 2017.

813 Pearman, J. K., Kürten, S., Sarma, Y., Jones, B., and Carvalho, S.: Biodiversity patterns of plankton assemblages at the extremes of the Red
814 Sea, *FEMS microbiology ecology*, 92, fiw002, 2016.

815 Qurban, M. A., Balala, A. C., Kumar, S., Bhavya, P. S., and Wafar, M.: Primary production in the northern Red Sea, *Journal of Marine
816 Systems*, 132, 75-82, 10.1016/j.jmarsys.2014.01.006, 2014.

817 Qurban, M. A., Wafar, M., Jyothibabu, R., and Manikandan, K. P.: Patterns of primary production in the Red Sea, *Journal of Marine Systems*,
818 169, 87-98, 10.1016/j.jmarsys.2016.12.008, 2017.

Deleted: Julkowska, M. M., Saade, S., Gao, G., Morton, M. J. L.,
wlia, M., and Tester, M. A., MVApp.pre-release_v2.0
mmjulkowska/MVApp: MVApp.pre-release_v2.0. DOI:
10.5281/zenodo.1067974

823 [Rahav, E., Herut, B., Mulholland, M. R., Belkin, N., Elifantz, H., and Berman-Frank, I.: Heterotrophic and autotrophic contribution to](#)
824 [dinitrogen fixation in the Gulf of Aqaba, *Marine Ecology Progress Series*, 522, 67-77, 2015.](#)
825 [Raitsos, D. E., Hoteit, I., Prihartato, P. K., Chronis, T., Triantafyllou, G., and Abualnaja, Y.: Abrupt warming of the Red Sea, *Geophysical*
826 \[Research Letters\]\(#\), 38, n/a-n/a, 10.1029/2011gl047984, 2011.](#)
827 [Raitsos, D. E., Pradhan, Y., Brewin, R. J., Stenchikov, G., and Hoteit, I.: Remote sensing the phytoplankton seasonal succession of the Red](#)
828 [Sea, *PLoS One*, 8, e64909, 10.1371/journal.pone.0064909, 2013.](#)
829 [Raitsos, D. E., Yi, X., Platt, T., Racault, M.-F., Brewin, R. J. W., Pradhan, Y., Papadopoulos, V. P., Sathyendranath, S., and Hoteit, I.:](#)
830 [Monsoon oscillations regulate fertility of the Red Sea, *Geophysical Research Letters*, 42, 855-862, 10.1002/2014gl062882, 2015.](#)
831 [Rasul, N. M., and Stewart, I. C.: The Red Sea: the formation, morphology, oceanography and environment of a young ocean basin, Springer,](#)
832 [2015.](#)
833 [Rasul, N. M., Stewart, I. C., and Nawab, Z. A.: Introduction to the Red Sea: its origin, structure and environment., in: *The Red Sea*, edited](#)
834 [by: Rasul, N. M., and Stewart, I. C., Springer, Berlin, 1-28, 2015.](#)
835 [Raven, J. A., and Geider, R. J.: Temperature and algal growth, *New phytologist*, 110, 441-461, 1988.](#)
836 [Regaudie-de-Gioux, A., and Duarte, C. M.: Temperature dependence of planktonic metabolism in the ocean, *Global Biogeochemical Cycles*,](#)
837 [26, 2012.](#)
838 [Regaudie-de-Gioux, A., and Duarte, C. M.: Global patterns in oceanic planktonic metabolism, *Limnology and Oceanography*, 58, 977-986,](#)
839 [doi:10.4319/lo.2013.58.3.0977, 2013.](#)
840 [Robinson, C., and Williams, P. J. I. B.: Plankton net community production and dark respiration in the Arabian Sea during September 1994,](#)
841 [Deep Sea Research Part II: Topical Studies in Oceanography, 46, 745-765, 1999.](#)
842 [Robinson, C., Serret, P., Tilstone, G., Teira, E., Zubkov, M. V., Rees, A. P., and Woodward, E. M. S.: Plankton respiration in the eastern](#)
843 [Atlantic Ocean, Deep Sea Research Part I: Oceanographic Research Papers, 49, 787-813, 2002.](#)
844 [Robinson, C., and Williams, P. I. B.: Respiration and its measurement in surface marine waters, *Respiration in aquatic ecosystems*, 147-180,](#)
845 [2005.](#)
846 [Robitzsch, V. S., Lozano-Cortes, D., Kandler, N. M., Salas, E., and Berumen, M. L.: Productivity and sea surface temperature are correlated](#)
847 [with the pelagic larval duration of damselfishes in the Red Sea, *Mar Pollut Bull*, 105, 566-574, 10.1016/j.marpolbul.2015.11.045, 2016.](#)
848 [Serret, P., Robinson, C., Fernández, E., Teira, E., and Tilstone, G.: Latitudinal variation of the balance between plankton photosynthesis and](#)
849 [respiration in the eastern Atlantic Ocean, *Limnology and Oceanography*, 46, 1642-1652, 2001.](#)
850 [Serret, P., Robinson, C., Fernández, E., Teira, E., Tilstone, G., and Pérez, V.: Predicting plankton net community production in the Atlantic](#)
851 [Ocean, Deep Sea Research Part II: Topical Studies in Oceanography, 56, 941-953, 2009.](#)
852 [Smith, S., and Mackenzie, F.: The ocean as a net heterotrophic system: implications from the carbon biogeochemical cycle, *Global*](#)
853 [Biogeochemical Cycles](#), 1, 187-198, 1987.
854 [Smith, S., and Hollibaugh, J.: Coastal metabolism and the oceanic organic carbon balance, *Reviews of Geophysics*, 31, 75-89, 1993.](#)
855 [Sofianos, S., and Johns, W. E.: Water mass formation, overturning circulation, and the exchange of the Red Sea with the adjacent basins, in:](#)
856 [The Red Sea, Springer, 343-353, 2015.](#)
857 [Sofianos, S. S.: An Oceanic General Circulation Model \(OGCM\) investigation of the Red Sea circulation, 1. Exchange between the Red Sea](#)
858 [and the Indian Ocean, *Journal of Geophysical Research*, 107, 10.1029/2001jc001184, 2002.](#)
859 [Sofianos, S. S., and Johns, W. E.: Observations of the summer Red Sea circulation, *Journal of Geophysical Research*, 112,](#)
860 [10.1029/2006jc003886, 2007.](#)
861 [Thomas, M. K., Kremer, C. T., and Litchman, E.: Environment and evolutionary history determine the global biogeography of phytoplankton](#)
862 [temperature traits, *Global Ecology and Biogeography*, 25, 75-86, 10.1111/geb.12387, 2016.](#)
863 [Tilstra, A., van Hoytema, N., Cardini, U., Bednarz, V. N., Rix, L., Naumann, M. S., Al-Horani, F. A., and Wild, C.: Effects of water column](#)
864 [mixing and stratification on planktonic primary production and dinitrogen fixation on a northern Red Sea coral reef, *Frontiers in*](#)
865 [microbiology](#), 9, 2018.
866 [Torfstein, A., and Kienast, S.: No Correlation Between Atmospheric Dust and Surface Ocean Chlorophyll-a in the Oligotrophic Gulf of](#)
867 [Aqaba, Northern Red Sea, *Journal of Geophysical Research: Biogeosciences*, 123, 391-405, 2018.](#)
868 [Wafar, M., Qurban, M. A., Ashraf, M., Manikandan, K., Flandez, A. V., and Balala, A. C.: Patterns of distribution of inorganic nutrients in](#)
869 [Red Sea and their implications to primary production, *Journal of Marine Systems*, 156, 86-98, 2016.](#)
870 [Williams, P., Raine, R. C. T., and Bryan, J. R.: Agreement between the c-14 and oxygen methods of measuring phytoplankton production-](#)
871 [reassessment of the photosynthetic quotient, *Oceanologica Acta*, 2, 411-416, 1979.](#)
872 [Williams, P.: On the definition of plankton production terms, *ICES marine science symposia*, 1993., 1993.](#)
873 [Williams, P. I. B.: The balance of plankton respiration and photosynthesis in the open oceans, *Nature*, 394, 55-57, 1998.](#)
874 [Williams, P. I. B., and del Giorgio, P. A.: Respiration in aquatic ecosystems: history and background, *Respiration in aquatic ecosystems*, 1-](#)
875 [17, 2005.](#)
876 [Zarokanellos, N., Papadopoulos, V. P., Sofianos, S., and Jones, B.: Physical and biological characteristics of the winter-summer transition](#)
877 [in the Central Red Sea, *Journal of Geophysical Research: Oceans*, 122, 6355-6370, <http://dx.doi.org/10.1002/2017jc012882>., 2017.](#)
878

Formatted: Font color: Text 1




Figure 1: Stations sampled along the Red Sea during (A) spring (2018), (B) summer 2018, (C) autumn (2016) and (D) winter 2016 and 2017




Figure 2: Overall seasonal and latitudinal variability of surface (A) temperature (SST), (B) salinity (C) and chlorophyll-a concentration (Chl-a) measured during spring (2018), summer (2017), autumn (2016) and winter (2016 and 2017) cruises along the Red Sea (~ 100 % of incident Photosynthetically Active Radiation, PAR).




Figure 3: Pearson correlations between environmental variables (temperature and the concentrations of nitrate+nitrite [NOx], phosphate and silicate) and their latitudinal distribution measured at selected depths: (A) the first optical depth (from the surface down to 37% of incident PAR) and (B) at the bottom of the photic layer (between 1–0.1 % of incident PAR values). The size of the squares is the magnitude, the color indicates the direction (green for positive correlations, purple for negative correlations). The value of the correlation coefficient (r) is shown in the color bar below the graphs. Non-significant correlations are denoted with a \times .




Figure 4: Ordinary least squares linear regression between gross primary production (GPP), planktonic community respiration (CR) and net community production rates (NCP) with (A, B, C) Chlorophyll-a concentration (Chl-a), (D, E, F) temperature and (G, H, I) latitude. The solid red line is the linear least square fit, while the shaded grey area represents the 95% confidence intervals. The coefficient of determination and the statistical significance are indicated for each regression.




Figure 5. Box plots illustrating the seasonal variability of (A) gross primary production (GPP), (B) community respiration (CR), and (C) net community production (NCP) measured along the Red Sea. On each box are the data layered over a 95% confidence interval (shaded in lighter color), and ± 1 SD (shaded in grey). The central horizontal white lines in the box mark the mean value for each season. The red dashed lines represent the overall mean while the red dotted line in (C) defines the limit between autotrophic from heterotrophic communities (NCP=0). Values inside the donut plots (C) indicate the percentage of heterotrophy (NCP<) for each season.




Figure 6: Ordinary least square linear regression between (A) planktonic community respiration and (B) net community production (NCP) with gross primary production (GPP) rates measured along the Red Sea. The ordinary least square regression parameters (slope and intercept) and the statistical significance of each regression are indicated. The solid red line represents the linear least square fit, the shaded grey area represents the 95% confidence interval.




Figure 7: Arrhenius plots indicating temperature dependence of planktonic metabolic rates plotted as the relationship between the natural logarithm of (A) chlorophyll-a normalised gross primary production, and (B) chlorophyll-a normalised planktonic community respiration with temperature as a function of 1/kT (lower axis), where k is the Boltzmann's constant (8.2×10^{-5} eV K⁻¹), and T denotes the absolute temperature (K). The corresponding temperatures in degree Celsius are shown in the upper axis for each graph. The solid red line is the linear least square fit, the shaded grey area represents the 95% confidence interval. E_a is the slope of each plot and represents the activation energy.




Figure 8: Arrhenius plots indicating the seasonal temperature dependence of planktonic metabolic rates plotted as the relationship between the natural logarithm of (A) chlorophyll-*a* normalised gross primary production, and (B) planktonic community respiration with temperature as a function of $1/kT$ (lower axis), where k is the Boltzmann's constant ($8.2 \times 10^{-5} \text{ eV K}^{-1}$), and T denotes the absolute temperature (K). Each line represents the linear least square fit. E_a is the slope of each regression line and represents the activation energy.

Table 1. Pearson correlation matrix between volumetric gross primary production (GPP), planktonic community respiration (CR) and net community production (NCP) with environmental variables (temperature; latitude; nitrite+nitrate, NO_x; and Chlorophyll-a concentration, Chl-*a*). Bold numbers indicate significant relationships and the significance level is indicated with *: p<0.05*, p<0.01** and p<0.001***.

	Temperature	Latitude	NO _x	Chl- <i>a</i>	GPP	CR	NCP
GPP	0.41**	-0.60***	0.01	0.69***		0.73***	0.70***
CR	0.40**	-0.37*	0.19	0.61***	0.73***		0.02
NCP	0.19	-0.49***	-0.19	0.37***	0.70***	0.02	
Chl- <i>a</i>	0.1	-0.41***	0.29*		0.71***	0.67***	0.37**