

1 **Shifts in dimethylated sulfur concentrations and microbiome composition in the red-**
2 **tide causing dinoflagellate *Alexandrium minutum* during a simulated marine heat wave**

3
4 Elisabeth Deschaseaux^{1*}, James O'Brien¹, Nachshon Siboni¹, Katherina Petrou^{1,2} and Justin
5 R. Seymour¹
6

7 ¹ University of Technology Sydney, Climate Change Cluster, Ultimo, NSW, 2007, Australia.

8 ² University of Technology Sydney, School of Life Sciences, Ultimo, NSW, 2007, Australia.

9
10 * **Corresponding author current address:** Dr Elisabeth Deschaseaux, elisabeth.deschaseaux@gmail.com, Centre for Coastal
11 Biogeochemistry, School of Environment Science and Engineering, Southern Cross University, Lismore, NSW, 2481,
12 Australia, Ph: (+61) 4 2360 2341.

13
14
15 **Abstract**

16
17 The biogenic sulfur compounds dimethyl sulfide (DMS), dimethyl sulfoniopropionate (DMSP)
18 and dimethyl sulfoxide (DMSO) are produced and transformed by diverse populations of
19 marine microorganisms and have substantial physiological, ecological and biogeochemical
20 importance spanning organism to global scales. Understanding the production and
21 transformation dynamics of these compounds under shifting environmental conditions is
22 important for predicting their roles in a changing ocean. Here, we report the physiological and
23 biochemical response of *Alexandrium minutum*, a dinoflagellate with the highest reported
24 intracellular DMSP content, exposed to a 6-day increase in temperature mimicking mild and
25 extreme coastal marine heatwave conditions (+ 4°C and + 12°C). Under mild temperature
26 increases (+ 4°C), *A. minutum* growth was enhanced, with no measurable physiological stress
27 response. However, under a very acute increase in temperature (+ 12°C), *A. minutum* growth
28 declined, photosynthetic efficiency (F_V/F_M) was impaired, and enhanced oxidative stress was
29 observed. These physiological responses indicative of thermal stress were accompanied by
30 increased DMS and DMSO concentrations followed by decreased DMSP concentrations. At
31 this temperature extreme, we observed a cascading stress response in *A. minutum*, which was
32 initiated 6h after the start of the experiment by a spike in DMS and DMSO concentrations and
33 a rapid decrease in F_V/F_M . This was followed by an increase in reactive oxygen species (ROS)
34 and an abrupt decline in DMS and DMSO on day 2 of the experiment. A subsequent decrease
35 in DMSP coupled with a decline in the growth rate of both *A. minutum* and its associated total
36 bacterial assemblage coincided with a shift in the composition of the *A. minutum* microbiome.
37 Specifically, an increase in the relative abundance of OTUs matching the genus *Oceanicaulis*
38 (17.0%), *Phycisphaeraceae* SM1A02 (8.8%) and *Balneola* (4.9%) as well as a decreased
39 relative abundance of *Maribacter* (24.4%), *Marinoscillum* (4.7%) and *Seohaecicola* (2.7%),

40 were primarily responsible for differences in microbiome structure observed between
41 temperature treatments. These shifts in microbiome structure are likely to have been driven by
42 either the changing physiological state of *A. minutum* cells, shifts in biogenic sulfur
43 concentrations, the presence of other solutes, or a combination of all. Nevertheless, we suggest
44 that these results point to the significant effect of heatwaves on the physiology, growth and
45 microbiome composition of the red-tide causing dinoflagellate *A. minutum*, as well as potential
46 implications for biogenic sulfur cycling processes and marine DMS emissions.

47

48 **Keywords:** DMS, DMSP, DMSO, oxidative stress, thermal stress

1. Introduction

Many marine phytoplankton produce the organic sulfur dimethyl sulfoniopropionate (DMSP) (Zhou et al., 2009; Berdalet et al., 2011; Caruana and Malin, 2014), for which it can function as an antioxidant, osmolyte, chemoattractant and currency in reciprocal chemical exchanges with heterotrophic bacteria (Stefels, 2000; Sunda et al., 2002; Kiene et al., 2000; Seymour et al., 2010). Phytoplankton-derived DMSP is in fact a major source of sulfur and carbon for marine heterotrophic bacteria (Kiene et al., 2000), which in turn play a major role in the cycling and turnover of organosulfur compounds in the ocean (Todd et al., 2007; Curson et al., 2011). The subsequent cycling of DMSP into other biogenic sulfur molecules including dimethyl sulfide (DMS) and dimethyl sulfoxide (DMSO) by a suite of microbial transformation pathways (Kiene et al., 2000; Sunda et al., 2002) and physical drivers (Brimblecombe and Shooter, 1986) have important ecological and biogeochemical implications spanning from cellular to global scales (Sunda et al., 2002; Charlson et al., 1987; DeBose et al., 2008; Van Alstyne et al., 2001; Knight, 2012; Nevitt et al., 1995).

Among DMSP-producing phytoplankton, the dinoflagellate *Alexandrium minutum*, has the highest recorded DMSP cell content, with an average concentration of 14.2 pmol cell⁻¹, compared with less than 1 pmol cell⁻¹ in most other dinoflagellates (Caruana and Malin, 2014). Blooms of *A. minutum* occur from the Mediterranean Sea to the South Pacific coast in sea surface waters within temperature ranges of 12°C to 25°C (Laabir et al., 2011). Notably, some strains of *Alexandrium*, including *A. minutum*, produce saxitoxins, which lead to paralytic shellfish poisoning (PSP) and are responsible for the most harmful algal blooms in terms of magnitude, distribution and consequences on human health (Anderson et al., 2012).

A. minutum commonly inhabits shallow coastal and estuarine waters (Anderson, 1998), which are globally experiencing substantial shifts in environmental conditions, including increases in sea surface temperature (SST) associated with climate change (Harley et al., 2006). Although generally less studied than chronic temperature rises associated with global climate change (Frölicher and Laufkötter, 2018), acute ephemeral temperature increases known as marine heatwaves (MHWs) (Hobday et al., 2016) have recently been demonstrated to be becoming more frequent and persistent as a consequence of climate change (Oliver et al., 2018). Increases in MHW occurrence are anticipated to become particularly frequent within the shallow coastal

83 and estuarine waters, where *A. minutum* blooms occur (Ummenhofer and Meehl,
84 2017;Anderson, 1998).

85

86 Coastal MHW events have recently had dramatic impacts on coastal environments. MHWs in
87 Western Australian (2011) and the Northeast Pacific (2013-2015) resulted in significant
88 ecosystem shifts with increases in novel species at the expenses of others (Frölicher and
89 Laufkötter, 2018). The 2016 MHW that was associated with El Niño Southern Oscillations
90 resulted in an 8°C increase in sea surface temperature leading to the mass coral bleaching of
91 more than 90% of the Great Barrier Reef (Hughes et al., 2017). While it is clear that MHWs
92 can have severe consequences on a variety of systems and organisms, their effects on marine
93 microbes and the biogeochemical processes that they mediate have rarely been investigated
94 (Joint and Smale, 2017).

95

96 While there is evidence that increases in seawater temperature can lead to increased DMSP
97 and/or DMS concentrations in phytoplankton (McLenon and DiTullio, 2012;Sunda et al.,
98 2002), it is not clear how a shift in DMSP net production by phytoplankton under acute
99 temperature stress will alter the composition and function of their associated microbiome and
100 how, in turn, this will influence biogenic sulfur cycling processes within marine habitats. There
101 is therefore a pressing need to understand the physiological and biogeochemical consequences
102 of thermal stress on phytoplankton-bacteria interactions within the context of events such as
103 MHWs. This is particularly important, given that a shift in the composition of the
104 phytoplankton microbiome could potentially dictate atmospheric DMS fluxes depending on
105 whether the bacterial community preferentially cleave or demethylate DMSP (Todd et al.,
106 2007;Kiene et al., 2000).

107

108 The aims of this study were to investigate how acute increases in temperature, such as those
109 associated with MHW events, alter the physiological state and biogenic sulfur cycling
110 dynamics of *A. minutum* and determine how these changes might influence the composition of
111 the *Alexandrium* microbiome. We hypothesized that an abrupt increase in temperature would
112 lead to physiological impairment (Falk et al., 1996;Robison and Warner, 2006;Iglesias-Prieto
113 et al., 1992;Rajadurai et al., 2005) and oxidative stress (Lesser, 2006) in *A. minutum*, leading
114 to an up-regulation of DMSP, DMS and DMSO production (McLenon and DiTullio,
115 2012;Sunda et al., 2002) in this high DMSP producer, which could ultimately lead to a shift in
116 the composition of the *A. minutum* microbiome.

117 **2. Methods**

118

119 *2.1. Culturing and experimental design*

120 Cultures of *Alexandrium minutum* (CS-324), isolated from Southern Australian coastal waters
121 (Port River, Adelaide, 11/11/1988, CSIRO, ANACC's collection) were grown in GSe medium
122 at 18°C and 50 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ under a 12:12 light:dark cycle. One month before the start
123 of each experiment, *A. minutum* cultures were acclimated over four generations to 20°C
124 (average summer temperature at Port River, IMOS) and 200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ using a
125 14:10 h light:dark cycle mimicking summer conditions. Light intensity was comparable to that
126 used in Berdalet et al. (2011) for *A. minutum* and conveniently allow to grow other algae
127 cultures using the same facilities. Cultures were grown to a cell concentration of $\sim 60,000$
128 mL^{-1} before cells were inoculated into fresh GSe medium. Six days prior to the start of each
129 experiment, 20 L of GSe medium was inoculated with a cell concentration of 1,140 mL^{-1}
130 (experiment 1, April 2016) and 680 mL^{-1} (experiment 2, June 2016) and aliquots of 500 mL
131 were transferred into 40 individual 750 mL sterile tissue culture flasks. Culture flasks were
132 incubated in four independent water baths (10 flasks in each) and maintained under control
133 conditions of 20°C and 200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. Temperature and light control was achieved
134 using circulating water heaters (Julabo, USA) and programmable LED lights (Hydra FiftyTwo,
135 AquaIllumination, USA). All cultures were mixed twice daily to keep cells in suspension by
136 gentle swirling.

137

138 On Day 1 (T_0), five culture flasks from each 20°C water bath were transferred to four new
139 water baths for exposure to experimental treatment temperatures (either 24°C experiment 1; or
140 32°C, experiment 2), so that each control and experimental water bath contained five flasks.
141 Experimental temperatures were carefully chosen based on preliminary experiments conducted
142 at 24°C, 28°C, 30°C and 32°C, where only a 12°C increase in temperature (32°C treatment)
143 led to a physiological stress response in this strain of *A. minutum* in culture. Although an
144 increase in temperature of this magnitude might be rare in coastal marine systems, this
145 presented a unique opportunity to investigate the consequences of MHW-induced thermal
146 stress on this relevant phytoplankton. One culture flask from each tank was immediately
147 sampled for baseline measurements of: DMS (2 mL), DMSP and DMSO (1 mL)
148 concentrations, photochemical efficiency (3 mL), algal and bacterial cell counts (1 mL), ROS
149 quantification (1 mL) and DNA extraction (~ 470 mL). The dissolved DMSP fraction was not
150 determined because preliminary investigations showed that gravity filtration was too time

151 consuming, potentially due to clogging of filters by the large *A. minutum* cells (30 μm
152 diameter), leading to filtration artefacts for DMSP analysis, as have previously been mentioned
153 by Berdalet et al. (2011). At 18:00 on Day 1 (T_6), 12:00 on Day 2 (T_{24}), 12:00 on Day 5 (T_{96})
154 and 12:00 on Day 6 (T_{120}), one flask from each of the eight water baths was removed from the
155 incubation conditions and sampled as described above.

156

157 *2.2. Photosynthetic efficiency measurements*

158 Subsamples for measurement of photosynthetic efficiency were dark adapted for 10 min under
159 aluminium foil and transferred to a quartz cuvette for Pulse Amplitude Modulated (PAM)
160 fluorometric analysis using a Water PAM (Walz GmbH, Effeltrich, Germany). Once the base
161 fluorescence (F_0) signal had stabilized (measuring light intensity 3, frequency 2s), a saturating
162 pulse (intensity 12, Width 0.8s) was used to measure the maximum quantum yield (F_v/F_m) of
163 photosystem II (PSII). As base fluorescence is dependent on cell density, the photomultiplier
164 gain was adjusted and recorded to maintain F_0 at a level of 0.2 a.u. before saturating the
165 photosystem. Samples were kept in suspension during measurements via continuous stirring at
166 minimal speed inside the quartz cuvette to avoid cells settling.

167

168 *2.3. Microalgal and bacterial cell counts*

169 Subsamples for bacterial cell counts were stained with SYBR Green at a final concentration of
170 1:10,000 and incubated in the dark for 15 min (Marie et al. 1997). Subsamples for microalgal
171 cell counts and stained subsamples for bacterial cell counts were diluted 1:10 and 1:100
172 respectively into sterile GSe medium prior to analysis with a BD Accuri C6 Flow Cytometer
173 (Becton Dickinson). Phytoplankton cells were discriminated using red auto-fluorescence and
174 side scatter (SSC), whereas bacterial populations were discriminated and quantified using
175 SYBR green fluorescence and SSC.

176

177 *2.4. Reactive oxygen species measurements*

178 The presence of reactive oxygen species (ROS) was detected within cultures using the
179 fluorescent probe 2,7-dichlorodihydrofluorescein-diacetate (CM-H2DCFDA; Molecular
180 Probes), which binds to ROS and other peroxides (Rastogi et al., 2010). The reagent was
181 thawed at room temperature for 10 min and activated using 86.5 μL of DMSO, with 5 μL of
182 activated reagent added to each sample (final concentration 5 μM). Samples were vortexed for
183 5 sec and incubated at room temperature for 30 min. Samples were then centrifuged at 2,000 g
184 for 2 min, the supernatant with reagent dye was discarded, and stained cells were resuspended

185 in 1 mL of PBS, prior to quantification of fluorescence by flow cytometry. Mean green
186 fluorescence was quantified from cytograms of forward light scatter (FSC) against green
187 fluorescence. A positive (+ 10 μ L of 30% H_2O_2 , final concentration 97mM) and negative (no
188 ROS added) control of PBS were run to ensure that detected cell fluorescence was completely
189 attributable to the ROS probe.

190

191 *2.5.Sulfur analysis by gas chromatography*

192 The preparation of all blanks and samples used in the dilution steps described below were
193 prepared with sterile (0.2 μ M filtered and autoclaved) phosphate-buffered saline (PBS, salinity
194 35ppt) to avoid cell damage from altered osmolarity and to maintain similar physical properties
195 as seawater during headspace analysis by gas chromatography. Aliquots for DMS analysis were
196 transferred into 14 mL headspace vials that were immediately capped and crimped using butyl
197 rubber septa (Sigma Aldrich Pty 27232) and aluminum caps (Sigma Aldrich Pty 27227-U),
198 respectively. DMSP aliquots were 1:1 diluted with sterile PBS and DMSP was cleaved to DMS
199 by adding 1 pellet of NaOH to each vial, which was immediately capped and crimped. Samples
200 were incubated for a minimum of 30 min at room temperature to allow for the alkaline reaction
201 and equilibration to occur prior to analysis by gas chromatography (Kiene and Slezak, 2006).

202

203 DMS and DMSP samples were analyzed by 500 μ L direct headspace injections using a
204 Shimadzu Gas Chromatograph (GC-2010 Plus) coupled with a flame photometric detector
205 (FPD) set at 180°C with instrument grade air and hydrogen flow rates set at 60 mL min^{-1} and
206 40 mL min^{-1} , respectively. DMS was eluted on a capillary column (30 m x 0.32 mm x 5 μ m)
207 set at 120°C using high purity Helium (He) as the carrier gas at a constant flow rate of 5 mL
208 min^{-1} and a split ratio of five. A six-point calibration curve and PBS blanks were run by 500
209 μ L direct headspace injections prior to subsampling culture flasks using small volumes of
210 concentrated DMSP.HCl standard solutions (certified reference material WR002, purity 90.3
211 \pm 1.8% mass fraction, National Measurement Institute, Sydney, Australia) that were diluted in
212 sterile PBS to a final volume of 2 mL. Detection limit was 50 nM for 500 μ L headspace
213 injections. Concentrations obtained in vials treated with NaOH accounted for both DMS and
214 DMSP. Consequently, DMSP concentration in each sample was obtained by subtracting the
215 corresponding DMS concentration.

216

217 Following DMS and DMSP analysis, alkaline samples used for DMSP analysis were uncapped
218 and left to vent overnight under a fume hood. On the next day, samples were purged for 10 min

219 with high purity N₂ at an approximate flow rate of 60 mL min⁻¹ to remove any remaining DMS
220 produced from the alkaline treatment. Samples were then neutralized by adding 80 µL of 32 %
221 HCl and DMSO was converted to DMS by adding 350 µL of 12 % TiCl₃ solution to each vial,
222 which was then immediately capped and crimped (Kiene and Gerard, 1994; Deschaseaux et al.,
223 2014b). Vials were then heated in a water bath at 50°C for 1h and cooled down to room
224 temperature prior to analysis by 500 µL direct headspace injections on the GC-FPD as
225 described above. A 5-point calibration curve was run prior to DMSO analysis using DMSO
226 standard solutions (Sigma Aldrich Pty, D2650) diluted in PBS to a final volume of 2 mL and
227 converted to DMS with TiCl₃ in the same manner as the experimental samples. PBS blanks
228 treated with NaOH and TiCl₃ were also run along with the calibration curves. All dimethylated
229 sulfur compounds were normalised to cell density, which best reflects biogenic production.

230

231 *2.6. DNA extraction*

232 Following sub-sampling for the physiological and biogenic sulfur measurements described
233 above, the remaining 400 mL within each culture flask was filtered onto a 47 mm diameter,
234 0.22 µm polycarbonate filter (Millipore) with a peristaltic pump at a rate of 80 rpm to retain
235 cells for DNA analysis. The filters were subsequently stored in cryovials, snap frozen with
236 liquid nitrogen and stored at -80°C until extraction. DNA extraction was performed using a
237 bead-beating and chemical lysis based DNA extraction kit (PowerWater DNA Isolation Kit,
238 MoBio Laboratories) following the manufacturer's instructions. DNA quantity and purity were
239 checked for each sample using a Nanodrop 2000 (Thermo Fisher Scientific, Wilmington, DE,
240 USA). Three replicate samples with the highest DNA quantity and purity from the control and
241 treatment tanks, collected at the beginning (T₀) and end (T₁₂₀) of the experiment, were
242 subsequently sequenced.

243

244 *2.7. 16S rRNA amplicon sequencing and bioinformatics*

245 To characterize the bacterial assemblage structure (microbiome) of *A. minutum* cultures, we
246 employed 16S rRNA amplicon sequencing. We amplified the V1-V3 variable regions of the
247 16S rRNA gene using the 27F (AGAGTTTGATCMTGGCTCAG, Lane, 1991) and 519R
248 (GWATTACCGCGGCKGCTG, Turner et al., 1999) primer pairing, with amplicons
249 subsequently sequenced using the Illumina MiSeq platform (Ramaciotti Centre for Genomics;
250 Sydney, NSW, Australia) following the manufacturer's guidelines. Raw data files in FASTQ
251 format were deposited in the National Center for Biotechnology Information (NCBI) Sequence
252 Read Archive (SRA) under the study accession number PRJNA486692.

253 Bacterial 16S rRNA gene sequencing reads were analysed using the QIIME pipeline (Caporaso
254 et al., 2010;Kuczynski et al., 2012). Briefly, paired-end DNA sequences were joined, de novo
255 Operational Taxonomic Units (OTUs) were defined at 97% sequence identity using UCLUST
256 (Edgar, 2010) and taxonomy was assigned against the SILVA v128 database (Quast et al.,
257 2012;Yilmaz et al., 2013). Chimeric sequences were detected using usearch61 (Edgar, 2010)
258 and together with chloroplast OTUs were filtered from the dataset. Sequences were then
259 aligned, filtered and rarefied to the same depth to remove the effect of sampling effort upon
260 analysis.

261

262 *2.8. Statistical analysis*

263 Repeated measures analysis of variance (rmANOVA) models were fitted to the data to quantify
264 the effects of temperature and time (fixed factors) on all response variables measured in this
265 experiment (cell density, F_V/F_M , ROS, DMS, DMSP and DMSO concentrations) using IBM
266 SPSS Statistics 20. Assumptions of sphericity were tested using Mauchly's test. In cases where
267 this assumption was violated, the degrees of freedom were adjusted using the Greenhouse-
268 Geisser correction factor. Bonferroni adjustments were used for pairwise comparisons. Each
269 variable was tested for the assumption of normality and log, ln or sqrt transformations were
270 applied when necessary.

271

272 For sequencing data, alpha diversity parameters of the rarefied sequences and Jackknife
273 Comparison of the weighted sequence data (beta diversity) were calculated in
274 QIIME (Caporaso et al., 2010). A two-way PERMANOVA with Bray-Curtis similarity
275 measurements was performed on abundance data of taxonomic groups that contained more
276 than 1% of total generated OTUs (represent 90.23% of the data) using PAST (Hammer et al.,
277 2008). In addition, PAST was used to perform non-metric multidimensional scaling (nMDS)
278 analysis and isolate the environmental parameters (normalised as follows: $(x-\text{mean})/\text{stdev}$) that
279 contributed the most to the differences between groups using the Bray-Curtis similarity
280 measure. SIMPER analysis performed with the White *t*-test was used to identify the taxonomic
281 groups that significantly contributed the most to the shift in bacterial composition in *A.*
282 *minutum* cultures over time and between temperature treatments.

283

284 **3. Results**

285

286 *3.1. Algal growth and physiological response*

287 *A. minutum* cell abundance exponentially increased over time in both the control (20°C) and
288 24°C temperature treatment, but a significantly faster growth rate ($p = 0.001$, t -test) occurred
289 at 24°C ($2.66 \pm 0.01 \text{ d}^{-1}$; average \pm SE) compared to the 20°C control ($2.57 \pm 0.01 \text{ d}^{-1}$), resulting
290 in significantly greater cell abundance at 96h ($p = 0.007$) and 120h ($p < 0.001$) (rmANOVA,
291 **Table 1, Fig. 1a**). On the other hand, the 32°C treatment resulted in decreased growth rates
292 ($2.40 \pm 0.02 \text{ d}^{-1}$ versus $2.58 \pm 0.02 \text{ d}^{-1}$; t -test) and significantly lower cell abundance, relative
293 to the 20°C control, at all time points from 6h after the start of the experiment ($p \leq 0.03$;
294 rmANOVA, **Table 1, Fig. 1b**). *A. minutum* abundance demonstrated a marked decline on day
295 5 in the 32°C treatment.

296

297 No significant difference in the maximum quantum yield (F_V/F_M) of *A. minutum* cultures
298 occurred between 20°C and 24°C until 120h after the start of the experiment, where a
299 significantly lower F_V/F_M occurred in the 24°C treatment ($p = 0.01$; rmANOVA, **Table 1, Fig.**
300 **2a**). In contrast, F_V/F_M was significantly lower in *A. minutum* cultures maintained at 32°C
301 compared to the 20°C control at all time points from 6h after the start of the experiment ($p \leq$
302 0.01 ; rmANOVA, **Table 1, Fig. 2b**). However, on days 5 and 6, the F_V/F_M of cultures kept at
303 32°C recovered to values (0.72 ± 0.008) close to those of the control (0.75 ± 0.004) (**Fig. 2B**),
304 although it remained significantly lower than at 20°C ($p < 0.01$ and $p < 0.001$ on day 5 and 6,
305 respectively.

306

307 3.2. Reactive oxygen species (ROS)

308 Significantly lower concentrations of ROS were measured at 24°C than at 20°C at 96h ($p =$
309 0.003) and 120h ($p = 0.03$) (rmANOVA, **Table 1, Fig. 2c**). In contrast, significantly greater
310 concentrations of ROS were measured at 32°C than at 20°C 24h ($p < 0.001$), 96h ($p = 0.001$)
311 and 120h ($p = 0.01$) after the start of the experiment (rmANOVA, **Table 1, Fig. 2d**). In-line
312 with the recovery in measured F_V/F_M , ROS concentrations in cultures kept at 32°C started to
313 decline to values closer to those of the control on days 5 and 6 of the experiment (**Fig. 2d**). A
314 significant negative correlation between F_V/F_M levels and ROS concentrations was observed
315 under the 32°C temperature treatment ($R^2 = 0.623$; $p = 0.02$, $n = 18$; **Fig. 3**).

316

317 3.3. Biogenic sulfur dynamics

318 Biogenic concentrations of DMSP, DMS and DMSO ranged from 424 ± 35 to $1629 \pm 170 \text{ fmol}$
319 cell^{-1} , from 13 ± 1.02 to $87 \pm 5 \text{ fmol cell}^{-1}$ and from 9 ± 1.41 to $94 \pm 24 \text{ fmol cell}^{-1}$, respectively,
320 over both experiments (**Fig. 4**). Concentrations of all three sulfur compounds slowly decreased

321 over time in all *A. minutum* cultures regardless of the temperature treatment. No significant
322 difference in DMSP concentration was recorded between 20°C and 24°C throughout the
323 experiment ($p > 0.05$; rmANOVA, **Table 1, Fig. 4a**), whereas significantly less DMSP was
324 measured in cells at 32°C than in the 20°C control at 96h ($p = 0.02$; rmANOVA, **Table 1, Fig.**
325 **4b**).

326

327 Significantly lower DMS concentrations were measured at 24°C compared to 20°C at 24h (p
328 < 0.001) and 120h ($p = 0.002$) (rmANOVA, **Table 1, Fig. 4c**). In contrast, DMS was
329 significantly higher at 32°C than 20°C 6h after the start of the experiment ($p = 0.008$;
330 rmANOVA, **Table 1, Fig. 4d**). A similar pattern was observed for DMSO, where relative to
331 the controls, it was significantly lower at 24°C 24h after the start of the experiment ($p = 0.001$;
332 rmANOVA, **Table 1, Fig. 4e**) and significantly greater at 32°C after 6h and 24h ($p < 0.05$, **Fig.**
333 **4f**).

334

335 *3.4. Bacterial abundance and composition*

336 Bacterial cell abundance exponentially increased over time at both 20°C and 24°C (**Fig. 5a**).
337 Bacterial abundance was significantly greater at 24°C than at 20°C 120 h after the start of the
338 experiment ($p = 0.05$; rmANOVA, **Table 1, Fig. 5a**). However, no significant difference ($p >$
339 0.05 , t -test) in bacterial growth rate was observed between 20°C ($4.15 \pm 0.05 \text{ d}^{-1}$) and 24°C
340 ($4.18 \pm 0.01 \text{ d}^{-1}$). In contrast, bacterial growth rate was significantly lower at 32°C than in the
341 20°C control ($4.05 \pm 0.01 \text{ d}^{-1}$ versus $4.23 \pm 0.02 \text{ d}^{-1}$; $p < 0.001$, t -test) (**Fig. 5b**), resulting in
342 significantly lower bacterial cell densities at 24h ($p = 0.002$), 96h ($p = 0.002$) and 120h ($p <$
343 0.001) relative to the control (rmANOVA, **Table 1, Fig. 5b**).

344

345 The composition of the initial (T_0) *A. minutum* microbiome was consistent across all samples,
346 but then diverged significantly with time and between temperature treatments (**Fig. 6a-b**; Bray-
347 Curtis similarity measurement, Shepard plot stress = 0.0587). A significant temporal shift in
348 bacterial composition occurred at both 20°C and 32°C, with dissimilarities in community
349 composition between T_0 and T_{120} of 27% and 42% occurring respectively (SIMPER analysis).
350 Notably, bacterial communities at 32°C differed significantly (two-way PERMANOVA; $p <$
351 0.05) to 20°C at T_{120} , with 32% dissimilarity in community composition. These differences
352 were primarily driven by increased relative abundance of bacterial Operational Taxonomic
353 units (OTUs) within the *Oceanicaulis* (17%), *Phycisphaeraceae SMIA02* (8.8%) and *Balneola*
354 (4.9%) genus along with a decline in the relative abundance of OTUs matching *Maribacter*

355 (24%), *Marinoscillum* (4.7%) and *Seohaecicola* (2.7%) (*Rhodobacter* family) in the 32°C
356 treatment (White test, **Fig. 6c**), with all taxa cumulatively contributing to 63% of the OTU
357 differences between temperature treatments at T₁₂₀ (SIMPER analysis). In the 32°C treatment,
358 differences in microbiome composition between T₀ and T₁₂₀ were aligned with the elevated
359 levels of ROS, while in the control (20°C) the community shift was principally aligned with
360 differences in bacterial and algal cell abundance (**Fig. 6a**; MDS analysis). Similarly, the
361 elevated concentration of ROS as well as the lower F_v/F_M, lower algal and bacterial cell
362 abundance and lower DMSP, DMS and DMSO concentrations at 32°C were aligned with the
363 differences in microbiome composition between the temperature treatments (**Fig. 6b**; MDS
364 analysis)

365

366 4. Discussion

367

368 Climate change induced shifts within marine ecosystems are predicted to fundamentally alter
369 the physiology of planktonic organisms and the biogeochemical transformations that they
370 mediate (Finkel et al., 2009; Tortell et al., 2008; Hallegraeff, 2010). Rising seawater
371 temperatures are one of the major impacts of climate change on marine ecosystems (Harley et
372 al., 2006), and can be manifested both as long-term gradual increases (IPCC, 2007, 2013) or
373 intense episodic marine heatwaves (Frölicher and Laufkötter, 2018; Hobday et al., 2016).
374 Although less examined to date than chronic temperature increases, MHWs are predicted to
375 become more frequent and severe (Oliver et al., 2018) and have been proposed as a mechanism
376 for triggering toxic algal blooms (Ummenhofer and Meehl, 2017). Against this backdrop of
377 changing environmental conditions, microbial production and cycling of dimethylated sulfur
378 compounds could be particularly relevant because they simultaneously play a role in the stress
379 response of marine phytoplankton (Berdalet et al., 2011; Deschaseaux et al., 2014a; Sunda et
380 al., 2002; Wolfe et al., 2002; Stefels and van Leeuwe, 1998) and have been predicted to have
381 biogeochemical feed-back effects that are relevant for local climatic processes (Charlson et al.,
382 1987).

383

384 This study investigated the biogenic sulfur cycling dynamics of *A. minutum*, and its
385 microbiome, in response to an intense, short-term thermal stress event, akin to the marine heat-
386 wave events occurring with increasing frequency within coastal habitats (Oliver et al., 2018).
387 Indeed, MHWs have been defined as an abrupt and ephemeral increase in temperature of at
388 least 3 to 5°C above climatological average that lasts for at least 3 to 5 days (Hobday et al.,
389 2016). Large increases in temperature of about 8°C above the monthly climatological average

390 led to red-tides of exceptional density in San Francisco Bay (Cloern et al., 2005). While a 12°C
391 increase in temperature constitutes an extreme scenario of MHWs, even for coastal habitats,
392 this experimental temperature was selected after preliminary investigations with the intention
393 to induce thermal stress in this strain of *A. minutum* in culture.

394

395 *A. minutum* has been targeted in this study as 1) an ecologically relevant phytoplankton
396 responsible for some of the most harmful algal blooms (Anderson et al., 2012) and 2) as
397 biochemically relevant for containing the highest DMSP concentrations ever reported in marine
398 dinoflagellates (Caruana and Malin, 2014). However, it is to be noted that DMSP
399 concentrations reported in this study were a degree of magnitude lower (0.42 ± 0.04 to $1.63 \pm$
400 1.70 pmol cell⁻¹) than that previously reported for *A. minutum* (14.2 pmol cell⁻¹; Caruana and
401 Malin, 2014; Jean et al., 2005). This is potentially because this culture of *A. minutum* had been
402 isolated from free-living *A. minutum* for a long time (1988) or because culturing conditions
403 failed to mimic the exact same biochemical conditions in which this strain of *A. minutum*
404 usually grow. This biochemical difference could potentially reflect that this strain of *A.*
405 *minutum* in culture was more robust than free-living dinoflagellates of the same species,
406 thereby potentially justifying the need of a 12°C increase in temperature to induce thermal-
407 stress.

408

409 4.1. Effects of thermal stress on *A. minutum* growth, physiology and ROS production

410 A 4°C increase in temperature resulted in faster algal growth and lower oxidative stress,
411 indicating that 24°C was close to a temperature optimum for this strain of *Alexandrium*. This
412 is perhaps not surprising considering that *Alexandrium* species are capable of growing under a
413 wide range of temperatures from 12°C to 25°C (Laabir et al., 2011). In contrast, a 12°C increase
414 in temperature resulted in a rapid and clear cascade of physiological responses, indicative of
415 an acute thermal stress response in *A. minutum*. Overall, *A. minutum* cells exposed to 32°C
416 immediately exhibited slower growth relative to the 20°C control, suggesting that a 12°C
417 increase in temperature rapidly led to either an increase in cell death rate or a decrease in cell
418 division (Rajadurai et al., 2005; Veldhuis et al., 2001). The slower growth rate at 32°C was
419 coupled with a drop in photosynthetic efficiency and an increase in ROS concentrations, which
420 are both common stress responses to thermal stress in marine algae (Lesser, 2006; Falk et al.,
421 1996; Robison and Warner, 2006; Iglesias-Prieto et al., 1992). In fact, these two physiological
422 responses are often interconnected as increased ROS production generally occurs in both the
423 chloroplast and mitochondria of marine algae exposed to thermal stress, causing lipid

424 peroxidation and ultimately leading to a loss in thylakoid membrane integrity (Falk et al., 1996)
425 and a decrease in the quantum yield of PSII (Lesser, 2006). This was reflected in the negative
426 correlation observed between the maximum quantum yield of PSII and ROS concentrations.

427

428 Although photosynthetic efficiency remained impaired and ROS concentrations remained high
429 under 32°C until the end of the experiment, both biomarkers of stress started to return to values
430 closer to those of the 20°C control by day 5 and 6 of the experiment. This was most likely at
431 the expense of a decline in algal abundance since slow growth often coincides with concurrent
432 cellular repair and photosystem activity recovery (Robison and Warner, 2006). The differential
433 physiological response between 24°C and 32°C indicates that although cultures of this strain
434 of *A. minutum* appear to be highly resistant to temperature changes, an abrupt increase in
435 temperature of 12°C simulating an extreme marine heatwave led to a clear stress response. The
436 physiological pattern at 32°C also suggested an acclimation period necessary for such an abrupt
437 shift in temperature, especially since recovery (in F_V/F_M and ROS levels) was observed towards
438 the end of the experiment.

439

440 *4.2. Biogenic sulfur cycling as a response to thermal stress in A. minutum*

441 Biogenic organic compounds are key compounds in the stress response of phytoplankton, with
442 evidence they can be used in responses to changes in temperature (Van Rijssel and Gieskes,
443 2002;Stefels, 2000). An up-regulation of the biogenic sulfur yield was expected as a stress
444 response to increased temperature in *A. minutum*, through either an increase in cellular DMSP
445 concentrations, or an increase in DMS via the cleavage of DMSP (McLenon and DiTullio,
446 2012;Berdalet et al., 2011;Wolfe et al., 2002;Sunda et al., 2002). No significant change in
447 DMSP concentrations was observed between the control and 24°C treatment, where, as
448 described above, physiological responses converged to indicate that 24°C was in fact a more
449 optimal growth temperature for this organism. This temperature optimum was generally
450 associated with lower DMS and DMSO concentrations than in the 20°C control, although this
451 was only evident 24h after the start of the experiment. Since algal stress responses often result
452 in increased cellular sulfur concentrations in dinoflagellates (McLenon and DiTullio,
453 2012;Berdalet et al., 2011), it is perhaps not surprising that DMS and DMSO concentrations
454 were lower under what appear to have been more optimal growth temperature conditions.

455

456 In contrast to the lower DMS and DMSO concentrations observed at 24°C compared to the
457 20°C control, exposure to 32°C resulted in spikes in DMS and DMSO 6h after the start of the

458 experiment, which accompanied decreased algal growth and impaired photosystem II.
459 Although sporadic, the increases in DMS and DMSO observed in the 32°C treatment may have
460 resulted from enhanced intracellular DMSP cleavage by phytoplankton (Del Valle et al., 2011)
461 or enhanced DMSP exudation from phytoplankton cells during cell lysis (Simó, 2001),
462 resulting in an increasing pool of dissolved DMSP made readily available to both bacteria and
463 phytoplankton DMSP-lyases (Riedel et al., 2015;Alcolombri et al., 2015;Todd et al.,
464 2009;Todd et al., 2007). However, it is notable that lower DMSP concentrations in the 32°C
465 treatment than in the control only occurred on day 4, whereas the spike in DMS and DMSO
466 were evident at the outset of the experiment (6h). Since this decrease in DMSP at 96h was not
467 coupled with an increase in DMS, this could alternatively be indicative of a decrease in
468 methionine synthase activity (McLenon and DiTullio, 2012) or assimilation of DMSP-sulfur
469 by bacterioplankton for *de novo* protein synthesis (Kiene et al., 2000), with this demethylation
470 pathway often accounting for more than 80% of DMSP turnover in marine surface waters. The
471 spike in DMSO measured 6h after the increase in temperature to 32°C most likely indicated
472 rapid DMS oxidation by ROS under thermal stress (Sunda et al., 2002;Niki et al., 2000). At
473 that time however, we found no evidence for ROS build up in *A. minutum* cultures, possibly
474 because ROS concentrations were kept in check by sufficient DMS synthesis and active DMS-
475 mediated ROS scavenging (Lesser, 2006;Sunda et al., 2002). In contrast, 24h after the start of
476 the experiment, increased ROS coincided with an abrupt decline in DMS and DMSO, perhaps
477 suggestive of serial oxidation via active ROS scavenging of both DMS to DMSO and DMSO
478 to methane sulfinic acid (MSNA) (Sunda et al., 2002), although it is always difficult to
479 confidently link DMS(O) and ROS dynamics unless using tracing techniques.

480

481 The only previous study that has examined sulfur responses to stress exposure in *A. minutum*
482 examined the effect of physical turbulence by shaking *A. minutum* cultures for up to four days
483 (Berdalet et al., 2011). While the authors of that study also observed slower cell growth as a
484 response to stress exposure, in contrast to our study, cellular DMSP concentrations increased
485 by 20%. Here, a drop in DMSP concentration was observed at 96h between the control and
486 temperature treatment. Therefore, even though DMSP concentrations were quantified with a
487 similar approach as in Berdalet et al. (2011) (no filtration of the samples with assuming that
488 particulate DMSP concentrations overrule dissolved DMSP and DMS concentrations), it seems
489 that heat stress and turbulence triggered a dissimilar sulfur response to stress in *A. minutum*.

490

491 Overall, a 12°C increase in temperature led to lower photosynthetic efficiency, increased
492 oxidative stress and slower cell growth in the red-tide mediating dinoflagellate *A. minutum*.
493 This physiological stress response was coupled with a differential biogenic sulfur cycling as
494 shown by spikes in DMS and DMSO as well as lower DMSP concentrations, most likely
495 translating ROS scavenging and DMSP uptake by bacterioplankton, respectively. Because the
496 turnover of DMS, DMSP and DMSO in biological systems can occur very quickly (Simo et al
497 2000), DMS and DMSO concentrations can change rapidly, which sometimes makes it difficult
498 to clearly establish cause-effect relationships between physiological stress and the biogenic
499 sulfur response.

500

501 4.3. A shift in *A. minutum* associated-bacteria composition triggered by thermal stress

502 In light of DMSP and related biogenic sulfur compounds constituting an important source of
503 carbon and sulfur to phytoplankton-associated bacteria (Kiene et al., 2000), it follows that any
504 shift in biogenic sulfur concentrations could influence the microbiome composition of *A.*
505 *minutum*. However, it is undeniable that a shift in the microbial community could also be driven
506 by a range of physiological and biochemical parameters that were not measured in this study.
507 Nevertheless, the most pronounced temporal shift in the composition of the bacterial
508 community associated with *A. minutum* occurred in the 32°C treatment. This shift was
509 primarily characterized by a statistically significant increase in the relative abundance of OTUs
510 classified as members of the *Oceanicaulis*, *Phycisphaeraceae* and *Balneola* and a significant
511 decrease in OTUs classified as members of the *Maribacter*, *Marinoscillum* and *Seohaecicola*.

512

513 To predict any potential role of these key OTUs in biogenic sulfur cycling processes, we
514 screened the genomes of members of these groups using BLAST for four genes commonly
515 involved in DMSP catabolism: *dmdA*, CP000031.2 (Howard et al., 2006); *dddP*, KP639186
516 (Todd et al., 2009); *tmm*, JN797862 (Chen et al., 2011); and *dsyB*, KT989543 (Kageyama et
517 al., 2018). A BLAST query of the sequences in the NCBI nucleotide collection (nr/nt) database
518 revealed that previously sequenced members of the genera *Maribacter* (taxid:252356, 357
519 sequences), *Oceanicaulis* (taxid:153232, 36 sequences), *Marinoscillum* (taxid:643701, 23
520 sequences), *Seohaecicola* (taxid:481178, 18 sequences) and *Balneola* (taxid:455358, 44
521 sequences) did not possess any homologs of these sulfur cycling genes. While no homologs
522 were found in the genus *SMIA02*, perhaps because very little genomic information is available
523 for this genus, a close phylogenetic relative to *SMIA02* (99% query cover, 80% identical, E-
524 value = 0.0), and also a member of the *Phycisphaeraceae* family (*P. mikurensis* 10266;

525 genbank accession numbers AP012338.1), possessed significant homologues to all four query
526 genes involved in DMSP metabolism: *dmdA* (92% identical, E-value < 0.001), *dddP* (87%
527 identical, E-value = 0.003), *tmm* (82% identical, E-value = 0.002) and *dsyB* (92% identical, E-
528 value < 0.001). It is thus possible that the spike in DMS and DMSO concentrations in the early
529 stage of the 32°C heat treatment was a consequence of (or contributed to) the preferential
530 recruitment of *Phycisphaeraceae* SM1A02.

531

532 Some members of the *Rhodobacter* family such as several members of the *Roseobacter* genus
533 and *Rhodobacter sphaeroides* are known to possess homologues of either or both *dmdA* and
534 *ddd* genes, which are responsible for DMSP demethylation and DMSP-to-DMS cleavage,
535 respectively (Howard et al., 2006;Curson et al., 2008). However, none of the available
536 reference genomes for *Seohaecicola*, a member of the *Rhodobacteraceae*, possessed any
537 homologs of targeted biogenic sulfur cycling. Similarly, members of the *Maribacter*, which
538 was the main contributor to the difference in microbiome structure between the control and
539 thermal stress treatment, are known not to possess DMSP/DMS transformation pathways
540 (Kessler et al., 2018). Hence, the decline of this taxa in the heat stress treatments, where an
541 upshift in biogenic sulfur availability occurred, is perhaps not surprising. However, this change
542 in microbial abundance could have also been triggered by a range of other parameters that were
543 not measured in this study.

544

545 Ultimately, the rapid changes in DMS and DMSO concentrations were potentially caused by
546 (or led to) a shift in microbiome composition towards the preferential growth of sulfur-
547 consuming bacteria (e.g. *Phycisphaeraceae* SM1A02) at the expense of other types of bacteria
548 (e.g. *Seohaecicola*). Alternatively, the observed shifts in microbiome structure may have
549 occurred independently to the biogenic sulfur cycling processes and was instead related to other
550 metabolic shifts in the heat-stressed *A. minutum*. Notably, the temporal shift in bacterial
551 composition under thermal stress was associated with increased cellular ROS at the end of the
552 experiment, indicating a potential link to oxidative stress.

553

554 **5. Conclusion**

555

556 Abrupt and intense increases in seawater temperatures associated with MHWs are predicted to
557 become more frequent and intense (Oliver et al., 2018) and have the potential to influence the
558 structure of coastal microbial assemblages and the nature of the important biogeochemical

559 processes that they mediate. Here, we hypothesized that a very acute increase in temperature,
560 mimicking extreme coastal MHWs, would trigger both a physiological and biochemical stress
561 response in the DMSP-producing dinoflagellate *A. minutum*. This response was indeed
562 observed following a 12°C-increase in temperature, with evidence for impaired photosynthetic
563 efficiency, oxidative stress, spikes in DMS and DMSO concentrations, a drop in DMSP
564 concentration and a shift in the composition of the *A. minutum* microbiome. These patterns are
565 indicative of a profound shift in the physiological state and biochemical function of this
566 ecologically relevant dinoflagellate in the context of MHWs and suggest that extreme thermal
567 stress has the potential to not only influence the composition and interactions of coastal
568 microbial food-webs, but re-shape sulfur budgets in coastal waters.

Acknowledgements

The work was funded by Australian Research Council grants FT130100218 and DP140101045 to JRS and KP. We thank Rendy Ruvindy and Associate Professor Shauna Murray for providing the *Alexandrium* cultures and soil extracts. Dr Bonnie Laverock provided advice on DNA extractions.

Author contribution:

ED, KP and JS devised the experimental design. ED and JOB conducted the thermal stress experiments, including sampling and sample analysis. NS and JOB processed sequencing data while ED processed the physiological and sulfur data. ED wrote the manuscript with significant contributions from all co-authors.

Competing interests:

The authors declare that they have no conflict of interest.

Figure captions

Figure 1 – Algal cell abundance in *A. minutum* cultures in experiment 1 (20°C and 24°C) (A) and experiment 2 (20°C and 32°C) (B); average \pm SE, $n = 4$.

Figure 2 – Photosynthetic efficiency (A, B) and reactive oxygen species (ROS) (C, D) in *A. minutum* cultures in experiment 1 (20°C and 24°C) (A, C) and experiment 2 (20°C and 32°C) (B, D); average \pm SE, $n = 4$.

Figure 3 – Correlation between the photosynthetic efficiency and reactive oxygen species (ROS) in *Alexandrium minutum* under the 32°C thermal stress treatment; $n = 18$.

Figure 4 – DMSP (A, B), DMS (C, D) and DMSO (E, F) concentrations in *A. minutum* cultures in experiment 1 (20°C and 24°C) (A, C, E) and experiment 2 (20°C and 32°C) (B, D, F); average \pm SE, $n = 4$.

Figure 5 – Bacterial cell abundance in *A. minutum* cultures in experiment 1 (20°C and 24°C) (A) and experiment 2 (20°C and 32°C) (B); average \pm SE, $n = 4$.

Figure 6 – Multi-dimensional scaling (MDS) of the three phylogenetic groups defined by 16s sequencing of the bacteria population associated with *A. minutum* cultures grown under control conditions (20°C) and acute thermal stress (32°C) at T₀ and T₁₂₀ (**A**) and MDS excluding the T₀ control (**B**). Vectors represent the factors that most likely drove the shift in bacterial composition between groups. The taxonomic groups that significantly contributed to the difference in bacterial composition between T₀ and T₁₂₀ at 32°C ⁽¹⁾, between T₀ and T₁₂₀ at 20°C ⁽²⁾ and between 32°C and 20°C at T₁₂₀ ⁽³⁾ appear in bold next to the heatmap (**C**), with scaling being based on relative abundance.

References

- Alcolombri, U., Ben-Dor, S., Feldmesser, E., Levin, Y., Tawfik, D. S., and Vardi, A.: Identification of the algal dimethyl sulfide-releasing enzyme: A missing link in the marine sulfur cycle, *Science*, 348, 1466-1469, 10.1126/science.aab1586, 2015.
- Anderson, D. M.: Physiology and bloom dynamics of toxic *Alexandrium* species, with emphasis on life cycle transitions, *Nato Asi Series G Ecological Sciences*, 41, 29-48, 1998.
- Anderson, D. M., Alpermann, T. J., Cembella, A. D., Collos, Y., Masseret, E., and Montresor, M.: The globally distributed genus *Alexandrium*: multifaceted roles in marine ecosystems and impacts on human health, *Harmful algae*, 14, 10-35, 2012.
- Berdalet, E., Llaveria, G., and Simó, R.: Modulation of dimethylsulfoniopropionate (DMSP) concentration in an *Alexandrium minutum* (Dinophyceae) culture by small-scale turbulence: A link to toxin production?, *Harmful Algae*, 11, 88-95, 10.1016/j.hal.2011.08.003, 2011.
- Brimblecombe, P., and Shooter, D.: Photooxidation of dimethylsulfide in aqueous-solution, *Mar. Chem.*, 19, 343-353, 1986.
- Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N., Peña, A. G., Goodrich, J. K., and Gordon, J. I.: QIIME allows analysis of high-throughput community sequencing data, *Nature methods*, 7, 335-336, 2010.
- Caruana, A. M. N., and Malin, G.: The variability in DMSP content and DMSP lyase activity in marine dinoflagellates, *Progress in Oceanography*, 120, 410-424, 10.1016/j.pocean.2013.10.014, 2014.
- Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulfur, cloud albedo and climate, *Nature*, 326, 655-661, 1987.
- Chen, Y., Patel, N. A., Crombie, A., Scrivens, J. H., and Murrell, J. C.: Bacterial flavin-containing monooxygenase is trimethylamine monooxygenase, *Proceedings of the National Academy of Sciences*, 201112928, 2011.
- Cloern, J. E., Schraga, T. S., and Lopez, C. B.: Heat wave brings an unprecedented red tide to San Francisco Bay, *Eos, Transactions American Geophysical Union*, 86, 66-66, 2005.
- Curson, A. R., Todd, J. D., Sullivan, M. J., and Johnston, A. W.: Catabolism of dimethylsulphoniopropionate: microorganisms, enzymes and genes, *Nat. Rev. Microbiol.*, 9, 849, 2011.
- Curson, A. R. J., Rogers, R., Todd, J. D., Brearley, C. A., and Johnston, A. W. B.: Molecular genetic analysis of a dimethylsulfoniopropionate lyase that liberates the climate-changing gas dimethylsulfide in several marine alpha-proteobacteria and *Rhodobacter sphaeroides*, *Environ. Microbiol.*, 10, 1099-1099, 10.1111/j.1462-2920.2008.01592.x, 2008.
- DeBose, J. L., Lema, S. C., and Nevitt, G. A.: Dimethylsulfoniopropionate as a foraging cue for reef fishes, *Science*, 319, 1356-1356, 10.1126/science.1151109, 2008.
- Del Valle, D.A., Slezak, D., Smith, C.M., Rellinger, A.N., Kieber, D.J., Kiene, R. P.: Effect of acidification on preservation of DMSP in seawater and phytoplankton cultures: Evidence for rapid loss and cleavage of DMSP in samples containing *Phaeocystis* sp, *Mar. Chem.*, 124 (1-4), 57-67, 2011.
- Deschaseaux, E. S. M., Jones, G. B., Deseo, M. A., Shepherd, K. M., Kiene, R. P., Swan, H. B., Harrison, P. L., and Eyre, B. D.: Effects of environmental factors on dimethylated sulphur compounds and their potential role in the antioxidant system of the coral holobiont, *Limnol. Oceanogr.*, 59, 758-768, 2014a.
- Deschaseaux, E. S. M., Kiene, R. P., Jones, G. B., Deseo, M. A., Swan, H. B., Oswald, L., and Eyre, B. D.: Dimethylsulphoxide (DMSO) in biological samples: A comparison of the TiCl₃

and NaBH₄ reduction methods using headspace analysis, *Mar. Chem.*, 164, 9-15, <http://dx.doi.org/10.1016/j.marchem.2014.05.004>, 2014b.

Edgar, R. C.: Search and clustering orders of magnitude faster than BLAST, *Bioinformatics*, 26, 2460-2461, 2010.

Falk, S., Maxwell, D. P., Laudenbach, D. E., and Huner, N. P.: Photosynthetic adjustment to temperature, in: *Photosynthesis and the environment*, Springer, 367-385, 1996.

Finkel, Z. V., Beardall, J., Flynn, K. J., Quigg, A., Rees, T. A. V., and Raven, J. A.: Phytoplankton in a changing world: cell size and elemental stoichiometry, *J. Plankton Res.*, 32, 119-137, 2009.

Frölicher, T. L., and Laufkötter, C.: Emerging risks from marine heat waves, *Nature Communications*, 9, 650, [10.1038/s41467-018-03163-6](https://doi.org/10.1038/s41467-018-03163-6), 2018.

Hallegraeff, G. M.: Ocean climate change, phytoplankton community responses, and harmful algal blooms: a formidable predictive challenge, *J. Phycol.*, 46, 220-235, [doi:10.1111/j.1529-8817.2010.00815.x](https://doi.org/10.1111/j.1529-8817.2010.00815.x), 2010.

Harley, C. D. G., Hughes, A. R., Hultgren, K. M., Miner, B. G., Sorte, C. J. B., Thornber, C. S., Rodriguez, L. F., Tomanek, L., and Williams, S. L.: The impacts of climate change in coastal marine systems, *Ecol. Lett.*, 9, 228-241, [10.1111/j.1461-0248.2005.00871.x](https://doi.org/10.1111/j.1461-0248.2005.00871.x), 2006.

Hobday, A. J., Alexander, L. V., Perkins, S. E., Smale, D. A., Straub, S. C., Oliver, E. C., Benthuyzen, J. A., Burrows, M. T., Donat, M. G., and Feng, M.: A hierarchical approach to defining marine heatwaves, *Progress in Oceanography*, 141, 227-238, 2016.

Howard, E. C., Henriksen, J. R., Buchan, A., Reisch, C. R., Bürgmann, H., Welsh, R., Ye, W., González, J. M., Mace, K., and Joye, S. B.: Bacterial taxa that limit sulfur flux from the ocean, *Science*, 314, 649-652, 2006.

Hughes, T. P., Kerry, J. T., Álvarez-Noriega, M., Álvarez-Romero, J. G., Anderson, K. D., Baird, A. H., Babcock, R. C., Beger, M., Bellwood, D. R., and Berkelmans, R.: Global warming and recurrent mass bleaching of corals, *Nature*, 543, 373, 2017.

Iglesias-Prieto, R., Matta, J. L., Robins, W. A., and Trench, R. K.: Photosynthetic response to elevated temperature in the symbiotic dinoflagellate *Symbiodinium microadriaticum* in culture, *Proc. Natl. Acad. Sci. U. S. A.*, 89, 10302-10305, 1992.

IPCC: Climate change 2007: The physical science basis. Contribution of working group II to the fourth assessment report of the Intergovernmental Panel on Climate Change, Cambridge, United Kingdom and New York, 2007.

IPCC: The physical science basis: working group I contribution to the fifth assessment report of the intergovernmental panel of climate change, Cambridge, United Kingdom and New York, 2013.

Jean, N., Boge, G., Jamet, J. L., Richard, S., Jamet, D.: Annual contribution of different plankton size classes to particulate dimethylsulfoniopropionate in a marine perturbed ecosystem, *J. Mar. Syst.*, 2005.

Joint, I., and Smale, D. A.: Marine heatwaves and optimal temperatures for microbial assemblage activity, *FEMS Microbiol. Ecol.*, 93, 2017.

Kageyama, H., Tanaka, Y., Shibata, A., Waditee-Sirisattha, R., and Takabe, T.: Dimethylsulfoniopropionate biosynthesis in a diatom *Thalassiosira pseudonana*: Identification of a gene encoding MTHB-methyltransferase, *Archives of biochemistry and biophysics*, 645, 100-106, 2018.

Kessler, R. W., Weiss, A., Kuegler, S., Hermes, C., and Wichard, T.: Macroalgal–bacterial interactions: Role of dimethylsulfoniopropionate in microbial gardening by *Ulva* (Chlorophyta), *Mol. Ecol.*, 27, 1808-1819, 2018.

Kiene, R. P., and Gerard, G.: Determination of trace levels of dimethylsulfoxide (DMSO) in seawater and rainwater, *Mar. Chem.*, 47, 1-12, 1994.

Kiene, R. P., Linn, L. J., and Bruton, J. A.: New and important roles for DMSP in marine microbial communities, *J. Sea. Res.*, 43, 209-224, 2000.

Kiene, R. P., and Slezak, D.: Low dissolved DMSP concentrations in seawater revealed by small-volume gravity filtration and dialysis sampling, *Limnol. Oceanogr. Meth.*, 4, 80-95, 2006.

Knight, K.: Hatchling loggerhead turtles pick up DMS, *J. Exp. Biol.*, 215, 2012.

Kuczynski, J., Stombaugh, J., Walters, W. A., González, A., Caporaso, J. G., and Knight, R.: Using QIIME to analyze 16S rRNA gene sequences from microbial communities, *Current protocols in microbiology*, 1E. 5.1-1E. 5.20, 2012.

Laabir, M., Jauzein, C., Genovesi, B., Masseret, E., Grzebyk, D., Cecchi, P., Vaquer, A., Perrin, Y., and Collos, Y.: Influence of temperature, salinity and irradiance on the growth and cell yield of the harmful red tide dinoflagellate *Alexandrium catenella* colonizing Mediterranean waters, *J. Plankton Res.*, 33, 1550-1563, 2011.

Lane, D. J.: 16S/23S rRNA sequencing, in: *Nucleic acid techniques in bacterial systematics*, edited by: Stackebrandt, E., and Goodfellow, M., John Wiley and Sons, New York, NY, 115-175, 1991.

Lesser, M. P.: Oxidative stress in marine environments: Biochemistry and physiological ecology, *Annu. Rev. Physiol.*, 68, 253-278, 10.1146/annurev.physiol.68.040104.110001, 2006.

McLenon, A. L., and DiTullio, G. R.: Effects of increased temperature on dimethylsulfoniopropionate (DMSP) concentration and methionine synthase activity in *Symbiodinium microadriaticum*, *Biogeochem.*, 110, 17-29, 10.1007/s10533-012-9733-0, 2012.

Nevitt, G. A., Veit, R. R., and Kareiva, P.: Dimethyl sulfide as a foraging cue for Antarctic procellariiform seabirds, *Nature*, 376, 680-682, 10.1038/376680ao, 1995.

Niki, T., Kunugi, M., and Otsuki, A.: DMSP-lyase activity in five marine phytoplankton species: its potential importance in DMS production, *Mar. Biol.*, 136, 759-764, 2000.

Oliver, E. C. J., Donat, M. G., Burrows, M. T., Moore, P. J., Smale, D. A., Alexander, L. V., Benthuyzen, J. A., Feng, M., Sen Gupta, A., Hobday, A. J., Holbrook, N. J., Perkins-Kirkpatrick, S. E., Scannell, H. A., Straub, S. C., and Wernberg, T.: Longer and more frequent marine heatwaves over the past century, *Nature Communications*, 9, 1324, 10.1038/s41467-018-03732-9, 2018.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O.: The SILVA ribosomal RNA gene database project: improved data processing and web-based tools, *Nucleic acids research*, 41, D590-D596, 2012.

Rajadurai, M., Poornima, E. H., Narasimhan, S. V., Rao, V. N. R., and Venugopalan, V. P.: Phytoplankton growth under temperature stress: Laboratory studies using two diatoms from a tropical coastal power station site, *Journal of Thermal Biology*, 30, 299-305, <https://doi.org/10.1016/j.jtherbio.2005.01.003>, 2005.

Rastogi, R. P., Singh, S. P., Häder, D.-P., and Sinha, R. P.: Detection of reactive oxygen species (ROS) by the oxidant-sensing probe 2', 7'-dichlorodihydrofluorescein diacetate in the cyanobacterium *Anabaena variabilis* PCC 7937, *Biochemical and biophysical research communications*, 397, 603-607, 2010.

Riedel, T., Spring, S., Fiebig, A., Scheuner, C., Petersen, J., Göker, M., and Klenk, H. P.: Genome sequence of the roseovarius mucosus type strain (Dsm 17069T), a bacteriochlorophyll a-containing representative of the marine roseobacter group isolated from the dinoflagellate *Alexandrium ostenfeldii*, *Standards in Genomic Sciences*, 10, 10.1186/1944-3277-10-17, 2015.

Robison, J. D., and Warner, M. E.: Differential impacts of photoacclimation and thermal stress on the photobiology of four different phylotypes of *Symbiodinium* (pyrrhophyta) 1, *J. Phycol.*, 42, 568-579, 2006.

Seymour J.R., Simó, R., Ahmed, T., Stocker, R. Chemoattraction to

dimethylsulfoniopropionate throughout the marine microbial food web, *Science*, 329 (5989), 342-345, 2010.

Simó, R., Pedrós-Alió, C., Malin, G., and Grimalt, J. O.: Biological turnover of DMS, DMSP and DMSO in contrasting open-sea waters, *Marine Ecology Progress Series*, 230, 1-11, 2000.

Simó, R.: Production of atmospheric sulfur by oceanic plankton: biogeochemical, ecological and evolutionary links, *Trends in Ecology & Evolution*, 16, 287-294, 2001.

Stefels, J., and van Leeuwe, M. A.: Effects of iron and light stress on the biochemical composition of Antarctic *Phaeocystis* sp. (Prymnesiophyceae). I. Intracellular DMSP concentrations, *J. Phycol.*, 34, 486-495, 10.1046/j.1529-8817.1998.340486.x, 1998.

Stefels, J.: Physiological aspects of the production and conversion of DMSP in marine algae and higher plants, *J. Sea. Res.*, 43, 183-197, 2000.

Sunda, W., Kieber, D. J., Kiene, R. P., and Huntsman, S.: An antioxidant function for DMSP and DMS in marine algae, *Nature*, 418, 317-320, 2002.

Todd, J. D., Rogers, R., Li, Y. G., Wexler, M., Bond, P. L., Sun, L., Curson, A. R. J., Malin, G., Steinke, M., and Johnston, A. W. B.: Structural and regulatory genes required to make the gas dimethyl sulfide in bacteria, *Science*, 315, 666-669, 10.1126/science.1135370, 2007.

Todd, J. D., Curson, A. R. J., Dupont, C. L., Nicholson, P., and Johnston, A. W. B.: The dddP gene, encoding a novel enzyme that converts dimethylsulfoniopropionate into dimethyl sulfide, is widespread in ocean metagenomes and marine bacteria and also occurs in some Ascomycete fungi, *Environ. Microbiol.*, 11, 1376-1385, 10.1111/j.1462-2920.2009.01864.x, 2009.

Tortell, P. D., Payne, C. D., Li, Y., Trimborn, S., Rost, B., Smith, W. O., Riesselman, C., Dunbar, R. B., Sedwick, P., and DiTullio, G. R.: CO₂ sensitivity of Southern Ocean phytoplankton, *Geophys. Res. Lett.*, 35, 2008.

Turner, S., Pryer, K. M., Miao, V. P., and Palmer, J. D.: Investigating deep phylogenetic relationships among cyanobacteria and plastids by small subunit rRNA sequence analysis, *Journal of Eukaryotic Microbiology*, 46, 327-338, 1999.

Ummenhofer, C. C., and Meehl, G. A.: Extreme weather and climate events with ecological relevance: a review, *Phil. Trans. R. Soc. B*, 372, 20160135, 2017.

Van Alstyne, K. L., Wolfe, G. V., Freidenburg, T. L., Neill, A., and Hicken, C.: Activated defense systems in marine macroalgae: evidence for an ecological role for DMSP cleavage, *Mar. Ecol. Prog. Ser.*, 213, 53-65, 2001.

Van Rijssel, M., and Gieskes, W. W. C.: Temperature, light, and the dimethylsulfoniopropionate (DMSP) content of *Emiliania huxleyi* (Prymnesiophyceae), *J. Sea. Res.*, 48, 17-27, 2002.

Veldhuis, M. J., Kraay, G. W., and Timmermans, K. R.: Cell death in phytoplankton: correlation between changes in membrane permeability, photosynthetic activity, pigmentation and growth, *European Journal of Phycology*, 36, 167-177, 2001.

Wolfe, G. V., Strom, S. L., Holmes, J. L., Radzio, T., and Olson, M. B.: Dimethylsulfoniopropionate cleavage by marine phytoplankton in response to mechanical, chemical, or dark stress, *J. Phycol.*, 38, 948-960, 10.1046/j.1529-8817.2002.t01-1-01100.x, 2002.

Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig, W., and Glöckner, F. O.: The SILVA and “all-species living tree project (LTP)” taxonomic frameworks, *Nucleic acids research*, 42, D643-D648, 2013.

Zhou, C. X., Xu, J. L., Yan, X. J., Hou, Y. D., and Jiang, Y.: Analysis of dimethylsulfide and dimethylsulfoniopropionate in marine microalgae culture, *Chin. J. Anal. Chem.*, 37, 1308-1312, 2009.

Table 1. Output of repeated measures analysis of variance (rmANOVA) for algal (CELLS_A) and bacterial (CELLS_B) cell abundance, photosynthetic efficiency (F_v/F_M), oxidative stress (ROS), dimethylsulfoniopropionate (DMSP), dimethylsulfide (DMS) and dimethylsulfoxide (DMSO) concentrations as a function of temperature (24°C or 32°C) and time. Numbers in bold indicate significant data based on the level of significance $p < 0.05$. df1 = numerator df; df2= denominator df.

Parameters	24°C – mild thermal stress			32°C – mild thermal stress			
	temperature	time	temperature × time	temperature	time	temperature × time	
CELLS _A	<i>F</i>	4.04	335	4.16	27.47	237.62	8.28
	<i>df1</i>	1	4	4	1	2.04	2.04
	<i>df2</i>	6	24	24	6	12.26	12.26
	<i>p</i>	0.91	< 0.001	0.01	< 0.001	< 0.001	0.005
CELLS _B	<i>F</i>	2.13	52.2	1.35	32.56	199.8	22.26
	<i>df1</i>	1	1.29	1.29	1	4	4
	<i>df2</i>	6	7.74	7.74	6	24	24
	<i>p</i>	0.2	< 0.001	0.3	0.001	< 0.001	< 0.001
F _v /F _M	<i>F</i>	0.42	33.43	6.90	48.79	12.58	13.11
	<i>df1</i>	1	4	4	1	1.19	1.19
	<i>df2</i>	6	24	24	5	5.93	5.93
	<i>p</i>	0.54	< 0.001	0.001	0.001	0.01	0.01
ROS	<i>F</i>	37.26	6.30	5.88	33.23	8.85	8.41
	<i>df1</i>	1	4	4	1	2.32	2.32
	<i>df2</i>	6	24	24	6	13.9	13.9
	<i>p</i>	0.001	0.001	0.002	0.001	0.003	0.003
DMSP	<i>F</i>	0.79	31.16	0.95	3.03	15.18	3.17
	<i>df1</i>	1	1.56	1.56	1	4	4
	<i>df2</i>	6	9.35	9.35	6	24	24
	<i>p</i>	0.41	< 0.001	0.4	0.13	< 0.001	0.03
DMS	<i>F</i>	51.5	38.73	2.01	5.08	30.77	5.23
	<i>df1</i>	1	2.14	2.14	1	4	4
	<i>df2</i>	6	12.87	12.87	6	24	24
	<i>p</i>	< 0.001	< 0.001	0.17	0.07	< 0.001	0.004
DMSO	<i>F</i>	36.56	26.64	7.21	4.68	14.74	7.14
	<i>df1</i>	1	4	4	1	4	4
	<i>df2</i>	6	24	24	6	24	24
	<i>p</i>	0.001	< 0.001	0.001	0.07	< 0.001	0.001

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

