

Dear Editor,

We deeply appreciate you for giving us an opportunity to improve our manuscript again. We would like to thank all of you and the two reviewers for the thoughtful and valuable suggestions on our manuscript entitled "**Plant functional traits determined the latitudinal variations in soil microbial functions: evidence from a forest transect in China**" [ID: bg-2018-499]. According to the comments and suggestions, we have carefully revised our manuscript. We have followed the formatting requirements as presented in the Guide for Authors. We have uploaded the document of the responses to reviewer and a clean manuscript. Here are the point-to-point responses (color-coded blue) to Editor's and **Reviewer's** comments (color-coded black). The page and line numbers mentioned here refer to our latest revision of our manuscript simultaneously submitted.

Reviewer #1:

Authors significantly improved this manuscript using reviewers' comments to clarify four hypotheses that motivate the work and explain numerous abbreviations. The large number of measurements made on each forest soil sample provides a great deal of correlative insight into factors that affect respiration potential. A few questions remain, which I expect the authors can address through clarification or some minor recalculations.

1) The Biolog and respiration rate experiments measured only aerobic activities. While these may be the most significant contributors to C mineralization, the authors should clarify that anaerobic microbial decomposition mechanisms may be important for intermittently saturated soils (such as CB at the time of sampling – Table 2) and sites with high mean annual precipitation. Such activities may have different effects on organic acid and DOC concentrations.

Response: The SOM decomposition and Biolog analysis were conducted in 2013-2014. We have no samples now to conduct the experiment about anaerobic microbial decomposition. We have added discussion about the anaerobic microbial decomposition mechanisms as “Some forest soils were intermittently saturated (such as CB, Table 2) or high with mean annual precipitation. Under the anaerobic conditions, soil organic decomposition is mediated by a complex suite of microbial processes (Meronigal et al., 2004). The fermentation products including low molecular weight alcohols, fatty acids, and dihydrogen can serve as substrates for anaerobic respiration using a variety of alternative terminal electron acceptors in place of oxygen to mineralize organic carbon to carbon dioxide (CO₂). Therefore, the soil organic matter decomposition rate might be slow in these anaerobic conditions. These results demonstrate that the reduction of organic matter is a key step of anaerobic decomposition (Keller and Takagi, 2013).” (P13-14, Line 356-364)

2) Were the significance tests of Pearson correlation coefficients (Figs 5-7 & S2) corrected for multiple hypothesis testing? Please explain in methods.

Response: All P values were adjusted using the Bonferroni correction to account for multiple comparisons. We have added the explanation in the methods. (P10, Line 266-267)

3) As explained in the cited Biolog methods paper by Guckert et al. Biolog results must be corrected for inoculum size. Others have used Average Well Color Development (AWCD) to normalize values, but that does not appear to be done here. Did the authors normalize values before integration for Figure 3 using microbial biomass or another indicator of inoculum size?

Response: To minimize the influence of cell density in comparisons among samples, results can be analyzed at constant average well color development (AWCD). The AWCD for each microplate was calculated by subtracting the control well optical density (OD) from the substrate well OD (blanked substrate wells), setting any resultant blanked substrate wells with negative values to 0 and taking the mean of the 95 blanked substrate wells.(P9,Line 238-242)

General comments

1. In previous comment I mentioned that “Please set more appropriate and clearer hypothesis”. And in this version, numbers of hypothesis increased from 3 to 4. I think multiple hypotheses might cause complex or complicated discussion and conclusion even in case two. In this version, authors added “information about the relationships between soil microbial communities and their functions in natural ecosystems is urgently needed” as necessity of this research after presenting 2 conflicting hypotheses. I agree with the opinion of

field data-oriented validation's necessity. But simple hypothesis or aim to conduct 4 testing is required. Moreover, in case conducting 4 testing, relation between these testing must be described. For example, in section 4.2, hypothesis 1 seemed to be described as counter hypothesis of 2. And these hypotheses should be integrated in conclusion. I could not find any comment about latitudinal variations in conclusion.

Response: we have rewritten our hypotheses as “(1) The profiles of soil microbial substrate use vary along a latitudinal gradient, (2) biogeographical patterns of soil microbial substrate use are constrained by climate and plant functional traits, and (3) different soil microbial communities may have substrate use profiles and SOM decomposition rates.” (P6, Line 165-168)

We have rewritten this part as “There was no significant latitudinal pattern in the soil total microbial metabolic intensity (S) in our study, which was inconsistent with hypothesis (1). However, there was significant latitudinal variation in the use of the six different carbon sources. The soil microbial use of carbon sources were relatively higher in tropical and subtropical climatic area than those in temperate climatic area.” (P14, Line 374-377)

We have added description on the latitudinal pattern of soil microbial use of different carbon sources in the conclusion part as “There was no obvious latitudinal pattern for the soil total microbial metabolic intensity. However, soil microbial use of the different carbon sources varied with the latitude except the

amines carbon source. Soil microbial use of carbon sources were relatively higher in tropical climatic areas.” (P18, Line 491-494)

2. Title of this MS include “latitudinal variations”. And I mentioned as “I could not know about latitudinal distribution in soil microbial functions but comparison between forest and forest types”. In this version authors mentioned that “plant traits have more influence on soil microbial functions than climate” and if it is correct, I think title of this MS can be “Plant functional traits determine soil microbial function”. However, in this case, careful discussion must be needed. Does only MAT represent variation of climate? How do authors count indirect effect of climate (at least climate condition regulate forest type)? Etc. For example, in the paper of Tu(2016), their conclusion was not “climate is not main regulator” but “This study suggests that even microbial subcommunities follow general biogeographic patterns”. And they used several indexes reflecting latitudinal gradient.

Response: we have added the description on the latitudinal pattern of soil microbial functions as “There was no significant latitudinal pattern in the soil total microbial metabolic intensity in our study, which was inconsistent with hypothesis (1). However, there was significant latitudinal variation in the use of different carbon sources. The soil microbial use of carbon sources were relatively higher in tropical and subtropical climatic area than those in temperate climatic area.” (P14, Line 374-377)

We have conducted the path analysis relevant environmental variables to soil microbial use of different carbon sources (Table S4-9, supporting information). We found that the MAT and MAP had indirect effect on the soil microbial use of different carbon sources through influencing the soil temperature, soil total nitrogen, and total phosphorus. (P14, Line 381-383)

We have written this part as “Consistent with hypothesis (2), soil microbial functions were similar in closely related to the CWM values of the tree species (Fig. 4). However, only MAP of climatic factors had a moderate effect on the soil microbial function (Fig. 4). Climate may have indirect effect on the latitudinal pattern of different carbon sources use. In our study, MAT and MAP affected the six different carbon sources use indirectly by influencing the soil temperature (ST), soil total nitrogen (TN), and total phosphorus (TP) (Table S4-9). It was reported that climate was the main environmental parameters driving the latitudinal patterns of community-level leaf traits through the regulation of species composition (Wang et al., 2016). Woody plant species with evergreen broadleaves dominated in the tropic regions characterized by hot, humid, and infertile habitats showed low CWM values of leaf N and SLA and high CWM values of LDMC (Table S2).” (P14, Line 378-387)

Technical comments

1. (P2, L36 37 38) Meaning of intensities are not clear.

Response: They means the soil microbial substrate use intensities. We have added the full meanings of them in the text. (P2, Line 36-38)

2. (P2, L39) What index do authors use as Plant functional traits and climate?

Response: the indexes included in the plant functional traits were leaf dry matter content, leaf C concentrations, and leaf N concentrations. The climate index was the mean annual precipitation. We have added them in the abstract. (P2, Line 39-41)

3. (P3) Abbreviations needs climate conditions.

Response: Done (P3, Line 65-67).

4. (P17, L462) Contribution of silt content is totally common.

Response: we have rewritten this part as “For the first time, we showed that CWM values of plant traits explained variations in soil microbial C substrate use at the regional scale. Additionally, the fine-grained soils with high silt contents were higher in the microbial C substrate use.” (P18, Line 494-498)

Thanks again to you and the two reviewers for the thoughtful and thorough comments.

We hope that our revisions will be satisfactory, and we are very happy to work with you and the reviewers to resolve any remaining problems.

Yours sincerely,

Zhiwei Xu, Guirui Yu, Qiufeng Wang, Xinyu Zhang, Ruili Wang, Ning Zhao,

Nianpeng He, Ziping Liu

A list of all relevant changes

1. Affiliations of the first authors (P1,line7-12)
2. Abstract: (P2,line36-41)
3. Abbreviations (P3,line65-67)
4. Introduction (P6,line165-168)
5. Material and methods(2.4(P9,line238-242); 2.6(P10,line266-267;
line275-278)
6. Result (P11,line281,287-289)
7. Discussion (4.1 (P13-14,line356-364); 4.2(P14,line374-387)
8. Conclusion (P18,line 491-498)
9. References (P20,line 574-575; 594-595; P21,line 673-674)

1 **Plant functional traits determine latitudinal variations** 2 **in soil microbial function: evidence from forests in** 3 **China** 4

5 Zhiwei Xu^{1,2,3}, Guirui Yu^{4,5,*}, Qiufeng Wang^{4,5,*}, Xinyu Zhang^{4,5}, Ruili Wang⁶, Ning Zhao⁷, Nianpeng
6 He^{3,4}, Ziping Liu^{1,2,3}

7 1. Key Laboratory of Geographical Processes and Ecological Security of Changbai Mountains,
8 Ministry of Education; School of Geographical Sciences, Northeast Normal University,
9 Changchun, 130024, China

10 2. Institute for Peat and Mire Research, Northeast Normal University, Changchun, 130024, China

11 3. Jilin Provincial Key Laboratory for Wetland Ecological Processes and Environmental Change in
12 the Changbai Mountains, Changchun, 130024, China

13 4. Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic
14 Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 10010, China.

15 5. College of Resources and Environment, University of Chinese Academy of Sciences, Beijing,
16 100190, China

17 6. College of Forestry, Northwest A&F University, Yangling, 712100, China

18 7. Key Laboratory of Remote Sensing of Gansu Province, Heihe Remote Sensing Experimental
19 Research Station, Cold and Arid Regions Environmental and Engineering Research Institute,
20 Chinese Academy of Sciences, Lanzhou 730000, China

21
22
23 * Corresponding author at: Key Laboratory of Ecosystem Network Observation and Modeling,
24 Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences,
25 Beijing 100101, China.No. 11A, Datun Road, Chaoyang District, Beijing, 100101, China. Tel.: +86-
26 10-64889268; fax: +86 10 64889432.

27 E-mail: yugr@igsnr.ac.cn (G. Y.), qfwang@igsnr.ac.cn (Q. W.)

28 **Abstract.** Plant functional traits have increasingly been studied as determinants of ecosystem
29 properties, especially for soil biogeochemical processes. While the relationships between biological
30 community structures and ecological functions are a central issue in ecological theory, these
31 relationships remain poorly understood at the large scale. We selected nine forests along the North–
32 South Transect of Eastern China (NSTEC) to determine how plant functional traits influence the
33 latitudinal pattern of soil microbial functions, and how soil microbial communities and functions are
34 linked at the regional scale. We found that there was considerable latitudinal variation in the profiles
35 of different substrate use along the NSTEC. Specifically, we found that the substrate use by
36 microorganisms was highest in the temperate forest soils (soil microbial substrate use intensities of
37 10–12), followed by the subtropical forest soils (soil microbial substrate use intensities of 7–10), and
38 was least in the coniferous forest soils (soil microbial substrate use intensities of 4–7). The latitudinal
39 variation in soil microbial function was more closely related to plant functional traits (leaf dry matter
40 content, leaf C concentrations, and leaf N concentrations, $P=0.002$) than climate (Mean annual
41 precipitation, $P=0.022$). The soil silt, leaf dry matter, and leaf C and N contents were the main
42 controls on the biogeographical patterns of microbial substrate use in these forest soils. The soil
43 microbial community structures and functions were significantly correlated along the NSTEC. Soil
44 carbohydrate and polymer substrate use were mainly related to soil G⁺ bacterial and actinomycic
45 PLFAs, while the use of amine and miscellaneous substrates were related to soil G⁻ bacterial and
46 fungal PLFAs. The enzyme production varied with changes in the soil microbial communities. The
47 soil enzyme activities were positively correlated with the bacterial PLFAs but were not correlated
48 with the fungal PLFAs. The soil organic matter (SOM) decomposition rates were significantly higher
49 in the temperate forests than in the subtropical and tropical forests, emphasizing the rapid
50 degradability of high-energy substrates, such as soil microbial biomass carbon, carbohydrates, and
51 amino acids. The SOM decomposition rates were significantly and negatively related to soil dissolved
52 organic carbon concentrations, carboxylic acids, polymers, and miscellaneous substrate use. The
53 relationships between soil PLFAs and microbial substrate use, enzyme activities, and SOM
54 decomposition rate show that, as the soil microbial community structure changes, soil
55 biogeochemical processes also change.

| | | |
|----|---------------------------------------|---------------------------------------|
| 56 | Abbreviations | |
| 57 | NSTEC | North-South Transect of Eastern China |
| 58 | AWCD | Average well color development |
| 59 | RDA | Redundancy analysis |
| 60 | Soil microbial community | |
| 61 | PLFAs | Phospholipid fatty-acids |
| 62 | G ⁺ | Gram positive bacteria |
| 63 | G ⁻ | Gram negative bacteria |
| 64 | F/B | Fungi/Bacteria |
| 65 | Climate conditions | |
| 66 | MAT: Mean annual temperature | |
| 67 | MAP: Mean annual precipitation | |
| 68 | Soil enzyme activities | |
| 69 | BG | β-glucosidase |
| 70 | NAG | N-acetylglucosaminidase |
| 71 | AP | Acid phosphatase |
| 72 | LAP | Leucine aminopeptidase |
| 73 | Soil properties | |
| 74 | SMC | Soil moisture content |
| 75 | SOM | Soil organic matter |
| 76 | SOC | Soil organic carbon |
| 77 | TN | Total Nitrogen |
| 78 | DOC | Dissolved organic carbon |
| 79 | MBC | Microbial biomass carbon |
| 80 | Silt | Soil silt fractions (<53 μm) |
| 81 | Plant functional properties: | |
| 82 | CWM | Community-weighted means |
| 83 | SLA | The specific leaf area |
| 84 | LDMC | Leaf dry matter content |
| 85 | Leaf C | Leaf C concentrations |
| 86 | Leaf N | Leaf N concentrations |

87 **1 Introduction**

88 The catabolic diversity of soil microbial communities is a useful indicator of how microbial functions
89 adapt to environmental stress. It can be used to test fundamental questions about soil biological
90 resistance and resilience (Jagadamma et al., 2014; Swallow and Quideau, 2015), and help us
91 understand the role of microbial communities in different environments (Preston-Mafham et al.,
92 2002). Biological community structure and function are intimately linked in ecological processes,
93 and their relationships are a central issue in ecological theory (Talbot et al., 2014). Therefore, a major
94 goal in ecological research is to identify and understand the mechanisms and relationships that control
95 the structure and function of microbial communities over large spatial scales.

96 Numerous studies have documented how environmental and anthropogenic perturbations
97 impact on the structure, diversity (Tu et al., 2016; Zhou et al., 2016), and enzyme activities (Peng
98 and Wang, 2016; Xu et al., 2017) of soil microbial communities, and have reported that forests in the
99 same climatic zone develop similar microbial communities. Other researchers have examined spatial
100 patterns in soil microbial function at different scales. For example, Tian et al. (2015), from their study
101 of Changbai Mountain, China, found that the soil microbial metabolic activity and functional
102 diversity were spatially dependent. Others reported that soil microbial activities varied by forest type,
103 with high local variation and significant separation along regional climate gradients (Brockett et al.,
104 2012; Cao et al., 2016). Soil microbes from different climatic zones have different affinities for
105 carbon substrates. For example, microorganisms from boreal pine forest soils used carboxylic acids
106 more efficiently, but decomposed amino acids much less efficiently, than microorganisms from
107 temperate forest soils (Klimek et al., 2016). The soil microbial metabolic abilities are also influenced
108 by the dominant tree species, through the production of chemically-unique litter and root exudates,
109 and the soil physico-chemical properties (Menyailo et al., 2002). Despite this, because of limitations
110 in analytical methods, questions still remain about how soil microbial functions vary at the regional
111 scale.

112 The functional diversity of soil microbial communities is regulated by physico-chemical soil
113 properties (Gartzia-Bengoetxea et al., 2016), climate (Cao et al., 2016), and the composition of plant
114 cover (Sherman and Steinberger, 2012). For example, the geographic patterns in soil microbial

115 activities mainly reflect the climate, soil pH, and total phosphorus concentrations over large
116 geographic scales (Cao et al., 2016). Research has shown that substrate-induced respiration rates
117 were higher in soil microbial communities that developed under beech and holm oak forests than
118 under oak and pine forests (Gartzia-Bengoetxea et al., 2016). Plant functional traits have increasingly
119 been studied as determinants of ecosystem properties, especially for soil biogeochemical processes
120 (De Vries et al., 2012; Pei et al., 2016). Soil bacteria phospholipid fatty-acids (PLFAs) were found
121 to be positively correlated with the community-weighted means (CWM) of plant functional traits
122 (leaf nitrogen (N) concentration) (De Vries et al., 2012). The plant leaf dry matter content and the
123 leaf carbon (C) to nitrogen (N) ratio both influence the multivariate soil microbial community
124 structure, and these factors positively promote the abundances of specific microbial functional groups
125 (Pei et al., 2016). Limited soil resources, particularly in tropical forests, mean that soil
126 microorganisms may be more reliant on plants than soil for C and nutrients via rhizosphere exudation
127 or litter production, which varies among plant species (Russell et al., 2007; Raich et al., 2014; Waring
128 et al., 2015). While soil functional diversity has been used as an indicator of microbial metabolic
129 potential, there have been few studies of the integrated effects of climate, vegetation, and soil
130 substrate availability on large-scale soil microbial functional diversity.

131 Although the functional characteristics of soil microorganisms are at least as important as their
132 patterns of community structure in biogeochemical studies, the links between microbial community
133 structure and microbial functions are poorly understood. There are two current hypotheses about how
134 microbes determine ecosystem process rates. In functional redundancy, different microbes perform
135 the same function and so changes in the microbial community structure do not necessarily lead to a
136 change in soil function (Balsler and Firestone, 2005; Strickland et al., 2009). For example, Banerjee
137 et al. (2016) showed that the abundance of different bacterial and fungal groups changed by up to
138 300-fold under straw- and nutrient-amended treatments but that the decomposition rate remained
139 similar, indicating possible functional redundancy. The functional redundancy hypothesis has
140 recently been challenged by a counter-hypothesis, referred to as functional dissimilarity, which
141 suggests that diversity brings stability, and that every species plays a unique role in ecosystem
142 function (Fierer et al., 2007; Waldrop and Firestone, 2006). Soil microbial community composition
143 therefore, combined with environmental variables, may ultimately determine ecosystem process rates.

144 Waldrop and Firestone (2006) showed that gram positive bacteria (G^+) were mainly responsible for
145 the decomposition of pine needles and soil organic matter, but gram negative bacteria (G^-) were
146 mainly responsible for the decomposition of starch and xylose, which are easy to break down.
147 Philippot et al. (2013), when studying the diversity of denitrifiers, showed that the loss of microbial
148 diversity could result in decreases of between 4- and 5-fold in denitrification activity. In the
149 Mediterranean, losses in the mass of decomposing leaf litter from shrub species accelerated as
150 detritivore assemblages became more functionally dissimilar (Coulis et al., 2015). Research to date
151 suggests that the different microbial communities will result in variations in soil microbial function
152 and soil biochemical processes, so information about the relationships between soil microbial
153 communities and their functions in natural ecosystems is urgently needed.

154 The North-South Transect of Eastern China (NSTEC) extends from a cold temperate coniferous
155 forest in the north to a tropical rainforest in the south, and includes almost all the forest types found
156 in the Northern Hemisphere (Zhang and Yang, 1995) (Fig. 1 and Table 1). This transect, therefore,
157 provides the optimal environment for investigating large-scale geographical patterns in microbial
158 communities and their responses to environmental changes. In this study, we examined spatial
159 patterns in soil labile C concentrations, soil organic matter (SOM) decomposition rates, and
160 metabolic activity and functional diversity of microbes in nine forest biomes along the NSTEC. We
161 assessed how abiotic factors, such as climate, soil physical and chemical properties, and biotic factors,
162 in the form of community-weighted means (CWM) of plant functional traits, contributed to soil
163 functional diversity at the regional scale. We also examined the links between soil microbial
164 community structure (PLFAs) and function (SOM decomposition rate, enzyme activities, and
165 microbial substrate use). We tested four hypotheses in this study, as follows: (1) The profiles of soil
166 microbial substrate use vary along a latitudinal gradient, (2) biogeographical patterns of soil
167 microbial substrate use are constrained by climate and plant functional traits, and (3) different soil
168 microbial communities may have substrate use profiles and SOM decomposition rates.

169 **2 Material and methods**

170 **2.1 Study area and soil sampling**

171 We selected nine forest ecosystems along the NSTEC, namely Huzhong (HZ), Liangshui (LS),
172 Changbai (CB), Dongling (DL), Taiyue (TY), Shennong (SN), Jiulian (JL), Dinghu (DH), and
173 Jianfeng (JF) (18°44'–51°46'N, 128°53'–108°51'E) (Fig. 1, Table 1). Further information about the
174 soil types and sites has been documented previously by Xu et al. (2017). Forest soils have been
175 classified following the U.S. soil taxonomy and are described in Table 1 (Soil Survey Staff, 2010),
176 where information about the climate and the dominant vegetation at each site is also presented.

177 Soil samples were collected from four random plots at each site in July and August 2013, as
178 described previously by Xu et al. (2017). Briefly, we established four sampling plots measured 30 ×
179 40 m and collected soil samples from a depth of between 0 and 10 cm at between 30 and 50 points
180 in each plot along an S-shape. On return to the laboratory, the fresh soil samples were immediately
181 sieved through a 2-mm mesh and subdivided into three subsamples. One subsample was stored briefly
182 at 4 °C until analysis for soil enzyme activities and soil pH. Another was stored briefly at –20 °C
183 until analysis for PLFAs and Eco-Biolog. The third was air-dried, sieved through a 0.25 mm mesh,
184 and analyzed for soil nutrients.

185 2.2 Soil chemical analyses

186 Soil pH was measured at a soil-to-water ratio of 1:2.5. The soil moisture content (SMC) was
187 measured gravimetrically on 20 g fresh soil that was oven-dried at 105 °C to constant weight
188 immediately on arrival at the study sites' laboratories (Liu et al., 2012a). Soil organic carbon (SOC)
189 and total N (TN) concentrations were determined by dry combustion of ground samples (100-mesh)
190 in a C/N analyzer (Elementar, Vario Max CN, Germany). Total phosphorus (TP) was determined
191 with a flow injection auto-analyzer following digestion with H₂SO₄-HClO₄ (Huang et al., 2011).
192 After extraction with distilled water at a soil:distilled water ratio of 1:5, dissolved organic carbon
193 (DOC) concentrations were determined by Liqui TOC II (Elementar, Liqui TOC II, Germany) (Jones
194 and Willett, 2006). Soil microbial biomass carbon (MBC) was measured using the chloroform
195 fumigation and direct extraction technique (Vance et al., 1987). A conversion factor of 2.64 was used
196 to convert extracted C to biomass C. The silt fractions (<53 μm) of the samples were separated by
197 wet-sieving and then were freeze-dried in the laboratory, as described by Six et al. (2000). The soil
198 properties are shown in Table 2. We followed the method described by Bååth et al. (2003) for PLFA

199 analysis and PLFAs are expressed in units of nmol g^{-1} . The four enzymatic activities of β -glucosidase
200 (BG), N-acetylglucosaminidase (NAG), acid phosphatase (AP), and leucine aminopeptidase (LAP)
201 responsible for soil C, N, and phosphorous cycling, were measured following the procedure outlined
202 in Saiya-Cork et al. (2002) and are expressed in units of $\text{nmol h}^{-1} \text{g}^{-1}$. Information about PLFA and
203 enzyme activities are presented in Table S1.

204 The Biolog-ECO plates were purchased from Biolog, US. The substrates for BG, NAG, AP, and
205 LAP were 4-MUB- β -D-glucoside, 4-MUB-N-acetyl-b-D-glucosaminide, 4-MUB-phosphate, and L-
206 Leucine-7-amino-4-methylcoumarin, and were stored at -20°C . An MUB standard was used for the
207 BG, NAG, and AP enzymes and an AMC standard was used for the LAP enzyme. The substrates and
208 standards were purchased Sigma. Analytical grade reagents were used for the soil nutrient analysis.

209 2.3 Vegetation data

210 We established four sampling plots (30×40 m) in each forest ecosystem. In each plot, we recorded all
211 the tree individuals, and measured the height and diameter-at-breast-height (DBH) of each woody
212 individual with a $\text{DBH}\geq 2$ cm. The diversity of the tree species in the sampling plots was represented
213 by H' , and the diversity (H' , Shannon-Wiener) of the tree species in the community was calculated
214 as follows:

$$215 \quad H' = \sum_{i=0}^n (P_i \ln P_i)$$

216 Where P_i was the importance value of the species i as a proportion of all species, and n was the
217 number of the species.

218 We also calculated the community-weighted means (CWM) values of the tree traits using the
219 cover of each tree. As described by Xu et al. (2018), we collected litter and sun-exposed and mature
220 leaves (leaf blades for trees) from between five and ten individuals of each plant species at each site
221 and determined their TN and TC concentrations. We calculated the specific leaf area (SLA, the one-
222 sided area of a fresh leaf divided by its oven-dried mass, $\text{m}^2 \text{kg}^{-1}$), leaf dry matter content (LDMC,
223 the oven-dried mass of a leaf divided by its water-saturated fresh mass, mg g^{-1}), leaf C concentrations
224 (leaf C, g kg^{-1}), and leaf N concentrations (leaf N, g kg^{-1}) for ten fully expanded leaves of each
225 sampled individual. To measure the leaf traits at the community level, we calculated the CWM of the

226 tree layer, as follows:

$$227 \quad CWM = \sum_{i=1}^n p_i \times \text{trait}_i$$

228 Where p_i is the relative contribution of the species i to the cover of the whole community, n is the
229 number of the most abundant species, and trait_i is the trait value of species i , as described by Garnier
230 *et al.* (2004). The diversity of the tree species and plant functional traits are summarized in Table S2.

231 2.4 Microbial substrate use

232 Microbial functional diversities were determined using a Biolog EcoPlate™ (Biolog Inc., Hayward,
233 California, USA) as described by Garland and Mills (1991). Briefly, approximately 10 g of fresh soil
234 was suspended in 100 ml saline solution (0.85% NaCl) and shaken on an orbital shaker for 30 min at
235 190 rpm. A 150 µl aliquot of supernatant from 1:1 000 dilutions of each soil sample was added to
236 each well. The plates were incubated at 25°C, and the absorbance at 590 nm was measured using a
237 microplate reader (GENios Pro™, Tecan Trading AG, Männedorf, Switzerland) every 24 h up to 240
238 h (0, 24, 48, 72, 96, 120, 144, 168, 192, 216, and 240 h). **To minimize the influence of cell density in
239 comparisons among samples, results can be analyzed at constant average well color development
240 (AWCD). The AWCD for each microplate was calculated by subtracting the control well optical
241 density (OD) from the substrate well OD (blanked substrate wells), setting any resultant blanked
242 substrate wells with negative values to 0 and taking the mean of the 95 blanked substrate wells.**

243 The Richness (R), Shannon-Weiner diversity index (H'), Shannon evenness index (E), and
244 Simpson dominance index (D) were calculated from the absorption values after EcoPlate™
245 incubation for 96 h (Gomez *et al.*, 2006). Additionally, the 31 C sources were divided into six groups,
246 namely carbohydrates, carboxylic acids, amines, amino acids, polymers, and miscellaneous, as
247 suggested by Zak *et al.* (1994). The average absorbance of all C sources within each group was
248 computed as the intensity of the single substrate use. The soil total microbial metabolic intensities (S)
249 of six carbon sources were estimated by the area underneath *AWCD* vs. t , and were obtained by
250 integrating the equation against time t (Guckert *et al.*, 1996):

$$251 \quad S = \sum [(v_i + v_{i-1})/2 \times (t_i + t_{i-1})]$$

252 Where v_i was the average optical density of the i th incubation time.

253 2.5 SOM decomposition rate

254 Four replicates from each sampling site with a 60% water-holding capacity were incubated at 20°C.
255 In brief, 40 g of each fresh soil sample were put into a 150-ml incubation bottle, and the samples
256 were then adjusted so that their moisture content corresponded to a water-holding capacity of 60%.
257 During the 4-week incubation period, the soil respiration rates were measured on days 1, 7, 14, 21,
258 and 28 using an automatic system. The SOM decomposition rates were calculated as described in the
259 study of Xu et al. (2015).

260 2.6 Statistical analysis

261 One-way analysis of variance (ANOVA) followed by a post hoc Tukey HSD test were used to test
262 the significance of the differences among the soil properties, C use, functional diversity, and SOM
263 decomposition rates in the different forest ecosystems. We tested the relationships between labile C,
264 soil microbial community structure, microbial function, and the SOM decomposition rates with the
265 Pearson correlation test. Differences were considered significant when $P < 0.05$, with the marginal
266 significance set at $P < 0.01$. All P values were adjusted using the Bonferroni correction to account for
267 multiple comparisons.

268 We used redundancy analysis (RDA) to examine the relationship between the environmental
269 variables and soil microbial substrate use. The environmental variables were the same as those
270 described in Xu et al. (2018), including climate, soil properties, litter properties, and plant functional
271 traits. Before RDA, we conducted forward selection of the environmental variables that were
272 significantly correlated with variations in the microbial substrate use profile using stepwise
273 regression and the Monte Carlo Permutation Test. We used CANOCO software 4.5 (Ter Braak and
274 Smilauer 2002) for the RDA and stepwise regression. The environmental properties, which were
275 significantly correlated with the microbial substrate use in the RDA, were stressed in the plots. Path
276 analysis was conducted to examine the direct and indirect effect of biotic and abiotic factors on soil
277 microbial use of carbon sources. All ANOVA, regression analyses, and path analysis were performed
278 using SPSS 19.0 for Windows. Data are reported as the mean \pm SE.

279 **3 Results**

280 3.1 Patterns in the microbial substrate use, soil labile carbon, and SOM decomposition rates

281 There was no obvious latitudinal pattern for the soil total microbial metabolic intensity (Fig.2). Of
282 the forests along the NSTEC, the C metabolic intensity of soil microbes was lowest in HZ and LS;
283 the C metabolic intensity of soil microbes differed significantly between JF and the other forests (Fig.
284 2), which indicates that the color development was significantly higher in the tropical forest soils
285 than in the subtropical and temperate forest soils and is consistent with the variations in the AWCD
286 (Fig. S1). The average values of R , H' , and D were significantly different among the nine forest soils
287 and were highest in JF, SN, and CB (Table 3). Generally, the soil microbial use of the six different
288 carbon sources were relatively higher in tropical and subtropical climatic area (low latitude) than
289 those in temperate climatic area (high latitude) (Fig.3).

290 Across the nine forests, soil microorganisms used the six substrate groups in the same order; the
291 carboxylic acid substrate was used most, followed by amino acids, carbohydrates, polymers, amines,
292 and miscellaneous substrates (Fig. 3). Microorganisms in the boreal and temperate forests mainly
293 metabolized carbohydrates, amino acids, and carboxylic acids, while those from the subtropical and
294 tropical forests used the substrates in equal proportions. The substrate microbial use ability was
295 highest in the coniferous broad-leaved mixed forest and tropical forest soils, and lowest in the
296 coniferous forest soil (Fig. 3).

297 Overall, soil MBC concentrations in the boreal and temperate forests were three to eight times
298 higher than those of the subtropical and tropical forests. In contrast, the average DOC concentrations
299 in the tropical and subtropical forest soils, which ranged from 311 to 458 mg kg⁻¹, were significantly
300 higher than the average concentrations in the temperate and boreal forest soils, where the average
301 concentrations ranged from 204 to 284 mg kg⁻¹ (Table 2). The average SOM decomposition rates in
302 the subtropical forests ranged from 0.64 to 2.42 μg C g⁻¹ d⁻¹, and were significantly lower than the
303 rates in the temperate forests, which ranged from 3.43 to 4.61 μg C g⁻¹ d⁻¹ (Table S3).

304 3.2 Effect of environmental properties on soil microbial substrate use

305 Redundancy analysis showed that the variations in soil microbial substrate use were strongly and
306 positively correlated with the CWM values of LDMC, leaf N, and leaf C, and strongly and negatively

307 correlated with the soil silt content and SMC (Fig. 4). The RDA2 of soil microbial substrate use was
308 strongly positively correlated with TN and SOC, but negatively correlated with the mean annual
309 precipitation (MAP) (Fig. 5). RDA1 mainly represented the plant functional traits, soil texture, and
310 micro-meteorological conditions, while RDA2 represented climate and soil nutrients. Overall, the
311 soil silt content and the CWM values of plant functional traits were the main predictors of the
312 latitudinal variation in the soil microbial substrate use along the NSTEC.

313 3.3 Relationships between soil microbial substrate use, enzyme activities, and PLFAs

314 Microbial carbohydrate use was positively related with bacterial biomass and actinomycic biomass
315 (Fig. 5). Microbial polymer use was negatively related with bacterial biomass and actinomycic
316 biomass. Microbial amines use was negatively related with G^- bacterial and fungal biomass.
317 Miscellaneous substrate use was positively related with fungal biomass and G^+/G^- bacterial biomass
318 (Fig. 5).

319 The abundance of G^- bacteria was positively associated first with the specific activities of BG,
320 whereas actinomycetes and G^+ bacteria were positively associated with BG and LAP. Soil fungi were
321 negatively associated with BG (Fig. 5).

322 3.4 Relationships between SOM decomposition rate, PLFAs, enzyme activity, and microbial 323 metabolic activities

324 The SOM decomposition rates were significantly and positively related to soil MBC concentrations
325 but significantly and negatively related to soil DOC concentrations (Fig. 6a and b). Except for amino
326 acid and amine substrates, the SOM decomposition rates were significantly and positively related to
327 microbial metabolic activities (AWCD) and carbohydrate substrate use (Fig. 6c and d) and negatively
328 related to carboxylic acid, polymer, and miscellaneous substrate use (Fig. 6e, g, and i).

329 The SOM decomposition rates were significantly and positively correlated with total PLFAs
330 ($r=0.456$, $P=0.005$), bacteria ($r=0.3836$, $P=0.021$), actinomycetes ($r=0.500$, $P=0.002$), and G^-
331 bacteria PLFAs ($r=0.520$, $P=0.001$) (Fig. 7a, b, d, and f) but were negatively correlated with fungal
332 PLFAs ($r=-0.370$, $P=0.026$), F/B ($r=-0.513$, $P=0.001$), and the G^+/G^- ($r=-0.496$, $P=0.002$) (Fig. 7c,
333 g, and h). Except for LAP activity, soil enzyme activities were significantly and positively correlated

334 with the SOM decomposition rates ($P < 0.01$) (Fig. 7i, j, and l).

335 **4 Discussion**

336 4.1 Response of soil labile C and SOM decomposition rates to variations in forest type

337 Soil organic matter is one of the most important C pools in terrestrial ecosystems. The concentrations
338 of soil DOC in the temperate forests were lower than those in subtropical forests but the soil MBC
339 concentrations were higher in temperate forests than in subtropical forests. This reflects the results
340 of previous regional and global studies (Tian et al., 2010; Xu et al., 2013), and shows that the
341 production/consumption ratio of soil DOC was lower, but that microbial C immobilization was higher,
342 in the high latitude forests than closer to the tropics (Fang et al., 2014). Soil DOC, as a labile SOM
343 fraction with a rapid turnover, is one of the primary energy sources for microorganisms. The higher
344 temperatures and precipitation in subtropical and tropical forests lead to higher turnover rates (Fang
345 et al., 2014), so soil DOC concentrations were highest in subtropical, and MBC concentrations were
346 lowest, in tropical forests. However, in temperate forests, more C is assimilated into microbial
347 biomass, so that less C is lost through chemical and physical processes (Liu et al., 2010). Also,
348 because the decomposition ability of different microbe groups varies, the differences in the soil
349 microbial communities in different forest ecosystems may also be responsible for the spatial
350 variations in the soil DOC and MBC concentrations along the NSTEC (Hagedorn et al., 2008).

351 Heterotrophic soil respiration is sustained by the decomposition of SOM. The SOM
352 decomposition rates along the NSTEC were greater in temperate forests than in subtropical forests,
353 which was consistent with the variations in the soil MBC and SOC concentrations. These results
354 indicate that, as found in other studies, large scale SOM decomposition rates are driven by the
355 amounts of substrate available (Yu et al., 2010). Changes in the availability of C in SOM may affect
356 the microbial resource strategies, which may in turn influence the SOM decomposition rate. **Some**
357 **forest soils were intermittently saturated (such as CB, Table 2) or high with mean annual precipitation.**
358 **Under the anaerobic conditions, soil organic decomposition is mediated by a complex suite of**
359 **microbial processes (Megonigal et al., 2004). The fermentation products including low molecular**
360 **weight alcohols, fatty acids, and dihydrogen can serve as substrates for anaerobic respiration using a**

361 variety of alternative terminal electron acceptors in place of oxygen to mineralize organic carbon to
362 carbon dioxide (CO₂). Therefore, the soil organic matter decomposition rate might be slow in these
363 anaerobic conditions. These results demonstrate that the reduction of organic matter is a key step of
364 anaerobic decomposition (Keller and Takagi, 2013).

365 4.2 Mechanisms driving latitudinal variations in microbial substrate use

366 The AWCD reflects the sole C source use ability of the soil microbial community (Garland and Mills,
367 1991). Of the six groups of C substrates, microbial communities in the temperate forests mainly used
368 carbohydrates, carboxylic acids, and amino acids, which suggests that microorganisms in temperate
369 forests probably use high-energy substrates that degrade easily (Kunito et al., 2009). The carbon
370 substrate use was lowest in the boreal coniferous forest (HZ). This shows that, compared with
371 coniferous species, broadleaved tree species produce root exudates and litter high in water-soluble
372 sugars, organic acids, and amino acids that are more favorable for microbial activity (Priha et al.
373 2001).

374 There was no significant latitudinal pattern in the soil total microbial metabolic intensity (S) in
375 our study, which was inconsistent with hypothesis (1). However, there was significant latitudinal
376 variation in the use of the six different carbon sources. The soil microbial use of carbon sources were
377 relatively higher in tropical and subtropical climatic area than those in temperate climatic area.
378 Consistent with hypothesis (2), soil microbial functions were similar in closely related to the CWM
379 values of the tree species (Fig. 4). However, only MAP of climatic factors had a moderate effect on
380 the soil microbial function (Fig. 4). Climate may have indirect effect on the latitudinal pattern of
381 different carbon sources use. In our study, MAT and MAP affected the six different carbon sources
382 use indirectly by influencing the soil temperature (ST), soil total nitrogen (TN), and total phosphorus
383 (TP) (Table S4-9). It was reported that climate was the main environmental parameters driving the
384 latitudinal patterns of community-level leaf traits through the regulation of species composition
385 (Wang et al., 2016). Woody plant species with evergreen broadleaves dominated in the tropic regions
386 characterized by hot, humid, and infertile habitats showed low CWM values of leaf N and SLA and
387 high CWM values of LDMC (Table S2).

388 A growing number of studies reported that vegetation type, land use, soil nutrients, and soil

389 organic matter quality and quantity can determine large scale patterns of microbial communities (de
390 Vries et al., 2012; Tu et al., 2016). Plant functional traits that are related to growth may determine a
391 tree species' ability to contribute to the soil carbon pool via leaf litter inputs. For example, it was
392 previously reported that plant traits such as the leaf N content, SLA, and LDMC could explain
393 variations in soil nutrients and litter decomposition rates (Eichenberg et al., 2014; Laughlin, 2011).
394 Therefore, we examined how these plant traits influenced the soil microbial function by latitude. We
395 found that changes in the soil microbial C substrate use with latitude were mainly related to the soil
396 silt contents and the CWMs of LDMC, and leaf C and leaf N concentrations, which indicates that the
397 quality of nutrients from plants had a major influence on microbial carbon use efficiency (Hypothesis
398 (2)). Plant species with high a SLA, high leaf N concentrations, and low LDMC can produce
399 bacterial-dominated soil microbial communities in grasslands (Orwin et al., 2010). Looking beyond
400 individual traits, related tree species may cultivate microbial communities with similar preferences
401 for carbon sources through coevolution of plants and microbes (Liu et al., 2012b; Buscot, 2015).

402 As hypothesized, the soil microbial community composition was explained by the CWMs of
403 plant traits at the regional scale. Carbon substrate use was negatively correlated with the CWM of
404 leaf N concentrations (Table S2, Fig. S2). Bacterially dominated soil microbial communities develop
405 from leaf litter comprised of N-rich leaves from fast growing species (De Vries et al., 2012), while
406 leaves with low N concentrations will promote fungal domination (Orwin et al., 2010; De Vries et
407 al., 2012). In line with this, fungal biomass decreased, and bacterial biomass increased, as the CWM
408 leaf N content increased, and is associated with fast-growing, N-exploitative plants (Xu et al., 2018).
409 Leaf N concentrations are considered as indicators of plant growth and resource uptake (Wright et
410 al., 2004). The results from this study show that, along the NSTEC, high leaf N restrained microbial
411 C substrate use (Fig.S2) and was a good indicator of the competition between plants for soil N (Pei
412 et al., 2016). Soil microbes and nearby plants may have been competing for N in the soil.

413 We also found that the C substrate use was negatively correlated with the CWM of leaf C
414 concentrations (Table S2, Fig. S2). Plants at high latitudes may have higher leaf C concentrations
415 than plants at lower latitudes so that they can balance the osmotic pressure of cells and resist freezing
416 (Millard et al., 2007; Hoch and Körner, 2012). The increased C was most likely in the form of an
417 increase in non-structural C, including starch, low molecular weight sugars, and storage lipids that

418 are easy to break down. Therefore, soil microorganisms from the temperate forests mainly
419 metabolized high-energy substrates, such as carbohydrates, carboxylic acids, and amino acids.

420 The LDMC is the ratio of the leaf dry weight to the fresh weight and has been used as a proxy
421 for the ratio of structural compounds to assimilatory tissue (mesophyll and epidermis, Van Arendonk
422 and Poorter, 1994). High values of LDMC indicate large amounts of vascular tissue, cellulose,
423 insoluble sugars, and leaf lignin that are difficult to decompose (Poorter and Bergkotte, 1992); C
424 substrates such as carbohydrates, carboxylic acid, and amino acid are, however, easy to decompose
425 (Myers et al., 2001). In line with this, the use of carbohydrate, carboxylic acid, and amino acid
426 substrates was negatively related to the CWMs of the LDMC (Table S2). Pei et al. (2016) reported
427 that the LDMC was an important driver of multivariate soil microbial community structure and G^-
428 bacterial abundance.

429 Soil texture regulates soil biological processes and so affects the soil microbial community
430 structure (Sessitsch et al., 2001). In the present study, microbial C substrate use was significantly and
431 positively related to the soil silt content. Soil types and textures varied along the NSTEC. Soil texture
432 influences how microbes use organic matter, and has a strong influence on soil moisture, nutrient
433 availability, and retention (Veen and Kuikman, 1990). Fine-textured soils with a higher silt content
434 are known to be more favorable for bacterial growth than soils with a lower silt content because of
435 their greater water-holding capacity and nutrient availability, and because they are better protected
436 from bacterial grazers (Carson et al., 2010). We found that the microbial C substrate use was higher
437 in LS, CB, SN, and JL than in the other forests, reflecting their fine-grained soils and high silt contents,
438 which ranged from 60% to 80%.

439 4.3 Links between soil microbial community structure and function

440 The soil microbial community structure and functions were significantly correlated along the NSTEC.
441 Soil carbohydrate and polymer substrate use were mainly related to soil G^+ bacterial and actinomycic
442 biomass, but amines and miscellaneous substrates were mainly related to soil G^- bacterial, fungal
443 biomass, and the F/B ratio. Soil bacteria mainly decomposed simple carbohydrates, organic acids,
444 and amino acids, whereas soil fungi mainly decomposed recalcitrant compounds (Myers et al., 2001;
445 Treonis et al., 2004). Consistent with this, soil bacterial PLFAs were positively correlated with the

446 carbohydrate substrate use and the fungal PLFAs were positively related with miscellaneous substrate
447 use. The results are similar to those reported by Sterner and Elser (2002), who found that fungi tended
448 to have higher C/N or C/P ratios while heterotrophic bacteria typically have lower C/N or C/P ratios.

449 Shifts in microbial community composition may influence enzyme production (DeForest et al.,
450 2012; Waldrop et al., 2000; Brockett et al., 2012). Different microbial groups require different
451 amounts of nutrients to construct biomass, or have enzymes that differ in their affinity for nutrients.
452 We found that the relative abundances of the G⁺ bacteria and actinomycetes communities were
453 associated with the specific activities of BG, AP, and LAP), whereas the relative abundance of the
454 G⁻ bacteria was correlated with soil NAG activities involved in chitin degradation. In agreement with
455 our study, numerous other researchers have reported significant correlations between PLFA profiles
456 and enzyme activities (Waldrop et al., 2000; DeForest et al., 2012; Brockett et al., 2012; Riah-Anglet
457 et al., 2015). Soil BG was mainly responsible for cellulose degradation and was involved in breaking
458 down complex organic compounds (cellobiose) into small molecule substrates (glucose) in favor of
459 acquiring C through microbial community growth (Waldrop et al., 2000). Soil NAG activities were
460 weakly and positively related with fungal biomass in the present study, and may have been mainly
461 produced by fungal populations (Valášková et al., 2007). Fungi are commonly considered as
462 producers of oxidative enzymes. Therefore, the influence of fungal biomass on variations in enzyme
463 activities was minimal (Kivlin and Treseder, 2014). The linkages between enzyme activity and
464 community composition may provide some insight into the microbial mechanisms that drive the
465 decomposition of macromolecular C compounds. These results suggest that that overall ecosystem
466 functioning may suffer if soil microbial groups are lost.

467 The quality and amounts of SOM are influenced by the biomass, vegetation coverage, root
468 distribution, and microbial species (Raich and Schlesinger, 1992). The SOM decomposition rates
469 were higher in temperate forests than in tropical forests and may reflect the higher soil microbial
470 biomass (Wang et al., 2016). In line with this, SOM decomposition rates were positively related with
471 soil MBC concentrations and different groups of PLFAs. The inverse relationships between SOM
472 decomposition rates and DOC, carboxylic acids, polymers, and miscellaneous along the NSTEC,
473 indicate a shift in the soil C turnover from open to closed with increases in the soil labile C
474 concentrations (Fang et al., 2014). Soil DOC and MBC influence SOM decomposition rates indirectly

475 by regulating microbial properties (Boberg et al., 2014; Wei et al., 2014). In our study, SOM
476 decomposition rates were positively related with bacterial PLFAs but negatively with fungal PLFAs.
477 Because different communities of microbes have different SOM use efficiencies (Balsler and Wixon,
478 2009; Lipson et al., 2009; Monson et al., 2006), changes in the microbial community structure may
479 influence the microbial activities and the decomposition rates of organic matter (Lipson et al., 2009;
480 Keiblinger et al., 2010). The functional dissimilarity of microbes and fungi may help explain these
481 results. However, we did not measure some key variables, such as the microbial competition and
482 interactions, and relationship between the microbial diversity and the decomposition rates. Therefore,
483 in the future, we will use different experimental techniques that will help us gain an improved
484 understanding of the mechanisms that drive the relationships between the structure and function of
485 microbial communities.

486 **5 Conclusions**

487 In this study, we examined the patterns in labile C concentrations, SOM decomposition rates,
488 microbial substrate use, and functional diversity and identified a combination of abiotic and biotic
489 factors that influenced soil microbial functional diversity at the regional scale. The MBC
490 concentration and SOM decomposition rates were significantly lower, and the soil DOC
491 concentrations were higher in the subtropical and tropical forests than in the temperate forests. **There**
492 **was no obvious latitudinal pattern for the soil total microbial metabolic intensity. However, soil**
493 **microbial use of the different carbon sources varied with the latitude except the amines carbon source.**
494 **Soil microbial use of carbon sources were relatively higher in tropical climatic areas. For the first**
495 **time, we showed that CWM values of plant traits explained variations in soil microbial C substrate**
496 **use at the regional scale. Additionally, the fine-grained soils with high silt contents were higher in**
497 **the microbial C substrate use. Climate factors affected the soil microbial uses of carbon sources**
498 **indirectly by influencing the soil temperature (ST) and soil nutrition.** Soil microbial community
499 structure and function were strongly related, which suggests that the loss of soil microbial groups
500 may have consequences for overall ecosystem functioning.

501 *Data accessibility.* Requests for data and materials should be addressed to N.H. (henp@igsnr.ac.cn) and G.Y.
502 (yugr@igsnr.ac.cn).

503

504 *Author contributions.* Z.W.X., G.R.Y. and X.Y.Z. planned and designed the research. Z.W.X., N.P.H., R.L.W., and
505 N.Z. conducted fieldwork. Z.W.X., G.R.Y., X.Y.Z., and Q.F.W wrote the manuscript. All authors contributed
506 critically to the drafts and gave final approval for publication.

507 *Competing interests.* The authors declare that they have no conflict of interest.

508

509 **Acknowledgements**

510 This research was jointly supported by the National Natural Science Foundation of China (41601084, 41571251), the
511 Fundamental Research Funds for the Central Universities (2412019FZ001), Science and Technology Research Project
512 of Jilin Province (JJKH20190283KJ), the National Key R&D Program of China (2016YFA0602301), and the China
513 Postdoctoral Science Foundation (2018M631850).

514 **References**

- 515 van Arendonk, J.J.C.M., Poorter, H. The chemical composition and anatomical structure of leaves of grass species
516 differing in relative growth rate, *Plant Cell Environ.*, 17, 963-970, 1994.
- 517 Bååth, E., Anderson, T.H. Comparison of soil fungal/bacterial ratios in a pH gradient using physiological and PLFA-
518 based techniques, *Soil Biol. Biochem.*, 35, 955-963, 2003.
- 519 Balsler, T.C., Firestone, M.K. Linking microbial community composition and soil processes in a California annual
520 grassland and mixed-conifer forest, *Biogeochemistry*, 73, 395-415, 2005.
- 521 Balsler, T.C., Wixon, D.L. Investigating biological control over soil carbon temperature sensitivity, *Global Change*
522 *Biol.*, 15, 2935-2949, 2009.
- 523 Banerjee, S., Kirkby, C.A., Schmutter, D., Bissett, A., Kirkegaard, J.A., Richardson, A.E. Network analysis reveals
524 functional redundancy and keystone taxa amongst bacterial and fungal communities during organic matter
525 decomposition in an arable soil, *Soil Biol. Biochem.*, 97, 188-198, 2016.
- 526 Boberg, J.B., Finlay, R.D., Stenlid, J., Ekblad, A., Lindahl, B.D. Nitrogen and carbon reallocation in fungal mycelia
527 during decomposition of boreal forest litter, *PLOS ONE*, 9, e92897, 2014.
- 528 Brockett, B.F.T., Prescott, C.E., Grayston, S.J. Soil moisture is the major factor influencing microbial community
529 structure and enzyme activities across seven biogeoclimatic zones in western Canada, *Soil Biol. Biochem.*, 44,
530 9-20, 2012.
- 531 Buscot, F. Implication of evolution and diversity in arbuscular and ectomycorrhizal symbioses, *J. Plant Physiol.*, 172,
532 55-61, 2015.
- 533 Cao, H., Chen, R., Wang, L., Jiang, L., Yang, F., Zheng, S., Wang, G., Lin, X. Soil pH, total phosphorus, climate and
534 distance are the major factors influencing microbial activity at a regional spatial scale, *Sci. Rep-UK*, 6, 25815,
535 2016.
- 536 Carson, J.K., Gonzalez-Quinones, V., Murphy, D.V., Hinz, C., Shaw, J.A., Gleeson, D.B. Low pore connectivity
537 increases bacterial diversity in soil, *Appl. Environ. Microb.*, 76, 3936-3942, 2010.
- 538 Coulis, M., Fromin, N., David, J.-F., Gavinet, J., Clet, A., Devidal, S., Roy, J., Hättenschwiler, S. Functional
539 dissimilarity across trophic levels as a driver of soil processes in a Mediterranean decomposer system exposed
540 to two moisture levels, *Oikos*, 124, 1304-1316, 2015..
- 541 De Vries, F.T., Manning, P., Tallwin, J.R., Mortimer, S.R., Pilgrim, E.S., Harrison, K.A., Hobbs, P.J., Quirk, H.,
542 Shipley, B., Cornelissen, J.H., Kattge, J., Bardgett, R.D. Abiotic drivers and plant traits explain landscape-scale
543 patterns in soil microbial communities, *Ecol. Lett.*, 15, 1230-1239, 2012.
- 544 Deforest, J., Smemo, K., Burke, D., Elliott, H., Becker, J. Soil microbial responses to elevated phosphorus and pH in
545 acidic temperate deciduous forests, *Biogeochemistry*, 109, 189-202, 2012.
- 546 Eichenberg, D., Trogisch, S., Huang, Y., He, J.S., Bruelheide, H. Shifts in community leaf functional traits are related
547 to litter decomposition along a secondary forest succession series in subtropical China. *Journal of Plant Ecology*,
548 2014. <http://dx.doi.org/10.1093/jpe/rtu021>.
- 549 Fang, H., Cheng, S., Wang, Y., Yu, G., Xu, M., Dang, X., Li, L., Wang, L. Changes in soil heterotrophic respiration,
550 carbon availability, and microbial function in seven forests along a climate gradient, *Ecol. Res.*, 29, 1077-1086,
551 2014.
- 552 Fiere, N., Bradford, M. A. Toward an ecological classification of soil bacteria, *Ecology*, 88, 1354-1364, 2007.
- 553 Garland, J.L., Mills, A.L. Classification and characterization of heterotrophic microbial communities on the basis of
554 patterns of community-level sole carbon source utilization, *Appl. Environ. Microb.*, 57, 2351-2359, 1991.

555 Garnier, E., Cortez, J., Billes, G., Navas, M.L., Roumet, C., Debussche, M., Gérard, L., Alain, B., David, A., Astrid,
556 B., Cathy, N., Jean-Patrick, T. Plant functional markers capture ecosystem properties during secondary
557 succession, *Ecology*, 85, 2630-2637, 2004.

558 Gartzia-Bengoetxea, N., Kandeler, E., Martínez de Arano, I., Arias-González, A. Soil microbial functional activity is
559 governed by a combination of tree species composition and soil properties in temperate forests, *Appl. Soil Ecol.*,
560 100, 57-64, 2016.

561 Gomez, E., Ferreras, L., Toresani, S. Soil bacterial functional diversity as influenced by organic amendment
562 application, *Bioresour. Technol.*, 97, 1484-1489, 2006.

563 Guckert, J.B., Carr, G.J., Johnson, T.D., et al. Community analysis by Biolog: curve integration for statistical analysis
564 of activated sludge microbial habitats, *J. Microbiol. Meth.*, 27, 183-197, 1996.

565 Huang, Z., Clinton, P.W., Baisden, W.T., Davis, M.R. Long-term nitrogen additions increased surface soil carbon
566 concentration in a forest plantation despite elevated decomposition, *Soil Biol. Biochem.*, 43, 302-307, 2011.

567 Jagadamma, S., Mayes, M.A., Steinweg, J.M., Schaeffer, S.M. Substrate quality alters the microbial mineralization
568 of added substrate and soil organic carbon, *Biogeosciences*, 11, 4665-4678, 2014.

569 Jones, D., Willett, V. Experimental evaluation of methods to quantify dissolved organic nitrogen (DON) and dissolved
570 organic carbon (DOC) in soil, *Soil Biol. Biochem.*, 38, 991-999, 2006.

571 Keiblinger, K.M., Hall, E.K., Wanek, W., Szukics, U., Hammerle, I., Ellersdorfer, G., Bock, S., Strauss, J., Sterflinger,
572 K., Richter, A., Zechmeister-Boltenstern, S. The effect of resource quantity and resource stoichiometry on
573 microbial carbon-use-efficiency. *FEMS Microb. Ecol.* 73, 430-440, 2010.

574 Keller, J. K. & Takagi, K. K. Solid-phase organic matter reduction regulates anaerobic decomposition in bog soil.
575 *Ecosphere*, 4(5), 1-12, 2013.

576 Kivlin, S.N. & Treseder, K.K. Soil extracellular enzyme activities correspond with abiotic factors more than fungal
577 community composition, *Biogeochemistry*, 117, 23-37, 2014.

578 Klimek, B., Chodak, M., Jaźwa, M., Niklińska, M. Functional diversity of soil microbial communities in boreal and
579 temperate Scots pine forests, *Eur. J. Forest Res.*, 135, 731-742, 2016.

580 Kunito, T., Akagi, Y., Park, H.-D., Toda, H. Influences of nitrogen and phosphorus addition on polyphenol oxidase
581 activity in a forested Andisol, *Eur. J. Forest Res.*, 128, 361-366, 2009.

582 Laughlin, D.C. Nitrification is linked to dominant leaf traits rather than functional diversity. *Journal of Ecology* 99,
583 1091-1099, 2011.

584 Lipson, D.A., Monson, R.K., Schmidt, S.K., Weintraub, M.N. The trade-off between growth rate and yield in
585 microbial communities and the consequences for under-snow soil respiration in a high elevation coniferous forest,
586 *Biogeochemistry*, 95, 23-35, 2009.

587 Liu, L., Gundersen, P., Zhang, T., and Mo, J. M.: Effects of phosphorus addition on soil microbial biomass and
588 community composition in three forest types in tropical China, *Soil Biol. Biochem.*, 44, 31-38, 2012a.

589 Liu, X., Liang, M., Etienne, R.S., Wang, Y., Staehelin, C., Yu, S. Experimental evidence for a phylogenetic Janzen-
590 Connell effect in a subtropical forest, *Ecol. Lett.*, 15, 111-118, 2012b.

591 Liu, Z., Liu, G., Fu, B., Wu, Y., Hu, H., Fu, S. Changes in the soil microbial community with a pine plantation
592 restoration in a dry valley of the upper reaches of the Minjiang River, southwest China, *Ann N Y Acad. Sci.*,
593 1195 Suppl 1, E82-95, 2010.

594 Megonigal, J. P., Hines, M. E., Visscher, P. T. *Anaerobic metabolism: linkages to trace gases and aerobic processes.*
595 *Pages 317-424 in W. H. Schlesinger, editor. Biogeochemistry. Elsevier-Pergamon, Oxford, UK, 2004.*

596 Menyailo, O.V., Hungate, B.A., Zech, W. Tree species mediated soil chemical changes in a Siberian artificial
597 Afforestation experiment. *Plant Soil*, 242, 171-182, 2002.

598 Millard, P., Sommerkorn, M., Grelet, G.A. Environmental change and carbon limitation in trees: a biochemical,
599 ecophysiological and ecosystem appraisal, *New Phytol.*, 175, 11-28, 2007.

600 Monson, R.K., Lipson, D.L., Burns, S.P., Turnipseed, A.A., Delany, A.C., Williams, M.W., Schmidt, S.K. Winter
601 forest soil respiration controlled by climate and microbial community composition, *Nature*, 439, 711-714, 2006.

602 Myers, R.T., Zak, D.R., White, D.C., Peacock, A. Landscape-level patterns of microbial community composition and
603 substrate use in upland forest ecosystems, *Soil Sci. Soc. Am. J.*, 65, 359-367, 2001.

604 Orwin, K.H., Buckland, S.M., Johnson, D., Turner, B.L., Smart, S., Oakley, S., Bardgett, R.D. Linkages of plant traits
605 to soil properties and the functioning of temperate grassland, *J. Ecol.*, 98, 1074-1083, 2010.

606 Pei, Z., Eichenberg, D., Bruelheide, H., Kröber, W., Kühn, P., Li, Y., von Oheimb, G., Purschke, O., Scholten, T.,
607 Buscot, F., Gutknecht, J.L.M. Soil and tree species traits both shape soil microbial communities during early
608 growth of Chinese subtropical forests, *Soil Biol. Biochem.*, 96, 180-190, 2016.

609 Peng, X., Wang, W. Stoichiometry of soil extracellular enzyme activity along a climatic transect in temperate
610 grasslands of northern China, *Soil Biol. Biochem.*, 98, 74-84, 2016.

611 Philippot, L., Spor, A., Henault, C., Bru, D., Bizouard, F., Jones, C.M., Sarr, A., Maron, P.A. Loss in microbial
612 diversity affects nitrogen cycling in soil, *ISME J.*, 7, 1609-1619, 2013.

613 Poorter, H., Bergkotte, M. Chemical composition of 24 wild species differing in relative growth rate, *Plant Cell*
614 *Environ.*, 15, 221-229, 1992.

615 Preston-Mafham, J., Boddy, L., Randerson, P.F. Analysis of microbial community functional diversity using sole-
616 carbon-source utilisation profiles-a critique, *FEMS Microb. Ecol.*, 42, 1-14, 2002.

617 Priha, O., Grayston, S.J., Hiukka, R., Pennanen, T., Smolander, A. Microbial community structure and characteristics
618 of the organic matter in soils under *Pinus sylvestris*, *Picea abies* and *Betula pendula* at two forest sites. *Biol*
619 *Fertil Soils*, 33, 17-24, 2001.

620 Raich, J.W., Clark, D.A., Schwendenmann, L., Wood, T.E. Aboveground tree growth varies with belowground carbon
621 allocation in a tropical rainforest environment, *PLoS one*, 9, e100275, 2014.

- 622 Riah-Anglet, W., Trinsoutrot-Gattin, I., Martin-Laurent, F., Laroche-Ajzenberg, E., Norini, M.-P., Latour, X., Laval,
623 K. Soil microbial community structure and function relationships: A heat stress experiment, *Appl. Soil Ecol.*, 86,
624 121-130, 2015..
- 625 Russell, A.E., Raich, J.W., Valverde-Barrantes, O.J., Fisher, R.F. Tree species effects on soil properties in experimental
626 plantations in tropical moist forest, *Soil Sci. Soc. Am. J.*, 71, 1389, 2007.
- 627 Saiya-Cork, K.R., Sinsabaugh, R.L., Zakb, D.R. The effects of long term nitrogen deposition on extracellular enzyme
628 activity in an *Acer saccharum* forest soil, *Soil Biol. Biochem.*, 34, 1309-1315, 2002.
- 629 Sessitsch, A., Weilharter, A., Gerzabek, M.H., Kirchmann, H., Kandeler, E. Microbial Population Structures in Soil
630 Particle Size Fractions of a Long-Term Fertilizer Field Experiment, *Appl. Environ. Microb.*, 67, 4215-4224, 2001.
- 631 Sherman, C., Steinberger, Y. Microbial functional diversity associated with plant litter decomposition along a climatic
632 gradient, *Microb. Ecol.*, 64, 399-415, 2012.
- 633 Six, J., Elliott, E. T., Paustian, K. Soil structure and soil organic matter: II. A normalized stability index and the effect
634 of mineralogy, *Soil Sci. Soc. Am. J.*, 64, 1042-1049, 2000.
- 635 Sterner, R. W. & Elser, J. J. Ecological stoichiometry: the biology of elements from molecules to the biosphere.
636 Princeton, New Jersey, USA: Princeton University Press, 2002.
- 637 Strickland, M.S., Lauber, C., Fierer, N., Bradford, M.A. Testing the functional significance of microbial community
638 composition, *Ecology*, 90, 441-451, 2009.
- 639 Soil Survey Staff Keys to Soil Taxonomy, 11th ed. USDA-Natural Resources Conservation Service, Washington, DC,
640 2010.
- 641 Swallow, M.J.B., Quideau, S.A. A method for determining community level physiological profiles of organic soil
642 horizons, *Soil Sci. Soc. Am. J.*, 79, 536-542, 2015.
- 643 Talbot, J.M., Bruns, T.D., Taylor, J.W., Smith, D.P., Branco, S., Glassman, S.I., Erlandson, S., Vilgalys, R., Liao, H.L.,
644 Smith, M.E., Peay, K.G. Endemism and functional convergence across the North American soil mycobiome, *P.*
645 *Natl. Acad. Sci. USA.*, 111, 6341-6346, 2014.
- 646 Ter Braak, C.J.F., Smilauer, P. CANOCO Reference manual and CanoDraw for Windows User's guide: Software for
647 Canonical Community Ordination (Version 4.5). Microcomputer, 2002.
- 648 Tian, H., Chen, G., Zhang, C., Melillo, J.M., Hall, C.A.S.. Pattern and variation of C:N:P ratios in China's soils: a
649 synthesis of observational data, *Biogeochemistry*, 98, 139-151 2010.
- 650 Tian, J., McCormack, L., Wang, J., Guo, D., Wang, Q., Zhang, X., Yu, G., Blagodatskaya, E., Kuzyakov, Y. Linkages
651 between the soil organic matter fractions and the microbial metabolic functional diversity within a broad-leaved
652 Korean pine forest, *Eur. J. Soil Biol.*, 66, 57-64, 2015.
- 653 Treonis, A.M., Ostle, N.J., Stott, A.W., Primrose, R., Grayston, S.J., Ineson, P. Identification of groups of
654 metabolically-active rhizosphere microorganisms by stable isotope probing of PLFAs, *Soil Biol. Biochem.*, 36,
655 533-537, 2004.
- 656 Tu, Q., Deng, Y., Yan, Q., Shen, L., Lin, L., He, Z., Wu, L., Van Nostrand, J.D., Buzzard, V., Michaletz, S.T., Enquist,
657 B.J., Weiser, M.D., Kaspari, M., Waide, R.B., Brown, J.H., Zhou, J. Biogeographic patterns of soil diazotrophic
658 communities across six forests in the North America, *Mol. Ecol.*, 25, 2937-2948, 2016.
- 659 Valášková, V., Šnajdr, J., Bittner, B., Cajthaml, T., Merhautová, V., Hofrichter, M., Baldrian, P. Production of
660 lignocellulose-degrading enzymes and degradation of leaf litter by saprotrophic basidiomycetes isolated from a
661 *Quercus petraea* forest, *Soil Biol. Biochem.*, 39, 2651-2660, 2007.
- 662 Vance, E.D., Brookes, P.C., Jenkinson, D.S. An extraction method for measuring soil microbial biomass C, *Soil Biol.*
663 *Biochem.*, 19, 703-707, 1987.
- 664 Veen, J.A.V., Kuikman, P.J. Soil structural aspects of decomposition of organic-matter by micro-organism,
665 *Biogeochemistry*, 11, 213-233, 1990.
- 666 Waldrop, M.P., Firestone, M.K. Response of microbial community composition and function to soil climate change,
667 *Microb. Ecol.*, 52, 716-724, 2006.
- 668 Waldrop, M. P., Balsler, T. C. & Firestone, M. K. Linking microbial community composition to function in a tropical
669 soil, *Soil Biol. Biochem.*, 32, 1837-1846, 2000.
- 670 Wang, Q., He, N., Yu, G., Gao, Y., Wen, X., Wang, R., Koerner, S.E., Yu, Q. Soil microbial respiration rate and
671 temperature sensitivity along a north-south forest transect in eastern China: Patterns and influencing factors, *J.*
672 *Geophys. Res.(Biogeosciences)*, 121, 399-410, 2016.
- 673 Wang, R., Yu, G., He, N., Wang, Q., Zhao, N., & Xu, Z. Latitudinal variation of leaf morphological traits from species
674 to communities along a forest transect in eastern China. *Journal of Geographical Sciences*, 26(1), 15-26, 2016.
- 675 Waring, B.G., Alvarez-Cansino, L., Barry, K.E., Becklund, K.K., Dale, S., Gei, M.G., Keller, A.B., Lopez, O.R.,
676 Markesteijn, L., Mangan, S., Riggs, C.E., Rodriguez-Ronderos, M.E., Segnitz, R.M., Schnitzer, S.A., Powers,
677 J.S. Pervasive and strong effects of plants on soil chemistry: a meta-analysis of individual plant 'Zinke' effects,
678 *Proc. R. Soc. B. The Royal Society*, 282, 20151001, 2015.
- 679 Wei, H., Guenet, B., Vicca, S., Nunan, N., AbdElgawad, H., Pouteau, V., Shen, W., Janssens, I.A. Thermal acclimation
680 of organic matter decomposition in an artificial forest soil is related to shifts in microbial community structure,
681 *Soil Biol. Biochem.*, 71, 1-12, 2014.
- 682 Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., et al. The worldwide leaf economics
683 spectrum, *Nature*, 428, 821-827, 2004.
- 684 Xu, X., Thornton, P.E., Post, W.M. A global analysis of soil microbial biomass carbon, nitrogen and phosphorus in
685 terrestrial ecosystems, *Global Ecol. Biogeogr.*, 22, 737-749, 2013.
- 686 Xu, Z., Yu, G., Zhang, X., Ge, J., He, N., Wang, Q., Wang, D. The variations in soil microbial communities, enzyme
687 activities and their relationships with soil organic matter decomposition along the northern slope of Changbai
688 Mountain, *Appl. Soil Ecol.*, 86, 19-29, 2015.

689 Xu, Z., Yu, G., Zhang, X., He, N., Wang, Q., Wang, S., Wang, R., Zhao, N., Jia, Y., Wang, C. Soil enzyme activity and
690 stoichiometry in forest ecosystems along the North-South Transect in eastern China (NSTEC), *Soil Biol.*
691 *Biochem.*, 104, 152-163, , 2017.

692 Xu, Z., Yu, G., Zhang, X., He, N., Wang, Q., Wang, S., Xu, X., Wang, R., Zhao, N. Biogeographical patterns of soil
693 microbial community as influenced by soil characteristics and climate across Chinese forest biomes, *Appl. Soil*
694 *Ecol.*, 124, 298-305, 2018.

695 Yu, G., Zheng, Z., Wang, Q., Fu, Y., Zhuang, J., Sun, X., Wang, Y. Spatiotemporal pattern of soil respiration of
696 terrestrial ecosystems in China: the development of a geostatistical model and its simulation., *Environ. Sci.*
697 *Technol.*, 44, 6074-6080, 2010.

698 Zak, J.C., Willig, M.R., Moorhead, D.L., Wildman, H.G. Functional diversity of microbial communities: a quantitative
699 approach, *Soil Biol. Biochem.*, 26, 1101-1108, 1994.

700 Zhang, X. S., Yang, D. A. Allocation and study on global change transects in China, *Quaternary Sci.*, 1, 43-52, 1995.
701 (In Chinese)

702 Zhou, J., Deng, Y., Shen, L., Wen, C., Yan, Q., Ning, D., Qin, Y., Xue, K., Wu, L., He, Z., Voordeckers, J.W., Nostrand,
703 J.D., Buzzard, V., Michaletz, S.T., Enquist, B.J., Weiser, M.D., Kaspari, M., Waide, R., Yang, Y., Brown, J.H.
704 Temperature mediates continental-scale diversity of microbes in forest soils, *Nat. Commun.*, 7, 12083, 2016.

705 **Figures legends**

706 **Fig. 1.** Distribution of typical forest ecosystems along the North-South Transect of eastern China (NSTEC). The
707 abbreviations for the sampling sites from north to south are as follows: HZ, Huzhong; LS, Liangshui; CB, Changbai;
708 DL, Dongling; TY, Taiyue; SN, Shennong; JL, Jiulian; DH, Dinghu; JF, Jiangfeng. These abbreviations are used for
709 the nine forests throughout.

710 **Fig.2.** Variations in soil microbial substrate use during a 240-h incubation for the nine forests. Different colors
711 represent different forest types: Yellow, coniferous forest; Dark yellow, coniferous broad-leaved mixed forest; Purple,
712 deciduous broad-leaved forest; Olive, subtropical evergreen broad-leaved forest; Orange, Tropical monsoon forest.
713 Different lowercase letters indicate significant differences among forests in the same climate zone. The abbreviations
714 of the sampling sites are given in Table 1.

715 **Fig. 3.** Characteristics of microbial use of (a) carbohydrates, (b) carboxylic acids, (c) amino acids, (d) polymers, (e)
716 amines, and (f) miscellaneous along the NSTEC. The representatives of different colors were showed in Figure 2.

717 **Fig.4.** Redundancy analysis (RDA) ordination biplot of soil microbial carbon sources use efficiency and
718 environmental properties. The representatives of different colors were showed in Figure 2. The dotted lines and solid
719 lines represent the environmental variables and lipid signatures and carbon sources. The abbreviations of the variables
720 in this figure are as follows: MAP, mean annual precipitation. The vegetation data: LDMC, leaf dry matter weight;
721 Leaf C, leaf carbon content; Leaf N, leaf nitrogen content; SLA, specific leaf area. Soil properties included SMC, soil
722 moisture content; Silt, soil silt content; TN, soil total nitrogen; SOC, soil organic carbon. The abbreviations of the
723 sampling sites were given in Table 1

724 **Fig.5.** The heatmap of the Pearson's correlation coefficients between the use of individual substrates and microbial
725 PLFAs and soil enzyme activities. Note: The abbreviations of the variables: Actino-, actinomycetes; F/B,
726 fungi/bacteria; G⁺, gram positive bacteria; G⁻, gram negative bacteria; G⁺/G⁻, Gram-positive bacteria/ Gram-negative
727 bacteria. BG, β -1, 4-glucosidase; NAG, β -1,4-N-acetylglucosaminidase; LAP, leucine aminopeptidase; AP, acid
728 phosphatase. ** $P < 0.01$, * $P < 0.05$.

729 **Fig. 6.** Relationships between soil carbon mineralization rates ($\mu\text{g C g}^{-1}\text{d}^{-1}$) and microbial biomass C (MBC), soil
730 dissolved organic C (DOC), average well color development (AWCD), and individual substrate use.

731 **Fig. 7.** Relationships between soil carbon mineralization rates ($\mu\text{g C g}^{-1}\text{d}^{-1}$) and different groups of soil microbial
732 PLFAs (a-h) and enzyme activities (i-l).

733
734

735 **Supporting information**

736 **Table S1** Average values of forest soil enzyme activities and different PLFA groups along the NSTEC.

737 **Table S2** Plant diversity and community weighted means of plant functional traits

738 **Table S3** Soil organic matter (SOM) decomposition rates during the 28 days of incubation time (Mean \pm SE) ($\mu\text{g C}$
739 $\text{g}^{-1}\text{d}^{-1}$)

740 **Fig. S1** Variations in the average well color development (AWCD) values during a 240-h incubation for the nine
741 forests. The abbreviations of the sampling sites are the same as those in Table 1.

742
743

Tables

Table 1. The main characteristics of the sampling sites along the North South Transect of East China

| Sampling Sites | Longitude (E) | Latitude (N) | Elevation (m) | MAT ^b (°C) | MAP ^b (mm) | Vegetation types | Soil type |
|-----------------|---------------|--------------|---------------|-----------------------|-----------------------|---|---------------------|
| HZ ^a | 123°01'12" | 51°46'48" | 850 | -3.7 | 473 | Cold temperate coniferous forest | Spodosols |
| LS | 128°53'51" | 47°11'06" | 401 | 0.01 | 648 | Temperate conifer broad-leaved mixed forest | Albi-Boric Argosols |
| CB | 128°05'27" | 42°24'16" | 758 | 2.8 | 691 | Temperate conifer broad-leaved mixed forest | Albi-Boric Argosols |
| DL | 115°25'24" | 39°57'27" | 972 | 6.6 | 539 | Warm temperate deciduous broad-leaved forest | Alfisols |
| TY | 112°04'39" | 36°41'43" | 1668 | 6.0 | 644 | Warm temperate deciduous broad-leaved forest | Alfisols |
| SN | 110°29'43" | 31°19'15" | 1510 | 8.5 | 1447 | Subtropical deciduous evergreen mixed forest | Inceptisols |
| JL | 114°26'28" | 24°35'05" | 562 | 18.2 | 1770 | Subtropical evergreen broad-leaved forest | Ultisols |
| DH | 112°32'14" | 23°10'25" | 240 | 21.8 | 1927 | Subtropical monsoon evergreen broad-leaved forest | Ultisols |
| JF | 108°51'26" | 18°44'18" | 809 | 23.2 | 2266 | Tropical monsoon forest | Ultisols |

744 a:HZ, Huzhong; LS, Liangshui; CB, Changbai; DL, Dongling; TY,Taiyue; SN, Shennong; JL, Jiulian; DH, Dinghu; JF, Jiangfeng.

745 b: MAT, mean annual temperature; MAP, mean annual precipitation.

746

Table 2. Soil properties of different sampling sites

| Sampling site | pH | ST (°C) | SMC (%) | Silt (%) | SOC (g kg ⁻¹) | MBC (mg kg ⁻¹) | DOC (mg kg ⁻¹) | TN (g kg ⁻¹) | TP (g kg ⁻¹) |
|---------------|------------|------------|-------------|-------------|------------------------------|-------------------------------|-------------------------------|-----------------------------|-----------------------------|
| HZ | 6.79±0.02a | 10.3±0.15g | 45.3±0.90c | 56±1.2c | 42.29±0.47b | 350±6.0a | 240±7.6e | 2.90±0.16d | 0.87±0.02b |
| LS | 6.17±0.02b | 15.9±0.02f | 46.9±0.76c | 64±0.3b | 62.08±7.20a | 316±0.7a | 204±4.9f | 4.59±0.29b | 0.59±0.02c |
| CB | 6.37±0.04b | 16.0±0.06f | 102.8±0.25a | 76±0.6a | 72.38±2.00a | 178±8.8b | 314±8.6c | 6.05±0.17a | 1.67±0.08a |
| DL | 6.87±0.02a | 17.8±0.14e | 32.4±0.30e | 6±2.4e | 38.83±0.41c | 43±0.8e | 284±2.6d | 3.17±0.04d | 0.56±0.01c |
| TY | 6.85±0.05a | 16.0±0.12f | 36.0±0.23d | 49±1.4d | 41.34±2.75c | 115±4.0c | 226±13.8f | 2.43±0.15e | 0.52±0.01c |
| SN | 6.93±0.01a | 18.4±0.12d | 50.5±0.63b | 74±0.3a | 36.13±1.26c | 72±13.1e | 311±13.2c | 3.76±0.05c | 0.81±0.01b |
| JL | 5.57±0.19b | 25.3±0.01a | 39.0±0.89d | 68±0.3b | 31.55±1.82c | 89±19.7d | 387±1.9b | 2.28±0.09e | 0.36±0.01d |
| DH | 5.43±0.03c | 24.4±0.04b | 37.8±0.38d | 50±1.8d | 28.47±0.54d | 38±0.1e | 334±7.7c | 1.77±0.02f | 0.20±0.01e |
| JF | 6.32±0.01c | 22.5±0.07c | 38.6±0.12d | 49±0.2d | 29.38±0.94d | 140±1.3c | 458±6.6a | 1.99±0.02e | 0.15±0.01e |

747

Note: ST=temperature of 0–10 cm soil; SMC=soil moisture content; Silt=soil silt content; SOC=soil organic carbon; MBC=microbial biomass carbon; DOC=dissolved organic carbon; TN=soil total nitrogen; TP=soil total phosphorus. Values were presented as means ± SE (n=4). The abbreviations of the sampling sites were given in the Table 1.

748

749

Table 3. Functional diversity of soil microbial communities in forest ecosystems along the NSTEC


| Sampling sites | Richness (R) | Shannon <i>H'</i> | Shannon <i>E</i> | Simpson <i>D</i> |
|-----------------------|---------------------|--------------------------|-------------------------|-------------------------|
| HZ | 14.08±0.34d | 2.65±0.03d | 1.01±0.007b | 0.91±0.002c |
| LS | 25.29±0.14b | 3.12±0.02b | 0.98±0.003c | 0.95±0.001a |
| CB | 27.00±0.27a | 3.22±0.01a | 0.98±0.001c | 0.95±0.001a |
| DL | 11.54±0.47e | 2.52±0.03e | 1.04±0.010a | 0.87±0.005d |
| TY | 22.33±0.87c | 3.02±0.02c | 0.98±0.002c | 0.94±0.001a |
| SN | 28.10±0.34a | 3.24±0.01a | 0.97±0.001c | 0.95±0.001a |
| JL | 23.54±0.07c | 3.04±0.01c | 0.96±0.001c | 0.93±0.003b |
| DH | 25.65±0.71b | 3.11±0.01b | 0.97±0.001c | 0.93±0.002b |
| JF | 27.63±0.68a | 3.19±0.02a | 0.96±0.001c | 0.95±0.002a |

750


Indices were calculated based on the optical density values after incubation for 96 h. Data are expressed as means±standard errors. Different lowercase letters indicate significant differences among forests. The abbreviations of the sampling sites are the same as those used in Table 1.

751

752




754 **Figure 1.** Distribution of typical forest ecosystems along the North-South Transect of eastern China (NSTEC).
 755 The abbreviations of sampling sites from north to south are as follows: HZ, Huzhong; LS, Liangshui; CB,
 756 Changbai; DL, Dongling; TY, Taiyue; SN, Shennong; JL, Jiulian; DH, Dinghu; JF, Jiangfeng.
 757




758
759
760
761
762
763

Figure 2. Variations in soil total microbial metabolic intensity during a 240-h incubation for the nine forests. Different colors represent different forest types: Yellow, coniferous forest; Dark yellow, coniferous broad-leaved mixed forest; Purple, deciduous broad-leaved forest; Olive, subtropical evergreen broad-leaved forest; Orange, Tropical monsoon forest. Different lowercase letters indicate significant differences among forests in the same climate zone. The abbreviations of the sampling sites are given in Table 1.



764
765
766
767

Figure 3. Characteristics of microbial use of (a) carbohydrates, (b) carboxylic acids, (c) amino acids, (d) polymers, (e) amines, and (f) miscellaneous along the NSTEC. The representatives of different colors were showed in Figure 2.



768
769
770
771
772
773
774
775

Figure 4. Redundancy analysis (RDA) ordination biplot of soil microbial carbon sources use efficiency and environmental properties. The representatives of different colors were showed in Figure 2. The dotted lines and solid lines represent the environmental variables and lipid signatures and carbon sources. The abbreviations of the variables in this figure are as follows: MAP, mean annual precipitation. The vegetation data: LDMC, leaf dry matter weight; Leaf C, leaf carbon content; Leaf N, leaf nitrogen content; SLA, specific leaf area. Soil properties included SMC, soil moisture content; Silt, soil silt content; TN, soil total nitrogen; SOC, soil organic carbon. The abbreviations of the sampling sites were given in Table 1.




Figure 5. The heatmap of the Pearson's correlation coefficients between the use of individual substrates and microbial PLFAs and soil enzyme activities. Note: The abbreviations of the variables: Actino-, actinomycetes; F/B, fungi/bacteria; G⁺, gram positive bacteria; G⁻, gram negative bacteria; G⁺/G⁻, Gram-positive bacteria/ Gram-negative bacteria. BG, β-1, 4-glucosidase; NAG, β-1,4-N-acetylglucosaminidase; LAP, leucine aminopeptidase; AP, acid phosphatase. ***P* < 0.01, **P* < 0.05.




Figure 6. Relationships between soil carbon mineralization rates ($\mu\text{g C g}^{-1} \text{d}^{-1}$) and microbial biomass C (MBC), soil dissolved organic C (DOC), average well color development (AWCD), and use of individual substrates.




Figure 7. Relationships between soil carbon mineralization rates ($\mu\text{g C g}^{-1} \text{d}^{-1}$) and different groups of soil microbial PLFAs (a-h) and enzyme activities (i-l).