

Author's response

Again, we thank all reviewers for their careful reviews. We acknowledge the issues brought up and, in the following, we reply to all of them point-by-point.

Reviewers comment

Our reply

Anonymous Referee #1

The work of Amann and Hartmann reviews the use of carbon negative technologies, particularly Enhanced Weathering (EW). The main view point expressed by the authors is the necessity for a “co-deployment” of several technologies, because no single approach will be able to reach the 1.5C target set by the IPCC. The manuscript is timely, and I support the overall goal.

However, the manuscript is largely qualitative rather than quantitative, and seems to focus on soil and EW, whereas the title, and partially the abstract, implied a much broader review. In my opinion this work can be published in BG, but some amendments are necessary.

This point is well taken. The synergies mainly apply to soil-based NETs, maybe we can reflect this in the title. However, “soil-based NET” is not a term established as such in the community. About the qualitiveness: This is a good point. There is some data out there on the processes mentioned, however the framework of research is very often so far off the focus of our manuscript, that it would introduce a level of detail that is misleading for the purpose of this manuscript. We explicitly chose the format of a “perspectives piece” to identify the main important processes that need to be considered in future research on NETs and their combined effects. This should provide a guideline for projects to come. Yet, we try to be at least a little more specific in the discussion of the processes.

Specific comments

- Figure 1 is not sufficiently clear. I would place the benefactor on top of the beneficiaries. Also I was not sure how to read the figure. For example, in the first block EW enhances the nutrient pool and maintain afforestation? I was not sure about the meaning of benefactor and beneficiary here;

We changed the entire figure on different levels. Benefactor/beneficiary were replaced by something more straightforward. The “connecting verbs” were amended with arrow boxes to indicate the reading direction. The CO₂ sequestration effect symbology was changed to avoid misunderstanding it as CO₂ release.

- P.2, line 11. I would appreciate if the authors would insert also a comment on the social and environmental effect of mining;

A short remark was added, to address the potential after-use of mines. The social effect was not address due to the fact that it touches a whole different topic.

- P.3, line 13 introduces the content of enhanced weathering but not all readers are familiar with what this is;

A brief explanation was added.

- P.3, line 26 mention for the first time dunite and basalt. Can the author specify why they choose these specific examples? References are reported but the reader is left wondering what's special about these rocks;

We extended the text to explain the background and we added some broader categories of rock geochemistry (see comment below) to be more general in our arguments.

- Figures 2-3 are interesting, but I am wondering about the overall availability of these resources. As an example, I am not sure about the relevance of komatiite. I understand this is an explicative diagram, but the context here is that of global-deployable technologies. I feel the text should explain better the abundance and distribution of some key resources, or at least provide relevant references/tabula data;

We removed the very specific selection of rocks, which was chosen as available in the database. We generalized the data now, by distinguishing classes via SiO₂ content of volcanic and plutonic ultrabasic/basic/intermediate/acid rocks. This classification enables us to give a broader and more general overview of what to look for in a rock. Additionally, we added the rock types dunite and basalt as commonly discussed types for reference. A map with the global distribution of the distinguished classes is now provided in the supplement (S5). As an example, basic volcanic rocks (this class contains basaltic rocks) covers about 3.5% of the land surface, according to the latest global lithologic map (Hartmann & Moosdorf 2012).

- P.5., line 20 would benefit of a reference. Current models do not consider nutrient availability?;

A reference was added, showing the importance of P cycling, which is commonly not included.

- The paragraph on soil hydrology does not specify the size of the grains that would decrease soil conductivity. I understand this would depend on the soil but can at least a range be specified? It would be important to specify size ranges for these rock flours;

While it isn't possible to be extremely specific, we added a sentence on how clayey, silty, and sandy grainsizes are expected to change hydrology.

- The paragraph on soil biota is extremely qualitative. At least one could point out which minerals are more susceptible to bioweathering, or which structural elements are more needed by soil bacteria.

We added a sentence on how weatherability of certain minerals depends on the type of bioinoculant.

It would also be important to point out the interaction between type of crops and bacteria. In fact I was surprised not to see a paragraph dedicated on how different crops may work synergistically with different type of rocks.

As this is a manuscript with a somewhat broad perspective, we deliberately did not go into details of synergies between crops and bacteria as the focus on NETs would be lost. However, we included the relationship between bacteria and weathering and crops and weathering.

The entire manuscript seem to focus on afforestation, but it would be interesting to point out how agriculture may also benefit from EW;

We do not fully understand this part of the comment. The manuscript approaches the synergies specifically with a focus on products released from weathering, while afforestation is merely addressed as a benefiting NET. How agriculture benefits from EW is specifically addressed in

the sections on nutrient release and retention. If we missed the connection to soil biota here, we kindly ask to be more specific.

- A general comment on nitrogen would benefit the manuscript (e.g., Nitrogen in rock: Occurrences and biogeochemical implications 2002 JoAnn and M. Holloway).

Rock N does play a minor role in agricultural systems. A remark on weathering derived N was added.

Technical comments

- Figure 1 caption can be improved. Land-based should be spelled with a dash and capital letters should be double-checked;

Checked and corrected.

- I may have missed it but I do not think EW was ever defined;

We took care of this added defined EW once in the beginning.

- I would switch the phrasing of the title to: “Synergies from co-deployment of negative emission technologies: Ideas and perspectives”;

We agree and like it better that way, however, the journals rules seem to demand it the other way round: https://www.biogeosciences.net/about/manuscript_types.html: “Manuscripts of this type should be short (a few pages only). The manuscript title must start with “Ideas and perspectives:”.“ We would be happy to change it, if allowed by the editor/journal.

- Latinism such as e.g. and i.e. should be italicized;

Agreed & changed.

- I would change the heading of the last paragraph from “synthesis” to “summary” or “conclusion”;

Agreed & renamed to conclusion.

Anonymous Referee #2

The authors present an overview of how different negative emission technologies might interact and thereby trigger additional carbon uptake. The work is timely and important as such synergies have not been addressed so far in great detail.

However, the authors focus on the beneficial effects of enhanced weathering and biochar on afforestation and BECCS while the title suggests a somewhat broader overview.

In this respect, you and your co-reviewer raise the same concern. We acknowledge this and will adapt the text, and maybe even the title to be more specific about the NETs discussed. Please look into our reply to reviewer #1 for an extended answer.

Additionally, some statements would benefit from describing synergies in a more quantitative way.

Also, here the concern is also raised by reviewer #1. We explicitly chose the format of a “perspectives piece” to identify the main important processes that need to be considered in future research on NETs and their combined effects. This should provide a guideline for projects to come. Yet, we try to be at least a little more specific in the discussion of the processes.’

Specific comments

- Title: As reviewer #1 I would also move “ideas and perspectives” to the end of the title. Maybe “Synergies from co-deployment of land-based negative emission technologies: Ideas and perspectives” to clarify this paper is about soil/land-based strategies.

Agreed, but difficult as described above in the answer to Reviewer #1.

- P2, line 1: “of” is repeated three times. Suggestion: “assessing the effects of combined. . .”.

Agreed and changed.

- P2, line8: “e.g.” implies that there are more nutrients so you can remove “and others”.

Agreed and removed.

- I agree with reviewer #1 that it’s not clear at all how to read Figure 1. Intuitively I would expect benefactors on top of beneficiaries and additional CO2 sequestration. I was also confused by the two verbs connecting benefactors and beneficiaries. Lastly, the additional CO2 sequestration is unclear. I assume the downward arrows mean that e.g. EW increases CO2 sequestration via BECCS or AFF (flux from the atmosphere to the land) but one could also interpret it as a decline.

As you both had similar issues with this figure, we modified it to have it more logically structured and less room for interpretation (also see answer to reviewer 1).

- P3, line 3: envisions.

✓

- P3, line 21: But only if these new forests are harvested.

Added a clause to be more specific.

- P3, line 25: Higher than other rock types?

Yes, this part was amended to complete the comparison.

- P3, line 29: types

✓

- Figure 2: I think it would be interesting to show the CO₂ capture potential of dunite in this figure as it seems to be a highly relevant rock. After all you show Komatiite which also has very low K and P contents.

As mentioned in the caption, dunite contains so little K and P that it wouldn't be easy to visualize. We added it now along with basalt and all the rock classes differentiated by SiO₂ content. For reasons of visibility, the axes had to be spread apart.

- Figure 3: So the extraction range was derived from min/max nutrient contents but what yields were assumed for Miscanthus (range is 40-4400 t/km² according to Table S5.-1)?

The minima of nutrient content and yield were multiplied to get the total minimum nutrient removal. Accordingly, the maximum was calculated. An explanation was added to the caption. Also a description was added next to the areas, to directly indicate their meaning.

In addition, Figure 3 seems to not be referred to in the text.

Good point. It is now.

- P4, line 3: I think this isolated sentence would fit better in the second paragraph of this section.

Yes. We moved it there.

- P5, line 14: nutrients limit tree growth?

Acknowledged. We reformulated the sentences to be precise. Geogenic nutrient supply can be the limiting component in an elevated CO₂ system.

- P6, line 17: Confusing, split this sentence into two sentences.

Agreed and rewritten.

- P6, line 24: Reference missing in the reference list.

Fixed.

- P7, line3: "where considered to be of concern" sounds awkward, I think it can be removed.

Agreed & removed.

- P7, line 10: Reference seems to be at the wrong position.

Agreed & corrected.

- Table 1: The reader is left wondering what values are typical. Can you also provide numbers for some other rock types for comparison?

We generalized the data now, by distinguishing classes via SiO₂ content of volcanic and plutonic ultrabasic/basic/intermediate/acid rocks. This classification enables us to give a broader and more general overview of what to look for in a rock. Additionally, we add the rock types dunite and basalt as commonly discussed types for reference. Due to the increased amount of data, we converted the table into a xy-plot, using median and 10/90 percentile data. The numerical data is provided in the supplement (S7).

In the following, we provide a detailed overview over all changes that have been done compared to the original document. It is document comparison generated by Microsoft Word.

Ideas and perspectives: Synergies from co-deployment of negative emission technologies

Thorben Amann¹ and Jens Hartmann¹

¹Institute for Geology, Center for Earth System Research and Sustainability, Universität Hamburg, Germany

5 Correspondence to: Thorben Amann (science@thorbenamann.de)

Abstract. Numerous publications propose the deployment of negative emission technologies, which intend to actively remove CO₂ from the atmosphere with the goal to reach the 1.5° target as discussed by the IPCC. The increasing amount of scientific studies on the individual potential of different envisaged technologies and methods indicates, that no single method has enough capacities to mitigate the issue by itself. It is thus expected that technology portfolios are deployed. As some of them utilize the same environmental compartment, co-deployment effects are expected. Those effects are particularly important to evaluate with respect to additional CO₂ uptake. Considering soils as one of the main affected compartments, we see a plethora of processes which can positively benefit from each other, cancelling out negative side effects or increasing overall CO₂ sequestration potentials. To derive more reliable estimates of negative emission potentials and to evaluate common effects on global carbon pools, it is now necessary to intensively study interrelated effects of negative emission technology deployment while minimizing side effects.

Introduction

As global mean temperatures are projected to increase further, strategies to mitigate climate change in time by decreasing CO₂ emissions seem to slowly take effect (Jackson et al., 2015). Some CO₂ emission pathways include negative carbon emission strategies (Fuss et al., 2014; Fuss et al., 2016; Rogelj et al., 2018), that essentially capture CO₂ from the atmosphere in different ways, storing them, in the long term, as CO₂ molecules, or as organic and inorganic compounds (Caldeira et al., 2013). All discussed options and technologies have yet to reach the large-scale deployment stage (Minx et al., 2018; Nemet et al., 2018). Most technologies are immature, lacking deep research on the global potential, technical feasibility, economics of deployment, and especially an assessment of the expected side effects (National Research Council, 2015; Fuss et al., 2018).

The proposed negative emission technologies (NET) encompass highly technical engineering solutions as well as methods that rely on natural processes, like growth of biomass (e.g., bioenergy with carbon capture and storage (BECCS), and afforestation), soil carbon increase, biochar, and chemical weathering (e.g., Enhanced Weathering (EW) and ocean liming). As these methods are aimed to be integrated in the global biogeochemical cycle and will redistribute carbon between reservoirs (Keller et al., 2018), their interaction is inevitable if NETs are deployed at the largest scale. As such, it must be assessed how the co-deployment of NETs will affect the individual and overall efficiency, since until now, publications

Formatvorlagendefinition ... [1]

hat formatiert: Englisch (Vereinigtes Königreich)

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: cancelling

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: CO₂ sequestration potentials

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: on a

hat formatiert ... [2]

hat gelöscht: .

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: ,

hat formatiert ... [3]

hat gelöscht: . enhanced weathering

hat formatiert ... [4]

hat gelöscht: cycles

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: within

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: ¶

focus generally on single NETs, disregarding any effects on concurrent deployment of additional technologies.

Findings from NET specific literature suggest that assessing the effects of combined NET rollout is advisable and future research should include CO₂ sequestration enhancing side effects that could increase the overall potential of NETs. However, the principal interaction between proposed methods needs to be studied in detail beforehand, to understand effects on the carbon pools (Fig. 1).





Fig. 1: Overview of effects from combining land-based negative emission technologies (bioenergy production coupled with carbon capture and storage (BECCS), afforestation (AFF), Enhanced Weathering (EW), and biochar). The additional CO₂ sequestration is a qualitative estimate based on the author's personal assessment. Technology symbols courtesy of W. Lamb (MCC Berlin).

While biomass-based NETs like afforestation and BECCS are widely discussed, EW is under-represented in this discussion (Minx et al., 2018). EW relies on the uptake of CO₂ via dissolution of minerals based on the natural process of chemical weathering. EW is facilitated by the application of finely ground rock on (agricultural) land, preferably in areas with elevated temperatures and rainfall. The resources for this NET have to be mined and, depending on the type and scale of rollout, the extraction of material can lead the creation of extensive mining areas. However, under a sustainable approach, affected environments could later be used to create biodiversity hotspots (e.g. Tropek et al., 2010; Benes et al., 2003).

It is unavoidable that the intended CO₂ sequestration effect by weathering is naturally accompanied by the release of elements with consequences for the environment (Kantola et al., 2017) and consequently the involved carbon pools. The release of elements that are important plant nutrients (e.g., potassium, phosphorus, magnesium) can be beneficial for additional CO₂ sequestration via organic carbon formation. In addition, the soil hydrology can be improved, and cation exchange capacity increased under optimal grain size distribution and mineral selection. In contrast, effects of potentially harmful trace element release (by choosing less suitable material) might be needed to be alleviated. However, an integrated framework to achieve optimization of interrelated effects between land-based NETs has yet to be developed, specifically for a global scale management of carbon pools.

To tackle the issue of climate change with negative carbon emission strategies on a global and comprehensive scale, it seems advisable to consider all proposed terrestrial biomass-based NETs, like e.g., BECCS, afforestation, and biochar to explore synergistic effects (Fig. 1). A scenario can be envisioned,

- hat gelöscht: solely
- hat formatiert: Englisch (Vereinigtes Königreich)
- Formatiert: Abstand Nach: 12 Pt.
- hat gelöscht: an assessment of
- hat gelöscht: of NETs
- hat formatiert: Englisch (Vereinigtes Königreich)
- hat formatiert: Englisch (Vereinigtes Königreich)
- hat formatiert: Englisch (Vereinigtes Königreich)



- hat gelöscht:
- Formatierte Tabelle
- hat formatiert: Englisch (Vereinigtes Königreich)
- hat gelöscht:
- hat gelöscht: Bioenergy
- hat gelöscht: Afforestation
- hat gelöscht: Biochar).
- hat gelöscht: Taking the example of Enhanced Weathering, the application of finely ground rock on agricultural land, it
- hat formatiert: Englisch (Vereinigtes Königreich)
- hat formatiert: Schriftart: Kursiv, Englisch (Vereinigtes Königreich)
- hat formatiert: Englisch (Vereinigtes Königreich)
- hat gelöscht: , and others
- hat formatiert: Englisch (Vereinigtes Königreich)
- hat formatiert: Englisch (Vereinigtes Königreich)
- hat gelöscht: by
- hat gelöscht:
- hat formatiert: Schriftart: Kursiv
- hat gelöscht: ¶

where rock powder and biochar are applied to agricultural land, which is used for bioenergy plant production (for further use in BECCS technology). Rock material would release geogenic nutrients and biochar could enhance the release of nutrients (Atkinson et al., 2010) and the overall crop productivity (Jeffery et al., 2011).

5 In combination with envisioned and deployed afforestation efforts, which often take place in tropical areas with depleted soils (Nilsson and Schopfhauser, 1995; Grainger, 1988; Zomer et al., 2008), **rock powder deployment for EW** could be an added, if not essential, benefit. The low capacity of these soils to retain highly soluble industrial fertilisers suggests the use of other forms of slow release fertiliser, like rock dust as a complement (Leonardos et al., 1987; Manning, 2015), or new emerging rock-based fertilizers (Ciceri and Allamore, 2019), which can, as a side effect, increase the retention of industrial fertilizers, that may still be needed in addition. The ultimate need for an intense management and design of a suitable soil to supply suitable conditions for tree growth can be deduced from a published extreme scenario, which **envisions** large scale afforestation of deserts (Ornstein et al., 2009).

10 It seems advisable to combine proposed NET methods to achieve an optimal carbon pool management for negative emissions and ensure food security over centuries at the global scale. To achieve this, interdisciplinary efforts are necessary (Fig. 1) and some of the key issues are reviewed here to point out the **future research directions**.

Nutrient pool

20 Increasing atmospheric CO₂ **concentrations** and an increasing world population will lead to challenges in the nutritional supply **for** large parts of Earth's population (Smith and Myers, 2018; Myers et al., 2014). In combination with partly declining resources **of natural mineral fertilisers** (Manning, 2015 and Suppl. Mat. S1), **alternative nutrient supplies, i.e.**, from rock products, are of high interest (Ciceri and Allamore, 2019). This idea has been discussed earlier (Van Straaten, 2006; van Straaten, 2002) and was recently revived in the context of **EW** (Beerling et al., 2018; **Hartmann et al., 2013**). However, this issue extends further, if biomass-based NETs are considered for large scale deployment. **While nutrients like P or K are normally supplied via mineral dissolution in natural systems, nitrogen is in general supplied via fixation of N from the atmosphere** (Graham and Vance, 2000). **In some ecosystems N supply via rocks might be a relevant source** (Houlton et al., 2018; Holloway and Dahlgren, 2002). **In general, and specifically under intensified demand scenarios created by enhanced biomass growth rock, rock N supply will not keep up with the demand.**

30 Many options of carbon dioxide removal rely on the production of biomass (*i.e.*, biochar, afforestation, carbon capture and storage from bio energy (BECCS), bio fuels). These CDR-methods demand, if driven to an optimum, more geogenic nutrients as typically available to plants **from** the soil-rock-systems in the long-term, specifically in humid, tropical areas, where soils are deeply weathered and show naturally low nutrient contents (Hamdan and Bumham, 1996) that could not supply an additional intense biomass growth. A study **on** commercially exploited forests in the U.S. points out that intensive harvest can withdraw more nutrients from the soils than can naturally be resupplied (de Oliveira Garcia et al., 2018).

hat gelöscht: .

hat gelöscht:

hat gelöscht: it

hat gelöscht: :

hat gelöscht:

hat gelöscht: envision

hat gelöscht: direction

hat gelöscht: of

hat gelöscht: (Manning, 2015 and Suppl. S1), alternative nutrient supplies, i.e.

hat gelöscht: Enhanced Weathering side effects

hat gelöscht:

hat formatiert: Schriftart: Kursiv

hat gelöscht: .

hat gelöscht: in

hat gelöscht: of

hat gelöscht: scenarios

hat gelöscht: , which would also apply to extensive afforestation scenarios...

hat gelöscht: ¶

The intensive withdrawal of nutrients should be included in a framework for biogeochemical cycle management under NET deployment. The withdrawal of each K and P from cropland amounts globally to more than 8 Mt a⁻¹ (Suppl. Mat. Fig. S2-2). For many ecosystems the natural resupply and potentially limiting effects under absence of deliberate fertilisation practices, is unknown or merely based on meta-analyses or model studies.

Due to desired global carbon sequestration goals (as in models for afforestation), growth rates will likely be driven to the maximum, which implies an increased demand of nutrients. Models show that N and P limit the global carbon sequestration potential for forests (Goll et al., 2012; Kracher, 2017). Nutrient release by EW can therefore play a relevant role in supporting the high demand. Particular rock classes contain, on average, higher K, P (Fig. 3) or micronutrients like Zn or Se, than others. To ensure a balanced supply of the needed elements, it is therefore necessary to consider not one specific rock type during EW application.

Considering a subtropical weathering scenario in combination with Miscanthus growth for BECCS, acid igneous rocks show a high potential to (partly) resupply extracted potassium, while (ultra-)basic rocks can (partly) resupply phosphorus (Fig. 3). Many earlier studies on EW focussed on dunite to maximize inorganic CO₂ sequestration, with the side effect of adding high levels of Ni and Cr to the system (e.g. Schuiling and Krijgsman, 2006; Hangx and Spiers, 2009). Later, basalt was added to the discussion (Beerling et al., 2018; Strefler et al., 2018; Hartmann et al., 2013). It is characterised by an elevated geogenic nutrient supply compared to ultrabasic rocks like dunite (Fig. 3) but still features a sufficiently high inorganic CO₂ sequestration potential (Fig. 2, and Strefler et al., 2018). Future application scenarios will likely use a mixture of locally available material to optimize both organic and inorganic carbon storage. Optimising the nutrient composition may come at the price of reducing the inorganic carbon sequestration potential, as some rock types with high nutrient content have low sequestration potentials (Fig. 2). If additional soil properties, like cation exchange capacity, water content/hydrology, and pH, are optimised, this reduction of inorganic carbon sequestration may be compensated by elevated biomass uptake and organic carbon storage.

hat verschoben (Einfügung) [1]

hat gelöscht: the

hat gelöscht: ,

hat gelöscht: already

hat gelöscht: Enhanced Weathering

hat gelöscht: types

hat gelöscht: (

hat formatiert: Englisch (Vereinigtes Königreich)

hat nach oben verschoben [1]: Fig.

hat gelöscht: 2),

hat gelöscht: .

hat gelöscht: (like before

hat gelöscht: or

hat gelöscht: (Strefler et al., 2018)

hat gelöscht: .

hat gelöscht: reduction

hat gelöscht: type

hat gelöscht: 2).

hat gelöscht: ¶





Fig. 2: The averaged relative K and P contents of igneous rocks (middle point: median, whiskers: 10/90%iles, some cut of at extreme ends for better graphic representation), classified by SiO₂ content (ultrabasic: <45%, basic: 45-52%, intermediate: 52-63%, acid: >63%). The circles indicate their potential to capture CO₂. Statistical data are from the GEOROC database (Sarbas, 2008), details in [Suppl. Mat. S3](#). Documentation on CO₂ capture potential calculation in [Suppl. Mat. S4](#). A map with the global distribution of all classes is available in [Suppl. Mat. S5](#). Basalt and dunit were added separately as reference for



hat gelöscht:

Gelöschte Zellen

Formatierte Tabelle

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Abstand Vor: 6 Pt.

hat gelöscht: : The averaged K and P contents of selected volcanic rocks in combination with their potential to capture CO₂. All shown rocks are selected for a net positive sequestration effect based on the optimistic emissions scenario from Moosdorf et al. (2014). The discussed dunit contains so little P and K that it cannot be shown in the figure (< 0.01 wt% each). Statistical data for shown rocks

hat gelöscht: Supplement S2.

hat gelöscht: Supplement S3.

hat gelöscht: ¶





Fig. 3. Weathering release rates (circles; bars as variability indicator) of P and K from selected rocks (assuming their full dissolution under a natural subtropical weathering scenario) and extraction of those nutrients by harvesting Miscanthus energy grass (blue/red areas indicate range between min. and max. nutrient content of different Miscanthus species multiplied with min. and max. yield reported in Brosse et al. (2012), Suppl. Mat. Tabs. S2-1 and S2-2). Details on rock dissolution and nutrient release rates in Suppl. Mat. S6 and on plant nutrient removal and additional to Miscanthus data on major crops in Suppl. Mat. S2. Dunite values for the 10%ile were cut off for better overall visibility

hat formatiert: Absatz-Standardschriftart, Schriftart: Times New Roman, Englisch (Vereinigtes Königreich)



hat gelöscht: hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Abstand Nach: 0 Pt.

hat gelöscht: 3:

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht:). Details on rock dissolution and nutrient release rates in Suppl. S4 and on plant nutrient removal and additional to Miscanthus data on major crops in Suppl. S5.

Formatiert: Beschriftung

hat gelöscht: Considering extreme scenarios of biomass plantations (e.g. in Boysen et al., 2016), the intense extraction of nutrients needs to be accounted for (globally more than 8 Mt P and K per year; c.f. nutrient removal from cropland Suppl. S4). ¶

hat gelöscht:

hat gelöscht: .

hat gelöscht: There are further potential effects that contribute to the overall amelioration effect of rock powder. It has been shown that the cation content

hat gelöscht: regions where

hat gelöscht: . in

hat formatiert: Schriftart: Kursiv

hat gelöscht: .

hat gelöscht: ¶

The introduction of additional nutrients to the soil system will not necessarily lead to an additional CO₂ uptake and increased CO₂ sequestration potentials of biomass-based NETs, if enough nutrients are supplied by traditional fertilisation. However, forest areas may benefit from slow release long term available nutrients as they may be less easy to be re-supplied on a regular basis by agrotechnical machinery. Also, industrial fertiliser may be unaffordable in low income regions, thus rock products could replace parts of the fertiliser (Ciceri and Allamore, 2019). A wider adoption of rock product utilisation may also lead to the development of new and optimised application techniques.

10 Nutrient retention

Nutrients released from industrial fertilizers or from natural rock products can be taken up by the plant, washed away, or retained by the properties of the soil. The latter is called retention capacity and is important to store nutrients in a plant available form. It has been shown that the weathering of basaltic material increases the cation exchange capacity, leading to an increased retention of nutrients (Anda et al., 2013, 2015). This is especially important for areas in which nutrients from industrial fertiliser material are quickly washed out, e.g. from the deeply weathered soils (e.g. oxisols) in tropical regions (Leonardos et al., 1987; Ciceri et al., 2017). In such settings, it will be favourable to establish improved soil conditions with optimised nutrient retention.

Another application case is the fertilisation of forests, specifically on areas which are re-forested after agricultural use. With increasing atmospheric CO₂ concentrations, an increase in biomass productivity on non-agricultural areas is expected through the CO₂ fertilisation effect (e.g. Ciais et al., 1995; Körner et al., 2007; Norby and Zak, 2011), especially with regard to afforestation efforts and general tree growth.

5 This effect has yet to be clearly shown (Leuzinger et al., 2011), and is likely limited by soil fertility (Oren et al., 2001; Bader et al., 2013). It can already be observed that nutrient supply by rock weathering, specifically P, K, Mg, and Ca, can be the limiting factor of tree growth under elevated atmospheric CO₂ (Jonard et al., 2015). Woodland soils might be amended with selected minerals or rocks to supply sufficient nutrients to keep up growth under elevated atmospheric pCO₂ conditions and organically bind carbon, a scenario that should be explored further for its potential to enlarge affected carbon pools. At some point, depending on the environmental setting, biomass growth will be limited by nutrient supply and as such, model outputs for CO₂ sequestration potentials of afforestation are likely to be overestimated, if geogenic nutrient cycles are not included in the assessment, as Goll et al. (2012) have shown exemplary for the C, N, and P cycles using a model.

10 The CO₂ sequestration effect of afforestation is even larger if soil organic carbon changes are taken into consideration: Depending on the underlying lithology, the organic carbon pools can be increased (Li et al., 2017), a process that may be optimised by the spreading of selected rock products.

Overall, specific element deficits in soils need to be mapped, since it can also affect the plant content of valuable, if not essential elements for human nutrition (Zhang et al., 2017; Hengl et al., 2017; White and Zasoski, 1999). It is necessary to be able to predict, which application amounts of elements causes a certain response in the biomass pool above and below ground. Such data are still scarce and inconclusive (Manning, 2010) but are needed if EW should be used as a method to help manage carbon and nutrient pools.

15 Biochar is another NET that has a beneficial effect on the retention of nutrients (Fig. 1). Due to its large surface area and increased cation exchange capacity, nutrients can be sustainably retained in soils (Lehmann, 2007; Liang et al., 2006), effectively saving applied fertiliser (Laird et al., 2010).

20 Increased nutrient retention may increase the overall CO₂ sequestration potential of biomass-based NETs through the long-term availability of nutrients. However, the order of magnitude of the effect remains to be shown.

30 Soil hydrology

35 The availability of water is essential for high crop yields (Rockstrom et al., 2007) and soil hydraulic properties fundamentally steer the availability of water to plants (Bodner et al., 2015; Pinheiro et al., 2019). The soil hydraulic conductivity is a measure of how easy water can percolate through the soil column. It depends largely on the grain size distribution of the soil. Roughly, coarse (sandy) soils have a higher hydraulic conductivity than fine (clayey) soils (Rawls et al., 1982). Spreading large amounts of rock products with very small grain sizes (silty to clayey) on land potentially leads to a decrease in soil hydraulic conductivity, which may lead to decreased weathering speeds due to local pore water oversaturation or enhanced surface runoff. However, there are some indications that the addition of biochar can be used to control hydraulic conductivity (Masiello et al., 2015; Barnes et al., 2014), which could enable

hat gelöscht: is

hat gelöscht: to be

hat gelöscht: nutrients provided

hat gelöscht: like

hat gelöscht: .

hat gelöscht: influenced

hat gelöscht: an issue that is increasingly addressed with a focus on...

hat gelöscht: Enhanced Weathering

hat gelöscht:

hat gelöscht:

hat gelöscht: Spreading large amounts of rock products with too small grain sizes on land potentially leads to a decrease in soil hydraulic conductivity, which may lead to decreased weathering speeds due to local pore water oversaturation or enhanced surface runoff. However, there are some indications that biochar can be used to control hydraulic conductivity (Masiello et al., 2015; Barnes et al., 2014), which could enable the use of smaller grain sizes for Enhanced Weathering, enhancing its potential, which strongly depends on the grain size (Strefler et al., 2018). ¶ Biochar could also

hat gelöscht: ¶

the use of smaller grain sizes for EW, enhancing its potential, which strongly depends on the grain size (Strefler et al., 2018).

As another hydraulic property, the water holding capacity determines how much water is kept in the soils and potentially is available to plants. This parameter becomes increasingly important with more frequently appearing droughts due to climate change (Kang et al., 2009). Biochar could be used to improve the water holding capacities of soils (Omondi et al., 2016; Liu et al., 2017), and also increase the plant available water in some cases (Masiello et al., 2015). This may render dryer regions or areas with unfavourable soil physical properties (Basso et al., 2013) usable for bio energy plants and/or afforestation. There are also indications that improvement of soil hydrology by biochar may increase yield potentials (Akhtar et al., 2014; Xu et al., 2015; Al-Wabel et al., 2018).

It is important to point out that all potential changes of soil physical properties due to biochar application strongly depend on its type, more specifically the feedstock and production temperature (Gul et al., 2015). The combination of rock product and biochar application however was not addressed in previous research, at all, but may provide a relevant potential to increase and maintain soil carbon.

hat gelöscht: yet

15 Soil pH

Soil pH steers the availability of elements to plants (Kabata-Pendias, 2010; Loomis and Morris, 1983). At pH values well below 7, nutrients become less available to plants and potentially harmful trace metals are successively mobilised. Nitrogen-fixing bacteria are also depending on a specific pH (Graham and Vance, 2000). Soil acidification on heavily used cropland is a problem (Helyar and Porter, 1989), which may lead to a decrease in crop yields. The main reason is the higher mobility of most exchangeable metals at low pH, which decreases logarithmically with increasing pH (Kabata-Pendias, 2010; Robinson et al., 1996; Tack et al., 1996; Harter, 1983).

Levels of pH 6 and higher generally ensures very low levels of exchangeable harmful metals, with the exception of arsenic, depending on the oxidation state (Dixit and Hering, 2003). The release of base cations from rock flour leads to a soil pH increase. Studies have demonstrated the effectiveness of basalt powder application in raising the soil pH up to 8 and higher (e.g. Gillman et al., 2001; Nunes et al., 2014). The effect is similar to agricultural liming, which is a common practice to counteract soil acidification on cropland (West and McBride, 2005). Despite the fast dissolution rate of carbonate minerals, they are in general, until today, not considered for EW scenarios, because of possible carbonate precipitation and subsequent CO₂ release in the ocean (Hartmann et al., 2013) or due to the potential release of CO₂ by excess fertiliser application (Semhi et al., 2000; Perrin et al., 2008).

The potential of carbonates in EW strategies remains to be studied, while silicate application is in the focus of recent research (Taylor et al., 2015). It could be a potential economic benefit to replace agricultural lime by silicate rock flour, bearing in mind that silicate dissolution rates are in general several orders of magnitude lower, with strong variability between different minerals (Lasaga, 1995; Brantley et al., 2008). Thus, the efficacy is decreased due to the slower release rate of cations, but other properties like nutrient retention or soil hydrology might be improved (cf. previous chapter). It remains to be investigated how (fast) the termination of pH stabilising silicate rock powder application will affect the soils. If relatively immobile potentially harmful metals accumulate at elevated pH values over the application period, an excessive and harmful release of toxic substances might occur in case of a future drop of pH due to

hat gelöscht: , establishing

hat gelöscht: at pH of 6 and higher

hat gelöscht: Arsenic

hat gelöscht: Despite the fast dissolution rate of carbonate minerals, they are in general until today not considered for Enhanced Weathering scenarios, because of possible carbonate precipitation and subsequent CO₂ release in the ocean (Hartmann et al., 2013)

hat gelöscht: Enhanced Weathering

hat gelöscht: cessation

hat gelöscht: to provide management suggestions

hat gelöscht: ¶

changes in pH controlling minerals, land use or general environmental conditions. Once the deployment of material rich in trace elements of concern is started, it is obligatory to maintain a stabilised pH environment, strengthening the need for material with low **harmful** trace element concentrations, (requirements may differ depending on ecosystem type).

- 5 Assuming that pH stabilisation and beneficial changes in soil hydrology (cf. previous section) are achievable by biochar and **EW** a significant additional CO₂ uptake can be expected, based on the effect that soils are made usable for biomass-based NETs, that couldn't support sustainable biomass growth before.

Soil biota

10 Chemical weathering of rocks can be significantly mediated by macro- and microbiota (Schwartzman and Volk, 1989; Uroz et al., 2009; Hoffland et al., 2004; Blouin et al., 2013), although the order of magnitude is a matter of debate (Drever, 1994). This is specifically the case for mycorrhizal fungi and microbes, which create physico-chemical conditions that **accelerate the dissolution of minerals** (Taylor et al., 2015). **The weatherability of minerals depends on the type of bioinoculant (Nishanth and Biswas, 2008; Benzerara et al., 2005; Cuadros, 2018), implicating that a supervision and management of the soil micro-**

15 **biota is necessary to optimise both crop yields and rock weathering.** Microbial populations in soils respond to the addition of biochar (Warnock et al., 2007) by providing a refuge for bacteria and fungi (Pietikainen et al., 2000; Saito, 1990), increasing nutrient availability, creating favourable pH conditions and other processes **discussed in Lehmann et al. (2011).**

20 Earthworms have been observed to thrive in biochar amended soils (Topoliantz and Ponge, 2005). Increased abundance of earthworms will likely increase bioturbation effects (Carcaillet, 2001; Major et al., 2010), leading to a better distribution of biochar and rock flour in deeper layers of the amended soils, increasing reactive surfaces of mineral grains. Bioturbation might also be a key process to achieve high CO₂ sequestration rates by weathering, **as, e.g., Earthworms can enhance mineral weathering (Carpenter et al., 2007; Carpenter et al., 2008) and contribute to** the downward transport of added rock products into

25 deeper soil layers (Taylor et al., 2015).

Trace metals

Soils are an important sink in the environmental cycling of trace metals (Kabata-Pendias, 1993). Besides naturally occurring concentrations, depending on the **underlying lithology**, the major source of trace metals to soils is agricultural practice, leading to an enrichment due to the application of manure, sewage sludge, fertilisers, and pesticides, which all contain metals to a certain extent (Gonnelli and Renella, 2013). Field studies using sewage sludge as fertiliser have shown a marked uptake by the crops and increased mobilisation of trace metals in the runoff water (Alloway, 2013). Adding to the anthropogenic input, the introduction of additional rock products with **elevated** levels of trace metals **Fig. 4** could lead to a critical exceedance of environmental thresholds if improper rock material is used **due to inconsiderate**

30 **management.** This is, however, relating to the solubility of minerals within the used rock type and the redox and pH conditions. **An EW soil incubation experiment using an olivine-rich rock product, with elevated Ni and Cr concentrations in the source material (Fig. 4),** showed only a few occurrences of

35

hat gelöscht: , where considered to be of concern

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: Enhanced Weathering

hat gelöscht:

hat gelöscht: accelerate the dissolution of minerals.

hat gelöscht: (Lehmann et al., 2011)

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: through

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: base rock

hat gelöscht: increased

hat gelöscht: .

hat gelöscht: A

hat gelöscht: Enhanced Weathering

hat gelöscht: with

hat gelöscht: application

hat gelöscht: ¶

elevated Cr levels but no Ni increase in the aquatic solution compared to a blank treatment, leading to the conclusion that the soil solid phase will be successively enriched in those elements (Renforth et al., 2015). The availability of heavy metals to biota remains an issue of ongoing discussion (Nagajyoti et al., 2010). The main elements of concern in source rocks with the highest sequestration potential (ultramafic rocks) are Ni and Cr. Especially the early discussed dunite application for EW must trigger the discussion about its high Cr and Ni contents (Fig. 4) and is therefore ruled out for large scale application on cropland.




Fig. 4: Contents of Ni and Cr in igneous rocks, classified by SiO₂ content (cf. Fehler! Verweisquelle konnte nicht gefunden werden.). Circles indicate median values, whiskers are the 10th and 90th percentiles. Extreme values of percentiles were cut off for better visibility of data. Extra figure for visualisation of dunite as it features extremely high values. For detailed statistics, cf. Suppl. Mat. S7.

If an application with rocks of high trace metal concentrations of concern is considered, it is necessary to stabilise the soil pH even after cessation of such actions in order to maintain the fixation of toxic elements, because of the strong pH control on metal mobility. A study of long-term sewage sludge application has shown that the pH had to be stabilised by liming in order to prevent phyto-toxicity of Cu and Zn (McBride et al., 2004). Additionally, the metal availability to plants has been shown to be influenced by the soil texture, with marked differences for different elements (Qian et al., 1996). This underlines the necessity to control or specifically design the grain size distribution of the soil to control water content, pH and oxygen content. To further ameliorate the issue, biochar, which has been shown to immobilise heavy metals in soils, depending on feedstock and production conditions (Ahmad et al., 2014; Beesley et al., 2011) could be jointly applied with rock powder. This would mean that potential limitations of fertiliser or rock spreading due to thresholds put in place for environmental protection could be overcome by a sensible management of biochar utilisation. Applying biochar products does not remove elements of

hat gelöscht: Enhanced Weathering

hat gelöscht: (Tab. 1)

hat gelöscht: Tab. 1 Statistical data for Ni and Cr in dunites from the GEOROC database (Sarbas, 2008).

Formatiert: Abstand Vor: 6 Pt.

Formatierte Tabelle

hat gelöscht: Element ... [5]

Formatiert: Abstand Vor: 0 Pt., Nach: 0 Pt.

hat gelöscht: ¶

concern, but the problem of heavy metal accumulation could be dampened bioremediation through heavy metal accumulating plants (Rajkumar et al., 2012). This in turn could be a potential new source of raw material for industrial use (Schuiling, 2013), though it is likely not applicable on a global scale, since this would compete against food and energy plant production, which is already an issue (Tilman et al., 2009).

5 The alleviation of trace metal effects does not directly affect CO₂ sequestration rates but could overall increase potential deployment areas for EW.

hat gelöscht: remediation

hat gelöscht: Enhanced Weathering

Conclusion

Looking forward it is likely that a portfolio of options will be established to optimize the sequestration effect and minimize negative impacts. The combination of previously separately studied NETs to increase the sequestered carbon pool should consider the management of biogeochemical cycles and optimize the combined application of Enhanced Weathering and biochar in context of biomass-based methods like BECCS and afforestation to maximize carbon capture as well as food production. It is therefore essential to address combined effects of NET co-deployment in future research projects.

hat gelöscht: Synthesis

hat formatiert: Englisch (Vereinigtes Königreich)

As all presented interactions take place in the soil, future research should put a focus on creating an optimized soil product for an optimal long-term sustainable carbon management. We propose that research around biomass-based NET interactions becomes the science of artificial soil products, which are most likely created on depleted and degraded soils especially in the sub(tropics). It may consist of the locally available "base soil" mixed with charcoal products to enhance hydraulic properties and nutrient retention, as well as rock powder, which raises the soil pH, provides nutrients and sequesters CO₂ at the same time. This engineered and managed soil could increase carbon pools and crop production, while contributing to tackle the issue of climate change. It remains to be studied where suitable material is available at the regional scale (Suppl. Mat. S5). The parameterisation of element release rates permitting a sustainable management are still subject to large uncertainties and the effects of massive rock product spreading will change the soil structure to an extent that remains to be explored.

hat gelöscht:

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht:

hat gelöscht: .

hat formatiert: Englisch (Vereinigtes Königreich)

25 The introduction of non-authigenic material into the environment, even if of bio- or geogenic origin, will increase the entropy of the system, making it difficult and expensive (energy and economic-wise) to quickly revert back into the "undisturbed" state, once large-scale deployment started. Thus, the continuous deployment of NETs at the global scale in an order of magnitude that would measurably impact atmospheric CO₂ levels must be seriously weighed. However, the high probability of NET adoption in the near future makes it imperative to create efficient cooperation networks across all involved disciplines in order to conceive the necessary knowledge on actual CO₂ sequestration potentials and century scale global carbon pool changes.

Competing interests

The authors declare that they have no conflict of interest.

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: ¶

Author contribution statements

This article was conceived by the joint work of J.H. and T.A. Both participated in discussions, planning and writing, with the lead of T.A..

hat formatiert: Englisch (Vereinigtes Königreich)

Acknowledgments

5 This research was executed with the financial support of the German Research Foundation's priority program DFG SPP1689 on "Climate Engineering—Risks, Challenges and Opportunities?" and specifically the CEMICS2 project. Further support came from the [Deutsche Forschungsgemeinschaft \(DFG\) under Germany's Excellence Strategy – EXC 20 2037 'Climate, Climatic Change, and Society' – Project Number: 390683824](#), contribution to the Center for Earth System Research and Sustainability (CEN) of [Universität Hamburg](#), and from the previous EXC177 'CLISAP2', [Universität](#).

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Standard (Web)

hat gelöscht: Cluster of

hat formatiert: Englisch (Vereinigtes Königreich), Muster: Transparent

hat gelöscht: 'CLISAP', EXC177, Universität

hat formatiert: Englisch (Vereinigtes Königreich), Muster: Transparent

hat gelöscht: ¶

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: ¶

References

- Ahmad, M., Rajapaksha, A. U., Lim, J. E., Zhang, M., Bolan, N., Mohan, D., Vithanage, M., Lee, S. S., and Ok, Y. S.: Biochar as a sorbent for contaminant management in soil and water: a review, *Chemosphere*, 99, 19-33, 10.1016/j.chemosphere.2013.10.071, 2014.
- 5 Akhtar, S. S., Li, G., Andersen, M. N., and Liu, F.: Biochar enhances yield and quality of tomato under reduced irrigation, *Agricultural Water Management*, 138, 37-44, 10.1016/j.agwat.2014.02.016, 2014.
- Al-Wabel, M. I., Hussain, Q., Usman, A. R. A., Ahmad, M., Abduljabbar, A., Sallam, A. S., and Ok, Y. S.: Impact of biochar properties on soil conditions and agricultural sustainability: A review, *Land Degradation & Development*, 29, 2124-2161, 10.1002/ldr.2829, 2018.
- 10 Alloway, B. J.: Sources of Heavy Metals and Metalloids in Soils, in: *Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their Bioavailability*, edited by: Alloway, B. J., Springer Netherlands, Dordrecht, 11-50, 2013.
- Anda, M., Shamshuddin, J., and Fauziah, C. I.: Increasing negative charge and nutrient contents of a highly weathered soil using basalt and rice husk to promote cocoa growth under field conditions, *Soil and Tillage Research*, 132, 1-11, 10.1016/j.still.2013.04.005, 2013.
- 15 Anda, M., Shamshuddin, J., and Fauziah, C. I.: Improving chemical properties of a highly weathered soil using finely ground basalt rocks, *CATENA*, 124, 147-161, 10.1016/j.catena.2014.09.012, 2015.
- Atkinson, C., Fitzgerald, J., and Hipps, N.: Potential mechanisms for achieving agricultural benefits from biochar application to temperate soils: a review, *Plant and Soil*, 337, 1-18, 10.1007/s11104-010-0464-5, 2010.
- Bader, M. K. F., Leuzinger, S., Keel, S. G., Siegwolf, R. T. W., Hagedorn, F., Schleppei, P., Körner, C., and Lee, J.: Central European hardwood trees in a high-CO₂ future: synthesis of an 8-year forest canopy CO₂ enrichment project, *Journal of Ecology*, 101, 1509-1519, 10.1111/1365-2745.12149, 2013.
- Barnes, R. T., Gallagher, M. E., Masiello, C. A., Liu, Z., and Dugan, B.: Biochar-Induced Changes in Soil Hydraulic Conductivity and Dissolved Nutrient Fluxes Constrained by Laboratory Experiments, *PLoS ONE*, 9, e108340, 10.1371/journal.pone.0108340, 2014.
- 25 Basso, A. S., Miguez, F. E., Laird, D. A., Horton, R., and Westgate, M.: Assessing potential of biochar for increasing water-holding capacity of sandy soils, *GCB Bioenergy*, 5, 132-143, 10.1111/gcbb.12026, 2013.
- Beerling, D. J., Leake, J. R., Long, S. P., Scholes, J. D., Ton, J., Nelson, P. N., Bird, M., Kantzas, E., Taylor, L. L., Sarkar, B., Kelland, M., DeLucia, E., Kantola, I., Muller, C., Rau, G., and Hansen, J.: Farming with crops and rocks to address global climate, food and soil security, *Nat Plants*, 10.1038/s41477-018-0108-y, 2018.
- 30 Beesley, L., Moreno-Jiménez, E., Gomez-Eyles, J. L., Harris, E., Robinson, B., and Sizmur, T.: A review of biochars' potential role in the remediation, revegetation and restoration of contaminated soils, *Environmental Pollution*, 159, 3269-3282, 10.1016/j.envpol.2011.07.023, 2011.
- [Benes, J., Kepka, P., and Konvicka, M.: Limestone quarries as refuges for European xerophilous butterflies, *Conservation Biology*, 17, 1058-1069, DOI 10.1046/j.1523-1739.2003.02092.x, 2003.](#)
- 35 [Benzerara, K., Yoon, T. H., Menguy, N., Tylliszczak, T., and Brown, G. E., Jr.: Nanoscale environments associated with bioweathering of a Mg-Fe-pyroxene, *Proc Natl Acad Sci U S A*, 102, 979-982, 10.1073/pnas.0409029102, 2005.](#)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

hat gelöscht: ¶

Blouin, M., Hodson, M. E., Delgado, E. A., Baker, G., Brussaard, L., Butt, K. R., Dai, J., Dendooven, L., Peres, G., Tondoh, J. E., Cluzeau, D., and Brun, J. J.: A review of earthworm impact on soil function and ecosystem services, *Eur. J. Soil Sci.*, 64, 161-182, 10.1111/ejss.12025, 2013.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

Bodner, G., Nakhforoosh, A., and Kaul, H.-P.: Management of crop water under drought: a review, *Agron. Sustain. Dev.*, 35, 401-442, 10.1007/s13593-015-0283-4, 2015.

hat gelöscht: Boysen, L. R., Lucht, W., Gerten, D., and Heck, V.: Impacts devalue the potential of large-scale terrestrial CO2 removal through biomass plantations, *Environmental Research Letters*, 11, 095010, 10.1088/1748-9326/11/9/095010, 2016.

Brantley, S. L., Kubicki, J. D., and White, A. F.: Kinetics of water-rock interaction, Springer, 2008.

hat formatiert: Englisch (Vereinigtes Königreich)

Brosse, N., Dufour, A., Meng, X., Sun, Q., and Ragauskas, A.: *Miscanthus*: a fast-growing crop for biofuels and chemicals production, *Biofuels, Bioproducts and Biorefining*, 6, 580-598, 10.1002/bbb.1353, 2012.

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

Caldeira, K., Bala, G., and Cao, L.: The Science of Geoengineering, *Annu Rev Earth Pl Sc*, 41, 231-256, 10.1146/annurev-earth-042711-105548, 2013.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

Carcaillet, C.: Soil particles reworking evidences by AMS 14C dating of charcoal, *Comptes Rendus de l'Académie des Sciences - Series IIA - Earth and Planetary Science*, 332, 21-28, 10.1016/s1251-8050(00)01485-3, 2001.

Carpenter, D., Hodson, M. E., Eggleton, P., and Kirk, C.: Earthworm induced mineral weathering: Preliminary results, *European Journal of Soil Biology*, 43, S176-S183, 10.1016/j.ejsobi.2007.08.053, 2007.

Carpenter, D., Hodson, M. E., Eggleton, P., and Kirk, C.: The role of earthworm communities in soil mineral weathering: a field experiment, *Mineralogical Magazine*, 72, 33-36, 10.1180/minmag.2008.072.1.33, 2008.

Ciais, P., Tans, P. P., Trolier, M., White, J. W. C., and Francey, R. J.: A Large Northern Hemisphere Terrestrial CO2 Sink Indicated by the 13C/12C Ratio of Atmospheric CO2, *Science*, 269, 1098-1102, 10.1126/science.269.5227.1098, 1995.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

Ciceri, D., de Oliveira, M., Stokes, R. M., Skorina, T., and Allamore, A.: Characterization of potassium agrominerals: Correlations between petrographic features, comminution and leaching of ultrapotassic syenites, *Minerals Engineering*, 102, 42-57, 10.1016/j.mineng.2016.11.016, 2017.

Ciceri, D., and Allamore, A.: Local fertilizers to achieve food self-sufficiency in Africa, *Sci Total Environ*, 648, 669-680, 10.1016/j.scitotenv.2018.08.154, 2019.

Cuadros, J.: Clay minerals interaction with microorganisms: a review, *Clay Minerals*, 52, 235-261, 10.1180/claymin.2017.052.2.05, 2018.

de Oliveira Garcia, W., Amann, T., and Hartmann, J.: Increasing biomass demand enlarges negative forest nutrient budget areas in wood export regions, *Scientific reports*, 8, 5280, 10.1038/s41598-018-22728-5, 2018.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

Dixit, S., and Hering, J. G.: Comparison of arsenic(V) and arsenic(III) sorption onto iron oxide minerals: implications for arsenic mobility, *Environ Sci Technol*, 37, 4182-4189, 10.1021/es030309t, 2003.

Drever, J. I.: The Effect of Land Plants on Weathering Rates of Silicate Minerals, *Geochim Cosmochim Acta*, 58, 2325-2332, Doi 10.1016/0016-7037(94)90013-2, 1994.

Fuss, S., Canadell, J. G., Peters, G. P., Tavoni, M., Andrew, R. M., Ciais, P., Jackson, R. B., Jones, C. D., Kraxner, F., Nakicenovic, N., Le Quere, C., Raupach, M. R., Sharifi, A., Smith, P., and Yamagata, Y.: Betting on negative emissions, *Nature Clim. Change*, 4, 850-853, 10.1038/nclimate2392, 2014.

hat gelöscht: ¶

Fuss, S., Jones, C. D., Kraxner, F., Peters, G. P., Smith, P., Tavoni, M., van Vuuren, D. P., Canadell, J. G., Jackson, R. B., Milne, J., Moreira, J. R., Nakicenovic, N., Sharifi, A., and Yamagata, Y.: Research priorities for negative emissions, *Environmental Research Letters*, 11, 10.1088/1748-9326/11/11/115007, 2016.

5 Fuss, S., Lamb, W. F., Callaghan, M. W., Hilaire, J., Creutzig, F., Amann, T., Beringer, T., de Oliveira Garcia, W., Hartmann, J., Khanna, T., Luderer, G., Nemet, G. F., Rogelj, J., Smith, P., Vicente, J. L. V., Wilcox, J., del Mar Zamora Dominguez, M., and Minx, J. C.: Negative emissions—Part 2: Costs, potentials and side effects, *Environmental Research Letters*, 13, 10.1088/1748-9326/aabf9f, 2018.

Gillman, G. P., Burkett, D. C., and Coventry, R. J.: A laboratory study of application of basalt dust to highly weathered soils: effect on soil cation chemistry, *Soil Research*, 39, 799-811, 10.1071/SR00073, 2001.

10 Goll, D. S., Brovkin, V., Parida, B. R., Reick, C. H., Kattge, J., Reich, P. B., van Bodegom, P. M., and Niinemets, Ü.: Nutrient limitation reduces land carbon uptake in simulations with a model of combined carbon, nitrogen and phosphorus cycling, *Biogeosciences*, 9, 3547-3569, 10.5194/bg-9-3547-2012, 2012.

Gonnelli, C., and Renella, G.: Chromium and Nickel, in: *Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their Bioavailability*, edited by: Alloway, B. J., Springer Netherlands, Dordrecht, 313-333, 2013.

15 [Graham, P. H., and Vance, C. P.: Nitrogen fixation in perspective: an overview of research and extension needs, *Field Crops Research*, 65, 93-106, Doi 10.1016/S0378-4290\(99\)00080-5, 2000.](#)

Grainger, A.: Estimating Areas of Degraded Tropical Lands Requiring Replenishment of Forest Cover, *International Tree Crops Journal*, 5, 31-61, 10.1080/01435698.1988.9752837, 1988.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

20 Gul, S., Whalen, J. K., Thomas, B. W., Sachdeva, V., and Deng, H.: Physico-chemical properties and microbial responses in biochar-amended soils: Mechanisms and future directions, *Agriculture, Ecosystems & Environment*, 206, 46-59, 10.1016/j.agee.2015.03.015, 2015.

Hamdan, J., and Bumham, C. P.: The contribution of nutrients from parent material in three deeply weathered soils of Peninsular Malaysia, *Geoderma*, 74, 219-233, 10.1016/s0016-7061(96)00062-6, 1996.

25 Hangx, S. J. T., and Spiers, C. J.: Coastal spreading of olivine to control atmospheric CO₂ concentrations: A critical analysis of viability, *International Journal of Greenhouse Gas Control*, 3, 757-767, 10.1016/j.ijggc.2009.07.001, 2009.

Harter, R. D.: Effect of Soil pH on Adsorption of Lead, Copper, Zinc, and Nickel, *Soil Science Society of America Journal*, 47, 47-51, 10.2136/sssaj1983.03615995004700010009x, 1983.

30 [Hartmann, J., West, A. J., Renforth, P., Köhler, P., De La Rocha, C. L., Wolf-Gladrow, D. A., Dürr, H. H., and Scheffran, J.: Enhanced chemical weathering as a geoengineering strategy to reduce atmospheric carbon dioxide, supply nutrients, and mitigate ocean acidification. *Rev Geophys*, 51, 113-149, 10.1002/rog.20004, 2013.](#)

Helyar, K. R., and Porter, W. M.: 2 - Soil Acidification, its Measurement and the Processes Involved, in: *Soil Acidity and Plant Growth*, edited by: Robson, A., Academic Press, 61-101, 1989.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

35 Hengl, T., Leenaars, J. G. B., Shepherd, K. D., Walsh, M. G., Heuvelink, G. B. M., Mamo, T., Tilahun, H., Berkhout, E., Cooper, M., Fegeus, E., Wheeler, I., and Kwabena, N. A.: Soil nutrient maps of Sub-Saharan Africa: assessment of soil nutrient content at 250 m spatial resolution using machine learning, *Nutr Cycl Agroecosys*, 109, 77-102, 10.1007/s10705-017-9870-x, 2017.

hat gelöscht: ¶

Hoffland, E., Kuyper, T. W., Wallander, H., Plassard, C., Gorbushina, A. A., Haselwandter, K., Holmstrom, S., Landeweert, R., Lundstrom, U. S., Rosling, A., Sen, R., Smits, M. M., van Hees, P. A., and van Breemen, N.: The role of fungi in weathering, *Frontiers in Ecology and the Environment*, 2, 258-264, 10.1890/1540-9295(2004)002[0258:TROFIW]2.0.CO;2, 2004.

Holloway, J. M., and Dahlgren, R. A.: Nitrogen in rock: Occurrences and biogeochemical implications, *Global Biogeochemical Cycles*, 16, 65-61-65-17, 10.1029/2002gb001862, 2002.

Houlton, B. Z., Morford, S. L., and Dahlgren, R. A.: Convergent evidence for widespread rock nitrogen sources in Earth's surface environment, *Science*, 360, 58-62, 10.1126/science.aan4399, 2018.

Jackson, R. B., Canadell, J. G., Le Quere, C., Andrew, R. M., Korsbakken, J. I., Peters, G. P., and Nakićenovic, N.: Reaching peak emissions, *Nature Clim. Change*, advance online publication, 10.1038/nclimate2892, 2015.

10 Jeffery, S., Verheijen, F. G. A., van der Velde, M., and Bastos, A. C.: A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis, *Agriculture, Ecosystems & Environment*, 144, 175-187, 10.1016/j.agee.2011.08.015, 2011.

15 Jonard, M., Fürst, A., Verstraeten, A., Thimonier, A., Timmermann, V., Potočić, N., Waldner, P., Benham, S., Hansen, K., Merilä, P., Ponette, Q., de la Cruz, A. C., Roskams, P., Nicolas, M., Croisé, L., Ingerslev, M., Matteucci, G., Decinti, B., Bascietto, M., and Rautio, P.: Tree mineral nutrition is deteriorating in Europe, *Global Change Biology*, 21, 418-430, 10.1111/gcb.12657, 2015.

Kabata-Pendias, A.: Behavioral Properties of Trace-Metals in Soils, *Applied Geochemistry*, Supplementary Issue No 2, January 1993, 3-9, 10.1016/S0883-2927(09)80002-4, 1993.

Kabata-Pendias, A.: Trace Elements in Soils and Plants, 4th ed., CRC Press, 548 pp., 2010.

20 Kang, Y., Khan, S., and Ma, X.: Climate change impacts on crop yield, crop water productivity and food security – A review, *Progress in Natural Science*, 19, 1665-1674, 10.1016/j.pnsc.2009.08.001, 2009.

Kantola, I. B., Masters, M. D., Beerling, D. J., Long, S. P., and DeLucia, E. H.: Potential of global croplands and bioenergy crops for climate change mitigation through deployment for enhanced weathering, *Biology Letters*, 13, 20160714, 10.1098/rsbl.2016.0714, 2017.

25 Keller, D. P., Lenton, A., Littleton, E. W., Oschlies, A., Scott, V., and Vaughan, N. E.: The Effects of Carbon Dioxide Removal on the Carbon Cycle, *Current Climate Change Reports*, 4, 250-265, 10.1007/s40641-018-0104-3, 2018.

Körner, C., Morgan, J., and Norby, R.: CO2 Fertilization: When, Where, How Much?, in: *Terrestrial Ecosystems in a Changing World. Global Change — The IGBP Series*, edited by: Canadell, J. G., Pataki, D. E., and Pitelka, L. F., Springer, Berlin, Heidelberg, 2007.

30 Kracher, D.: Nitrogen-Related Constraints of Carbon Uptake by Large-Scale Forest Expansion: Simulation Study for Climate Change and Management Scenarios, *Earth's Future*, 5, 1102-1118, 10.1002/2017ef000622, 2017.

Laird, D., Fleming, P., Wang, B., Horton, R., and Karlen, D.: Biochar impact on nutrient leaching from a Midwestern agricultural soil, *Geoderma*, 158, 436-442, 10.1016/j.geoderma.2010.05.012, 2010.

35 Lasaga, A. C.: Fundamental approaches in describing mineral dissolution and precipitation rates, *Reviews in Mineralogy and Geochemistry*, 31, 23-86, 1995.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

hat gelöscht: ¶

Lehmann, J.: [Bio-energy in the black](#), *Frontiers in Ecology and the Environment*, 5, 381-387, Doi 10.1890/1540-9295(2007)5[381:Bitb]2.0.Co:2, 2007.

[Lehmann, J., Rillig, M. C., Thies, J., Masiello, C. A., Hockaday, W. C., and Crowley, D.](#): Biochar effects on soil biota – A review, *Soil Biology and Biochemistry*, 43, 1812-1836, 10.1016/j.soilbio.2011.04.022, 2011.

- 5 Leonardos, O. H., Fyfe, W. S., and Kronberg, B. I.: The use of ground rocks in laterite systems: An improvement to the use of conventional soluble fertilizers?, *Chemical Geology*, 60, 361-370, 10.1016/0009-2541(87)90143-4, 1987.

Leuzinger, S., Luo, Y., Beier, C., Dieleman, W., Vicca, S., and Körner, C.: Do global change experiments overestimate impacts on terrestrial ecosystems?, *Trends in Ecology & Evolution*, 26, 236-241, 10.1016/j.tree.2011.02.011, 2011.

- 10 Li, D., Wen, L., Zhang, W., Yang, L., Xiao, K., Chen, H., and Wang, K.: Afforestation effects on soil organic carbon and nitrogen pools modulated by lithology, *Forest Ecology and Management*, 400, 85-92, 10.1016/j.foreco.2017.05.050, 2017.

[Liang, B., Lehmann, J., Solomon, D., Kinyangi, J., Grossman, J., O'Neill, B., Skjemstad, J. O., Thies, J., Luizão, F. J., Petersen, J., and Neves, E. G.](#): Black Carbon Increases Cation Exchange Capacity in Soils, *Soil Science Society of America Journal*, 70, 1719, 10.2136/sssaj2005.0383, 2006.

- 15 Liu, Z., Dugan, B., Masiello, C. A., and Gonnermann, H. M.: Biochar particle size, shape, and porosity act together to influence soil water properties, *PLoS One*, 12, e0179079, 10.1371/journal.pone.0179079, 2017.

[Loomis, R. S., and Morris, J. G.](#): *Handbook of Agricultural Productivity, Vol 1 Plant Productivity, Vol 2 Animal Productivity - Rochcigl, M.*, *Bioscience*, 33, 338-339, Doi 10.2307/1309328, 1983.

[Major, J., Lehmann, J., Rondon, M., and Goodale, C.](#): Fate of soil-applied black carbon: downward migration, leaching and soil respiration, *Global Change Biology*, 16, 1366-1379, 10.1111/j.1365-2486.2009.02044.x, 2010.

- 20 Manning, D. A. C.: Mineral sources of potassium for plant nutrition. A review, in: *Agron. Sustain. Dev.*, 2, Springer Netherlands, 281-294, 2010.

Manning, D. A. C.: How will minerals feed the world in 2050?, *Proceedings of the Geologists' Association*, 126, 14-17, 10.1016/j.pgeola.2014.12.005, 2015.

- 25 Masiello, C., Dugan, B., Brewer, C., Spokas, K., Novak, J.-J., Liu, Z., and Sorrenti, G.: Biochar effects on soil hydrology, in: *Biochar for Environmental Management Science, Technology and Implementation*, edited by: Lehmann, J., and Joseph, S., Routledge, 2015.

McBride, M. B., Richards, B. K., and Steenhuis, T.: Bioavailability and crop uptake of trace elements in soil columns amended with sewage sludge products, *Plant and Soil*, 262, 71-84, 10.1023/B:Plso.0000037031.21561.34, 2004.

- 30 Minx, J. C., Lamb, W. F., Callaghan, M. W., Fuss, S., Hilaire, J., Creutzig, F., Amann, T., Beringer, T., de Oliveira Garcia, W., Hartmann, J., Khanna, T., Lenzi, D., Luderer, G., Nemet, G. F., Rogelj, J., Smith, P., Vicente Vicente, J. L., Wilcox, J., and del Mar Zamora Dominguez, M.: Negative emissions—Part 1: Research landscape and synthesis, *Environmental Research Letters*, 13, 10.1088/1748-9326/aabf9b, 2018.

- 35 [Myers, S. S., Zanobetti, A., Kloog, I., Huybers, P., Leakey, A. D., Bloom, A. J., Carlisle, E., Dietterich, L. H., Fitzgerald, G., Hasegawa, T., Holbrook, N. M., Nelson, R. L., Ottman, M. J., Raboy, V., Sakai, H., Sartor, K. A., Schwartz, J., Seneweera, S., Tausz, M., and Usui, Y.](#): Increasing CO2 threatens human nutrition, *Nature*, 510, 139-142, 10.1038/nature13179, 2014.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

hat gelöscht: Moosdorf, N., Renforth, P., and Hartmann, J.: Carbon dioxide efficiency of terrestrial enhanced weathering, *Environ Sci Technol*, 48, 4809-4816, 10.1021/es4052022, 2014.

hat formatiert: Englisch (Vereinigtes Königreich)

hat gelöscht: ¶

Nagajyoti, P. C., Lee, K. D., and Sreekanth, T. V. M.: Heavy metals, occurrence and toxicity for plants: a review, *Environ Chem Lett*, 8, 199-216, 10.1007/s10311-010-0297-8, 2010.

National Research Council: Climate Intervention: Carbon Dioxide Removal and Reliable Sequestration, The National Academies Press, Washington, DC, 154 pp., 2015.

- 5 Nemet, G. F., Callaghan, M. W., Creutzig, F., Fuss, S., Hartmann, J., Hilaire, J., Lamb, W. F., Minx, J. C., Rogers, S., and Smith, P.: Negative emissions—Part 3: Innovation and upscaling, *Environmental Research Letters*, 13, 10.1088/1748-9326/aabff4, 2018.

Nilsson, S., and Schopfhauser, W.: The carbon-sequestration potential of a global afforestation program, *Climatic Change*, 30, 267-293, 10.1007/BF01091928, 1995.

- 10 [Nishanth, D., and Biswas, D. R.: Kinetics of phosphorus and potassium release from rock phosphate and waste mica enriched compost and their effect on yield and nutrient uptake by wheat \(*Triticum aestivum*\), *Bioresource technology*, 99, 3342-3353, 10.1016/j.biortech.2007.08.025, 2008.](#)

Norby, R. J., and Zak, D. R.: Ecological Lessons from Free-Air CO₂ Enrichment (FACE) Experiments, *Annual Review of Ecology, Evolution, and Systematics*, 42, 181-203, 10.1146/annurev-ecolsys-102209-144647, 2011.

- 15 Nunes, J. M. G., Kautzmann, R. M., and Oliveira, C.: Evaluation of the natural fertilizing potential of basalt dust wastes from the mining district of Nova Prata (Brazil), *Journal of Cleaner Production*, 84, 649-656, 10.1016/j.jclepro.2014.04.032, 2014.

Omond, M. O., Xia, X., Nahayo, A., Liu, X., Korai, P. K., and Pan, G.: Quantification of biochar effects on soil hydrological properties using meta-analysis of literature data, *Geoderma*, 274, 28-34, 10.1016/j.geoderma.2016.03.029, 2016.

- 20 Oren, R., Ellsworth, D. S., Johnsen, K. H., Phillips, N., Ewers, B. E., Maier, C., Schafer, K. V. R., McCarthy, H., Hendrey, G., McNulty, S. G., and Katul, G. G.: Soil fertility limits carbon sequestration by forest ecosystems in a CO₂-enriched atmosphere, *Nature*, 411, 469-472, 10.1038/35078064, 2001.

Ornstein, L., Aleinov, I., and Rind, D.: Irrigated afforestation of the Sahara and Australian Outback to end global warming, *Climatic Change*, 97, 409-437, 10.1007/s10584-009-9626-y, 2009.

- 25 Perrin, A. S., Probst, A., and Probst, J. L.: Impact of nitrogenous fertilizers on carbonate dissolution in small agricultural catchments: Implications for weathering CO₂ uptake at regional and global scales, *Geochimica et Cosmochimica Acta*, 72, 3105-3123, 2008.

Pietikainen, J., Kiikkila, O., and Fritze, H.: Charcoal as a habitat for microbes and its effect on the microbial community of the underlying humus, *Oikos*, 89, 231-242, 10.1034/j.1600-0706.2000.890203.x, 2000.

- 30 [Pinheiro, E. A. R., de Jong van Lier, Q., and Šimůnek, J.: The role of soil hydraulic properties in crop water use efficiency: A process-based analysis for some Brazilian scenarios, *Agricultural Systems*, 173, 364-377, 10.1016/j.agsy.2019.03.019, 2019.](#)

Qian, J., Shan, X.-q., Wang, Z.-j., and Tu, Q.: Distribution and plant availability of heavy metals in different particle-size fractions of soil, *Sci. Total Environ.*, 187, 131-141, 10.1016/0048-9697(96)05134-0, 1996.

Rajkumar, M., Sandhya, S., Prasad, M. N. V., and Freitas, H.: Perspectives of plant-associated microbes in heavy metal phytoremediation, *Biotechnology Advances*, 30, 1562-1574, 10.1016/j.biotechadv.2012.04.011, 2012.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

hat gelöscht: ¶

[Rawls, W. J., Brakensiek, D. L., and Saxton, K. E.: Estimation of Soil Water Properties, Transactions of the ASAE, 25, 1316-1320, 10.13031/2013.33720, 1982.](#)

Renforth, P., Pogge von Strandmann, P. A. E., and Henderson, G. M.: The dissolution of olivine added to soil: Implications for enhanced weathering, *Applied Geochemistry*, 61, 109-118, 10.1016/j.apgeochem.2015.05.016, 2015.

- 5 Robinson, B. H., Brooks, R. R., Kirkman, J. H., Gregg, P. E. H., and Gremigni, P.: Plant-available elements in soils and their influence on the vegetation over ultramafic ("serpentine") rocks in New Zealand, *Journal of the Royal Society of New Zealand*, 26, 457-468, 10.1080/03014223.1996.9517520, 1996.

[Rockstrom, J., Lannerstad, M., and Falkenmark, M.: Assessing the water challenge of a new green revolution in developing countries, Proc Natl Acad Sci U S A, 104, 6253-6260, 10.1073/pnas.0605739104, 2007.](#)

- 10 Rogelj, J., Popp, A., Calvin, K. V., Luderer, G., Emmerling, J., Gernaat, D., Fujimori, S., Strefler, J., Hasegawa, T., Marangoni, G., Krey, V., Kriegler, E., Riahi, K., van Vuuren, D. P., Doelman, J., Drouet, L., Edmonds, J., Fricko, O., Harmsen, M., Havlik, P., Humpenöder, F., Stehfest, E., and Tavoni, M.: Scenarios towards limiting global mean temperature increase below 1.5 °C, *Nature Climate Change*, 8, 325-332, 10.1038/s41558-018-0091-3, 2018.

- 15 Saito, M.: Charcoal as a micro-habitat for VA mycorrhizal fungi, and its practical implication, *Agriculture, Ecosystems & Environment*, 29, 341-344, 10.1016/0167-8809(90)90298-r, 1990.

Sarbas, B.: The GEOROC Database as Part of a Growing Geoinformatics Network, *Geoinformatics 2008—Data to Knowledge*, Potsdam, 2008.

Schuiling, R. D., and Krijgsman, P.: Enhanced weathering: An effective and cheap tool to sequester CO₂, *Climatic Change*, 74, 349-354, DOI 10.1007/s10584-005-3485-y, 2006.

- 20 Schuiling, R. D.: Farming nickel from non-ore deposits, combined with CO₂ sequestration, *Natural Science*, 05, 445-448, 10.4236/ns.2013.54057, 2013.

Schwartzman, D. W., and Volk, T.: Biotic Enhancement of Weathering and the Habitability of Earth, *Nature*, 340, 457-460, 10.1038/340457a0, 1989.

- 25 Semhi, K., Suchet, P. A., Clauer, N., and Probst, J. L.: Impact of nitrogen fertilizers on the natural weathering-erosion processes and fluvial transport in the Garonne basin, *Applied Geochemistry*, 15, 865-878, Doi 10.1016/S0883-2927(99)00076-1, 2000.

Smith, M. R., and Myers, S. S.: Impact of anthropogenic CO₂ emissions on global human nutrition, *Nature Climate Change*, 8, 834-839, 10.1038/s41558-018-0253-3, 2018.

Strefler, J., Amann, T., Bauer, N., Kriegler, E., and Hartmann, J.: Potential and costs of carbon dioxide removal by enhanced weathering of rocks, *Environmental Research Letters*, 13, 10.1088/1748-9326/aaa9c4, 2018.

- 30 Tack, F. M., Callewaert, O. W. J. J., and Verloo, M. G.: Metal solubility as a function of pH in a contaminated, dredged sediment affected by oxidation, *Environmental Pollution*, 91, 199-208, 10.1016/0269-7491(95)00049-6, 1996.

Taylor, L. L., Quirk, J., Thorley, R. M. S., Kharecha, P. A., Hansen, J., Ridgwell, A., Lomas, M. R., Banwart, S. A., and Beerling, D. J.: Enhanced weathering strategies for stabilizing climate and averting ocean acidification, *Nature Climate Change*, 6, 402-406, 10.1038/nclimate2882, 2015.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

hat gelöscht: ¶

Tilman, D., Socolow, R., Foley, J. A., Hill, J., Larson, E., Lynd, L., Pacala, S., Reilly, J., Searchinger, T., Somerville, C., and Williams, R.: Beneficial Biofuels—The Food, Energy, and Environment Trilemma, *Science*, 325, 270-271, 10.1126/science.1177970, 2009.

5 Topoliantz, S., and Ponge, J.-F.: Charcoal consumption and casting activity by *Pontoscolex corethrurus* (Glossoscolecidae), *Applied Soil Ecology*, 28, 217-224, 10.1016/j.apsoil.2004.08.003, 2005.

[Tropek, R., Kadlec, T., Karesova, P., Spitzer, L., Kocarek, P., Malenovsky, I., Banar, P., Tuf, I. H., Hejda, M., and Konvicka, M.: Spontaneous succession in limestone quarries as an effective restoration tool for endangered arthropods and plants, *Journal of Applied Ecology*, 47, 139-147, 10.1111/j.1365-2664.2009.01746.x, 2010.](#)

10 Uroz, S., Calvaruso, C., Turpault, M.-P., and Frey-Klett, P.: Mineral weathering by bacteria: ecology, actors and mechanisms, *Trends in Microbiology*, 17, 378-387, 10.1016/j.tim.2009.05.004, 2009.

van Straaten, P.: *Rocks for Crops: Agrominerals of sub-Saharan Africa*, ICRAF, Nairobi, Kenya, 338 pp., 2002.

Van Straaten, P.: Farming with rocks and minerals: challenges and opportunities, *Anais da Academia Brasileira de Ciencias*, 78, 731-747, 2006.

15 Warnock, D. D., Lehmann, J., Kuyper, T. W., and Rillig, M. C.: Mycorrhizal responses to biochar in soil – concepts and mechanisms, *Plant and Soil*, 300, 9-20, 10.1007/s11104-007-9391-5, 2007.

West, T. O., and McBride, A. C.: The contribution of agricultural lime to carbon dioxide emissions in the United States: dissolution, transport, and net emissions, *Agriculture, Ecosystems & Environment*, 108, 145-154, 10.1016/j.agee.2005.01.002, 2005.

20 White, J. G., and Zasoski, R. J.: Mapping soil micronutrients, *Field Crops Research*, 60, 11-26, 10.1016/s0378-4290(98)00130-0, 1999.

Xu, C. Y., Hosseini-Bai, S., Hao, Y., Rachaputi, R. C., Wang, H., Xu, Z., and Wallace, H.: Effect of biochar amendment on yield and photosynthesis of peanut on two types of soils, *Environ Sci Pollut Res Int*, 22, 6112-6125, 10.1007/s11356-014-3820-9, 2015.

25 Zhang, G.-l., Liu, F., and Song, X.-d.: Recent progress and future prospect of digital soil mapping: A review, *Journal of Integrative Agriculture*, 16, 2871-2885, 10.1016/s2095-3119(17)61762-3, 2017.

Zomer, R. J., Trabucco, A., Bossio, D. A., and Verchot, L. V.: Climate change mitigation: A spatial analysis of global land suitability for clean development mechanism afforestation and reforestation, *Agriculture, Ecosystems & Environment*, 126, 67-80, 10.1016/j.agee.2008.01.014, 2008.

hat formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm, Abstand Nach: 12 Pt.

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm

hat gelöscht: ¶

Seite 1: [1] Formatvorlagendefinition	revised	01.05.19 20:33:00
EndNote Bibliography: Englisch (Vereinigtes Königreich), Zeilenabstand: einfach		
Seite 1: [2] hat formatiert	revised	01.05.19 20:33:00
Englisch (Vereinigtes Königreich)		
Seite 1: [2] hat formatiert	revised	01.05.19 20:33:00
Englisch (Vereinigtes Königreich)		
Seite 1: [3] hat formatiert	revised	01.05.19 20:33:00
Englisch (Vereinigtes Königreich)		
Seite 1: [3] hat formatiert	revised	01.05.19 20:33:00
Englisch (Vereinigtes Königreich)		
Seite 1: [4] hat formatiert	revised	01.05.19 20:33:00
Englisch (Vereinigtes Königreich)		
Seite 1: [4] hat formatiert	revised	01.05.19 20:33:00
Englisch (Vereinigtes Königreich)		
Seite 10: [5] hat gelöscht	revised	01.05.19 20:33:00