

## POINT BY POINT RESPONSE TO THE REVIEWERS' COMMENTS

### REFEREE #1

**Comment 9.** The reply does not address the point made by the referee: is there robust evidence that charcoal affects the background albedo during a harvest decades after a fire? The referee seems to question this hypothesis because within days to weeks after a harvest the residual charcoal may be covered by snow, litter, herbs, mosses and/or ferns.

#### *Response to comment 9:*

Ground vegetation in recently disturbed sites in the study area is characteristically sparse (see photo below of a stand one-year post harvest). In older stands (45+ years post-disturbance) typical ground vegetation cover values, including non-vascular plants, are ~45% (Kumar et al., 2018).

Distinguishing charcoal from other dark material (e.g., leaf litter, organic matter, and fungi) is extremely difficult at a large spatial scale; we are aware of no existing practical methods that would permit such measurements. We did not quantify the percentage of surface charcoal present at the spatial scale measured by sensors in this study (~10-100 m<sup>2</sup>, depending on tower height). However, we did report our general field observations during data collection, in which we observed legacy charcoal as a ubiquitous component of soil surface material in both recent post-harvest and post-fire stands that could potentially affect surface albedo. Prior studies have also found that legacy charcoals can be visually observed on the forest floor and within the top 0.5 m soil many years (decades to 100+ years) following wildfire in boreal forests in Scandinavia (Ohlson et al., 2009) and Russia (Wallenius, 2002). Simulations from Tinker and Knight (2001) suggest that residuals from a fire event returning at a 100-year interval can cover ~25% of the forest floor.

In response to this criticism, we visually assessed our retained surface samples of intact forest floor (used for spectroscopic measurements: i.e., Fig. 3) for visually detectable charcoal fragments. Among post-fire stands, 100% (n = 30; surface area of each sample: 78.5 cm<sup>2</sup>) samples had visually detectable charcoal; among post-harvest stands, 70% (19 of n = 27 samples – with a similar

percentage in 4-year-old post-harvest stands, and 11-19-year-old stands) had visually detectable charcoal. These data are briefly reported in results in the revised manuscript, and confirm our general field observations that charcoal is a ubiquitous component of the soil surface in the study system.



*Stand at 1-year post harvest (Photography: Md Abdul Halim, 4 June 2014)*



*Examples of surface soil samples from harvested stands with visible charcoal fragments outlined.*

**Comment 19.** This confused the referee and is likely to confuse other readers as well. Rewrite this sentence (it requires adding a single word which would have been less work than the reply and my subsequent comment).

*Response to comment 19:*

DONE: thank you. We have added the word “basal” in the sentence which now reads “... proportion of deciduous-broadleaf basal area (%), ...”.

**Comment 20.** Add this information to the caption of the relevant figures. Readers should easily understand why different figures have different numbers of observations.

*Response to comment 20:*

DONE: thank you. We have added a clarification note on why Figure 6 has different number of observations in the figure caption.

**Comment 26.** Replace “... aspen may account for this effect” by “aspen may cause for this effect”.

*Response to comment 26:*

DONE: thank you.

**Comment 27.** Neither the reply nor the manuscript addresses why the albedo values seem 50% higher as noted by the referee.

“Another thing that caught my attention is the high albedo (approx. 0.3) of some stands in summertime (Figure 4c, d). Albedo values 0.2 (at solar noon) are rarely reported in boreal forests. Your values are approx. 50% higher than that. Is this because of the limited spectral range of the pyranometer?”

*Response to comment 27:*

Sorry, we missed this part of the comment in our prior response. As addressed in the first response, the cross-calibration results reported in the prior response indicate there was no detectable bias in albedo values estimated by the Si-based instruments used compared to a thermopile instrument.

The high summer albedo values observed correspond to a subset of post-fire stands with a high proportion (>60%) of deciduous species (mostly aspen): see Fig. 6f. There are only a few ground-based estimates of boreal forest albedo for deciduous-dominated stands, and one can expect variability from stand to stand. Although we did not quantify this effect, aspen in some of our sites had high levels of aspen serpentine leaf miner. Leaves affected by this insect turn nearly white in color (see photo below), and we strongly suspect this is responsible for the high summer albedo values observed in two of the stands.



*Aspen leaves impacted by aspen serpentine leaf miner (*Phyllocnistis populiella*)*

We have added a note on this pattern in the Results section.

**Comment 30.** Add this information to the caption of the relevant figures. Readers should easily understand why different figures have different numbers of observations.

***Response to comment 30:***

DONE: thank you. We have added a clarification note in figure caption on why the figure has different number of observations compared to other figures.

**Comment 31, 32, 35, 36.** This confused the referee and is likely to confuse other readers as well. Address this issue in the manuscript.

***Response to comment 31–32:***

DONE: we have rewritten the text in sections 3.3.3 and 2.5 (now 2.6) to enhance clarity.

***Response to comment 35:***

DONE: we have clarified “how we determined the effect” in the Discussion section (L335–337).

***Response to comment 36:***

DONE: we have edited the text to avoid confusion on energy balance in late spring (L350–355).

**Comment 33.** “Dramatic” is a subjective interpretation. Remove from the manuscript or define when and why a change in albedo should be considered dramatic.

***Response to comment 33:***

DONE: we have replaced the word “dramatic” with “significant”.

## REFEREE # 2

**Comment 1.** Based on the presented material the referee concluded that the manuscript lacks novel ideas or data. The authors could add a paragraph in which they show how their work goes beyond the work presented in the papers listed by the referee. Such a paragraph should avoid that other readers come to the same conclusion as the referee.

### *Response to Comment 1:*

We have responded on the data quality previously: in the revised manuscript we added results from a field calibration study that showed there was no statistically significant difference in albedo measured from Si-based pyranometers compared to that from thermopile pyranometers.

In this revision, we have included a paragraph (first paragraph of the Conclusions) that elaborates on the novel contributions of this study. In addition, we have added/modified a few sentences in the Introduction to clarify the novel points addressed.

## EDITOR

**Comment 1.** The discussion is still very much centered on stand age and does not even touch on the issue that age is not a driving process. The age-oriented approach is a shortcoming of the study and limits the value of the results and its applicability for data-model comparison. It is not because there is long tradition of studying age-related patterns in forestry that this tradition is well justified and should not be questioned.

### *Response to comment 1:*

We strongly disagree that the focus on age-dependent patterns in albedo is a “shortcoming” of the study: it is rather the main point of the study. We emphasize this in the paper not because there is a long tradition of studying age-related patterns (in forestry and other disciplines), but because assessing the larger-scale implications of ground-based measurements demand this approach. We do not see how it is possible to provide any integrated assessment of forest management impacts on albedo (or other critical climate feedbacks, such as forest carbon sequestration), without quantitatively assessing patterns through stand age. It is reasonable to describe albedo as (for example) a function of stand structure and composition (we do this explicitly in the paper). But such descriptions would only be sufficient (for purposes of modeling or integrated assessment over larger spatial or temporal scales) if one could strictly assume that the measured parameters describing stand structure and composition completely account for variation in albedo. We do not think such an assumption is warranted. Moreover, even if one did have a comprehensive physical process model that would allow prediction of albedo from, say, forest structure, understanding and modeling the implications of this from a forest management perspective (or in the context of a natural disturbance regime) would demand analyses based on temporal dynamics following disturbance.

We include the following sentences in the first part of the last paragraph of the Introduction to more explicitly justify our focus on temporal dynamics:

*“The temporal dynamics of stand albedo following disturbance regimes have critical implications to interactions between climate forcing, forest management, and disturbance regimes. For*

*example, if harvested stands converge in albedo with older stands within a few years (a small fraction of total rotation length), forest management is expected to have little impact on albedo at the landscape scale. Conversely, slow recovery in albedo or persistent effects of harvest compared to natural disturbance would indicate that the forest management regime fundamentally alters albedo-related climate feedbacks.”*

We have also added a sentence in the Discussion (sub-section 4.2) that touches upon this issue.

## **REFERENCES**

Kumar, P., Chen, H. Y. H., Thomas, S. C. and Shahi, C.: Epixylic vegetation abundance, diversity, and composition vary with coarse woody debris decay class and substrate species in boreal forest, *Can. J. For. Res.*, 48(4), 399–411, doi:10.1139/cjfr-2017-0283, 2018.

Ohlson, M., Dahlberg, B., Økland, T., Brown, K. J. and Halvorsen, R.: The charcoal carbon pool in boreal forest soils, *Nature Geosci*, 2(10), 692–695, doi:10.1038/ngeo617, 2009.

Tinker, D. B. and Knight, D. H.: Temporal and spatial dynamics of coarse woody debris in harvested and unharvested lodgepole pine forests, *Ecological Modelling*, 141(1), 125–149, doi:10.1016/S0304-3800(01)00269-1, 2001.

Wallenius, T.: Forest age distribution and traces of past fires in a natural boreal landscape dominated by *Picea abies*, *Silva Fenn.*, 36(1), doi:10.14214/sf.558, 2002.

# Stand age and species composition effects on surface albedo in a mixedwood boreal forest

Mohammad Abdul Halim<sup>1,2</sup>, Han Y. H. Chen<sup>3</sup>, Sean C. Thomas<sup>1</sup>

5 <sup>1</sup>University of Toronto, Faculty of Forestry, 33 Willcocks Street, M5S 3B3, ON, Canada

<sup>2</sup>Shahjalal University of Science and Technology, Dept. of Forestry and Environmental Science, Sylhet-3114, Bangladesh

<sup>3</sup>Lakehead University, Natural Resources Management, 955 Oliver Road, Thunder Bay, P7B 5E1, ON, Canada

*Correspondence to:* Mohammad Abdul Halim (abdul.halim@mail.utoronto.ca)

10

**Abstract.** Surface albedo is one of the most important processes governing climate forcing in the boreal forest and is directly affected by management activities such as harvesting and natural disturbances such as forest fires. Empirical data on the effects of these disturbances on boreal forest albedo are sparse. We conducted ground-based measurements of surface albedo from a series of instrument towers over four years in a replicated chronosequence of mixedwood boreal forest sites differing in stand age (to 19 years since disturbance) in both post-harvest and post-fire stands. We investigated the effects of stand age, canopy height, tree species composition, and ground vegetation cover on surface albedo through stand development. Our results indicate that winter and spring albedo values were 63 and 24 % higher, respectively, in post-harvest stands than in post-fire stands. Summer and fall albedo values were similar between disturbance types, with summer albedo showing a transient peak at ~10 years stand age. The proportion of deciduous broadleaf species showed a strong positive relationship with seasonal averages of albedo in both post-harvest and post-fire stands. Given that stand composition in mixedwood boreal forests generally shows a gradual replacement of deciduous trees by conifers, our results suggest that successional changes in species composition are likely a key driver of age-related patterns in albedo. Our findings also suggest the efficacy of increasing the proportion of deciduous broadleaf species as a silvicultural option for climate-friendly management of the boreal forest.

25

## Keywords

Stand age, species composition, canopy height, ground vegetation cover, harvesting, forest fire, succession, surface albedo, boreal forest.

## 1 Introduction

30

Surface albedo, the fraction of incoming solar energy reflected from the surface in all directions, is one of the most important biophysical factors affecting both local and global climates. In boreal forest, the magnitude of albedo-related forcing on climate is even more important than in other ecosystems because of snow-related feedbacks, low sensible heat flux, and the relative stability of the atmospheric temperature profile due to weak latent-heat-driven convection (Bright et al., 2015a; Hansen et al., 2005). Even though albedo is increasingly used as an important state variable in climate models (Brown and Caldeira, 2017; Bala et al., 2007; Betts, 2000), forest disturbance effects on net radiative forcing due to local albedo changes and related feedbacks with regional/global mean surface temperature remain highly uncertain (Bright et al., 2015a; Lee et al.,

35

Deleted: (0-19 years)

Deleted: (7-19 years)

Deleted: Winter and summer albedos saturated at roughly 50 years stand age in both post-harvest and post-fire stands.

2011). Harvest and fire suppression may differ substantially in their effects on albedo, but empirical data on albedo responses to disturbance type remain particularly sparse.

45 Following disturbance events, albedo of boreal forests is expected to change with stand age due to changing surface properties, and forest structure and composition. Age-related stand structural attributes (e.g., tree species composition, leaf area index [LAI], canopy height, and ground vegetation cover) can substantially influence surface albedo of a stand throughout the year. Studies have generally found higher albedos in young stands than in mature stands in the boreal forest (Bright et al., 2015a; Kuusinen et al., 2014; Amiro et al., 2006b), but the dynamic patterns with stand age remain unclear. The main study using ground-based measurements fit functions describing a linear decrease in summer albedo, and an exponential decrease in winter albedo, with stand age (Amiro et al., 2006b); however, in both cases the variability in young stands (<25 years) was much greater than in older stands and poorly described by fitted models. Early in stand development boreal mixedwood forests are commonly dominated by deciduous broadleaf species (Madoui et al., 2015; Brassard and Chen, 2010; Johnstone et al., 2010), which have higher leaf and canopy reflectance than conifers (Lukeš et al., 2013a; Linacre, 2003), contributing to high summer albedo in young stands (Lukeš et al., 2013b; Betts and Ball, 1997). These deciduous species shed leaves in the winter, which increases canopy openness (lowers LAI) and allows snow albedo to dominate, contributing to the high winter albedo in young stands. Available data suggest that at this stage both LAI and ground vegetation cover usually increase with stand age, depending on site quality and silvicultural practices (Amiro et al., 2006b; Uotila and Kouki, 2005). Low LAI can increase canopy background reflectance both in snow-covered and snow-free conditions, and thus can contribute to the high albedos in young stands (Amiro et al., 2006b). LAI effects on albedo in young stands may be highly modulated by ground vegetation cover in the summer, but probably not much in the winter as ground vegetation is generally leafless or covered with snow (Kuusinen et al., 2015; Lukeš et al., 2013b; Betts and Ball, 1997). In conjunction with other factors, surface albedo tends to decrease with increasing canopy height (Hovi et al., 2016; Linacre, 2003). In the later stages of stand development, albedo is expected to saturate non-linearly as conifers dominate the stand and canopy cover and stand attributes change gradually, but data describing this pattern remain sparse (Amiro et al., 2006b).

65 Harvesting and fire are the major stand-replacing disturbances in the boreal forest (Brassard et al., 2008). These disturbances may differentially affect surface albedo of post-disturbance stands in complex ways by altering ground surface spectral properties, species composition, and stand structure (Lukeš et al., 2013b; Liu et al., 2005), but field data directly addressing this issue are essentially limited to a single study in Europe (Kuusinen et al., 2016). Structure and composition of post-fire stands are generally more heterogeneous than post-harvest stands; for example, post-fire stands are more likely to show a bimodal vertical structure and a mixture of conifer and hardwood species during early stand development stages (Brassard and Chen, 2010; Chen et al., 2009). Charcoal residues may also strongly reduce albedo in snow-free conditions in the first years following fire disturbances (Amiro et al., 2006b). Both charcoal effects and stand heterogeneity might be expected to reduce surface albedo in post-fire stands relative to post-harvest stands. However, the magnitude of this difference in surface albedo might be less than expected due to the presence of legacy charcoals from historical fires in post-harvest stands (Hart and Luckai, 2013). Immediately after harvesting, the albedo of a post-harvest stand can also be reduced because of the presence of coarse woody debris (CWD) and high soil moisture content (Linacre, 2003). In the years following a disturbance event, CWD might be expected to further reduce albedo by becoming darker in color due to decomposition processes (Brassard and Chen, 2008) and plant colonization (Kumar et al., 2018).

**Deleted:** , and it has been suggested that albedo stabilizes ~ 25 years after a disturbance event

**Deleted:** (during the 'stem exclusion' phase in the ecological succession trajectory)

85 Despite the important roles of stand age, and stand structure and composition as determinants of boreal forest albedo, field  
measurements are scarce (Kuusinen et al., 2014) and particularly limited for early stand ages that show high variability in  
surface properties (Bright et al., 2013). This has contributed to poorly constrained estimates of the local albedo changes on  
net global radiative forcing (Bright et al., 2015a). Although some recent studies (e.g., Luysaert et al., 2018; Naudts et al.,  
90 2016) have incorporated vegetation structure and composition in albedo estimation for land surface models, scarcity of field  
measurements is still a challenge for proper attribution of boreal forest albedo in climate models (Li et al., 2016; Thackeray  
et al., 2019). Thus, to estimate the net change in surface temperature as a function of albedo change from deforestation in  
boreal forests, a number of climate models (e.g., Bala et al. 2007, Betts 2000) have used a 'biome replacement' approach  
(replacing boreal forests with grassland or agricultural land cover types) and approximated boreal forests' albedo as a single  
value from mature stands (~60-year old). Early stand dynamics is reported to determine which mechanism, albedo vs. carbon  
95 storage, dominates the net forcing for the boreal forest (Kirschbaum et al., 2011). Simplifications in climate models that do  
not explicitly consider stand age and successional effects on albedo will likely result in strongly biased estimations of boreal  
forests' albedo over the rotation (harvesting/fire) period (Bright et al., 2018).

Deleted: Such s

100 Temporal dynamics of stand albedo following disturbance events have critical implications to interactions between climate  
forcing, forest management, and disturbance regimes. For example, if harvested stands converge in albedo with older stands  
within a few years (a small fraction of total rotation length), forest management is expected to have little impact on albedo at  
the landscape scale. Conversely, slow recovery in albedo or persistent effects of harvest compared to natural disturbance  
would indicate that the forest management regime fundamentally alters albedo-related climate feedbacks. Better  
105 understanding of post-disturbance patterns and of the mechanisms that account for variation in albedo will not only enhance  
global climate models (e.g., by improving the land-surface model: Bright et al., 2018), but also help to design climate-friendly  
silvicultural practices (Astrup et al., 2018; Matthies and Valsta, 2016; Bright et al., 2015a). In the present study, we set up  
micrometeorological towers with pyranometers in a replicated chronosequence of post-harvest and post-fire sites to study  
stand age, disturbance type, and species composition effects on albedo in a mixedwood boreal forest in northwestern Ontario,  
Canada. We focused on the early stand development (0-19 years post harvest), where dynamics is expected to be most rapid and  
110 where ground-based data from boreal forests is most sparse. We hypothesized: (1) that post-fire stands would show lower  
albedo values than post-harvest stands as a consequence of stand composition, legacy structures, and fire residues; (2) that all  
stands would approach albedo values similar to mature stands within 20-25 years, soon after crown closure; and (3) that stands  
with higher dominance of deciduous broadleaf species would show higher albedo than conifer-dominated stands, with this  
effect being most pronounced under snow-covered conditions.

Deleted: Given the complex nature and limited understanding of underlying processes (Lukeš et al., 2013b), it is important to conduct field studies to better quantify forest albedo in relation to stand age and disturbance type in boreal forests. Deeper

Deleted: local

Deleted: of

Deleted: ~

## 115 2 Materials and Methods

### 2.1 Study area

The study was conducted in the boreal forest of the Lake Nipigon region (49.55° N and 89.5° W), Ontario, Canada, approximately 200 km north of Thunder Bay. A series of circular (10-m radius) chronosequence plots were established in the post-harvest (full-tree harvest) and post-fire stands in the study area. Three plots were set up in each of three cutblocks (in  
120 separate stands) harvested in 1998, 2006, and 2013. Selected stands were at least 5 ha in size, and plots were established at  
least 100 m from any older or taller stand to avoid edge effects. Recent (2013) post-fire stands were not present, so we set up

three plots only in post-fire stands dating from 1998 and 2006 fire events (Fig. 1). Replicate stands were spatially interspersed to the extent feasible. For each of the 15 plots, albedo and stand attributes (stand age, percentage of deciduous broadleaf species, canopy height, and percentage of ground vegetation cover) were measured from July 2013 to June 2017.

135 The mesic mixedwood study area is dominated by jack pine (*Pinus banksiana* Lamb.), black spruce (*Picea mariana* (Mill.)  
BSP), white spruce (*P. glauca* (Moench) Voss), trembling aspen (*Populus tremuloides* Michx.), eastern white cedar (*Thuja*  
*occidentalis* L.), balsam fir (*Abies balsamea* (L.) Mill.), and paper birch (*Betula papyrifera* Marsh.) (Chen and Popadiouk,  
2002). The management regime in the region is based on clearcut silviculture modified to include live tree retention in  
140 harvested stands (OMNRF, 2015); typical rotation lengths are 80 years (Colombo et al., 2005). In study plots over the study  
period canopy height ranged from 0–7.7 m, ground vegetation cover ranged from 1.8–96.7 %, LAI from 0–2.1, and the  
proportion of deciduous broadleaf basal area from 10.6–100 %, and stand density from 0–11556 stem/ha (Table 1). The study  
area has an average elevation of 416 m a.s.l. The soil is a moderately deep brunisol (coarse-loamy texture) with 1–15 cm thick  
organic layer (i.e., the total litter, fermented, and humic [LFH] layers). The area remains snow covered for 5–6 months with  
an average snow depth of ~10 cm (Environment Canada, 2018; Sims et al., 1997) and the mean annual air temperature of the  
145 study plots was -1.1 °C (Halim and Thomas, 2018).

## 2.2 Experimental setup

In the center of each circular plot a pair of upward- and downward-facing pyranometers (Silicon [Si] pyranometer; Onset,  
Massachusetts, USA; measurement range 0–1280 Wm<sup>-2</sup> over a spectral range of 300–1100 nm, accuracy ± 5 %, resolution  
150 1.25 Wm<sup>-2</sup>) were set up on a mast 3.5 m above the canopy (above the ground for 2013 post-harvest stands) to measure incident  
and reflected solar radiation every 10 minutes. The plot and tower locations were selected to avoid trees from surrounding  
stands falling within the footprints of the pyranometers or blocking incoming solar radiation. Instrument masts consisted of  
extendible galvanized steel poles and were set in concrete bases and guyed to mitigate instrument sway. At least once a year  
pyranometer heads were cleaned and realigned to make sure they were normal to the ground. Average daily albedo was  
155 calculated as the ratio of daily total incident and reflected radiation for each plot. The average daily albedo was used to  
calculate average monthly albedo, which was finally used to calculate mean seasonal albedo for each year in each plot.

Quality control for the irradiance and reflected solar radiation measurements was conducted following guidelines of the World  
Meteorological Organization (WMO). Any unusually high/low values were replaced by interpolated values by taking the  
160 average of preceding and subsequent measurements. Daily total irradiance data were compared against the WMO-provided  
maximum possible daily sums of clear-sky irradiance for 50°N latitudes (WMO, 1987, p.26). If the measured daily total  
irradiance was higher than the maximum possible value, we excluded the measurements for that day. For reflected solar  
radiation, if the daily total of reflected solar radiation was higher than the daily total irradiance, we also excluded the  
measurements for that day. In addition, we excluded measurements for any snowy day; snowfall was detected using data from  
165 the closest available weather station (Environment Canada, 2018).

In addition to albedo, winter (December–February)/spring (March–May) and summer (June–August)/fall (September–  
November) proportion of deciduous broadleaf basal area (%), canopy height (m), and ground vegetation cover (%) were  
measured every year in late October and early July, respectively, in each plot. The proportion of deciduous broadleaf species

Deleted: ranged

Deleted: ranged

Deleted: range

Deleted: Brunisol

Deleted: the

Deleted:

(%) were determined for trees with diameter at breast height  $\geq 5$  cm and height  $> 1$  m. Canopy height was determined as the mean height of all trees sampled; the young stands sampled were at stages of development prior to and just after canopy closure, so essentially all trees were “canopy dominants”. The proportion of deciduous broadleaf species of a plot was calculated as the ratio of basal area of the deciduous species to the total basal area of the plot. In each plot, four 1-m<sup>2</sup> subplots were set up and percent ground vegetation cover was determined visually (Kumar et al., 2018). Stand age was determined as the time (year) since the last disturbance (fire/harvesting) for each plot. Fire maps (from the Ontario Ministry of Natural Resources, Canada) and forest management plans (from Resolute Forest Products, Canada) were used to verify type and year of disturbances.

### 2.3 Sources of secondary data

Since we did not have recent post-fire stands (0–6-year old) in the study area, we used secondary albedo data from studies in post-fire boreal forests with similar stand characteristics in Saskatchewan and Manitoba (Canada) (Fig. 1). We also used secondary albedo data for old stands ( $> 70$  years) from these sites along with primary data to develop regression models for both post-fire and post-harvest stands. Here we assumed that at this *late* stage of stand development, there is negligible difference in stand attributes (e.g., species composition, height, LAI) between post-fire and post-harvest stands (Moussaoui et al., 2016). We did not use satellite-based albedo data as secondary sources as they tend to diverge from field measurements depending on a number of factors including stand age, latitude, and cloud cover effects (Halim and Thomas, 2017; Bright et al., 2015b).

Data for Saskatchewan sites were retrieved from Amiro et al. (2006a), and for Manitoba sites from Amiro et al., (2006b) by digitizing data points from relevant figures using the WebPlotDigitizer software (Rohatgi, 2018). These stands were dominated by jack pine and black spruce with some intermixing of trembling aspen. All post-fire sites (including this study) had severe fires that completely killed previous vegetation. There were a few burned snags in the Saskatchewan and Manitoba sites and none in the present study sites. These areas remain snow covered for ~6 months with average snow depths of 10–15 cm (Environment Canada, 2018). Pyranometers were located in Saskatchewan sites at 18–20 m, and in Manitoba sites at 6 m heights. There was no detailed information on how proportions of broadleaf deciduous species were calculated for these sites; however, we assumed they were basal-area based. A detailed description of the study areas and methods can be found in the respective articles.

### 2.4 Accuracy assessment of albedo measurements

To test the relative accuracy of albedo measurements from Si-based pyranometers (Onset Computers' Hobo, used in this study) in comparison to thermopile pyranometers (Kipp and Zonen's CNR1, used in the studies providing secondary data), we conducted a field calibration study over nine days under variable sky and ground conditions (see Supplementary Materials). Results from this study showed a very close agreement between the measurements of Hobo and CNR1 pyranometers (Fig. S1). The difference (CNR1 – Hobo) in daily albedo over the study period ranged from –0.0601 to 0.064, and the mean difference in daily albedo was 0.0028 ( $\pm 0.031$ ). The mean difference was negligible and the range in differences was well within the previously reported error ranges (~5–7 %) for similar pyranometers (Myers, 2010; Stroeve et al., 2005). We also did not observe any detectable pattern in deviations between the pyranometers under different sky and ground conditions. We therefore concluded that albedo measurements from Si-based Hobo and thermopile-

Deleted: n

Deleted: s

Deleted: n

Deleted: n

220 based CNR1 pyranometers are closely comparable, and corrections to the Si-based albedo estimates presented are not warranted.

## 2.5 Measurements of ground surface reflectance

225 To examine effects of disturbance type on ground surface reflectance, three soil samples (top 10 cm including LFH layer, surface area 78.5 cm<sup>2</sup>) from each plot were collected in fall 2017 to measure the ground surface reflectance. Samples were all collected within a two-day precipitation-free period, and were brought to the lab in airtight packaging without disturbing the top surface. Surfaces of these samples were visually assessed for presence of visible charcoal fragments. A spectrometer (SD 230 2000; Ocean Optics, Florida, USA; measurement spectral range 338.7–1001.8 nm) equipped with an integrating sphere was used to measure the directional-hemispherical reflectance factor of the top surface of the soil samples. As there were no recent post-fire stands in the study area, we collected charcoal samples (of twigs, branches, barks, and stems) from the forest floor of a jack pine dominated post-fire (fire occurred in 2011) stand in summer 2015 from near the Musselwhite mine (52.61° N and 90.37° W), Ontario, Canada. Every sample was measured ten times in ten different locations (each 0.84 cm<sup>2</sup> in area), and each measurement was performed by scanning 10 times (with Boxcar width 5 [spatial averaging of 5 pixels] and 100 millisecond integration time) to get an average reflectance for each location of a sample. Details of the spectrometer and integrating sphere used can be found in the Materials and Methods section of Baltzer and Thomas (2005). Forest floor 235 reflectance values from the Musselwhite stand (4-year old) were compared to soil sample reflectance values from recent (2013) post-harvest stands (4-year old). For older stands (1998 and 2006 post-harvest and post-fire stands), soil sample reflectance data were compared using samples from the main study plots.

## 2.6 Data analysis

240 Robust t-tests (Wilcox, 2016) were used to compare mean differences in ground surface reflectance (in visible [400–700 nm] and near-infrared [ $> 700$ –1000 nm] spectral bands) and seasonally averaged albedo between post-harvest and post-fire stands. Mean seasonal albedo values of post-harvest and post-fire stands were also compared using analysis of covariance (ANCOVA) controlling for the effects of stand age as a covariate. Secondary albedo data for 0–6-year-old post-fire stands were only available for winter and summer seasons. Therefore, in the t-tests (and in ANCOVA) for winter and summer albedo, 245 data from 0–19-year old post-harvest and post-fire stands were used. For spring and fall, albedo data from recent (0–6-year-old) post-harvest stands were omitted (since there were no data from post-fire stands for these seasons), and data from 7–19-year-old stands were used to make the comparisons unbiased. Secondary data from old stands ( $> 70$  years) were not used in the t-tests/ANCOVA. These analyses treat seasonally averaged albedo values from the same stands as independent. We also conducted parallel analyses using linear mixed models that included plot as a random variable; in all cases, the random effect 250 was not significant, and thus only the simpler linear model results are presented.

Generalized linear models (GLMs) with the log-linked gaussian family (additive-observation-error model with constant variance) were found to be the best fitted to model seasonal albedo as a function of stand attributes (stand age, proportion of deciduous broadleaf species, canopy height, ground vegetation cover, and their interactions) for both post-harvest and post- 255 fire stands. Best models were chosen using an AIC-based stepwise algorithm. Asymptotic chi-square statistics based on deviance were calculated for each best-fit model to test if the model was significantly better than its counterpart null model. We could not use a GLM to predict fall albedo because some stand attributes were only nonlinearly (double exponentially)

Deleted: Analysis

Deleted: Covariance

Deleted: In fall months,

related to albedo. If we included these nonlinearly-related stand attributes with other attributes in a GLM, the model structure became very complex (a mixture of nonlinear and linear families) and defining an appropriate GLM family became a statistical challenge. To avoid modeling complexity, for each of these nonlinearly-related fall attributes a separate nonlinear model was fitted, and for other attributes GLMs with identity-linked gaussian family were found to be the most suitable. The  $\Delta$ AIC for each best-fit model is calculated as its AIC difference with the corresponding null model (AIC of the best-fit model – AIC of the corresponding null model). Sample-size corrected AIC values were used in all cases.

Data were analyzed using the R platform (R Core Team, 2018) and graphs were prepared using the ‘ggplot2’ package (Wickham, 2016). Robust t-tests were done by 10,000 bootstrapped samples considering mean as an estimator for group comparison, and implemented by the *pb2gen* function of the WRS2 R-package (Mair and Wilcox, 2018). Adjusted  $R^2$  values for GLMs were calculated using the *rsq* function of the R-package ‘rsq’ (Zhang, 2018).

### 3 Results

#### 3.1 Seasonal albedo in post-harvest and post-fire stands

Albedo differences between post-harvest and post-fire stands varied among seasons. Albedo values in periods of the year with appreciable snow cover were significantly higher in post-harvest stands than in post-fire stands (for winter: 0.56 vs. 0.34,  $p < 0.01$ ; for spring: 0.32 vs. 0.24,  $p = 0.11$ ). Summer albedo values were also marginally higher in post-harvest stands ( $p = 0.24$ ), and fall albedos were similar between disturbance types ( $p = 0.73$ ) (Fig. 2). Considering stand age as a covariate, ANCOVA results also indicate higher albedo of post-harvest stands in winter ( $p = 0.02$ ), spring ( $p = 0.15$ ), summer ( $p = 0.04$ ), and similar in fall ( $p = 0.77$ ) compared to post-fire stands. Data also suggest higher variability in albedo in post-harvest stands than in post-fire stands (Fig. 2).

#### 3.2 Ground surface reflectance in post-harvest and post-fire stands

Surface charcoal fragments were visually observed in all post-fire soil core samples, and in 70% of post-harvest samples. Specular-included reflectance measurements of ground surface samples suggest that differences in ground surface characteristics contribute to overall surface albedo in the study sites. Summer ground surface reflectance was generally higher in old stands (Fig. 3b) than in young stands (Fig. 3a) particularly in the 600–1000 nm range. Young (4-year old) post-harvest stands showed significantly lower mean ground reflectance values (74.3 %,  $p < 0.01$ ) in the visual spectrum (400–700 nm) and higher (32.3 %,  $p < 0.01$ ) in the near-infrared spectrum ( $> 700$ –1000 nm) than those of young post-fire stands (Fig. 3a). Older (11- and 19-year old) post-harvest stands however showed higher mean ground reflectance in both visible (31.7%,  $p < 0.01$ ) and near-infrared (4.6%,  $p < 0.01$ ) spectra compared to post-fire stands (Fig. 3b).

#### 3.3 Seasonal albedo in relation to stand attributes in post-harvest and post-fire stands

##### 3.3.1 Winter albedo in post-harvest and post-fire stands

Results from the best-fit GLM ( $p < 0.01$ , adj.  $R^2 = 0.97$ ) for post-harvest stands indicated that stand age, proportion of deciduous broadleaf species, canopy height, and interactions among these variables were significant predictors of winter albedo (Table 2). Stand age was related to winter albedo via an exponential decay model with a horizontal asymptote ( $\Delta$ AIC

= - 25.5), and all estimated model parameters were significant (for 0.19 and 0.55:  $p < 0.01$ ; for - 0.06:  $p < 0.05$ ) (Fig. 4a). The proportion of deciduous broadleaf species (Fig. 6a) and canopy height (Fig. 7a) were also related to winter albedo via negative exponential models with horizontal asymptotes ( $\Delta AIC = - 6.7$  and  $- 100.4$ , respectively), and all estimated parameters for both models were significant ( $p < 0.01$ ).

300

For post-fire stands the best-fit GLM ( $p < 0.01$ , adj.  $R^2 = 0.75$ ) indicated that stand age and proportion of deciduous broadleaf species were significant predictors of winter albedo (Table 2). Stand age was related to winter albedo via an exponential decay model with a horizontal asymptote ( $\Delta AIC = - 38.5$ ), and all estimated model parameters were significant ( $p < 0.01$ ) (Fig. 4b). Proportion of deciduous broadleaf species was related to winter albedo via a negative exponential model with horizontal asymptote ( $\Delta AIC = - 16.3$ ), and all estimated model parameters were significant (for - 0.27:  $p < 0.06$ ; for 1.02 and 0.45:  $p < 0.01$ ) (Fig. 6b).

305

### 3.3.2 Spring albedo in post-harvest and post-fire stands

For post-harvest stands the best-fit GLM ( $p < 0.01$ , adj.  $R^2 = 0.99$ ) indicated that stand age, proportion of deciduous broadleaf species, height, and the interaction of stand age and proportion of deciduous broadleaf species were significant predictors of spring albedo (Table 2). Stand age (Fig. 5a) and canopy height (Fig. 7c) were related to spring albedo via exponential decay models with horizontal asymptotes ( $\Delta AIC = - 15.1$  and  $- 31.2$ , respectively). Estimated parameters of stand age-albedo (for 0.26:  $p < 0.01$ ; for 0.72 and - 0.72:  $p < 0.05$ ) and canopy height-albedo (for 0.16 and 0.33:  $p < 0.01$ ; for - 1.84:  $p = 0.07$ ) models were likewise significant. The proportion of deciduous broadleaf species was related to spring albedo via a negative exponential model ( $\Delta AIC = - 6.72$ ), and all estimated parameters were significant ( $p < 0.01$ ) (Fig. 6c).

310

315

The best-fit GLM ( $p < 0.01$ , adj.  $R^2 = 0.99$ ) for post-fire stands indicated that stand age and proportion of deciduous broadleaf species were the only significant predictors of spring albedo (Table 2). Stand age (Fig. 5b) and proportion of deciduous broadleaf species (Fig. 6d) were related to spring post-fire stand albedo via exponential negative growth models ( $\Delta AIC = - 7.0$  and  $- 7.5$ , respectively), and all estimated parameters for both models were significant ( $p < 0.01$ ).

320

### 3.3.3 Summer albedo in post-harvest and post-fire stands

The best-fit GLM ( $p < 0.01$ , adj.  $R^2 = 0.97$ ) for post-harvest stands indicated that stand age, proportion of deciduous broadleaf species, ground vegetation cover and its interaction with stand age and proportion of deciduous broadleaf species were significant predictors of summer albedo (Table 2). Stand age **alone (not with other stand attributes as in the GLM)** was related to summer albedo via a double exponential model ( $\Delta AIC = - 73.1$ ), and all estimated model parameters were significant ( $p < 0.01$ ) (Fig. 4c). The pattern described by this function indicates a sharp peak in albedo with a maximum at 10–15 years of stand age. Proportion of deciduous broadleaf species is related to summer albedo via a 3-parameter sigmoid model ( $\Delta AIC = - 48.6$ ), and all the estimated parameters were significant ( $p < 0.01$ ) (Fig. 6e). Ground vegetation cover was related to summer albedo via an exponential model with a Gumbel distribution without a horizontal asymptote ( $\Delta AIC = - 25.8$ ), and all estimated parameters were significant ( $p < 0.01$ ) (Fig. 8e).

325

330

For post-fire stands the best-fit GLM ( $p < 0.01$ , adj.  $R^2 = 0.95$ ) indicated that stand age, proportion of deciduous broadleaf species, canopy height, and their interactions with stand age were significant predictors of summer albedo (Table 2). Stand

335 age (Fig. 4d) and canopy height (Fig. 7f) were related to summer post-fire stand albedo via exponential models with Gumbel  
distributions with horizontal asymptotes ( $\Delta AIC = -49.3$  and  $-5.3$ , respectively). As in the case of post-harvest stands, peak  
albedo was found at  $\sim 10$ – $15$  years of stand age. All estimated parameters of stand age-albedo and canopy height-albedo  
models were significant ( $p < 0.01$ ). Proportion of deciduous broadleaf species was related to summer albedo via a negative  
exponential growth model ( $\Delta AIC = -6.8$ ), and all estimated model parameters were significant ( $p < 0.01$ ) (Fig. 6f). Two  
340 instances of particularly high summer albedo measurements ( $> 0.2$ ) were found in aspen-dominated stands affected by damage  
from aspen serpentine leaf miner (*Phyllocnistis populiella*).

### 3.3.4 Fall albedo in post-harvest and post-fire stands

The best-fit GLM ( $p < 0.01$ , adj.  $R^2 = 0.94$ ) for post-harvest stands indicated that stand age, canopy height, ground vegetation  
345 cover, and their interactions were significant predictors of fall albedo (Table 2). Proportion of deciduous broadleaf species  
was also an important predictor that was modelled separately via a double exponential model (and was not added to the GLM  
to avoid modelling complexities) ( $\Delta AIC = -0.9$ ); all estimated model parameters were significant (for 28.9 and 45.4:  $p <$   
 $0.05$ ; for 67.6:  $p < 0.01$ ) (Fig. 6g). Stand age (Fig. 5c) and ground vegetation cover (Fig. 8g) were related to albedo via  
exponential decay models with horizontal asymptotes ( $\Delta AIC = -36.8$  and  $-28.38$ , respectively), and all estimated parameters  
350 for both models were significant ( $p < 0.01$ ). Canopy height was also related to albedo via a negative exponential model ( $\Delta AIC$   
 $= -11.2$ ), and all estimated parameters were significant ( $p < 0.01$ ) (Fig. 7g).

To avoid modelling complexities, stand age and proportion of deciduous broadleaf species were fitted individually with fall  
albedo of post-fire stands (Table 2). Stand age was related to albedo via a double exponential model ( $\Delta AIC = -3.1$ ), and all  
355 estimated model parameters were significant ( $p < 0.01$ ) (Fig. 5d). Proportion of deciduous broadleaf species was generally  
related to albedo via a simple exponential model ( $\Delta AIC = -25.4$ ), and all estimated model parameters were significant ( $p <$   
 $0.01$ ) (Fig. 6h). In the case of fall albedo in post-harvest stands, there is an apparent decline in nearly pure stands (Fig. 6g),  
with a better fit of the double exponential model. We speculate that very dark post-senescence leaf litter of aspen may be the  
main cause for this effect.

Deleted: account

## 360 4 Discussion

Our results provide evidence for significant effects of disturbance type on the albedo of boreal forest systems, with post-  
harvest stands showing much higher albedo values in winter and spring months than post-fire stands. Stands of both  
disturbance types also showed strongly age-dependent patterns in albedo, with a transient peak in summer albedo at  $\sim 10$  years;  
however, analyses also suggest that later post-disturbance changes are more gradual than anticipated, with dynamics  
365 continuing for decades following stand closure. The proportion of deciduous species also had large effects on stand albedo –  
generally larger than stand age effects as indicated by overall lower residual standard errors of deciduous broadleaf species  
(%) regression models (Fig. 4–5 vs. 6). – and showing a positive response in all seasons and for both disturbance types.

Deleted: dramatic

Deleted: past

Deleted: , up to  $\sim 50$  years post-disturbance

Deleted: saturating

### 4.1 Albedo in post-harvest and post-fire stands

370 Mean albedo in post-harvest stands was significantly higher than in post-fire stands in winter and spring, marginally higher  
in summer, and similar in fall (Fig. 2). A similar pattern in albedo differences was also observed when the stand age effects  
on albedo were statistically controlled. The magnitude of differences in winter and spring values (0.22 and 0.08, or 63% and

34% increases relative to post-fire values) is large – comparable to albedo differences observed between biomes (Stephens et al., 2015). During snow-covered seasons (winter and spring), charcoal residues in post-fire stands are usually covered with snow, and thus stand structure and composition act as dominant drivers of albedo (Lyons et al., 2008; Amiro et al., 2006b; Liu et al., 2005). Deciduous broadleaf species made up 37.8 % of basal area in post-fire stands and 55.4 % in post-harvest stands: the higher percentage of dark conifer leaves is expected to result in lower winter/spring albedos in post-fire stands compared to post-harvest stands (Betts and Ball, 1997). However, immediately after a stand-replacing fire, the presence of black carbon (charcoal and soot) in the snow can reduce early winter albedo and possibly enhance spring snowmelt by absorbing solar radiation (Qian et al., 2009; Conway et al., 1996). During late spring when snow cover is shallow, it is also likely that charred branches and stems protrude through the snow and reduce albedo. Additionally, by the time of snowmelt, snow generally has accumulated particulate matter and has lower albedo compared to fresh snow (Conway et al., 1996). During this time of the year, latent heat flux from the melting snow is usually very high. Thus, from an energy balance perspective, it is important to note that albedo differences in late spring may be less important as turbulent and latent fluxes likely dominate (Conway et al., 1996).

Deleted: melting snow, due to the high incoming solar radiation, is usually dirty

Deleted: F

In snow-free seasons (summer and fall) the marginal differences in mean albedo between post-harvest and post-fire stands can partly be attributed to recovery of ground vegetation in post-fire stands (0–5 years old) compared to post-harvest stands (Bartels et al., 2016), and to the vegetation covering dark charcoals in older (> 5 year) post-fire stands (Randerson et al., 2006). Soon after a fire, the presence of early-successional plants (Johnstone et al., 2010) can increase surface albedo of post-fire stands because of their higher albedo relative to charcoal (Amiro et al., 2006b; Betts and Ball, 1997). This effect is expected to offset the albedo difference between post-harvest and post-fire stands. In the first year following disturbance events, we might expect lower snow-free albedo in post-fire stands than post-harvest stands because of high charcoal occurrence on the soil surface (Lyons et al., 2008; Chambers and Chapin, 2002). However, our soil reflectance data indicate that soils from 4-year-old post-fire stands unexpectedly showed significantly higher reflectance in the visible spectrum than did post-harvest stands (with the pattern reversed in the NIR spectrum) (Fig. 3a). Similar patterns in spectral response were recently observed in a biochar-amended agricultural soil relative to the control (Zhang et al., 2013). Soils from older post-harvest stands (11- and 19-year old), as expected, showed higher reflectance in the visible and NIR spectra compared to post-fire stands of similar age (Fig. 3b). Most post-harvest stands exhibited patches of charcoal in surface soils, presumably originating from historical fires. Legacy charcoals have similarly been visually observed on the forest floor and within upper mineral soils even after a hundred years following wildfire in Scandinavian (Ohlson et al., 2009) and Russian (Wallenius, 2002) boreal forests. The importance of “legacy” soil charcoal on surface albedo of harvested stands has not been considered previously to our knowledge. Charcoal reflectance is highly dependent on charring conditions (e.g., temperature, oxygen content) (Hudspith et al., 2015), and may possibly change with weathering; these processes require additional study in the context of albedo and surface energy balance.

Deleted: rapid

Deleted: year

Deleted: (personal observations)

Although charcoal residues likely have some influence on post-disturbance albedo, our results from both snow-covered and snow-free seasons strongly suggest that fire residues on the ground cannot explain the observed differences in albedo between post-harvest and post-fire stands. This result is consistent with the generalization that stand structure and composition are the main drivers of surface albedo and energy balance in the boreal forest (Amiro et al., 2006a).

Deleted: Our

#### 4.2 Albedo convergence with stand age in post-harvest and post-fire stands

425 Compiled data for winter and summer albedos from post-harvest and post-fire stands indicate that **changes in surface albedo**  
430 **continue for some decades following disturbance** (Figs. 4a–d). This finding does not support our second hypothesis that albedo  
**shows an early saturation near the onset of the ‘stem exclusion’ phase**. The rationale behind this hypothesis was that **high**  
**productivity of mixedwood conifer stands would result in more rapid canopy closure and attainment of peak LAI**. Studies  
435 using remote sensing techniques suggest that albedo in both post-harvest and post-fire stands **commonly saturates at ~40-80**  
years after harvest/fire (Bright et al., 2015a; Kuusinen et al., 2014; Lyons et al., 2008; McMillan and Goulden, 2008),  
consistent with our findings. These results thus suggest that gradual changes in species composition through later stages of  
succession are an important driver of stand albedo. Stand structural features such as canopy height (in winter) and ground  
vegetation cover (in summer) usually increase with stand age (Bartels et al., 2016) and might additionally contribute to a  
gradual reduction in albedo (Hovi et al., 2016) after ~25 years (Table 2, Figs. 7 and 8).

435 The shape of best-fit curves for winter albedo vs. stand age (exponential decay) of post-harvest (Fig. 4a) and post-fire (Fig.  
4b) stands are similar to other studies (Bright et al., 2015b; Kuusinen et al., 2014; Lyons et al., 2008; McMillan and Goulden,  
2008; Amiro et al., 2006b); however our results diverge markedly for summer albedo. Our best-fit curves for summer albedo  
440 vs. stand age for both post-harvest and post-fire stands showed pronounced peaks in early albedo described by double  
exponential functions (Fig. 4c–d), whereas Amiro et al (2006b) described data with a negative linear relationship, and other  
remote sensing-based studies have used exponential decay curves (e.g., Kuusinen et al., 2014). However, Lyons et al. (2008)  
and Randerson et al. (2006) found summer albedo of post-fire stands were related to stand age via a humped-shape curve, and  
albedo reached peak at ~20 years and gradually levelled off at ~50 years after fire, which closely corresponds to our findings  
445 (although our observed peak is at ~10 years post-disturbance: Fig. 4c–d). We suggest that most prior studies with sparser or  
more noisy data sets may have missed this early peak pattern. Immediately after fires and harvesting (because of high soil  
moisture, decaying CWD, legacy charcoal etc.) the summer albedo of post-harvest and post-fire stands is expected to show a  
low value (also see section 4.1 and Fig. 3) which sharply increases as dark ground is covered with early successional pioneer  
species (Lyons et al., 2008; Randerson et al., 2006; Amiro et al., 2006b; Betts and Ball, 1997). This sharp increase continues  
450 until ~20 years of stand age but then decreases slowly until ~50 years and saturates – consistent with the patterns found in  
other seasons.

455 We did not have albedo data **from** late-seral post-harvest or post-fire stands for spring and fall. In post-harvest (Fig. 5a) and  
post-fire (Fig. 5b) stands spring albedos did not show strong patterns with stand age, and the patterns were disturbance-specific  
(exponential decrease vs. negative exponential growth, respectively). Results from Kuusinen et al. (2014), Lyons et al. (2008),  
and Randerson et al. (2006) also suggest that patterns of spring albedo as a function of stand age can be disturbance-specific.  
In post-harvest stands, Kuusinen et al. (2014) found that spring albedo was high immediately after harvest and decreased  
exponentially until ~50 years and then saturated. However, in post-fire stands, Lyons et al. (2008) and Randerson et al.  
(2006), found hump-shaped patterns with a peak at ~10–15 years, and subsequent declines, similar to the winter albedo pattern.  
As discussed in Section 4.1, disturbance-specific responses may partially be attributed to the presence of black carbon  
460 (charcoal/soot) **on** snow immediately after a fire, which can substantially reduce snow albedo (Qian et al., 2009). Trends in  
fall albedo values with stand age in post-harvest (Fig. 5c) and post-fire (Fig. 5d) stands showed stronger patterns than spring,  
but similar disturbance-specific responses. Immediately after harvesting fall albedo was high, and exponentially decreased as

**Deleted:** saturates at ~50 years of stand age in the boreal forest

**Deleted:** saturates after ~

**Deleted:** 25 years of stand age (during

**Deleted:** )

**Deleted:** a high cover of deciduous species until ~25 years contributes to high albedo in both snow-covered (due to leaf shedding) and snow-free (due to high canopy albedo) seasons; after this age, deciduous species are replaced by conifers and albedo tends to saturate (Kuusinen et al., 2014; Amiro et al., 2006b).

**Deleted:** also

**Deleted:** 50

**Deleted:** or

**Deleted:** in

stand age increased. Increased fall albedo in recent post-harvest stands may be due to contributions to senescing leaves and to snow in the late fall (Amiro et al., 2006b; Liu et al., 2005). In contrast, fall albedo immediately after a fire was low (possibly because of charcoal or soot residues as discussed above), and increased with stand age.

The ability to predict forest surface albedo from stand age using the models presented in this study might be useful to the forest managers to easily incorporate albedo in forest management practice to develop climate-sensitive forestry practice.

Predicting albedo has been a long-standing problem in climate simulations (Bright et al., 2015a; Kuusinen et al., 2012; Qu and Hall, 2007). Our findings indicate that there are important qualitative differences in the post-disturbance albedo patterns between seasons in boreal forests. These differences need to be considered in enhancing albedo predictability of land surface models.

#### 4.3 Deciduous broadleaf species as a key determinant of surface albedo in the post-harvest and post-fire stands

Our results indicate that the proportion of deciduous broadleaf species is a strong predictor of albedo irrespective of disturbance type, and in most cases a better predictor than stand age (Figs. 4–6). Using remote sensing techniques Kuusinen et al. (2014) also found that stand age alone was not consistently the best predictor of stand albedo in the boreal forest. We found a similar mean model residual sum of errors for snow-covered seasons and snow-free seasons (Figs. 4–5 vs. Fig. 6), indicating that the proportion of deciduous broadleaf species are similarly important in both cases. These findings strongly support our third hypothesis that stands with a higher proportion of deciduous broadleaf species show higher albedo than conifer-dominated stands, but also that this effect is pronounced under both snow-covered and snow-free conditions. Except for fall post-fire stands, the relationship between albedo and proportion of deciduous broadleaf species approximated by an exponential saturating curve in which albedo declined rapidly where the proportion of deciduous broadleaf species fell below 25–50 %. Fall albedo in post-fire stands, on the other hand, was found to be even more sensitive, with a drop in fall albedo at a proportion of deciduous broadleaf species below 80%. We speculate that this sensitivity was related to exposure of fire residues in early stand development.

Overall our results indicate a strong dependency of seasonal albedo on the proportion of deciduous broadleaf species both in post-harvest and post-fire stands. This effect provides a strong link between albedo and successional patterns in mixedwood boreal forests. Prior studies addressing this relationship (e.g., Lyons et al., 2008; Amiro et al., 2006b) have suggested that increasing deciduous tree cover results in increased albedo values from stand initiation to ~ 25 years of stand age; thereafter, conifers start dominating the canopy, canopy height increases, and albedo decreases gradually until ~ 50 years of stand age before reaching a steady state. The data presented in the current study provide a somewhat different picture of these trends, in that patterns show important quantitative differences depending on the season and disturbance type. The importance of deciduous broadleaf species in the albedo signal over ~ 50 years of stand development suggests that slow successional changes in species composition are likely the main driver of the age-related patterns in mixedwood boreal forests albedo. The dynamics of this pattern is likely to depend on the intensity and frequency of disturbance, edaphic conditions, species abundance, and climate (Taylor and Chen, 2011). For example, in dry nutrient-poor boreal stands, deciduous broadleaf species-driven albedo might never occur, as such stands are commonly dominated by jack pine (Taylor and Chen, 2011); however, in mesic moderate-nutrient-rich stands, deciduous broadleaf species can dominate for ~ 100 years (Cogbill, 1985). Future studies

Deleted: classical

should prioritize robust modelling of boreal succession pathways under different biotic/abiotic conditions to properly characterize stand albedo.

## 5 Conclusions

520 This study presents the first available data on albedo patterns in boreal forests for all four seasons as well as the first  
comparisons of albedo in post-fire and post-harvest stands in the mixedwood boreal forest. The new data presented here are  
from 15 instrumented sites each monitored for four years, providing 60 site-years of measurements, all in mixedwood boreal  
forests that the most important forest from a forest management perspective, but for which there are almost no prior albedo  
525 measurements. Analyses of this unique dataset indicate that: i) winter and spring albedo values are substantially higher in  
post-harvest than in post-fire stands; ii) post-disturbance patterns of albedo recovery in boreal mixedwood stands are strongly  
influenced by changes in species composition; iii) there are important stand-age-related dynamics in albedo in the first 20  
years following disturbance events that have been missed by prior studies.

530 These findings have important implications for climate-friendly forest management practices. Since the proportion of  
deciduous broadleaf species is a strong predictor of seasonal albedo, stand-level albedo can be increased by enhancing the  
proportion of deciduous broadleaf species in a stand. Precisely this approach has recently been suggested as an adaptation and  
mitigation strategy to counter negative climate forcings of boreal forest (Astrup et al., 2018), but empirical data from actual  
managed stands have been lacking. Historically, forest managers have commonly sought to decrease or eliminate deciduous  
species and enhance conifers. However, there is strong evidence that local tree diversity enhances productivity in boreal  
535 forests as in other systems (Paquette and Messier, 2011), and in particular that mixedwood boreal forests including both  
conifers and deciduous trees show high productivity (MacPherson et al., 2001; Zhang et al., 2012). Management to increase  
the proportion of deciduous broadleaf species in managed boreal forests (for example, simply by avoiding chemical herbicide  
used to kill deciduous broadleaf species or retaining deciduous broadleaf species seed-trees) could thus be a “win-win”  
scenario for enhanced carbon sequestration via primary productivity, and climate mitigation via enhanced albedo.

540 In climate modeling studies albedo estimation for boreal forests have commonly been achieved by highly simplified  
representations of vegetation dynamics (Thackeray et al., 2019). In a recent study, Bright et al. (2018) pointed out that  
overlooking stand structural and compositional properties over the successional trajectory is likely to substantially bias  
radiative forcing estimates in the boreal forest. Ground-based estimates such as those presented are essential: at high latitudes  
545 when solar zenith angle is high ( $> 70^\circ$ ), satellites such as MODIS often provide poor-quality albedo data due to spatial  
heterogeneity of the landscape pixel signature and performance degradation of atmospheric correction algorithms (Bright et  
al., 2015b; Wang et al., 2012). Our findings based on field data are thus important in evaluating and potentially improving  
albedo predictions in land surface characterizations with climate models, and in improving albedo estimates derived from  
remote sensing. In addition, our results point to the importance of slow ecological succession as a driver of age-related patterns  
550 in albedo, suggesting that future models should explicitly incorporate these ecological processes to better predict long-term  
trends in climate forcings in boreal forests.

Deleted: Our

## 6 Code availability

R (version 3.5.1) codes used in the data analysis can be requested to M.A.H. (abdul.halim@mail.utoronto.ca).

## 555 7 Data availability

Data used to produce the graphs are also available via requesting corresponding author (abdul.halim@mail.utoronto.ca).

## 8 Author contribution

M.A.H., H.Y.H.C., and S.C.T. designed the experiment. M.A.H. analysed data with inputs from S.C.T. and H.Y.H.C. M.A.H. wrote the manuscript with edits and comments from H.Y.H.C. and S.C.T.

Deleted: o

## 560 9 Competing interests

The authors declare no competing interests.

## 10 Acknowledgements

565 Authors acknowledge assistances from Jillian Bieser, Kyle Gaynor, Lutchmee Sujeeun, Jack Richard, Jad Murtada, Anna Almero, and Hiro Sato during experiment setup and data collection. We thank Mark Horsburgh for his helpful comments on an earlier version of this manuscript and Shannon Brown for her help during the field comparison of pyranometers. We would also like to thank two anonymous reviewers and the editor for their critical comments, which helped to improve the manuscript. This study was funded by the NSERC (Natural Sciences and Engineering Research Council of Canada) Strategic Grant (grant number: STPGP 428641) and the NSERC Discovery Grant (grant number: RGPIN 06209).

## 11 References

- 570 Amiro, B., Barr, A., Black, T., Iwashita, H., Kljun, N., Mccaughey, J., Morgenstern, K., Murayama, S., Nestic, Z. and Orchansky, A.: Carbon, energy and water fluxes at mature and disturbed forest sites, Saskatchewan, Canada, *Agric. For. Meteorol.*, 136(3–4), 237–251, doi:10.1016/j.agrformet.2004.11.012, 2006a.
- 575 Amiro, B. D., Orchansky, A. L., Barr, A. G., Black, T. A., Chambers, S. D., Chapin III, F. S., Goulden, M. L., Litvak, M., Liu, H. P., McCaughey, J. H., McMillan, A. and Randerson, J. T.: The effect of post-fire stand age on the boreal forest energy balance, *Agric. For. Meteorol.*, 140(1), 41–50, doi:10.1016/j.agrformet.2006.02.014, 2006b.
- Astrup, R., Bernier, P. Y., Genet, H., Lutz, D. A. and Bright, R. M.: A sensible climate solution for the boreal forest, *Nat. Clim. Change*, 8(1), 11–12, doi:10.1038/s41558-017-0043-3, 2018.
- Avery, T. E. and Burkhart, H. E.: *Forest measurements*, 5th ed., Waveland Press., 2015.
- 580 Bala, G., Caldeira, K., Wickett, M., Phillips, T. J., Lobell, D. B., Delire, C. and Mirin, A.: Combined climate and carbon-cycle effects of large-scale deforestation, *Proc. Natl. Acad. Sci.*, 104(16), 6550–6555, doi:10.1073/pnas.0608998104, 2007.
- Baltzer, J. L. and Thomas, S. C.: Leaf optical responses to light and soil nutrient availability in temperature deciduous trees, *Am. J. Bot.*, 92(2), 214–223, 2005.
- Bartels, S. F., Chen, H. Y. H., Wulder, M. A. and White, J. C.: Trends in post-disturbance recovery rates of Canada's forests following wildfire and harvest, *For. Ecol. Manag.*, 361, 194–207, doi:10.1016/j.foreco.2015.11.015, 2016.

- Betts, A. K. and Ball, J. H.: Albedo over the boreal forest, *J. Geophys. Res. Atmospheres*, 102(D24), 28901–28909, doi:10.1029/96JD03876, 1997.
- Betts, R. A.: Offset of the potential carbon sink from boreal forestation by decreases in surface albedo, *Nature*, 408(6809), 187–190, doi:10.1038/35041545, 2000.
- 590 Brassard, B. W. and Chen, H. Y. H.: Effects of Forest Type and Disturbance on Diversity of Coarse Woody Debris in Boreal Forest, *Ecosystems*, 11(7), 1078–1090, doi:10.1007/s10021-008-9180-x, 2008.
- Brassard, B. W. and Chen, H. Y. H.: Stand Structure and Composition Dynamics of Boreal Mixedwood Forest: Implications for Forest Management, Sustainable Forest Management Network., 2010.
- 595 Brassard, B. W., Chen, H. Y. H., Wang, J. R. and Duinker, P. N.: Effects of time since stand-replacing fire and overstory composition on live-tree structural diversity in the boreal forest of central Canada, *Can. J. For. Res.*, 38(1), 52–62, doi:10.1139/X07-125, 2008.
- Bright, R. M., Astrup, R. and Strømman, A. H.: Empirical models of monthly and annual albedo in managed boreal forests of interior Norway, *Clim. Change*, 120(1–2), 183–196, doi:10.1007/s10584-013-0789-1, 2013.
- 600 Bright, R. M., Zhao, K., Jackson, R. B. and Cherubini, F.: Quantifying surface albedo and other direct biogeophysical climate forcings of forestry activities, *Glob. Change Biol.*, 21(9), 3246–3266, doi:10.1111/gcb.12951, 2015a.
- Bright, R. M., Myhre, G., Astrup, R., Antón-Fernández, C. and Strømman, A. H.: Radiative forcing bias of simulated surface albedo modifications linked to forest cover changes at northern latitudes, *Biogeosciences*, 12(7), 2195–2205, doi:10.5194/bg-12-2195-2015, 2015b.
- 605 Bright, R. M., Bogren, W., Bernier, P. and Astrup, R.: Carbon-equivalent metrics for albedo changes in land management contexts: relevance of the time dimension, *Ecol. Appl.*, 26(6), 1868–1880, doi:10.1890/15-1597.1, 2016.
- Bright, R. M., Eisner, S., Lund, M. T., Majasalmi, T., Myhre, G. and Astrup, R.: Inferring Surface Albedo Prediction Error Linked to Forest Structure at High Latitudes, *J. Geophys. Res. Atmospheres*, 123(10), 4910–4925, doi:10.1029/2018JD028293, 2018.
- 610 Brown, P. T. and Caldeira, K.: Greater future global warming inferred from Earth’s recent energy budget, *Nature*, 552(7683), 45–50, doi:10.1038/nature24672, 2017.
- Chambers, S. D. and Chapin, F. S.: Fire effects on surface-atmosphere energy exchange in Alaskan black spruce ecosystems: Implications for feedbacks to regional climate, *J. Geophys. Res.*, 108(D1), doi:10.1029/2001JD000530, 2002.
- Chen, H. Y. and Popadiouk, R. V.: Dynamics of North American boreal mixedwoods, *Environ. Rev.*, 10(3), 137–166, doi:10.1139/a02-007, 2002.
- 615 Chen, H. Y. H., Vasiliauskas, S., Kayahara, G. J. and Ilisson, T.: Wildfire promotes broadleaves and species mixture in boreal forest, *For. Ecol. Manag.*, 257(1), 343–350, doi:10.1016/j.foreco.2008.09.022, 2009.
- Cogbill, C. V.: Dynamics of the boreal forests of the Laurentian Highlands, Canada, *Can. J. For. Res.*, 15(1), 252–261, doi:10.1139/x85-043, 1985.
- 620 Colombo, S. J., Parker, W. C., Luckai, N., Dang, Q. and Cai, T.: The Effects of Forest Management on Carbon Storage in Ontario’s Forests, Climate Change Research Report, Ontario Forest Research Institute, Ministry of Natural Resources, Canada, Toronto: Queens Printer for Ontario. [online] Available from: [http://www.climateontario.ca/MNR\\_Publications/276922.pdf](http://www.climateontario.ca/MNR_Publications/276922.pdf), 2005.
- Conway, H., Gades, A. and Raymond, C. F.: Albedo of dirty snow during conditions of melt, *Water Resour. Res.*, 32(6), 1713–1718, doi:10.1029/96WR00712, 1996.
- 625 Environment Canada: Historical climate data, [online] Available from: [http://climate.weather.gc.ca/historical\\_data/search\\_historic\\_data\\_e.html](http://climate.weather.gc.ca/historical_data/search_historic_data_e.html) (Accessed 11 January 2018), 2018.


- Halim, M. A. and Thomas, S. C.: Surface albedo in relation to disturbance and early stand dynamics in the boreal forest: Implications for climate models., p. abstract #B21F-2025, American Geophysical Union Fall Meeting, 11-15 December 2017, New Orleans, USA., 2017.
- 630 Hansen, J., Sato, M., Ruedy, R., Nazarenko, L., Lacis, A., Schmidt, G. A., Russell, G., Aleinov, I., Bauer, M., Bauer, S., Bell, N., Cairns, B., Canuto, V., Chandler, M., Cheng, Y., Del, G. A., Faluvegi, G., Fleming, E., Friend A., Hall T., Jackman C., Kelley M., Kiang N., Koch D., Lean J., Lerner J., Lo K., Menon S., Miller R., Minnis P., Novakov T., Oinas V., Perlwitz Ja., Perlwitz Ju., Rind D., Romanou A., Shindell D., Stone P., Sun S., Tausnev N., Thresher D., Wielicki B., Wong T., Yao M. and Zhang, S.: Efficacy of climate forcings, *J. Geophys. Res. Atmospheres*, 110(D18), doi:10.1029/2005JD005776, 2005.
- 635 Hart, S. and Luckai, N.: Charcoal function and management in boreal ecosystems, edited by P. Brando, *J. Appl. Ecol.*, 1197–1206, doi:10.1111/1365-2664.12136, 2013.
- Hovi, A., Liang, J., Korhonen, L., Kobayashi, H. and Rautiainen, M.: Quantifying the missing link between forest albedo and productivity in the boreal zone, *Biogeosciences*, 13(21), 6015–6030, doi:10.5194/bg-13-6015-2016, 2016.
- 640 Hudspeth, V. A., Belcher, C. M., Kelly, R. and Hu, F. S.: Charcoal Reflectance Reveals Early Holocene Boreal Deciduous Forests Burned at High Intensities, *PLoS ONE*, 10(4), doi:10.1371/journal.pone.0120835, 2015.
- Johnstone, J. F., Hollingsworth, T. N., Chapin, F. S. and Mack, M. C.: Changes in fire regime break the legacy lock on successional trajectories in Alaskan boreal forest, *Glob. Change Biol.*, 16(4), 1281–1295, doi:10.1111/j.1365-2486.2009.02051.x, 2010.
- 645 Kirschbaum, M. U. F., Whitehead, D., Dean, S. M., Beets, P. N., Shepherd, J. D. and Ausseil, A.-G. E.: Implications of albedo changes following afforestation on the benefits of forests as carbon sinks, *Biogeosciences*, 8(12), 3687–3696, doi:10.5194/bg-8-3687-2011, 2011.
- Kumar, P., Chen, H. Y. H., Thomas, S. C. and Shahi, C.: Epixylic vegetation abundance, diversity, and composition vary with coarse woody debris decay class and substrate species in boreal forest, *Can. J. For. Res.*, 48(4), 399–411, doi:10.1139/cjfr-2017-0283, 2018.
- 650 Kuusinen, N., Kolari, P., Levula, J., Porcar-Castell, A., Stenberg, P. and Berninger, F.: Seasonal variation in boreal pine forest albedo and effects of canopy snow on forest reflectance, *Agric. For. Meteorol.*, 164, 53–60, doi:10.1016/j.agrformet.2012.05.009, 2012.
- Kuusinen, N., Tomppo, E., Shuai, Y. and Berninger, F.: Effects of forest age on albedo in boreal forests estimated from MODIS and Landsat albedo retrievals, *Remote Sens. Environ.*, 145, 145–153, doi:10.1016/j.rse.2014.02.005, 2014.
- 655 Kuusinen, N., Stenberg, P., Tomppo, E., Bernier, P. and Berninger, F.: Variation in understory and canopy reflectance during stand development in Finnish coniferous forests, *Can. J. For. Res.*, 45(8), 1077–1085, doi:10.1139/cjfr-2014-0538, 2015.
- Kuusinen, N., Stenberg, P., Korhonen, L., Rautiainen, M. and Tomppo, E.: Structural factors driving boreal forest albedo in Finland, *Remote Sens. Environ.*, 175, 43–51, doi:10.1016/j.rse.2015.12.035, 2016.
- 660 Lee, X., Goulden, M. L., Hollinger, D. Y., Barr, A., Black, T. A., Bohrer, G., Bracho, R., Drake, B., Goldstein, A., Gu, L., Katul, G., Kolb, T., Law, B. E., Margolis, H., Meyers, T., Monson, R., Munger, W., Oren, R., Paw U, K. T., Richardson, A. D., Schmid, H. P., Staebler, R., Wofsy, S. and Zhao, L.: Observed increase in local cooling effect of deforestation at higher latitudes, *Nature*, 479(7373), 384–387, doi:10.1038/nature10588, 2011.
- 665 Li, Y., Wang, T., Zeng, Z., Peng, S., Lian, X. and Piao, S.: Evaluating biases in simulated land surface albedo from CMIP5 global climate models: Albedo Evaluation in CMIP5, *J. Geophys. Res. Atmospheres*, 121(11), 6178–6190, doi:10.1002/2016JD024774, 2016.
- Linacre, E.: *Climate Data and Resources: A Reference and Guide*, e-book., Routledge, London., 2003.
- Liu, H., Randerson, J. T., Lindfors, J. and Chapin, F. S.: Changes in the surface energy budget after fire in boreal ecosystems of interior Alaska: An annual perspective, *J. Geophys. Res.*, 110(D13), doi:10.1029/2004JD005158, 2005.

- 670 Lukeš, P., Stenberg, P., Rautiainen, M., Möttö, M. and Vanhatalo, K. M.: Optical properties of leaves and needles for boreal tree species in Europe, *Remote Sens. Lett.*, 4(7), 667–676, doi:10.1080/2150704X.2013.782112, 2013a.
- Lukeš, P., Stenberg, P. and Rautiainen, M.: Relationship between forest density and albedo in the boreal zone, *Ecol. Model.*, 261–262, 74–79, doi:10.1016/j.ecolmodel.2013.04.009, 2013b.
- 675 Luysaert, S., Marie, G., Valade, A., Chen, Y.-Y., Njakou Djomo, S., Ryder, J., Otto, J., Naudts, K., Lansø, A. S., Ghattas, J. and McGrath, M. J.: Trade-offs in using European forests to meet climate objectives, *Nature*, 562(7726), 259–262, doi:10.1038/s41586-018-0577-1, 2018.
- Lyons, E. A., Jin, Y. and Randerson, J. T.: Changes in surface albedo after fire in boreal forest ecosystems of interior Alaska assessed using MODIS satellite observations, *J. Geophys. Res. Biogeosciences*, 113(G02012), doi:10.1029/2007JG000606, 2008.
- 680 MacPherson, D. M., Loeffers, V. J. and Blenis, P. V.: Productivity of aspen stands with and without a spruce understory in Alberta's boreal mixedwood forests, *For. Chron.*, 77(2), 351–356, doi:10.5558/tfc77351-2, 2001.
- Madoui, A., Gauthier, S., Leduc, A., Bergeron, Y. and Valeria, O.: Monitoring Forest Recovery Following Wildfire and Harvest in Boreal Forests Using Satellite Imagery, *Forests*, 6(12), 4105–4134, doi:10.3390/f6114105, 2015.
- Mair, P. and Wilcox, R.: WRS2: Wilcox robust estimation and testing, R package. [online] Available from: <https://r-forge-project.org/projects/psychor/>, 2018.
- 685 Matthies, B. D. and Valsta, L. T.: Optimal forest species mixture with carbon storage and albedo effect for climate change mitigation, *Ecol. Econ.*, 123, 95–105, doi:10.1016/j.ecolecon.2016.01.004, 2016.
- McMillan, A. M. S. and Goulden, M. L.: Age-dependent variation in the biophysical properties of boreal forests: Biophysical Properties of boreal forests, *Glob. Biogeochem. Cycles*, 22(2), n/a-n/a, doi:10.1029/2007GB003038, 2008.
- 690 Moussaoui, L., Fenton, N., Leduc, A. and Bergeron, Y.: Can Retention Harvest Maintain Natural Structural Complexity? A Comparison of Post-Harvest and Post-Fire Residual Patches in Boreal Forest, *Forests*, 7(12), 243, doi:10.3390/f7100243, 2016.
- Myers, D. R.: Comparison of direct normal irradiance derived from silicon and thermopile global hemispherical radiation detectors, edited by N. G. Dhere, J. H. Wohlgemuth, and K. Lynn, p. 77730G, San Diego, California., 2010.
- 695 Naudts, K., Chen, Y., McGrath, M. J., Ryder, J., Valade, A., Otto, J. and Luysaert, S.: Europe's forest management did not mitigate climate warming, *Science*, 351(6273), 597–600, doi:10.1126/science.aad7270, 2016.
- [Ohlson, M., Dahlberg, B., Økland, T., Brown, K. J. and Halvorsen, R.: The charcoal carbon pool in boreal forest soils, \*Nat. Geosci.\*, 2\(10\), 692–695, doi:10.1038/ngeo617, 2009.](#)
- 700 OMNRF: Forest Management Guide to Silviculture in the Great Lakes-St. Lawrence and Boreal Forests of Ontario., Ontario Legislative Library (OGDC) Open Access Electronic Government Documents, Ministry of Natural Resources and Forestry, Toronto: Queens Printer for Ontario. [online] Available from: <https://docs.ontario.ca/documents/4125/revised-silvguide-mar-2015-aoda-compliant.pdf>, 2015.
- Paquette, A. and Messier, C.: The effect of biodiversity on tree productivity: from temperate to boreal forests: The effect of biodiversity on the productivity, *Glob. Ecol. Biogeogr.*, 20(1), 170–180, doi:10.1111/j.1466-8238.2010.00592.x, 2011.
- 705 Qian, Y., Gustafson, W. I., Leung, L. R. and Ghan, S. J.: Effects of soot-induced snow albedo change on snowpack and hydrological cycle in western United States based on Weather Research and Forecasting chemistry and regional climate simulations, *J. Geophys. Res.*, 114(D3), doi:10.1029/2008JD011039, 2009.
- Qu, X. and Hall, A.: What controls the strength of snow-albedo feedback?, *J. Clim.*, 20(15), 3971–3981, doi:10.1175/JCL14186.1, 2007.

- 710 R Core Team: R: A language and environment for statistical computing., R Foundation for Statistical Computing, Vienna, Austria. [online] Available from: URL <https://www.R-project.org/>, 2018.
- Randerson, J. T., Liu, H., Flanner, M. G., Chambers, S. D., Jin, Y., Hess, P. G., Pfister, G., Mack, M. C., Treseder, K. K., Welp, L. R., Chapin, F. S., Harden, J. W., Goulden, M. L., Lyons, E., Neff, J. C., Schuur, E. A. G. and Zender, C. S.: The Impact of Boreal Forest Fire on Climate Warming, *Science*, 314(5802), 1130–1132, doi:10.1126/science.1132075, 2006.
- Rohatgi, A.: WebPlotDigitizer, Austin, Texas, USA. [online] Available from: <https://automeris.io/WebPlotDigitizer>, 2018.
- 715 Sims, R. A., Towill, W. D., Baldwin, K. A. and Wickware, G. M.: Field guide to the forest ecosystem classification for northwestern Ontario., 2nd ed., Ontario Ministry of Natural Resources, Northwest Science and Technology, Thunder Bay, Ontario., 1997.
- Stephens, G. L., O'Brien, D., Webster, P. J., Pilewski, P., Kato, S. and Li, J.: The albedo of Earth, *Rev. Geophys.*, 53(1), 141–163, doi:10.1002/2014RG000449, 2015.
- 720 Stroeve, J., Box, J. E., Gao, F., Liang, S., Nolin, A. and Schaaf, C.: Accuracy assessment of the MODIS 16-day albedo product for snow: comparisons with Greenland in situ measurements, *Remote Sens. Environ.*, 94(1), 46–60, doi:10.1016/j.rse.2004.09.001, 2005.
- Taylor, A. R. and Chen, H. Y. H.: Multiple successional pathways of boreal forest stands in central Canada, *Ecography*, 34(2), 208–219, doi:10.1111/j.1600-0587.2010.06455.x, 2011.
- 725 Thackeray, C. W., Fletcher, C. G. and Derksen, C.: Diagnosing the Impacts of Northern Hemisphere Surface Albedo Biases on Simulated Climate, *J. Clim.*, 32(6), 1777–1795, doi:10.1175/JCLI-D-18-0083.1, 2019.
- Uotila, A. and Kouki, J.: Understorey vegetation in spruce-dominated forests in eastern Finland and Russian Karelia: Successional patterns after anthropogenic and natural disturbances, *For. Ecol. Manag.*, 215(1–3), 113–137, doi:10.1016/j.foreco.2005.05.008, 2005.
- 730 [Wallenius, T.: Forest age distribution and traces of past fires in a natural boreal landscape dominated by \*Picea abies\*, \*Silva Fenn.\*, 36\(1\), doi:10.14214/sf.558, 2002.](#)
- Wang, Z., Schaaf, C. B., Chopping, M. J., Strahler, A. H., Wang, J., Román, M. O., Rocha, A. V., Woodcock, C. E. and Shuai, Y.: Evaluation of Moderate-resolution Imaging Spectroradiometer (MODIS) snow albedo product (MCD43A) over tundra, *Remote Sens. Environ.*, 117, 264–280, doi:10.1016/j.rse.2011.10.002, 2012.
- 735 Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, R, Springer-Verlag, New York. [online] Available from: <http://ggplot2.org>, 2016.
- Wilcox, R. R.: *Introduction to robust estimation and hypothesis testing*, 4th edition., Elsevier, Waltham, MA., 2016.
- WMO: Guidelines on the quality control of data from the World Radiometric Network, World Meteorological Organization, Leningrad. [online] Available from: [https://library.wmo.int/pmb\\_ged/wmo-td\\_258\\_en.pdf](https://library.wmo.int/pmb_ged/wmo-td_258_en.pdf), 1987.
- 740 Zhang, D.: *rsq: R-Squared and Related Measures*, R. [online] Available from: <https://CRAN.R-project.org/package=rsq>, 2018.
- Zhang, Q., Wang, Y., Wu, Y., Wang, X., Du, Z., Liu, X. and Song, J.: Effects of Biochar Amendment on Soil Thermal Conductivity, Reflectance, and Temperature, *Soil Sci. Soc. Am. J.*, 77(5), 1478–1487, doi:10.2136/sssaj2012.0180, 2013.
- Zhang, Y., Chen, H. Y. H. and Reich, P. B.: Forest productivity increases with evenness, species richness and trait variation: a global meta-analysis: Diversity and productivity relationships, *J. Ecol.*, 100(3), 742–749, doi:10.1111/j.1365-2745.2011.01944.x, 2012.
- 745


## 12 Figures and Tables

### 12.1 Figures




750

**Figure 1.** Map of the study area. Labels in the rectangular boxes indicate disturbance types (H: harvest and F: fire) and years (98: 1998, 06: 2006, and 13: 2013) for each plot (grey circles). **Inset:** black squares indicate locations of all data sources including the current study area (LN: Lake Nipigon area, Ontario, Canada; SK: Saskatchewan, Canada; MB: Manitoba, Canada).



755 **Figure 2.** Comparison of seasonal albedo (mean  $\pm$  SE) in post-harvest and post-fire stands. Winter (no. of observations for post-fire stands,  $n_F = 35$ ; no. of observations for post-harvest stands,  $n_H = 48$ ) and summer ( $n_F = 44$ ,  $n_H = 41$ ) albedo data were from 0–19-year-old stands, and spring ( $n_F = 30$ ,  $n_H = 30$ ) and fall ( $n_F = 30$ ,  $n_H = 30$ ) albedo data were from 7–19-year-old stands. Albedo of 0–6-year-old post-fire stands were from secondary sources. \* and \*\*\* indicate significant mean albedo differences between post-harvest and post-fire stands with  $p = 0.11$  and  $p < 0.01$ , respectively.


760



**Figure 3.** Specular-included ground surface reflectance (400–1000 nm) of post-harvest and post-fire stands. Lines indicate mean reflectance (number of sample (n) × 10 replicated measurements/sample) in the corresponding wavelengths, and shades indicate SE. **[a]** ground surface reflectance of young (4-year old) post-harvest stands (n = 9) and a post-fire stand (n = 12). **[b]** ground surface reflectance of old (11- and 19-year old) post-harvest (n = 18) and post-fire (n = 18) stands.

765


770



775 **Figure 4.** Stand age affecting mean seasonal albedo ( $\pm$  SE) in boreal forest over 0–150 years of stand development. Mean winter albedo as  
 a function of stand age in **[a]** post-harvest stands (n = 42) and **[b]** post-fire stands (n = 36). Mean summer albedo as a function of stand age  
 in **[c]** post-harvest stands (n = 41) and **[d]** post-fire stands (n = 30). Each field-data point is the average seasonal albedo (error bars indicate  
 standard errors) of three plots from each stand-age category over the study period.

780

785




**Figure 5.** Stand age affecting mean seasonal albedo ( $\pm$  SE) in boreal forest in the early seral stage. Mean spring albedo as a function of stand age in [a] post-harvest stands ( $n = 26$ ) and [b] post-fire stands ( $n = 14$ ). Mean fall albedo as a function of stand age in [c] post-harvest stands ( $n = 29$ ) and [d] post-fire stands ( $n = 22$ ). Each field-data point is the average seasonal albedo (error bars indicate standard errors) of three plots from each stand-age category over the study period.

790

795

800





**Figure 6.** Mean seasonal albedo ( $\pm$  SE) as a function of deciduous broadleaf species (%) (proportion of deciduous broadleaf species) in the boreal forest. Proportion of deciduous broadleaf species affecting mean winter albedo in [a] post-harvest stands ( $n = 17$ ) and [b] in post-fire stands ( $n = 20$ ), mean spring albedo in [c] post-harvest stands ( $n = 8$ ) and [d] post-fire stands ( $n = 6$ ), mean summer albedo in [e] post-harvest stands ( $n = 20$ ) and [f] post-fire stands ( $n = 15$ ), and mean fall albedo in [g] post-harvest stands ( $n = 8$ ) and [h] post-fire stands ( $n = 7$ ). *Note: albedo values of some 0–4 years old stands were omitted from this analysis because these young sites had only a few seedlings of*

810


deciduous broadleaf species. If we included them here, the percentage of deciduous broadleaf species for these sites became 100%, which was misleading compared to other sites; they were not zero either. Thus, the percentage of deciduous broadleaf species of these sites were excluded from this analysis and were considered as the ground vegetation cover (%) (Figure 8). Additionally, percentage deciduous broadleaf species of some secondary-data sites were not reported, so were excluded from this analysis.

815





**Figure 7.** Mean seasonal albedo ( $\pm$  SE) as a function of canopy height (m) in the boreal forest. Canopy height affecting [a] mean winter albedo in post-harvest stands ( $n = 31$ ), [c] mean spring albedo in post-harvest stands ( $n = 16$ ), [f] mean summer albedo in post-fire stands ( $n = 23$ ), and [g] mean fall albedo in post-harvest stands ( $n = 20$ ). In [b, d, e, h] canopy height is not a significant predictor of the corresponding mean seasonal albedo; thus, no model is fitted to the data points.



**Figure 8.** Mean seasonal albedo ( $\pm$  SE) as a function of ground vegetation cover (%) in the boreal forest. Ground vegetation cover affecting [e] mean summer albedo in post-harvest stands ( $n = 22$ ) and [g] mean fall albedo in post-harvest stands ( $n = 18$ ). In [a–d, f, h] ground vegetation cover is not a significant predictor of the corresponding mean seasonal albedo; thus, no model is fitted to the data points.

## 12.2 Tables

Table 1. Structural characteristics of post-harvest and post-fire stands sampled. Mean values ( $\pm$  SE) are reported across all sites of a given disturbance type.

Stand type	Stand age (year)	DBS (%)	LAI	Stem density (stems ha <sup>-1</sup> $\geq$ 5 cm DBH)	Height (m)	GCV (%)
Post-harvest	0–19	55.4 $\pm$ 11.2	0.4 $\pm$ 0.3	6472 $\pm$ 3060	1.7 $\pm$ 1.3	51.8 $\pm$ 20.1
Post-fire	7–19	37.8 $\pm$ 9.1	0.7 $\pm$ 0.4	8400 $\pm$ 1902	2.9 $\pm$ 1.5	62.5 $\pm$ 14.1

830

Notes: DBS, LAI, and GCV indicate deciduous broadleaf species (% by basal area), leaf area index, and ground cover vegetation.

835

840

845

850

855

860

865

870

**Table 2. Regression model coefficients and fit statistics for albedo as a function of stand attributes in different seasons in the boreal forest.**

Season	Post-harvest stands				Post-fire stands						
	Parameter Estimates		Model fit		Parameter Estimates		Model fit				
	Coefficient	Estimate	$\Delta AIC$	Adj. $R^2$	Coefficient	Estimate	$\Delta AIC$	Adj. $R^2$			
Winter	Intercept	<b>1.722</b>			Intercept	<b>-1.25</b>					
	SA	<b>-0.031</b>			SA	-0.004					
	PDBS	<b>-0.021</b>	-69.2	0.97	PDBS	<b>0.005</b>	-5.3	0.75			
	CH	<b>-0.079</b>									
	SA:CH	<b>0.002</b>									
	PDBS:CH	<b>-0.007</b>									
Intercept	<b>-7.195</b>					Intercept			<b>-1.747</b>		
SA	<b>1.298</b>	-495.4			0.99	SA			0.016	-18.8	0.92
PDBS	<b>0.116</b>										
CH	<b>-1.264</b>										
SA: PDBS	<b>-0.024</b>										
Intercept	<b>-1.377</b>					Intercept	<b>-2.996</b>				
SA	<b>0.032</b>		-24.9	0.97		SA	<b>-0.012</b>	-48.3	0.95		
PDBS	<b>-0.003</b>										
GVC	<b>-0.01</b>										
SA: GVC	<b>-0.0004</b>										
PDBS: GVC	<b>0.0001</b>										
Intercept	<b>0.398</b>					Intercept	$\frac{4.5}{6.87} e^{(-\frac{SA-13.2}{6.87} - e^{-\frac{SA-67.62}{45.39}})}$			-3.1	0.045 <sup>1</sup>
SA	0.013	-6.1	0.94	SA:CH: PDBS	$0.099 e^{0.013 PDBS}$	-25.4	0.008 <sup>1</sup>				
CH	<b>-0.182</b>										
GVC	<b>-0.007</b>										
SA:CH	<b>0.007</b>										
CH: GVC	<b>0.005</b>										
SA:CH: GVC	<b>-0.0002</b>										
Intercept	$\frac{28.86}{45.39} e^{(\frac{PDBS-67.62}{45.39} - e^{-\frac{PDBS-67.62}{45.39}})}$			Intercept		-0.9	0.049 <sup>1</sup>				

**Notes:** SA, PDBS, CH, and GVC indicate stand age (year), proportion of deciduous broadleaf species (%), canopy height (m), and ground vegetation cover (%), respectively. Parameter estimates for GLMs in bold and regular fonts indicate statistical significance at 1% and 5% level, respectively. For fall nonlinear regression models, 28.86 and 45.39 coefficients of post-harvest stands are significant at 5% level and the rest is significant at 1% level. <sup>1</sup> indicates residual standard error of the nonlinear regression model.  $\Delta AIC$  = AIC of the best-fit model – AIC of the corresponding null model.