We received three referees' reports. Referees 2 and 3 recommended minor revisions but Referee 1 recommended reject. We address each point made by the referees below.

Our responses are in red, new text added to the manuscript is indented and italicised.

Anonymous Referee #1

General comments: The manuscript by van der Horst et al., poses an interesting question about the FLUXNET data and about the representativeness of the flux measurements during temperature extremes. While the topic is of interest in particular to the modelling community, I have a major concern about their approach. The authors explore data availability at each measurement site based on the availability of the temperature, sensible and latent heats, and NEE data. They take the ratio of the available data for heat (latent or sensible) or NEE, relative to the available temperature data, also accessed through FLUXNET, to compare sites.

Yes, this is correct: our aim with this study was to independently assess the availability of the measured surface energy and carbon fluxes during extreme conditions.

This way data availability is biased by the availability of the temperature data. My questions is why did the authors not use complete temperature records (from meteorological or remote sensing products) for each site to compare with the absolute availability instead of taking a relative proxy that is a biased by the quality of temperature measurements and is not comparable between sites?

First, we clearly did not express our goals clearly enough and to resolve this we have added several new pieces of text in the introduction to make this clear including:

We use measurements of temperature and Q_{le} , Q_h and NEE from FLUXNET sites because ultimately we seek to identify those sites useful for developing, evaluating and benchmarking LSMs. To do this requires the LSM to be provided meteorological forcing measured concurrently with the fluxes. We therefore cannot blend the measured fluxes with meteorological observations taken elsewhere.

By providing simultaneous and co-located measurements of both the meteorological forcing of the surface, and the associated turbulent energy fluxes, FLUXNET provides a critical resource for the development, evaluation and benchmarking of land surface models.

We use measurements of temperature and Q_{le} , Q_h and NEE from FLUXNET sites because ultimately we seek to identify those sites useful for developing, evaluating and benchmarking LSMs. To do this requires the LSM to be provided meteorological forcing measured concurrently with the fluxes. We therefore cannot blend the measured fluxes with meteorological observations taken elsewhere.

Our goal is to identify those FLUXNET sites with data useful to explore land surface processes under extreme temperature conditions.

We agree that longer records could be achieved by using alternative data sources.

However, weather stations at other locations need not reflect the conditions at the flux tower (and therefore the measured fluxes), whereas remotely sensed products do not directly measure the air temperatures and depending on the product, will have a mismatch in spatial scale compared to the tower footprint (\sim 1 km²).

Second, we did check for bias and we have added some additional text to note that there is no bias – the temperature data from each FLUXNET site is much more freely available than any of the actual fluxes. We have added two new pieces of text to highlight this.

Overall, temperature observations were available 86% of the time, $Q_{\text{le},}\,62\%$ of the time, $Q_h\,68\%$ of the time and NEE 30% of the time.

We examined the availability of measured temperature relative to the potential availability after we excluded sites with less then 8 months of data, less than 50% of data being measured and nighttime. We note 88% of all sites reported measurements for more than 80% of the time. Only 6% of sites had measurements for 50%-70% of the time and we excluded sites with less than 50% from subsequent analysis.

We can display the frequency histogram to demonstrate this – we do not think this should be added to the paper however as the sentences above provide the required information. Clearly, virtually all the sites we use have 80-100% data cover for temperature:



The authors themselves suggest this approach to the modelling community in lines 31-33.

There are often many ways to approach a problem and each method has different strengths and weaknesses. In our approach, we were particularly motivated by how these data may be used to develop, run and evaluate models. Given this motivation,

there is no need to generate a complete temperature record, instead our aim was to assess whether during conditions one could run a model (i.e. because we have meteorological forcing data including observed temperature), whether we have matching surface energy and carbon fluxes with which we could evaluate the model output.

The text the reviewer is highlighting was:

One way forward to resolve how to choose FLUXNET data for extremes is to combine an analysis of meteorological sites with FLUXNET sites. Using sites maintained by meteorological agencies to identify extreme events (e.g. heatwaves) and then interrogate the FLUXNET sites near to the meteorological site for the availability of measurements of Q_{le} , Q_h and NEE could enable a modeller to choose suitable sites for land surface model development and evaluation.

We have added the following text to help clarify the issues here:

While one possible way forward, inconsistencies between observations from meteorological agencies relative to FLUXNET (location, geographical distribution, height of measurements, standardisation of measurements over short grass) highlight the challenges in using meteorological observations that are physically separate from the FLUXNET observations.

Specific comments:

Page 1 Line 17: Why not using the Tier 2 dataset that is more complete, if this study is focusing on data availability?

We did not use Tier 2 for the simple reason that the Tier 2 data set is not freely available to the community. Those with access to Tier 2 data can use our codes to reproduce our results if they wish but we are committed to using freely available data. We have added a statement in the paper to clarify this (Section 2.1):

Only freely-available site datasets from each release were used.

Page 1 Line 22: Perhaps they mean the "availability" of temperature and not "measurement ratio". Measurement ratio for temperature would be 1 based on their description.

No, we meant the measurement ratio, but we have modified this sentence from:

We showed that the measurement ratios for both sensible and latent heat fluxes are generally lower (0.79 and 0.73, respectively) than for temperature measurements, and the ratio of net ecosystem exchange measurements are appreciably lower (0.42)

to

We showed that the measurement ratios for both sensible and latent heat fluxes are generally lower (0.79 and 0.73, respectively) than for temperature

measurements, and the measurement ratio of net ecosystem exchange measurements are appreciably lower (0.42)

(we added the word "measurement" after "temperature")

Page 3 Lines 5-6: Exactly for this reason, the measurement ratio is relative to each site and cannot be compared across sites.

We acknowledge the reviewers statement but we are not clear what the reviewer really means here. The statement we made was:

While eddy-covariance data have been widely used to examine the impact of temperature extremes, the measurement of temperature and the measurement of Q_{le} , Q_h and NEE are independent.

We think that there may be a misunderstanding based on our explanation and to accommodate this we have added the following:

While eddy-covariance data have been widely used to examine the impact of temperature extremes, the measurement of temperature and the measurement of Q_{le} , Q_h and NEE are independent in terms of the instrumentation used. However, the measured temperature is provided in published data, along with measurements for the site of net radiation, wind speed, humidity etc alongside measurements of Q_{le} , Q_h and NEE. A land surface modeller requires all these data to drive a land surface model for evaluation or process-based studies. We are therefore interested in the relationship between measurements of temperature, and in particular extreme temperatures and concurrent measurements of Q_{le} , Q_h and NEE.

Page 9 line 8: Indeed. But, in my opinion, the authors should have assessed the quality of the flux data independently of the quality of temperature data since the two are measured separately.

We accept that this is the reviewer's opinion and we do of course respect that different researchers can approach a question in different ways. From our perspective, while temperature and fluxes are measured by separate instruments, they are not independent of each other in terms of ecosystem behaviour, due to feedbacks between the land and the atmosphere. To build process-based understanding of land surface dynamics during temperature extremes, the co-occurring measurements at flux towers offer a unique opportunity to explore this because the temperature and fluxes are measured at a specific location and then published within a freely accessible data set together. Similarly, a modeler has to use the observed temperature and the observed meteorology together to then use the measured fluxes because the temperature, radiation, wind are all used to force a land model. In the context of how these FLUXNET data are actually used by modellers, using other temperature data would have led to an incompatibility between our results and how the results can actually be used.

We think the reviewer's comment is directed at our statement:

Our analysis poses interesting questions about the FLUXNET data that deserve further exploration.

We think this is a valid statement and it is not clear how we might modify it to reflect the reviewer's opinion given the rest of this paragraph communicates some of these specific questions.

Anonymous Referee #2

The manuscript by van der Horst et al. addresses the availability of eddy covariance (EC) flux measurements under extreme temperature conditions, whereby 'extreme' is defined relative to each site. The analysis in this manuscript is, in my opinion, very well conducted and the results are described and presented in a very concise and clear manner. Potential caveats and misinterpretation of the results are well explained. This study will certainly be of interest to both the eddy covariance and land surface model community, and it will be very helpful for researchers selecting sites that experience temperature extremes.

Thank-you.

I do not have major criticism concerning the overall approach of this study, but some (mostly minor) suggestions on how the results are presented and discussed:

Second paragraph of the introduction section: I agree that land surface models need a better representation of physiological processes under extreme conditions, but I do not think that eddy covariance data are the key to provide these formulations. In this paragraph it reads a bit as if the authors try to suggest that with the right selection, EC data could be used to parameterize photosynthesis/respiration response curves in land surface model, which would be a huge stretch. EC data provide information on a bigleaf photosynthesis (provided that flux partitioning is correct), which cannot be directly used to parameterize LSMs which require information at leaf level. In addition, I am not sure how well NEE partitioning algorithms are tested/evaluated in extreme conditions. That being said, EC data are of course not useless for land surface modelers, but will be most useful (at least for physiology) in model evaluation/benchmarking, in which emergent canopy-level model results can be directly compared to EC data. The use of these data is mentioned at several points in the manuscript, and I would simply suggest some rewording here to not give the reader a wrong impression of what this study could be used for.

We had to choose to either delete the text or improve it a little to address this reviewer's comment and we concluded on balance to add a little further detail. We added:

Although model improvements in the representation of physiological responses to temperatures are likely to be driven by data from leaf-level and manipulation experiments, data from eddy-covariance is also of value. For example, Keenan et al. (2019) recently quantified an apparent inhibition of daytime ecosystem respiration, showing that the diurnal pattern differed from expectations using the global FLUXNET network.

Figure 1: as the authors explain in the results section, the sudden increase in the availability of flux measurements relative to temperature measurements above approx. 40 degrees is mostly caused by a few sites that experience such high temperatures, thus it could be a site selection effect and not a robust pattern. In order to make this clearer in this Figure (and not just in the text), one could present e.g. dashed lines (instead of solid lines), wherever the number of sites/site years is below a certain value, just to give the reader an idea about the robustness of the results at

different temperatures.

We have taken this advice on board and have revised the figure to use dashed lines where the sample size is less than 1000. The choice of 1000 is somewhat arbitrary of course.

page 2, I.35: is 'high-frequency' really the right term here?

We have modified this text to avoid the term while retaining the important text:

From each FLUXNET site, measurements of the exchange of latent heat flux (Q_{le}) , sensible heat flux (Q_h) and net ecosystem exchange (NEE) are available at 30- to 60-minute resolution, alongside meteorological variables (including air temperature, net radiation, precipitation and relative humidity).

page 2, I. 40ff: I suggest to list some more relevant studies here. It would be useful to get a better idea of what and how EC measurements taken under extreme conditions can be used for.

We have taken this advice on board and added further examples.

page 3, I. 7ff: '...whether the availability of direct observations...' would be better here?

We modified this section to accommodate Reviewer 1's criticisms and so we have not made this edit.

page 3, I. 33f: I suggest to move this sentence to the last paragraph of the introduction section.

We agree that we should move this sentence – but feel it is better at the beginning of the last paragraph of the introduction, rather than at the very end.

page 4, I. 10: what exactly do you mean by 'lack enough measurements'?

We mean there are sometimes very few observations to base analyses on if we use a 2 standard deviation threshold.

We have modified the text to read:

We did repeat our analysis using exactly the two standard deviations; this led to some qualitative differences in our results because some sites lack enough measurements to provide reliable results where the temperature distribution was not normally distributed.

page 5, I. 10f.: so where do the rainfall amounts in Figure 5 come from? Were they calculated from the same measurements that were used in this analysis? Please describe this in more detail here. Please add 'mean annual' precipitation and temperature to axis labels of Figure 5.

The data were sourced from FLUXNET and are from the same measurements as used in the other analysis. We have added appropriate text. We have also amended the axes as suggested.

page 5, I. 12ff: For me the absence of any relationship is the most striking aspect of

Figure 5, and it would make sense to describe this first.

We have restructured this paragraph to follow the advice of the reviewer and now talk about the absence of any relationship first.

page 5, I. 29ff: presentation of results is very detailed here and could be shortened for the sake of readability.

We agree it is quite detailed – it is actually a lot shorter than we originally drafted to try to help readability. We have not further reduced this part of the text in the revised version as on balance we are keen to provide the detail, but we are open to further advice from the editor.

page 6, I. 4ff: I think these results are also useful for observation-based studies, not only for modeling, which could be mentioned here or later in the paper.

Yes – we fully agree and it was not our intention to suggest otherwise. We have edited this sentence on the basis that we to touch on the value of our results for observational programs later. Specifically, at the end of the discussion, but we are also reluctant to be seen to "direct" the observational community, so we are attempting to retain balance.

The edit introduced is minor – we have simply edited this sentence from:

The FLUXNET eddy covariance flux measurements are among the most valuable observations available for developing, evaluating and benchmarking land surface models.

to

The FLUXNET eddy covariance flux measurements are among the most valuable observations available for understanding processes, and for developing, evaluating and benchmarking land surface models.

We hope this suffices but if we have misunderstood the reviewer we are happy to take further advice on board.

page 7, I. 4: if there is no 'second', remove the 'first'

We have made this correction

page 7, l. 27f.: maybe clarify that the removed data points are gap filled and hence not used for this analysis, which focuses on measured data only.

Yes – we have followed the reviewer's advice and made this clear by adding a sentence at the end of the paragraph:

We avoid the consequences of these procedures in quality controlling and gap-filling data by only using those data that are directly observed.

page 8, I. 6: delete 'are excluded'

Yes – thanks, we have corrected this.

page 8, I. 35ff.: this paragraph belongs mostly to the Results section.

We have followed the reviewer's advice and moved this paragraph to the results section.

Please do not present new results in the Conclusions (I.30ff.)

These are not results from our paper, they are contextualizing our results and so would not be appropriate in the results section. We have now moved this to the discussion section.

Figure 9 caption: 'separately', not 'separatly'

Thank-you – corrected.

Table S4: Please add whether the data come from the LaThuile or the FLUXNET2015 data set

Yes – we have added a column with this information to Table S4.

Anonymous Referee #3

General comments:

The manuscript by van der Hort et al. gives a very interesting overview of global FLUXNET data availability - within an objectively pre-selected subset of FLUXNET stations - with a focus on how well temperature extremes are represented. The results are of good use for, above all, the modelling community. It might provide help for selecting suitable sites for validating models in the context of climate projections. This is what the authors propose.

Thanks for these positive comments

I want to highlight that the results are interesting mainly because of their clear yet counter-intuitive character. For instance, the low correlation between the amount of precipitation as well as mean temperature and the data availability are rather unexpected.

Yes – and in responding to Reviewer 2 some changes in the text's order has helped highlight this.

Knowing that many factors other than (micro)meteorological play into the overall data availability, I think this study objectively summarizes the results using their own metric for an easy interpretation, without going into too much detail about the reasons (which is not the focus of this manuscript).

Yes – thank-you. That was our intent.

As a general comment on the used methodology I want to mention that using temperature measurements as the reference for data availability is confusing at first (as was mentioned by Referee 1). It seems arbitrary and is clearly biased by the availability of temperature measurements. However, I understand from the author's response that they rely on actually measured data available at the respective FLUXNET site as one possible method. It would help, though, if this is expressed in more detail. After all, temperature is used as 'reference length' of time series, since the metric of measurement ratio requires the temperature to be available. This simple

circumstance is not clearly described in the manuscript.

These comments relate to the comments by Reviewer 1. We have responded in detail to Reviewer 1 and hope that these changes will satisfy reviewer 3 too.

Also, a lot of possible comments related to shortcomings of the study (i.e., the qualitative character of the results) are dispersed with the caveat and comments in anticipation of criticism to certain points in the conclusion.

Yes – we obviously agree with the reviewer here.

Minor comments:

p9 l08: It is important to always make clear whether the mean and tails refer to persite or all sites

Yes - we have followed the reviewer's advice and added clarification.

p3 I18: 'most observations' should be something like 'longest time series' or similar

Yes – "most" is too vague. We have modified the last two sentences to read:

We therefore seek to identify which FLUXNET site data are most suitable for analysing processes under extreme temperature conditions with the goal of identifying those sites most useful for land surface model development and evaluation of the surface energy, water and carbon budgets during extreme temperatures.

p3 l38: A more widely used threshold is 10 Wm2, why is 1 used (instead of straightforwardly zero)? Also, how are erroneous measurements of shortwave radiation at night identified? Was this condition applied only if it seems erroneous?

Our text reads:

Given our focus on Q_{le} , Q_h and NEE we excluded night-time data where incoming shortwave radiation was < 1 W m⁻². In case of erroneous measurements of shortwave radiation at night we excluded all data between 11 pm and 6 am.

We did not use 10 W m⁻² for identifying night time because this seems a very high flux to use to exclude night time. That said, we see that our text was not clear and we have modified it to:

Given our focus on Q_{le} , Q_h and NEE we excluded night-time data using two criteria. We first excluded all data between 11 pm and 6 am local time. In addition, if shortwave radiation was < 1 W m⁻² for an individual time period then associated measurements were also excluded. This did not exclude many measurements as shortwave radiation was rarely reported as non-zero at night but there were occasional shortwave radiation > 1 W m⁻² in observations at night.

p4 I21ff: The numbers given on the following lines referring to Fig. 1, while given as approximates, are still inaccurate. E.g., the maximum ratio for Qh, Qle in 1b is clearly rather at 30 degC than 20 degC.

We agree the precision on our approximations could have been better and we have modified these in the text. However, the maximum ratio for Qh and Qle in Figure 1b is at 20° C – and this then remains very similar through to about 30° C. The modified text now reads:

At the lowest temperatures, ratios for Q_{le} and Q_h range from ~0 to ~0.3 but these increase as temperatures increase to maximum ratios of ~0.8 at around 20° C and remain at ~0.8 through to 30° C. For NEE ratios increase to ~0.6 at around 35° C for NEE. A minor dip in Q_{le} and Q_h ratios occurs at 0° C associated with the phase change of water, which most likely affects the operation of instrumentation. At the upper extreme of the temperature distribution, ratios decline between 30° C and 45° C from ~0.8 to ~0.6 for Q_{le} and Q_h and from ~0.6 to ~0.5 for NEE.

p4 I23: 'affect instrumentation' is a very unprecise wording, as the measurements are affected

Yes - we have modified this to

"which most likely affects the operation of instrumentation"

p4 I26: What is meant by '68 individual measurements'?

We mean, there are 68 actual reported measurements at this site at a temperature in excess of the 44°C threshold. We have revised the sentence to make this clear.

This peak is associated with temperatures > 44°C, which are rare and associated with measurements at Au-Cpr (there are only 68 individual measurements in excess of 44°C at this site), AU-GWW (23 individual measurements), AU-Stp (24 individual measurements) and SN-Dhr (33 individual measurements).

p5 I10f: This phrase seems odd and unnecessarily complicated, if meaning not representing the total amount of annual precipitation, please re-phrase.

We agree and have re-written the sentence:

Note the amounts of rainfall shown in Figure 5 are accumulated only over times when temperature data are selected and therefore cannot be compared with observations taken at meteorological stations.

p5 I14: from 'Qle and Qh, but for NEE, most are in cool sites...' I don't get this phrase. Should it read 'Qle and Qh but for NEE and most are in cool sites...'?

We agree that this was unclear. In part to help simplify our results section we decided this was a sentence that could be deleted.

Figure 1: Please show more ticks on the y-axis.

We have followed the reviewer's suggestion.

How representative are FLUXNET measurements of surface fluxes during temperature extremes?

Sophie V.J. van der Horst¹, Andrew J. Pitman^{2,3}, Martin G. De Kauwe^{2,3}, Anna Ukkola^{2,4}, Gab Abramowitz^{2,3}, Peter Isaac⁴

¹Meteorology and Air Quality, Wageningen University, 6700 HB, Wageningen, the Netherlands.
 ²ARC Centre of Excellence for Climate Extremes
 ³Climate Change Research Centre, University of New South Wales, Sydney, NSW, 2052, Australia
 ⁴Research School of Earth Sciences, Australian National University, Canberra, ACT, 2601 Australia

⁴OzFlux Central Node, TERN Ecosystem Processes, Melbourne, VIC 3159, Australia

10

5

Correspondence to: Andrew J. Pitman (a.pitman@unsw.edu.au)

Abstract. In response to a warming climate, temperature extremes are changing in many regions of the world. Therefore,

- 15 understanding how the fluxes of sensible heat, latent heat and net ecosystem exchange respond and contribute to these changes is important. We examined 216 sites from the open access Tier 1 FLUXNET2015 and Free-Fair-Use La Thuile datasets, focussing only on observed (non-gap filled) data periods. We examined the availability of sensible heat, latent heat and net ecosystem exchange observations coincident in time with measured temperature for all temperatures, and separately for the upper and lower tail of the temperature distribution and expressed this availability as a measurement ratio. We
- 20 showed that the measurement ratios for both sensible and latent heat fluxes are generally lower (0.79 and 0.73 respectively) than for temperature <u>measurements</u>, and the measurement ratio of net ecosystem exchange measurements are appreciably lower (0.42). However, sites do exist with a high proportion of measured sensible and latent heat fluxes, mostly over the United States, Europe and Australia. Few sites have a high proportion of measured fluxes at the lower tail of the temperature distribution over very cold regions (e.g. Alaska, Russia) or at the upper tail in many warm regions (e.g. Central America and
- 25 the majority of the Mediterranean region), and many of the world's coldest and hottest regions are not represented in the freely available FLUXNET data at all (e.g. India, the Gulf States, Greenland and Antarctica). However, some sites do provide measured fluxes at extreme temperatures suggesting an opportunity for the FLUXNET community to share strategies to increase measurement availability at the tails of the temperature distribution. We also highlight a wide discrepancy between the measurement ratios across FLUXNET sites that is not related to the actual temperature or rainfall

1

30 regimes at the site, which we cannot explain. Our analysis provides guidance to help select eddy covariance sites for researchers interested in <u>understanding and/or modelling</u> responses to temperature extremes. Martin De Kauwe 21/2/2019 12:28 PM Deleted: ³

Martin De Kauwe 21/2/2019 12:27 PM Deleted: and Martin De Kauwe 21/2/2019 12:27 PM Deleted: ³ARC Centre of Excellence for Climate

Extremes, and

Andy Pitman 26/2/2019 9:39 AM

Andy Pitman 26/2/2019 9:39 AM Deleted: exploring

1 Introduction

Changes in the upper and lower tails of the temperature distribution are key characteristics of how global warming will impact climate (Hartmann et al. 2013). These expected changes in temperature are in line with a series of recent high-profile extremes witnessed across Europe (2003, 2010; Coumou and Rahmstorf, 2012; Schär et al. 2004; Barriopedro et al. 2011),

- 5 western North America (van Mantgem et al. 2009), the Amazon (2005, 2010; Philips et al. 2009; Lewis et al. 2011) and Australia (2012/2013; van Gorsel et al. 2018). Changes in temperature extremes are not only limited to the warm tail; the cold tail has also seen a notable change, with observed decreases in cold extremes particularly across North America (Wolter et al. 2015). Given the wide-ranging impacts of temperature on vegetation function (Berry and Björkman, 1980; Gunderson et al., 2009; Valladares et al., 2014; Van Gorsel et al., 2016; Kumarathunge et al., in press), health (McMichael and
- 10 Lindgren, 2011), socio-economics (McEvoy et al., 2012; Colombo et al., 1999; Zander et al., 2015) and land-atmosphere feedbacks (Fischer et al., 2007; Teuling et al., 2010; Miralles et al., 2012; Kala et al., 2016; Donat et al. 2017), projecting the impact of changes in temperature extremes is critical.

Our understanding of how temperature extremes will change is based on simulations using coupled climate models, e.g. the Coupled Model Intercomparison Project (CMIP5) (Eyring et al., 2016). To build confidence in these projections, models should be consistent with our understanding of changing temperature extremes, the impact on the vegetation and the associated feedback on the climate. However, current models are known to have key weaknesses in simulating both temperature extremes (Sillmann et al., 2013; Sippel et al., 2017) and the response of the vegetation to these extremes. For example, most climate models represent broad geographic regions with a single photosynthetic temperature response

- 20 function, which varies only with plant functional type (Smith and Dukes 2013; Lombardozzi et al. 2015; Mercado et al. 2018). This assumption would seemingly contradict empirical evidence showing that the temperature response of photosynthesis varies as a function of climate (Berry & Björkman, 1980; Gunderson *et al.*, 2009). Furthermore, studies show that plants adjust their temperature response of photosynthesis and respiration to changes in ambient temperature (Way and Sage 2008; Lombardozzi et al. 2015). Although model improvements in the representation of physiological responses to
- 25 temperatures are likely to be driven by data from leaf-level and manipulation experiments, data from eddy-covariance is also of value. For example, Keenan et al. (2019) recently quantified an apparent inhibition of daytime ecosystem respiration, showing that the diurnal pattern differed from expectations using the global FLUXNET network.

Improving how well models simulate temperature extremes, and how vegetation responds to these extremes requires
empirical data. The global network of eddy-covariance towers (commonly known as FLUXNET), which includes over 900 sites and over 7000 site years (fluxdata.org, 2018) provides measurements of the exchange of carbon, energy and water between the land and the atmosphere. Therefore, eddy covariance measurements provide our best ecosystem-scale estimate of the vegetation's response to heat extremes (Ciais et al. 2005; Teuling et al. 2010; Wolf et al. 2013; von Buttlar et al. 2018; Flach et al. 2018; De Kauwe et al. 2019) although some limitations inevitably remain (*e.g.* lack of energy closure, see

35 Wilson et al, 2002). Although the length of the temporal records varies across sites, some sites extend back several decades allowing estimates of the impact of natural variability and climate trends on carbon, energy and water fluxes to be examined.

From each FLUXNET site, measurements of the exchange of latent heat flux (Q_{ie}), sensible heat flux (Q_b) and net ecosystem exchange (NEE) are available at 30- to 60-minute resolution, alongside meteorological variables (including air temperature, net radiation, precipitation and relative humidity). By providing simultaneous and co-located measurements of both the meteorological forcing of the surface, and the associated turbulent energy fluxes, FLUXNET provides a critical resource for understanding ecosystem responses to temperature extremes and, for the development, evaluation and benchmarking of land surface models. Importantly, the scale of recorded flux measurements (~km²) is directly relevant for evaluating land surface

Andy Pitman 26/2/2019 9:40 AM Deleted: model

Andy Pitman 27/2/2019 2:39 PM Deleted: 8

Andy Pitman 20/2/2019 1:43 PM Deleted: high-frequency (30 or 60 minute)

Andy Pitman 26/2/2019 9:42 AN Deleted: T

schemes used in CMIP type climate models (e.g. Krinner et al., 2005; Abramowitz et al., 2008; Blyth et al., 2011). As a result, land surface modellers routinely use these data to parameterise and evaluate models for extreme conditions. For example, van Gorsel et al. (2016) synthesised eddy-covariance data from seven Australian sites alongside a land surface model, to investigate the impact of heat extremes on the exchange of carbon and water fluxes during the record-breaking

- 5 heatwave in 2012/2013. They found that water-limited woodlands and energy-limited forest ecosystems responded differently to the heatwave, with the forests showing greater resilience to short-term heat than the woodlands. <u>Ukkola et al.</u> (2016) used FLUXNET data to show systematic errors in how well models captured land-atmosphere feedbacks during periods of water stress as a landscape transitioned into drought. In general, as the land surface dries, the surface energy balance tends to partition available energy increasingly towards Q_h and less towards Q_{le}, which has important implications
- 10 for atmospheric temperature, moisture and atmospheric boundary layer depth (Seneviratne et al., 2010). This understanding of land-atmosphere processes was used by Miralles et al. (2014) to link soil desiccation to the amplification of extreme heatwaves via land surface feedbacks.
- While eddy-covariance data have been widely used to examine the impact of temperature extremes, the measurement of15temperature and the measurement of Q_{le} , Q_h and NEE are independent in terms of the instrumentation used. However, the15measured temperature is provided in published data, along with measurements for the site of net radiation, wind speed,humidity etc alongside measurements of Q_{le} , Q_h and NEE. A land surface modeller requires all these data to drive a landsurface model for evaluation or process-based studies. We are therefore interested in the relationship between measurementsof temperature, and in particular extreme temperatures, and concurrent measurements of Q_{le} , Q_h and NEE. Our aim is to
- 20 characterise, for example, whether direct observations of Q_{le}, Q_h and NEE are biased towards the temperature mean and lacking at the tails of the temperature distribution, or whether they are biased to one tail of the distribution. If biases exist, is this true for all FLUXNET sites, or are there specific regions or climates where the tails of the temperature distribution are
 rich with measurements of Q_{le}, Q_h and NEE? We use measurements of temperature and Q_{le}, Q_h and NEE from FLUXNET sites because they provide co-located measurements of meteorological variables and land surface fluxes. We seek to identify
- 25 those sites with data useful to explore land surface processes under extreme temperature conditions, and potentially those sites with the meteorological forcing measured concurrently with the fluxes required to drive LSMs, We therefore do not blend the measured fluxes with meteorological observations taken elsewhere to ensure the land surface fluxes are fully representative of the concurrent meteorological conditions.
- 30 Our goal is to identify those FLUXNET sites with data useful to explore land surface processes under extreme temperature conditions. We therefore first investigate which parts of the temperature distribution have simultaneous measurements of Q_{le}, Q_h and NEE for a given site. We then aggregate the answers to this question to ask which sites contain the most measured Q_{le}, Q_h and NEE relative to measured temperatures. This question is posed separately for the flux measurements over the whole temperature distribution and for the upper and lower tails of the distribution. We therefore seek to identify which
- 35 FLUXNET site data are most suitable for analysing processes under extreme temperature conditions with the goal of identifying those sites most useful for land surface model development and evaluation of the surface energy, water and carbon budgets during extreme temperatures.

2 Methods

2.1 FLUXNET data

40 We use 165 site-based data sets from the FLUXNET2015 (November 2016 release; http://fluxnet.fluxdata.org/data/fluxnet2015-dataset/) and an additional 51 data sets from the FLUXNET La Thuile

Andy Pitman 26/2/2019 10:12 AM Moved (insertion) [2]

Andy Pitman 26/2/2019 10:12 AM Deleted: thisFor example, the land surface is important in simulating Andy Pitman 26/2/2019 10:12 AM Moved up [2]: Ukkola et al. (2016) used FLUXNET data to show systematic errors in how well models captured this

Andy Pitman 26/2/2019 10:12 AM

Deleted: s

Andy Pitman 26/2/2019 10:12 AM **Deleted:** transition of the vegetation during periods of water stress.

Andy Pitman 20/2/2019 12:42 PM **Deleted:** In this study, we focus on

Deleted. In this study, we locus on

Andy Pitman 27/2/2019 2:35 PM Deleted:

Anna Ukkola 26/2/2019 3:02 PM Deleted: ultimately we seek to identify those sites useful for developing, evaluating and benchmarking LSMs. To do this requires the LSM to be provided meteorological forcing measured concurrently with the fluxes

Anna Ukkola 26/2/2019 3:02 PM Deleted: c Anna Ukkola 26/2/2019 3:02 PM Deleted: an

Andy Pitman 27/2/2019 2:35 PM **Deleted:** Andy Pitman 20/2/2019 1:50 PM

Deleted: Here

Andy Pitman 20/2/2019 1:50 PM

Deleted: , w

Andy Pitman 21/2/2019 10:41 AM

Andy Pitman 21/2/2019 10:39 AM

Deleted: . In particular, we

Andy Pitman 21/2/2019 10:39 AM

Deleted: y Andy Pitman 21/2/2019 10:40 AM

Deleted: with the most observations of Q_{le}, Q_h and NEE since these would

Andy Pitman 21/2/2019 10:40 AM

Deleted: enable

Andy Pitman 21/2/2019 10:41 AM **Deleted:** elements of

(http://fluxnet.fluxdata.org/data/la-thuile-dataset/) data release. <u>Only freely-available site datasets from each release were</u> <u>used.</u> Overall, our analysis is therefore based on 216 different site data sets. A list of all sites used and associated information including vegetation type, location, the period of observations and references are provided in Supplementary Table 4. The data were pre-processed using the FluxnetLSM package (Ukkola et al., 2017). Variables LE F MDS, H F MDS and

- 5 NEE_VUT_REF and TA_F_MDS were used from FLUXNET2015 for Q_{le}, Q_h, NEE and air temperature respectively and LE_f, H_f, NEE_f and Ta_f from La Thuile. These variables were accompanied by quality control (QC) flags to indicate whether the data were observed or gap-filled. These QC flags facilitate the selection of data based on measurement quality. In this study, we focus only on the observed data, which is marked by the quality control flag 0 and exclude all other data.
- To be representative a site requires a reasonable sample of measured data. We therefore first excluded any FLUXNET and La Thuile sites with less than 8 months of observed data. We also excluded any sites with less than 50% of the temperature data having been measured (i.e. QC = 0) as distinct from gap filled or missing data (this excluded 14 sites). We also tested the sensitivity of our conclusions to data length. Given our focus on Qle, Qh and NEE, we excluded night-time data using two criteria. We first excluded all data between 11 pm and 6 am local time. In addition, if shortwave radiation was < 1 W m₂⁻² for an individual time period then associated measurements were also excluded. This did not exclude many measurements as
- shortwave radiation was rarely reported as non-zero at night but there were occasional shortwave radiation > 1 W m_{h}^{-2} in observations at night. Thus, discussion of the availability of measured fluxes at the lower tail of the temperature distribution focuses on daytime minimum temperatures. Overall, temperature observations were available 86% of the time, $Q_{le_{h}}$ 62% of the time, Q_{b} 68% of the time and NEE 30% of the time.
- 20

We examined the availability of measured temperature relative to the potential availability after we excluded sites with less than 8 months of data, and less than 50% of data being measured, and nighttime. We note 88% of all sites reported measurements for more than 80% of the time. Only 6% of sites had measurements for 50%-70% of the time and we excluded sites with less than 50% from subsequent analysis.

25

2.2 Data processing

For each site, we first determine which time steps have measurements of temperature. If an observation of temperature is available (i.e. QC = 0) we explore if, for this same time step₂ there are measurements of Q_{le} , Q_h and NEE with a QC flag of 0. We then calculate the ratio of the number of measurements of each of the three fluxes relative to the number of temperature measurements. For each site, this ratio was first calculated over the whole temperature distribution. Thus, per flux, the total number of measurements for Q_{le} , Q_h and NEE were each divided by the total number of measured temperatures. In addition, this ratio was calculated for only the temperatures in the highest 2.275% of the temperature distribution, and separately for the lowest 2.275% of temperatures. These ranges approximate the data above and below two standard deviations from the mean. We did repeat our analysis using exactly the two standard deviations; this led to some

35 qualitative differences in our results because some sites lack enough measurements to provide reliable results where the temperature distribution was not normally distributed.

3 Results

Figure 1a shows the normalised frequency distribution of temperature, aggregated over all sites. Values range from about -40° C to 40° C and are approximately normally distributed. However, the upper tail ends more abruptly than the lower tail. Figure 1a also shows the normalised frequency of Q_{le} , Q_h and NEE for different values of temperature. The shapes of the

4

Andy Pitman 20/2/2019 1:49 PM

Deleted: Our goal is to identify those FLUXNET sites with data useful to explore land surface processes under extreme temperature conditions.

Andy Pitman 21/2/2019 10:53 AM Formatted: Superscript

Andy Pitman 21/2/2019 10:53 AM Formatted: Superscript Andy Pitman 21/2/2019 10:53 AM Deleted: Given our focus on Q_{les} Q_h and NEE we excluded night-time data where incoming shortwave

radiation was $< 1 \text{ W m}^2$. In case of erroneous measurements of shortwave radiation at night we excluded all data between 11 pm and 6 am.

Anna Ukkola 26/2/2019 3:06 PM Deleted: e distributions for Q_{le} and Q_{h} are similar and measurements exist across the entire range of sampled temperatures. Not surprisingly, the normalised frequency of measurements for both Q_{le} and Q_{h} are lower than for measured temperature. Notably, the frequency of NEE is much lower than for Q_{le} and Q_{h} . Figure 1b shows the ratio of the number of measurements of Q_{le} , Q_{h} and NEE relative to the number of measurements of temperature. In all cases, the ratios increase as a function of

- 5 increasing temperature, indicating that fluxes are better sampled for warmer than colder temperatures. At the lowest temperatures, ratios for Q_{lc} and Q_{h} range from ~0 to ~0.3 but these increase as temperatures increase to maximum ratios of ~0.8 at around 20°C and remain at ~0.8 through to 30°C. For NEE ratios increase to ~0.6 at around 35°C for NEE. A minor dip in Q_{lc} and Q_{h} ratios occurs at 0°C associated with the phase change of water, which most likely affects the operation of instrumentation. At the upper extreme of the temperature distribution, ratios decline between 30°C and 45°C from ~0.8 to
- 10 ~0.6 for Q_{le} and Q_h and from ~0.6 to ~0.5 for NEE. However, in each case a secondary peak of high ratios occurs for the very highest temperatures. This peak is associated with temperatures > 44°C, which are rare and associated with measurements at Au-Cpr (there are only 68 individual measurements in excess of 44°C at this site), AU-GWW (23 individual measurements), AU-Stp (24 individual measurements) and SN-Dhr (33 individual measurements). Of these, the Australian sites tend to have high measurement ratios and this peak at very high temperatures almost entirely reflects observations from Australian sites.
 15 Figure 1b highlights where there are less than 1000 measurements in an individual bin and as expected they occur at the upper and lower tails of the distribution.

Figure 2 shows the geographic distribution of measurement ratios for Q_{le} (Figure 6 provides the actual ratio values associated with each site and temperature range). The ratio over the whole temperature distribution shows most sites (63%) exceed 0.7 and some sites (5%) exceed 0.9 (Figure 2a). These ratios drop considerably if the lower tail (Figure 2b) is examined. Since the lower tail is calculated for each site independently this result is not surprising for mid- and high-latitude sites where

- snow, freezing and frosts would affect measurements. However, this result is more surprising in southern Europe and south-eastern Australia where the lower tail is warm relative to some sites with higher ratios that are colder (e.g. Japan, northern China, Scandinavia). In contrast, for the upper tail, Figure 1c shows many (67) sites with ratios exceeding 0.9 (see also
 Figure 6). While we focus on the US, Europe and Australia, we note sites in Japan, China, South America and Russia with
- ratios exceeding 0.9. We also note few sites with measurement ratios > 0.8 over some regions with very high temperatures, including Africa and the Middle East, and no sites in e.g. India, Pakistan and Greece. Figure 3 shows a broadly similar result for Q_h although overall the ratios are higher (on average 0.79) than for Q_{le} (on average 0.73). This is most apparent for the upper tail (Figure 3c) where many of the sites with ratios of 0.8-0.9 for Q_{le} are above 0.9 for Q_h .
- 30

20

Figure 4 shows the geographic distribution of measurement ratios for NEE (see also Figure 8). There is a sharp contrast with the maps of Q_{le} (Figure 2) and Q_h (Figure 3) and the overall average is 0.42 compared to 0.79 for Q_h and 0.73 for Q_{le} . In terms of the overall metric (Figure 4a), no sites exist with a ratio exceeding 0.9, only one exceeds 0.8 but 18 exceed 0.7. Two sites exceed 0.7 for the lower tail (Figure 4b) located in the eastern US (US-Orv, US-Wi0). Multiple sites (11) over North

35 America exceed 0.9 for NEE at the upper tail of temperatures (Figure 4c) together with isolated sites over Europe (IT-Tor, ES-Ln2), China (CN-HaM, CN-Cha, CN-Dan) and Australia (AU-Ade).

To examine these results further, Figure 5 shows the measurement ratios as a function of mean annual precipitation and mean annual temperature. Note the amounts of rainfall shown in Figure 5 are accumulated only over times when temperature

40 data are selected and therefore cannot be compared with observations taken at meteorological stations. Figure 5 shows little relationship between temperature or rainfall and the measurement ratios. For example, some cool dry sites have high measurement ratios whereas others have low ratios. Similarly, some hot wet sites have high and some have low ratios for both Q_{le} and Q_h and for the upper tail of NEE. Few sites have high ratios for the overall temperature distribution or for the Andy Pitman 21/2/2019 11:05 AM Deleted: , Andy Pitman 21/2/2019 11:05 AM Deleted: Q Andy Pitman 21/2/2019 11:05 AM Deleted: and NEE Andy Pitman 21/2/2019 11:05 AM Deleted: for Q_k and Q_h Andy Pitman 21/2/2019 11:05 AM Deleted: and Andy Pitman 21/2/2019 11:06 AM Deleted: 0 Andy Pitman 21/2/2019 11:06 AM Deleted: 0

Andy Pitman 21/2/2019 11:15 AM **Deleted:** Since only a sub-set of Andy Pitman 21/2/2019 11:15 AM **Deleted:** and associated rainfall data meet the quality control criteria, the rainfall shown in Figure 5 Andy Pitman 21/2/2019 11:13 AM **Deleted:** standard Andy Pitman 21/2/2019 11:13 AM

Deleted: observations at these locations



lower tail of NEE. In other words, the temperature or rainfall at a specific FLUXNET sites does not explain why some sites have a high frequency of flux measurements while other sites rarely observe Q_{le} , Q_{h} , and NEE. For Q_{le} , Q_{h} , and NEE, Figure 5 also shows the lack of high ratios for the lower tail relative to the upper tail and the low ratios for NEE compared to Q_{le} and Q_{h} . At the upper tail, many sites (e.g. AU-Cpr, DE-Akm and US-NR1) exceed measurement ratios of > 0.9 for Q_{le} and Q_{h} . Overall, Figure 5 shows 5-10 sites with high measurement ratios at temperatures above ~25°C for the upper tail and for

5

Q_{lex} Q_h, and (to a lesser degree) for NEE; these are predominantly FLUXNET sites located over Australia,

We finally aggregate our analyses for the overall ratio, the lower tail and the upper tail separately for Q_{le} , Q_h and NEE (Figures 6-8) and we identify each FLUXNET site in terms of the measurement ratio. Figures 6-8 are then combined in

- Figure 9 to highlight those sites with high measurement ratios for all of Q_{le}, Q_h and NEE and for just Q_{le} and Q_h for the overall metric (Figure 9a), the lower tail (Figure 9b) and the upper tail (Figure 9c). Taking the overall statistic first (Figure 9a, additional details are listed in Supplementary Table 1), no sites are found with Q_{le}, Q_h and NEE ratios exceeding 0.9. Only two sites, both in the US (US-Whs, US-WiO) have measurement ratios above 0.8. If NEE is omitted, 19 sites are selected where both Q_{le} and Q_h ratios exceed 0.9 (Figure 9a, listed in Supplementary Table 1). These include eight sites over
- 15 the US, four sites over Australia, two over China, and a single site from Denmark, Germany, France, Italy, and Portugal. Even if the threshold is reduced to only Q_{le} and Q_h ratios exceeding 0.8, still no sites over South America, Africa and perhaps critically, for high temperatures Central America and the majority of the Mediterranean region. The freely available FLUXNET datasets provides no data over India, Pakistan or the Gulf States.
- 20 If we are interested in the lower tail of temperatures and we seek sites with measurement ratios exceeding 0.8 for each of Q_{le}, Q_h and NEE, we have two choices (US-Orv, US-Wi0). If only Q_{le} and Q_h are needed, the choice widens to 18 sites with seven sites in Australia, four in the US, one in each of China, Canada and France (Figure 9b, Supplementary Table 2). Here, we note that very cold regions are poorly sampled with no sites in Alaska, Russia, the Himalaya, Greenland or Antarctica.
- At the upper tail, 16 sites have ratios exceeding 0.9 for each of Q_{le}, Q_h and NEE and are in Canada (7), the US (6), China (3), Spain (1), Australia (1) and Italy (1) (Figure 7c, Supplementary Table 3). If only Q_{le} and Q_h are required above 0.9 there are many sites (32) and above 0.8 there are three sites in South America, one in Botswana, several in the southern US and southern Europe and one in Israel. No sites remain in India, Pakistan, the Gulf States, Central America and the majority of the Mediterranean region.
- 30

We also examined whether the measurement ratio varied by time of day for each site (Supplementary Figure 2). These examples are provided to illustrate individual site behaviour and to emphasise that major variations at each site are present. At Au-ASM, a weak diurnal cycle is visible in the measurement ratio with very similar and consistently high ratios of Q_{le}, Q_h and NEE being slightly lower. At a second Australian site, AU-Tum measurement ratios increase from dawn through the

- 35 day, and then drop off just before dusk. At CA-NS4 behaviour is similar to AU-ASM until late in the day when the measurement ratios drop sharply. At DE-Hai there is little variation though the day and Q_b is much higher than Q_{bc}, and only NEE shows any diurnal variation. DE-Meh shows Q_{bc} and Q_b are consistent through the day and are almost identical. DK-NuF shows Q_{bc} and Q_b falling from dawn to around 10am, then stabilising at low value (~0.3-0.4) and then increasing strongly from 2pm to ratios ≥ 0.7 while NEE increases weakly from ~0.2 gradually though the day. It-Tor shows little
- diurnal variation in Q_{lc} and Q_b, but there is a strong diurnal variation in NEE. Finally, US-Whs shows high measurement ratios for Q_b and Q_{lc}, but falling slightly through the day with NEE increasing strongly from dawn to 11am, and then slowly declining through the day. If we assume that the hottest part of the day is around 1pm, those sites that provide useful observations of Q_b and Q_{lc} coincident with these temperatures clearly require site-by-site evaluation. Thus, if sites are being

6

Deleted: For Q_{le} , Q_h , and NEE, Figure 5 shows the lack of high ratios for the lower tail relative to the upper tail and the low ratios for NEE compared to Q_k and Q_h . At the upper tail, many sites (e.g. AU-Cpr, DE-Akm and US-NR1) exceed measurement ratios of > 0.9 for Q_{le} and Q_h , but for NEE, most are in cool sites with low precipitation. In addition

Andy Pitman 21/2/2019 9:58 AM

Deleted: . Figure 5 shows little relationship between temperature or rainfall and the measurement ratios. For example, some cool dry sites have high measurement ratios whereas others have low ratios. Similarly, some hot wet sites have high and some have low ratios for both Qle and Qh and for the upper tail of NEE. Few sites have high ratios for the overall temperature distribution or for the lower tail of NEE. In other words, the temperature or rainfall at a specific FLUXNET sites does not explain why some sites have a high frequency of flux measurements while other sites rarely observe Q_{le} , Q_h , and NEE. In figure 5, we observe 5-10 sites with high measurement ratios at temperatures above ~25°C for the upper tail and for Q_{le}, Q_h, and (to a lesser degree) for NEE; these are predominantly FLUXNET sites located over Australia.

Andy Pitman 21/2/2019 10:19 AN Moved (insertion) [1] composited, the knowledge that different sites sample different parts of the diurnal cycle, and sample Q_{le}, Q_b and NEE differently across the diurnal cycle needs to be taken into account.

4 Discussion

- 5 The FLUXNET eddy covariance flux measurements are among the most valuable observations available for <u>understanding</u> processes, and for developing, evaluating and benchmarking land surface models. Under future climate change, warming driven by radiative forcing is likely to amplify by changes in the partitioning of available energy between latent and sensible heat at the surface (e.g. Seneviratne et al., 2010; Miralles et al., 2014; Donat et al., 2018; Ukkola et al., 2018). This change in the partitioning, linked with soil desiccation or changes in stomatal conductance under higher CO₂ provides an amplification
- 10 of the large-scale meteorology and can lead to more extreme conditions via the coupled land-boundary layer system (Seneviratne et al., 2010; Miralles et al., 2014). As the continental surface warms, some regions will experience temperatures beyond the historical record. Building land models for CMIP-type climate models that properly capture mechanisms and processes occurring in a region experiencing higher temperatures is helped if observations from other regions already experiencing those temperatures are available (so called climate analogues, or space for time substitutions). In this context,
- 15 observations from FLUXNET are particularly valuable if they sample existing hot locations, and if they actually measure fluxes at those locations at the upper tail of temperature.

Our results highlight multiple positives for those wishing to probe vegetation responses to temperate extremes and/or evaluate land surface models. Figure 9 shows many sites with high measurement ratios for Q_{le} and Q_h at the upper and lower 20 tail, indicating a rich source of available observations. Conversely, if we seek observations of Q_{le}, Q_h and NEE, these data are more limited with only two sites with a measurement ratio > 0.8, and none > 0.9 at the lower tail and 16 sites at the upper tail

- (see Supplementary Tables 2 and 3). Of course, the > 0.9 measurement ratio is arbitrary and more sites become available at lower ratios; however, it is somewhat confronting that at > 0.8, 87% of the sites in Supplementary Table 2 are located in Europe, North America and Australia and for the upper tail, 88% of the sites in Supplementary Table 3 are located in these
 three regions. The non-Europe, North America and Australia sites are not distributed globally: Figure 9 shows virtually no
- sites with high (> 0.8) measurement ratios in the tropics, Africa or South America for Q_{le} and Q_{h} , and no sites at all in India or the Gulf states. These typically hot regions may be surrogates for how continental surfaces behave under future climate scenarios in the mid-latitudes and it is unfortunate that FLUXNET lacks observations in these regions.
- 30 In the absence of measurements from hot regions, the availability of observations from Australia becomes particularly important because these sites cover a wide rainfall gradient, ranging from water through to energy limited sites. We note two possible reasons for the lack of freely available data in many regions. First, there may be a lack of sites, or sites that exist, may have low measurement ratios. Second, the high number of sites identified in our analysis with high measurement ratios located in Europe, North America and Australia largely reflects the high number of sites in the FLUXNET data. Similarly,
- 35 the low number of sites in Africa, South America, India and the Gulf States reflects the rarity of FLUXNET sites in these regions. There are, however, four sites in Africa, three in South America and one in Israel in FLUXNET, but these are excluded due to the shortness of the data record, and the low temperature measurement ratios. This is not intended as a criticism; it is a consequence of history (where groups grew with the capacity to maintain measurements and the common desire to run measurement sites near home institutions).

40

One result from our analysis is that overall, measurement ratios for Q_h are higher than for Q_{le} and both of these are much higher than NEE. This is true for the overall distribution of temperatures, and for the lower and upper tails of the distribution. This result can be quickly visualised by comparing Figures 6, 7 and 8. In part, this is associated with the actual temperatures at the sites influencing the measurement ratios once aggregated. Figure 5 shows that the measurement ratios are generally

- 5 lower at the lower tail than the higher tail for Q_{le} and Q_{h} . Furthermore, for the lower tail, the ratios are generally lower at colder temperatures than warmer temperatures. We propose multiple reasons explaining these findings Q_{h} , Q_{le} and NEE are all products of turbulent transport. While there have been significant improvements in instrumentation over the last 20 years, measurements of these fluxes over long periods and across a range of weather conditions remains challenging.
- 10 Measurement ratios of < 1 for Q_h, Q_{le} and NEE are expected due to data loss caused by instrument failure, precipitation, ambient conditions that violate the assumptions of the eddy covariance method (particularly low- or non-stationary turbulence) and other artefacts (Foken et al 2010; Burba 2013). The lower ratios for Q_{le} in comparison to Q_h are likely to be associated with measurement methods. The majority of sites use a sonic anemometer and an open-path gas analyser to measure Q_h, Q_{le} and NEE. Both devices use measurement techniques over a physical path (sound waves for the sonic and 15 infra-red for the open-path gas analyser). Anything that partially obscures the measurement path (condensation, mist, drizzle, snow, ice, etc) can interfere with the measurements. The sonic anemometers are robust to all but very intense rain but the

open-path gas analysers are more sensitive to anything that blocks the optical path (Foken et al., 2010). The Q_h measurements only involve the sonic anemometer while Q_{le} and NEE use measurements from the sonic (for vertical velocity component) and from the open-path gas analyser (for water and CO₂ concentration). Measurements for Q_{le} and NEE are

 $20 \quad \text{therefore inherently more complex than for Q_h, which explains the lower measurement ratio for Q_{le} relative to Q_{h}.}$

The lower ratios at lower temperatures are likely to be associated with the occurrence of condensation (dew), which is more common at cooler temperatures; hence the observed dependence of the ratio on measured air temperature. However, the assumptions underpinning the measurement of surface fluxes using the eddy covariance method are violated in low turbulence conditions, which occurs mostly at night (excluded in our analysis) and low temperatures (e.g. at dawn where

- 25 turbulence conditions, which occurs mostly at night (excluded in our analysis) and low temperatures (e.g. at dawn where radiative cooling leads to a stable surface layer). For fluxes that are significantly different from 0 at night (e.g. NEE due to ecosystem respiration) this leads to an overwhelming bias in the measurements unless low turbulence conditions, where the assumptions of the eddy covariance method fail, are excluded from the analysis. Therefore, friction velocity (u*) is used as a proxy for turbulence, by finding the site-specific value for u* above which NEE is independent of u* and removing all
- 30 observations when u* is below this threshold (Aubinet et al., 2012). This often results in less than 20% of NEE data being available for estimating ecosystem respiration. The application of this turbulence filter causes the ratio for NEE to be much lower than the ratio for Q_h and Q_{le}. The occurrence of these conditions is more likely in lower temperature conditions, contributing to the slope in Figure 1b. We avoid the consequences of these procedures in quality controlling and gap-filling data by only using those data that are directly observed.

35

Our analysis has a specific weakness, which requires consideration when interpreting our results. There may be a temptation to interpret the ratios we report as a metric linked with measurement quality. To discourage such a temptation we draw attention to two hypothetical FLUXNET sites, one with ratios around 0.9 and another around 0.3. In the former, the efforts around measurement quality are superficial and data are included unless a specific problem identified. At the latter, the

40 efforts around measurement quality are rigorous and any doubts whatsoever about the data leads to it being discarded. For the latter case, one would suggest that the resulting data reported to the FLUXNET2015 or La Thuile archive are likely of the highest quality and most reliable to use in process-level examination of models or understanding of the surface energy and carbon balance. The more complete data in the former example could in fact be misleading. In short, our analysis does

8

Andy Pitman 19/2/2019 3:34 PM

Deleted: First,

not report on data quality, it only relates to coincident data availability and identifies those sites where measurements are available with high frequency and with a QC = 0.

Our methodology contained several assumptions, for example we excluded sites with less than 8 months of data. We tested

- 5 the sensitivity to this assumption, examining whether the sites identified with high measurement ratios changed if we required 12 months of data. If we set a minimum length of record as 12 months, US-Wi0 (one of two sites with Q_{le}, Q_h and NEE > 0.8), US-SP1, US-Orv and ES-Ln2 are excluded in Supplementary Table 1. The only sites with Q_{le}, Q_h and NEE > 0.8 are excluded from the lower tail (US-Orv and US-Wi0), along with DK-Fou, US-SP1, and NL-Lan. At the upper tail multiple sites (AU-Rob, PT-Mi1, NL-Lan, Es-Ln2, US-Wi0, US-SP1 and US-Bar) are excluded. Therefore, requiring a 12-
- 10 month data set has a significant impact on some of the otherwise most useful sites. Given the purpose of our analysis is to examine the tails of the distributions at each site, we suggest that imposing longer measurement periods than absolutely required may prove counterproductive. In addition, we examined two other attributes of the FLUXNET data – whether our measurement ratio changes between the first half of the data and the second half (i.e. to examine whether the measurement ratio improved over time) and whether any relationship exists between the total number of QC=0 observations and the
- 15 measurement ratio. The first analysis found no evidence that higher measurement ratios were apparent in the first or second half of the data, something that might have been expected if the ability to sustain measurements improved over time. The second analysis also found no evidence of a relationship between the measurement ratio and the length of data (Supplementary Figure 1).
- 20 One obvious criticism of our measurement ratio metric is the temptation to interpret the results as a way to select FLUXNET sites for model development and evaluation without further thought. Clearly, a high measurement ratio is only one aspect of a valuable data set. A modeller might, for example, prefer a large number of actual measurements with a low overall measurement ratio rather than a site with few measurements but a high overall measurement ratio. We have noted above that we find no correlation between data length and measurement ratio but some sites (see Supplementary Tables 1-3) have both
- 25 high measurement ratios and large amounts of data and others have high measurement ratios and low amounts of data. For example, the two sites with the highest measurement ratios overall (US-Whs and US-Wi0) sharply contrast on the amount of data (63,619 and 4621 temperature measurements respectively). In this case, US-Whs covers 2922 days of measurement and 93% of the time temperature data are reported (Supplementary Table 1), whereas US-Wh0 only measures for 365 days and only 62% of the time temperature data are reported. In contrast, sites such as CA-NS1 and CA-NS3 display very similar
- 30 measurement ratios for Q_{le}, Q_h and NEE, both cover 1826 days but CA-NS1 includes 30,269 temperature measurements while CA-NS3 includes only 22,689 temperature measurements. Clearly, many characteristics of a data set make it valuable for model development or model evaluation and our analysis should be viewed only as one of these characteristics. One way forward to resolve how to choose FLUXNET data for extremes is to combine an analysis of meteorological sites with FLUXNET sites. Using sites maintained by meteorological agencies to identify extreme events (e.g. heatwaves) and then
- 35 interrogate the FLUXNET sites near to the meteorological site for the availability of measurements of Q_{le}, Q_h and NEE could enable a modeller to choose suitable sites for land surface model development and evaluation. <u>While one possible way</u> forward, inconsistencies between observations from meteorological agencies relative to FLUXNET (location, geographical distribution, height of measurements, standardisation of measurements over short grass) highlight the challenges in using meteorological observations that are physically separate from the FLUXNET observations.

40

Our analysis poses interesting questions about the FLUXNET data that deserve further exploration. Why do sites with a similar climate vary so greatly in terms of their frequency of reporting of Q_{le}, Q_h and NEE in comparison to temperature?

9

Andy Pitman 19/2/2019 3:35 PM **Deleted:** are excluded

Andy Pitman 21/2/2019 10:19 AM Deleted:

Andv Pitman 21/2/2019 10:19 AM Moved up [1]: We also examined whether the measurement ratio varied by time of day for each site (Supplementary Figure 2). These examples are provided to illustrate individual site behaviour and to emphasise that major variations at each site are present. At Au-ASM, a weak diurnal cycle is visible rement ratio with very similar and in the meas consistently high ratios of Qle, Qh and NEE being slightly lower. At a second Australian site, AU-Tum measurement ratios increase from dawn through the day, and then drop off just before dusk. At CA-NS4 behaviour is similar to AU-ASM until late in the day when the measurement ratios drop sharply. At DE-Hai there is little variation though the day and Qh is much higher than Q_{le} , and only NEE shows any diurnal variation. DE-Meh shows Q_{le} and Q_h are consistent through the day and are almost identical. DK-NuF shows Q_{le} and Q_h falling from dawn to around 10am, then stabilising at low value (~0.3-0.4) and then increasing strongly from 2pm to ratios 0.7 while NEE increases weakly from ~0.2 gradually though the day. It-Tor shows little diurnal variation in Q_{le} and Q_{h} , but there is a strong diurnal variation in NEE. Finally, US-Whs shows high measurement ratios for Qh and Qle, but falling slightly through the day with NEE increasing strongly from dawn to 11am, and then slowly declining through the day. If we assume that the hottest part of the day is around 1pm, those sites that provide useful observations of Qh and Qle coincident with these temperatures clearly require site-by-site evaluation. Thus, if sites are being composited, the knowledge that different sites sample different parts of the diurnal cycle, and ample Qle, Qh and NEE differently across the diurnal cycle needs to be taken into account Andy Pitman 21/2/2019 10:20 AN

Formatted: Justified

Why are some sites able to do this routinely while others cannot, and can expertise be shared to resolve this? What are the implications of aggregating FLUXNET data given the large variations in which parts of the temperature distribution are sampled? Why are there major variations in the measurement ratios between sites over the diurnal cycle and what does this mean in terms of using site data from FLUXNET? Clearly, the FLUXNET data do provide our best ecosystem-scale estimate

- of the vegetation's response to heat extremes (Ciais et al. 2005; Teuling et al. 2010; Wolf et al. 2013; von Buttlar et al. 2018;
 Flach et al. 2018; De Kauwe et al. 2012) but given the need to build land models representing extreme conditions these data cannot be used without further evaluation of the specific site data. We do not know if there are opportunities for the global community to prioritise new sites in regions that currently lack data, or directly support those measurements in regions with low measurement ratios. However, we suggest value in either new sites, or investment in existing sites, in countries that
- 10 experience temperatures that are higher than those experienced across North America and Europe to enable land models to be developed in anticipation of further warming. <u>Virtually all sites (~90%) with high measurement metrics for Q_{le}. Q_h and NEE, or just Q_{le} and Q_{ha} whether examining the whole distribution or just the lower tail or just the upper tail, are located in North America, Western Europe and Australia. There are no sites in India, South America, Africa, the Middle East and few sites in China. In terms of vulnerability, the freely available FLUXNET data therefore cover regions representing 12-14% of</u>
- 15 the global population. Indeed, the poorest country with measurements (based on Gross Domestic Product, Portugal) suggests all countries ranked from Portugal (ranked 47^{th}) to the poorest country (ranked 211^{th}) lack any measurements. Another perspective is if countries are ranked on average temperature, none of the warmest 98 countries contain a site and Australia is the hottest country with sites with high measurement ratios. Conversely, North America, Western Europe and Australia have multiple sites with observations of Q_{te} and Q_{h} and some with NEE with high measurement ratios for both the lower and
- upper tail of the temperature distribution. For these three regions therefore, FLUXNET data provide a rich source of data for understanding how fluxes of energy, water and carbon behave under extreme temperature conditions. Overall, we have noted more frequent observations of Q_h than Q_{lc} and both these fluxes are much more common than NEE. An implication of this is that some regions, particularly very hot regions that will be first to experience novel climates require observations. We also highlight a wide discrepancy between the measurement ratios across FLUXNET sites that is not related to the actual temperature or rainfall at the site.

5. Conclusions

We have examined the FLUXNET data by evaluating the availability of Q_{le} , Q_h and NEE observations at time steps where temperature is measured (with a quality control flag QC = 0). We have analysed this spatially to identify those sites with a high availability of flux measurements, relative to temperature measurements, across the whole temperature distribution, and

at the upper and lower tails of the distribution.

Virtually all sites (~90%) with high measurement metrics for Q_{le}, Q_h and NEE, or just Q_{le} and Q_h, whether examining the whole distribution or just the lower tail or just the upper tail, are located in North America, Western Europe and Australia.

There are no sites in India, South America, Africa, the Middle East and few sites in China. <u>This</u> discrepancy between the measurement ratios across FLUXNET sites is not related to the actual temperature or rainfall at the site. Clearly, some sites seem able to retrieve Q_{le}, Q_h and NEE reliably at extreme temperatures while others cannot. This may provide an opportunity for the FLUXNET community to share best practice strategies to identify ways to ensure measurements at the tails of the temperature distribution.

40

30

Deleted: 8

Andy Pitman 21/2/2019 10:21 AM

Deleted: In terms of vulnerability, the freely available FLUXNET data therefore cover regions representing 12-14% of the global population. Indeed, the poorest country with mea urements (based on Gross Domestic Product, Portugal) suggests all countries ranked from Portugal (ranked 47th) to the poorest country (ranked 211th) lack any measurements. Another perspective is if countries are ranked on average temperature, none of the warmest 98 countries contain a site and Australia is the hottest country with sites with high measurement ratios. Conversely, North America, Western Europ and Australia have multiple sites with observations of Q_{le} and Q_h and some with NEE with high measurement ratios for both the lower and upper tail of the temperature distribution. For these three regions therefore, FLUXNET data provide a rich source of data for understanding how fluxes of energy, water and carbon behave under extreme temperature conditions. Overall, we have noted more frequent observations of Qh than Qle and both these fluxes are much more common than NEE. This leads to two conclusions: first, some region particularly very hot regions that will be first to experience novel climates require observations. Less obviously, we highlight a wide

Andy Pitman 21/2/2019 10:22 AM Deleted: that

Finally, we restate a key caveat to our paper to avoid any misunderstanding. Our analysis does not highlight the "best data". A site might have high ratios because of poor QC control, or low metrics because of strict controls. However, our paper does highlight sites with frequent observations of Q_{le} , Q_h and NEE coincident with temperature observations where all have a QC = 0. A modeller might of course reject some of these sites for reasons of data record length, vegetation type, soil type or a

- 5 multitude of other reasons. However, we suggest that our analysis provides one way for modellers to identify sites from the FLUXNET archive that warrant closer scrutiny for development and evaluation of land surface models under extreme temperature conditions.
- *Author contributions.* The ideas for this study originated in discussions with all authors. SH carried out the analysis supported by all authors. The manuscript was prepared with contributions from all authors.

Code availability. All code is freely available from: https://github.com/sophievanderhorst/FLUXNET

Data availability. All eddy covariance data are available from http://fluxnet.fluxdata.org/data/fluxnet2015-dataset/ and http://fluxnet.fluxdata.org/data/la-thuile-dataset/.

Competing interests. The authors declare no competing financial interests.

Acknowledgements. AJP, MDK, AU and GA acknowledge support from the Australian Research Council Centre of
 Excellence for Climate Extremes (CE170100023). SH would like to thank Prof. Bert Holtslag of Wageningen University for
 his comments on the manuscript and his help in arranging the internship. This work used eddy covariance data acquired by
 the FLUXNET community and in particular by the following networks: AmeriFlux (U.S. Department of Energy, Biological
 and Environmental Research, Terrestrial Carbon Program (DE–FG02–04ER63917 and DE–FG02–04ER63911)), AfriFlux,
 AsiaFlux, CarboAfrica, CarboEuropeIP, CarboItaly, CarboMont, ChinaFlux, Fluxnet–Canada (supported by CFCAS,

- 25 NSERC, BIOCAP, Environment Canada, and NRCan), GreenGrass, KoFlux, LBA, NECC, OzFlux, TCOS–Siberia, USCCC. We acknowledge the financial support to the eddy covariance data harmonization provided by CarboEuropeIP, FAO–GTOS–TCO, iLEAPS, Max Planck Institute for Biogeochemistry, National Science Foundation, University of Tuscia, Université Laval and Environment Canada and US Department of Energy and the database development and technical support from Berkeley Water Center, Lawrence Berkeley National Laboratory, Microsoft Research eScience, Oak Ridge
- 30 National Laboratory, University of California, University of Virginia.

References

15

Abramowitz, G., Leuning, R., Clark, M., and Pitman, A.: Evaluating the performance of land surface models, J. Climate, 21, 468-5481, https://doi.org/10.1175/2008JCLI2378.1, 2008.

35 Aubinet, M., Feigenwinter, C., Heinesch, B., Laffineur, Q., Papale, D., Reichstein, M., Rinne, J. and van Gorsel, E.: Nighttime Flux Correction, Chapter 5 in Eddy Covariance; A Practical Guide to Measurement and Data Analysis, Springer Atmospheric Sciences, 438, ISBN 978-94-007-2350-4, 2012.

Barriopedro, D., Fischer, E. M., Luterbacher, J., Trigo, R. M. and García-Herrera, R.: The hot summer of 2010: redrawing the temperature record map of Europe, Science, 332 (6026), 220–224, https://doi.org/10.1126/science.1201224, 2011.

40 Berry, J. A. and Björkman, O.: Photosynthetic Response and Adaptation to Temperature in Higher Plants, Annu.

Rev. Plant Physio., 31(1), 491-543, 1980.

Blyth, E., Clark, D. B., Ellis, R., Huntingford, C., Los, S., Pryor, M., Best, M., and Sitch, S.: A comprehensive set of benchmark tests for a land surface model of simultaneous fluxes of water and carbon at both the global and seasonal scale, Geosci. Model Dev., 4, 255–269, https://doi.org/10.5194/gmd-4-255-2011, 2011.

5 Burba, G.: Eddy Covariance Method for Scientific, Industrial, Agricultural and Regulatory Applications, LI-COR Biosciences, 331, 2013.

Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., et al.: Europe-wide reduction in primary productivity caused by the heat and drought in 2003, Nature, 437(7058), 529, https://doi.org/10.1038/nature039722005, 2005.

Colombo, A. F., Etkin, D. and Karney, B.W.: Climate variability and the frequency of extreme temperature events for nine
sites across Canada: implications for power usage, J. Clim., 12, 2490–2502, https://doi.org/10.1175/1520-0442(1999)012%3C2490:CVATFO%3E2.0.CO;2, 1999

Coumou, D. and Rahmstorf, S.: A decade of weather extremes, Nat. Clim. Chang., 2 (7), 491–496, https://doi.org/10.1038/nclimate1452, 2012.

De Kauwe, M. G., Medlyn, B. E., Pitman, A. J., Drake, J. E., Ukkola, A., Griebal, A., Pendall, E., Prober, S., and Roderick,
 M.: Examining the evidence for sustained transpiration during heat extremes, <u>Biogeosciences</u>, 16, 903–916, 2019, doi: 10.5194/bg-16-903-2019, 2019,

Donat, M. G., Pitman, A. J. and Angelil, O.: Understanding and reducing future uncertainty in mid-latitude heat extremes, Geophysical Research Letters, 45, 10,627-10,636, https://doi.org/10.1029/2018GL079128, 2018.

Donat, M. G., Pitman, A. J., and Seneviratne, S. I.: Regional warming of hot extremes accelerated by surface energy fluxes,
20 Geophys. Res. Lett., 44(13), 7011-7019, https://doi.org/10.1002/2017GL073733, 2017.

Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, Geosci. Model Dev., 9, 1937-1958, doi: 10.5194/gmd-9-1937-2016, 2016.

Fischer, E. M., Seneviratne, S. I., Vidale, P. L., Lüthi, D. and Schär, C.: Soil moisture–atmosphere interactions during the 2003 European summer heat wave, J. Clim, 20 (20), 5081–5099, 2007.

Flach, M., Sippel, S., Gans, F., Bastos, A., Brenning, A., Reichstein, M., and Mahecha, M. D.: Contrasting biosphere responses to hydrometeorological extremes: revisiting the 2010 western Russian heatwave, Biogeosciences, 15, 6067-6085, https://doi.org/10.5194/bg-15-6067-2018, 2018.

Foken, T., Gockede, M., Mauder, M., Mahrt, L., Amiro, B. and Munger, W.: Post-field Data Quality Control, Chapter 9 in
Handbook of Micrometeorology; A Guide for Surface Flux Measurement and Analysis, Kluwer Academic Publishers, 250, 2010.

Gunderson, C. A., O'Hara, K. H., Campion, C. M., Walker, A. V. and Edwards N. T.: Thermal plasticity of photosynthesis: the role of acclimation in forest responses to a warming climate. Glob. Change Biol., 16(8), 2272-2286, https://doi.org/10.1111/j.1365-2486.2009.02090.x, 2009.

12

Andy Pitman 27/2/2019 2:41 PM Deleted: Biogeosciences Discuss., Andy Pitman 27/2/2019 2:41 PM Deleted: https://doi.org/10.5194/bg-2018-399, in review, 2018. Kala, J., De Kauwe, M. G., Pitman, A. J., Medlyn, B. E., Wang, Y. P., Lorenz, R., and Perkins-Kirkpatrick, S. E.: Impact of the representation of stomatal conductance on model projections of heatwave intensity, Sci. Rep-UK, 6, 23418, https://doi.org/10.1038/srep234182016, 2016.

Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S. and Prentice, I.
C.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, Global Biogeochem. Cy., 19, GB1015, https://doi.org/1029/2003GB002199, 2005.

Kumarathunge, D. P., Medlyn, B. E., Drake, J. E., Tjoelker, M. G., Aspinwall, M. J., Battaglia, M., Cano, F. J, Carter, K. R.,
Cavaleri, M. A., Cernusak, L. A., Chambers, J. Q., Crous, K. Y., De Kauwe, M. G., Dillaway, D. N., Dreyer, E., Ellsworth,
D. S., Ghannoum, O., Han, Q., Hikosaka, K., Jensen, A. M., Kelly, J. W. G., Kruger, E. L., Mercado, L. M., Onoda, Y.,

10 Reich, P. B., Rogers, A., Slot, M., Smith, N. G., Tarvainen, L., Tissue, D. T., Togashi, H. F., Tribuzy, E. S., Uddling, J., Vårhammar, A., Wallin, G., Warren, J. M. and Way, D. A.: Acclimation and adaptation components of the temperature dependence of plant photosynthesis at the global scale, New Phytol., in press, 2018.

Lewis, S. L., Brando, P. M., Phillips, O. L., van der Heijden, G. M., and Nepstad, D.: The 2010 amazon drought, Science, 331(6017), 554-554, 2011.

15 Lombardozzi, D. L., Bonan, G. B., Smith, N. G., Dukes, J. S. and Fisher, R. A.: Temperature acclimation of photosynthesis and respiration: A key uncertainty in the carbon cycle-climate feedback, Geophys. Res. Lett., 42(20), 8624-8631, https://doi.org/10.1002/2015GL065934, 2015.

McEvoy, D., Ahmed, I. and Mullett, J.: The impact of the 2009 heat wave on Melbourne's critical infrastructure, Local Environ., 17, 783–796, https://doi.org/10.1080/13549839.2012.678320, 2012.

20 McMichael, A. J. and Lindgren, E.: Climate change: present and future risks to health, and necessary responses, J. Intern. Med, 270, 401–413, https://doi.org/10.1111/j.1365-2796.2011.02415.x, 2011.

Mercado, L. M., Medlyn, B. E., Huntingford, C., Oliver, R. J., Clark, D. B., Stephen, S., Przemyslaw, Z., Kattge, J., Harper, A. B. and Cox, P. M.: Large sensitivity in land carbon storage due to geographical and temporal variation in the thermal response of photosynthetic capacity, New Phytol., 218(4), 1462-1477, https://doi.org/10.1111/nph.15100, 2018.

25 Miralles, D. G., den Berg, M. V., Teuling, A. J. and Jeu, R. D.: Soil moisture temperature coupling: a multiscale observational analysis. Geophys. Res. Lett., 39 (21), https://doi.org/10.1029/2012GL053703, 2012.

Miralles, D. G., Teuling, A. J., van Heerwaarden, C. C. and de Arellano, J. V. G.: Megaheatwave temperatures due to combined soil desiccation and atmospheric heat accumulation, Nat. Geosci, 7 (5), 345–349, https://doi.org/10.1038/ngeo2141, 2014.

30 Phillips, O. L., Aragão, L. E., Lewis, S. L., Fisher, J. B., Lloyd, J., López-González, G., et al.: Drought sensitivity of the Amazon rainforest, Science, 323(5919), 1344-1347, 2009.

Schär, C., Vidale, P. L., Lüthi, D., Frei, C., Häberli, C., Liniger, M. A., and Appenzeller, C.: The role of increasing temperature variability in European summer heatwaves, Nature, 427(6972), 332, https://doi.org/10.1038/nature02300, 2004.

Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B. and Teuling, A. J.: Investigating soil moisture-climate interactions in a changing climate: A review, Earth-Sci. Rev., 99(3-4), 125-161, https://doi.org/10.1016/j.earscirev.2010.02.004, 2010.

Sillmann, J., Kharin V. V., Zhang, X., Zwiers, F. W. and Bronaugh, D.: Climate extremes indices in the CMIP5 multimodel ensemble: Part 1. Model evaluation in the present climate, J. Geophys. Res. Atmos., 118, 1716–1733, https://doi.org/10.1002/jgrd.50203, 2013.

Sippel, S., Zscheischler, J., Mahecha M. D., Orth, R., Reichstein, M., Vogel, M., and Seneviratne, S. I.: Refining multimodel projections of temperature extremes by evaluation against land-atmosphere coupling diagnostics, Earth Syst. Dynam., 8, 387–403, https://doi.org/10.5194/esd-8-387-2017, 2017.

10 Smith, N. G. and Dukes, J. S.: Plant respiration and photosynthesis in global-scale models: incorporating acclimation to temperature and CO₂, Glob. Change Bio., 19(1), 45-63, https://doi.org/10.1111/j.1365-2486.2012.02797.x, 2013.

Teuling, A. J., Seneviratne, S. I., Stöckli, R., Reichstein, M., Moors, E., Ciais, P., et al.: Contrasting response of European forest and grassland energy exchange to heatwaves, Nat. Geosci., 3(10), 722, https://doi.org/10.1038/ngeo950, 2010.

Ukkola, A. M., De Kauwe, M. G., Pitman, A. J., Best, M. J., Abramowitz, G., Haverd, V., Decker, M. and Haughton, N.:
Land surface models systematically overestimate the intensity, duration and magnitude of seasonal-scale evaporative droughts, Environ. Res. Lett., 11(10), 104012, https://doi.org/10.1088/1748-9326/11/10/104012, 2016.

Ukkola, A. M., Haughton, N., Kauwe, M. G. D., Abramowitz, G., and Pitman, A. J.: FluxnetLSM R package (v1. 0): a community tool for processing FLUXNET data for use in land surface modelling, Geosci. Model Dev., 10(9), 3379-3390, https://doi.org/10.5194/gmd-10-3379-2017, 2017.

20 Ukkola, A. M., Pitman, A. J., Donat, M. G., De Kauwe, M. G., and Angélil, O.: Evaluating the contribution of landatmosphere feedbacks to heat extremes in CMIP5 models, Geophys. Res. Lett., 45, 9003–9012, https://doi.org/10.1029/2018GL079102, 2018.

Valladares, F., Matesanz, S., Guilhaumon, F., Araújo, M. B., Balaguer, L., Benito-Garzón, M., Cornwell, W., Gianoli, E., van Kleunen, M., Naya, D.E., et al.: The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change, Ecol. Lett., 17(11), 1351-1364, https://doi.org/10.1111/ele.12348, 2014.

25

van Gorsel, E., Cleverly, J., Beringer, J., Cleugh, H., Eamus, D., Hutley, L. B., Isaac, P., and Prober, S.: Preface: OzFlux: a network for the study of ecosystem carbon and water dynamics across Australia and New Zealand, Biogeosciences, 15, 349-352, https://doi.org/10.5194/bg-15-349-2018, 2018.

van Gorsel, E., Wolf, S., Cleverly, J., Isaac, P., Haverd, V., Ewenz, C., Arndt, S., Beringer, J., Resco de Dios, V., Evans,
B. J., Griebel, A., Hutley, L. B., Keenan, T., Kljun, N., Macfarlane, C., Meyer, W. S., McHugh, I., Pendall, E., Prober, S. M., and Silberstein, R.: Carbon uptake and water use in woodlands and forests in southern Australia during an extreme heat wave event in the "Angry Summer" of 2012/2013, Biogeosciences, 13, 5947-5964, https://doi.org/10.5194/bg-13-5947-2016, 2016.

Van Mantgem, P. J., Stephenson, N. L., Byrne, J. C., Daniels, L. D., Franklin, J. F., Fulé, P. Z., Harmon, M. E., Larson, A. J., Smith, J. M., Taylor, A. H. and Veblen, T. T.: Widespread increase of tree mortality rates in the western United States, Science, 323(5913), 521-524, https://doi.org/10.1126/science.1165000, 2009.

von Buttlar, J., Zscheischler, J., Rammig, A., Sippel, S., Reichstein, M., Knohl, A., Jung, M., Menzer, O., Arain, M. A.,

- 5 Buchmann, N., Cescatti, A., Gianelle, D., Kiely, G., Law, B. E., Magliulo, V., Margolis, H., McCaughey, H., Merbold, L., Migliavacca, M., Montagnani, L., Oechel, W., Pavelka, M., Peichl, M., Rambal, S., Raschi, A., Scott, R. L., Vaccari, F. P., van Gorsel, E., Varlagin, A., Wohlfahrt, G., and Mahecha, M. D.: Impacts of droughts and extreme-temperature events on gross primary production and ecosystem respiration: a systematic assessment across ecosystems and climate zones, Biogeosciences, 15, 1293-1318, https://doi.org/10.5194/bg-15-1293-2018, 2018.
- 10 Way, D. A. and Sage, R. F.: Thermal acclimation of photosynthesis in black spruce [Picea mariana (Mill.) B.S.P.], Plant Cell Environ., 31(9), 1250-1262, https://doi.org/10.1111/j.1365-3040.2008.01842.x, 2008.

Wilson, K., Goldstein, A., Falge, E., Aubinet, M., Baldocchi, D. D., Berbigier, P., Bernhofer, C., et al.: "Energy Balance Closure at FLUXNET Sites." Agricultural and Forest Meteorology, FLUXNET 2000 synthesis, 113 (14), 223–43. https://doi.org/10.1016/S0168-1923(02)00109-0, 2002.

15 Wolter, K., Eischeid, J. K., Quan, X. W., Chase, T. N., Hoerling, M., Dole, R. M., van Oldenborgh, G., and Walsh, J. E.: How Unusual was the Cold Winter of 2013/14 in the Upper Midwest?, B. Am. Meteorol. Soc., 96(12), S10-S14, https://doi.org/10.1175/BAMS-D-15-00126.1, 2015.

Zander, K. K., Botzen, W. J. W, Oppermann, E., Kjellstrom, T., and Garnett, S. T.: Heat stress causes substantial labour productivity loss in Australia, Nat. Clim. Change, 5, 647-652, https://doi.org/10.1038/nclimate2623, 2015.

20