

A novel isotope pool dilution approach to quantify gross rates of key abiotic and biological processes in the soil phosphorus cycle

Wolfgang Wanek*, David Zezula, Daniel Wasner, Maria Mooshammer, Judith Prommer

5 Division of Terrestrial Ecosystem Research, Department of Microbiology and Ecosystem Science, Research Network „Chemistry meets Microbiology“, University of Vienna, Althanstraße 14, 1090 Vienna, Austria

Correspondence to: Wolfgang Wanek (wolfgang.wanek@univie.ac.at)

Abstract.

10 Efforts to understand and model the current and future behavior of the global phosphorus (P) cycle are limited by the availability of global data on rates of soil P processes, as well as their environmental controls. We here present a novel isotope pool dilution approach using ^{33}P labelling of live and sterile soils, which allows to obtain high quality data on gross fluxes of soil inorganic P (P_i) sorption and desorption, as well as of gross fluxes of organic P mineralization and microbial P_i uptake. At the same time, net immobilization of $^{33}\text{P}_i$ by soil microbes and abiotic
15 sorption can be easily derived and partitioned. Compared to other approaches, we used short incubation times (up to 48 h), avoiding tracer re-mineralization, which was confirmed by separation of organic P and P_i using isobutanol fractionation. This approach is also suitable for strongly weathered and P impoverished soils, as sensitivity is increased by extraction of exchangeable bio-available P_i (Olsen P_i ; 0.5 M NaHCO_3) followed by P_i measurement using the malachite green assay. Biotic processes were corrected for desorption/sorption processes by using
20 adequate sterile abiotic controls that exhibited negligible microbial and extracellular phosphatase activities. Gross rates are calculated using analytical solutions of tracer kinetics, which also allows to study gross soil P dynamics under non-steady-state conditions. Finally, we present major environmental controls of gross P cycle processes that were measured for three P-poor tropical forest and three P-rich temperate grassland soils.

25 **Keywords:** phosphorus, organic P mineralization, sorption, desorption, isotope pool dilution, ^{33}P ;

1 Introduction

Phosphorus (P) is a major limiting nutrient to terrestrial primary production, particularly so on highly weathered soils, as, e.g. found in the tropics. Globally, increasing imbalances between nitrogen (N) and P inputs (i.e.
30 increasing N:P stoichiometry of inputs) caused by human activities and land-use changes through increased emissions of reactive N are suggested to lead to progressive P limitation of terrestrial ecosystems, and first signs thereof have been identified (Penuelas et al., 2013). A decrease in the relative P availability might have strong repercussions on future nutrient limitations of natural ecosystems, on food production and on carbon (C) sequestration (Penuelas et al., 2013; Penuelas et al., 2012; Yang et al., 2013). Efforts to understand and model the
35 current and future global P cycle and its coupling to the global C and N cycles have been intensified, but are

strongly limited by the availability of global data on soil gross P processes and their environmental controls (Reed et al., 2015). Large investments into new projects, experiments and models have therefore been recently undertaken to advance our understanding of the terrestrial P cycle, and to fill data gaps, e.g. IMBALANCE-P (<http://imbalancep-erc.creaf.cat>) and Ngee-TROPICS (<http://ngee-tropics.lbl.gov>).

40 Soil P_i availability is governed by transfers between pools of exchangeable P, immobilized/fixed P and occluded P, by slow release of P_i from mineral P through weathering of primary minerals, and by mineralization of organic P (P_o) (Fig. 1) (Bünemann, 2015; Turner et al., 2007). In strongly weathered soils, primary mineral P pools are depleted, and the largest fraction of P is found in occluded and fixed pools, as well as in P_o (Vitousek and Farrington, 1997; Yang and Post, 2011). Phosphorus limitation in such soils is further aggravated by their high
45 P sorption potentials caused by high contents of Fe-Al (hydr)oxides (Goldberg and Sposito, 1985). Most of the immediate P needs of plants (and microbes) in natural and agricultural systems is supplied by P_o mineralization, catalyzed by extracellular phosphatases that are released by soil microbes and plant roots (Richardson and Simpson, 2011), as well as by abiotic P_i desorption. Soil microbes and plant roots can promote the release of P from primary and secondary minerals by accelerating mineral dissolution and P_i desorption, through exudation of
50 (phyto)siderophores and organic acids (Mander et al., 2012; Ryan et al., 2001).

Soil P cycling processes such as soil P_i sorption/desorption fluxes and gross P_o mineralization rates, as well as the size of the exchangeable soil P_i pool have been measured by isotope exchange (IEK) techniques using ^{32}P or ^{33}P . These techniques are based on recurrent measurements of radiotracer recovery and P_i concentration in soil water extracts (Di et al., 1997; Frossard et al., 2011; Bünemann, 2015) (Table 1). A variety of IEK procedures
55 and protocols are in use, and optimizations in methodology have been called for, particularly for P_o mineralization (Bünemann, 2015). Only during the last decade common, accepted protocols have become adopted and are currently used to measure soil P processes following Oehl et al. (2001b). In this IEK approach in short-term experiments abiotic sorption/desorption processes from an isotopically exchangeable P_i pool are measured over a short time period in batch experiments (100 min, 1:10 (w:v) soil: water slurry, \pm microbicides).
60 This assumes that no microbial tracer uptake (blocked by microbicides) and no organic P mineralization occurs, and that soils are in a steady state i.e. don't show changes in P_i concentration (Table 1). In such short-term IEK experiments the decrease in radioactivity (radiotracer recovery) in soil water is described by a power function:

$$r(t)/R = r_{1min}/R \times t^{-n}$$

R is the added radioactivity and $r(t)$ the radioactivity recovered at any time t in soil water extracts. The
65 parameters r_{1min}/R and n (slope of the regression indicating speed of isotopic exchange) are derived from the log-log regression of $r(t)$ versus time. This is based on steady state assumptions, i.e. that P_i concentration in soil water extracts (C_p) is constant. In some soils an extended version of this equation needs to be applied:

$$r(t)/R = m \times (t + m^{1/n})^{-n} + r_{inf}/R$$

Here, r_{inf}/R is the maximum possible dilution of the added radiotracer, approximated as the ratio of C_p to total inorganic P in soils. n and m are derived from non-linear fitting procedures. Assuming that tracer and tracee
70 behave similarly in the system, the specific activity of P_i in soil solution should reflect the specific activity of isotopically exchangeable P – termed E -value (in $mg\ P\ kg^{-1}$ soil).

$$E_{(t)} = C_p / (r(t)/R).$$

Isotopic dilution ($E'_{(t)}$) is further measured over the full length of a moist soil incubation experiment lasting for
75 several days to weeks, constituting the total amount of exchangeable P_i or isotope dilution caused by concurrent biological processes (P_o mineralization) and physicochemical processes. Short-term exchange kinetics are then

extrapolated over the full time period of the moist soil incubation ($E_{(t)}$). The difference between $E'_{(t)}$ and the extrapolated $E_{(t)}$ -value provides then the measure of gross P_o mineralization.

The isotope pool dilution approach (IPD) of Kirkham and Bartholomew (1954) was developed as a general tracer approach to measure gross rates of soil element cycle processes, but was most frequently applied to nitrogen cycling processes such as organic N mineralization and nitrification (Booth et al., 2005). The IPD approach can however also be transferred to measure gross rates of P cycle processes. It then also relies on the labelling of the P_i pool with ^{33}P or ^{32}P and on subsequent time-resolved measurements of concentrations and specific activities of P_i (Table 1, Figure 1B). However, in contrast to IEK techniques, changes in P_i concentrations and specific activities are then solved by mass balance equations developed specifically for gross rate calculations based on tracer studies (Kirkham and Bartholomew, 1954). In the following we list the criteria that have to be met by the IPD method to correctly determine gross rates of soil P_o mineralization and soil P_i sorption/desorption (Di et al., 2000; Murphy et al., 2003; Kirkham and Bartholomew, 1954).

1. The tracer ($^{32}\text{P}_i$ or $^{33}\text{P}_i$) and tracee (unlabeled $^{31}\text{P}_i$) behave identically and are well mixed. This is given for the different isotopes of P as long as radiotracer solution is homogeneously distributed in the soil and sufficient time is provided for isotope equilibration between added radiotracer and the native P_i pool.
2. The influx into the target (P_i) pool (i.e. the product of P_o mineralization) has to be unlabeled (i.e. no tracer remineralization), in order for it to dilute the tracer: tracee ratio over time (Figure 1B and C). Tracer remineralization via microbial tracer assimilation, mortality and subsequent remineralization of labelled P_o would result in an underestimation of P_o mineralization, but can be avoided by short incubation times (1-2 days).
3. Abiotic release of P_i from a non-extractable pool (P_i desorption) causes an influx of unlabeled P_i into the target pool, resulting in an overestimation of the biotic process, P_o mineralization, and has to be determined in parallel abiotic incubations of sterile soils. However, adequate abiotic controls with no contribution of biological processes has remained a major obstacle in measuring soil P dynamics with radiotracers, both in IEK and IPD experiments. Procedures in earlier studies ranged from short-term assays with no inhibitor addition as often performed in IEK assays (Spohn et al., 2013; Oehl et al., 2001b), to amendments of HgCl_2 , sodium azide, toluene or chloroform, and gamma irradiation or repeated autoclaving (Kellogg et al., 2006; Bünemann, 2015; Bünemann et al., 2007; Oehl et al., 2001b; Achat et al., 2010).
4. The soil extraction should target the bio-available exchangeable P_i pool. P_i in soil solution undergoes rapid equilibration with easily adsorbed P_i . An incomplete extraction of this pool causes an underestimation of P_o mineralization rates, due to desorption from this pool, causing an influx of unlabeled tracer (together with unlabeled P_i) into the target pool, and thus violates assumption #2 of IPD assays. The commonly used soil water extractions target only a small fraction of this target pool, whereas standard soil P extractants, such as Olsen, Mehlich-3 or Bray-1, extract a larger fraction (Kleinman et al., 2001) and, therefore, are suggested to be better suited to extract the rapidly exchanging P_i pool (Kellogg et al., 2006).
5. The efflux from the isotopically labelled pool (i.e. microbial P_i immobilization and P_i sorption into a non-extractable pool) occurs at the ratio of tracer: tracee as present in the P_i pool at any specific time, with no discrimination between native P_i and added radiotracer (Figure 1B). A short pre-incubation time is therefore needed to allow for full mixing and isotopic equilibration of tracer and tracee (see point #1).
6. Changes in specific activity need to be measured specifically in the target pool, i.e. in extractable P_i for measurements of gross rates of P_o mineralization and P_i sorption/desorption. However, most current

approaches do not separate extractable P_i and P_o but measure radioactivity in unfractionated extracts, including radiolabeled P_o formed during the incubation, leading to an eventual overestimation of P_o mineralization.

- 120
7. The rates of P_i influx (P_o mineralization, abiotic P_i release) and P_i efflux (biotic and abiotic P_i immobilization) need to be constant over the duration of incubation: (i) the initial phase of fast immobilization by sorption, microbial uptake and isotopic equilibration of radiotracer is excluded from calculations of gross rates, and (ii) incubation takes place within a suitable timeframe to avoid microbial turnover and $^{33}P_o$ remineralization (see point #2). The minimum two time points necessary to measure concentration and specific activity of P_i for the IPD calculations should therefore lie in between the initial phase and the start of re-mineralization but it is recommendable to test more time points in the beginning to test time linearity of IPD rates for specific soil types.

130 Mooshammer et al. (2012) adopted such a protocol for measurements of gross P_o mineralization in decomposing plant litter, following the knowledge of IPD processes based on ^{15}N additions to study gross rates of soil N cycling (Hart et al., 1994; Murphy et al., 2003; Wanek et al., 2010; Braun et al., 2018). However, in plant litter P sorption and the abiotic release of P_i from sorbed P pools do not interfere. Consequently, the litter protocol cannot be directly transferred to soil studies. In the present study we developed an IPD protocol to assess soil P dynamics, based on the previous work for litter by Mooshammer et al. (2012) and soils by Kellogg et al. (2006). The protocol is based on IPD theory (Kirkham and Bartholomew, 1954; Di et al., 2000) applied to parallel incubations of live and sterile soil with $^{33}P_i$ tracer addition. Gross rates of P_i sorption (abiotic immobilization) and P_i desorption are determined in sterile soils, and allow correction of gross P_o mineralization and microbial P_i immobilization rates in live soils. We used bicarbonate extractions to target the bio-available exchangeable P_i pool. To avoid tracer re-mineralization, we used short incubation periods (up to 2 days). To confirm that no significant amount of $^{33}P_o$ was formed during incubation, P_i was also separated from P_o based on isobutanol fractionation (Jayachandran et al., 1992). P_i concentrations were measured based on the phosphomolybdate blue protocol. At very low P_i concentrations, e.g. in tropical soils, that are below the detection limit of the phosphomolybdate blue method, P_i was determined by parallel measurements of P_i in bicarbonate extracts using the more sensitive malachite green assay (D'Angelo et al., 2001; Ohno and Zibilske, 1991). The protocol was tested rigorously with two different soils, and then applied to in total six soils (three tropical forest and three temperate grassland soils) to explore environmental controls on gross soil P dynamics.

2 Materials and methods

2.1 Soil materials and basic characterization

150 Soils (0-15 cm depth) were collected in summer 2015 from three temperate grassland sites in Austria and in spring 2015 from three tropical lowland forest sites in Costa Rica (Table 2). The grassland soils were extensively managed meadows, collected in Lower Austria (48° 13-20' N, 16° 12-17' E) in the vicinity of Vienna, at elevations between 170 and 320 m. The tropical forest soils were collected along a topographic gradient (ridge-slope-valley bottom) in wet evergreen old-growth forests in SW Costa Rica close to the National Park Piedras Blancas (8° 41' N, 83° 12' W, 110-250 m a.s.l.). Soils were sieved to 2 mm and stored in an air-dried state. Soil pH was measured in a 1:5 (w:v) mixture of air dried soil in water after 60 min of equilibration using an ISFET electrode (Sentron SI600 pH Meter). Soil texture was quantified using a miniaturized pipette/sieving protocol for 2-4 g air dried soils (Miller

and Miller, 1987), using 4% sodium metaphosphate as a dispersant. Soil total C and total soil N content were determined after grinding oven dried soil in a ball mill, using an elemental analyzer (EA 1110, CE Instruments, Thermo Scientific). Temperate grassland soils were treated with 2 M HCl to remove carbonates, re-dried, ground and then analyzed by elemental analyzer for soil organic C. Total soil P and total soil P_i were measured after 0.5 M H_2SO_4 extraction of ignited soils (5 h at 450 °C in a muffle furnace; (O'Halloran and Cade-Menun, 2008)) and of untreated soils, respectively, by the malachite green method (Ohno and Zibilske, 1991;D'Angelo et al., 2001). Total organic P was estimated by calculating the difference between total soil P and total soil P_i . We must however submit that ignition methods tend to overestimate soil organic P in highly weathered tropical soils (Condon et al., 1990).

2.2 Soil pre-treatment and assay of sterilization efficiency (abiotic controls)

Before starting the experiments, the soils were re-equilibrated from an air-dried state by rewetting to 60% water holding capacity for 6 days at 20 °C. Gravimetric soil water content and water holding capacity were determined prior to the experiment. Soils were then either sterilized twice, 48 and 2 h before start of the IPD experiments, by autoclaving at 121 °C for 60 min (sterile soils), or were kept at 20 °C (live soils, Fig. 2). Sterilization efficiency was checked based on soil enzyme activity measurements. Fluorescein diacetate (FDA) hydrolysis in soils was measured as a proxy of viable, active microbial biomass (Green et al., 2006;Schnurer and Rosswall, 1982), and the activity of acid phosphomonoesterases, which are extracellular enzymes involved in P_o mineralization, was determined using methylumbelliferyl (MUF)-phosphate (Sirova et al., 2013;Marx et al., 2001).

2.3 ^{33}P IPD assay

A schematic representation of the final IPD protocol can be found in Figure 2. Duplicate soil aliquots (2 g fresh weight) of sterile and live soil were amended each with 20 kBq $^{33}P_i$ (dilution of orthophosphoric acid phosphorus-33 radionuclide, 5 mCi mL^{-1} , i.e. 185 MBq mL^{-1} HCl-free water at specified date, Perkin NEZ080002MC). Between 0.15-0.2 mL of ^{33}P -label solution was added to each sample (Fig. 2); the volume added was adjusted for each soil type to obtain an optimal water content in each soil (~75 % water holding capacity). Soils were extracted with 30 mL (temperate soils) or 15 mL (tropical soils) of 0.5 M $NaHCO_3$ (pH 8.5) after 4 and 24 h of incubation for 30 min on a horizontal shaker and filtered through ash-free cellulose filters. Lower extractant volumes in tropical and other P poor soils are used to reach higher P_i concentrations in the bicarbonate extracts for better quantification.

Thereafter isobutanol fractionation of the bicarbonate extracts was performed, separating P_i (into the organic phase) from P_o (into the acidic aqueous phase) allowing measurement of the kinetics and specific activity of the P_i pool without interference of P_o (Kellogg et al., 2006;Mooshammer et al., 2012). Isobutanol partitioning enables 100% recovery of P_i with no hydrolysis of P_o (Jayachandran et al., 1992). For isobutanol fractionation each 1.5 mL of soil extracts, standards and blanks were amended by sequential addition of 1.5 mL acidified molybdate, 3 mL deionized water and 3 mL isobutanol. The acidified molybdate reagent consists of 5 g ammonium molybdate tetrahydrate ($(NH_4)_6Mo_7O_{24}.4H_2O$) dissolved in 0.1 L 2.3 M H_2SO_4 (stable at room temperature for at least three months) and causes strong CO_2 outgassing from the bicarbonate extracts. After addition of all reagents the vials were shaken overhead for 1 min and then rested for 10 min for phase separation. For later photometric quantification of P_i in the isobutanol phase, standards ranging from 320 to ~1 μM P_i (1:2 dilution series) and blanks, both of the same matrix as soil extracts (i.e. 0.5 M $NaHCO_3$), were prepared and underwent isobutanol

fractionation together with the samples. ^{33}P recovery standards were also prepared and processed through the isobutanol fractionation protocol, consisting of the same volume of extractant (15 or 30 mL) and ^{33}P tracer activity as added to soils (Fig. 2).

P_i in the isobutanol phase was quantified using the phosphomolybdate blue color reaction according to Murphy and Riley (1962). Briefly, each 1.5 mL of the upper organic phase were transferred to vials and amended with 2.1 mL molybdate free reducing agent, consisting of 1.32 g ascorbic acid dissolved in 250 mL antimony potassium tartrate (APT) solution (145.4 mg APT in 0.5 M H_2SO_4). The APT solution is stable at room temperature for >4 weeks, whereas the molybdate free reducing agent has to be prepared fresh daily. Thereafter samples were shaken overhead for 1 min and rested for 20 min for phase separation and color development. A volume of 250 μL of the blue isobutanol phase was then pipetted into a microtiter plate and absorbance was read at 725 nm with a microplate photometer (Tecan Infinite M200, Tecan Austria GmbH, Grödig, Austria).

In parallel to the phosphomolybdate blue reaction of P_i in the isobutanol phase, P_i concentrations were also determined directly in acidified bicarbonate extracts using the malachite green approach (D'Angelo et al., 2001). This method is 4-10 times more sensitive than the commonly used phosphomolybdate blue method and was chosen to account for the expectedly low P_i concentrations of the tropical soils. Standards for calibration of the malachite green method were prepared in 0.5 M NaHCO_3 , ranging from 50 to 0.039 μM P_i . Acidification of bicarbonate extracts and standards (blanks) was performed on 2.5 mL sample aliquots by adding 250 μL 2.75 M H_2SO_4 (Fig. 2). Of the acidified samples and standards, 200 μL were pipetted into a microtiter plate, 40 μL malachite green reagent A were added and incubated for 10 min. Then 40 μL reagent B were added and absorbance was read after 45 min at 610 nm with a microplate reader. Reagent A was prepared by adding 50 mL deionized water in an amber 0.1 L glass bottle, adding 16.8 mL concentrated H_2SO_4 , stirring and dissolving 1.76 g ammonium heptamolybdate tetrahydrate ($(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}\cdot 4 \text{H}_2\text{O}$). Reagent B was prepared by heating 0.25 L of distilled H_2O to 80°C in an amber 0.5 L glass bottle, dissolving 0.875 g PVA (polyvinyl alcohol, MW = 72000 g/mol) whilst continuously stirring, cooling to room temperature, and finally dissolving 87 mg malachite green oxalate in this solution. Both reagents are stable for >6 months at room temperature.

Radioactivity (^{33}P activity) was measured in 0.25 mL aliquots of acidified bicarbonate extracts and in 0.4 mL aliquots of the isobutanol phase, after addition of each 4 mL scintillation cocktail (Ultima Gold, Perkin Elmer), by liquid scintillation counting (Tri-Carb 1600 TR, Packard, Perkin Elmer) (Fig. 2).

2.4 Experiments

- (i) Time kinetics: high resolution time kinetics of tracer and tracee dynamics ($^{33}\text{P}_i$, $^{31}\text{P}_i$) were measured in two soils (temperate grassland, soil 4; tropical forest, soil 3; Table 2). After tracer addition to live and sterile soils in triplicates IPD assays were stopped by extraction with 0.5 M NaHCO_3 after 0, 1, 2, 4, 8, 24, and 48 h.
- (ii) Microbial ^{33}P immobilization: the procedure outlined in chapter 2.3 can be combined with direct determination of microbial P by extraction with liquid chloroform-enriched salt solutions (Setia et al., 2012). We here tested a sequential extraction-liquid chloroform extraction (sECE) procedure. After 24 h of soil incubation in experiment (i), soil samples (2 g fresh weight) were first extracted with 15 (soil 4) or 30 (soil 3) mL 0.5 M NaHCO_3 for 30 min, centrifuged for 15 min at 10.000 g, and the supernatant was decanted. The soil residue was then re-extracted with 15 (30) mL 0.5 M NaHCO_3 containing 3% (v:v) chloroform for 30 min and finally filtered through ash-free cellulose filters.

240 Volume corrections were applied for extractant absorption by the soil pellet after centrifugation. Volume corrections were calculated as soil wet weight after centrifugation minus fresh weight weighed into each tube in grams, divided by the density of the bicarbonate solution (in g/mL), providing the carry-over of extractant from the first extraction (in mL).

245 (iii) Soil effects on tracer dynamics: live and sterile soils (2 g aliquots) of all 6 soils (Table 2) were measured in triplicates for $^{33}\text{P}_i$ activity and P_i concentrations, and assays were stopped after 0, 4 and 24 h. Net immobilization of ^{33}P and gross process rates were calculated for the time interval 4 to 24 h, and relationships between gross and net soil P processes and soil physicochemical parameters were tested.

250 2.5 Calculations of abiotic and biotic net ^{33}P immobilization

Additionally to the measurement of gross rates, abiotic net ^{33}P immobilization (net soil P_i fixation) and biotic net ^{33}P immobilization (net soil microbial P_i immobilization) were calculated based on the determination of the recovery of added tracer in soil extracts of live and autoclaved soils (see above) after 0, 1, 2, 4, 8, 24 and 48 hours. Abiotic immobilization (in % added tracer) was estimated as 100 percent minus the percent ^{33}P recovery in autoclaved soils. Total immobilization was estimated as 100 minus the percent ^{33}P recovery in live soils. Biotic immobilization was calculated as the difference between total and abiotic immobilization. These data provide a rapid assessment of the abiotic versus microbial sink strengths for P_i , but do not represent gross rates.

260 2.6 Calculations of gross rates of soil P dynamics

Calculation of gross IPD rates followed the mass balance equations of Kirkham and Bartholomew (1954), as later applied by others for soil gross P fluxes (Kellogg et al., 2006; Mooshammer et al., 2012). In these previous studies abiotic processes were not corrected from the final data. However, to calculate gross P_o mineralization, gross rates of P_i desorption have to be corrected for in live soils. This is due to gross rates of P_i influx in live soils being the integral of abiotic processes (desorption) and biological processes (P_o mineralization). This abiotic correction was performed by applying IPD calculations for influx (GI, gross influx; equation 1) for sterile soils (abiotic influx by P_i desorption) and live soils (total P_i influx), and taking the difference as biotic influx (i.e. gross P_o mineralization). The same procedure was performed for tracer efflux (GE=gross efflux; equation 2) calculating gross immobilization fluxes for live soils (total P_i efflux) and sterile soils (P_i sorption), the difference providing gross rates of microbial P_i immobilization. Both abiotic corrections are based on the assumption that abiotic sorption/desorption processes are not affected by autoclaving, i.e. that these processes act similarly in sterile and in live soils.

$$\text{Gross influx: } GI = \frac{c_{t2} - c_{t1}}{t_2 - t_1} \times \frac{\ln\left(\frac{SA_{t1}}{SA_{t2}}\right)}{\ln\left(\frac{c_{t2}}{c_{t1}}\right)} \quad (\text{Eq. 1})$$

$$\text{Gross efflux: } GE = \frac{c_{t1} - c_{t2}}{t_2 - t_1} \times \left(1 + \frac{\ln\left(\frac{SA_{t2}}{SA_{t1}}\right)}{\ln\left(\frac{c_{t2}}{c_{t1}}\right)}\right) \quad (\text{Eq. 2})$$

275 where t_1 and t_2 represent incubation time (4 and 24 h; in days), C the soil P_i concentration (in $\mu\text{g } \text{P}_i \text{ g}^{-1}$ soil dry weight), SA the specific activity (in $\text{Bq } \mu\text{g}^{-1} \text{P}_i$) and LN the natural logarithm. Gross rates are therefore in $\mu\text{g } \text{P}_i \text{ g}^{-1}$ soil dry weight d^{-1} . Net organic P mineralization rates can easily be derived by subtracting gross microbial P_i uptake from gross P_o mineralization rates.

Due to the relatively rapid decline in ^{33}P activity by radioactive decay, all data were decay corrected back to the start of each experiment, i.e. the time point of tracer addition to the soil. This was done according to equation 3.

$$N_{\text{to}} = \frac{N_t}{e^{-\lambda t}} \quad (\text{Eq. 3})$$

where N_{to} is the decay corrected ^{33}P activity in a sample (in Bq), N_t the measured ^{33}P activity at time of liquid scintillation counting, t is time (in days) elapsed between tracer addition and ^{33}P activity measurement, $e=2.71828$ and λ the decay constant of ^{33}P (0.0273539).

285 2.7 Statistics

Regressions were performed in Sigmaplot 13.0 (Systat Software, Inc.) and group differences were tested by one-way and two-way ANOVA followed by Tukey's HSD test in Statgraphics Centurion XVIII (Statpoint Technologies, Inc.). Variance homogeneity was tested by Levene's test and if necessary data were log, square root or rank transformed to meet assumptions of homoscedasticity and normal distribution.

290

3 Results

3.1 Soil characterization

Temperate grassland soils had a pH between 6.3 and 6.8, with a silt loam to sandy loam texture (Table 2). Soil organic C contents ranged between 48 and 127 mg C g⁻¹, soil N from 2.3 to 5.0 mg N g⁻¹ and soil total P from 0.44 to 0.82 mg P g⁻¹. Tropical forest soils had a pH between 4.1 and 4.2, and soil texture varied between silt, silt loam and sandy loam. Soil organic C contents were lower, at 26 to 31 mg C g⁻¹, soil N ranged from 2.2 to 2.6 mg N g⁻¹, and soil total P from 0.09 to 0.17 mg P g⁻¹. Organic P comprised a larger fraction of total P in tropical forest soils (64-76%) than in temperate grassland soils (22-57%). Extractable soil P_i was higher in temperate grasslands (4.2-13.1 μg P g⁻¹ soil dry weight) compared to tropical forest soils (0.07-0.13 μg P g⁻¹ soil dry weight). Acid phosphomonoesterase activities of tropical forest soils (1396-2346 nmol MUF released g⁻¹ dry weight h⁻¹) markedly exceeded those in temperate grasslands (233-256 nmol MUF released g⁻¹ dry weight h⁻¹).

300

3.2 Abiotic controls: soil sterilization efficiency

A separation of biotic and abiotic processes is based on the comparison of gross rates using the IPD assay in live versus autoclaved soils, where the latter should not exhibit any microbial activity (no FDA hydrolysis activity) and no extracellular enzyme activities (no MUF-phosphatase activity), in order to serve as abiotic controls. An incomplete inhibition of extracellular phosphatase activities would lead to an underestimation of biological processes and therefore of gross P_o mineralization. Our results show that two consecutive treatments of the soils by autoclaving, with a 48 hours incubation in between, effectively reduced microbial metabolic activity as shown by the reduction in soil FDA hydrolysis by 90% in soil 4 and by 97-99% in all other soils (Fig. 3). Autoclaved soils did not show any increase in soil microbial activity during the two days of incubation. On the contrary, the inhibition of FDA hydrolysis even increased from 1 hour (all soil average: 94%) towards 24 and 48 hours after sterilization (average: 97-99%). The inhibition of extracellular acid phosphatase activity was almost complete in tropical soils (95-97%) and strongly reduced in temperate soils (79-80%). Similar to FDA hydrolysis the extent of inhibition of phosphatase activity increased from day 0 (average: 86%) to day 1 and 2 (average: 88-89%, Fig. 3). However, autoclaving increased available P_i by 1.86 ± 0.32-fold (mean ± 1SD) in temperate soils and by 1.65 ± 0.36-fold in the tropical soils (Fig. S1).

315

3.3 Comparison of isobutanol fractionation and direct measurements of P_i and ^{33}P activity

320 Soil P_i concentrations measured by the malachite green method directly in acidified bicarbonate extracts were compared to those measured after isobutanol fractionation by phosphomolybdate blue reaction, including both live and sterile soils. Both approaches yielded similar soil P_i concentrations, and the relationship showed no bias (slope = 0.979 ± 0.033 , mean \pm 1SE), with a coefficient of determination of 0.92 (Fig. 4B). The malachite green method is much more sensitive and therefore produced more reliable results for the low-P soils from the three tropical forests.

325 Moreover, the relationship between ^{33}P recoveries by isobutanol fractionation and by direct measurements in acidified bicarbonate extracts had a slope less than 1 (slope = 0.875 ± 0.010 ; Fig. 4A), indicating no significant formation of $^{33}P_o$ during soil incubations. We also found no $^{33}P_o$ formation in other soils using the same measurement protocols, e.g. from the Jena biodiversity experiment (82 plots of temperate grassland varying in soil texture and plant biodiversity, slope = 0.891 ± 0.017) and from French Guyana (24 soils from two primary forest

330 regions, with soils sampled across topographic gradients, slope = 1.043 ± 0.020) (same regression types as in Fig. 3A; data not shown). The specific activities of P_i were indistinguishable between both approaches for temperate soils (slope = 0.977 ± 0.064 , $R^2 = 0.93$, $P < 0.0001$; Fig. 4C) but varied strongly for the tropical soils, where soil P_i measurements in the isobutanol fraction were at or below the limit of detection of the phosphomolybdate blue method.

335

3.4 Sensitivity of the IPD assay

The sensitivity of this assay is greatly improved relative to traditional ones, by using a combination of bicarbonate extractions and malachite green P_i measurements. The detection limit of the IPD approach was $0.12 \mu\text{g P g}^{-1} \text{ soil dw d}^{-1}$, based on three times the standard deviation of gross P_o mineralization, measured for the three tropical soils

340 (each measured in triplicates), and therefore fully suitable across all soil types tested so far. However, the precision suffers from IPD equations that combine uncertainties from four measurements, two P_i concentrations and two radioactivity measurements for the two time points in live as well sterile soils. The coefficients of variation (CV) ranged between 1.0 and 22.1% (average 10.0%) for P_i concentration across temperate and tropical soils, and between 1.5 and 22.1% (average 9.6%) for SA, the two major input variables into the IPD equation. CVs increased

345 towards lower P_i concentrations and higher SA values, i.e. closer to the detection limit of the malachite green method. The CVs might be reduced by working with larger soil aliquots (increase from 2 to 5 or 10 g soil fresh weight) and by duplicate measurements of all samples. Purely methodological CVs were lower, at about 2.5 and 0.9% for P_i measurements by malachite green in the range 3-12 and 12-120 μM , respectively, and 0.8% for liquid scintillation counting. Therefore, much of the shown variability derived from differences between biological soil

350 replicates. However, the variability found here compares well with CVs published for soil P_i concentrations of 2-10% (Bünemann et al., 2007) and 20-25% (Bünemann et al., 2012), and CVs for measured E values that are calculated from P_i concentrations and ^{33}P recoveries analogous to SA values ranged between 6-16% (Bünemann et al., 2007), 8-19% (Bünemann et al., 2012) and 9-10% (Randriamanantsoa et al., 2015) across a range of cultivated and non-cultivated soils from temperate to tropical regions. These variations naturally propagate into

355 higher errors in the measured rates of soil P cycling and increase the limit of detection and the limit of quantification of the various methods.

3.5 Time kinetics

During the first hour of the incubation, we found a rapid drop in ^{33}P recovery and in the SA of P_i (Fig. 4), while soil P_i concentrations increased slightly (Fig. S1). Thereafter a dynamic equilibrium between added ^{33}P tracer and the soil P_i pool was reached and concentrations of extractable P_i remained constant. A plot of $\ln(^{33}\text{P}$ recovery) versus time of both live and sterile soils showed that the consumption of ^{33}P occurred linearly between 4 and 48 h in the temperate soil and between 2 and 24 h in the tropical soil (Fig. 5). Similarly, the plot of $\ln(\text{SA of } \text{P}_i)$ versus time showed a linear relationship from 4 to 48 h in the temperate soil and for 2 to 48 h in the tropical soil, particularly in live soils (Fig. 5), showing constant dilution of the isotopic signature of the pool over time. The regressions became insignificant in the sterile tropical soil, as ^{33}P recovery and SA declined more slowly. The data clearly show that abiotic ^{33}P processes (i.e. decreases in ^{33}P recovery and SA of P_i over time in sterile soils) occurred, particularly in the temperate soil, and this over a prolonged period of time. More importantly, the dynamics of abiotic ^{33}P processes changed over time: rapid abiotic immobilization during the initial 0-4 h was followed by a period of slower but linear tracer immobilization.

3.6 Net ^{33}P immobilization by abiotic and biotic processes

Abiotic net ^{33}P immobilization (net soil P fixation) increased markedly from 0 to 48 h in the grassland soil (17 to 58% of added tracer), while it reached 83% almost instantaneously in tropical soil and further increased to 90% after 48 h (Fig. 6A). Similar patterns were found across all 6 soils, with significantly higher abiotic net immobilization in tropical than temperate soils, increasing in both with time from 0 to 4 and 24 h (Fig. 6C). Biotic (microbial) net ^{33}P immobilization ranged from 3 to 8% in the tropical soil and 8 to 17% in the temperate soil in the time kinetics experiment, with a significant increase in the temperate but not in the tropical soil (Fig. 6B). Similarly, biotic net ^{33}P immobilization was low but increased with time in all three tropical soils (3 to 6%), while it was significantly higher in temperate soils but increased (soil 6) or decreased (soil 2 and 4) with time (Fig. 6D). Microbial immobilization was very fast, with almost instantaneous ^{33}P uptake by microbes (sampling at 0 h), ranging between 3% (tropical soils) and 15-38% (temperate soils). Given the strong changes in both abiotic and biotic net ^{33}P immobilization, we suggest that it is best to measure them after 24 (up to 48) h.

Sequential extraction-liquid chloroform-extraction (sECE) was applied to directly follow net ^{33}P uptake by microbes, whereas biotic net ^{33}P immobilization was estimated indirectly as the difference in net ^{33}P immobilization by live and sterile soils. In the two measured soils, sECE estimates of microbial net ^{33}P uptake were higher than the microbial net ^{33}P immobilization estimates (temperate soil: 24.6% vs. 16.0%, and tropical soil: 16.8% vs. 7.5%, for direct and indirect estimates, respectively). This indicates incomplete extraction of exchangeable P_i prior to microbial lysis with chloroform and re-extraction.

3.7 ^{33}P pool dilution rates of abiotic and biotic processes

We calculated gross P_i influx and efflux rates for live and sterile soils. Calculated rates of sterile soils provide estimates of gross rates of soil P_i sorption and desorption, and the difference between live and sterile soils give the biotic influx (gross P_o mineralization) and efflux (gross microbial P_i uptake). Gross P_o mineralization significantly differed between soils, with two out of three temperate soils (0.48 to $2.03 \mu\text{g P g}^{-1} \text{ dw d}^{-1}$) exhibiting higher rates than two out of three tropical soils (0.08 to $0.15 \mu\text{g P g}^{-1} \text{ dw d}^{-1}$) (Fig. 7A). Gross rates of P_i sorption in temperate soils (2.06 to $6.14 \mu\text{g P g}^{-1} \text{ dw d}^{-1}$) were higher than in tropical soils (0.15 to $0.32 \mu\text{g P g}^{-1} \text{ dw d}^{-1}$), and a similar trend was found for gross rates of microbial P_i uptake (temperate: 0.44 to $1.13 \mu\text{g P g}^{-1} \text{ dw d}^{-1}$, tropical: 0.05 to

0.12 $\mu\text{g P g}^{-1} \text{ dw d}^{-1}$; Fig. 7B). Gross rates of soil P_i desorption were significantly higher in temperate soils (1.44 to 3.63 $\mu\text{g P g}^{-1} \text{ dw d}^{-1}$) than in tropical soils (0.04-0.14 $\mu\text{g P g}^{-1} \text{ d}^{-1}$, Fig. 7A). The relative contribution of P_o mineralization to total P_i release into the soil P_i pool ranged between 25.0 and 73.8%, with two tropical P-poor soils showing the highest contributions (Fig. 7C). Contributions of biological processes to gross P_i immobilization did not differ between soils (range 11.5% to 34.9%).

3.8 Physicochemical and biological controls on soil P_i processes

Gross P_o mineralization was strongly positively correlated with total soil P ($R^2=0.87$, $P<0.01$, Fig. 8A) and to total as well as extractable soil P_i concentration ($R^2>0.83$, $P<0.05$, Fig. 8B) but not to soil organic P or its contribution to soil total P, nor to soil organic C, soil texture or soil acid phosphatase activity (Table S1). Gross abiotic P_i release rates through desorption and dissolution were strongly positively related to total soil P and bicarbonate soil P_i ($R^2=0.97$ and 0.98 , respectively, both $P<0.001$, Fig. 8C and Table S1), but not to other parameters such as soil pH, soil texture, and soil organic C content. Gross P_i sorption rates exceeded gross P_i desorption rates approximately 2-fold, but both were strongly related ($R^2=0.99$, $P<0.001$, Fig. 8E). Gross P_i sorption rates were strongly positively related to soil total P ($R^2=0.96$, $P<0.001$, Fig. 8D), soil total P_i ($R^2=0.88$, $P<0.05$, Table S1) and bicarbonate soil P_i ($R^2=0.99$, $P<0.001$, Table S1), but neither to soil pH, soil organic C, nor to clay content or soil texture. Abiotic net P_i immobilization was most strongly and negatively related to soil pH ($R^2=0.95$, $P<0.001$, Fig. 8L) and weakly to soil P_i sorption ($R^2=0.59$, $P=0.073$, Fig. 8J). Gross microbial P_i uptake rates were directly proportional to microbial biomass P measured by sECE ($R^2=0.95$, $P<0.01$, Fig. 8G), and positively related to net microbial P_i immobilization ($R^2=0.85$, $P<0.01$, Fig. 8I). We found a negative curvilinear relationship between net immobilization rates by sorption and microbes ($R^2=0.97$, $P<0.001$, Fig. 8F).

420

4 Discussion

About a decade ago Kellogg et al. (2006) compared two IEK techniques with an IPD approach, identifying several biases of the different approaches and making recommendations for further development. The authors highlighted IPD approaches with soil extractions using 0.5 M sodium bicarbonate as best suited, for potentially any type of soil. However, this approach is currently underused and had issues with abiotic controls. IPD methods are state-of-the-art to measure gross processes of soil N cycling (Murphy et al., 2003), but have rarely been applied to soil P cycling processes (Mooshammer et al., 2012; Di et al., 2000; Kellogg et al., 2006). We here present a novel and versatile approach to derive quantitative estimates of soil P cycling processes that drive soil P availability in low to high P soils. The approach quantifies gross rates of soil P_o mineralization and the abiotic release of P_i from non-extractable soil P_i pools (P_i desorption and dissolution), both causing gross influx of P_i into the soil available P_i pool. Furthermore, gross rates of P_i immobilization by soil sorption and precipitation and by microbial uptake processes are derived from the same data by calculating the efflux from the soil P_i pool in sterile soils (abiotic) and in live minus sterile soils (biotic processes), respectively.

In contrast to many earlier IEK assays the IPD approach presented here is based on real isotope pool dilution theory (Kirkham and Bartholomew, 1954), and not on curvilinear extrapolation of E values (Table 1). Moreover, IEK assays of P_o mineralization necessitate steady-state conditions (constant P_i and microbial biomass P pools, and constant rates of isotope exchange and respiration) to allow extrapolation of short-term exchange processes to the full length of the moist soil incubation experiments. IPD approaches can accommodate non-steady

state conditions as caused by flush effects and disturbances (Mooshammer et al., 2017) or as induced by addition
440 of organic matter. The equations to estimate IPD rates can easily be solved for soils where target pool
concentrations increase (net mineralization) or decrease (net immobilization) over time and where microbial
biomass P changes (Kirkham and Bartholomew, 1954), and do not necessitate constant pool sizes as wrongly
suggested previously (Di et al., 2000; Randhawa et al., 2005).

445 **4.1 Soil sterilization**

³³P IPD experiments in soils differ from the more common ¹⁵N IPD variants for gross N processes (Murphy et al.,
2003), since the persistence of abiotic P processes over time (Figs. 5 and 6) needs to be accounted for via the use
of sterile soils. Our data clearly show that the dynamics of abiotic ³³P processes change over time. Therefore, the
IPD rates in the sterile soils need to be measured over the same time period and under similar environmental
450 conditions as in the live soils. It is likely insufficient to extrapolate from short-term (100 min) batch incubations
run under very different conditions to correct for abiotic processes in the respective live moist soil incubations
over weeks. Bünemann et al. (2007) indicated that batch incubations (1:10 (w:v) soil: water suspensions) have
higher water-soluble and isotopically exchangeable P_i concentrations (measured as extractable P_i and as E values)
and tended to have higher tracer recoveries (measured as r/R, i.e. water-soluble ³³P_i recovered relative to total ³³P_i
455 added) compared to moist soil incubations. Incubation conditions should therefore also match between live and
sterile soils.

We chose autoclaving as the sterilization procedure as other procedures only reduce or eliminate
microbial activity (gamma irradiation, azide, mercuric chloride, toluene or chloroform treatment) but do not curtail
extracellular enzyme activities (Blankinship et al., 2014; Wolf et al., 1989; Tiwari et al., 1988; Oehl et al., 2001b).
460 Given that P_o mineralization is mediated by extracellular phosphatases, previous isotope experiments using short-
term batch experiments with or without microbicides or γ -irradiation therefore did not inhibit phosphatases and
therefore did not allow to separate abiotic and biotic processes of P_i release in soils. While application of
phosphatase inhibitors might be another viable option, we are only aware of one study testing this; application of
silver nanoparticles to soils showed a general inhibitory effect on soil enzymes (Shin et al., 2012). Previous tests
465 in our laboratory with two commercial phosphatase inhibitor cocktails (Sigma-Aldrich) at 10-fold of the
recommended final concentration did not significantly decrease IPD rates in two soils (data not shown), indicating
an insufficient inhibition of extracellular phosphatases. However, more rigorous tests of soil enzyme activities
with synthetic substrates (e.g. MUF-Pi) and of P mineralization based on ³³P-IPD using increasing concentrations
and different types of commercial phosphatase inhibitor cocktails might make clear whether this approach is viable
470 or not. In contrast, autoclaving soils twice was highly efficient in suppressing biological activities, and those soils
had no or very low extracellular enzyme activity and no residual microbial metabolic activity. Previous studies
showed (almost) total inhibition of hydrolytic enzyme activities (including phosphomonoesterases) by autoclaving,
in a wide range of arable, grassland and forest soils (Serrasolsas et al., 2008; Kedi et al., 2013; Blankinship et al.,
2014; Tiwari et al., 1988). Other studies demonstrated successful killing of bacterial and fungal cells in soils by
475 autoclaving (Carter et al., 2007; Blankinship et al., 2014; Serrasolsas and Khanna, 1995b; Alpehi and Scheu, 1993)).
Most importantly, the final step in P_o mineralization is catalyzed by phosphomonoesterases, which were fully
inactivated by autoclaving in all soils tested so far.

It must be noted that autoclaving could potentially alter the physicochemical properties of soils, thereby
affecting abiotic sorption-desorption kinetics. Despite this, in previous studies autoclaving up to two times and

480 steam sterilization did neither affect the cation exchange capacity, nor base saturation, soil surface area, contents
of total organic carbon and total nitrogen, and only slightly soil pH (Wolf et al., 1989; Tanaka et al.,
2003; Serrasolsas and Khanna, 1995b). Autoclaving might however weaken soil aggregates and therefore increase
the number of sites accessible for sorption-desorption processes that were previously hidden in aggregates.
However, we did not find clear support for or against this in the literature as autoclaving only weakly affected soil
485 aggregate size distribution, causing a 0.5 to 1% increase in clay-sized compared to silt-sized aggregates (Berns et
al., 2008). In contrast, aggregate stability and aggregation increased upon autoclaving in two other studies (Lostrario
et al., 1995; Salonijs et al., 1967). Effects of autoclaving on soil aggregation and soil P dynamics could be tested
by measuring P processes rates on intact aggregates <2 mm and after destroying them by ultrasonication or
grinding. In our study autoclaving caused a pulse of labile P into the available soil P pool due to the lysis of
490 microbial biomass (Fig. S1), as has also been demonstrated for P and N by Serrasolsas and Khanna (1995a, b).
Soil P_i concentrations increased significantly in the autoclaved soils studied here, but only by an average of 1.86-
fold in the three temperate soils and by 1.65-fold in the three tropical forest soils, which was in the range found by
others, e.g. 1.3- to 1.6-fold (Skipper and Westermann, 1973) and 1.5- to 1.6-fold (Anderson and Magdoff, 2005)
but lower than reported elsewhere, e.g. 2.6- to 11-fold (Serrasolsas and Khanna, 1995a). Autoclaving was also
495 demonstrated to increase the tracer recovery (r/R) and decrease the velocity of its decline over time as expected
due to loss of microbial biomass (Bünemann et al., 2007). Autoclaving therefore slightly affects the soil P_i pool,
but most likely has minor effects on its abiotic sorption/desorption dynamics while it inhibits biological reactions.
Nonetheless, the effects of increased P_i mobilization due to microbial lysis on P_i sorption-desorption could be
tested in sterile soils by adding increasing concentrations of non-labelled P_i alongside the $^{33}P_i$ tracer and then could
500 be corrected for in future ^{33}P -IPD experiments. As stated earlier, changes in P_i concentration caused by autoclaving
can easily be accounted for in IPD approaches, as long as abiotic process rates remain unaffected by the treatment.
However, the estimation of the contribution of abiotic and biotic processes is based on calculating the difference
in P fluxes between sterile and non-sterile soils. This assumes that biotic and abiotic fluxes are additive while there
is potential that both processes compete for available P_i . In this case we would overestimate abiotic process rates
505 in autoclaved soils, due to lack of competition by biotic processes. This could effectively cause an underestimation
of biotic processes i.e. organic P mineralization and microbial P_i uptake. To date we have no approach at hand to
cope with this potential bias. Overall, there is therefore potential for method improvement, particularly in terms of
using abiotic controls circumventing autoclaving (e.g. bacteriostats combined with phosphomonoesterase
inhibitors) or correcting for autoclave-induced changes in aggregation and in soil P_i content.

510

4.2 Soil P_i extraction using bicarbonate

Similar to ^{15}N IPD assays, where salt extractions are employed to target the available inorganic or organic N pool
(Murphy et al., 2003; Wanek et al., 2010; Hu et al., 2017), we focused on the potentially bio-available, salt-
extractable P_i pool that reflects the plant- (and microbial) accessible amount of soil P_i better (Fardeau et al.,
515 1988; Olsen et al., 1954; Horta and Torrent, 2007) than the water extractable P_i pool that is mostly assessed with
soil IEK methods. The applied 0.5 M $NaHCO_3$ extraction (pH 8.5, Olsen P) promotes the displacement of P_i (and
the extraction of labile P_o), particularly from Al-Fe (hydr)oxides and soil organic matter, by competition of
bicarbonate anions with P_i . The underlying process is an increase of the negative charge on surfaces and a decrease
of the concentration and activity of Ca^{2+} and Al^{3+} , thereby increasing P solubility in acid to alkaline soils (Horta
520 and Torrent, 2007; Schoenau and O'Halloran, 2008; Demaria et al., 2005). Several studies compared soil P tests like

Bray III, resin P, and Olsen-P to soil water P_i and plant P uptake in order to assess how well they reflect the available P_i pool. These studies demonstrated that soil tests like bicarbonate extractions (Olsen-P), resin P and DGT (diffusive gradients in thin films technique) closely resembled the SA values of P_i extracted by water or 10 mM $CaSO_4$ or from plants (Six et al., 2012;Fardeau et al., 1988;Demaria et al., 2005). Others further showed that isotopically exchangeable P_i in soil water extracts (E values) and those extracted by plant roots in plant growth experiments (L values) also were strongly related (Bühler et al., 2003;Frossard et al., 1994). Bicarbonate extracted 8- to 22-fold greater amounts of exchangeable P_i compared to water and SA of P_i in bicarbonate extracts reached 66-90% of the SA values measured in soil water extracts (Demaria et al., 2005). IPD approaches require fast extractions to quickly terminate the assay after 4 and 24 h, which renders water extractions (generally 16 h), resin P (16 h) and DGT (up to 48 h in low P soils; (Six et al., 2012)) impossible. Bicarbonate extractions only take 30-60 min and therefore represent a viable alternative. Moreover, it makes the IPD assay on average 8-fold more sensitive as a greater amount of exchangeable P_i is extracted by bicarbonate than with water (Kleinman et al., 2001). Underestimation of this labile P_i pool - even if specific activities thereof are correctly measured - also causes underestimation of IPD rates given that P_i concentrations linearly affect IPD rates according to IPD equations 1 and 2 above.

4.3 Microbial P dynamics

We observed very fast microbial P_i immobilization in live soils (within minutes; extraction started directly after tracer addition), causing net immobilization of ^{33}P by 3-38%. Similar results were reported within 1.5 to 4 h by others, ranging from 6-37% (Bünemann et al., 2012;Kellogg et al., 2006). This has two major repercussions: (i) rapid uptake might cause microbial P_i assimilation and efflux or exudation of $^{33}P_o$ metabolites without microbial death and turnover. However, the comparison between specific activities and ^{33}P recoveries of the direct measurement and after isobutanol fractionation (see below, and Fig. 3) showed that no significant release of microbial $^{33}P_o$ occurred during the 24 and 48 h incubations. The short extraction times used in this study also decrease the likelihood of significant hydrolysis of P_o compounds. (ii) Rapid microbial $^{33}P_i$ uptake clearly rules out the use of P_o mineralization assays that measure abiotic IEK in short-term batch experiments (100 min) without addition of a microbicide or without prior sterilization and then extrapolate these “abiotic” process rates to the full experimental duration.

Microbial P_i uptake can be derived indirectly as the difference in ^{33}P recovery between live and sterile soils (Fig. 5, this study), more directly by sECE (this study), or by parallel water or bicarbonate extraction with and without addition of liquid chloroform or hexane (measuring resin strip or extractable P_i), or by chloroform fumigation extraction (Bünemann et al., 2012;Oberson et al., 2001;Oehl et al., 2001a;Spohn and Kuzyakov, 2013). Microbial net ^{33}P immobilization measured by direct sECE was higher relative to the difference in ^{33}P immobilized in live minus sterile soils, pointing towards (i) overestimation of microbial net ^{33}P immobilization by sECE due to incomplete extraction of non-microbial $^{33}P_i$ by one-time bicarbonate extraction prior to sECE, or (ii) overestimation of abiotic sorption processes by autoclaving. In favor of (i) repeated extractions of soils with Bray I-extractant showed that soils continued to release P at lower rates in subsequent extractions after readily extractable P was removed by the first extraction (Serrasolsas et al., 2008;Messiga et al., 2014). Repeated extractions with bicarbonate also showed that the first extraction only removed 67-78% of the $^{33}P_i$ that was extractable with three consecutive extractions (D. Wasner, data not shown). In favor of (ii) (Kellogg et al., 2006) found higher net ^{33}P immobilization or sorption in sterile compared to live soils. This was interpreted as a lack of

microbial competition for P in sterile soils. However, we found a weak positive relationship ($R=0.749$, $P=0.087$; Table S1) between gross microbial P_i uptake and gross P_i sorption. This opposes the idea of strong competition between sorption and microbial uptake on the basis of gross process measurements. Another possible mechanism underlying (ii) could be changes in soil structure and reactive surfaces enhancing soil P sorption. Delineation of the causes could be performed by a comparison of sECE with liquid chloroform-fumigation extraction (CFE) i.e. parallel assessments of microbial ^{33}P uptake, using a comparison of ^{33}P in bicarbonate versus bicarbonate+liquid chloroform or bicarbonate+liquid hexane extracts. Given the continued extraction of P_i from exchangeable P_i pools in serial extraction tests, we propose that parallel determination of microbial P and ^{33}P by CFE is favorable relative to sequential extractions by sECE.

4.4 Comparison of isobutanol fractionation with direct measurements of P_i and ^{33}P activity

We showed that ^{33}P IPD assays can be performed specifically on the P_i pool using isobutanol fractionation in high P soils. However, due to low production or persistence of $^{33}P_o$, results closely conformed with measurements run without P_i - P_o fractionation by malachite green and direct $^{33}P_{total}$ estimates. This was ascertained for forest soils from French Guyana and Costa Rica, and for grassland soils from Austria and Germany (data not shown for French Guyana and Germany). Isobutanol fractionation has previously been applied in radiotracer studies on P dynamics in soils (Kellogg et al., 2006) and litter (Mooshammer et al., 2012), to ascertain the separation of P_i from any possible radiolabeled P_o contaminant, however without comparison to SA in unfractionated bicarbonate extracts. Oehl et al. (2001a) also applied isobutanol fractionation to water extracts of fumigated and control soils, demonstrating that with long extraction times (16 h), $^{33}P_i$ activities in water extracts with and without isobutanol fractionation were comparable. It was suggested that $^{33}P_o$ possibly released during fumigation was cleaved by soil phosphatases during extraction. This may not apply for short-term extractions (e.g. 0.5 M $NaHCO_3$ for 30 min, as used in this study) where hydrolysis by phosphatases would not necessarily occur due to short contact times. Measurements of ^{33}P isotope pool dilution in soils based on bicarbonate extracts can therefore be interchangeably be performed by (i) direct measurements of $^{33}P_{tot}$ and P_i in acidified bicarbonate extracts and after (ii) isobutanol fractionation on $^{33}P_i$ and P_i . However, this needs to be validated for other types of soil, and may change significantly after longer incubation periods (weeks), when microbial $^{33}P_i$ uptake, assimilation and turnover causes the release of $^{33}P_o$ into the soil. The short cut by performing direct measurements of P_i concentration and ^{33}P in acidified bicarbonate extracts comes along with 4- to 10-fold greater sensitivity of the malachite green assay relative to phosphomolybdate blue measurements of soil P_i . Another option to increase the measurement sensitivity for P_i (and possibly also for $^{33}P_i$) for strongly sorbing low-P soils has been adopted by Randriamanantsoa et al. (2013), based on concentration of the phosphomolybdate blue complex from a large volume of extract into a smaller volume of hexane, with subsequent phase separation (Murphy and Riley, 1962). This allowed to decrease limits of quantification of P_i by 66-fold compared to the classical Murphy-Riley protocol, and 14-fold compared to the malachite green procedure (Randriamanantsoa et al., 2013) but involves the handling of large volumes of radiolabeled extracts.

4.5 Time kinetics

During the first minutes, equilibration between tracer and tracee was not achieved, indicated by the enhanced extractability of added tracer ($^{33}P_i$) relative to more strongly bonded native tracee (soil exchangeable P_i). The fast process of equilibration caused very rapid declines in SA of P_i during the first few minutes. Thereafter, microbial

uptake and soil P fixation caused a rapid draw down of extractable $^{33}\text{P}_i$ and thereby a further decrease in the SA of soil P_i while soil P_i concentrations did not change after the initial phase of tracer-tracee equilibration (Fig. 4).
605 These processes slowed down within the first 1-2 h but did not cease, and declines in ^{33}P recoveries and in the SA of P_i occurred over the whole incubation period, in sterile as well as live soils. Thereafter time kinetics of IPD were relatively constant between 4 and 24 h for both, temperate and tropical soils, as shown by the linearity of the relationship in a plot of $\ln(\text{SA of } \text{P}_i)$ versus time. This linear relationship is conceptually different from the plot of $\log(\text{recovery, } r/R)$ versus $\log(\text{time})$ in short-term IEK batch experiments, that provides the parameter “n”, i.e.
610 the slope or the rate of decline in tracer recovery due to sorption over time (Bünemann, 2015). Based on constant IPD rates in the above-mentioned time interval we advise to run ^{33}P pool dilution experiments for an incubation period of 4 to 24 h. This time frame is well within the linear range, as it lies after the rapid abiotic equilibration, and is long enough to allow significant pool dilution to occur for sensitive measurements of organic P mineralization. Longer incubation times are not recommended due to the risk of $^{33}\text{P}_o$ release from dying microbes,
615 potentially causing a $^{33}\text{P}_i$ reflux through remineralization, violating a major assumption of IPD theory.

4.6 Comparison of P_o mineralization rates with published values

The detection limit of the IPD approach was $0.12 \mu\text{g P g}^{-1} \text{ soil dw d}^{-1}$. In comparison, the detection limits for gross P_o mineralization by the IEK approach were $0.20 \mu\text{g P g}^{-1} \text{ soil d}^{-1}$ by the modified protocol including hexane
620 concentration of phosphomolybdate blue for tropical soils (Randriamanantsoa et al., 2015) and $0.6\text{-}2.6 \mu\text{g P g}^{-1} \text{ soil d}^{-1}$ by the traditional IEK approach on temperate soils (Bünemann et al., 2007). Values of gross P_o mineralization measured via IPD in this study ranged between $0.08\text{-}0.15 \mu\text{g P g}^{-1} \text{ soil dw d}^{-1}$ in tropical forest soils and $0.48\text{-}2.03 \mu\text{g P g}^{-1} \text{ soil dw d}^{-1}$ in temperate grassland soils and were therefore well in the range of those compiled for IEK measurements by Bünemann (2015) for 14 different soils, including temperate arable, grassland
625 and forest soils ($0.1\text{-}12.6 \mu\text{g P g}^{-1} \text{ soil dw d}^{-1}$) and one tropical arable soil ($0.8 \mu\text{g P g}^{-1} \text{ soil dw d}^{-1}$). To date, highest gross P_o mineralization rates were reported for decomposing beech litter, i.e. $22.5\text{-}86.3 \mu\text{g P g}^{-1} \text{ soil dw d}^{-1}$ (Mooshammer et al., 2012). A direct comparison of the present IPD and the IEK approaches on the same soils might help to clarify how far the approaches really deviate or converge in their gross P_o mineralization rate estimates.

630

4.7 Physicochemical and biological controls on soil P_i processes

We found that gross P_o mineralization was strongly positively correlated to total soil P but not to soil organic P, soil organic C, soil texture or soil acid phosphatase activity. This indicates that gross P_o mineralization might rather be driven by total P than by soil enzyme activity, and that total soil P_o does not well represent the P_o fraction
635 accessible to soil phosphatases. A few studies demonstrated positive correlations between gross P_o mineralization and soil P_o (Lopez-Hernandez et al., 1998) or litter P_o (or its inverse C:P; (Mooshammer et al., 2012)). However, Wyngaard et al. (2016) did not find this relationship of gross P_o mineralization with total soil P_o but with the P_o content of the coarse soil fraction only, which points into a similar direction as our results. Moreover, P_o mineralization might be controlled rather by soil phosphodiesterases targeting DNA, RNA, teichoic acids and phospholipids, than by phosphomonoesterases that are responsible for the final extracellular dephosphorylation of
640 P_o . In contrast to our results, positive relationships were found between gross P_o mineralization and phosphomonoesterase activities in two studies (Spohn et al., 2013; Oehl et al., 2004), however not across studies (Bünemann, 2015). A larger set of soils varying in soil pH, texture and mineralogy might therefore provide better

insights into the controls of soil P_o mineralization, such as effects by extracellular phosphatase activity (phosphomonoesterases and phosphodiesterases), and the availability, stabilization and accessibility of organic P in soils, among others. Moreover, high P_i availability (i.e., bicarbonate P_i) strongly suppressed phosphomonoesterase activity in soils, causing a negative correlation between the enzyme activity and extractable P_i . In contrast, extractable P_i was positively related to gross P_o mineralization, indicating that high- P_i conditions suppressed phosphatase production but not P_o mineralization across these soils. This was also found as a positive correlation between gross P_o mineralization and water-extractable P_i by others (Schneider et al., 2017).

The contribution of gross P_o mineralization to total P_i supply including P_i desorption from exchangeable P_i pools and dissolution ranged between 25 and 74%, with a trend towards larger contributions in low-P tropical soils (35-74%) compared to temperate soils (25-51%). This clearly demonstrates that biological processes contribute importantly to the P_i supply in soils, particularly in low-P soils, as also pointed out by (Bünemann, 2015). In low-P forest soils biological processes were shown to dominate over physicochemical processes, while in P-rich forest soils abiotic processes controlled gross P_i supply rates (Bünemann et al., 2016). It was also found that the contributions of microbial processes decreased with soil depth, where in deep soils diffusive fluxes (i.e. gross P_i desorption) dominated the soil P_i supply due to low total P_o contents relative to total P (Achat et al., 2012;Achat et al., 2013).

Gross abiotic P_i release rates through desorption and dissolution were strongly positively related to total soil P and bicarbonate P_i , but not to other parameters such as soil pH, soil texture, and soil organic C content. In contrast to the weak effects of soil pH and texture on gross soil P_i supply, soil mineralogy and particularly oxalate-extractable Fe and Al as proxy for Fe-Al (hydr)oxides play a major role in controlling abiotic dynamics of phosphate ions in soils, across the full range from acidic to alkaline soils (Achat et al., 2016). Fe-Al (hydr)oxides provide large positively-charged surface areas in weathered soils that are highly reactive to phosphate ions, more so than clay minerals such as kaolinite, illite and others (Hinsinger, 2001;Regelink et al., 2015). Soil mineralogy might therefore provide further interesting insights into the controls of abiotic processes as demonstrated by (Achat et al., 2011;Achat et al., 2016), but can also affect P_o mineralization through strong effects on the sorption strength of organic matter and of P_o compounds. Moreover, the elsewhere reported positive relations of P_i availability and P_i desorption with soil organic C contents was explained by competitive sorption of P_i and SOC or DOC to reactive surfaces such as positively charged metal (hydr)oxides (Regelink et al., 2015;Achat et al., 2016).

Gross P_i sorption rates exceeded gross P_i desorption rates approximately 2-fold but both were strongly related, indicating close and rapid cycling of available P_i through sorption-desorption processes. The observed rates indicate that soils immobilized more P_i than they mobilized by abiotic processes, causing an intermediate draw down of available P_i pools. Two processes work against this draw down of P_i in soils, i.e. P_o mineralization and microbial P release through turnover and lysis. Moreover, plants (and microbes) might also desorb this sorbed P_i by release of phytosiderophores and organic acids and thereby replenish P_i and re-inject it in the organic P cycle. Similar to the soil C-N cycle we might also expect an active “bank mechanism” regulating nutrient and C sequestration in soils (Fontaine et al., 2011). At high nutrient availability priming effects are low, allowing the sequestration of nutrients and SOC build-up. At low nutrient availability microbes (and plants) release nutrients from SOM and from mineral surfaces stimulated by root exudates, effectively mining inorganic and organic P stored in soils.

The strong positive relationship between gross P_i sorption rates and soil total P, soil total P_i and bicarbonate soil P_i , and the lack of relationship with soil pH, soil organic C, clay content and soil texture highlights

685 again that specific soil minerals, particularly metal (hydr)oxides and to a lesser extent clay minerals such kaolinite,
factors not fully captured by soil pH and soil texture alone, are responsible for P_i sorption in soils (Regelink et al.,
2015). In IEK experiments it was found that the rate of abiotic P_i depletion from soil solution through sorption
was positively related to Al-Fe (hydr)oxide content and negatively to soil organic C divided by Al and Fe oxide
content (Achat et al., 2016; Tran et al., 1988). The strong negative relation between abiotic net P_i immobilization
690 and soil pH re-confirms that strongly weathered, acid tropical soils have a higher P sorption and fixation capacity
than temperate soils.

Finally, gross microbial P_i uptake rates were directly proportional to microbial biomass P measured by
sECE. We also found greater P_i immobilization potentials through sorptive reactions (28-92%) than through
biological sinks (5-37%) in the soils studied here. The importance of rapid net uptake of tracer by soil microbes
695 has been demonstrated also by other studies, e.g. (Bünemann et al., 2012). However, the presented IPD approach
for the first time allowed to estimate gross rates of microbial P_i uptake in addition to net microbial P_i
immobilization. Gross rates of microbial uptake were calculated from the IPD approach, not necessitating the
application of any extraction factor to calculate microbial biomass P from chloroform-labile P (k_{EP}-factor). The
use of extraction factors becomes necessary when studying net P_i uptake over prolonged time periods in tracer
700 experiments and for correction of net P_o mineralization rates (Bünemann, 2015; Bünemann et al., 2007).

4.8 Application and modeling

The combination of this IPD assay with advanced numerical modeling approaches, as applied by Müller and
Bünemann (2014), might further enhance the precision of estimates of simultaneously occurring soil P cycle
705 processes and thereby advance the knowledge of major controls of the transformations and fluxes of this important
nutrient in terrestrial ecosystems. There is an ever-increasing need of high quality data on soil P processes, even
more so to calibrate terrestrial biogeochemical models and incorporate nutrient controls on plant productivity in
global models. This IPD approach may provide highly important quantitative data to implement soil P cycling
processes into global biogeochemical models. This will further enhance our current understanding of nutrient
710 controls on the global terrestrial C cycle and improve our capabilities to predict future changes by increasing
discrepancies in N and P inputs into the terrestrial biosphere.

Data availability. The data of the different experiments are freely available upon request from the corresponding
author.

715 **Author contributions.** The project was conceived and supervised by WW. DZ, JP and DW performed the
measurements and data evaluation. WW wrote the manuscript with contributions from all coauthors.

Competing interests. The authors declare that they have no conflict of interest.

720 **Acknowledgements.** We are indebted to the Isotope Laboratory managers for access and training (Virginie
Canoine, Markus Schmid).

725 **References**

- Achat, D. L., Bakker, M. R., Saur, E., Pellerin, S., Augusto, L., and Morel, C.: Quantifying gross mineralisation of P in dead soil organic matter: Testing an isotopic dilution method, *Geoderma*, 158, 163-172, 10.1016/j.geoderma.2010.04.027, 2010.
- 730 Achat, D. L., Augusto, L., Morel, C., and Bakker, M. R.: Predicting available phosphate ions from physical-chemical soil properties in acidic sandy soils under pine forests, *J. Soils Sediments*, 11, 452-466, 10.1007/s11368-010-0329-9, 2011.
- Achat, D. L., Augusto, L., Bakker, M. R., Gallet-Budynek, A., and Morel, C.: Microbial processes controlling P availability in forest spodosols as affected by soil depth and soil properties, *Soil Biol. Biochem.*, 44, 39-48, 10.1016/j.soilbio.2011.09.007, 2012.
- 735 Achat, D. L., Bakker, M. R., Augusto, L., and Morel, C.: Contributions of microbial and physical-chemical processes to phosphorus availability in Podzols and Arenosols under a temperate forest, *Geoderma*, 211, 18-27, 10.1016/j.geoderma.2013.07.003, 2013.
- Achat, D. L., Pousse, N., Nicolas, M., Bredoire, F., and Augusto, L.: Soil properties controlling inorganic phosphorus availability: general results from a national forest network and a global compilation of the literature, *Biogeochemistry*, 127, 255-272, 10.1007/s10533-015-0178-0, 2016.
- 740 Alpei, J., and Scheu, S.: Effects of biocidal treatments on biological and nutritional properties of a mull-structured woodland soil, *Geoderma*, 56, 435-448, 10.1016/0016-7061(93)90125-5, 1993.
- Anderson, B. H., and Magdoff, F. R.: Autoclaving soil samples affects algal-available phosphorus, *J. Environ. Qual.*, 34, 1958-1963, 10.2134/jeq2005.0024, 2005.
- 745 Berns, A. E., Philipp, H., Narres, H. D., Burauel, P., Vereecken, H., and Tappe, W.: Effect of gamma-sterilization and autoclaving on soil organic matter structure as studied by solid state NMR, UV and fluorescence spectroscopy, *European Journal of Soil Science*, 59, 540-550, 10.1111/j.1365-2389.2008.01016.x, 2008.
- Blankinship, J. C., Becerra, C. A., Schaeffer, S. M., and Schimel, J. P.: Separating cellular metabolism from exoenzyme activity in soil organic matter decomposition, *Soil Biol. Biochem.*, 71, 68-75, 10.1016/j.soilbio.2014.01.010, 2014.
- Booth, M. S., Stark, J. M., and Rastetter, E.: Controls on nitrogen cycling in terrestrial ecosystems: A synthetic analysis of literature data, *Ecological Monographs*, 75, 139-157, 2005.
- 755 Braun, J., Mooshammer, M., Wanek, W., Prommer, J., Walker, T. W. N., Rütting, T., and Richter, A.: Full N-15 tracer accounting to revisit major assumptions of N-15 isotope pool dilution approaches for gross nitrogen mineralization, *Soil Biol. Biochem.*, 117, 16-26, 10.1016/j.soilbio.2017.11.005, 2018.
- Bühler, S., Oberson, A., Sinaj, S., Friesen, D. K., and Frossard, E.: Isotope methods for assessing plant available phosphorus in acid tropical soils, *Eur. J. Soil Sci.*, 54, 605-616, 10.1046/j.1365-2389.2003.00542.x, 2003.
- 760 Bünemann, E. K., Marschner, P., McNeill, A. M., and McLaughlin, M. J.: Measuring rates of gross and net mineralisation of organic phosphorus in soils, *Soil Biol. Biochem.*, 39, 900-913, 10.1016/j.soilbio.2006.10.009, 2007.
- Bünemann, E. K., Oberson, A., Liebisch, F., Keller, F., Annaheim, K. E., Huguenin-Elie, O., and Frossard, E.: Rapid microbial phosphorus immobilization dominates gross phosphorus fluxes in a grassland soil with low inorganic phosphorus availability, *Soil Biol. Biochem.*, 51, 84-95, 10.1016/j.soilbio.2012.04.012, 2012.
- 765 Bünemann, E. K.: Assessment of gross and net mineralization rates of soil organic phosphorus - A review, *Soil Biol. Biochem.*, 89, 82-98, 10.1016/j.soilbio.2015.06.026, 2015.
- Bünemann, E. K., Augstburger, S., and Frossard, E.: Dominance of either physicochemical or biological phosphorus cycling processes in temperate forest soils of contrasting phosphate availability, *Soil Biol. Biochem.*, 101, 85-95, 10.1016/j.soilbio.2016.07.005, 2016.
- 770 Carter, D. O., Yellowlees, D., and Tibbett, M.: Autoclaving kills soil microbes yet soil enzymes remain active, *Pedobiologia*, 51, 295-299, 10.1016/j.pedobi.2007.05.002, 2007.
- Condon, L. M., Moir, J. O., Tiessen, H., and Stewart, J. W. B.: CRITICAL-EVALUATION OF METHODS FOR DETERMINING TOTAL ORGANIC PHOSPHORUS IN TROPICAL SOILS, *Soil Science Society of America Journal*, 54, 1261-1266, 10.2136/sssaj1990.03615995005400050010x, 1990.
- 775 D'Angelo, E., Crutchfield, J., and Vandiviere, M.: Rapid, sensitive, microscale determination of phosphate in water and soil, *J. Environ. Qual.*, 30, 2206-2209, 2001.
- Demaria, P., Flisch, R., Frossard, E., and Sinaj, S.: Exchangeability of phosphate extracted by four chemical methods, *Journal of Plant Nutrition and Soil Science-Zeitschrift Fur Pflanzenernahrung Und Bodenkunde*, 168, 89-93, 10.1002/jpin.200421463, 2005.
- 780 Di, H. J., Condon, L. M., and Frossard, E.: Isotope techniques to study phosphorus cycling in agricultural and forest soils: A review, *Biol. Fertil. Soils*, 24, 1-12, 10.1007/bf01420213, 1997.
- Di, H. J., Cameron, K. C., and McLaren, R. G.: Isotopic dilution methods to determine the gross transformation rates of nitrogen, phosphorus, and sulfur in soil: a review of the theory, methodologies, and limitations, *Aust. J. Soil Res.*, 38, 213-230, 2000.

- 785 Fardeau, J. C., Morel, C., and Boniface, R.: Why the Olsen method should be used to estimate available soil phosphorus?, *Agronomie*, 8, 577-584, 10.1051/agro:19880702, 1988.
- Fardeau, J. C.: Le phosphore assimilable des sols : sa représentation par un modèle fonctionnel à plusieurs compartiments *Agronomie*, 13, 317-331, 1993.
- 790 Fontaine, S., Henault, C., Aamor, A., Bdioui, N., Bloor, J. M. G., Maire, V., Mary, B., Revaillet, S., and Maron, P. A.: Fungi mediate long term sequestration of carbon and nitrogen in soil through their priming effect, *Soil Biology & Biochemistry*, 43, 86-96, 10.1016/j.soilbio.2010.09.017, 2011.
- Frossard, E., Fardeau, J. C., Brossard, M., and Morel, J. L.: Soil isotopically exchangeable phosphorus - a comparison between E and L-values, *Soil Sci. Soc. Am. J.*, 58, 846-851, 1994.
- 795 Frossard, E., Achat, D. L., Bernasconi, S. M., Bünemann, E. K., Fardeau, J. C., Jansa, J., Morel, C., Rabeharisoa, L., Randriamanantsoa, L., Sinaj, S., Tamburini, F., and Oberson, A.: The Use of Tracers to Investigate Phosphate Cycling in Soil-Plant Systems, in: *Phosphorus in Action: Biological Processes in Soil Phosphorus Cycling*, edited by: Bunemann, E. K., Oberson, A., and Frossard, E., *Soil Biology*, 59-91, 2011.
- Goldberg, S., and Sposito, G.: On the mechanism of specific phosphate-adsorption by hydroxylated mineral surfaces - a review, *Commun. Soil Sci. Plant Anal.*, 16, 801-821, 10.1080/00103628509367646, 1985.
- 800 Green, V. S., Stott, D. E., and Diack, M.: Assay for fluorescein diacetate hydrolytic activity: Optimization for soil samples, *Soil Biol. Biochem.*, 38, 693-701, 10.1016/j.soilbio.2005.06.020, 2006.
- Hart, S. C., Nason, G. E., Myrold, D. D., and Perry, D. A.: Dynamics of gross nitrogen transformations in an old-growth forest: the carbon connection, *Ecology*, 75, 880-891, 1994.
- 805 Hinsinger, P.: Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: a review, *Plant Soil*, 237, 173-195, 10.1023/a:1013351617532, 2001.
- Horta, M. D., and Torrent, J.: The Olsen P method as an agronomic and environmental test for predicting phosphate release from acid soils, *Nutrient Cycling in Agroecosystems*, 77, 283-292, 10.1007/s10705-006-9066-2, 2007.
- 810 Hu, Y. T., Zheng, Q., and Wanek, W.: Flux Analysis of Free Amino Sugars and Amino Acids in Soils by Isotope Tracing with a Novel Liquid Chromatography/High Resolution Mass Spectrometry Platform, *Analytical Chemistry*, 89, 9192-9200, 10.1021/acs.analchem.7b01938, 2017.
- Jayachandran, K., Schwab, A. P., and Hetrick, B. A. D.: Partitioning dissolved inorganic and organic phosphorus using acidified molybdate and isobutanol, *Soil Sci. Soc. Am. J.*, 56, 762-765, 1992.
- 815 Kedi, B., Sei, J., Quiquampoix, H., and Staunton, S.: Persistence of catalytic activity of fungal phosphatases incubated in tropical soils, *Soil Biol. Biochem.*, 56, 69-74, 10.1016/j.soilbio.2012.02.005, 2013.
- Kellogg, L. E., Bridgham, S. D., and Lopez-Hernandez, D.: A comparison of four methods of measuring gross phosphorus mineralization, *Soil Sci. Soc. Am. J.*, 70, 1349-1358, 10.2135/sssaj2005.0300, 2006.
- Kirkham, D., and Bartholomew, W. V.: Equations for following nutrient transformations in soil, utilizing tracer data, *Soil Science Society of America Proceedings*, 18, 33-34, 1954.
- 820 Kleinman, P. J. A., Sharpley, A. N., Gartley, K., Jarrell, W. M., Kuo, S., Menon, R. G., Myers, R., Reddy, K. R., and Skogley, E. O.: Interlaboratory comparison of soil phosphorus extracted by various soil test methods, *Communications in Soil Science and Plant Analysis*, 32, 2325-2345, 10.1081/css-120000376, 2001.
- Lopez-Hernandez, D., Brossard, M., and Frossard, E.: P-isotopic exchange values in relation to Po mineralisation in soils with very low P-sorbing capacities, *Soil Biol. Biochem.*, 30, 1663-1670, 10.1016/s0038-0717(97)00255-1, 1998.
- 825 Lotrario, J. B., Stuart, B. J., Lam, T., Arands, R. R., Oconnor, O. A., and Kosson, D. S.: EFFECTS OF STERILIZATION METHODS ON THE PHYSICAL CHARACTERISTICS OF SOIL - IMPLICATIONS FOR SORPTION ISOTHERM ANALYSES, *Bull. Environ. Contam. Toxicol.*, 54, 668-675, 1995.
- 830 Mander, C., Wakelin, S., Young, S., Condron, L., and O'Callaghan, M.: Incidence and diversity of phosphate-solubilising bacteria are linked to phosphorus status in grassland soils, *Soil Biol. Biochem.*, 44, 93-101, 10.1016/j.soilbio.2011.09.009, 2012.
- Marx, M. C., Wood, M., and Jarvis, S. C.: A microplate fluorimetric assay for the study of enzyme diversity in soils, *Soil Biol. Biochem.*, 33, 1633-1640, 2001.
- 835 Messiga, A. J., Ba, Y. X., Ziadi, N., Belanger, G., and Lafond, J.: Assessing the depletion of soil P following sequential extractions with Mehlich-3 and Olsen solutions, *Arch. Agron. Soil Sci.*, 60, 1445-1458, 10.1080/03650340.2014.884709, 2014.
- Miller, W., and Miller, D.: A micropipette method for soil mechanical analysis, *Communications in Soil Science and Plant Analysis*, 18, 1-15, 1987.
- 840 Mooshammer, M., Wanek, W., Schneckner, J., Wild, B., Leitner, S., Hofhansl, F., Blochl, A., Hammerle, I., Frank, A. H., Fuchslueger, L., Keiblinger, K. M., Zechmeister-Boltenstern, S., and Richter, A.: Stoichiometric controls of nitrogen and phosphorus cycling in decomposing beech leaf litter, *Ecology*, 93, 770-782, 2012.
- Mooshammer, M., Hofhansl, F., Frank, A. H., Wanek, W., Hammerle, I., Leitner, S., Schneckner, J., Wild, B., Watzka, M., Keiblinger, K. M., Zechmeister-Boltenstern, S., and Richter, A.: Decoupling of microbial carbon, nitrogen, and phosphorus cycling in response to extreme temperature events, *Sci. Adv.*, 3, 10.1126/sciadv.1602781, 2017.
- 845

- Müller, C., and Bünemann, E. K.: A P-33 tracing model for quantifying gross P transformation rates in soil, *Soil Biol. Biochem.*, 76, 218-226, 10.1016/j.soilbio.2014.05.013, 2014.
- 850 Murphy, D. V., Recous, S., Stockdale, E. A., Fillery, I. R. P., Jensen, L. S., Hatch, D. J., and Goulding, K. W. T.: Gross nitrogen fluxes in soil: Theory, measurement and application of 15N pool dilution techniques, *Advances in Agronomy*, 79, 69-118, 2003.
- Murphy, J., and Riley, J. P.: A modified single solution method for determination of phosphate in natural waters, *Analytica Chimica Acta*, 26, 31-&, 1962.
- 855 O'Halloran, I. P., and Cade-Menun, B. J.: Chapter 24. Total and Organic Phosphorus, in: *Soil Sampling and Methods of Analysis*. Second Edition., edited by: Carter, M. R., and Gregorich, E. G., CRC Press, Taylor & Francis, Boca Raton, FL, U.S.A., 265-291, 2008.
- Oberson, A., Friesen, D. K., Rao, I. M., Buhler, S., and Frossard, E.: Phosphorus Transformations in an Oxisol under contrasting land-use systems: The role of the soil microbial biomass, *Plant and Soil*, 237, 197-210, 10.1023/a:1013301716913, 2001.
- 860 Oehl, F., Oberson, A., Probst, M., Fliessbach, A., Roth, H. R., and Frossard, E.: Kinetics of microbial phosphorus uptake in cultivated soils, *Biol. Fertil. Soils*, 34, 31-41, 2001a.
- Oehl, F., Oberson, A., Sinaj, S., and Frossard, E.: Organic phosphorus mineralization studies using isotopic dilution techniques, *Soil Sci. Soc. Am. J.*, 65, 780-787, 2001b.
- 865 Oehl, F., Frossard, E., Fliessbach, A., Dubois, D., and Oberson, A.: Basal organic phosphorus mineralization in soils under different farming systems, *Soil Biol. Biochem.*, 36, 667-675, 10.1016/j.soilbio.2003.12.010, 2004.
- Ohno, T., and Zibilske, L. M.: Determination of low concentrations of phosphorus in soil extracts using malachite green, *Soil Sci. Soc. Am. J.*, 55, 892-895, 1991.
- 870 Olsen, S. R., Cole, C. V., Watanabe, F. S., and Dean, L. A.: Estimation of Available Phosphorus in Soils by Extraction with Sodium Bicarbonate, USDA Circ. 939, US. Government Printing Office, Washington, DC, 1954.
- Penuelas, J., Sardans, J., Rivas-Ubach, A., and Janssens, I. A.: The human-induced imbalance between C, N and P in Earth's life system, *Global Change Biology*, 18, 3-6, 10.1111/j.1365-2486.2011.02568.x, 2012.
- 875 Penuelas, J., Poulter, B., Sardans, J., Ciais, P., van der Velde, M., Bopp, L., Boucher, O., Godderis, Y., Hinsinger, P., Llusia, J., Nardin, E., Vicca, S., Obersteiner, M., and Janssens, I. A.: Human-induced nitrogen-phosphorus imbalances alter natural and managed ecosystems across the globe, *Nature Communications*, 4, 2934, 10.1038/ncomms3934, 2013.
- 880 Randhawa, P. S., Condron, L. M., Di, H. J., Sinaj, S., and McLenaghan, R. D.: Effect of green manure addition on soil organic phosphorus mineralisation, *Nutr. Cycl. Agroecosyst.*, 73, 181-189, 10.1007/s10705-005-0593-z, 2005.
- Randriamanantsoa, L., Morel, C., Rabeharisoa, L., Douzet, J. M., Jansa, J., and Frossard, E.: Can the isotopic exchange kinetic method be used in soils with a very low water extractable phosphate content and a high sorbing capacity for phosphate ions?, *Geoderma*, 200, 120-129, 10.1016/j.geoderma.2013.01.019, 2013.
- 885 Randriamanantsoa, L., Frossard, E., Oberson, A., and Bünemann, E. K.: Gross organic phosphorus mineralization rates can be assessed in a Ferralsol using an isotopic dilution method, *Geoderma*, 257, 86-93, 10.1016/j.geoderma.2015.01.003, 2015.
- Reed, S. C., Yang, X. J., and Thornton, P. E.: Incorporating phosphorus cycling into global modeling efforts: a worthwhile, tractable endeavor, *New Phytol.*, 208, 324-329, 10.1111/nph.13521, 2015.
- 890 Regelink, I. C., Weng, L., Lair, G. J., and Comans, R. N. J.: Adsorption of phosphate and organic matter on metal (hydr)oxides in arable and forest soil: a mechanistic modelling study, *Eur. J. Soil Sci.*, 66, 867-875, 10.1111/ejss.12285, 2015.
- Richardson, A. E., and Simpson, R. J.: Soil Microorganisms Mediating Phosphorus Availability, *Plant Physiology*, 156, 989-996, 10.1104/pp.111.175448, 2011.
- 895 Ryan, P. R., Delhaize, E., and Jones, D. L.: Function and mechanism of organic anion exudation from plant roots, *Annu. Rev. Plant Physiol. Plant Molec. Biol.*, 52, 527-560, 10.1146/annurev.arplant.52.1.527, 2001.
- Salonius, P. O., Robinson, J. B., and Chase, F. E.: A COMPARISON OF AUTOCLAVED AND GAMMA-IRRADIATED SOILS AS MEDIA FOR MICROBIAL COLONIZATION EXPERIMENTS, *Plant and Soil*, 27, 239-&, 10.1007/bf01373392, 1967.
- 900 Schneider, K. D., Voroney, R. P., Lynch, D. H., Oberson, A., Frossard, E., and Bünemann, E. K.: Microbially-mediated P fluxes in calcareous soils as a function of water-extractable phosphate, *Soil Biol. Biochem.*, 106, 51-60, 10.1016/j.soilbio.2016.12.016, 2017.
- Schnurer, J., and Rosswall, T.: Fluorescein diacetate hydrolysis as a measure of total microbial activity in soil and litter, *Appl. Environ. Microbiol.*, 43, 1256-1261, 1982.
- 905 Schoenau, J. J., and O'Halloran, I. P.: Sodium Bicarbonate-Extractable Phosphorus, in: *Soil Sampling and Methods of Analysis*, edited by: Carter, M. R., and Gregorich, E. G., CRC Press, Boca Raton, FL, 89-94, 2008.

- Serrasolsas, I., and Khanna, P. K.: Changes in heated and autoclaved forest soils of SE Australia. 2. Phosphorus and phosphatase activity, *Biogeochemistry*, 29, 25-41, 1995a.
- 910 Serrasolsas, I., and Khanna, P. K.: Changes in heated and autoclaved forest soils of SE Australia. 1. Carbon and nitrogen, *Biogeochemistry*, 29, 3-24, 1995b.
- Serrasolsas, I., Romanya, J., and Khanna, P. K.: Effects of heating and autoclaving on sorption and desorption of phosphorus in some forest soils, *Biol. Fertil. Soils*, 44, 1063-1072, 10.1007/s00374-008-0301-7, 2008.
- 915 Setia, R., Verma, S. L., and Marschner, P.: Measuring microbial biomass carbon by direct extraction - Comparison with chloroform fumigation-extraction, *European Journal of Soil Biology*, 53, 103-106, 10.1016/j.ejsobi.2012.09.005, 2012.
- Shin, Y.-J., Kwak, J. I., and An, Y.-J.: Evidence for the inhibitory effects of silver nanoparticles on the activities of soil exoenzymes, *Chemosphere*, 88, 524-529, 10.1016/j.chemosphere.2012.03.010, 2012.
- 920 Sirova, D., Rejmankova, E., Carlson, E., and Vrba, J.: Current standard assays using artificial substrates overestimate phosphodiesterase activity, *Soil Biol. Biochem.*, 56, 75-79, 10.1016/j.soilbio.2012.02.008, 2013.
- Six, L., Pypers, P., Degryse, F., Smolders, E., and Merckx, R.: The performance of DGT versus conventional soil phosphorus tests in tropical soils - An isotope dilution study, *Plant Soil*, 359, 267-279, 10.1007/s11104-012-1192-9, 2012.
- 925 Skipper, H. D., and Westermann, D. T.: Comparative effects of propylene oxide, sodium azide, and autoclaving on selected soil properties, *Soil Biol. Biochem.*, 5, 409-414, 1973.
- Spohn, M., Ermak, A., and Kuzyakov, Y.: Microbial gross organic phosphorus mineralization can be stimulated by root exudates - A P-33 isotopic dilution study, *Soil Biol. Biochem.*, 65, 254-263, 10.1016/j.soilbio.2013.05.028, 2013.
- 930 Spohn, M., and Kuzyakov, Y.: Phosphorus mineralization can be driven by microbial need for carbon, *Soil Biol. Biochem.*, 61, 69-75, 10.1016/j.soilbio.2013.02.013, 2013.
- Tanaka, S., Kobayashi, T., Iwasaki, K., Yamane, S., Maeda, K., and Sakurai, K.: Properties and metabolic diversity of microbial communities in soils treated with steam sterilization compared with methyl bromide and chloropicrin fumigations, *Soil Sci. Plant Nutr.*, 49, 603-610, 2003.
- 935 Tiwari, S. C., Tiwari, B. K., and Mishra, R. R.: Enzyme activities in soils - effects of leaching, ignition, autoclaving and fumigation, *Soil Biol. Biochem.*, 20, 583-585, 10.1016/0038-0717(88)90079-x, 1988.
- Tran, T. S., Fardeau, J. C., and Giroux, M.: Effects of soil properties on plant-available phosphorus determined by the isotopic dilution phosphorus-32 method, *Soil Sci. Soc. Am. J.*, 52, 1383-1390, 1988.
- 940 Turner, B. L., Condon, L. M., Richardson, S. J., Peltzer, D. A., and Allison, V. J.: Soil organic phosphorus transformations during pedogenesis, *Ecosystems*, 10, 1166-1181, 2007.
- Vitousek, P. M., and Farrington, H.: Nutrient limitation and soil development: Experimental test of a biogeochemical theory, *Biogeochemistry*, 37, 63-75, 1997.
- 945 Wanek, W., Mooshammer, M., Blöchl, A., Hanreich, A., and Richter, A.: Determination of gross rates of amino acid production and immobilization in decomposing leaf litter by a novel ¹⁵N isotope pool dilution technique, *Soil Biology and Biochemistry*, 42, 1293-1302, 2010.
- Wolf, D. C., Dao, T. H., Scott, H. D., and Lavy, T. L.: Influence of sterilization methods on selected soil microbiological, physical, and chemical properties, *J. Environ. Qual.*, 18, 39-44, 1989.
- 950 Wyngaard, N., Cabrera, M. L., Jarosch, K. A., and Bünemann, E. K.: Phosphorus in the coarse soil fraction is related to soil organic phosphorus mineralization measured by isotopic dilution, *Soil Biol. Biochem.*, 96, 107-118, 10.1016/j.soilbio.2016.01.022, 2016.
- Yang, X., and Post, W. M.: Phosphorus transformations as a function of pedogenesis: A synthesis of soil phosphorus data using Hedley fractionation method, *Biogeosciences*, 8, 2907-2916, 10.5194/bg-8-2907-2011, 2011.
- 955 Yang, X., Post, W. M., Thornton, P. E., and Jain, A.: The distribution of soil phosphorus for global biogeochemical modeling, *Biogeosciences*, 10, 2525-2537, 10.5194/bg-10-2525-2013, 2013.

Table 1. Comparison of traditional isotope exchange kinetic (IEK) experiments and the novel isotope pool dilution (IPD) approach to measure organic P mineralization.

Factor and approach	Isotope exchange (IEK)	Isotope pool dilution (IPD)
Tracer addition and incubation period	^{33}P , ^{32}P ; Several time points across several days to weeks and months	^{33}P , (^{32}P); Two time points at 4 and 24 hours
Measured P pool	Water-extractable P_i	Bicarbonate-extractable P_i and P_o
Abiotic controls	Abiotic controls measured in batch experiment with live soil: 100 min P_i exchange experiment in soil suspension 1:10 (soil: water), $\pm\text{HgCl}_2$ or sodium azide; microbial contributions in short-term experiment often not accounted for	Duplicate autoclaving for abiotic controls to kill microbial biomass and extracellular enzymes; treatment of abiotic controls similar as live soils in terms of tracer addition, incubation period and extraction
Microbial processes in abiotic controls	Microbial biomass active in abiotic controls if no microbicide added, extracellular phosphatases fully active (causing organic P mineralization in abiotic controls)	Microbial biomass and phosphatases deactivated by autoclaving (no/almost no P mineralization occurring in abiotic controls)
Pre-incubation of soils to equilibrate to moisture and temperature	Yes (to constant respiration – equilibrium conditions necessary)	Yes (not necessary)
Change in soil structure and P availability	No (if no microbicide is added)	Potentially yes, as autoclaving might increase available P by death of microbial biomass and soil structure might change by autoclaving
Numerical solution for P_o mineralization	Isotopically exchangeable P within t minutes ($E_{(t)}$) derived as the inverse of the relative specific activity of phosphate in soil solution (water extractable P_i) over time in live soils. $E'_{(t)}$ derived for abiotic controls extrapolated from 100 min to length of full experiment, graphical solution of corrected data following (Fardeau, 1993). Differences in $E'_{(t)}$ and $E_{(t)}$ estimate gross P_o mineralization	Calculation of IPD influx rates based on mass/isotope balance equations derived by (Kirkham and Bartholomew, 1954) for tracer: tracee experiments. Gross P_o mineralization calculated as difference of IPD influx rates of live soils minus abiotic controls

965 Table 2. Soil characterization of three temperate grassland soils (soil 2, 4, and 6) and three tropical lowland forest soils (soil 3, 5, and 7).

Parameter	Unit	Temperate soils			Tropical soils		
		2	4	6	3	5	7
Soil pH (10 mM CaCl ₂)		6.30	6.25	6.80	4.15	4.15	4.10
Clay	(%)	16.8	14.1	2.76	4.12	19.6	26.2
Silt	(%)	59.2	24.4	40.6	88.0	72.8	70.1
Sand	(%)	24.0	61.4	56.6	7.92	7.61	3.74
Total organic C	(mg g ⁻¹ DW)	48.3	126.7	60.3	26.4	30.8	28.5
Total N	(mg g ⁻¹ DW)	3.35	5.03	2.32	2.17	2.57	2.27
Total P (TP)	(mg g ⁻¹ DW)	0.82	0.44	0.51	0.14	0.17	0.09
Total organic P (TP _o)	(mg g ⁻¹ DW)	0.40	0.25	0.11	0.09	0.13	0.07
Soil P _i	(μg g ⁻¹ DW)	15.1	4.23	5.59	0.56	0.49	0.37
TP _o of TP	(%)	49.1	56.5	22.3	64.2	75.7	76.4
Soil C:N		14.4	25.2	26.0	12.1	12.0	12.5
Soil C:TP _o		121	507	548	293	237	406
Soil N:TP _o		8.4	20.1	21.1	24.1	19.8	32.5
Phosphatase	(nmol MUF g ⁻¹ DW h ⁻¹)	256	316	233	1396	1698	2346

Figure 1. Schematic representation of (A) major fluxes of soil P processes controlling the availability of inorganic P (P_i) in soils, and of (B) the isotope pool dilution principle showing influxes of unlabeled P_i (^{31}P) into the available P_i pool labelled by a spike of $^{33}P_i$, and efflux of P_i at the ratio of $^{33}P_i:^{31}P_i$ as present in the target pool. Biotic and abiotic processes of influx and efflux are presented. This causes (C) a decline in the specific activity of P_i i.e. $^{33}P_i:^{31}P_i$ declines over time in sterile soils (abiotic processes only) and live soils (biotic plus abiotic processes), allowing to derive biotic contributions to overall gross fluxes. TP_i ...total soil P_i , TP_o ...total organic P, TP_i includes occluded and fixed P as well as primary mineral P, TP_o includes occluded P_o in aggregates. Avail...available. P_i desorption includes P_i dissolution from minerals, and P_i sorption includes P_i precipitation.








Figure 2. Schematic overview of the final isotope pool dilution (IPD) procedure. Abbreviations: MG, malachite green procedure and MR, Murphy-Riley procedure to measure P_i concentrations; LSC, liquid scintillation counting to measure radioactivity in extracts.



985 Figure 3. Response of soil enzyme activities to autoclaving: Percentage inhibition of (A) fluorescein diacetate (FDA) hydrolysis as a proxy for the inhibition of live, cell-bound microbial enzyme activity and of (B) MUF-phosphomonoesterase activity as a proxy for the inhibition of extracellular enzyme activity. Temperate grassland soils (2, 4, 6) and tropical forest soils (3, 5, 7) were tested. Two-way ANOVA was calculated to test for the factors soil, time (1, 24 and 48 hours after second autoclaving cycle, in open, grey and black bars, respectively) and their interaction. P values are presented.



995 Figure 4. Relationship between (A) ^{33}P recoveries as measured directly in acidified bicarbonate extracts and after isobutanol fractionation, relative to the added tracer amount, and between (B) P_i concentrations measured by the malachite green method in acidified bicarbonate extracts and after isobutanol fractionation following the phosphomolybdate blue approach. (C) Comparison of specific activities (SA) of P_i measured in acidified bicarbonate extracts and after isobutanol fractionation. Regression in (C) is only for temperate grassland soils (closed circles) as for tropical forest soils (open circles) P_i concentrations were close to the detection limit of the phosphomolybdate method, impairing calculations of SA for isobutanol fractionation. Linear regressions are given (slopes and intercepts $\pm 1\text{SD}$).





Figure 5. Test for linearity of change in ^{33}P recoveries (A, B) and in specific activities of P_i (C, D) over time, for a temperate grassland soil (A, C) and a tropical forest soil (B, D). Data presented are for ^{33}P measured directly in bicarbonate extracts of live soils (closed circles) and sterile soils (open circles), and are shown on y-axes in a logarithmic manner (LN). Regression lines follow exponential decay which in this linear – LN plot appears as straight line; dashed lines represent sterile soils and solid lines live soils. Regressions were calculated for the time interval 2 to 24 hours (tropical soil) and 4 to 48 hours (temperate soil).


1005



1010

1015


Figure 6. Net immobilization rates of $^{33}\text{P}_i$ by abiotic processes (sorption; A, C) measured in sterile soils and biotic processes (microbial uptake; B, D) measured in live soils of a temperate grassland (soil 4) and a tropical forest (soil 3) after 0, 1, 2, 4, 8, 24 and 48 hours (A, B) and for six soils measured after 0, 4 and 24 hours (C, D). Temperate grassland soils (2, 4, 6) and tropical forest soils (3, 5, 7) were investigated in C and D. Curvilinear regressions following the function “exponential rise to maximum” were performed on the data in (A, B). Statistical analyses of data in (C, D) were run by two-way ANOVA for the factors soil and time (0, 4 and 24 hours after tracer addition), and the interaction of both factors.



1020

1025


Figure 7. Gross influx rates into the available soil P_i pool (A) and gross efflux rates from this pool (B) measured by ^{33}P isotope pool dilution for six soils over the time period 4 to 24 hours and assessed in sterile and live soils. Abiotic and biotic process rates are indicated by open and closed bars, respectively. Temperate grassland soils (2, 4, 6) and tropical forest soils (3, 5, 7) were studied. Presented are means $\pm 1SD$ of triplicate live and sterile soils per time point and soil type. One-way ANOVA was performed on transformed data as indicated in brackets. Different lower case letters indicate significant differences between soils for abiotic processes (open bars), upper case letters for biological processes (black bars).



1030

Figure 8. Relationship between selected soil physicochemical parameters, net abiotic and microbial immobilization fluxes, gross P_i influx rates by biological processes (gross P_o mineralization) and abiotic processes (gross P_i desorption), and gross P_i efflux rates by biological processes (gross microbial P_i uptake) and abiotic processes (gross P_i sorption). Regression lines are for linear or power function fits, and P and R^2 values for these are shown. Open circles (\circ) depict tropical forest soils and closed circles (\bullet) temperate grassland soils. Units are provided in Table 2 for soil physicochemical parameters and phosphomonoesterase, are % of added tracer for net processes, and $\mu\text{g P g}^{-1}$ soil dw d^{-1} for gross process rates.

1035



1040