

Multiyear methane ebullition measurements from water and bare peat surfaces of a patterned boreal bog

Elisa Männistö¹, Aino Korrensalo¹, Pavel Alekseychik², Ivan Mammarella², Olli Peltola², Timo Vesala^{2,3}, Eeva-Stiina Tuittila¹

5 ¹Peatland and soil ecology research group, School of Forest Sciences, University of Eastern Finland, PO Box 111, FIN-80101 Joensuu, Finland

²Institute for Atmospheric and Earth System Research/Physics, Faculty of Science, University of Helsinki, P.O. Box 68, 00014 Helsinki, Finland

10 ³Institute for Atmospheric and Earth System Research/Forest Sciences, Faculty of Agriculture and Forestry, University of Helsinki, P.O. Box 27, 00014 Helsinki, Finland

Correspondence to: elisa.mannisto@uef.fi

Abstract

15 We measured methane ebullition from a patterned boreal bog situated in the Siikaneva wetland complex in southern Finland. Measurements were conducted on water (W) and bare peat surfaces (BP) in three growing seasons 2014–2016 using floating gas traps. The volume of the trapped gas was measured weekly, and methane and carbon dioxide (CO₂) concentrations of bubbles were analyzed from fresh bubble samples collected separately. We applied a mixed effects model to quantify the effect of the environmental controlling factors on the ebullition.

20 Ebullition was higher from W than from BP, and more bubbles were released from open water (OW) than from water's edge (EW). On average, ebullition rate was the highest in the wettest year 2016 and ranged between 0–253 mg m⁻²d⁻¹ with median 2 mg m⁻²d⁻¹, 0–147 mg m⁻²d⁻¹ with median 3 mg m⁻²d⁻¹ and 0–186 mg m⁻²d⁻¹ with median 28 mg m⁻²d⁻¹ in 2014, 2015 and 25
2016, respectively. Ebullition increased together with increasing peat temperature, weekly air temperature sum and atmospheric pressure, and decreasing water table (WT). Methane concentration in the bubbles released from W was 15–20 times higher and from BP 10 times higher than their CO₂ concentration. The proportion of ebullition fluxes upscaled to ecosystem level for the peak season was 2–8 % and 2–5 % of the total flux measured with eddy covariance technique and with chambers and gas traps, respectively. Thus, the contribution of methane
30 ebullition from wet non-vegetated surfaces of the bog to the total ecosystem-scale methane emission appeared to be small.

Keywords: CH₄, ebullition, peatland, peat temperature, water table, atmospheric pressure

1 Introduction

35 Historically, bogs were commonly feared as people saw mysterious lights which gave rise to
the tales of the “will o’ the wisps” that lure travelers from their paths to sink into bog holes
(Meredith, 2002). Nowadays, these lights are thought to be spontaneous combustion of
peatland gases, such as methane, bubbling to the atmosphere, rather than deceptive fairies.
However, the widespread folklore indicates that the phenomenon is well known around the
40 world in peatland rich areas. Although currently peatlands are more known for their climate
cooling impact as small carbon sinks and the storage of a third of the global soil carbon stock
(Strack, 2008), they are also a major natural source of methane, a potent climate warming
greenhouse gas (IPCC, 2014). The same high water table (WT) conditions that support
accumulation of organic material as peat by slowing down aerobic decomposition also favor
45 methane production by anaerobic microbes, methanogens (Archaea) (Hanson and Hanson,
1996). It has been predicted that carbon dioxide (CO₂) uptake typically offsets sustained
methane emissions in natural ecosystems in the long term (i.e., several centuries), although
with large spatiotemporal variability (Petrescu et al., 2015).

Methane is emitted from peatlands into the atmosphere via three routes: by diffusion from peat,
50 transport through aerenchymatous vascular plants and by episodic bubble release i.e. ebullition
(LeMer and Roger, 2001; Raghoebarsing et al., 2005). Large part of the produced methane is
oxidized by methanotrophic bacteria in the aerobic peat layer above water level (Hanson and
Hanson, 1996; LeMer and Roger, 2001; Larmola et al., 2010), and thus methane flux rate of a
peatland depends on the rates of methane production and consumption, in addition to
55 transportation within the peat to the atmosphere. It is known that part of methane can be
oxidized also in plants, such as rice (Bosse and Frenzel, 1997), but so far significant methane
oxidation has not been detected in bog plants, such as *Eriophorum* spp. (Frenzel and Rudolph,
1998). As methane emitted through vascular plants or by ebullition bypasses the oxidation in
the aerobic peat layer, these pathways can potentially release high amounts of methane into the
60 atmosphere. Diffusion through peat and vascular plants have been regarded to be the dominant
pathways of methane emissions and those emission pathways have been largely targeted with
chamber measurements (e.g. Bubier et al., 2005; Ström et al., 2005; Turetsky et al., 2014).
Alternatively, eddy covariance (EC) technique is used used to estimate the integrated

ecosystem scale methane flux (e.g. Brown et al., 2014; Rinne et al., 2018), but are unable to
65 differentiate the emission pathways.

Current models of global methane budget are still uncertain due to limited knowledge of the
relative contribution of different factors controlling methane fluxes (Riley et al., 2011). The
largest source of uncertainty is the quantity of methane emissions from natural wetlands, such
as peatlands (Riley et al., 2011; Melton et al., 2013). Within peatland emissions, the largest
70 uncertainty is related to the magnitude of ebullition (Peltola et al., 2018). We are aware of only
few studies that have directly measured ebullition from boreal peatlands with gas traps. In the
first one, Hamilton et al. (1994) carried out measurements over 24 hours and found no bubbles.
In the three other studies conducted in a fen (Starck et al., 2005; Strack and Waddington, 2008)
and a bog (Stamp et al., 2013) ebullition fluxes between 7 – 96 mg CH₄ m⁻² d⁻¹ were detected
75 but the importance of ebullition for the ecosystem flux remained unrevealed. Ebullition has
also been measured in the field by separating peak methane releases from steady chamber flux
(Riutta et al., 2007; Tokida et al., 2007; Goodrich et al., 2011) with emissions varying from
49–152 mg CH₄ m⁻² d⁻¹ (Goodrich et al., 2011) to 48–1440 mg CH₄ m⁻² d⁻¹ (Tokida et al.,
2007). These studies show contrasting results on the contribution of ebullition to the total
80 emission. While Riutta et al. (2007) estimated the role of ebullition to be small in the two study
years, Tokida et al. (2007) (with two sample plots) found that the proportion of ebullition may
constitute up to 50% of the total flux. Results on mesocosm studies in laboratory conditions
are similarly disparate as they show that the proportion of ebullition in the total emission varies
from 3% (Green and Baird, 2013) to 50% (Christensen et al., 2013).

85 Similarly to chamber and EC measurements (Rinne et al., 2007; Jackowicz-Krczyński et al.,
2010; Turetsky et al., 2014; Mikhaylov et al., 2015; Rinne et al., 2018), direct ebullition studies
have connected the rate of methane emission to peat temperature (Strack et al., 2005) relating
to increasing microbial activity (Conrad et al., 1997). Noteworthy, incoming energy flux has
been shown to primarily control the methane production and ebullition in shallow subarctic
90 lakes (Wik et al., 2014) that could be contrasted to peatland pools. Ebullition in peatlands has
additionally been linked to decreasing WT and falling atmospheric pressure: the decrease of
hydrostatic pressure increases the volume of the gas phase of methane in peat and releases it
into the atmosphere (Tokida et al., 2007). Also, an increase in atmospheric pressure can trigger
ebullition by decreasing the bubble size due to compression and thus increasing the bubble
95 mobility in shallow peat (Comas et al., 2011; Chen and Slater, 2015). Furthermore, peat
structure has been shown to affect bubble sizes and determine whether ebullition is steady or

erratic (Ramirez et al., 2016). However, the importance of these factors for ebullition is still based on the few studies, of which the longest covers two growing seasons (Strack and Waddington, 2008).

100 In this study, we measured methane ebullition from open water pools (W) and bare peat surfaces (BP) with gas traps in three consecutive growing seasons 2014 – 2016 in a boreal bog where methane fluxes were measured also with EC and static chambers techniques. We aimed to: (1.) quantify the spatial and temporal variation of methane ebullition from wet bog surfaces; (2.) study the controlling factors; and (3.) assess the contribution of ebullition from wet surfaces
105 to the ecosystem level emission.

2 Materials and methods

The study was conducted in the ombrotrophic bog that is part of Siikaneva peatland complex situated in southern Finland (61°50'N, 24°12'E), 160 m a.s.l., within the southern boreal
110 vegetation zone (Ahti et al., 1968). Annual rainfall of the area is 707 mm, the snow depth in March (with the thickest snow cover) is 36 cm, the annual cumulative temperature is 1318, the length of growing season is 168 days, the average annual temperature is 4.2 °C and the average temperatures in January and July are -7.2 °C and 17.1 °C, respectively (30-year averages from the nearby Juupajoki-Hyytiälä weather station, except snow depth as 20-year average). The
115 microtopography of the studied bog site varies from W and BP to hollows, lawns and hummocks. W and BP cover together approximately one fourth of the site (W 11.6% and BP 15.3% within 30 m radius from the EC tower of the site). The bottom layer is formed by *Sphagnum* mosses, except in W and BP that are devoid of moss. Sedges are the dominating vascular plants in hollows and lawns, whereas vascular plant vegetation on hummocks is
120 dominated by dwarf shrubs. In BP, *Rhynchospora alba* is often the only plant species (Korrensalo et al., 2018a).

In order to measure methane ebullition from the studied bog, floating gas traps were placed in W and BP in three consecutive years 2014–2016. Only W and BP microforms were chosen because we expected high ebullition from these waterlogged surfaces that have almost no
125 vegetation, and because the sampling method required gas traps to be easily filled with water. BP are patches of visible peat that have WT at or near surface. For example in 2014, WT in BP was on average -1.8 cm. W are without a clear bottom but have on average one meter of water over very loose peat slurry and their water area starts directly from the edge of the surrounding

moss cover. As it is difficult to determine what is the bottom of the pools, we did not measure
130 the water depth or temperature in the bottom of the W.

The gas traps were constructed from inverted plastic funnels with diameters ranging between
14.3 cm and 24.5 cm (Fig. 1). A syringe with a three-way stopcock was attached to the narrow
end of each funnel and the joint was covered with sealant to make it airtight. Piece of metallic
netting coated with filter fabric was glued inside the funnels to prevent litter and small animals
135 from entering the gas traps in the open water pools. The gas traps on W were attached to a
floating styrofoam raft and placed in the pools in lines of two or three gas traps, anchored to
the opposing shores of the pool with strings (Fig. 1). To study the potential difference in
availability of substrate for methanogenesis some gas traps were anchored further away from
the surrounding moss cover at the center of the pools (open water, OW), while the other gas
140 traps were anchored at the water's edge (EW) right next to the moss (Fig. 2). The gas traps on
BP were placed next to boardwalks in the study site (Fig. 2). The air was sucked out of the gas
traps with an extra syringe until they were filled with water. The rate of ebullition was measured
weekly by sampling the gas volume that had replaced water in each gas trap.

16 gas traps were used (11 in W and 5 in BP) from 3 June to 25 September in 2014, 20 gas
145 traps (13 in W and 7 in BP) from 13 May to 24 September in 2015, and 18 gas traps (12 in W
and 6 in BP) from 27 May to 9 September in 2016.

Methane concentrations of the gas caught in the gas traps during the weekly sampling periods
were measured in 2014 and compared with methane concentration of fresh ebullition samples.
We found methane concentrations in the gas traps to be clearly lower than in the fresh ebullition
150 samples (Table A1), and thus methane concentration of the gas caught in the traps was assumed
to dilute during the weekly sampling periods due to diffusion. Therefore, methane
concentration of the releasing bubbles was not measured from the weekly samples, but instead
by collecting fresh ebullition samples from W without disturbing the gas traps, and from BP
that had no gas traps. Ebullition was triggered manually from the sampled surfaces and the
155 formed bubbles were caught in an extra gas trap, from where 20 ml samples were taken into
vacuumed glass vials. The samples were analyzed with an Agilent Technologies HP 8690 gas
chromatograph in Natural Resources Institute Finland (LUKE), Vantaa. Fresh ebullition
samples were collected four times during the measurement season in 2014 and 2016, and 13
times in 2015. Average methane concentration was interpolated linearly from the fresh
160 ebullition samples for each weekly measurement day.

Average methane emission by ebullition as $\text{ml m}^{-2} \text{d}^{-1}$ was calculated based on the area of the gas trap, number of days and volume of gas collected in each measurement period and the average methane concentration of each measurement period. In order to convert the emissions to $\text{mg m}^{-2} \text{d}^{-1}$, methane density in each measurement period was calculated based on the average
165 air temperature of the measurement period in $^{\circ}\text{C}$ and the standard atmospheric air pressure, 1.01325 bar. Average methane emission ($\text{mg m}^{-2} \text{d}^{-1}$) was calculated separately for ebullition from OW, EW and BP.

In order to compare the ebullition fluxes to EC and chamber measurements (Korrensalo et al., 2018b), the ebullition flux was upscaled to ecosystem level by interpolating linearly the total
170 average ebullition that was calculated as a sum of average ebullition fluxes from W and BP weighted with their relative surface areas.

Air pressure and temperature data from 2014–2016 were received from the Juupajoki-Hyytiälä weather station that is situated about 6 km from the study site in Siikaneva. The data on WT, water temperature and peat temperatures at the depths of 5, 20 and 50 cm were received from
175 data loggers installed in a lawn about 1.5 m away from the EC raft. Photosynthetically active radiation (PAR) data was measured in the site.

Linear mixed-effects models were used to analyze the effect of measured environmental variables (peat temperature in different depths, WT, atmospheric pressure and cumulative PAR and effective temperature sum of a measurement period as variables of incoming energy flux)
180 on log-transformed ebullition flux rates. The gas trap was included as a random effect in the model. We also tested which of the four peat temperature variables explained the variation in ebullition fluxes the best. The data were analyzed with the function lme of the package nlme of the R software (version 3.3.2).

185 **3 Results**

Among the three studied years, the year 2014 was the warmest, driest and with the highest amount of cumulative photosynthetically active radiation (PAR) (Finnish Meteorological Institute open data) (Table 1). It was also warmer than 30y average. Year 2015 was the coolest, with a lowered annual rainfall and PAR, while 2016 was the wettest and the cloudiest year
190 (Table 1). All three years were significantly drier than the average (Table 1).

Measured methane ebullition ranges were 0–253 mg m⁻²d⁻¹, 0–147 mg m⁻²d⁻¹ and 0–186 mg m⁻²d⁻¹ with medians 2, 3 and 28 mg m⁻²d⁻¹ in 2014, 2015 and 2016, respectively (A2). Weekly medians of individual gas traps were 0–57 mg m⁻²d⁻¹ in 2014, 0–33 mg m⁻²d⁻¹ in 2015 and 10–67 mg m⁻²d⁻¹ in 2016. The three years differed (degrees of freedom (DF) = 2, 746; *p* <.0001) as slightly higher ebullition fluxes were generally obtained in 2015 than 2014, while, on average, the ebullition fluxes were on their highest in the wettest year 2016.

Higher ebullition was observed on W than on BP (Fig. 3) (DF = 1, 746; *p* <.0001). Ebullition from OW was significantly higher than ebullition from EW, except in the middle of the growing season 2015 (Fig. 3). Although BP showed lower ebullition with fewer peaks than W, all the surfaces had the same seasonal ebullition pattern each year with highest fluxes observed in August (Fig. 3). However, in 2015 the highest ebullition was measured later than in other years after relatively low ebullition in late summer (Fig. 3).

Ebullition increased with increasing average peat temperature at the depth of 5 cm (DF = 1, 746; *p* <.0001) that explained ebullition better than the other peat temperature variables measured. Seasonal pattern of ebullition followed temperature in each year (Fig. 4). Higher ebullition rates were also explained with decreasing average WT (DF = 1, 746; *p* = 0.0001). The highest ebullition peaks were associated with the lowest WT in each year (Fig. 4). A prolonged depression of WT further explained the late peak of ebullition in 2015, as well as the increase in ebullition in the autumn 2016 (Fig. 4). Change in atmospheric pressure during the measurement period further explained ebullition: more bubbles were released with higher increase in pressure (DF = 1, 746; *p* = 0.001). Some events of ebullition might be directly related to decreasing atmospheric pressure such as the small peak in ebullition in mid-August 2014 that appears to be better explained by long decrease in atmospheric pressure than by peat temperature or WT (Fig.5). After including peat temperature, WT and change in atmospheric pressure, the effective temperature sum of a measurement period still had a positive effect on ebullition (DF = 1, 746; *p* = 0.0351). Finally, the cumulative PAR had no significant effect on ebullition and was excluded from the final model.

Fresh ebullition samples analyses showed that the released gas bubbles contained more methane than CO₂. Methane concentration of bubbles released from W was 15–20 times higher than their CO₂ concentration, while bubbles from BP had tenfold higher methane than CO₂ concentration (Table 2).

The average ebullition flux upscaled to ecosystem level was an order of a magnitude lower than the net methane flux measured by EC in each year (Fig. 6). The sum of ebullition and upscaled chamber flux in 2014 was higher than the one measured with EC, but the two estimates followed the same seasonal trend (Fig. 6). The contribution of ebullition to the total methane flux measured with chambers and bubble traps during the peak season in 2014 was 2, 3 and 5 % in June, July and August, respectively (Table 3). The contribution of ebullition to EC flux during the peak season varied from 2 % in June 2014 to 8 % in August 2015 (Table 3).

230

4 Discussion

4.1 The magnitude of ebullition

The methane ebullition measured in this study ranged from 0 to 253 mg m⁻² d⁻¹ and the seasonal weekly median of ebullition for different surfaces ranged from 0 (measured from BP in 2014) to 37 mg m⁻² d⁻¹ (measured from OW in 2016). Our results are of the same magnitude as ebullition fluxes previously measured in boreal peatlands with gas traps, ranging from 7 to 96 mg m⁻² d⁻¹ (Strack et al., 2005; Strack and Waddington, 2008; Stamp et al., 2013), and with automatic chambers, ranging from 9 to 152 mg m⁻² d⁻¹ (Goodrich et al., 2011). In addition to field measurements some of the laboratory-based experiments have shown similar ebullition flux rates in the range of 0-270 mg m⁻² d⁻¹ (Christensen et al., 2004; Kellner et al., 2006; Yu et al., 2014) but also higher fluxes up to 784 mg m⁻² d⁻¹ (Green and Baird, 2012). Some laboratory studies have shown potential for even much higher ebullition rates up to 33 000 mg m⁻² d⁻¹ (*Sphagnum* surface samples from bog in Tokida et al., 2005; fen lawn samples in Waddington et al., 2009). So far, only Tokida et al. (2007) have estimated ebullition fluxes reaching 1440 mg m⁻² d⁻¹ in the field based on methane fluxes measured with the static chamber method from two sample plots showing high episodic fluxes during 30 min measurements. Generally, there is a difference in temporal resolution between the two methods as chamber measurements usually cover only short time periods (from minutes to hours) while gas traps show estimates of cumulative bubble flux over several days.

250 The fact that the ebullition rates measured with gas traps are lower than in laboratory studies might be partly explained by the process of bubbles stacking in the gas traps instead of automatically gathering in the headspace. In this study, we tried to overcome this potential error source by gently shaking and tapping the gas traps before sampling, simultaneously trying

to avoid causing more ebullition by this disturbance. However, methane ebullition fluxes of up
255 to 1683 mg m⁻² d⁻¹ have been previously measured with the same method from subarctic lakes
(Wik et al., 2013), which shows the potential of this method to measure also higher ebullition
fluxes.

4.2 Temporal and spatial variation

260 Our study conducted over three growing seasons showed interannual variation. The highest
ebullition on average was measured in 2016, whereas the average flux rates of 2014 and 2015
did not differ significantly from each other. More ebullition was measured especially from BP
in 2016 that was the wettest year with the highest WT. This indicates that despite higher WT
increases hydrostatic pressure in peat, wetter conditions in BP facilitate gas release as bubbles.
265 Although 2015 was almost as wet year as 2016, it was much cooler, which decreases methane
production. The warmest year 2014 again was much drier than 2016, and although there was
high ebullition with sharp drop in WT during the peak season, the general ebullition level from
BP was low. The only other peatland study with gas traps covering more than one growing
season (Strack and Waddington, 2008) also found the ebullition level to differ between the
270 study years. Similarly, Wik et al. (2013) found differences in bubble methane concentrations
and fluxes in subarctic lakes among the four summers studied. These results point out the need
for multi-year studies in order to include inter annual variation of ebullition fluxes in methane
models. Furthermore, the higher ebullition rate from W than from BP in our study indicates
that balanced sampling in a bog should cover microform variability, although in some studies
275 no spatial variation in ebullition were found (Green and Baird, 2012 and 2013; Stamp et al.,
2013). However, drier and wetter conditions can change the proportions of water and bare peat
surfaces, and according to our results, such changes may have an impact on ebullition.

4.3 Controlling factors and their importance

280 The measured ebullition rates increased together with peat temperature as shown also earlier
(Strack and Waddington, 2008). Increasing temperature generally increases the activity of
methanogens, and thus more methane is produced in the peat when it gets warmer until the
temperature optimum of the microbes around 20-30 °C is reached (Dunfield et al., 1993). Peat
temperature affects also the solubility of methane according to Henry's law as gas solubility

285 decreases with increasing temperature (Strack, 2005). Thus, increasing peat temperature may lead to transfer of methane from aqueous to gaseous phase, which increases bubble formation (Strack, 2005). In our study, the peat temperature at the depth of 5 cm showed the highest correlation with ebullition but temperature in all depths were highly intercorrelated. The effect of peat temperature was reflected in the seasonal pattern of ebullition.

290 As expected, ebullition fluxes increased when WT decreased as found in previous studies (Strack and Waddington, 2008). Bubbles may accumulate in peat under barriers, such as pieces of wood, and they are suppressed by high hydrostatic or air pressure (Rosenberry et al., 2003; Strack and Waddington, 2008; Chen and Slater, 2015). Decreasing WT lowers the hydrostatic pressure releasing newly formed and the accumulated bubbles. Many studies have also shown
295 that the falling atmospheric pressure can trigger high rates of ebullition (Tokida et al., 2005; Tokida et al., 2007). Although some weeks showed higher ebullition rates when atmospheric pressure was falling, this pattern was not consistent as increasing ebullition rates were measured also during periods of rising atmospheric pressure. After including WT as explanatory variable, we still found the weekly change in atmospheric pressure to significantly
300 affect ebullition as bigger increase in weekly pressure was related to more ebullition. Previously, Comas et al. (2011) used ground penetrating radar (GPR) to study the vertical distribution of free-phase gas in a northern peatland and found that increasing atmospheric pressure caused rapid ebullition by releasing gas from shallow peat, whereas decreasing pressure released gas from deeper peat to shallow layers. Also Chen and Slater (2015) showed
305 that increasing pressure can trigger ebullition as it increases the bubble mobility in peat.

Furthermore, higher ebullition rates were measured with higher effective temperature sum of the measurement period. This indicates the importance of energy input as a driver of methane production and release as shown by Wik et al. (2014). They found strong positive correlations between seasonal bubble methane flux from subarctic lakes and four proxies of energy flux,
310 such as average short wave radiation and maximum water sediment temperature (Wik et al., 2014). We tried also to compare cumulative PAR (i.e. short wave radiation) to seasonal cumulative ebullition fluxes but could not find clear correlation between the two in the three study years. However, the positive effect of the measurement period temperature sum on ebullition shows that increasing energy input can increase ebullitive methane flux rates in the
315 studied bog site.

4.4 Importance for the ecosystem level flux

When measured ebullition fluxes were upscaled to the ecosystem level, they showed much lower methane emissions than measured with chamber and EC techniques. In our previous study (Korrensalo et al., 2018b), we measured diffusive methane fluxes with the static chamber technique from six different plant community types, including BP, at the same bog site in 2014. We found higher methane fluxes from BP than from high hummocks (HHU), but otherwise all the studied plant community types had similar methane fluxes. When chamber fluxes were upscaled to ecosystem level they were similar to the EC flux (Korrensalo et al., 2018b). Although laboratory incubation studies have shown that the contribution of ebullition to the total methane flux may reach up to 50 % (Christensen et al., 2013; Tokida et al., 2007), the ebullition contribution in this study was only 3-5 % during the peak season of 2014. Here, ebullition is considered only from waterlogged surfaces as we did not measure ebullition from vegetated surfaces. Previously, Riutta et al. (2007) measured methane fluxes from different plant communities with static chambers in Siikaneva fen site situated 1.3 km south-east from our studied bog site and calculated results for both diffusive and ebullition fluxes. They found ebullition from all communities but showed that its contribution to the total flux (diffusive flux + ebullition) was negligible or very small (Riutta et al., 2007). We estimated ebullition to occur only twice in 210 measurements on moss cover surfaces, i.e. in 0.8 %, of our 2014 chamber data. Therefore, we assume that ebullition from vegetated surfaces would not greatly contribute to the total flux in the bog site either. Earlier, similarly to our study, Green and Baird (2013) have found ebullition to contribute less than 3.3 % of total methane fluxes when incubating peat samples collected from hollows and lawns from two raised bogs in laboratory study. As the measured bubble methane fluxes in our study were of the same magnitude in each year, ebullition did not contribute significantly to the ecosystem methane emissions in any studied growing season as seen in the comparison with the EC flux. While the same seasonal trend and peaks can be seen in both fluxes in each year, the total flux measured with EC is constantly at least an order of magnitude higher than the ebullition flux rate.

345 **Conclusions**

More methane ebullition was found from W than from BP, and within the pools more bubbles were released from OW than from EW. We found also variation between the three studied growing seasons as ebullition rate was generally higher in the wettest year 2016. Due to this

spatial and temporal variation, differences between years in wet/dry conditons may have an
350 effect on ebullition. As expected, ebullition increased together with increasing peat
temperature, that facilitates methane production, and with decreasing WT, that reduces
hydrostatic pressure on peat. Additionally, more bubbles were released with bigger weekly
increase in atmospheric pressure, which has been related to rapid ebullition from shallow peat.
Futhermore, higher weekly temperature sum had a positive effect on ebullition, which shows
355 that increasing energy input can increase ebullitive methane flux rates in the studied bog site.
Therefore, the growing season lengthening and increase in the average temperatures due to
climate change may increase the methane emissions in the peatland ecosystem as long as
waterlogged anoxic conditions in the peat for methane production persist. Ebulliton flux
upscaled to the ecosystem level showed similar seasonal pattern as methane fluxes measured
360 with EC and chamber techniques but was an order of magnitude lower and had a very small
contribution to the total ecosystem flux. Our study includes ebullition only from the
waterlogged surfaces as we did not expect ebullition from all the plant community types to be
substantial based on the previous study in the nearby fen site and our chamber measurements
in 2014. However, estimating the amount of ebullition from all the plant community types
365 would be needed to fully understand the spatial variation of ebullition in the future. In addition,
measurements with e.g. time-lapse cameras are needed to study the short term temporal
variation of ebullition and to estimate the frequency and magnitude of rapid ebullition events
that may contribute to the total ecosystem flux.

Data availability

370 Data is available upon request from the corresponding author.

Author contribution

AK, EST, PA and TV came up with the idea and design. AK and EM conducted the ebullition
measurements and processed the data. Eddy covariance data was collected and analyzed by PA
and IM. EM fitted the mixed-effects models. The manuscript was written by EM, AK and EST
375 and commented by all the other authors.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

This work is supported by faculty of Science and Forestry, University of Eastern Finland, the
380 Finnish Cultural Foundation, Academy of Finland (project code 287039), Academy research
project CLIMOSS (41007-00086900), Strategic Research Council research project SOMPA
(41007-00114600), the National Centre of Excellence (272041), ICOS-Finland (281255),
Academy professor project (284701) funded by the Academy of Finland and AtMath funded
by the University of Helsinki. OP was supported by the Postdoctoral Researcher project
385 (315424) funded by the Academy of Finland. We thank Hyytiälä Forest Research Station and
its staff for research facilities and Pauli Karppinen from Natural Resources Institute Finland
(LUKE) for analyzing methane concentration of the fresh bubble samples. We also thank Salli
Uljas, Janne Sormunen, Franziska Rossocha, and Laura Kettunen for the help in the field.

390 **References**

- Ahti, T., Hämet-Ahti, L., & Jalas, J.: Vegetation zones and their sections in
northwestern Europe, *Ann. Bot. Fenn.*, 5, 169–211, 1968.
- Bosse, U. and Frenzel, P.: Activity and Distribution of Methane-Oxidizing Bacteria in
Flooded Rice Soil Microcosms and in Rice Plants (*Oryza sativa*), *Appl. Environ. Microb.*, 63,
395 1199–1207, 1997.
- Brown, M. G., Humphreys, E. R., Moore, T. R., Roulet, N. T., & Lafleur, P. M.:
Evidence for a nonmonotonic relationship between ecosystem-scale peatland methane
emissions and water table depth, *J. Geophys. Res. Biogeosci.*, 119, 826–835, 2014,
doi:10.1002/2013JG002576
- 400 Bubier, J. Moore, T., Savage, K., & Crill, P.: A comparison of methane flux in a
boreal landscape between a dry and a wet year *Global Biogeochem. Cy.*, 19, 2005 GB1023
doi:10.1029/2004GB002351.
- Chen, X., & Slater, L.: Gas bubble transport and emissions for shallow peat from a
northern peatland: The role of pressure changes and peat structure, *Water Resour. Res.*, 51,
405 151–168, 2015, doi:10.1002/2014WR016268.
- Christensen, T. R., Panikov, N., Mastepanov, M., Joabsson, A., Stewart, A., Öquist, M.,
Sommerkorn, M., Reynald, S., & Svensson, B.: Biotic controls on CO₂ and CH₄ exchange
in wetlands –a closed environment study, *Biogeochemistry*, 64, 337–254, 2003.

- 410 Comas, X., Slater, L., & Reeve, A. S.: Atmospheric pressure drives changes in the vertical distribution of biogenic free-phase gas in a northern peatland, *J. Geophys. Res.*, 116, G04014, 2011, doi:10.1029/2011JG001701.
- Conrad, R., Schultz, H., & Babel, M.: Temperature imitation of hydrogen turnover and methanogenesis in anoxic paddy soil, *FEMS Microbiol. Ecol.*, 45, 281-289, 1997.
- 415 Dunfield, P., Knowles, R., Dumont, R., & Moore, T. R.: Methane production and consumption in temperate and subarctic peat soils: Response to temperature and pH, *Soil Biol. Biochem.*, 25, 321–326, 1993.
- Finnish Meteorological Institute open data: <https://en.ilmatieteenlaitos.fi/open-data>, last access: 08 May 2019.
- 420 Frenzel, P. and Rudolph, J.: Methane emission from a wetland plant: the role of CH₄ oxidation in *Eriophorum*, *Plant Soil*, 202, 27–32, 1998.
- Goodrich, J. P., Varner, R. K., Frohking, S., Duncan, S. B., & Crill, P. M.: High-frequency measurements of methane ebullition over a growing season at a temperate peatland site, *Geophys. Res. Lett.*, 38, L07404, 2011, doi:10.1029/2011GL046915.
- 425 Green, S. M., & Baird, A. J.: A Mesocosm study of the role of the sedge *Eriophorum angustifolium* in the efflux of methane –including that due to episodic ebullition –from peatlands, *Plant Soil*, 351, 207–218, 2012.
- Green, S. M., & Baird, A. J.: The importance of episodic ebullition methane losses from three peatland microhabitats: a controlled-environment study, *Eur. J. Soil. Sci.*, 64, 27–36, 2013.
- 430 Hamilton, J. D., Kelly, C. A., Rudd, J. W. M., Hesslein, R. H., & Roulet, N. T.: Flux to the atmosphere of CH₄ and CO₂ from wetland ponds on the Hudson Bay lowlands (HBLs), *J. Geophys. Res.*, 99, 1495–1519, 1994.
- Hanson, R. S., and Hanson, T. E.: Methanotrophic Bacteria, *Microbiol. Rev.*, 60, 439–471, 1996.
- 435 IPCC, Climate Change 2014: Synthesis Report, Contribution of Working Groups I, and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151

pp., 2014.

Jackowicz-Korczyński, M., Christensen, T. R., Bäckstrand, K., Crill, P., Friborg, T.,
440 Mastepanov, M., & Ström, L.: Annual cycle of methane emission from a subarctic
peatland, *J. Geophys. Res.*, 115, 2010, G02009 doi:10.1029/2008JG000913.

Korrensalo, A., Kettunen, L., Laiho, R., Alekseychik, P., Vesala, T., Mammarella, I., &
Tuittila, E.-S.: Boreal bog plant communities along a water-table gradient differ by their
standing biomass but not by their biomass production, *J. Veg. Sci.*, 29, 136–146, 2018a,
445 DOI: 10.1111/jvs.12602.

Korrensalo, A., Männistö, E., Alekseychik, P., Mammarella, I., Rinne, J., Vesala, T., &
Tuittila, E.-S.: Small spatial variability in methane emission measured from a wet patterned
boreal bog, *Biogeosciences*, 15, 1749–1761, 2018b.

Kellner, E., Baird, A. J., Oosterwoud, M., Harrison, K., & Waddington, J. M.: Effect of
450 temperature and atmospheric pressure on methane (CH₄) ebullition from near-surface peats,
Geophys. Res. Lett., 33, 2006, doi:10.1029/2006GL027509.

Larmola, T., Tuittila, E.-S., Tirola, M., Nykänen, H., Martikainen, P. J., Yrjälä, K., Tuomivirta,
T., & Fritze, H.: The role of Sphagnum mosses in the methane cycling of a boreal mire, *Ecology*,
91, 2356–2365, 2010.

455 LeMer, J., & Roger, P.: Production, oxidation, emission and consumption of methane by soils:
A review, *Eur. J. Soil Biol.*, 37, 25–50, 2001.

Melton, J. R., Wania, R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis,
C. A., Beerling, D. J., Chen, G., Eliseev, A., Denisov, S. N., Hopcroft, P. O., Lettenmaier, D.
P., Riley, W. J., Singarayer, J. S., Subin, Z. M., Tian, H., Zürcher, S., Brovkin, V., van
460 Bodegom, P. M., Kleinen, T., Yu, Z. C., & Kaplan, J. O.: Present state of global wetland extent
and wetland methane modelling: conclusions from a model inter-comparison project
(WETCHIMP), *Biogeosciences*, 10, 753–788, 2013.

Meredith, M.: Hazards in the bog -real and imagined, *Geogr. Rev.*, 92, 319–332, 2002.

Mikhaylov, O. A., Miglovets, M. N., & Zagirova, S. V.: Vertical Methane Fluxes in
465 Mesooligotrophic Boreal Peatland in European Northeast Russia, *Contemp. Probl. Ecol.*, 8,
368–375, 2015.

- Peltola, O., Raivonen, M., Li, X., & Vesala, T.: Technical note: Comparison of methane ebullition modelling approaches used in terrestrial wetland models, *Biogeosciences*, 15, 937–951, 2018.
- 470 Petrescu, A. M. R., Lohila, A., Tuovinen, J.-P., Baldocchi, D. D., Desai, A. R., Roulet, N. T., Vesala, T., Dolman, A. J., Oechel, W. C., Marcolla, B., Friborg, T., Rinne, J., Matthes, J. H., Merbold, L., Meijide, A., Kiely, G., Sottocornola, M., Sachs, T., Zona, D., Varlagin, A., Lai, D. Y. F., Veenendaal, E., Parmentier, F.-J. W., Skiba, U., Lund, M., Hensen, A., van Huissteden, J., Flanagan, L. B., Shurpali, N. J., Grünwald, T., Humphreys, E. R., Jackowicz-
475 Korczyński, M., Aurela, M. A., Laurila, T., Grüning, C., Corradi, C. A. R., Schrier-Uijl, A. P., Christensen, T. R., Tamstorf, M. P., Mastepanov, M., Martikainen, P. J., Verma, S. B., Bernhofer, C., & Cescatti, A.: The uncertain climate footprint of wetlands under human pressure, *PNAS*, 112, 4594–4599, 2015.
- Raghoebarsing, A. A., Smolders, A. J., Schmid, M. C., Rijpstra, W. I. C., Wolters-Arts,
480 M., Derksen, J., Jetten, M. S., Schouten, S., Sinninghe Damsté, J. S., Lamers, L. P., Roelofs, J. G., Op den Camp, H. J., & Strous, M.: Methanotrophic symbionts provide carbon for photosynthesis in peat bogs, *Nature*, 436, 1153–1156, 2005.
- Ramirez, J. A., Baird, A. J., & Coulthard, T. J.: The effect of pore structure on ebullition from peat, *J. Geophys. Res. Biogeo.*, 121, 1646–1656, 2016, doi:10.1002/2015JG003289.
- 485 Riley, W. J., Subin, Z. M., Lawrence, D. M., Swenson, S. C., Torn, M. S., Meng, L., Mahowald, N. M., & Hess, P.: Barriers to predicting changes in global terrestrial methane fluxes: analyses using CLM4Me, a methane biogeochemistry model integrated in CESM, *Biogeosciences*, 8, 1925–1953, 2011.
- Rinne, J., Riutta, T., Pihlatie, M., Aurela, M., Haapanala, S., Tuovinen, J.-P., Tuittila, E.-S., &
490 Vesala, T.: Annual cycle of methane emission from a boreal fen measured by the eddy covariance technique, *Tellus Ser. B*, 59, 449–457, 2007.
- Rinne, J., Tuittila, E.-S., Peltola, O., Li, X., Raivonen, M., Alekseychik, P., Haapanala, S., Pihlatie, M., Aurela, M., Mammarella, I., & Vesala, T.: Temporal variation of ecosystem scale methane emission from a boreal fen in relation to temperature, water table position, and carbon
495 dioxide fluxes, *Global Biogeochem. Cy.*, 32, 1087–1106, 2018.

- Riutta, T., Laine, J., Aurela, M., Rinne, J., Vesala, T., Laurila, T., Haapanala, S., Pihlatie, M., & Tuittila, E.-S.: Spatial variation in plant community functions regulates carbon gas dynamics in a boreal fen ecosystem, *Tellus B*, 59, 838–852, 2007.
- 500 Rosenberry, R. O., Glaser, O. H., Siegel, D. I., & Weeks, E. P.: use of hydraulic head to estimate volumetric gas content and ebullition flux in northern peatlands, *Water Resour Res.*, 39, 2003, doi:10.1029/2002WR001377.
- Stamp, I., Baird, A. J., & Heppell, C. M.: The importance of ebullition as a mechanism of methane (CH₄) loss to the atmosphere in a northern peatland, *Geophys. Res. Lett.*, 40, 2087–2090, 2013.
- 505 Strack, M., Kellner, E., & Waddington, J. M.: Dynamics of biogenic gas bubbled in peat and their effects on peatland biogeochemistry, *Global Biogeochem. Cy.*, 19, 2005, doi:10.1029/2004GB002330.
- Strack, M.: Peatlands and climate change. Jyväskylä, Finland: International Peat Society, 2008.
- Strack, M., & Waddington, J. M.: Spatiotemporal variability in peatland subsurface methane dynamics, *J. Geophys. Res.*, 113, G02010, 2008, doi:10.1029/2007JG000472.
- 510 Ström, L., Ekberg, A., Mastepanov, M., & Christensen, T. R.: The effect of vascular plants on carbon turnover and methane emissions from tundra wetland, *Glob. Change Biol.*, 9, 1185–1192, 2003.
- Tokida, T., Miyazaki, T., & Mizoguchi, M.: Ebullition of methane from peat with falling atmospheric pressure, *Geophys. Res. Lett.*, 32, 2005, doi:10.1029/2005GL022949.
- 515 Tokida, T., Miyazaki, T., Mizoguchi, M., Nagata, O., Takakai, F., Kagemoto, A., & Hatano, R.: Falling atmospheric pressure as a trigger for methane ebullition from peatland, *Global Biogeochem. Cy.*, 21, 2007, doi:10.1029/2006GB002790.
- Turetsky, M. R., Kotowska, A., Bubier, J., Dise, N. B., Crill, P., Hornibrook, E. R. C., 520 Minkinen, K., Moore, T. R., Myers-Smith, I. H., Nykänen, H., Olefeldt, D., Rinne, J., Saarnio, S., Shurpali, N., Tuittila, E.-S., Waddington, J. M., White, J. R., Wickland, K. P., & Wilmsking, M.: A synthesis of methane emissions from 71 northern, temperate, and subtropical wetlands, *Global Change Biol.*, 2014, doi: 10.1111, gcb.12580.
- Waddington, J. M., Harrison, K., Kellner, E., & Baird, A. J.: Effect of atmospheric pressure and temperature on entrapped gas content in peat, *Hydrol. Process.*, 23, 2970–2980, 2009.
- 525

Wik, M., Crill, P. M., Varner, R. K., & Bastviken, D.: Multiyear measurements of ebullitive methane flux from three subarctic lakes, *J. Geophys. Res-Bioge.*, 118, 1307–1321, 2013.

Wik, M., Thorton. B. F., Bastviken, D., MacIntyre, S., Varner, R. K., & Crill, P. M.: Energy input is primary controller of methane bubbling in subarctic lakes, *Geophys. Res. Lett.*, 2014, doi: 10.1002/2013GL058510.

Yu, Z., Slater, L. D., Schäfer, K. V. R., Reeve, A. S., & Varner, R. K.: Dynamics of methane ebullition from a peat monolith revealed from a dynamic flux chamber system, *J. Geophys. Res. Biogeosci.*, 119, 178F9–1806, 2014, doi:10.1002/2014JG002654.

535

Tables

Table 1. Effective temperature sum of the growing season, annual rainfall and the cumulative photosynthetically active radiation (PAR) in the three studied years 2014–2016 and compared to the 30 years averages of the area. Data for the Hyytiälä weather station from Finnish Meteorological Institute open data.

Year	Temp. sum °C	Annual rainfall mm	PAR $\mu\text{mol m}^{-2}$
2014	1 349	579	70 800
2015	1 166	658	69 180
2016	1 280	660	67 996
30 year mean	1 318	707	-

545

550

555

Table 2. Average methane (CH₄) and carbon dioxide (CO₂) concentrations (ml/l) with standard deviation (SD) of releasing gas bubbles from pools (W) and bare peat surfaces (BP) in the three studied years 2014–2016.

	2014		2015		2016	
	Average	SD	Average	SD	Average	SD
W CH ₄	380.0	50.7	285.0	93.0	423.0	103.9
W CO ₂	23.8	4.9	18.0	8.8	20.8	6.5
BP CH ₄	274.2	64.5	273.6	56.1	364.2	123.7
BP CO ₂	29.4	16.0	26.9	9.4	31.8	8.9

560

565

570

575

Table 3. Monthly cumulative methane fluxes ($\text{mg m}^{-2} \text{mo}^{-1}$) measured as ebullition and with eddy covariance (EC) technique for June–August in the three studied years 2014–2016.

	2014			2015			2016		
	Ebullition	EC	% of ebullition	Ebullition	EC	% of ebullition	Ebullition	EC	% of ebullition
June	27	1668	2	73	1139	6	117	2530	5
July	155	3423	5	112	2277	5	314	4216	7
August	176	3447	5	223	2657	8	249	3448	7


580

585

590

595

Figures




600

Figure 1. Floating gas traps in open water peatland pool (OW).



Figure 2. Aerial photo of the study site in Siikaneva bog. Red lines with dots mark the floating gas traps in open water (OW) and water's edge (EW). Red circles mark the area within what the gas traps were placed on bare peat surfaces (BP) that are seen as brownish-grey in the photo. The eddy covariance (EC) raft is marked with the red x.



610 Figure 3. Mean methane ebullition measured weekly in Siikaneva bog in three consecutive years a) 2014, b) 2015 and c) 2016 over different surfaces: bare peat surfaces, open water and water's edge of pools.




Figure 4. Mean weekly methane ebullition with standard error of the means from all surfaces compared to water table (WT) (left panel), and air, water and peat temperatures in the depths of 5, 25 and 50 cm (right panel) measured in Siikaneva bog in years a-b) 2014, c-d) 2015 and e-f) 2016.

615





Figure 5. Mean weekly methane ebullition from all surfaces compared to atmospheric pressure measured in Siikaneva bog in a) 2014, b) 2015 and c) 2016.



620 Figure 6. Ecosystem level methane fluxes measured with the eddy covariance (EC) technique and upscaled from ebullition measurements in a) 2014, b) 2015 and c) 2016. In 2014, ecosystem level methane fluxes are compared also to upscaled chamber fluxes.

Appendix A.

625 Table A1. Mean methane concentrations of the gas caught in the gas traps and of fresh ebullition samples in 2014. Concentration samples were collected four times from gas traps on water (W) and bare peat (BP) surfaces and by triggering fresh ebullition from similar surfaces.

Date	Mean CH ₄ concentration ml/l			
	W gas trap	BP gas trap	W fresh	BP fresh
10-Jun	17	45	293	268
16-Jul	61	<i>245</i>	420	313
13-Aug	86	152	399	172
05-Sep	49	63	379	221

Numbers in Italic indicate the concentration in a single measured gas trap, not a mean of many gas traps.

630




Figure A2. Frequency distribution of methane ebullition ($\text{mg m}^{-2} \text{d}^{-1}$) per gas trap from open water pools (OW), water's edge (EW) and bare peat surfaces (BP) in a-b) 2014, c-d) 2015 and e-f) 2016. Note differences in scales between the years.