

1 **Patterns of soil respiration and its temperature sensitivity in** 2 **grassland ecosystems across China**

3
4 Jiguang Feng¹, Jingsheng Wang², Yanjun Song³, Biao Zhu¹

5
6 ¹Institute of Ecology, College of Urban and Environmental Sciences, and Key Laboratory for Earth Surface
7 Processes of the Ministry of Education, Peking University, Beijing 100871, China

8 ²Qianyanzhou Ecological Research Station, Key Laboratory of Ecosystem Network Observation and
9 Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences,
10 Beijing 100101, China

11 ³Forest Ecology and Forest Management Group, Wageningen University and Research Centre, PO Box 47,
12 6700 AA Wageningen, the Netherlands

13
14 *Correspondence to:* Biao Zhu (biaozhu@pku.edu.cn)

15
16 **Abstract.** Soil respiration (R_s), a key process in the terrestrial carbon cycle, is very sensitive
17 to climate change. In this study, we synthesized 54 measurements of annual R_s and 171
18 estimates of Q_{10} value (the temperature sensitivity of soil respiration) in grasslands across
19 China. We quantitatively analyzed their spatial patterns and controlling factors in five
20 grassland types, including temperate typical steppe, temperate meadow steppe, temperate
21 desert steppe, alpine grassland, and warm-tropical grassland. Results showed that the mean
22 (\pm SE) annual R_s was 582.0 ± 57.9 g C m⁻² yr⁻¹ across Chinese grasslands. Annual R_s
23 significantly differed among grassland types, and positively correlated with mean annual
24 temperature, mean annual precipitation, soil temperature, soil moisture, soil organic carbon
25 content and aboveground biomass, but negatively correlated with soil pH ($p < 0.05$). Among
26 these factors, mean annual precipitation was the primary factor controlling the variations of
27 annual R_s among grassland types. Based on the overall data across Chinese grasslands, the
28 Q_{10} values ranged from 1.03 to 8.13, with a mean (\pm SE) of 2.60 ± 0.08 . Moreover, the Q_{10}
29 values varied largely within and among grassland types and soil temperature measurement

30 depths. Among grassland types, the highest Q_{10} derived by soil temperature at the depth of
31 5 cm occurred in alpine grasslands. In addition, the seasonal variation of soil respiration in
32 Chinese grasslands generally cannot be well explained by soil temperature using the van't
33 Hoff equation. Overall, our findings suggest that the combined factors of soil temperature
34 and moisture would better predict soil respiration in arid and semi-arid regions, highlight
35 the importance of precipitation in controlling soil respiration in grasslands, and imply that
36 alpine grasslands in China might release more carbon dioxide to the atmosphere under
37 climate warming.

38

39 **Keywords:** Soil respiration, Soil carbon emission; Temperature sensitivity; Grassland
40 ecosystem; Q_{10}

41

42 **1 Introduction**

43 Soil respiration (R_s) represents carbon dioxide (CO_2) efflux from the soil surface, including
44 autotrophic/root respiration, and heterotrophic/microbial respiration associated with soil
45 organic matter and litter decomposition (Boone et al., 1998; Kuzyakov, 2006; Schindlbacher
46 et al., 2009). As one of the largest fluxes in the global carbon cycle, R_s plays an important
47 role in regulating ecosystem carbon cycling, carbon-climate feedback, and climate change
48 (Raich and Schlesinger, 1992; Davidson et al., 2002; Luo and Zhou, 2006; Bond-Lamberty
49 and Thomson, 2010). The temperature sensitivity of R_s (Q_{10}), the factor by which R_s is
50 multiplied when temperature increases by 10 °C, is a key parameter to evaluate the feedback
51 intensity between soil carbon efflux and climate warming (Reichstein et al., 2005; Davidson
52 and Janssens, 2006). Knowledge on patterns and controls of R_s and Q_{10} variation at a large
53 scale is crucial for better understanding and modeling soil carbon cycle in a warmer world
54 (Peng et al., 2009; Wang et al., 2010).

55

56 Temperature and precipitation are commonly believed to be the most important climatic
57 factors controlling R_s at the large scale, as suggested by a number of studies (Raich and
58 Schlesinger, 1992; Raich and Potter, 1995; Chen et al., 2014; Hursh et al., 2017). Soil
59 properties, such as soil organic carbon (SOC), soil total nitrogen (STN) and soil pH, can
60 also affect R_s in that they can directly or indirectly affect substrate quality and quantity,

61 which strongly control soil microbial activity and heterotrophic respiration (Ryan and Law,
62 2005; Chen et al., 2010a, 2014; Song et al., 2014). Additionally, biotic factors including
63 decomposer microbes and roots (together with associated mycorrhizal fungi) can directly
64 influence soil respiration via heterotrophic and autotrophic respiration, respectively (Ryan
65 and Law, 2005; Bahn et al., 2010). Previous studies have shown that R_s increased with total,
66 aboveground and belowground net/gross primary production, aboveground biomass (AGB),
67 belowground biomass (BGB), and leaf area index (Raich and Schlesinger, 1992; Hibbard et
68 al., 2005; Bahn et al., 2008; Chen et al., 2014; Zhao et al. 2017).

69

70 As the response of R_s to temperature is controlled by temperature effects on autotrophic
71 respiration from roots and heterotrophic respiration from SOC decomposition, the
72 temperature sensitivity of R_s should be regulated by plant-related biotic variables and soil-
73 related environmental variables. Several studies have shown that climatic factors had strong
74 controls on the spatial variation of Q_{10} , and Q_{10} generally decreased with mean annual
75 temperature (MAT) and mean annual precipitation (MAP) (Raich and Schlesinger, 1992;
76 Kirschbaum, 2000; Peng et al., 2009; Song et al., 2014). In addition to climatic variables,
77 the spatial variation of Q_{10} could be affected by seasonality of plant activity. Previous studies
78 suggested that plant growth plays an important role in the seasonal variation of R_s , and
79 thereby the seasonal dynamic changes in plant activity affect seasonal Q_{10} (Yuste et al., 2004;
80 Wang et al., 2010). Furthermore, Q_{10} is also affected by soil properties, such as soil
81 temperature, soil moisture, soil pH, SOC and STN, which can directly influence root and
82 microbial activities, substrate availability and nutrient supply (Zhou et al., 2009; Song et al.
83 2014; Zhao et al., 2017).

84

85 Grasslands in China cover 29–41% of its total land area (Shen et al., 2016) and have
86 significant effects on regional climate and carbon cycle (Ni, 2002). In China, grasslands are
87 widely distributed throughout the country, and the different climate gradients and landforms
88 in China support a number of grassland types, including tropical, warm, temperate, and
89 alpine grassland, etc. (Chen et al., 2002; Shen et al., 2016). Specifically, the temperate arid
90 and semi-arid grasslands in Inner Mongolia, and the alpine meadow and steppe in Qinghai-
91 Tibet Plateau comprise the main body of temperate and alpine grasslands, respectively
92 (Shen et al., 2016). In the past two decades, a large number of case studies on R_s have been
93 widely conducted in grasslands across China. However, few have been included in global
94 R_s and Q_{10} syntheses (Raich and Schlesinger, 1992; Wang et al., 2010; Bond-Lamberty and

95 Thomson, 2010; Chen et al., 2014; Hursh et al., 2017), largely because most studies were
96 published in Chinese journals. Given the diverse grassland types, especially alpine
97 grasslands distributed in China, R_s and Q_{10} may vary among grassland types due to the
98 differences in abiotic and biotic factors, and the patterns of R_s and Q_{10} across Chinese
99 grasslands may differ from global terrestrial ecosystems and grasslands. However, how the
100 spatial variation of R_s and Q_{10} varies with abiotic and biotic factors across Chinese
101 grasslands and their differences among grassland types still remain poorly understood.

102

103 In this study, we synthesized all the available data relating to R_s and Q_{10} in grasslands across
104 China. Our main objectives were to: (1) analyze the spatial patterns of R_s and Q_{10} across
105 various grassland ecosystems in China; (2) compare the differences in R_s and Q_{10} among
106 grassland types; (3) identify how abiotic and biotic factors drive R_s and Q_{10} among sites at
107 the regional scale, including geographic variables, climatic factors, soil properties and biotic
108 factors; and (4) compare the R_s and Q_{10} in Chinese grasslands with those from previous
109 syntheses at the global and regional scale.

110

111 **2 Materials and Methods**

112 **2.1 Data collection**

113 Peer-reviewed journal articles and published theses (including available online) before 1
114 December, 2017 were searched using Web of Science and China National Knowledge
115 Infrastructure (CNKI, available online: <http://epub.cnki.net>) with the following search term
116 combinations: (soil respiration OR soil CO₂ flux OR soil CO₂ efflux OR soil CO₂ emission
117 OR soil carbon flux OR soil carbon efflux OR soil carbon emission) AND (grassland OR
118 steppe OR meadow OR grass). Additional searches with the same keywords were conducted
119 on ScienceDirect (Elsevier Ltd., Amsterdam, Nederland), Springer Link (Springer
120 International Publishing AG, Berlin, Germany), and Wiley Online Library (John Wiley &
121 Sons Ltd., Hoboken, USA). Furthermore, previous global and regional syntheses on the
122 similar topic were also screened to check Chinese grassland data, such as Peng et al. (2009),
123 Wang and Fang (2009), Bond-Lamberty and Thomson (2010), Wang et al. (2010) and Chen
124 at al. (2010, 2014).

125

126 To ensure data consistency and accuracy, the following six criteria were applied to select
127 appropriate studies: (1) experimental studies were conducted in the field; (2) experiments
128 with the treatments of nitrogen (fertilizer) addition, increased or decreased precipitation,
129 warming, elevated CO₂, simulated acid rain, clipping, and grazing were removed; (3) the
130 study must contain soil respiration or Q_{10} with a clear record of grassland type and
131 experimental duration; (4) the investigation time for measuring Rs was not less than twelve
132 months so that the annual Rs can be obtained, and modeled annual Rs based on the
133 relationships between Rs rate and temperature were not included; (5) the investigation time
134 for estimating Q_{10} value was not less than four months; and (6) Q_{10} values were calculated
135 by the van't Hoff equation (Van's Hoff, 1898).

$$136 \quad SR = \alpha \times \exp(\beta \times T) \quad (1)$$

137 where SR is the measured soil respiration rate, T is the measured soil temperature at the
138 given depth, and coefficient α and β are fitted parameters. The Q_{10} values were calculated
139 as:

$$140 \quad Q_{10} = \exp(10\beta) \quad (2)$$

141 Several studies measured Rs and its temperature sensitivity at different years, and then these
142 Rs and Q_{10} values were averaged across years. In this case, only the highest R_Q^2 (coefficient
143 of determination for calculating Q_{10} using Eq. (1)) was extracted if more than one R_Q^2 were
144 available in the same study. In addition, the Q_{10} values were estimated by Rs measured at
145 different plant growth stages, and they were further categorized into three types according
146 to the Rs measurement period, including growing season Q_{10} , non-growing season Q_{10} and
147 annual Q_{10} . If these three types of Q_{10} were all available, only the annual Q_{10} was selected
148 in our database. Within these constraints, 54 measurements of annual Rs rate and 171
149 estimates of Q_{10} value were obtained from 108 published experimental studies across
150 Chinese grasslands (Table S1). Our database contained 14 variables associated with Rs,
151 including annual Rs, growing and non-growing season Rs and their proportions to annual
152 Rs, the proportion of autotrophic and heterotrophic respiration to annual Rs, Q_{10} values of
153 Rs and their corresponding R_Q^2 . Here, the growing season was from May to October, and
154 the non-growing season was from November to April in the next year. The Q_{10} values were
155 divided into five soil depths with different soil temperature (ST0, soil surface temperature;
156 ST5, soil temperature at 5 cm; ST10, soil temperature at 10 cm; ST15, soil temperature at
157 15 cm; and ST20, soil temperature at 20 cm) for the same site. In one study, the Q_{10} was
158 derived by soil temperature at the depth of 6 cm, and then it was treated as Q_{10-ST5} because
159 of little difference in soil temperature between 5 cm and 6 cm.

160

161 In most of publications, the R_s , Q_{10} and its RQ^2 of the model were presented, and they were
162 incorporated into our database directly. The R_s , Q_{10} and RQ^2 values were recalculated
163 according to the available information if these values were not directly provided in some
164 publications. The growing season, non-growing season and annual R_s were obtained by
165 interpolating measured R_s rate between respective sampling dates for each seasonal
166 measurement period of the year, and then computing the sum to obtain the corresponding
167 values (Frank and Dugas, 2001; Sims and Bradford, 2001) as follows:

$$168 \quad CSR = \sum (\Delta t_k \times F_{m,k}) \quad (3)$$

169 where CSR is cumulative soil respiration during the season, $\Delta t_k (= t_k - t_{k-1})$ is the time
170 interval between each field measurement within the season, and $F_{m,k}$ is the average R_s rate
171 over the interval $t_k - t_{k-1}$.

172

173 In addition, for each study site, we also recorded other supporting information from the
174 original publications, including grassland type, geographic variables (longitude and
175 latitude), climatic factors (MAT and MAP), soil properties (soil temperature, soil moisture,
176 soil pH, SOC and STN), and biotic factors (microbial biomass carbon (MBC), AGB and
177 BGB). Missing climatic information was obtained using NASA Surface meteorology and
178 Solar Energy-Location, and the other missing information was obtained from the related
179 references according to the study site and described experiment design. Several studies
180 provided the soil organic matter content, which was converted to SOC by multiplying a
181 conversion factor of 0.50 (Pribyl 2010). In case of gravimetric soil moisture being
182 provided, it was converted to volumetric soil moisture according to soil bulk density. Given
183 that BGB were measured in different soil depths, only BGB measured in 0–40 and 0–50 cm
184 soil depths were selected because roots were mainly distributed in 0–50 cm and there were
185 minor difference between 0–40 and 0–50 cm. The distributions of selected experimental
186 sites were shown in Fig. 1.

187

188 **2.2 Data analysis**

189 In this study, grasslands were divided into five groups, including temperate typical steppe,
190 temperate meadow steppe, temperate desert steppe, alpine grassland, and warm-tropical
191 grassland. If grassland type was not provided directly, it was determined according to the
192 dominant plant species reported in selected publications and the Classification of Grassland

193 Ecosystem in China (Chen et al., 2002). Detailed statistical parameters for the five grassland
194 types were presented in Table S2.

195

196 One-way analysis of variance (ANOVA) was used to examine whether annual R_s and Q_{10}
197 values differed among grassland types, measuring periods or measuring methods. In case of
198 homogeneity of variances, the least significant differences (LSD) test was applied;
199 otherwise, the Dunnett T3 test was applied. Paired-samples t -test was performed to compare
200 the differences between growing season and non-growing season R_s , and between
201 autotrophic respiration and heterotrophic respiration. The reason for using paired-samples
202 t -test was that these two corresponding variables were interconnected as they were from the
203 same study sites. In addition, we used two statistical methods to explore the differences for
204 Q_{10} among measurement depths. The paired-samples t -test was used to compare Q_{10} among
205 different measurement depths from same sites, whereas the ANOVA was used to compare
206 Q_{10} among different measurement depths from all sites. Compared with ANOVA, the
207 advantages of paired-samples t -test was that it avoided the effects of unequal spatial
208 distribution of samples from different depths on Q_{10} and only compared the effects of
209 measurement depth. The univariate regression analysis was used to identify the relationships
210 between annual R_s , Q_{10} , and a given biotic or abiotic factor mentioned above, except for
211 MBC because of its limited sample size. The multiple linear regression analyses were also
212 performed to identify the comprehensive effects of environmental variables (including
213 MAT, MAP, soil temperature and moisture as they had relatively enough sample sizes) on
214 annual R_s , and Q_{10} derived by ST5 and ST10. Correlations among these factors were
215 calculated with the Pearson correlation coefficients. All statistical analyses were performed
216 using the software IBM SPSS Statistics 20.0 (IBM Corporation, New York, USA).

217

218 **3 Results**

219 **3.1 Soil respiration and its controlling factors**

220 **3.1.1 Patterns of annual soil respiration across Chinese grasslands**

221 The annual R_s ranged from 122.9 to 2407.1 g C m⁻² yr⁻¹, with the total mean (\pm SE) of
222 582.0 \pm 57.9 g C m⁻² yr⁻¹. There were significant differences in annual R_s between grassland
223 types ($p < 0.001$), with the highest annual R_s in the warm-tropical grassland and the lowest

224 annual Rs in the temperate desert steppe (Table 1). The proportions of growing season or
225 non-growing season Rs varied slightly among different grassland types ($p > 0.05$), but the
226 proportion of Rs in growing season was significantly higher than that in non-growing season
227 ($p < 0.001$). Overall, growing season and non-growing season Rs consisted of 78.7% and
228 21.3% of the annual Rs, respectively, across all grasslands in China (Table 1). In addition,
229 growing season Rs was significantly positively correlated with the annual Rs based on linear
230 regression model ($r^2 = 0.923$, $p < 0.001$, Fig. S1). At the annual scale, the mean contribution
231 of heterotrophic respiration to Rs was 72.8% across Chinese grasslands, which was
232 significantly larger than that of autotrophic respiration with the mean of 27.2% ($p < 0.01$,
233 Fig. S2).

234

235 **3.1.2 Spatial controls of abiotic and biotic factors over soil respiration**

236 In the univariate linear regressions, annual Rs significantly increased with MAT, MAP, soil
237 temperature, soil moisture, SOC, and AGB across all grasslands in China, but decreased
238 with soil pH ($p < 0.05$, Fig. 2). In contrast, annual Rs did not correlate well with STN and
239 BGB ($p > 0.05$). The single factor of MAT, MAP, soil temperature, soil moisture, SOC, soil
240 pH, and AGB accounted for 22.4%, 31.3%, 20.9%, 32.0%, 29.6%, 20.6%, and 36.1% of the
241 spatial variation of annual Rs, respectively (Fig. 2).

242

243 **3.2 Temperature sensitivity of soil respiration and its controlling factors**

244 **3.2.1 Distributions of Q_{10} values and its coefficient of determination**

245 Most of the Q_{10} values (83.0%) were distributed between 1.5 and 3.8. However, the
246 distributions of Q_{10} values derived by the five soil depths were different (Fig. 3a-e). The
247 largest relative frequency for Q_{10-ST5} and $Q_{10-ST10}$ values was within the range of 1.5 to 3.0
248 (68.5%) and 1.5 to 3.5 (83.1%), respectively, while that of Q_{10-ST0} was mainly within 1.0–
249 2.0 (88.2%, Fig. 3). In addition, the distribution of $Q_{10-ST15}$ and $Q_{10-ST20}$ were relatively
250 uniform (Fig. 3d and e).

251

252 Similarly, the distributions of R_{Q^2} for Q_{10} derived by the five soil depths also differed from
253 each other (Fig. 3f-g). The R_{Q^2} values for Q_{10-ST5} and $Q_{10-ST10}$ were mainly distributed in
254 0.6–0.9 and 0.5–0.7, respectively, while those for $Q_{10-ST15}$ and $Q_{10-ST20}$ were mainly

255 distributed in 0.3–0.6. The R_Q^2 value for Q_{10-ST0} was distributed uniformly (Fig. 3f). Overall,
256 only 35.6% of R_Q^2 values for Q_{10} were within the range of 0.7–1.0.

257

258 3.2.2 Patterns of Q_{10} values across Chinese grasslands

259 Across all grasslands, the overall Q_{10} values ranged from 1.03 to 8.13, with the mean (\pm SE)
260 of 2.60 ± 0.08 . Specifically, the mean (\pm SE) of Q_{10} values derived by ST0, ST5, ST10,
261 ST15, and ST20 was 1.73 ± 0.08 , 2.80 ± 0.14 , 2.56 ± 0.12 , 2.64 ± 0.33 , and 2.81 ± 0.31 ,
262 respectively (Fig.3 a-e). Paired t -test demonstrated that Q_{10} significantly differed between
263 two adjacent depths in the top 15 cm soil ($p < 0.05$), whereas no difference occurred below
264 15 cm depth ($p > 0.05$; Fig. 4). Generally, the overall Q_{10} and paired Q_{10} increased with soil
265 temperature measurement depth (Fig. 4; Fig. S3). In terms of grassland types, there were
266 significant differences for Q_{10} derived by ST5 and ST10 among grassland types, respectively
267 ($p < 0.05$, Fig. 4b and c). For Q_{10} derived by ST5, it was highest in alpine grassland, while
268 for Q_{10} derived by ST10, the highest value was in warm-tropical grassland. In addition, Q_{10}
269 values derived by ST0, ST15 and ST20 were not enough to meet the demand of statistical
270 analysis, so their differences among grassland types were not examined.

271

272 3.2.3 Spatial controls of environmental factors over Q_{10}

273 The relationships of Q_{10-ST5} and $Q_{10-ST10}$ with abiotic and biotic factors were presented in
274 Fig. 5. Among these abiotic and biotic factors, Q_{10-ST5} significantly positively correlated
275 with SOC, AGB and BGB, whereas negatively correlated with soil temperature ($p < 0.05$,
276 Fig. 5). In contrast, $Q_{10-ST10}$ significantly correlated with MAP, soil temperature and SOC (p
277 < 0.05 , Fig. 5). In addition, combined MAP and MAT, and combined soil temperature and
278 soil moisture affected Q_{10-ST5} , and jointly accounted for 22.1% and 13.9% of the spatial
279 variation of Q_{10-ST5} across Chinese grasslands (Table S4).

280

281 **4 Discussion**

282 **4.1 Spatial patterns and controlling factors of annual soil respiration**

283 **4.1.1 Annual soil respiration among grassland types**

284 In arid and semi-arid ecosystems, such as grassland and desert, MAP might play a key role
285 in controlling carbon cycling. Our results also suggested that MAP had significant controls
286 on mean annual Rs among various grassland types in China ($p < 0.01$, Fig S4). The
287 significant difference in mean annual Rs might be mainly attributed to the differences in
288 AGB, BGB and microbial activity induced by precipitation across various grassland types.
289 Previous studies suggested that grasslands with higher MBC had larger heterotrophic
290 respiration (Colman and Schimel, 2013; Ding et al., 2016). Meanwhile, a regional study
291 demonstrated that microbial biomass increased with MAP in grasslands (Chen et al., 2016b),
292 which was also found in this study (Table S3). Altogether, these suggested that the regions
293 with high MAP would have larger heterotrophic respiration. Additionally, a previous study
294 demonstrated that both AGB and BGB increased with MAP across Chinese grasslands (not
295 including warm-tropical grasslands) (Ma et al., 2014). In this case, autotrophic respiration
296 would be higher in the grasslands with high plant biomass. Collectively, the grasslands with
297 high MAP would have relatively higher Rs rate.

298

299 **4.1.2 Controls of environmental factors on annual Rs**

300 Across Chinese grasslands, annual Rs was strongly related to MAT, MAP, soil temperature
301 and soil moisture, which was consistent with previous results obtained from global
302 terrestrial ecosystems (Raich and Schlesinger, 1992; Raich and Potter 1995; Chen et al.,
303 2014; Hursh et al. 2017), global grasslands (Wang and Fang, 2009), and Chinese forests
304 (Song et al., 2014; Xu et al., 2015). Compared with MAT and soil temperature, MAP and
305 soil moisture explained more spatial variations of annual Rs, suggesting that these two
306 factors are more important in predicting Rs in arid and semi-arid ecosystems under climate
307 change.

308

309 In addition, spatial variations of annual Rs were also controlled by soil properties, such as
310 SOC and soil pH. The relationships between annual Rs and SOC as well as pH were also
311 observed in global, regional and local terrestrial ecosystems (Chen et al., 2010b, 2014; Song

312 et al., 2014; Xu et al., 2016). Since R_s involves the process of converting organic carbon
313 into inorganic carbon, the soil CO_2 emission from microbial decomposition of soil organic
314 carbon is ultimately determined by the supply of C substrate (Wan et al., 2007).
315 Additionally, soil pH can directly regulate the activities of microbes and C-acquiring
316 enzymes (Turner, 2010). In neutral and alkaline soils, microbial biomass tended to decrease
317 with soil pH (Ding et al., 2016). Therefore, this led to a negative correlation between R_s and
318 soil pH in Chinese grasslands because most of grasslands in China are distributed in neutral
319 and alkaline soils. Further, Chen et al. (2010b) demonstrated that annual R_s significantly
320 increased with soil total nitrogen at global scale. Meanwhile, some case studies revealed the
321 similar relationship between growing season R_s and soil total nitrogen among different
322 grassland types and vegetation communities (Chen et al., 2010a; Wang et al., 2015; Xu et
323 al., 2016) at local scales, while annual R_s did not correlate well with STN in this study.
324 Given that SOC and STN are closely associated with one another (Table S3), the
325 insignificant correlation of R_s with STN might be due to the fact that soil total nitrogen
326 might not well represent nitrogen availability for plants and microbes. Therefore, how STN
327 influence R_s across Chinese grasslands at regional scale should be further studied.

328

329 Furthermore, as the source of autotrophic respiration, BGB can directly influence R_s , which
330 has been observed in ecosystems at global and local scale (Chen et al., 2010a, 2014).
331 However, no significant correlation between BGB and R_s was observed in the present study,
332 which might be attributed to the limited sample size ($n = 6$) and the uncertainty in measuring
333 BGB (due to inconsistent or insufficient sampling depth). In grassland ecosystems, BGB
334 generally increased with AGB (Ma et al., 2014), and this relationship was also observed in
335 this study ($p < 0.10$, Fig. S5). Therefore, given the significant correlation between AGB and
336 R_s in Chinese grasslands (Fig. 2), BGB may also have the potential to control annual R_s
337 across Chinese grasslands, although this should be further investigated based on accurate
338 quantification of BGB and R_s across a large number of sites.

339

340 **4.2 Spatial patterns and controlling factors of Q_{10} values**

341 **4.2.1 R_Q^2 for Q_{10} in Chinese grasslands**

342 In this study, only 37.3% of R_Q^2 for Q_{10} was larger than 0.7, indicating that most of the
343 seasonal variation of R_s rate cannot be well explained by soil temperature using the van't

344 Hoff equation (Eq. 2). Compared with the results obtained from Chinese forests (Xu et al.,
345 2015), the van't Hoff equation (Eq. 2) was not very suitable to describe the relationships
346 between Rs rate and soil temperature in most of Chinese grasslands. This might be
347 associated with the difference in soil moisture between these two ecosystems because
348 besides temperature, soil moisture may strongly influence the apparent Q_{10} (Subke and
349 Bahn, 2010). Previous studies have suggested that in humid and semi-humid regions the
350 effect of soil moisture on Rs is weak, whereas in arid and semi-arid regions, Rs is
351 significantly influenced by soil moisture (Jia et al., 2006; Li et al., 2011; Wang et al., 2014a,
352 2014b). Moreover, some studies showed that soil moisture and temperature had an
353 interactive effect on the seasonal variations of Rs rate (Davidson et al., 1998; Jia et al., 2006;
354 Wang et al., 2014b; Liu et al., 2016), indicating that the two-variable equations could better
355 explain the variation in Rs than the single variable of temperature. Our results also showed
356 that, in general, R_Q^2 for Q_{10} closely increased with MAP and soil moisture ($p < 0.05$, Fig.
357 S6), indicating that the R_Q^2 for Q_{10} tended to be larger in the regions with abundant
358 precipitation. Collectively, for ecosystems (e.g., grassland and desert) in arid and semi-arid
359 regions, Rs could be better estimated by the combined factors of soil temperature and
360 moisture. By comparison, 46.6% of R_Q^2 for Q_{10-ST5} was distributed in 0.7–1.0, which was
361 higher than those derived by soil temperature at other depths, suggesting that the seasonal
362 variation of Rs can be better explained by soil temperature at the depth of 5 cm across
363 Chinese grasslands.

364

365 **4.2.2 Q_{10} among soil depths and grassland types**

366 In Chinese grasslands, the estimated Q_{10} generally increased with soil temperature
367 measurement depth, which was consistent with previous synthesis study about Chinese
368 ecosystems (Peng et al., 2009). The differences for Q_{10} among measurement depths might
369 be due to the seasonal amplitudes of temperature at different soil depths (Pavelka et al.,
370 2007; Graf et al., 2008).

371

372 In terms of grassland types, the highest Q_{10-ST5} was in the alpine grassland and the lowest in
373 the temperate desert steppe and typical steppe (Fig. 4). This difference could be associated
374 with soil properties and climatic conditions. For example, it is well known that the alpine
375 grasslands are usually distributed in high altitude regions (above 3000 m), where the climate
376 is relatively colder and SOC is relatively higher than the other grassland types (Table S2).

377 However, the temperate desert steppes and typical steppes are mainly distributed in north
378 China, with relatively high MAT and low MAP that may lead to low Q_{10} . Moreover, as
379 shown in Fig. 4, the highest $Q_{10-ST10}$ occurred in warm-tropical grassland, which might be
380 associated with the abundant substrate supply in this grassland type because high substrate
381 availability can enhance apparent Q_{10} of soil respiration (Davidson et al., 2006; Zhu and
382 Cheng, 2011).

383

384 **4.2.3 Controls of environmental factors on Q_{10}**

385 Generally, the Q_{10} derived by either ST5 or ST10 did not correlate well with climatic factors,
386 which was inconsistent with previous results at global and regional scales (Chen and Tian,
387 2005; Peng et al., 2009; Wang et al., 2010; Song et al., 2014; Xu et al., 2015). However, we
388 found that Q_{10} derived by soil temperature at the depth of 5 and 10 cm decreased closely
389 with increasing soil temperature, partly supporting the idea that Q_{10} tends to be higher in
390 colder regions. Additionally, the positive relationships of Q_{10-ST5} with SOC, AGB and BGB
391 indicated that soil properties and plant biomass can also profoundly influence the spatial
392 variation of Q_{10} . Previous studies suggested higher plant biomass and SOC can lead to more
393 substrate supply for soil respiration and then result in higher Q_{10} values, because apparent
394 Q_{10} increased with increasing substrate availability (Gershenson et al., 2009; Zhao et al.,
395 2017).

396

397 The extremely low R^2 value for the relationship of Q_{10} with climatic variables suggested
398 that the single factor of temperature, precipitation or soil moisture poorly control the spatial
399 variations of Q_{10} in Chinese grasslands. Therefore, the variation of Q_{10} in Chinese grasslands
400 should be controlled by multiple factors due to the complex and diverse environments
401 among grasslands at large scale. Multiple linear regression analyses also showed that
402 combined MAT and MAP, and combined soil temperature and moisture could better explain
403 the variations of Q_{10} derived by ST5 (Table S4), indicating their integrative effects on the
404 spatial variation of Q_{10-ST5} . Additionally, both univariate and multiple regression analyses
405 demonstrated that there were no significant relationships between $Q_{10-ST10}$ and abiotic and
406 biotic factors (not shown), indicating that the $Q_{10-ST10}$ might not have clear spatial pattern or
407 its variation might be controlled by other factors.

408

409 In addition to the environment variables discussed above, seasonality of plant activity could

410 also affect the spatial variation of Q_{10} at large scale. Plant activity can directly affect R_s via
411 controlling root respiration, and can indirectly affect SOC decomposition by microbes via
412 regulating rhizosphere priming effect (see Wang et al., 2010). In this study, the dataset
413 covered various climatic regions, and accordingly seasonal amplitudes of plant activity
414 among grassland types were also different. A previous global synthesis using NDVI
415 (normalized difference vegetation index) as an indicator of plant activity demonstrated that
416 seasonal amplitude of plant activity dominated the variation of seasonal Q_{10} among different
417 sites (Wang et al., 2010). Therefore, the seasonal amplitude of plant activity might be an
418 important factor explaining the spatial variation of Q_{10} across Chinese grasslands, and
419 should be further studied.

420

421 **4.3 Comparisons of R_s and Q_{10} between Chinese grasslands and the global ecosystems**

422 **4.3.1 Comparisons of annual R_s**

423 The annual R_s varied largely within and among the grassland types across China (Table 1),
424 with the mean value of $582.0 \text{ g C m}^{-2} \text{ yr}^{-1}$, which was much lower than those in global
425 terrestrial ecosystems and in Chinese forests (Table 2). For these results, the main biomes
426 in their dataset were forests, which had relatively high precipitation and net primary
427 productivity (Hursh et al., 2017), leading to relatively higher R_s than grasslands (Table S2).
428 Compared with global grasslands, our result was much lower or higher than the results
429 obtained from Chen et al. (2010b, 2014) and Wang and Fang (2009), but approximately
430 consistent with Hursh et al. (2017). These differences might be associated with data sources
431 and distributions of case study sites. In general, the mean annual R_s rate across Chinese
432 grasslands was within the lowest and highest R_s across global grasslands.

433

434 Across Chinese grasslands, the proportions of R_s during growing season ranged from 76.2–
435 86.8%, which were 2.2–5.6 times higher than those during non-growing season. Microbial
436 activity and plant growth is constrained by temperature and precipitation during non-
437 growing season, leading to lower decomposition of soil organic carbon and root respiration.
438 In addition, as a whole, heterotrophic respiration contributed 72.8% of the annual R_s , 2.7
439 times of autotrophic respiration, which was close to that of global terrestrial ecosystems and
440 grasslands (Wang and Fang, 2009; Chen et al., 2014) and Chinese forests (Song et al., 2014).
441 Previous studies suggested that the proportions of heterotrophic respiration to total R_s varied

442 with ecosystem types and depended on the magnitude of total Rs (Subke et al., 2006).
443 However, the limited samples ($n = 7$) limited our comparisons among these grassland types.
444 Generally, our findings and previous studies suggested that both Rs during growing season
445 and heterotrophic respiration were an important part of the annual Rs in Chinese grasslands,
446 respectively, and should be given enough attention.

447

448 **4.3.2 Comparisons of Q_{10}**

449 The overall mean Q_{10} of 2.60 derived by soil temperature at all measurement depths was
450 similar to 2.40 and 2.54 in global terrestrial ecosystems (Raich and Schlesinger, 1992;
451 Lenton and Huntingford, 2003). The Q_{10} derived by ST5 varied from 1.39 to 8.13, with the
452 mean of 2.80, which was higher than that of global and Chinese terrestrial ecosystems,
453 Chinese forests, and particularly global grasslands (Table 2). The difference may be partly
454 due to the distribution of grasslands in China and the grassland types. Chinese grasslands
455 are mainly distributed in the high latitude (temperate grassland) and high altitude (Qinghai-
456 Tibet Plateau alpine grassland) regions, and Q_{10} takes relatively higher values in cold
457 regions than in warm regions (Chen and Tian, 2005; Wang et al., 2010). In addition, in this
458 study, averaged Q_{10-ST5} was highest in alpine grassland with the mean of 3.30, implying that
459 grasslands in alpine regions may release more carbon dioxide under climate warming.
460 However, there were no alpine grasslands in the global database. Collectively, this may lead
461 to higher Q_{10} value in Chinese grasslands. In terms of Q_{10} derived by ST10, the mean value
462 for Chinese grasslands was close to Chinese terrestrial ecosystems, but much lower than the
463 global ecosystems (Table 2).

464

465 **4.4 Uncertainties**

466 In order to ensure data consistency and minimize the error, only field experiments in
467 accordance with the six aforementioned criteria were selected. However, the inter-annual
468 variation in Rs and Q_{10} might be very large for grassland at one site, which was associated
469 with the inter-annual variations in annual precipitation and mean temperature (Peng et al.,
470 2014; Wang et al., 2016). Therefore, the inter-annual variation of Rs would impact the
471 accuracy of the results. Additionally, three methods including static closed chamber,
472 dynamic closed chamber, and alkali absorption were widely applied to measure Rs in the

473 selected experiments, and previous studies have suggested that measurement methods
474 affected the results of Rs rate and Q_{10} value (Bekku et al., 1997; Yim et al., 2002; Peng et
475 al., 2009). However, in this study, there were generally no significant differences for Rs,
476 Q_{10-ST5} and $Q_{10-ST10}$ among the three measurement methods (Fig. S7). Given that only one
477 sample of annual Rs was measured by alkali absorption, the effects of measurement methods
478 on Rs could be neglected. Therefore, including data measured by the alkali absorption
479 method in our synthesis does not meaningfully change the results of Rs and Q_{10} .

480

481 Furthermore, Q_{10} values measured during three periods, including growing season, non-
482 growing season and the whole year, were selected as long as the investigation time was
483 longer than four months. The seasonal dynamics of plant growth and microbial activity may
484 influence autotrophic and heterotrophic respiration, thus the Q_{10} of Rs. Our results showed
485 that measurement period did not significantly affect $Q_{10-ST10}$, but significantly affected Q_{10-}
486 $ST5$ (Fig. S7). In terms of Q_{10-ST5} , the significant differences between annual Q_{10} and non-
487 growing season Q_{10} across all sites was mainly caused by alpine grasslands, in which annual
488 Q_{10} was much higher than non-growing season Q_{10} (Fig. S7). Likely, the seasonal amplitude
489 of plant activity at annual scale is much greater than that at non-growing season scale in
490 alpine regions. Therefore, the different investigation time and measurement period for
491 estimating Q_{10} would inevitably affect the accuracy of results.

492

493 In this study, the selected experiments were mainly conducted in temperate and alpine
494 grasslands, so the limited data obtained from desert, tropical and subtropical grasslands
495 might lead to some uncertainties in these ecosystems. Moreover, grassland management
496 practices such as land use/cover change, intensity and pattern of livestock grazing, and
497 fencing can have significant effect on soil carbon emission (Chen et al., 2013; Zhang et al.,
498 2015b; Chen et al., 2015; Chen et al., 2016a). In the past three decades, several ecological
499 projects relating to grassland have been implemented in China, and have observably
500 increased the grassland area and altered the land cover (Zhang et al., 2015a). To some extent,
501 these changes can also impact our findings.

502

503 **5 Conclusion**

504 Chinese grasslands cover vast area, have high spatial heterogeneity, and include various

505 grassland types. By synthesizing all the available data relating to R_s and Q_{10} , we analyzed
506 their spatial patterns and driving factors in grasslands across China. Our results showed that
507 annual R_s and Q_{10} varied greatly within and among grassland types. Across Chinese
508 grasslands the mean annual R_s and Q_{10} were $582.0 \text{ g C m}^{-2} \text{ yr}^{-1}$ and 2.60, respectively. MAT,
509 MAP, soil temperature, soil moisture, SOC and AGB all significantly positively affected
510 annual R_s , whereas soil pH negatively affected annual R_s . Among these environmental
511 factors, MAP played an important role in controlling R_s variations across Chinese
512 grasslands. Moreover, the combined factors of MAP and MAT accounted for 22.1% of the
513 variation of Q_{10-ST5} across Chinese grasslands. The Q_{10-ST5} in Chinese grasslands was much
514 higher than that in global ecosystems, mainly attributed to the higher Q_{10} value in alpine
515 grasslands. These findings together advance our understanding of the spatial variation and
516 environmental control of R_s and Q_{10} across Chinese grasslands, and also improve our ability
517 to predict soil carbon efflux under climate change at regional scale. However, the few
518 experiments measuring soil and microbial variables, R_s and Q_{10} at annual scale, especially
519 measuring autotrophic and heterotrophic respiration separately, limit our in-depth
520 knowledge on the key drivers of R_s and Q_{10} in grasslands across China. Therefore, more
521 field measurements are needed to verify the relationships found here and reveal how
522 environmental variables control R_s and its temperature sensitivity in relatively arid
523 grassland ecosystem.

524

525 **Acknowledgement**

526 This research was financially supported by the National Key Research and Development
527 Program of China (2017YFC0503903 and 2016YFC0502006), the National Natural
528 Science Foundation of China (31621091 and 31622013), and the Key Science and
529 Technology Plan Projects of Tibet Autonomous Region (Z2016C01G01/08-004 and
530 Z2016C01G01/03). We thank all the scientists whose data and work were included in this
531 synthesis. We are also grateful to the editor (Dr. Zhongjun Jia) and two anonymous
532 reviewers for their insightful comments which improved the manuscript.

533

534 **References**

- 535 Bahn, M., Janssens, I. A., Reichstein, M., Smith, P., and Trumbore, S. E.: Soil respiration across scales:
536 towards an integration of patterns and processes, *New Phytol.*, 186, 292–296, doi: 10.1111/j.1469-
537 8137.2010.03237.x, 2010.
- 538 Bahn, M., Rodeghiero, M., Andersondunn, M., Dore, S., Gimeno, C., Dr ösler, M., Williams, M., Ammann, C.,
539 Berninger, F., and Flechard, C.: Soil respiration in european grasslands in relation to climate and assimilate
540 supply, *Ecosystems*, 11, 1352–1367, doi: 10.1007/s10021-008-9198-0, 2008.
- 541 Bekku, Y., Koizumi, H., Oikawa, T., and Iwaki, H.: Examination of four methods for measuring soil respiration,
542 *Appl. Soil Ecol.*, 5, 247–254, doi: 10.1016/S0929-1393(96)00131-X, 1997.
- 543 Bond-Lamberty, B. and Thomson, A.: A global database of soil respiration data. *Biogeosciences*, 7, 1915–
544 1926, doi: 10.5194/bg-7-1915-2010, 2010.
- 545 Boone, R. D., Nadelhoffer, K. J., Canary, J. D., and Kaye, J. P.: Roots exert a strong influence on the
546 temperature sensitivity of soil respiration, *Nature*, 396, 570–572, doi: 10.1038/25119, 1998.
- 547 Chen, H. and Tian, H. Q.: Does a general temperature-dependent Q_{10} model of soil respiration exist at biome
548 and global scale? *J. Integr. Plant Biol.*, 47, 1288–1302, doi: 10.1111/j.1744-7909.2005.00211.x, 2005.
- 549 Chen, J., Hou, F., Chen, X., Wan, X., and Millner, J.: Stocking rate and grazing season modify soil respiration
550 on the Loess Plateau, China, *Rangel. Ecol. Manag.*, 68, 48–53, doi: 10.1016/j.rama.2014.12.002, 2015.
- 551 Chen, J., Zhou, X., Wang, J., Hruska, T., Shi, W., Cao, J., Zhang, B., Xu, G., Chen, Y., and Luo, Y.: Grazing
552 exclusion reduced soil respiration but increased its temperature sensitivity in a meadow grassland on the
553 Tibetan Plateau, *Ecol. Evol.*, 6, 675–687, doi: 10.1002/ece3.1867, 2016a.
- 554 Chen, Q., Wang, Q., Han, X., Wan, S., and Li, L.: Temporal and spatial variability and controls of soil
555 respiration in a temperate steppe in northern China, *Global Biogeochem. Cycle*, 24, doi:
556 10.1029/2009GB003538, 2010a.
- 557 Chen, S., Huang, Y., Zou, J., Shen, Q., Hu, Z., Qin, Y., Chen, H., and Pan, G.: Modeling interannual variability
558 of global soil respiration from climate and soil properties, *Agr. Forest Meteorol.*, 150, 590–605, doi:
559 10.1016/j.agrformet.2010.02.004, 2010b.
- 560 Chen, S., Zou, J., Hu, Z., Chen, H., and Lu, Y.: Global annual soil respiration in relation to climate, soil
561 properties and vegetation characteristics: Summary of available data, *Agri. Forest Meteorol.*, 198–199,
562 335–346, doi: 10.1016/j.agrformet.2014.08.020, 2014.
- 563 Chen, W., Wolf, B., Zheng, X., Yao, Z., Butterbach-Bahl, K., Brueggemann, N., Han, S., Liu, C., and Han, X.:
564 Carbon dioxide emission from temperate semiarid steppe during the non-growing season, *Atmos. Environ.*,
565 64, 141–149, doi: 10.1016/j.atmosenv.2012.10.004, 2013.
- 566 Chen, Y., Ding, J., Peng, Y., Li, F., Yang, G., Liu, L., Qin, S., Fang, K., and Yang, Y.: Patterns and drivers of
567 soil microbial communities in Tibetan alpine and global terrestrial ecosystems, *J Biogeogr.*, 43, 2027–
568 2039, doi: 10.1111/jbi.12806, 2016b.
- 569 Chen, Z., Wang, Y., Wang, S., and Zhou, X.: Preliminary studies on the classification of grassland ecosystem
570 in China, *Acta Agrestia Sinica*, 10, 81–86, 2002. (in Chinese with English abstract)
- 571 Colman, B. P. and Schimel, J. P.: Drivers of microbial respiration and net N mineralization at the continental
572 scale, *Soil Biol. Biochem.*, 60, 65–76, doi: 10.1016/j.soilbio.2013.01.003, 2013.

573 Davidson, E. A., Belk, E., Boone, R. D.: Soil water content and temperature as independent or confounded
574 factors controlling soil respiration in a temperate mixed hardwood forest, *Glob. Change Biol.*, 4, 217–227,
575 doi: 10.1046/j.1365-2486.1998.00128.x, 1998.

576 Davidson, E. A. and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to
577 climate change, *Nature*, 440, 165–173, doi: 10.1038/nature04514, 2006.

578 Davidson, E. A., Janssens, I. A., and Luo, Y. Q.: On the variability of respiration in terrestrial ecosystems:
579 moving beyond Q_{10} , *Glob. Change Biol.*, 12, 154–164, doi: 10.1111/j.1365-2486.2005.01065.x , 2006.

580 Davidson, E. A., Savage, K., Verchot, L. V., and Navarro, R.: Minimizing artifacts and biases in chamber-
581 based measurements of soil respiration, *Agr. Forest Meteorol.*, 113, 21–37, doi: 10.1016/S0168-
582 1923(02)00100-4, 2002.

583 Ding, J., Chen, L., Zhang, B., Liu, L., Yang, G., Fang, K., Chen, Y., Li, F., Kou, D., Ji, C., Luo, Y., and Yang,
584 Y.: Linking temperature sensitivity of soil CO₂ release to substrate, environmental, and microbial
585 properties across alpine ecosystems, *Global Biogeochem. Cycles*, 30, 1310–1323, doi:
586 10.1002/2015GB005333, 2016.

587 Frank, A. B. and Dugas, W. A.: Carbon dioxide fluxes over a northern, semiarid, mixed-grass prairie, *Agr.*
588 *Forest Meteorol.*, 108, 317–326, doi: 10.1016/S0168-1923(01)00238-6, 2001.

589 Gershenson, A., Bader, N. E., and Cheng, W.: Effects of substrate availability on the temperature sensitivity
590 of soil organic matter decomposition, *Glob. Change Biol.*, 15, 176–183, doi: 10.1111/j.1365-
591 2486.2008.01827.x, 2009.

592 Graf, A., Weihermueller, L., Huisman, J. A., Herbst, M., Bauer, J., and Vereecken, H.: Measurement depth
593 effects on the apparent temperature sensitivity of soil respiration in field studies, *Biogeosciences*, 5, 1175–
594 1188, doi: 10.5194/bg-5-1175-2008, 2008.

595 Hibbard, K. A., Law, B. E., Reichstein, M. and Sulzman, J.: An analysis of soil respiration across northern
596 hemisphere temperate ecosystems, *Biogeochemistry*, 73, 29–70, doi: 10.1007/s10533-004-2946-0, 2005.

597 Hursh, A., Ballantyne, A., Cooper, L., Maneta, M., Kimball, J., and Watts, J.: The sensitivity of soil respiration
598 to soil temperature, moisture, and carbon supply at the global scale. *Glob. Change Biol.*, 23, 2090–2103,
599 doi: 10.1111/gcb.13489, 2017.

600 Jia, B., Zhou, G., Wang, Y., Wang, F., and Wang, X.: Effects of temperature and soil water-content on soil
601 respiration of grazed and ungrazed *Leymus chinensis* steppes, Inner Mongolia. *J. Arid Environ.*, 67, 60–
602 76, doi: 10.1016/j.jaridenv.2006.02.002, 2006.

603 Kirschbaum, M. U. F.: Will changes in soil organic carbon act as a positive or negative feedback on global
604 warming? *Biogeochemistry*, 48, 21–51, doi: 10.1023/A:1006238902976, 2000.

605 Kuzyakov, Y.: Sources of CO₂ efflux from soil and review of partitioning methods, *Soil Biol. Biochem.*, 38,
606 425–448, doi: 10.1016/j.soilbio.2005.08.020, 2006.

607 Lenton, T. M. and Huntingford, C.: Global terrestrial carbon storage and uncertainties in its temperature
608 sensitivity examined with a simple model, *Glob. Change Biol.*, 9, 1333–1352, doi: 10.1046/j.1365-
609 2486.2003.00674.x, 2003.

610 Li, Z., Wang, X., Zhang, R., Zhang, J., and Tian, C.: Contrasting diurnal variations in soil organic carbon
611 decomposition and root respiration due to a hysteresis effect with soil temperature in a *Gossypium s.*
612 (cotton) plantation, *Plant Soil*, 343, 347–355, doi: 10.1007/s11104-011-0722-1, 2011.

613 Liu, X., Zhang, W., Zhang, B., Yang, Q., Chang, J., and Hou, K.: Diurnal variation in soil respiration under
614 different land uses on Taihang Mountain, North China, *Atmos. Environ.*, 125, 283–292, doi:
615 10.1016/j.atmosenv.2015.11.034, 2016.

616 Luo, Y. and Zhou, X.: *Soil respiration and the environment*, Academic Press/Elsevier, San Diego, USA, 257–
617 305 pp, 2006.

618 Ma, A., Yu, G., He, N., Wang, Q., and Peng, S.: Above- and below-ground biomass relationships in China's
619 grassland vegetation, *Quaternary Sci.*, 34, 769–776, doi: 10.3969/j.issn.1001-7410.2014.04.09, 2014. (in
620 Chinese with English abstract)

621 Ni, J.: Carbon storage in grasslands of China, *J. Arid Environ.*, 50, 205–218, doi: 10.1006/jare.2201.0902,
622 2002.

623 Pavelka, M., Acosta, M., Marek, M. V., Kutsch, W., and Janous, D.: Dependence of the Q_{10} values on the
624 depth of the soil temperature measuring point, *Plant Soil*, 292, 171–179, doi: 10.1007/s11104-007-9213-
625 9, 2007.

626 Peng, F., You, Q., Xu, M., Guo, J., Wang, T., and Xue, X.: Effects of warming and clipping on ecosystem
627 carbon fluxes across two hydrologically contrasting years in an alpine meadow of the Qinghai-Tibet
628 Plateau, *Plos One*, 9, e109319, doi: 10.1371/journal.pone.0109319, 2014.

629 Peng, S., Piao, S., Wang, T., Sun, J., and Shen, Z.: Temperature sensitivity of soil respiration in different
630 ecosystems in China, *Soil Biol. Biochem.*, 41, 1008–1014, doi: 10.1016/j.soilbio.2008.10.023, 2009.

631 Pribyl, D. W.: A critical review of the conventional SOC to SOM conversion factor. *Geoderma*, 156, 75–83,
632 doi: 10.1016/j.geoderma.2010.02.003, 2010.

633 Raich, J. W. and Potter, C. S.: Global patterns of carbon dioxide emissions from soils, *Global Biogeochem.*
634 *Cycle*, 9, 23–36, doi: 10.1029/94GB02723, 1995.

635 Raich, J. W. and Schlesinger, W. H.: The global carbon dioxide flux in soil respiration and its relationship to
636 vegetation and climate, *Tellus*, 44, 81–99, doi: 10.1034/j.1600-0889.1992.t01-1-00001.x, 1992.

637 Reichstein, M., Subke, J. A., Angeli, A. C., and Tenhunen, J. D.: Does the temperature sensitivity of
638 decomposition of soil organic matter depend upon water content, soil horizon, or incubation time? *Glob.*
639 *Change Biol.*, 11, 1754–1767, doi: 10.1111/j.1365-2486.2005.001010.x, 2005.

640 Ryan, M. G. and Law, B. E.: Interpreting, measuring, and modeling soil respiration, *Biogeochemistry*, 73, 3–
641 27, doi: 10.1007/s10533-004-5167-7, 2005.

642 Schindlbacher, A., Zechmeister-Boltenstern, S., and Jandl, R.: Carbon losses due to soil warming: Do
643 autotrophic and heterotrophic soil respiration respond equally? *Glob. Change Biol.*, 15, 901–913, doi:
644 10.1111/j.1365-2486.2008.01757.x, 2009.

645 Shen, H., Zhu, Y., Zhao, X., Geng, X., Gao, S., and Fang, J.: Analysis of current grassland resources in China,
646 *Chin. Sci. Bull.*, 61, 139–154, doi: 10.1360/N972015-00732, 2016. (in Chinese with English abstract)

647 Song, X., Peng, C., Zhao, Z., Zhang, Z., Guo, B., Wang, W., Jiang, H., Zhu Q.: Quantification of soil
648 respiration in forest ecosystems across china. *Atmos. Environ.*, 94, 546–551, doi:
649 10.1016/j.atmosenv.2014.05.071, 2014.

650 Sims, P. L. and Bradford, J. A.: Carbon dioxide fluxes in a southern plains prairie, *Agr. Forest Meteorol.*, 109,
651 117–134, doi: 10.1016/S0168-1923(01)00264-7, 2001.

652 Subke, J. A. and Bahn, M.: On the ‘temperature sensitivity’ of soil respiration: Can we use the immeasurable

653 to predict the unknown? *Soil Biol. Biochem.*, 42, 1653-1656, doi: 10.1016/j.soilbio.2010.05.026, 2010.

654 Subke, J. A., Inglema, I. and Cotrufo, M. F.: Trends and methodological impacts in soil CO₂ efflux partitioning:
655 A metaanalytical review. *Glob. Change Biol.*, 12: 921–943, doi: 10.1111/j.1365-2486.2006.01117.x , 2006.

656 Turner, B. L.: Variation in pH optima of hydrolytic enzyme activities in tropical rain forest soils, *Appl. Environ.*
657 *Microb.*, 76, 6485–6493, doi: 10.1128/AEM.00560-10, 2010.

658 van's Hoff, J. H.: Lectures on Theoretical and Physical Chemistry. Part 1: Chemical Dynamics. Edward Arnold,
659 London, UK, 224–229 pp, 1898.

660 Wan, S., Norby, R. J., Ledford, J., and Weltzin, J. F.: Responses of soil respiration to elevated CO₂, air warming,
661 and changing soil water availability in a model old-field grassland, *Glob. Change Biol.*, 13, 2411–2424,
662 doi: 10.1111/j.1365-2486.2007.01433.x, 2007.

663 Wang, B., Zha, T. S., Jia, X., Wu, B., Zhang, Y. Q., and Qin, S. G.: Soil moisture modifies the response of soil
664 respiration to temperature in a desert shrub ecosystem, *Biogeosciences*, 11, 259–268, doi: 10.5194/bg-11-
665 259-2014, 2014a.

666 Wang, M., Liu, X., Zhang, J., Li, X., Wang, G., Li, X., and Chen, W.: Soil respiration associated with plant
667 succession at the meadow steppes in Songnen Plain, Northeast China, *J. Plant Ecol.*, 8, 51–60, doi:
668 10.1093/jpe/rtu006, 2015.

669 Wang, M., Liu, X., Zhang, J., Li, X., Wang, G., Li, X., and Lu, X.: Diurnal and seasonal dynamics of soil
670 respiration at temperate *Leymus chinensis* meadow steppes in western Songnen Plain, China, *Chin. Geogr.*
671 *Sci.*, 24, 287–296, doi: 10.1007/s11769-014-0682-5, 2014b.

672 Wang, W. and Fang, J.: Soil respiration and human effects on global grasslands, *Global Planet. Change*, 67,
673 20–28, doi: 10.1016/j.gloplacha.2008.12.011, 2009.

674 Wang, X., Piao, S., Ciais, P., Janssens, I.A., Reichstein, M., Peng, S., and Wang, T.: Are ecological gradients
675 in seasonal Q_{10} of soil respiration explained by climate or by vegetation seasonality? *Soil Biol. Biochem.*,
676 42, 1728–1734, doi: 10.1016/j.soilbio.2010.06.008, 2010.

677 Wang, Z., Ji, L., Hou, X., and Schellenberg, M. P.: Soil respiration in semiarid temperate grasslands under
678 various land management, *Plos One*, 11, e0147987, doi: 10.1371/journal.pone.0147987, 2016.

679 Xu, W., Li, X., Liu, W., Li, L., Hou, L., Shi, H., Xia, J., Liu, D., Zhang, H., Chen, Y., Cai, W., Fu, Y., and Yuan,
680 W.: Spatial patterns of soil and ecosystem respiration regulated by biological and environmental variables
681 along a precipitation gradient in semi-arid grasslands in China, *Ecol. Res.*, 31, 505–513, doi:
682 10.1007/s11284-016-1355-x, 2016.

683 Xu, Z., Tang, S., Xiong, L., Yang, W., Yin, H., Tu, L., Wu, F., Chen, L., and Tan, B.: Temperature sensitivity
684 of soil respiration in china's forest ecosystems: patterns and controls, *Appl. Soil Ecol.*, 93, 105–110, doi:
685 10.1016/j.apsoil.2015.04.008, 2015.

686 Yim, M. H., Joo, S. J., and Nakane, K.: Comparison of field methods for measuring soil respiration: a static
687 alkali absorption method and two dynamic closed chamber methods, *Forest Ecol. Manage.*, 170, 189–197,
688 doi: 10.1016/S0378-1127(01)00773-3, 2002.

689 Yuste, J. C., Janssens, I. A., Carrara, A., and Ceulemans, R.: Annual Q_{10} of soil respiration reflects plant
690 phenological patterns as well as temperature sensitivity, *Glob. Change Biol.*, 10, 161-169, 10.1111/j.1529-
691 8817.2003.00727.x, 2004.

692 Zhang, H., Fan, J., and Shao, Q.: Land use/land cover change in the grassland restoration program areas in

693 China, 2000–2010, *Prog. Geogr.* 34(7), 840–853, doi: 10.18306/dlkxjz.2015.07.006, 2015a. (in Chinese
694 with English abstract)

695 Zhang, Y., Guo, S., Liu, Q., Jiang, J., Wang, R., and Li, N.: Responses of soil respiration to land use
696 conversions in degraded ecosystem of the semi-arid Loess Plateau, *Ecol. Eng.*, 74, 196–205, doi:
697 10.1016/j.ecoleng.2014.10.003, 2015b.

698 Zhao, J., Li, R., Li, X. and Tian, L.: Environmental controls on soil respiration in alpine meadow along a large
699 altitudinal gradient on the central Tibetan Plateau. *Catena*, 159, 84–92, doi: 10.1016/j.catena.2017.08.007,
700 2017.

701 Zhou, T., Shi, P., Hui, D., and Luo, Y.: Global pattern of temperature sensitivity of soil heterotrophic respiration
702 (Q_{10}) and its implications for carbon-climate feedback. *J. Geophys. Res-Bioge.*, 114, 271–274, doi:
703 10.1029/2008JG000850, 2009.

704 Zhu, B. and Cheng, W. X.: Rhizosphere priming effect increases the temperature sensitivity of soil organic
705 matter decomposition. *Glob. Change Biol.*, 17, 2172–2183, doi: 10.1111/j.1365-2486.2010.02354.x, 2011.

706 **Table 1** The annual soil respiration (Rs) and the proportions of growing season, non-
 707 growing season Rs to annual Rs in different grassland ecosystems across China. There was
 708 no sample for annual Rs in temperate desert steppe, so the data were not presented in this
 709 table. The different lowercase letters in each column indicate the significant difference at p
 710 = 0.05, and different uppercase letters indicate the significant difference between growing
 711 and non-growing season at $p = 0.001$. n represents the number of samples.


| Grassland types | n | Annual Rs (g C m ⁻² yr ⁻¹) | | | Rs proportion (%) | |
|--------------------------|-----|---------------------------------------------------|-------|--------|-------------------|--------------------|
| | | Mean ±SE | Min. | Max. | Growing season | Non-growing season |
| Temperate typical steppe | 16 | 371.3±94.8 a | 122.9 | 1670.0 | 79.6±2.9 a | 20.4±2.9 a |
| Temperate meadow steppe | 6 | 442.1±83.4 a | 218.8 | 784.7 | 86.8±2.7 a | 13.2±2.7 a |
| Alpine grassland | 20 | 581.5±62.3 a | 246.3 | 1161.1 | 77.3±2.5 a | 22.7±2.5 a |
| Warm-tropical grassland | 12 | 933.6±161.8 b | 428.8 | 2407.1 | 76.2±2.5 a | 23.8±2.5 a |
| Total | 54 | 582.0±57.9 | 122.9 | 2407.1 | 78.7±1.5 A | 21.3±1.5 B |

712

713 **Table 2** Comparisons of annual soil respiration and Q_{10} between Chinese grasslands and
 714 other syntheses. The numbers in parentheses represent the number of samples.

| Scope | Annual Rs (g C m ⁻² yr ⁻¹) | Q_{10-ST5} | $Q_{10-ST10}$ | Reference source |
|--------------------------------|------------------------------------------------------|--------------|---------------|---------------------|
| Global terrestrial ecosystems | 910.0 (657) | | | Chen at al., 2010 |
| | 870.0 (1195) | | | Chen at al., 2014 |
| | 791.2 (1741) | | | Hursh et al., 2017 |
| | | 2.40 (77) | 3.10 (46) | Wang et al., 2010 |
| Global grasslands | 448.9 (46) | 2.13 (41) | | Wang and Fang, 2009 |
| | 745.0 (179) | | | Chen at al., 2010 |
| | 840.0 (113) | | | Chen at al., 2014 |
| | 599.1 (163) | | | Hursh et al., 2017 |
| Chinese terrestrial ecosystems | | 2.03 (64) | 2.61 (33) | Peng et al., 2009 |
| Chinese forests | 919.7 (139) | 2.46 (107) | | Song et al., 2014 |
| | | 2.51 (145) | | Xu et al., 2015 |
| Chinese grasslands | 582.0 (54) | 2.80 (73) | 2.56 (59) | This study |


715



716

717 **Figure 1.** The site location of soil respiration studies selected in this study across Chinese
 718 grasslands.


719



720

721 **Figure 2.** Relationships of annual soil respiration (Rs) with abiotic and biotic factors. The
 722 dash lines represent the 95% confidence interval. When p value was greater than 0.05,
 723 the regression lines were not drawn.


724



725

726 **Figure 3.** Histogram plots for Q_{10} values (a-e) and the coefficient of determination for
 727 Q_{10} (R_{Q^2} , f-j) across Chinese grasslands. (a) and (f): soil surface temperature; (b) and (g):
 728 soil temperature at the depth of 5 cm; (c) and (h): soil temperature at the depth of 10 cm;
 729 (d) and (i): soil temperature at the depth of 15 cm; (e) and (j): soil temperature at the depth
 730 of 20 cm. n represents the number of samples.


731



732

733 **Figure 4.** Comparisons of Q_{10} values among soil temperature measurement depths (a)
 734 and among grassland types (b, c). (a) Q_{10} values derived by soil temperature at the depth
 735 of 0, 5, 10, 15, and 20 cm, respectively. (b) Q_{10} values derived by soil temperature at the
 736 depth of 5 cm. (c) Q_{10} values derived by soil temperature at the depth of 10 cm. TTS,
 737 TMS, TDS, ALG, and WTG represent temperate typical steppe, temperate meadow
 738 steppe, temperate desert steppe, alpine grassland, and warm-tropical grassland,
 739 respectively. Error bars in (a) represent standard errors. In the box plot (b and c), the “+”
 740 represent mean values, horizontal lines inside box represent medians, box ends represent
 741 the 25th and the 75th quartiles, vertical lines represent 2.5th and 97.5th percentiles,
 742 hollow circles represent outliers, and n represents the number of samples. Different
 743 lowercase letters indicate significant differences among soil depths or grassland types at
 744 $p = 0.05$.

745



746

747 **Figure 5.** Relationships of temperature sensitivity of soil respiration (Q_{10}) with abiotic
 748 and biotic factors. The black and gray points represent Q_{10} derived by soil temperature at
 749 the depth of 5 and 10 cm, respectively, and the black and gray lines represent their
 750 corresponding relationships with environmental factors. When p value was greater than
 751 0.05, the regression lines were not drawn.

752