

# Soil total phosphorus and nitrogen explain vegetation community composition in a northern forest ecosystem near a phosphate massif

Laura Matkala<sup>1</sup>, Majja Salemaa<sup>2</sup>, Jaana Bäck<sup>1</sup>

<sup>1</sup>Institute for Atmospheric and Earth System Research / Forest Sciences, Faculty of Agriculture and Forestry, PO Box 27,  
5 00014, University of Helsinki, Helsinki, Finland

<sup>2</sup> Natural Resources Institute Finland (Luke), Latokartanonkaari 9, 00790 Helsinki, Finland

*Correspondence to:* Laura Matkala (laura.matkala@helsinki.fi)

**Abstract.** We studied how the dominant tree species, tree age, rock parent material and soil layer affect soil nutrient contents, and how the community composition of forest vegetation relates to soil nutrient contents near Sokli phosphate ore in northern  
10 Finland. For this purpose, we established 16 study plots on different distances from the phosphate ore along four transects. Phosphate mining may take place in Sokli in the future, and the vegetation surveys and soil sampling that we conducted at the plots can be used as a baseline status for following the possible changes that the mining may cause to the surrounding ecosystem. We found, that the total phosphorus (P) and nitrogen (N) contents of the soil humus layer were positively related with species number and abundance of the understorey vegetation, and the correlation was slightly higher with P than N. This  
15 is interesting as usually N has the most important growth-limiting role in boreal ecosystems. The spatial variation in the content of soil elements was high both between and within plots, emphasizing the heterogeneity of the soil. Dominant tree species and the soil layer were the most important environmental variables affecting soil nutrient content. High contents of P in the humus layer (maximum 2.60 g kg<sup>-1</sup>) were measured from the birch-dominated plots. As the P contents of birch leaves and leaf litter were also rather high (2.58 g kg<sup>-1</sup> and 1.28 g kg<sup>-1</sup>, respectively), this may imply that the leaf litter of birch forms an important  
20 source of P for the soil.

## 1 Introduction

Climate and availability of soil nutrients are important factors controlling the species composition of tree stand and understorey vegetation in boreal forests (Cajander 1909, 1949, Kuusipalo 1985, Økland and Eilertsen 1996). High latitude forest ecosystems are characteristically cold, have a short growing season, and are nutrient poor. Organic matter decomposition and  
25 nutrient release are usually slow in cold climates (Hobbie et al. 2002). The edaphic conditions are reflected in the growth and chemical composition of plant species, as well as in species composition of vegetation (Vinton and Burke 1995, Salemaa et al. 2008). In addition, tree cover affects the species composition and abundance in the understorey by shading (Verheven et al. 2012, Tonteri et al. 2016) and regulating nutrient input in throughfall precipitation (Salemaa et al. 2019) and litterfall (Ukonmaanaho et al. 2008).

30

Nitrogen (N) and phosphorus (P) are generally the main growth-limiting nutrients for plants (Koerselman and Meuleman 1996). Boreal forests are mostly N-limited (Tamm 1991), and fertilization with N usually speeds up forest growth (Saarsalmi and Mälikönen 2001). Nitrogen is bound in organic material, and only a little is directly available for plants as inorganic ammonium ( $\text{NH}_4^+$ ) and nitrate ( $\text{NO}_3^-$ ) (Marschner 1995) or as organic forms like amino acids (Näsholm et al. 2008 and references within). Phosphorus deficiency occurs in temperate and tropical forest ecosystems, but P is rarely a limiting factor in boreal forests on mineral soil (Augusto et al. 2017). However, P can be growth limiting on boreal peatlands (Moilanen et al. 2010, Brække and Salih 2002). The ratio of soil N to soil P is significant for forest growth on a global scale (Augusto et al. 2017). Hedwall et al. (2017) found that the species richness of vascular plants in a temperate forest doubled with combined NP fertilization in southern Sweden but not when either of the nutrients was added alone. This positive effect was strongest in grass species. In boreal N-limited forests, the number of vascular plant species (grasses and forbs) increased with increasing N concentration of the organic layer (Salemaa et al. 2008). Hofmeister et al. (2009) noticed that in a temperate forest the species richness of the herb layer was higher in P-rich than P-poor soils but only if strong N limitation occurred simultaneously in the P-rich soils. However, in many regions where humans have enhanced atmospheric N deposition the number of plant species has decreased (Dirnböck et al. 2014). For instance, high soil N was related to decreased herb layer species richness in deciduous forests in Sweden (Dupré et al. 2002).

In this study, we analysed whether plant species composition and nutrient levels of tree leaves indicate soil total N and P in a northern boreal (Hämet-Ahti 1981) research site in Sokli, Finland. At this site, the soil contains naturally large variations in P content. In Sokli, there is a large deposit of phosphate rock, a carbonatite complex mainly consisting of apatite ( $\text{Ca}_5(\text{PO}_4)_3\text{F}$ ), which was discovered by the Mining and Steel Company of Rautaruukki Oy in 1967 (Vartiainen and Paarma 1979). Plans to open a phosphate mine in Sokli have been on display for decades and will possibly be realized in the future. The vegetation at the carbonatite complex differs from that of the typical forests of the region (Talvitie 1979, Pöyry Environment 2009). Downy birch (*Betula pubescens*) is dominating and often the single tree species, whereas the more typical forests of the region are dominated by Scots pine (*Pinus sylvestris*) or Norway spruce (*Picea abies*). Understorey vegetation at Sokli is slightly richer in herb and grass species compared with surrounding forests, where dwarf shrubs, bryophytes and lichen dominate the understorey. However, vegetation similar to that in Sokli can be found as patches elsewhere in the region.

The general aim of this study was to determine the undisturbed baseline status of the forest ecosystem in terms of soil, understorey vegetation and tree layers in the Sokli area in case there is a need to monitor the effects of phosphate mining. Phosphate mining can cause, for instance, aerial deposition of heavy metals and phosphate onto the surroundings of the mine (Reta et al. 2018), which can lead to changes in the abundance and species composition of the understorey. Vegetation, soil and foliage chemistry surveys provide data on the current state of the ecosystem (from the year 2015) that can be used as a reference level for the changes. Our specific aim was to identify which factors in the soil and tree layer explain the composition

and abundance of plant species. In addition, we studied which environmental variables could explain soil nutrient contents, especially total P content.

We hypothesize that there are positive relationships between

- 5 a) N and P contents of the soil humus layer and the abundance and species composition of the understory vegetation
- b) N and P contents in the topmost soil layers and the N and P contents of needle and leaf biomass
- c) N and P contents in the topmost soil layers and the occurrence of birch trees in the research plots

## 2 Material and methods

### 2.1 Site description

10 We established 16 study plots along four transects (A–D) around the planned Sokli mining district (67° 48′ N, 29° 16′ E) in Savukoski, eastern Lapland in 2014 and 2015 (Fig. 1). The plots were located different distances from the phosphate ore in four transects, enabling evaluation of the possible effects of the mine in the future. No plots were located inside the mining district, as accessing and doing research at the mining district would have required a permit from the mining company. The carbonatite massif of Sokli belongs to the Devonian Kola Alkaline Province (Tuovinen et al. 2015). Nine of the plots were  
15 located in Natura 2000 conservation areas. Plots A4, A5 and A6 were in Värriö, A1 and A2 in Yli-Nuortti, B1, B2 and B3 in Törmäoja and D5 in the UK-puisto – Sompio – Kemihaara Natura area. In terms of topography, Törmäoja is a valley, reminiscent of the form of a kettle (*kattilalaakso* in Finnish) (NATURA 2000 - Standard Data Forms FI1301512 and FI1301513). The mid-parts of the valley are treeless, or the trees are at sapling stage, because cold winds blowing through the valley kill the new buds in the spring. Thus, our plots at Törmäoja were on the edge of the less steep western part, where some  
20 mature trees grow. The majority of the plots had a mixed composition of at least two tree species, while in some plots the tree cover consisted of only one species. An additional factor affecting vegetation cover and species composition at our research site is reindeer herding. Since all the plots were located in areas where reindeer roam freely, we assume that the pressure caused by grazing and trampling is equal in all plots. The plots of this study together with SMEAR 1 (Station for Measuring Ecosystem-Atmosphere Relations) at the Värriö Subarctic Research Station (67° 46′ N, 29° 35′ E) (Hari et al. 1994) serve as  
25 a gradient type network for monitoring the current status and the possible, mining-induced, changes of the environment in the future.

Meteorological parameters from the years of data collection and for the climatological normal period of 1980–2010 are presented in Table 1. The wind blows almost equally from the south-west and north-east during spring and summer, whereas  
30 in winter and autumn the prevailing wind direction is south-west (Ruuskanen et al. 2003). The growing season, when daily average temperature exceeds 5 °C, lasts from June to September. Soils are haplic Podzols with sandy tills (FAO 1988).

## 2.2 Plot setup and vegetation characterization

The distance between two plots depended on the topography and existing roads, but generally it was about 2 km. A plot consisted of four clusters, each including three square-shaped subplots sized 1 m<sup>2</sup> for observations and sampling (Fig. 2). The size of the whole plot was 30 × 30m. We recorded all tree species growing on the plots, measured their heights and diameter at breast height (dbh) (equivalent to a height of 1.3 m) (Table 2). Stem volumes were estimated using the equations of Laasasenaho (1982). We estimated tree age by measuring dbh and examining the existing approximated tree age from plot A6 at SMEAR1, where the mature trees are about 70 years old. We considered trees with dbh 1–9.9 cm to be young, 10–14.9 cm as middle-aged and >15 cm as old. We visually assessed the cover (% surface area) and counted the number of plant species in the understorey vegetation in all 12 subplots per plot in the summers of 2014 and 2015 (Appendix A). We used a 1 m<sup>2</sup> square frame to delineate the subplot (Salemaa et al. 1999). All species in the bottom layer (bryophytes and lichens) and field layer (dwarf shrubs, tree seedlings, grasses, sedges and forbs, height < 50 cm) were included.

### 2.2.1 Sampling of soil

Altogether, 256 soil samples were collected from sixteen plots using a soil corer (inner diameter 5 cm) in June 2015. The soil was sampled within a 1m distance from the subplots (cf. Liski 1995). The soil cores were separated by visual criteria into four soil horizons: the top layer, which is a mixture of litter and decomposing organic layer (F), the humus layer (O), the eluvial layer (A), and the illuvial layer (B) (cf. Köster et al. 2014). The rocky soil and shallow humus layer made it impossible to sample the mineral soil layers in some clusters. The soil samples from each horizon were combined into composite samples in each cluster in the field. The composite samples were air-dried except for the organic F and O horizons, which were dried at 60 °C for 48 hours. Dried mineral soils were sieved with a 2 mm sieve and the F and O horizons were milled before storing in a dry place for further analyses.

### 2.2.2 Sampling of needles and leaves

Five pines and five spruces per plot were chosen for needle sampling in September 2015, when the needle growth had ended. If less than five trees per species were present, all of them were chosen. Three branches (length approximately 50 cm) were taken from the upper third of the canopy using a branch saw. We took only second order branches because cutting of first order branches would have been too destructive to the trees (cf. Helmisaari 1990). Needle age classes (C = current year, C+1 = one-year-old needles, C+2 = two-year-old needles) were separated from each branch and dried at 65 °C for 48 hours, milled and stored in a dry place for further analyses. The samples were combined so that there was one C, one C+1 and one C+2 composite needle sample per tree.

We sampled green birch leaves in July and leaf litter in September 2015. Approximately 10 green leaves from 10 different trees were picked and combined (Rautio et al. 2010). Only mature, undamaged leaves were chosen. Birch litter was collected

under the same tree canopies from which the green leaves had been taken and in approximately the same number as the green leaf samples. We aimed to take litter leaves shed in the current year, so that they were decomposed as little as possible. Green and litter leaves were dried at 65 °C for 48 h and manually cleaned of extra material, such as soil particles and needles. The needles and the few soil particles attached on the litter leaves were removed with tweezers. The green leaves did not need cleaning. The litter leaves were also rather clean, as it had rained at the time of sampling. After cleaning, the leaves were milled and stored in a dry place for further analyses. Needles and leaves were sampled at a different time than the soil. Both needle (e.g. Helmisaari 1990) and soil nutrient contents vary between the seasons. However, as all soil and all needle sampling was conducted at the same time of the season, the comparison between the plots was not hindered.

### 2.3 Laboratory analyses

10 Total element contents of potassium (K) and P were analysed from soil and foliar samples by inductively coupled plasma optical emission spectrometry. For this analysis, the samples were first wet combusted. One gram of mineral soil samples and 0.3 g of organic samples were combusted with 1 ml of H<sub>2</sub>O<sub>2</sub> and 10 ml HNO<sub>3</sub>, and heated in a microwave oven. The samples were then filtered with Whatman Grade 589/3 filter paper and stored in plastic bottles in a cooler until analysed.

15 Total carbon (C) and N were analysed directly from dried and milled foliar samples as well as from the F and O soil layers. Samples of 2–3 mg were measured and analysed with an Element Analyser, which uses a high temperature combustion method with subsequent gas analysis of CN (VarioMax, Elementar Analysensysteme GmbH, Germany). Soil pH was measured from two O layer samples per plot and their average value used. 20 mg of dried sample was mixed together with ultrapure water (50ml). The suspension was covered and left standing for 24 hours, and pH was measured with a glass electrode.

### 20 2.4 Statistical analyses

We used one-way analysis of variance (ANOVA) and Tukey's honestly significant difference post-hoc test for analysing the plot-wise differences in needle element contents. Plot-averages of needle elemental contents were calculated across all needle age classes and both conifer species. One-way ANOVA was also used in analysing differences between the needle age classes. We grouped the plots based on their dominant tree species into pine, birch and spruce plots and calculated the average soil nutrient contents in each horizon in these plots. We then compared the nutrient contents in each soil horizon with one-way ANOVA.

We tested the effects of environmental variables on soil total P and N contents and C:N with linear mixed-effect models. We used dominant tree species, estimated age class, rock parent material (Fig. 1) and soil horizon as fixed effects and plot as random effect. Soil total P needed to be log-transformed, while for N and C:N the visual inspection of residual plots (Fig. B1) did not reveal obvious deviations from homoscedasticity or normality. We obtained *p*-values for the fixed effects by likelihood ratio tests, where the full model with all the fixed effects was tested against a model where each fixed effect was removed in

turn. We used package lme4 (Bates et al. 2015) in R programme 3.4.3 (R Development Core Team 2017) for building the models. Pseudo  $R^2$ -values for the models were calculated by using package r2glmm (Jaeger 2017). The models took the form:

$$SC_{P,N,CN} = B_0 + B_{dt} + B_{ta} + B_g + B_h + \epsilon, \quad (1)$$

5

where  $SC_{P,N,CN}$  is the soil nutrient content (total P, N or C:N ratio),  $B_0$  denotes a fixed intercept parameter,  $B_{dt}$  denotes the fixed unknown parameters associated with the dominant tree species,  $B_{ta}$  denotes the fixed unknown parameters associated with the age of the dominant tree species,  $B_g$  denotes the fixed unknown parameters associated with the rock parent material, and  $B_h$  denotes the fixed unknown parameters associated with the soil horizon. The random effect  $\epsilon$  is assumed to take the

10 form:

$$\epsilon = \alpha_p + u, \quad (2)$$

where  $\alpha_p$  denotes the random parameters related to the research plot and  $u$  is an unobservable error term. Random effect parameters and the random error term are assumed to follow normal distributions  $\alpha_p \sim N(0, \sigma_p^2)$  and  $u \sim N(0, \sigma_u^2)$ .

15

We calculated plot-wise averages from the percentage covers of the plant species in the subplots. We ordinated this vegetation data by global non-metric multidimensional scaling (Minchin 1987) using the Vegan package (Oksanen et al. 2018) in R programme 3.4.3 (R Development Core Team 2017). Ordination pattern of the plots based on the Bray-Curtis dissimilarity indices in floristic composition was analysed to find the main environmental gradients behind the vegetation variation. We analysed the data in three-dimensional space but present the results in 1 vs 2 and 1 vs 3 dimensions (the results in 2 vs 3 dimensions did not give any new information). We then fit the plot-wise data of soil elements (O horizon), needle element contents, volume of birch (% of total tree volume), species cover (% of the surface area) as well as plot distance from the phosphate ore as linear vectors to the ordination pattern of the plots. The correlation between the environmental variables and the ordination was calculated by a linear vector procedure (envfit in Vegan). The total P in the soil O horizon was also fitted as a smooth surface to the ordination pattern in order to analyse the form of the relationship (linear or non-linear). The fit was done by a generalized additive model (Gaussian distribution error).

25

### 3 Results

#### 3.1 Soil element contents

The outlying points in Fig 3. as well as the high standard deviations of P showed rather high variation between and within plots (Table B1, Fig. B2). Birch-dominated plots had the highest P and N contents and lowest C:N ratio compared with coniferous plots in all soil layers where the elements were measured (Figs. 3 and 4), and these differences were mostly

30

statistically significant (Table B2). We found no statistical evidence for differences in soil N:P ratio or total C content based on the dominant tree species of the plots. There was only one spruce-dominated plot and thus only four soil samples from each soil horizon from the spruce plot, which may have affected these results. In general, topsoil had the highest P content, but in many plots deeper soil layers also had high P content. Certain plots (A1, A3, B1, D2) had clearly different P and N contents and C:N ratio than most other plots. The N contents and C:N ratios were in most cases higher in F than O horizon. The majority of the plots had higher N:P ratio in the F horizon than O horizon.

### 3.2 Needle and leaf element contents

Needle P contents were highest in the C needles and significantly different from other age classes in both pine and spruce (Table C1). Against our expectations, the needle P contents of both conifer species were rather similar across plots (Table C2). On the other hand, N and C contents, as well as the C:N ratio of the conifers showed some between-plot variation ( $p < 0.05$ ), but no significant variation was found in the foliar N:P ratio in either species. Spruce had slightly higher needle P contents than pine in all age classes, whereas N contents were higher in pine than in spruce needles (Table 3). Birch had higher P contents of green leaves than the conifers. Also leaf litter of birch had quite high P contents, and in general litter leaves showed more variation in element contents than the green ones. Nitrogen contents were lower in birch leaf litter than in green leaves, but the contents of C increased slightly from green leaves to litter. However, no differences between the plots were detected in either of the elements. According to the correlation matrix between the elements in the soil O horizon, tree leaves and needles, and number of species in the understory (Fig. 5) birch K (green leaves) correlated with soil K and pH, birch litter N with soil N:P and birch litter K with soil N. Otherwise no significant correlations between foliar element contents and soil element contents were found. The number of mosses and lichen correlated negatively with soil total C ( $p < 0.01$ ) and C:N ( $p < 0.05$ ), the number of sedges and grasses positively with soil total P ( $p < 0.001$ ) and pH ( $p < 0.01$ ) and negatively with soil C:N ( $p < 0.01$ ) and N:P ( $p < 0.001$ ) and the number of dwarf shrubs and trees positively with soil total K ( $p < 0.05$ ). No significant correlations were found between number of species and foliar element contents. The cover (% of surface) of grass and sedge species correlated positively with soil P ( $p < 0.001$ ) and number of sedges and grasses ( $p < 0.05$ ), while the cover of dwarf shrubs and trees correlated positively with the P content of green birch leaves ( $p < 0.05$ ).

### 3.3 Mixed-effect model results

We used linear mixed-effect models for determining which environmental factors can best explain soil total P and N contents and the C:N ratio. The dominant tree species and soil horizon explained 45 % of the total P of soil, and the soil horizon explained 20 % of the total N of soil as well as the C:N ratio of soil (Table 5). The other tested fixed effects had  $p$ -values  $> 0.05$  and were for that reason excluded from the models. The highest estimates of P were produced with birch as the dominant tree species and F as the soil layer, and the highest estimates of N with F as the soil layer in the final models.

### 3.4 Ordination analysis of understorey vegetation

The closer the plots were to each other in the ordination space, the more similar their vegetation was (Fig. 6a,b). Plots positioned more on the left-hand side in the figure had a higher number of forbs and grasses growing on them than the plots positioned on the right-hand side in the figure (Fig. 6a,b). Species such as *Calamagrostis epigejos*, *Carex* spp., *Rubus arcticus* and *Luzula pilosa* had relatively high coverage in the plots on the left. The plots further on the right in the figure had more species which tolerate poor and dry growing conditions, such as *Cladonia* and *Cladina* lichens. Also the tree species changed from right to left, as the plots on the right were dominated by pine, whereas furthest on the left in plot D2 birch was the only tree species present. In general, the fertility trend in the vegetation followed the first dimension, while the moisture gradient followed the second dimension. Moisture demanding species, such as *Equisetum sylvaticum* and *Rhododendron tomentosum* are located in the upper part of Fig. 6b, and those tolerating drier conditions, such as *Peltigera rufescens* and *Stereocaulon tomentosum*, are located in the lower part of the ordination space in Fig. 6b. Another moisture gradient, expressing specific paludified conditions, seemed to follow the third dimension. Peatland species like *Sphagnum angustifolium* and *Aulacomnium palustre* are located in the upper part, and species preferring dry conditions, such as *Cetraria islandica*, are in the lower part of Fig 6d. Considering all three dimensions of ordination space, the generalist species, such as *Polytrichum commune*, *Pleurozium schreberi* and *Vaccinium myrtillus* are located in the middle.

The vector arrows fitted to the ordination space (Fig. 7c,d) depict the maximum correlations between environmental variables and plot ordination. The length of an arrow indicates the magnitude and direction of the polarity (plus–minus) of the correlation. The highest correlations occurred between the plot-wise average P content of the soil O horizon and the ordination pattern of the plots. The isocline gradient of soil P in relation to the ordination pattern was almost linear (Fig. 7a). Vectors of soil pH, N and P content all increased towards the more fertile plots, but the vectors of soil C:N and N:P went to the opposite directions (Fig. 7b) indicating poor soil conditions. The average total number of grass, forb and sedge species as well as their coverage in the study plots also increased towards the more fertile plots (Fig. 7d).

### 4 Discussion

All the plant species growing in the study plots were common forest species in Finland (e.g., Reinikainen et al. 2000, Finnish Biodiversity Information Facility <https://laji.fi/en>) (Appendix A). However, in some plots the structure and abundance of species in the understorey clearly differed from the surrounding, more typical northern boreal forests. We found evidence that the number of species in the group of grasses and sedges as well as the cover (% of surface) of the same plant group had a higher positive correlation with humus P content than N content (Fig. 5). However, both of these nutrients were important factors explaining the vegetation composition in the ordination configuration (Fig. 7), which supports our first hypothesis. We also found that the humus C:N ratio correlated negatively with the abundance and species composition in the understorey. Also, Salemaa et al. (2008) have observed that total N and the C:N ratio of the humus layer explained most large-scale



vegetation variation, across several forest sites in Finland. They also measured extractable soil P, which seemed to have more power in explaining vegetation patterns in northern than southern Finland. Soil P availability was one of the key factors in plant community variation in alpine habitats in Troms, northern Norway (Arnesen et al. 2007), where a higher variety of lichen species and the frequency of occurrence of *Salix herbacea* and certain sedge and grass species were explained by higher availability of P in soil. We conclude that the possible aerial deposition of phosphate from the mine in Sokli could lead to changes in plant species composition and abundance if high amounts of P are deposited into the ecosystem surrounding the mining region.

Our second hypothesis stated that the N and P contents of the topmost soil layers correlate with the N and P contents of foliar biomass, but our results (Table B2) did not support this hypothesis. The reason could be that we measured total contents of N and P in soil instead of the plant available contents of these nutrients. The plant available contents of these nutrients might have given different results. Perhaps plant species composition in ground vegetation is sensitive to even small additions of available N and P in the upper soil layers where the roots occur, whereas higher contents of these elements are required to have any effect on the needles. The P and N levels of our needle samples were similar to those previously measured in Finland (Helmisaari 1990, Merilä and Derome 2008, Moilanen et al. 2013). The higher P contents of C needles compared with older needles is common for conifers and occurs because the dry weight in recently matured needles increases faster than the transportation of P to the needles (Helmisaari 1990). The N contents of both green birch leaves and leaf litter agreed with those reported by Ferm and Markkola (1985). The P contents of the green leaves were higher than they measured (approximately 2.0 g kg<sup>-1</sup>). Although the foliar N and P contents were not reflected in the uppermost soil layers, our results support the third hypothesis, and the occurrence of birch correlates positively with the N and P content of the top layers of soil (Table B2). The plots dominated by birch had significantly higher total P content in all but the B layer compared with plots dominated by the conifers (Figs. 3 and 4). Birch leaves were a major source of litter in the plots where soil P was high. These findings are supported by the study of Lukina et al. (2019), which found that the extractable P content of organic soil layers was significantly higher in birch- and spruce-dominated forest sites than in sites dominated by pine in North-Western Russia. Viro (1955) found that the leaf litter of birch had remarkably high P content compared with other Finnish tree species. Our litter P contents were near the approximate 1.50 g kg<sup>-1</sup> that Ferm and Markkola (1985) measured from a 40-year-old forest but much less than those reported from younger forests. In a litter experiment in Abisko (northern Sweden), the addition of birch litter increased both the total P (Sorensen and Michelsen 2011) and the available P (Rinnan et al. 2008) contents in the organic soil layer in those subarctic heaths, where *Hylocomium splendens* dominated the moss layer. These results imply that birch is an important factor in recycling and providing P to the soil in certain types of northern forest sites.

In general, the spatial variation in soil element contents between plots was high, emphasizing the heterogeneity of soil fertility level (Figs. A1 and A2). As our results showed, this heterogeneity can partially be explained by the dominant tree species of the research plot, which especially affects the topmost soil layers. According to the nutrient-uplifting hypothesis (Jobbágy and

Jackson 2004), trees and other vegetation can transport minerals such as P and K from the deep soil layers to the surface of soils. The P contents of soil samples (Table B1) in our study (1.80–2.60 g kg<sup>-1</sup> in the O horizon) fell mostly in the category we could expect based on the literature. The P content of the humus layer in southern Finnish forest soil has been observed to vary between 0.80 and 2.10 g kg<sup>-1</sup> (Mäkipää 1999), whereas different studies in northern Finland have found the P contents of  
5 0.39–3.00 g kg<sup>-1</sup> in the organic topsoil (Mikkola and Sepponen 1986, Reimann et al. 1997). Most of our plots had highest P content in the organic soil layers, implying that decaying plant parts were a major source of P added to the soil. Low Arctic soils tend to have organic P as the primary form of P (Weintraub 2011). The content of organic P usually gets smaller in the deeper soil (Achat et al. 2009). Thus, if P content is high in deep soil layers, as it was in some of our plots, the source of P in  
10 these plots is most likely to be the underlying bedrock. The plot-wise average pH of our soil samples agreed with that measured by Köster et al. (2014), who conducted their study at the same site, albeit not in the same plots. The pH of the soil humus layer correlated positively with the number of grass, herb and sedge species, which is reasonable, since higher pH usually implies a more fertile site. The soil N contents from our plots agreed with the reported values from Finnish forest sites (Merilä and Derome 2008, Salemaa et al. 2008), ranging between 9.8 and 12.8 g kg<sup>-1</sup>. Salemaa et al. (2008) reported a soil C:N ratio of 40 from a northern Finnish forest site, which is higher than we measured.

15

Our study area does not represent typical northern boreal forest, as it was located near the phosphate massif, the effect of which needs to be considered. Talvitie (1979), who used remote sensing for a geobotanical survey of the Sokli massif, found that the density of occurrence of birch, juniper and grass species increased when carbonatite was the underlying rock material. The surveys related to Natura 2000 (Standard Data Forms FII301512 and FII301513) stated that the Törmäoja and Yli-Nuortti  
20 areas, where plots B1-B3 and A1-A2 were, have a high occurrence of grass species and a sparse birch-dominated tree cover due to carbonatite in the soil. According to the geological map, only small parts in the western ends of both the Törmäoja and Yli-Nuortti Natura areas are located on top of carbonatite rock. Similarly, the map shows that those of our plots where the vegetation community was reminiscent of Sokli have something other than carbonatite as the rock parent material. However, all of our plots have metamorphic (tonalitic migmatite and amphibolite) or igneous (mafic volcanic and ultramafic) rock as the  
25 parent material, and phosphate mineral apatite can occur in such rocks (Walker and Syers 1976). It is likely that these types of rock materials leach more phosphate than other types of bedrock (Arnesen et al. 2007). Thus, the rocks outside of the carbonatite massif may also have locally high P content, which affects the P content of the soil. The mixed-effect model factor ‘geology’ did not consider this, which could be the reason why it was not important in explaining soil P content.

30 The baseline status and the current vegetation composition of our research site was worth studying for several reasons. We conducted our study in a region which has for decades been under more or less heated discussion related to whether mining activities will begin or not. The site is very remote and the plan is to move the material from the mine to the locations of further production by trucks (Pöyry Environment 2009). In addition to the aerial deposition from the mine, this could increase the dust and pollution caused by transportation, the amount of which is currently minimal. The effects of mining on the surrounding

ecosystem and its vegetation composition can be unpredictable when combined with the changes caused by climate change. High latitude regions are considered more vulnerable to climate change than more southern regions (Hartmann et al. 2013). Soil microbial activity may change due to a warmer climate, and so N may become more available from organic sources (Rustad et al. 2001). This, together with high soil P may induce growth and affect vegetation dynamics. Climate change has already caused variation in the vegetation at high latitudes, as deciduous shrub coverage has expanded in the Arctic region (Sturm et al. 2001, Park et al. 2016). Greater deciduous shrub cover causes increased leaf litter input, which in turn may bring more nutrients that are recyclable to the ecosystem.

## 5 Conclusions

We found that the total P content of the soil humus layer was an important factor explaining the community composition of forest understorey vegetation near the Sokli phosphate ore in Finnish Lapland. The plots with high soil total P in the humus layer had birch as the dominating tree species. As also green birch leaves and leaf litter high high contents of P, we suggest that the leaf litter from birch caused the high total P contents in the humus layer, but the rock parent material As climate change and the possible mining activities may affect the nutrient and vegetation dynamics in the studied region, the research that we carried out has an important role in both clarifying the current situation and forming a baseline for evaluating the magnitude of changes in the future.

## Data availability

We have made all data used in the analyses publicly available. All can be downloaded at: <https://doi.org/10.23728/b2share.615b46018cef40fe8c9d9245c56f0547>.

## Author contributions

LM and JB planned the study set-up; LM conducted all fieldwork, laboratory analyses and statistical analyses and led the writing process; MS had a substantial role in guiding the ordination analyses and the writing process; all authors contributed to the writing.

## 25 Competing interests

The authors declare that they have no conflict of interest.

## Acknowledgements

We thank the staff at the Värriö Subarctic Research Station for providing full board during the field-work, Marjut Wallner for guidance in the laboratory, Jarkko Isotalo for commenting on the statistical analyses, Jukka Pumpanen and Kajar Köster for advice and equipment for the soil sampling and Olli Peltola for helping with field-work. The study was supported by the Maj and Tor Nessling Foundation and Finnish Centre of Excellence grants 272041 and 307331.

## References

- Achat D. L., Bakker M. R., Augusto L., Saur E., Dousseron L., and Morel C.: Evaluation of the phosphorus status of P-deficient podzols in temperate pine stands: combining isotopic dilution and extraction methods, *Biogeochemistry*, 92, 183–200, doi:10.1007/s10533-008-9283-7, 2009.
- 10 Arnesen G., Beck P. S. A., and Engelskjøn T.: Soil Acidity, Content of Carbonates, and Available Phosphorus Are the Soil Factors Best Correlated with Alpine Vegetation: Evidence from Troms, North Norway, *Arct. Antarct. Alp. Res.*, 39, 189–199, doi:10.1657/1523-0430(2007)39[189:SACOCA]2.0.CO;2, 2007.
- Augusto L., Achat D. L., Jonard M., Vidal D., and Ringeval B.: Soil parent material—A major driver of plant nutrient limitations in terrestrial ecosystems, *Glob. Change Biol.*, 23, 3808–3824, doi:10.1111/gcb.13691, 2017.
- 15 Bates D., Maechler M., Bolker B., Walker S: Fitting Linear Mixed-Effects Models Using lme4, *J. Stat. Soft.*, 67, 1 – 48, doi:10.18637/jss.v067.i01, 2015.
- Brække F. H. and Salih N.: Reliability of Foliar Analyses of Norway Spruce Stands in a Nordic Gradient, *Silva Fenn.*, 36, 489–504, <http://dx.doi.org/10.14214/sf.540>, 2002.
- Cajander A.K.: Über Waldtypen, *Acta For. Fenn.*, 1,1–175, 1909.
- 20 Cajander A.K.: Forest types and their significance, *Acta For. Fenn.*, 56, 1–71, 1949.
- Dirnböck T., Grandin U., Bernhardt-Römermann M., Beudert B., Canullo R., Forsius M., Grabner M.-T., Holmberg M., Kleemola S., Lundin L., Mirtl M., Neumann M., Pompei E., Salemaa M., Starlinger F., Staszewski T., and Uzieblo A.K.: Forest floor vegetation response to nitrogen deposition in Europe, *Glob. Change Biol.*, 20, 429–440, doi:10.1111/gcb.12440, 2014.
- 25 Dupré C., Wessberg C., and Diekmann M.: Species richness in deciduous forests: Effects of species pools and environmental variables, *J. Veg. Sci.*, 13, 505–516, doi:10.1111/j.1654-1103.2002.tb02077.x, 2002.
- FAO: FAO/Unesco Soil map of the world, revised legend, *World Soil Resources Report 60*, FAO, Rome, 140 pp., 1988.
- Ferm A. and Markkola A.: Hieskoivun lehtien, oksien ja silmujen ravinnepitoisuuksien kasvukautinen vaihtelu. Abstract: Nutritional variation of leaves, twigs and buds in *Betula pubescens* stands during the growing season, *Folia For.*, 613, 1–28, 1985.
- 30 Finnish Biodiversity Information Facility: <https://laji.fi/en>, last access March 20<sup>th</sup> 2019.
- Hakku service: <https://hakku.gtk.fi/en/locations/search>, last access February 28<sup>th</sup> 2019.

- Hari P., Kulmala M., Pohja T., Lahti T., Siivola E., Palva L., Aalto P., Hämeri K., Vesala T., Luoma S., and Pulliainen E.: Air pollution in Eastern Lapland: Challenge for an environmental measurement station, *Silva Fenn.*, 28, 29–39, doi: 10.14214/sf.a9160, 1994.
- Hartmann D.L., Klein Tank A.M.G., Rusticucci M., Alexander L.V., Brönnimann S., Charabi Y., Dentener F.J., Dlugokencky E.J., Easterling D.R., Kaplan A., Soden B.J., Thorne P.W., Wild M. and Zhai P.M.: Observations: Atmosphere and Surface. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T.F., Qin D., Plattner G.-K., Tignor M., Allen S.K., Boschung J., Nauels A., Xia Y., Bex V. and Midgley P.M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 5 Hedwall P.-O., Bergh J., and Brunet J.: Phosphorus and nitrogen co-limitation of forest ground vegetation under elevated anthropogenic nitrogen deposition, *Oecologia*, 185, 317–326, doi:10.1007/s00442-017-3945-x, 2017.
- Helmisaari H.-S.: Temporal Variation in Nutrient Concentrations of *Pinus sylvestris* Needles, *Scand. J. of Forest Res.*, 5, 177–193, doi:10.1080/02827589009382604, 1990.
- Hobbie S. E., Nadelhoffer K. J., and Högberg P.: A synthesis: The role of nutrients as constraints on carbon balances in boreal and arctic regions, *Plant Soil*, 242, 163–170, doi:10.1023/A:1019670731128, 2002.
- 15 Hofmeister J., Hošek J., Modrý M., and Roleček J.: The influence of light and nutrient availability on herb layer species richness in oak-dominated forests in central Bohemia, *Plant Ecol.*, 205, 57–75, doi:10.1007/s11258-009-9598-z, 2009.
- Hämet-Ahti L.: The boreal zone and its biotic subdivision, *Fennia*, 159, 69–75, 1981.
- 20 Jaeger B.: *r2glmm: Computes R Squared for Mixed (Multilevel) Models*. R package version 0.1.2. available at: <https://CRAN.R-project.org/package=r2glmm>, 2017.
- Jaeger B., Edwards L. J., Das K., and Sen P. K.: An  $R^2$  statistic for fixed effects in the generalized linear mixed model, *J. Appl. Stat.*, 44, 1086 – 1105, doi: 10.1080/02664763.2016.1193725, 2016.
- Jobbágy E. G. and Jackson R.B.: The uplift of soil nutrients by plants: biogeochemical consequences across scales. *Ecology*, 25 85, 2380 – 2389, doi: 10.1890/03-0245, 2004.
- Johnson P. C. D.: Extension of Nakagawa & Schielzeth's  $R^2$ GLMM to random slopes models, *Methods Ecol. Evol.*, 5, 944 – 946, doi: 10.1111/2041-210X.12225, 2014.
- Koerselman W. and Meuleman A. F. M.: The vegetation N:P ratio: a new tool to detect the nature of nutrient limitation, *J. Appl. Ecol.*, 33, 1441–1450, doi:10.2307/2404783, 1996.
- 30 Kuusipalo J.: An ecological study of upland forest site classification in southern Finland, *Acta For. Fenn.*, 192, 77 pp., doi:10.14214/aff.7638, 1985.
- Köster K., Berninger F., Lindén A., Köster E., and Pumpanen J.: Recovery in fungal biomass is related to decrease in soil organic matter turnover time in a boreal fire chronosequence, *Geoderma*, 235–236, 74–82, doi:10.1016/j.geoderma.2014.07.001, 2014.

- Laasasenaho J.: Taper curve and volume functions for pine, spruce and birch, *Commun. Inst. For. Fenn.*, 108, 1–74, available at: <http://jukuri.luke.fi/handle/10024/522516>, 1982.
- Liski J.: Variation in soil organic carbon and thickness of soil horizons within a boreal forest stand – effect of trees and implications for sampling, *Silva Fenn.*, 29, 255–266, doi:10.14214/sf.a9212, 1995.
- 5 Lukina N., Tikhonova E., Danilova M., Bakhmet O., Kryshen A., Tebenkova D., Kuznetsova A., Smirnov A., Braslavskaya T., Gornov A., Shashkov M., Knyazeva S., Kataev A., Isaeva L., and Zukert N.: Associations between forest vegetation and the fertility of soil organic horizons in northwestern Russia, *Forest Ecosyst.*, 6, 34, doi: 10.1186/s40663-019-0190-2, 2019.
- Marschner H.: *Mineral Nutrition of Higher Plants*, Academic Press Limited, Cambridge, Great Britain, 1995.
- 10 Merilä P. and Derome J.: Relationships between needle nutrient composition in Scots pine and Norway spruce stands and the respective concentrations in the organic layer and in percolation water, *Boreal Env. Res.*, 13 (suppl. B), 35–47, 2008.
- Mikkola K. and Sepponen P.: Kasvupaikkatekijöiden ja kasvillisuuden suhteet Luoteis-Enontekiön tunturikoivikossa – Relationships between site factors and vegetation in mountain birch stands in northwestern Enontekiö, *Folia For.*, 674, 1–30, 1986.
- 15 Minchin P. R.: An evaluation of the relative robustness of techniques for ecological ordination, *Vegetatio*, 69, 89–107, 1987.
- Moilanen M., Saarinen M., and Silfverberg K.: Foliar nitrogen, phosphorus and potassium concentrations of Scots pine in drained mires in Finland, *Silva Fenn.*, 44, 583–601, doi:10.14214/sf.129, 2010.
- Moilanen M., Saarsalmi A., Kukkola M., and Issakainen J.: Effects of stabilized wood ash on nutrient status and growth of Scots pine – Comparison between uplands and peatlands, *Forest Ecol. Manag.*, 295, 136–144, doi:10.1016/j.foreco.2013.01.021, 2013.
- 20 Mäkipää R.: Response Patterns of *Vaccinium myrtillus* and *V. vitis-idaea* along Nutrient Gradients in Boreal Forest, *J. Veg. Sci.*, 10, 17–26, doi:10.2307/3237156, 1999.
- Nakagawa S. and Schielzeth H.: A general and simple method for obtaining  $R^2$  from generalized linear mixed-effects models, *Methods. Ecol. Evol.*, 4, 133 – 142, doi: 10.1111/j.2041-210x.2012.00261.x, 2013.
- 25 NASA: <https://worldview.earthdata.nasa.gov/>, last access December 17th 2019.
- NATURA 2000 - Standard Data Form FI1301512: <http://natura2000.eea.europa.eu/Natura2000/SDF.aspx?site=FI1301512>, last access February 28th 2019.
- NATURA 2000 - Standard Data Form FI1301513: <http://natura2000.eea.europa.eu/Natura2000/SDF.aspx?site=FI1301513>, last access February 28th 2019.
- 30 Näsholm T., Kielland K., and Ganeteg U.: Uptake of organic nitrogen by plants, *New Phytol.*, 182, 31–48, doi: 10.1111/j.1469-8137.2008.02751.x, 2008.
- Økland R. H. and Eilertsen O.: Dynamics of understory vegetation in an old-growth boreal coniferous forest, 1988 -1993, *J. Veg. Sci.*, 7, 747–762, doi:10.2307/3236386, 1996.

- Oksanen J., Blanchet F. G., Friendly M., Kindt, R., Legendre P., McGlenn D., Minchin P. R., O'Hara R. B., Simpson G. L., Solymos P., Stevens M. H. H., Szoecs E., and Wagner H.: *vegan: Community Ecology Package*, R package version 2.4-6, available at: <https://CRAN.R-project.org/package=vegan>, 2018.
- 5 Park T., Sangram G., Tømmervik H., Euskirchen E. S., Høgda K-A., Karlsen S. R., Brovkin V., Nemani R. R., and Myneni R. B.: Changes in growing season duration and productivity of northern vegetation inferred from long-term remote sensing data, *Environ. Res. Lett.*, 11, 1–11, doi:10.1088/1748-9326/11/8/084001, 2016.
- Pirinen P., Simola H., Aalto J., Kaukoranta J-P., Karlsson P., and Ruuhela R.: *Tilastoja Suomen ilmastosta 1981-2010. Ilmatieteen laitos, Raportteja 2012:1*, Finnish Meteorological Institute. 94 pp, available at: <https://helda.helsinki.fi/handle/10138/35880>, 2012.
- 10 Pöyry Environment: *Soklin kaivoshankkeen YVA-selostus*, 332 pp., 2009.
- R Development Core Team: *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, available at: <http://www.R-project.org/>, 2017.
- Rautio P., Fürst A., Stefan K., Raitio H., and Bartels U.: *Sampling and Analysis of Needles and Leaves, Manual Part XII*, in: *Manual on methods and criteria for harmonized sampling, assessment, monitoring and analysis of the effects of air pollution on forests*, edited by: UNECE, ICP Forest Programme Co-ordinating Centre, Hamburg, 19 pp., available at: <http://icp-forests.net/page/icp-forests-manual>, 2010.
- 15 Reimann C., Boyd R., de Caritat P., Halleraker J. H., Kashulina G., Niskavaara H., and Bogatyrev I.: *Topsoil (0-5 Cm) Composition In Eight Arctic Catchments In Northern Europe (Finland, Norway And Russia)*, *Environ. Pollut.*, 1, 45–56, doi: 10.1016/S0269-7491(96)00102-9, 1997.
- 20 Reinikainen A., Mäkipää R., Vanha-Majamaa I., and Hotanen J.-P.: *Kasvit muuttuvassa metsäluonnossa (Changes in the frequency and abundances of forest and mire plants in Finland since 1950)*, Kustannusyhtiö Tammi, Helsinki, 2000.
- Reta G., Dong X., Li Z., Su B., Hu X., Bo H., Yu D., Wan H., Liu J., Li Y., Xu G., Wang K., and Xu S.: *Environmental impact of phosphate mining and beneficiation: review*, *Int. J. Hydro.*, 2, 424–431, doi: 10.15406/ijh.2018.02.00106, 2018.
- Rinnan R., Michelsen A., and Jonasson S.: *Effects of litter addition and warming on soil carbon, nutrient pools and microbial communities in a subarctic heath ecosystem*, *Appl. Soil Ecol.*, 39, 271–281, doi:10.1016/j.apsoil.2007.12.014, 2008.
- 25 Rustad L.E., Campbell J.L., Marion G.M., Norby R.J., Mitchell M.J., Hartley A.E., Cornelissen J.H.C., Gurevitch J., and GCTE-NEWS: *A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming*, *Oecologia*, 126, 543–562, doi:10.1007/s004420000544, 2001.
- Ruuskanen T. M., Reissel A., Keronen P., Aalto P. P., Laakso L., Grönholm T., Hari P., and Kulmala M.: *Atmospheric trace gas and aerosol particle concentration measurements in Eastern Lapland, Finland 1992-2001*, *Boreal Env. Res.*, 8, 335–349, 2003.
- 30 Saarsalmi A. and Mälkönen E.: *Forest Fertilization Research in Finland: A Literature Review*, *Scand. J. Forest Res.*, 16, 514–535 doi:10.1080/02827580152699358, 2001.

- Salemaa M., Monni S., Royo Peris F., and Uhlig C.: Sampling strategy for the assessment of temporal change in ground vegetation in boreal forests, in: Forest condition monitoring in Finland, edited by: Raitio, H. and Kilponen, T., National report 1998, The Finnish Forest Research Institute, Research Papers, 743,117–127, 1999.
- Salemaa M., Derome J., and Nöjd P.: Response of boreal forest vegetation to the fertility status of the organic layer along a climatic gradient, *Boreal Env. Res.*, 13 (suppl. B), 48–66, 2008.
- Salemaa, M., Lindroos, A.-J., Merilä, P., Mäkipää, R., and Smolander, A: N<sub>2</sub> fixation associated with the bryophyte layer is suppressed by low levels of nitrogen deposition in boreal forests, *Sci. Total Environ.*, 653, 995–1004, doi: 10.1016/j.scitotenv.2018.10.364, 2019.
- Sorensen P. L. and Michelsen A.: Long-term warming and litter addition affects nitrogen fixation in a subarctic heath, *Glob. Change Biol.*, 17, 528–537, doi: 10.1111/j.1365-2486.2010.02234.x, 2011.
- Sturm M., Racine C., and Tape K.: Increasing shrub abundance in the Arctic, *Nature*, 411, 546–547, doi:10.1038/35079180, 2001.
- Talvitie J.: Remote sensing and geobotanical prospecting in Finland, *Bull. Geol. Soc. Finland*, 51, 63–73, 1979.
- Tamm C. O.: Nitrogen in terrestrial ecosystems: Questions of productivity, vegetational changes, and ecosystem stability, *Ecological Studies* 81, Springer-Verlag, Berlin, Heidelberg, 116 pp., 1991.
- Tonteri T., Salemaa M., Rautio P., Hallikainen V., Korpela L., and Merilä P.: Forest management regulates temporal change in the cover of boreal plant species, *Forest Ecol. Manag.*, 381, 115–124, doi:10.1016/j.foreco.2016.09.015, 2016.
- Tuovinen H., Vesterbacka D., Lempinen J., Pohjolainen. E., Read D., Solatie D., and Lehto J.: Radionuclide and heavy metal redistribution at the former pilot-scale apatite plant in Sokli, northern Finland, *Boreal Env. Res.*, 20, 350–362, available at: <https://helda.helsinki.fi/handle/10138/228212>, 2015.
- Ukonmaanaho L., Merilä P., Nöjd P. and Nieminen T. M.: Litterfall production and nutrient return to the forest floor in Scots pine and Norway spruce stands in Finland, *Boreal Env. Res.*, 13 (suppl. B), 67–91, 2008.
- Vartiainen H., and Paarma H.: Geological Characteristics of the Sokli Carbonatite Complex, Finland, *Econ. Geol.*, 74, 1296–1306, doi:0.2113/gsecongeo.74.5.1296, 1979.
- Verheven K., Baeten L., De Frenne P., Bernhardt-Romermann M., Brunet J., Cornelis J., Decocq G., Dierschke H., Eriksson O., Hedl R., Heinken T., Hermy M., Hommel P., Kirby K., Naaf T., Peterken G., Petrik P., Pfadenhauer J., Van Calster H., Walther G.R., Wulf M., and Verstraeten G.: Driving factors behind the eutrophication signal in understorey plant communities of deciduous temperate forests, *J. Ecol.*, 100, 352–365, doi: 10.1111/j.1365-2745.2011.01928.x, 2012.
- Vinton M. A. and Burke I.C.: Interactions between individual plant species and soil nutrient status in shortgrass steppe, *Ecology*, 76, 1116–1133, doi:10.2307/1940920, 1995.
- Viro P. J.: Investigations on forest litter, *Commun. Inst. For. Fenn.*, 45, 60 pp., 1955.
- Walker T. W. and Syers J. K.: The fate of phosphorus during pedogenesis, *Geoderma*, 15, 1–19, doi:10.1016/0016-7061(76)90066-5, 1976.




Weintraub M. N.: Biological Phosphorus Cycling in Arctic and Alpine Soils, in: Phosphorus in action , edited by: Bünemann E. K., Oberson A., and Frossard E., Soil Biol., 26, 295–316, Springer, Berlin, Heidelberg, doi:10.1007/978-3-642-15271-9\_12, 2011.

5


10

15

20



**Figure 1:** Left: Geological map of the research area. Plots are marked with black stars. The easternmost plot is located at the SMEAR1 station. (Source: Hakku Service, <https://hakku.gtk.fi/en/locations/search>.), Right: Satellite image showing the study location (source: NASA).



- ◆ = centre of the plot
- = understorey vegetation subplot (1 m<sup>2</sup>)
- = soil sampling point
- ↖ ↗ ↘ ↙ = outer corner of the whole research plot

20 **Figure 2: Set-up of each research plots with clusters and subplots within clusters. Trees were measured from the whole 30 × 30 m area.**







Figure 3. Soil total phosphorus (P) contents of the research plots (based on dominant tree species) in different soil horizons. Species 1 = pine, 2 = spruce and 3 = birch.







Figure 4. Soil total N content (upper panel) and soil C:N ratio (lower panel) of the research plots (based on dominant tree species) in soil horizons F and O. Species 1 = pine, 2 = spruce and 3 = birch.



5 Figure 5. Correlation (Pearson) including soil elements (K, P, C:N, pH, C, N:P), number of species (with *\_n* in the end of the name) and total % cover of plant species in different layers (moss and lichen, grasses, herbs and sedges, dwarf shrubs and trees), needle elements (N, P, K), plot distance from Sokli, green birch elements (N, P, K) and birch litter elements (N, P, K). Levels of significance \* = 0.05, \*\* = 0.01, \*\*\* = 0.001. Positive correlations are displayed in blue and negative correlations in red. Colour intensity and size of the circle are proportional to the correlation coefficients.



5 **Figure 6: Ordination pattern of the research plots in dimensions 1 and 2 (a and b), and dimensions 1 and 3 (c and d). Figures (a) and (c) give plot ordinations and b and d weighted averages of the most abundant species (highest cover % of surface). Less abundant species are marked with light-coloured crosses. The names of species are combinations of the first four letters of genera and species names (e.g. Solivirg = *Solidago virgaurea*). The tree species mentioned in the figure are at seedling stage. In (a) and (c) plots D4 and B3 were located on top of each other and only D4 is shown.**



5 **Figure 7: Ordination pattern with smooth surface fit and linear vector fit of soil phosphorus (P) in the O layer (a), linear vector fits of soil element contents in the O layer (b), linear vector fits of foliar data as well as plot distance from Sokli phosphate ore (c), and linear vector fits of number of species in different layers of understory (d). Moss\_lichen includes moss and lichen species, Grass\_sedge includes forb, grass, and sedge species and D.shrubs\_trees includes dwarf shrubs and tree seedlings. In (a) and (c) plots D4 and B3 were located on top of each other and only D4 is shown.**



5 **Table 1. Meteorological parameters from Värriö. SWE=snow water equivalent, \*= data from SMEAR 1 station, \*\*= data from SMEAR station (only 2009–2015), otherwise data are collected from Värriö Subarctic Research Station by the Finnish Meteorological Institute. Values for the climatological normal period are from Pirinen et al. 2012. Growing degree day sum was calculated as the average daily temperature (average of daily maximum and minimum temperatures) above 5 °C base temperature, accumulated on a daily basis over the year. Negative values are treated as zeros and ignored.**

	2014	2015	Climatological normal period (1981-2010)
Average annual temperature (°C)	0.84	0.95	−0.5
Average min. temperature (°C)	−2.09	−1.7	−3.5
Average max. temperature (°C)	3.9	3.8	2.6
Growing degree day sum	860	640	680
Total precipitation (mm)	610	660	601
· Snowfall (mm, SWE)*	390	420	400 **
· Rainfall (mm)	220	240	190 **

**Table 2. Tree species composition of the research plots. dbh = diameter at breast height.**

Plot	Trees/ha	Basal area of trees (m <sup>2</sup> /ha)	Total volume of trees (m <sup>3</sup> /ha)	Volume of pine (m <sup>3</sup> /ha)	Volume of spruce (m <sup>3</sup> /ha)	Volume of birch (m <sup>3</sup> /ha)	Average dbh of pine (cm)	Average dbh of spruce (cm)	Average dbh of birch (cm)
A1	1300	10	75	44	0.02	30.7	10	1.8	7
A2	1200	12	78	69.2	2.3	6.9	9.1	6.5	5.7
A3	900	8	46	-	0.5	45.5	-	6.4	9.1
A4	600	16	130	83.7	35.5	11.2	21.5	11.3	9
A5	1200	17	125	94.5	20.9	10.0	18.7	8.4	6.2
A6	500	10	54	53.7	-	0.7	16.7	-	3.9
B1	300	1	3	2.8	0.07	-	13.4	5.6	-
B2	300	5	33	32.6	-	-	18.2	-	-
B3	500	17	128	127.9	0.2	0.4	19.2	6.8	5.7
C1	800	3	14	14.3	-	-	6.1	-	-
C2	1100	14	105	47.1	41.1	17.7	21.6	12.1	7
D1	700	14	99	99.3	-	0.03	11.9	-	3.3
D2	1100	9	48	-	-	48	-	-	9.8
D3	500	4	18	15.3	-	2.5	8.5	-	9.2
D4	300	7	43	13.6	17.2	12.3	21.5	22	9.9
D5	700	11	98	98.3	-	-	11.4	-	-

**Table 3. Average foliar element contents (g kg<sup>-1</sup>) of the three major nutrients and C, and the relationships of C:N and N:P with standard deviations. Needle age classes: C = current year, C+1 = one-year-old needles, C+2 = two-year-old needles.**

	<b>C</b>	<b>N</b>	<b>P</b>	<b>K</b>	<b>C:N</b>	<b>N:P</b>	5
Pine C	510.0 (3.70)	14.10 (0.80)	1.40 (1.40)	4.50 (9.10)	36.0 (1.90)	10.0 (1.10)	
Pine C+1	510.0 (7.30)	13.80 (0.90)	1.20 (0.80)	3.70 (3.70)	38.0 (2.40)	11.80 (1.10)	
Pine C+2	480.0 (140.0)	12.10 (3.70)	1.2 (0.90)	3.50 (3.30)	36.6 (11.20)	10.60 (3.30)	10
Spruce C	500 (3.80)	12.0 (1.00)	1.70 (0.20)	6.40 (0.90)	42.0 (3.50)	7.10 (0.70)	
Spruce C+1	440.0 (170.0)	10.70 (4.20)	1.50 (0.20)	4.20 (0.90)	37.0 (14.50)	7.30 (2.90)	
Spruce C+2	440.0 (165.0)	10.10 (3.90)	1.40 (0.20)	3.70 (0.80)	39.2 (14.90)	7.50 (3.00)	15
Birch, green	470.0 (3.40)	25.0 (1.20)	2.60 (0.30)	8.20 (1.50)	19.0 (1.0)	9.80 (1.0)	
Birch, litter	490.0 (6.0)	10.10 (1.40)	1.30 (0.50)	2.40 (1.00)	50.0 (7.40)	8.50 (2.10)	

**20 Table 4. Linear correlations of element contents of soil, needles and leaves, number of**

**25 species in different vegetation layers and plot distance from Sokli with the non-metric multidimensional scaling ordination pattern. The group of ‘grasses and sedges’ includes forb, grass and sedge species and ‘d. shrubs and trees’ includes dwarf shrubs and tree seedlings. Levels of significance: ° = 0.1, \* = 0.05, \*\* = 0.01. The first seven rows are soil values.**

	Variable	R <sup>2</sup>	p <
	K	0.225	0.220
	P	0.717	0.002**
	N	0.368	0.087 <sup>o</sup>
5	C	0.137	0.461
	C:N	0.576	0.012*
	N:P	0.431	0.045*
	pH	0.386	0.075 <sup>o</sup>
	Needle P	0.440	0.033*
	Needle N	0.010	0.927
10	Needle K	0.465	0.029*
	Birch P	0.247	0.213
	Birch N	0.104	0.569
	Birch K	0.346	0.099 <sup>o</sup>
	Moss and lichen, species number	0.249	0.223
15	Grasses and sedges, species number	0.738	0.003**
	D. shrubs and trees, species number	0.181	0.341
	Moss and lichen, % cover of surface	0.180	0.325
	Grasses and sedges, % cover of surface	0.248	0.196
	D.shrubs and trees, % cover of surface	0.250	0.198
20	Plot distance from Sokli	0.183	0.344

25

**Table 5. Results from the mixed-effect models, testing the effects of environmental variables on soil total P and N content and C:N ratio. The tested variables were dominant tree species of the research plot, estimated tree age, rock parent material (geology) and soil layer. Random effect was related to plot number. Pseudo-R<sup>2</sup> was calculated based on Nakagawa and Schielzeth (2013), Johnson (2014) and Jaeger et al. (2016).**

5			
Soil total P content			
Fixed effects	Chisq value	p-value	Pseudo-R <sup>2</sup>
factor (dominant tree species)	7.9009	0.01925	0.45
factor (tree age)	4.0408	0.1326	
factor (geology)	4.8171	0.08995	
factor (soil layer)	155.97	2.20 x 10 <sup>-16</sup>	
Soil total N content			
Fixed effects	Chisq value	p-value	Pseudo-R <sup>2</sup>
factor (dominant tree species)	4.9146	0.08567	0.2
factor (tree age)	2.1769	0.3367	
factor (geology)	2.2291	0.3281	
factor (soil layer)	53.408	2.71 x 10 <sup>-13</sup>	
Soil total C:N ratio			
Fixed effects	Chisq value	p-value	Pseudo-R <sup>2</sup>
factor (dominant tree species)	4.2076	0.122	0.2
factor (tree age)	1.3484	0.5096	
factor (geology)	0.3339	0.8462	
factor (soil layer)	60.036	9.31 x 10 <sup>-15</sup>	

**Appendix A: Average coverage (% of surface area) of understory plant species per plot**

Species	A1	A1	A3	A4	A5	A6	B1	B2	B3	C1	C2	D1	D2	D3	D4	D5
<i>Pleurozium schreberi</i>	43.3	51.3	39.6	59.3	44.4	57.1	25.0	24.7	64.7	9.0	38.8	0.3	1.2	53.5	14.5	38.7
<i>Hylocomium splendens</i>	38.7	8.3	22.8	9.6	3.3	1.7	–	3.8	1.3	–	28.9	–	44.6	–	14.5	–
<i>Dicranum scoparium</i>	–	12.6	–	4.2	5.9	10.7	1.6	2.8	–	8.3	–	72.9	–	1.0	9.7	4.1
<i>Dicranum polysetum</i>	–	–	–	–	0.2	0.1	–	0.9	0.8	–	–	–	–	–	0.6	0.4
<i>Dicranum majus</i>	–	–	–	1.1	–	–	–	–	–	–	–	–	–	–	–	1.3
<i>Barbilophozia barbata</i>	–	–	–	0.1	0.7	0.6	–	–	–	–	–	–	–	0.2	0.2	5.8
<i>Polytrichum strictum</i>	–	–	2.8	–	–	–	5.0	5.3	–	0.8	–	–	–	2.1	–	–
<i>Polytrichum commune</i>	3.4	0.4	0.7	0.1	–	–	0.4	–	–	–	1.4	–	23.4	4.0	4.9	0.1
<i>Aulacomnium palustre</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	3.7	–	–
<i>Sphagnum angustifolium</i>	–	–	–	–	–	–	–	–	–	–	1.0	–	–	–	–	–
<i>Sphagnum girgensohnii</i>	–	–	–	–	–	–	–	–	–	–	0.3	–	–	–	–	–
<i>Sphagnum capillifolium</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	0.2	–	–
<i>Ptilidium ciliare</i>	–	–	–	–	–	1.1	–	–	–	–	–	–	–	–	–	–
<i>Peltigera aphthosa</i>	1.1	–	–	–	1.0	–	0.6	0.7	0.4	–	–	–	–	–	0.1	–
<i>Peltigera rufescens</i>	–	–	0.2	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Peltigera neopolydactyla</i>	–	–	–	–	–	–	10.9	0.1	–	–	–	–	–	–	–	–
<i>Nephroma arcticum</i>	–	2.3	1.0	3.9	–	–	6.1	1.5	0.5	–	–	–	–	0.5	0.1	1.3
<i>Umbilicaria deusta</i>	–	–	–	–	–	–	–	0.1	–	–	–	–	–	–	–	–
<i>Cladonia rangiferina</i>	0.3	1.0	0.3	1.6	1.5	2.0	6.2	2.8	1.6	31.0	–	3.8	–	5.2	0.3	5.3
<i>Cladonia cornuta</i>	–	–	0.1	–	0.2	–	0.1	0.2	–	0.1	–	0.2	–	–	–	0.3
<i>Cladonia stellaris</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Cladonia deformis</i>	–	–	–	–	–	–	–	0.1	–	0.6	–	–	–	–	–	0.1
<i>Cladonia crispata</i> var. <i>crispata</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Cetraria islandica</i>	–	0.1	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Stereocaulon tomentosum</i>	–	–	–	–	–	–	4.7	2.2	–	–	–	–	–	–	–	–
<i>Linnaea borealis</i>	–	–	–	–	0.3	0.2	–	–	0.1	–	–	–	–	–	–	–
<i>Vaccinium myrtillus</i>	1.2	4.3	–	10.7	14.7	27.7	–	5.2	2.3	1.4	8.4	1.3	2.7	18.8	6.0	6.9
<i>Vaccinium vitis-idaea</i>	15.3	26.8	5.6	22.2	11.9	5.2	1.3	5.5	21.6	5.7	3.4	28.2	25.8	4.0	7.4	4.4
<i>Vaccinium uliginosum</i>	7.1	0.8	–	–	–	0.5	15.5	5.3	2.3	2.9	34.3	–	11.6	41.7	6.3	1.5

<i>Empetrum nigrum</i>	2.7	9.8	0.8	8.4	14.1	12.4	–	9.6	32.8	9.3	14.5	15.0	4.1	40.4	7.0	6.9
<i>Arctostaphylos uva-ursi</i>	–	–	–	–	–	–	10.7	0.2	0.4	–	–	–	–	–	–	–
<i>Arctostaphylos alpina</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Betula nana</i>	23.9	6.3	1.3	–	–	–	–	–	–	–	2.2	–	–	2.7	–	–
<i>Calluna vulgaris</i>	–	–	–	–	–	0.5	–	–	–	0.2	–	–	–	0.6	–	0.4
<i>Rhododendron tomentosum</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	0.8	3.6	–
<i>Juniperus communis</i>	0.9	–	0.8	0.6	–	–	5.7	0.3	2.2	–	2.5	–	3.7	–	–	–
<i>Picea abies</i>	–	–	–	0.5	–	–	–	–	–	–	–	–	–	–	–	–
<i>Pinus sylvestris</i>	–	–	–	–	–	–	1.6	0.8	–	2.1	0.1	0.9	–	0.8	–	0.7
<i>Betula pubescens</i>	1.1	0.2	1.4	–	–	0.4	2.5	0.7	–	–	–	–	–	0.2	0.1	–
<i>Populus tremula</i>	–	–	–	–	–	0.4	–	–	–	–	–	–	–	–	–	–
<i>Diphasiastrum complanatum</i>	–	–	–	–	–	0.1	0.7	–	–	–	–	–	–	–	–	–
<i>Trientalis europaea</i>	–	0.1	0.7	–	–	–	–	–	–	–	–	–	3.4	–	–	–
<i>Melampyrum sylvaticum</i>	–	–	0.3	–	–	–	0.1	–	–	–	–	–	–	0.1	–	–
<i>Solidago virgaurea</i>	–	–	1.1	–	–	–	1.2	–	–	–	0.2	–	0.7	–	–	–
<i>Rubus arcticus</i>	–	–	0.6	–	–	–	–	–	–	–	–	–	3.2	–	–	–
<i>Rubus chamaemorus</i>	–	–	–	–	–	–	–	–	–	–	0.4	–	–	–	–	–
<i>Antennaria dioica</i>	–	–	–	–	–	–	1.2	–	–	–	–	–	–	–	–	–
<i>Orthilia secunda</i>	–	–	–	–	–	–	–	–	–	–	0.1	–	–	–	–	–
<i>Epilobium angustifolium</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.7	–	–	–
<i>Galium uliginosum</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.1	–	–	–
<i>Geranium sylvaticum</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.2	–	–	–
<i>Chelidonium majus</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.1	–	–	–
<i>Comarum palustre</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.3	–	–	–
<i>Equisetum sylvaticum</i>	0.1	–	–	–	–	–	–	–	–	–	–	–	0.8	–	0.2	–
<i>Luzula pilosa</i>	0.6	–	0.4	–	–	–	1.8	–	–	–	0.2	–	2.4	0.3	–	–
<i>Elymus repens</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Deschampsia flexuosa</i>	3.5	0.6	10.3	0.6	–	0.2	2.1	0.3	0.4	–	2.8	–	15.9	7.1	2.2	0.2
<i>Festuca rubra</i>	–	–	–	–	–	–	–	–	–	–	–	–	1.9	–	–	–
<i>Calamagrostis epigejos</i>	–	–	–	–	–	–	–	–	–	–	–	–	5.4	–	–	–
<i>Carex digitata</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.6	–	–	–
<i>Carex nigra</i>	–	–	–	–	–	–	–	0.2	–	–	–	–	–	–	–	–

<i>Carex canescens</i>	-	-	-	-	-	-	-	-	-	-	0.5	-	3.4	-	-	-
<i>Carex globularis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	1.7	-	-



## Appendix B1. Soil element contents within and across plots

**Table B1. Average total contents of elements (g kg<sup>-1</sup>) in soil layers and their standard deviations in parenthesis. All plots are included.**

	<b>K</b>	<b>P</b>	<b>N</b>	<b>C</b>	<b>C:N</b>	<b>N:P</b>	<b>pH</b>
F layer	0.83 (0.30)	0.81 (0.32)	13.3 (2.9)	420 (98)	32 (6.8)	17.8 (5.5)	-
O layer	0.49 (0.22)	0.72 (0.50)	9.9 (3.7)	260 (106)	26 (5.7)	17.2 (7.3)	3.7 (0.2)
A layer	0.32 (0.22)	0.38 (0.51)	-	-	-	-	-
B layer	0.59 (0.23)	0.10 (0.13)	-	-	-	-	-

5

10

15

**Table B2. Statistical differences of soil elements in each soil layer of different plots, grouped by their dominant tree species. Levels of significance: \* = 0.05, \*\* = 0.01, \*\*\* = 0.001.**

	P	N	C:N
F layer	birch & pine*** birch & spruce**	birch & pine*** birch & spruce**	birch & pine** birch & spruce**
O layer	birch & pine*** birch & spruce***	birch & pine**	birch & pine* birch & spruce**
A layer	birch & pine*** birch & spruce*	–	–
B layer	–	–	–

5




Figure B1. The fitted vs. residuals, q-q plots and histograms of residuals from the mixed effect models (upper = phosphorus (P), lower = nitrogen (N)).







Figure B2. Soil total P content within plots in different soil layers.







Figure B3. Soil total N content and C:N ratio within plots in the F and O layers.

## Appendix C: Needle and leaf nutrient contents per plot

5 **Table C1. Statistically significant differences between needle age group by species (C = youngest needles, C+1 = one-year-old needles, C+2 = two-year-old needles),  $p < 0.05$  one-way analysis of variance, with Tukey's honestly significant difference post hoc test.**

	P	N	C	C:N	N:P
Pine	C & C+1, C & C+2	C & C+2	C & C+1, C & C+2	C & C+2	C & C+1, C & C+2
Spruce	C & C+1, C & C+2	No differences between age classes	No differences between age classes	No differences between age classes	C & C+1, C & C+2

10

15

20

25

30

**Table C2. Statistically significant differences of needle nutrient contents between plots,  $p < 0.05$ .**

	P	N	C	C:N	N:P
Pine	No differences between plots	Plots: A5&B1, A4&D3, A1&D3, B1&D3, B2&D3, B1&D1, B2&D1	No differences between plots	Plots: D3&A1, D3&B1, D3&B2	No differences between plots
Spruce	Plots: C2&A2	Plots: A3&A5, A2&A5, B3&A5, B3&A4, B3&B1, C2&B3, D5&B3, D4&B3	Plots: A4&A5, A3&A5, B1&A5, D4&A5, A2&A4, D5&A4, D5&A3, B1&A2, D4&A2, D5&B1	Plots: A3&A5, A2&A5, B3&A5, B3&A4, B3&B1, D5&B3, D4&B3	No differences between plots