

Interactive comment on “Ocean carbonate system variability in the North Atlantic Subpolar surface water (1993–2017)” by Coraline Leseurre et al.

Are Olsen (Referee)

are.olsen@uib.no

Received and published: 25 April 2019

This contribution presents the trends in a very impressive set of data from the western subpolar North Atlantic, collected within the framework of the SURATLANT program. While the paper presents the data and trends more or less adequately, it fails in properly attributing them to (climatic + oceanographic + biological) drivers. These aspects are unclear, speculative, and somewhat confused. As such, major revisions are required.

My major comments are:

1. Samples are collected two to four times a year. My sense is that with such rare sampling one might be very prone to getting false trends because of differences in the timing of the data collection, also in combination with timing of the spring bloom. For

Printer-friendly version

Discussion paper



example, the strong increase in region C+D+E summer DIC in 2001 -2008 might be generated by earlier sampling/later spring bloom.

While some effort seems to have been taken to deal with this; reconstructed data; it is not quite clear how this is done, and how well it works. Some improvement is needed in this description (page 5 lines 1-19). Is it possible to test this scheme by using data from a year with sampling in all of the months used (e.g. Jan-March)? At least for SST and SSS this can be done as there should be continuous TSG data available, and for fCO₂ there are data available from the VOSs Nuka Arctica and Atlantic Companion, which crosses the study area. These can be retrieved from SOCAT and used to test the method.

2. More broadly, it would be interesting to know how representative the data are for large-scale interannual phenomena. This can be tested. For example, the trends in SST can be compared with objectively analysed SST from NOAA. The trends in SSS can be compared with some reanalysis model output. And again, for the time representativeness, the continuous data from the TSG, which collects data on all crossings, should be used. It might also be worthwhile to look at remotely sensed Chl a, to evaluate if there are concomitant trends in surface ocean primary production, e.g. a loss in production from 2001 -2008 such as the DIC data seem to indicate and was suggested in the manuscript.

3. The attribution section is not well done. In particular, I strongly suggest the authors to explicitly account for salinity changes in the pCO₂ driver decomposition following the method by Keeling et al. (2004) and recently used in the subpolar North Atlantic by Fröb et al. (2019). The reason is that dilution may completely overwhelm increase of DIC expected from uptake of anthropogenic CO₂.

4. Further, showing trends in salinity normalised DIC and TA is worthwhile, but make sure to use the correct method for salinity normalising as described in Friis et al. (2003).

5. Also, there is a lot of mentioning of the change in the air-sea CO₂ difference. But this

[Printer-friendly version](#)[Discussion paper](#)

is not illustrated, which leaves a lot to the readers imagination. I therefore recommend to show the actual air-sea $f\text{CO}_2$ difference in Fig. 3. You can make room for this by removing one of the panels for Omega (Calcite/Aragonite), as there is no need to show both.

6. Uncertainties are not properly dealt with. For example, the uncertainty in the SST of 0.1 degrees C results in an $f\text{CO}_2$ uncertainty of 2 microatmospheres. There are also uncertainties in DIC and TA. These errors need to be propagated to calculated $f\text{CO}_2$, pH etc. This can be done using the most recent CO2SYS from James Orr as this includes error propagation. It is available from GitHub. The errors in $f\text{CO}_2$ (and the others) can be propagated to the trends using Monte Carlo.

7. I am in particular concerned with the fact that the large summer $f\text{CO}_2$ increase in region B 2001-2008 are basically caused by the 'reconstructed' data of 2001 and 2008. The actual observations are pretty steady. How confident are you in these reconstructed values?

8. As the other reviewer, I think this contribution confuses anthropogenic and natural CO_2 . The trends that are observed in $p\text{CO}_2$ does not say any on anthropogenic carbon uptake. Further, the North Atlantic is not a big sink of anthropogenic CO_2 because the air-sea flux is large. The air-sea flux is a combination of natural and anthropogenic CO_2 fluxes. Horizontal advection is likely a big source of anthropogenic CO_2 to the North Atlantic. Hence, the data that are presented only informs about the changes in surface $p\text{CO}_2$ and air-sea CO_2 flux, not on the North Atlantic sink for anthropogenic CO_2 . This needs to be considered both in the introduction and in the discussion.

9. I think the dicussion is pretty dissapointing. It is basically a recap of the results + some more exploration of these, combined with some speculation based on published litterature. There are no attempts to analyse the relation between the observed trends and likely drivers (for which data exist) – such as NAO or AMV indices, winter/summer mixed layer depths (Argo data), primary production (remotely sensed ocean color), and

[Printer-friendly version](#)[Discussion paper](#)

SGP strenght (SPG index). This should be done.

10. Finally, Metzl et al (2010) suggested deep mixing as the cause for the sharp winter fCO₂ trend in 2001-2008. Fröb et al (2019) demonstrated unequivocally that deep mixing leads to strong increases in winter fCO₂ in NAO positive years because of intensified deep mixing (bringing remineralised DIC to the surface). In light of this, it is interesting that the smallest winter fCO₂ trends from SURATLANT are observed in the period 2008-2017, as it is well known that this has been a period of rather large deep mixing compared to 2001-2008. I am not sure how this can be reconciled, i.e. the large fCO₂ trend in the period of little deep mixing 2001-2008 vs the small fCO₂ trend in the period of frequent deep mixing in 2008-2017. It seems worthwhile to dvelve into this. While doing so, keep in mind that deep-mixing events seemed (in Fröb et al., 2019) to cause year-to-year anomalies and not so much an anomalous trend. Therefore, I recommend to reconsider the use of three periods - the trend in each of these can be strongly affected by the fCO₂ in the start and/or end year, which might just be an anomalous year. Therefore the authors might be doing themselves a disfavor by sticking to the three periods, which are largely defined because of 'historical'/'traditional' reasons related to sampling and previous SURATLANT papers.

Other comments:

Page 2, line 4. Takahashi's estimates of air-sea CO₂ flux cannot be equated to anthropogenic CO₂ flux.

Page 2, line 6. I don't think any has ascribed the variations in the North Atlantic CO₂ sink to climate change. Careful with such statements.

Page 2, line 21. 'based on Total Alkalinity (TA) and Dissolved Inorganic Carbon (DIC) observations,..'

Page 3, line 7. 'which', not 'witch'.

Page 3, line 14-15. 'to this end', not 'to this aim'.

BGD

Interactive
comment

Printer-friendly version

Discussion paper



Page 3, line 15. 'Iceland', not 'Island'

Page 3, line 17. These acronyms have already been defined.

Page 4, equation 1. I think it would be useful to recap the accuracy of this relationship, and how well it works for the data that are presented here.

Page 4. Atmospheric fCO₂ calculation. What atmospheric pressure was used?

Page 4, decomposition equation. As mentioned above, please use the method that explicitly accounts for the effects of dilution/concentration. Note also, that the equation as written is wrong. The dX/dt term should be dX/dz , where z is the driver in question. You further need to explain what values you used for these sensitivities, and how they were derived.

Page 5, line 5. How many times were data excluded from box B because of SSS being outside the 34-35 range?

Page 5. Also, please consider the grouping of these regions for the trend analysis. From Figure 2 SSS and TA, E and D appears quite similar while C and B both appear different. My sense is that combining E and D, while keeping C and B separate might be the best approach.

Page 5, lines 10-19. Please collect this information in one section. Please also write that these are the reconstructed values in Fig. 3 (If I understand correctly the e.g. Jan/March values adjusted to February are the 'reconstructed' values in Fig 3).

Page 6, lines 2-3. As mentioned above I am not convinced that these are 'pluriannual' trends. You might be doing yourself a disfavor by splitting the data into three time periods. It might make more sense to look at the timeseries as a whole and instead look at anomalies from the long-term trend. In particular this should be done from 2001 onwards. Consider to relate anomalies to mixed layer depths similar to Fröb et al. (2019).

[Printer-friendly version](#)

[Discussion paper](#)



Page 6, lines 7 onwards. Here a panel showing air-sea fCO₂ difference, as suggested above would help, as it will make the discussion more quantitative.

Page 6, line 20. I think 'near-stagnation' is the wrong word here. In winter the increase appears significant, at 1 $\mu\text{atm}/\text{yr}$. Generally, please make the summary of the results quantitative.

Page 7, line 11. Remove 'Thereafter'

Page 7, line 15-16. Enclose 'much faster than the atmospheric signal', with commas. And... 'suggest larger productivity in the beginning of the period than at the end'. BTW this can and should be checked with Chl a data.

Page 7, line 19, replace 'than' with 'to'

Page 7, line 21-22. It is interesting that the slower increase in fCO₂ is associated with strengthening of the winds and enhanced deep mixing. As noted above, Metzl et al., (2010) suggested deep mixing as the cause for the larger increase in the 2001-2008 period. Recently Fröb et al. (2019) found anomalously high fCO₂ during years of deep mixing. What is suggested here, is thus at odds with these papers. This needs to be explored or revised.

Page 7, line 25. Set the '2' as subscript.

Page 8, line 9 -10. The link between the trends in the carbon system and NAO+AMV that are described here are not backed up with any statistics. It comes across as very speculative. The statements need to be backed up with for example correlation analyses.

Page 8, line 11. Fröb et al (2018) shows in particular a large increase of anthropogenic DIC inventory in deep mixing years, and tendencies for a loss of natural carbon. Fröb et al. (2019) shows a outgassing of CO₂ during deep mixing years. The coupling between the inventory changes and the variability in fluxes has yet to be made. Some discussion around this would be interesting.

[Printer-friendly version](#)[Discussion paper](#)

Page 9, line 5. 'Makes it difficult to predict the evolution of CO₂ uptake..' I suggest to read the paper by Li et al 2016, on decadal predictions of North Atlantic CO₂ uptake, this might provide some relevant information.

References (not cited in the paper) Fröb, F., Olsen, A., Becker, M., Chafik, L., Johannessen, T., Reverdin, G., and Omar, A.: Wintertime fCO₂ Variability in the Subpolar North Atlantic Since 2004, *Geophys Res Lett*, 46, 1580-1590, 2019.

Li, H. M., Ilyina, T., Muller, W. A., and Sienz, F.: Decadal predictions of the North Atlantic CO₂ uptake, *Nat Commun*, 7, 2016.

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2019-119>, 2019.

BGD

Interactive
comment

Printer-friendly version

Discussion paper

