

Dear editor,

We hereby submit a revision of the paper “Regulation of N<sub>2</sub>O emissions from acid organic soil drained for agriculture” (bg-2019-14) by Taghizadeh-Toosi et al. Major revisions were requested, and we are grateful to be granted the extension needed to complete this work.

Perhaps the most critical point raised was the assumptions about land use and site conditions that were necessary to test for these effects. In the revised statistical analyses, individual combinations of site and crop were instead tested separately to avoid such assumptions. Surprisingly, the effects identified in the previous analysis were more or less confirmed, indicating that the relationships were robust.

Review comments were overall constructive and helpful in clarifying many aspects. With the exception of some proposed reorganization of Figures 3-6 we believe that we have acted on all points raised, as specified below in the point by point response. In this light, we hope that it will now be possible to accept the manuscript for final publication.

Kind regards,

Arezoo Taghizadeh-Toosi

*Detailed response to reviewer comments (in blue):*

I raised several issues in my previous review, and thank the authors for their efforts at addressing them. Overall, I think the discussion is much improved. Unfortunately, I believe my concern regarding the experimental design needs to be further addressed. I do believe that revision is possible, and although it will likely require new statistical analyses, it shouldn't be overly difficult once the appropriate design is identified.

Below I list four general comments, of which the first two are most important, followed by responses to some of the authors' responses to my earlier comments, followed by specific comments. Apologies again for the rather verbose review.

*Response: We appreciate the reviewer's critical eye on the data analyses. We have made an effort to meet the different concerns, and in particular assumptions regarding experimental design have been revised, as explained below.*

#### GENERAL COMMENTS

1. One of my main concerns was regarding the robustness of the conclusions that can be drawn from this experimental design (Author response 17). The authors have removed the seasonality analysis, which I appreciate, but I believe that important issues remain. Some of the problem is in the language used to present and discuss results rather than in the underlying science, but science issues remain.

- This remains a study using only one year of data, though I don't think this is in itself a problem, it is a limitation. The authors argue in their response 18 that a previous 14-month study (Petersen et al. 2012) should also be considered, and that study does provide context for the results presented here (though note that any results from Petersen et al 2012 that are important for understanding the rationale for this study or for understanding its results should be summarized in this manuscript). However, Petersen 2012 is not part of the analysis, and the current manuscript doesn't provide any insight into the effects of interannual variability (also it seems that this was perhaps an abnormally warm and wet year). That's not a problem--I think the authors have done a good job of removing generalizing statements about seasonality from the revised manuscript--but I do think that a statement noting that the analyses of spring and fall are based on data from a single year should be included in the conclusion as part of an acknowledgment of potential limitations of the study.

*Response: There are five references to the study of Petersen et al. 2012 in the Introduction, and we have added additional detail to explain the context:*

*"Among sites with arable crops in three regions, two sites had N<sub>2</sub>O emissions corresponding to, respectively, 38 and 61 kg N ha<sup>-1</sup> yr<sup>-1</sup>; both of these sites showed distinct seasonal patterns with the highest emissions in spring and autumn periods, whereas emissions at the third site were lower (6.4 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>) and much less variable. Notably, WT depth at the two sites with seasonal patterns of N<sub>2</sub>O emission fluctuated between 10-30 and 10-120 cm depth, whereas WT depth at the third site remained at 90-125 cm depth throughout the 14-month monitoring period." (l. 53 in cleaned manuscript version)*

*Also, we now begin the conclusion by stressing that this is a one-year study:*

” In this one-year study, N<sub>2</sub>O emissions were...” (l. 600 in cleaned manuscript version)

- This remains a study that lacks traditional replication of treatments, and I think this issue requires substantial additional revision.

- o The authors argue that RG1 and AR1 should be considered as independent units because they differ in land use history, and were treated as such in Petersen et al. 2012. I’m afraid I’m not convinced—the design is an illustration of the traditional definition of a split plot: two adjacent fields with different treatments (or, as in this case, treatment histories), but a shared geographic location, and thus conditions that have been formed by a history of shared state factors (*sensu* Hans Jenny). As such they are simply not independent sites. The map in figure 1a, which presents AR1/RG1 as a single site, illustrates the point rather well, and as I understand, sampling positions in the two fields were only 10-20m from one another. One of the clearest issues here is the nearly identical temporal patterns of the water table depth—the kind of shared environmental variable that split plot analyses were created to accommodate. I think it’s fine for the authors to describe patterns at each of the RG and AR locations, but I think the experimental design makes any formal comparison between land uses impossible. If the authors want to examine site differences, I think it would have to be done using an *n* of 3, treating RG1/AR1 as a single site with higher within-site replication, but it would be a good idea to consult a statistician—there may be other options. Because of this issue, I think the graphical analysis may also need to be redone—perhaps organized around sites rather than land uses—if I am correct in my understanding that independence is an underlying assumption (as I mentioned in my previous review, I am not an expert in graphical analysis). In any event, I recommend consulting with a statistician before proceeding with additional revisions.

*Response: We acknowledge that site conditions and distribution of treatments represent factors that warrant great caution in data interpretation, and we have indeed acted to meet the concern raised here. Note that the effect on N<sub>2</sub>O emission of the fertilisation treatment was not the same at the four sites; therefore, one of the greatest advantages of using split plot designs (estimating a common effect of the fertilization treatment) vanishes. In the revised analyses we have instead used a single model describing the effect of fertilisation for each site. Therefore, we now report the N<sub>2</sub>O emissions separately per site (both for the temporal dynamics and the cumulative emissions). We believe that this approach documents the results of our experiment better. The conclusions reached previously still stand, with a few modifications which are described.*

- o A related note: the very big site differences between RG2 and AR2 (e.g., in carbon stocks) highlight the rationale for using split plot approaches (such as the one the authors use for their fertilizer treatments): such approaches avoid treatment or land use history differences being confounded with site differences.

*Response: See previous comment. In the revised Table 2, results are now presented for each combination of site, crop, fertiliser treatment and season (n=8).*

- o The authors also argue that differences between land use was evident without any statistical support, which is a fair argument. I think it’s fine to discuss possible

effects of land use qualitatively, but important to include and clearly highlight the caveat that the study lacks independent replicates for the land use treatments.

*Response: We have re-analysed cumulative N<sub>2</sub>O emissions for individual site-crop combinations by season (see revised Table 2). Results are now expressed as average daily rates, which allowed us to test for differences during spring and autumn, respectively. Furthermore, the temporal dynamics were analysed with a suitable mixed model. Results are now presented and discussed with reference to statistical support where possible, and otherwise wording has been changed to be more qualitative where relevant.*

o I don't think these issues mean that the graphical model cannot be applied or provide novel results, but the model needs to be applied using the appropriate experimental design, and can't treat RG1 and AR1 as independent sites.

*Response: As for cumulative N<sub>2</sub>O emissions, graphical models were reanalysed for individual site-crop combinations by season, see revised Fig. 7 (now split into Figure 7 (spring) and 8 (autumn)). Notably, even without the statistical power of treatment replication, the effects remained largely unchanged with respect to treatment differences.*

o In Response 17, the authors note that land use was not analyzed statistically. However, the generalized linear mixed model includes land use ("crop") in the interaction term used as the fixed effect (line 248-249), which is inappropriate given the replication concerns regarding RG1 and AR1.

*Response: In the revised analysis, land use treatments were analysed separately.*

2. This leads me to the language presenting results. The qualitative results need to be more clearly communicated as such—the way many results are presented here implies that there has been a quantitative statistical analysis conducted. A very simple example is line 328: "Mineral N concentrations were greater at AR1 compared to AR2". In the scientific literature, such a statement implies a significant statistical difference in mineral N concentrations between the two sites. If a statistical test was conducted, a P value should be included (in general, there should be much more inclusion of p values in the text). If it wasn't, small changes to the language need to be made that make that clear, e.g., "Mineral N concentrations appeared to be greater at AR1 compared to AR2". Another small example might be line 334-35: "There was variation at depth in the soil, which could not be explained by fertilization." The phrase "could not be explained" implies that a statistical test was attempted, but was not significant. If, instead, this is qualitative interpretation of data, it should include more qualifying language, e.g., "There was variation at depth in the, which did not appear to be related to fertilization."

In fact, there are numerous instances of this kind of presentation of results throughout the manuscript; here are just a few examples (there are many more involving site comparisons, temporal comparisons within sites, fertilizer effects, etc., and all should be addressed):

L316: "There were only minor differences. . . between seasons" implies that seasonal differences were tested; can be changed to something like "If there were any differences, they were likely minor. . ."

L 326: "The residence time for mineral N in the soil solution was generally longer at AR compared to the RG sites" implies that site differences were tested; can be changed to

“The residence time for mineral N in the soil solution generally appeared to be longer at AR compared to the RG sites”

L328: “Mineral N concentrations were greater at AR1 compared to AR2” implies that site differences were tested; can be changed to “It appeared that mineral N concentrations may have been greater at AR1 than AR2”

L 329: “Fertilisation increased  $\text{NH}_4^+$ -N and  $\text{NO}_3^-$ -N concentrations” implies that differences between fertilized and unfertilized split plots were tested. Can be changed to “It appeared as though  $\text{NH}_4^+$ -N and  $\text{NO}_3^-$ -N concentrations increased following fertilization”

One quick way around this issue would be to state in the methods section that all interpretations of results are qualitative unless accompanied by a p value, and then to include p values where statistical tests were conducted (in any event, I encourage the authors to include P values for all statements reflecting the results of a statistical test). But a clearer solution would be 1) to test differences statistically if they have not been and include p values, and 2) to change the language describing results to make it qualitative, using words and phrases such as “may,” “appears to,” “might,” etc. to describe apparent patterns that cannot be tested or aren’t important enough to test. I don’t think it’s necessarily a problem to present results qualitatively (though for any results that are important to the central hypotheses being tested, it’s always good to include statistical analyses), but it needs to be very clear when differences were tested and when they were not.

*Response: Regrettably, with limited resources we had made the decision to not collect and analyse soil samples at the block level, but instead pool soil samples from each block (by depth interval) for analysis of mineral N. For this reason, we have been unable to formally test effects of fertilisation or temporal dynamics.*

*We agree with the reviewer that the text should have reflected this more clearly. We have adjusted wording throughout the manuscript and further stress the qualitative nature of the mineral N dynamics with the following sentence:*

*“Since subsamples were pooled for analysis, only a qualitative description of the effects of treatments and temporal dynamics is possible.” (l. 346 in cleaned manuscript version)*

3. I think there could be more detail in the discussion of figures 3-7, which represent a lot of the data presented in this manuscript. Briefly, the text often invokes a relationship between the water table and a)  $\text{N}_2\text{O}$  concentrations in the soil or b)  $\text{N}_2\text{O}$  fluxes at the surface, but the relationships described in the text are not obvious to me from examining the figures—the spatiotemporal relationship between the water table depth and  $\text{N}_2\text{O}$  concentrations seems quite different in different sites and seasons. Additional text walking the reader through the authors’ interpretation of these figures would be helpful. I made related specific comments in my last review, and detail some points below in my response to the authors’ response #46, as well as in specific comments for lines 494, 501, and thereabouts.

*Response: In the presentation of results, we gave priority to the comparison of land uses at the four sites, which is why fluxes and subsurface  $\text{N}_2\text{O}$  concentrations (and seasons) are presented separately. We have revised the description of these results to better link observations to specific treatments or days, and to better link fluxes with subsurface  $\text{N}_2\text{O}$  dynamics.*

*However, due to the delay in gas exchange through saturated and/or highly tortuous peat soil (a point now stressed in the Discussion l. 489f in cleaned manuscript version), a close relationship between fluxes*

*and subsurface concentrations can not be expected. We have added a new paragraph to the discussion explaining this (see below).*

4. It seems to me that the two conclusions that can be clearly reached from this study are the rejection of the hypothesis that that FeS<sub>2</sub> oxidation coupled with NO<sub>3</sub><sup>-</sup> reduction was an important driver of N<sub>2</sub>O emissions, and the identification of the capillary fringe as an important predictor of surface N<sub>2</sub>O fluxes. If the authors agree, the conclusion should be revised; as it currently stands, the second of these findings is not discussed at all in the conclusion, and the description of the first conclusion gives the reader the impression that the hypothesized mechanism is possibly and maybe even probably not trivial.

*Response: N mineralisation from decomposing peat after WT drawdown was mentioned, but we now specifically refer to the capillary fringe. With respect to FeS<sub>2</sub>, we have strengthened the wording:*

*“The hypothesis that NO<sub>3</sub><sup>-</sup> reduction coupled with FeS<sub>2</sub> oxidation was an important source of N<sub>2</sub>O could not be confirmed.” (l. 603 in cleaned manuscript version)*

*Also, the Discussion now includes a proper reference to the laboratory study confirming this conclusion – this study has now been published ([doi.org/10.1080/01490451.2019.1666192](https://doi.org/10.1080/01490451.2019.1666192)).*

Minor comment: the response to reviewers letter was sloppy – a number (maybe all?) of the line numbers of revised text referred to in the response to reviewers are incorrect, there are typos in the quoted text revisions (e.g., response 50), and sometimes there’s no indication of what revisions were made or where they can be found in the text (e.g., response 49). I empathize with these kinds of errors, especially after getting a fresh batch of comments from a new reviewer late in the publication process, but anything you can do to make the job of the reviewers easier is appreciated.

*Response: An annotated copy with tracked changes was in fact submitted together with the revision, and we specifically referred to in the response letter. The critical comments above therefore appears to be based on a misunderstanding.*

Responses to specific author responses

Response 20: I think the authors missed my point here, which was that the manuscript seems to only conduct a qualitative analysis of the relationship between changes in water table depth and changes in surface N<sub>2</sub>O flux, rather than including a statistical analysis of the relationship between the two variables that could provide quantitative insight into the importance of variation in water table depth for N<sub>2</sub>O surface fluxes. I agree that the manuscript does provide a graphical analysis showing that capillary fringe N<sub>2</sub>O is the only significant predictor of N<sub>2</sub>O surface fluxes at several sites, but being statistically significant and being important are not necessarily the same thing, and the graphical analysis does not include any analysis of the relationship between water table depth per se and N<sub>2</sub>O. I don’t think that the authors need to do what I am suggesting here for the manuscript to be publishable, but it seems like a missed opportunity.

*Response: We did not test possible direct effects of WT depth on N<sub>2</sub>O flux. However, WT depth was indirectly, in the analysis of N<sub>2</sub>O flux patterns with graphical models, where N<sub>2</sub>O concentration above the WT table carried information about WT depth at the time of sampling. As pointed out by the reviewer in a comment to the earlier Response no. 46 (see below), in early autumn the N<sub>2</sub>O fluxes were greatly*



*stimulated at a time where WT depth was still low, presumably due to wetting of the upper soil layers in which nitrate had accumulated. This exemplifies that WT depth per se may not be a strong predictor of emissions.*

Response 21: Thank you for the response, though part of my concern was not addressed. This concern is largely related to my general comment about the language used to describe results. My concern here had been that there were multiple instances in the manuscript (I cited them in the original comment as including lines 380 and lines 411-412) where temporal variation in surface N<sub>2</sub>O flux at a site was attributed to fertilizer effects, or where fertilizer effects were excluded as a cause of variation in surface N<sub>2</sub>O flux. However, the relationship between temporal variability in surface N<sub>2</sub>O fluxes and fertilizer applications appears not to have been tested statistically.

*Response: We have analysed the temporal dynamics of N<sub>2</sub>O emissions, as well as WT depth. Information about statistical significance has been included where relevant.*

Response 25: The authors argue that “The graphical model results (Figure 7) did show increasing N<sub>2</sub>O emissions with declining, as well as increasing WT depth that depended on soil N status.” I do not think that the graphical analysis included a water table depth variable?

A separate point: I also don’t understand what is intended by “increasing WT depth that depended on soil N status” but if the point is that N<sub>2</sub>O<sub>WT</sub> depends on soil N status, that would be an interesting result that deserves more discussion. If instead it’s referring to AR-autumn, where the N<sub>2</sub>O flux is determined in part by soil N status, then ignore this second comment of mine.

*Response: This wording was not too well-considered. In fact the effect of WT depth was indirect, by shifting the position and magnitude of soil volume (in the capillary fringe) with a potential for N<sub>2</sub>O production. It is true, as stated, that we were thinking about the NO<sub>3</sub> accumulation and subsequent wetting of the top soil. No changes made.*

Response 39: I’m glad this was caught. I’d encourage the authors to carefully review the script for producing all figures again if they haven’t already.

*Response: All Figures were double-checked.*

Response 46: I wonder whether this explanation may also hold for DOY 252 and 259: it looks to me as though there are elevated N<sub>2</sub>O concentrations in surface soils, and no clear connection to the capillary fringe. Wouldn’t that suggest a topsoil source for all of the highest surface fluxes at AR1 during Autumn—a finding supported by the graphical analysis? The text implies that the high WT is responsible for the high emissions for DOY 252 and 259 (“The subsequent decline in N<sub>2</sub>O emissions at AR sites coincided with WT withdrawal”), but I would think that anoxia in topsoil related to elevated precipitation during this period could be a more likely explanation. The variability in topsoil N<sub>2</sub>O concentrations looks fairly physically separated from the water table dynamics, and if capillary fringe is not a significant predictor for surface N<sub>2</sub>O fluxes at this site during autumn, why invoke a relationship between WT withdrawal and surface N<sub>2</sub>O emission declines?

I have further questions about the interpretation of the subsoil N<sub>2</sub>O concentration dynamics discussed in my comment on line 494 below.

*Response: We largely agree with the interpretation presented above, and the reference to WT dynamics did not imply a causal relationship. We have modified the text to put more emphasis on the wetting of the soil:*

*“High fluxes were observed on the first day of this monitoring period, DOY 246, at a time where WT depth was still at 40 to 80 cm depth. Instead the high N<sub>2</sub>O fluxes may have been triggered by saturation of the top soil after 10 and 22 mm rainfall the previous two days. Additional rainfall during the following days then was accompanied by a rise in WT. The subsequent decline in N<sub>2</sub>O emissions at AR sites coincided with WT draw-down and drainage of the top soil.” (l. 437 in cleaned manuscript version)*

Response 54: I think it's a good move to consider N<sub>2</sub>O as a time-integrated measure. My previous concern here was very much regarding the limitations of using point measurements of NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> to infer N transformation rates. I think the framing of the revised discussion addresses my concerns.

*Response: Thank you.*

Response 56: So the data presented in figure 2 are means (and standard errors) of intact cores containing either fertilized and unfertilized soils in RG1, but for AR1, it's a mean of intact cores containing only unfertilized soils? I don't understand the rationale for presenting the data this way (instead of, for example, presenting fertilized and unfertilized soils separately where appropriate). At the very least, the caption should clearly detail this odd fertilization treatment situation.

*Response: The management of grassland required that fertilisation took place before we had an opportunity to collect intact soil cores at the remote field site. Subsequent inspection of NO<sub>2</sub><sup>-</sup> results upon sampling one week after fertilisation, however, did not suggest any effect of fertilisation, and this was confirmed by a paired-sample t-test across the six depths. We therefore went ahead and presented soil profiles across both fertiliser treatments.*

*As recommended, we have revised the Figure caption to describe this procedure.*

Response 57: It's your decision, but there isn't much fine-scale temporal variation in these figures; I'd think it's worth graphing up the full year and seeing what it looks like.

*Response: Whether seasons or land use treatments should be shown side by side is not an easy decision. However, since grassland sites, and arable sites, are mostly discussed together, we wanted to facilitate a direct comparison with this side by side presentation. Although a case could also be made for other solutions, we have decided not to change the format of Figures 3-6.*

Response 59: Again, it's your decision, but would think it's worth including in SI as figures as well as data tables—it's just very difficult to read patterns in a data table. But thank you for including the raw data for researchers who may be interested in using it for their analyses.

*Response: Thank you for this suggestion. Given the many variables involved (sites, crops, fertiliser treatments, soil depths and N species), we believe graphic presentations could be either too busy or too disaggregated to have much added value, and therefore we prefer to focus on the Table format.*

New specific comments:

Line 23: perhaps define capillary fringe here: “...in the capillary fringe— [definition here]—was.



. . “

*Response: Done – the following text was added:*

“(i.e, the soil volume above the water table influenced by tension saturation)” (l. 23 in cleaned manuscript version)

Line 62: it would be helpful to readers if you could provide an explicit definition of capillary fringe here, given how important it is to the manuscript.

*Response: text was added (l. 67) to stress that this zone is defined by capillary rise of groundwater, but that it can still be partly unsaturated depending on pore size distribution:*

“The capillary fringe of organic soils represents an interface between saturated and unsaturated soil conditions, in which the extent of tension saturation depends on pore size distribution (Gillham, 1984).” (l. 68 in cleaned manuscript version)

Line 105: what ridges? Probably need to provide more agronomic details

*Response: It has been specified that these are established around potato rows.*

Line 124: Perhaps this should be “method of fertilizer application”? (as N fertilization was noted as an exception in the next sentence)

*Response: OK, has been included to clarify this distinction. (l. 131 in cleaned manuscript version)*

Line 127: I believe the erroneous NS fertilizer application in RG2 was made on the same date as the second slurry application; if correct I would clarify that fact by starting this sentence with “On the same date as the second slurry . . . “ or “Immediately after the second slurry. . . “ If the NS application was made on a different date, indicate the date.

*Response: Done.*

Line 137: here it would be helpful to explain the choices of spring and fall as emerging from the patterns observed in Petersen 2012, and in the introduction briefly describe those patterns. It could be just 1-2 sentences in the introduction, and an introductory clause to the first sentence here.

*Response: The following was added to the Introduction, and is referred to in section 2.4, as requested:*

“Among sites with arable crops in three regions, two sites had N<sub>2</sub>O emissions corresponding to, respectively, 38 and 61 kg N ha<sup>-1</sup> yr<sup>-1</sup>; both of these sites showed distinct seasonal patterns with the highest emissions in spring and autumn periods, whereas emissions at the third site were lower (6.4 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>) and much less variable. Notably, WT depth at the two sites with seasonal patterns of N<sub>2</sub>O emission fluctuated between 10-30 and 10-120 cm depth, whereas WT depth at the third site remained at 90-125 cm depth throughout the 14-month monitoring period.” (l. 53 in cleaned manuscript version)

Line 144: it would give a more complete picture if the dates of these exceptions were enumerated here.

*Response: The following was added:*

“With a few exceptions (DOY 169 and 265 at site *RG1*, DOY 132 and 314 at site *AR2*), each campaign was initiated between 9:00 and 12:00; the order of sites visited in each trip alternated from week to week.” (l. 151 in cleaned manuscript version)

Line 165: revise to “...at -20C until analysis, described in section 2.4.5.”

*Response: Done.*

Line 170-1: I’m afraid this sentence is hard to follow. It reads as though the diluted soil gas was first transferred to the exetainer, and then from the exetainer to the glass syringe. Perhaps also specify whether exetainers were filled with 10ml gas as they were in the chamber measurements.

*Response: Section 2.4.3 has been edited to clarify the procedure. (l. 176 in cleaned manuscript version)*

Line 174: As a general rule I’d argue it’s best to explicitly include all applicable methodological detail, rather than referring readers to an additional manuscript (within reason). In this case, my institution does not have a subscription to the European Journal of Soil Science, so my library had to obtain 24-hour access to Petersen 2014 so I could see the dilution calculations; I would include them in the manuscript.

*Response: Equation 5 from Petersen 2014 is now shown and explained in section 2.5.*

Line 240-245: it would be helpful to specify if/how field temperature and pressure corrections were made to obtain surface flux estimates.

*Response: The following sentence was added:*

“Nitrous oxide mixing ratios were converted to units of mass per volume using the ideal gas law and values of pressure and air temperature recorded by the weather station.” (l. 257 in cleaned manuscript version)

Line 297: It should be noted here that these characterizations were conducted after the fertilizer treatment of RG1

*Response: We did not locate the text referred to above, but we have assumed that it was soil N characterisation. We have added the following sentence:*

“At site *RG1*, fertilisation had taken place one week earlier, but a two-sample *t* test did not find evidence for an effect on NO<sub>2</sub><sup>-</sup> availability (*p* = 0.19), and therefore the results from both *AR1* and *RG1* subplots are presented together.” (l. 355 in cleaned manuscript version)

Line 380: Instead of “independent of fertilization” which suggests a formal analysis, how about “this variation appeared to be broadly similar between fertilized and unfertilized subplots”

*Response: OK, paragraph has been revised.*

Line 394 and following: this is one of many examples related to my general comment on presenting qualitative results, but the generalized linear mixed model appears to include only time, and not fertilizer, as a factor in the fixed effect, so statements regarding the effects of fertilization should be made qualitatively (e.g., rather than “with no effect of fertilizer

amendment,” use something like “and did not appear to vary in response to fertilization.” Similarly, “no effect of N fertilisation was observed. Hence, the higher emissions were associated with site differences other than fertilization” could be changed to “fertilisation did not appear to influence N<sub>2</sub>O fluxes. Hence, the higher emissions were likely associated with site differences other than fertilization”). The paragraph starting with line 406 does a good job of qualitatively describing results.

*Response: We have revised the text to make sure that results are presented in a qualitative way where necessary.*

Line 467: change to “In accordance with this effect of rewetting...” (in general, it’s good to give “this” an object to make it clear what is being referred to)

*Response: Done.*

Line 490: here and maybe elsewhere there are still instances where time of year are referred to without DOY (here parenthetical DOYs would be particularly helpful since its referring to figures using DOY on the x-axis). In addition, please specify that this accumulation is in the fertilized plots of RG2.

*Response: OK, several instances were amended with DOY information.*

Line 491: change “significant” to “substantial” if the meaning is “a lot.” Avoid using “significant” except when alluding to the result of a statistical test.

*Response: Done.*

Line 492: if the contrast is that in RG2 there was only accumulation at certain times and depths but accumulation was everywhere and all the time in the AR sites, change “..accumulation of N<sub>2</sub>O in the soil” to “...accumulation of N<sub>2</sub>O across all soil depths throughout the spring” or something similar. Maybe include a reference to Figure 4.

*Response: OK, the following wording is now used:*

“In contrast, at AR sites N<sub>2</sub>O consistently accumulated in the soil throughout spring” (l. 536 in cleaned manuscript version)

\*Line 494: as was the case with the previous version of the manuscript (see my original comment that elicited the authors’ response 46 and my general comment 3), I’m not sure I see the pattern so clearly. The location of elevated N<sub>2</sub>O above 40cm in AR2 is attributed to the water table being higher than in AR1, where elevated N<sub>2</sub>O tends to be in deeper soils, and the authors argue that this points to a capillary fringe source for N<sub>2</sub>O. But in RG2 the water table is higher than it is in RG1 during the spring, but the N<sub>2</sub>O distributions are reversed: they are higher at depth in in RG2, where the water table is higher, and higher in shallower soils in RG1, where the water table is lower. The fertilized AR2 also has an inversion of N<sub>2</sub>O concentrations during spring that seems entirely unrelated to water table depth. In autumn, the N<sub>2</sub>O at depth declines as the water table depth increases, and the patterns in the two blocks in RG1 seem somewhat opposite of one another even though the water table is just about identical. With so much apparent variation in how N<sub>2</sub>O concentrations vary with the water table, how do these results all

relate to capillary fringe as the N<sub>2</sub>O source? What am I missing? I suppose that this comment, in combination with my new response to the authors response 46, suggests an opportunity for the authors to revise and expand their presentation (and perhaps interpretation) of these results. It would be a very satisfying read if the paper spent a little more time on the subtleties in the links the graphical analysis identification of the significant drivers of surface N<sub>2</sub>O fluxes to the spatiotemporal dynamics presented in figures 3-6 (these are most of the data being presented in the paper, so feel free to give them more attention in the text).

*Response: We agree that patterns of soil N<sub>2</sub>O concentration profiles were complex and not readily aligned with emissions. It is probably important to acknowledge the importance of soil wetness as a barrier towards transport, and we have added a new paragraph to the discussion (section 4.1) to motivate the focus on N<sub>2</sub>O concentration above the water table, which was a potential site of N<sub>2</sub>O production:*

“A limited number of potential drivers (rainfall, temperature, soil mineral N and soil concentrations of N<sub>2</sub>O) were monitored to help explain N<sub>2</sub>O emission dynamics. Soil N<sub>2</sub>O concentration profiles showed complex patterns where, for example, the highest concentrations were sometimes observed above, and sometimes below the WT depth at both RG (Figure 3) and AR sites (Figure 4). Fertilisation in spring was associated with higher concentrations of N<sub>2</sub>O below the WT depth at sites RG2 and AR2, which indicated downward transport of fertiliser N, but this was not reflected in elevated N<sub>2</sub>O emissions. The reason may be that in wet soil the time required to reach a steady state between N<sub>2</sub>O production and emissions from the soil surface can be significant and increases with distance (Jury et al., 1982). In accordance with this, Clough et al. (1999) observed a delay of 11 days before <sup>15</sup>N enriched N<sub>2</sub>O produced at 80 cm depth was released from the soil surface at a corresponding rate. Since the presence of air-filled porosity is critical for the exchange of gases between soil and the atmosphere (Jury et al., 1982), the soil N<sub>2</sub>O concentration closest to, but above the WT depth (N<sub>2</sub>O<sub>WT</sub>), was taken to represent “subsoil” processes stimulating N<sub>2</sub>O emissions.” (l. 484 in cleaned manuscript version)

Line 501: I’m not sure why water table dynamics are invoked in the discussion of N<sub>2</sub>O emissions at AR1, if capillary fringe was not a significant predictor of surface N<sub>2</sub>O fluxes in the graphical analysis, and, as noted previously (and in my comment on the authors’ response 46), surface fluxes were already elevated before the rise in WT depth? A simpler explanation consistent with the graphical analysis might be that topsoil was the source of surface N<sub>2</sub>O fluxes in AR1 during autumn, apparently independent of water table depth. If this discussion is intended to be descriptive only of dynamics within the soil (and does not have any relation to surface N<sub>2</sub>O emissions), that needs to be made clear.

*Response: We agree, and we have adjusted the text accordingly, as explained above.*

Line 528: It would be helpful to have a definition of “dead-end pores”

*Response: These are associated with plant cell remains. This information has been added.*

Line 567-559: These are all the kinds of statements that need to be made much more qualitatively in the absence of clear statistical support, and issues with experimental design and strength of statistical support for conclusions need to be highlighted in the conclusion.

*Response: The Conclusion section has been rewritten to reflect where statistical support for statement was available, and to reflect better this is our interpretation of the results.*

Line 560: remove the comma between “hypothesis” and “that”

*Response: Done – and sentence edited.*

Line 561: my impression from the results is that NO<sub>3</sub><sup>-</sup> reduction coupled with FeS<sub>2</sub> reduction is likely to be a trivial mechanism of N<sub>2</sub>O production in these soils (a conclusion apparently supported by the related manuscript that has been submitted elsewhere)—this statement makes it sound as though it could be non-trivial.

*Response: We agree that this conclusion should be expressed more clearly. Has been rephrased as:*

“The hypothesis that NO<sub>3</sub><sup>-</sup> reduction coupled with FeS<sub>2</sub> oxidation was an important source of N<sub>2</sub>O was not confirmed.” (l. 603 in cleaned manuscript version)

# Regulation of N<sub>2</sub>O emissions from acid organic soil drained for agriculture

Arezoo Taghizadeh-Toosi<sup>1</sup>, Lars Elsgaard<sup>1</sup>, Tim J. Clough<sup>2</sup>, Rodrigo Labouriau<sup>3</sup>, Vibeke Ernstsén<sup>4</sup> and Søren O. Petersen<sup>1</sup>

<sup>1</sup> Department of Agroecology, Aarhus University, Tjele, Denmark

<sup>2</sup> Faculty of Agriculture and Life Sciences, Lincoln University, Christchurch, New Zealand

<sup>3</sup> Applied Statistics Laboratory, Department of Mathematics, Aarhus University, Aarhus, Denmark

<sup>4</sup> Geological Survey of Denmark and Greenland, Copenhagen, Denmark

Correspondence to: Arezoo Taghizadeh-Toosi (Arezoo.Taghizadeh-Toosi@agro.au.dk)

## Abstract

Organic soils drained for crop production or grazing land are agroecosystems with potentially high, but variable emissions of nitrous oxide (N<sub>2</sub>O). The present study investigated the regulation of N<sub>2</sub>O emissions in a raised bog area drained for agriculture, which ~~was~~<sup>is</sup> classified as potentially acid sulfate soil. ~~Here, we~~<sup>We</sup> hypothesised that pyrite (FeS<sub>2</sub>) oxidation was a potential driver of N<sub>2</sub>O emissions through microbially mediated reduction of nitrate (NO<sub>3</sub><sup>-</sup>). Two sites with rotational grass, and two sites with a potato crop, were equipped for monitoring of N<sub>2</sub>O emissions and soil N<sub>2</sub>O concentrations at 5, 10, 20, 50 and 100 cm depth during weekly field campaigns in spring and autumn 2015. Further data acquisition included temperature, precipitation, soil moisture, water table (WT) depth, and soil NO<sub>3</sub><sup>-</sup> and ammonium (NH<sub>4</sub><sup>+</sup>) concentrations. At all sites, the soil was acidic with pH ranging from 4.7 to 5.4. Spring and autumn monitoring periods together represented between 152 and 174 d, with cumulative emissions of ~~3-64-5~~ kg N<sub>2</sub>O-N ha<sup>-1</sup> at sites with rotational grass and ~~19-21-20-50~~ kg N<sub>2</sub>O-N ha<sup>-1</sup> at sites with a potato crop. Equivalent soil gas phase concentrations of N<sub>2</sub>O ~~mostly~~<sup>mostly</sup> ranged ~~from up to~~<sup>from up to</sup> around ~~4015~~<sup>4015</sup> μL L<sup>-1</sup> at grassland sites, ~~and up~~<sup>and up</sup> to several hundred μL L<sup>-1</sup> at potato sites, in accordance with lower soil mineral N ~~concentrations~~<sup>availability</sup> at grassland sites. Statistical analyses using graphical models showed that soil N<sub>2</sub>O concentration in the capillary fringe (~~i.e., the soil volume above the water table influenced by tension saturation~~) was the strongest predictor of N<sub>2</sub>O emissions in spring and, for grassland sites, also in the autumn. For potato sites in autumn, ~~the analysis found~~<sup>there was evidence</sup> that NO<sub>3</sub><sup>-</sup> availability in the top soil, ~~together with~~<sup>and</sup> temperature, were the main controls on N<sub>2</sub>O emissions. Chemical analyses of intact soil cores ~~from 0-1 m depth~~<sup>collected to 1 m depth</sup> at adjacent grassland and potato sites ~~in spring and autumn~~, showed that the total reduction capacity of the peat soil (assessed by cerium (IV) reduction) was much higher than that represented by FeS<sub>2</sub>, and the concentrations of total reactive iron (TRFe) were higher than those of FeS<sub>2</sub>. Based on the statistical graphical models and the tentative estimates of reduction capacities, FeS<sub>2</sub> oxidation was ~~found~~<sup>unlikely</sup> to be important for N<sub>2</sub>O emissions. Possible pathways of N<sub>2</sub>O production in spring and autumn periods, and the potential sources of N, are further discussed.

**Key words:** Drained peat, potentially acid sulfate soil, rotational grass, potato, nitrous oxide, reactive iron




## 35 1 Introduction

Worldwide, 25.5 million ha of organic soils have been drained for agricultural use, mainly as cropland (Tubiello et al., 2016), and this accelerates decomposition of soil organic matter and net carbon (C) and nitrogen (N) mineralisation above the water table (WT) (Schothorst, 1977). Drained organic soils are significant net sources of greenhouse gas (GHG) emissions as carbon dioxide (CO<sub>2</sub>) and nitrous oxide (N<sub>2</sub>O) (Goldberg et al., 2010; Maljanen et al., 2003). A recent supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories on Wetlands (IPCC, 2014) proposed average annual emission factors of 4.3 and 8.2 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup> for temperate grassland on drained organic soil with low and high nutrient status, respectively, and an emission factor of 13 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup> for cropland. For soil C losses, the emission factors proposed for ~~these~~ three land use categories were between 5.3 and 7.9 Mg CO<sub>2</sub>-C ha<sup>-1</sup> yr<sup>-1</sup> (Hiraishi et al., 2014). Thus, while CO<sub>2</sub> emissions are overall more important, site conditions appear to be more critical for N<sub>2</sub>O.


Site conditions are defined by land use, management, inherent soil properties and climate (Mander et al., 2010; Leppelt et al., 2014). Both WT drawdown (Aerts and Ludwig, 1997) and WT rise (Goldberg et al., 2010) may enhance N<sub>2</sub>O emissions, but such effects depend on soil N status (Martikainen et al., 1993; Aerts and Ludwig, 1997). Maljanen et al. (2003) found that WT, CO<sub>2</sub> emissions and temperature at 5 cm depth explained 55% of the observed variability in N<sub>2</sub>O emissions during a two-year field study on a drained organic soil, whereas the response to N fertilisation was limited, and they suggested that N released by soil organic matter mineralisation was the main source of N<sub>2</sub>O. In a study comparing GHG emissions from organic soil with different land uses in three regions of Denmark (in total eight site-years), Petersen et al. (2012) also found that site conditions such as WT, pH and precipitation contributed significantly to explain N<sub>2</sub>O emission dynamics. Among sites with arable crops in three regions, two sites had N<sub>2</sub>O emissions corresponding to, respectively, 38 and 61 kg N ha<sup>-1</sup> yr<sup>-1</sup>; both of these sites showed distinct seasonal patterns with the highest emissions in spring and autumn periods, whereas emissions at the third site were lower (6.4 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>) and much less variable. Notably, WT depth at the two sites with seasonal patterns of N<sub>2</sub>O emission fluctuated between 10-30 and 10-120 cm depth, whereas WT depth at the third site remained at 90-125 cm depth throughout the 14-month monitoring period.

~~In the study by Petersen et al. (2012), extremely high N<sub>2</sub>O emissions corresponding to 38 and 61 kg N ha<sup>-1</sup> were observed from arable sites in two of the three regions investigated.~~ Several processes can lead to N<sub>2</sub>O formation in acid organic soil: biotic processes include ammonia (NH<sub>3</sub>) oxidation to nitrite (NO<sub>2</sub><sup>-</sup>) by archaea or bacteria (Herrmann et al., 2012; Herold et al., 2012; Stieglmeier et al., 2014), as well as nitrifier denitrification and heterotrophic denitrification by bacteria or fungi (Liu et al., 2014; Maeda et al., 2015; Wrage-Mönnig et al., 2018). The recently discovered process comammox by Nitrospira sp. is also a potential, but probably minor, source of N<sub>2</sub>O (Kits et al., 2019; Palomo et al., 2019). Abiotic N<sub>2</sub>O production can occur through chemodenitrification (Van Cleemput and Samater, 1996; Jones et al., 2015) or abiotic codenitrification (Spott et al., 2011). The two regions showing extreme N<sub>2</sub>O emissions from arable soil had both developed from marine forelands and were categorised as potentially acid sulfate soil, i.e., saturated to poorly drained soil containing pyrite (FeS<sub>2</sub>) that, upon oxidation, may lead to acid production in excess of the soil's

neutralising capacity (Madsen and Jensen, 1988). The capillary fringe of organic soils represents an interface between saturated and unsaturated soil conditions, and in which the extent of tension saturation depends on pore size distribution (Gillham, 1984). Previously, it was has been speculated that oxidation and reduction of iron sulfides could have interacted with influence N transformations during periods with changing groundwater level (Petersen et al., 2012). Drainage will promote promotes oxidation of FeS<sub>2</sub>, a process which may be linked to microbially mediated nitrate (NO<sub>3</sub><sup>-</sup>) reduction (Jørgensen et al., 2009; Torrento et al., 2010). The complete reduction of NO<sub>3</sub><sup>-</sup> to dinitrogen (N<sub>2</sub>) can proceed as follows:



However, in the capillary fringe residual oxygen (O<sub>2</sub>) or, alternatively, the acidification produced by FeS<sub>2</sub> oxidation, could favour incomplete denitrification with accumulation of the intermediate N<sub>2</sub>O (Torrento et al., 2010):



Nitrate reduction *via* the reaction described in Eq. 2 could potentially have contributed to the very high N<sub>2</sub>O emissions reported by Petersen et al. (2012) previously from ~~the~~ two arable sites, where groundwater sulfate concentrations were also consistently high: (Petersen et al., 2012).

We Here, we studied four agricultural sites within one of the regions previously studied investigated by Petersen et al. (2012), i.e., a raised bog area with acid soil conditions. The This study included two sites with rotational grass and two sites with a potato crop, and the study covered monitoring took place in spring and autumn periods, where high emissions of N<sub>2</sub>O occurred in previous studies (Petersen et al., 2012; Kandel et al., 2018). We hypothesised that FeS<sub>2</sub> oxidation coupled with NO<sub>3</sub><sup>-</sup> reduction was a possible driver of N<sub>2</sub>O emissions. It was further hypothesised that N<sub>2</sub>O emissions would vary with site conditions affecting denitrification (mineral N availability, rainfall, WT depth and temperature).

## 2 Materials and methods

### 2.1 Study sites

The sites investigated in this study were located in Store Vildmose, which is a 5,000 ha raised bog in northern Jutland, Denmark. The area was, until 150 years ago, the largest raised bog in Denmark, and largely unaffected by human activity. The bog overlies a marine plain formed by the last marine transgression; the sea retreated around 8000 BC, and peat later developed in wet parts of the landscape, attaining a maximum depth of 4.5 to 5.3 m in central parts of the bog (Kristensen, 1945). Between 1880 and 2010, the peat has generally subsided by at least 2 m due to drainage for agriculture or peat excavation (Regina et al., 2016), and today the peat depth is mostly 1-2 m, but in some locations even less (Kandel et al., 2018). The peat and underlying sand is acidic and has been categorised as a potentially acid sulfate soil (Madsen and Jensen, 1988). According to Kandel et al. (2018), the peat at 0-25 cm depth in arable soil in this area has a high degree of humification at, which corresponds to H8 on the Von Post scale.

Four sites were selected along an ~~east-west~~East-West transect (Figure 1a). One arable site (*ARI*) was in a field cropped with second-year potato in 2015, while an adjacent site (*RG1*) in a neighbouring field had second-year rotational grass; these two sites were also represented in the study of Petersen et al. (2012) as sites *N-AR* and *N-RG*, respectively. Land use treatments (i.e., potato and rotational grass) were replicated at sites in other fields; ~~and will be~~ referred to as *AR2* and *RG2*; ~~site~~. Site *AR2* was located 4.6 km to the west, and site *RG2* was located 1.7 km to the east of the paired *ARI-RG1* sites (Figure 1a and S1).

## 2.2 Experimental design

In January 2015, an area of 10 m × 24 m was ~~defined at the location of~~selected each site. Sampling positions were georeferenced using a Topcon HiPer SR geopositioning system (Livermore, CA). On 25 February 2015, each site was fenced, and three 10 m × 8 m experimental blocks were defined (Figure 1b). Each site was further divided along its longitudinal axis to establish two 5 m × 24 m fertilisation subplots.

For monitoring of WT depth, piezometer tubes (Rotek A/S, Sdr. Felding, Denmark) were installed to 150 cm depth at the centre of each block. On either side of the piezometers, at 2.7 m distance, collars of white PVC (base area: 55 cm × 55 cm, height: 12 cm [*RG*] or 24 cm [*AR*]) were installed to between 5 and 10 cm depth (Figure 1). The higher collars used at *AR* sites were ~~at~~level with the ridges ~~established around potato rows~~ during the growth period. The collars, which were fixed to the ground by four 40 cm pegs, had a 4 cm wide flange extending outwards 2 cm from the top to support gas flux chambers. To prevent soil disturbance during gas sampling, platforms (60 cm × 100 cm) of perforated PVC were placed in front of each collar to create a boardwalk. The exact headspace of each collar was determined from 16 individual measurements of distance from the upper rim; this procedure was repeated whenever collars had been removed and reinstalled to accommodate field operations.

Sets of five stainless steel diffusion probes for soil gas sampling at 5, 10, 20, 50 and 100 cm depth were installed vertically within 0.5 m of the flux measurement positions in two blocks (Block 2 and 3) at sites *ARI* and *RG1*, while at sites *AR2* and *RG2* diffusion probes were installed only in Block 2. The stainless steel probes were constructed as described in detail by Petersen (2014), with a 10 cm<sup>3</sup> diffusion cell having a 3 mm diameter opening at the sampling depth covered by a silicone membrane, which was connected to the soil surface via two 18G steel tubes with Luer Lock fittings (Figure S1).

A HOBO Pendant Temperature Data Logger (Onset Computer Corp., Bourne, MA) was installed at 5 cm depth in Block 2 at each site. A mobile weather station (Kestrel 4500; Nielsen-Kellerman, Boothwyn, PA) was mounted at 170 cm height at site *RG1* for hourly recording of air temperature, barometric pressure, wind speed and direction, and relative humidity. Daily precipitation was recorded at <10 km distance from the monitoring sites at a meteorological station, from where data to fill a gap in air temperature were also obtained.

## 2.3 Management

Management within the fenced experimental sites followed the practices adopted by the respective farmers, e.g., with respect to method of fertiliser application, grass cuts, potato harvest and soil tillage. One exception to this was N fertilisation, ~~which rate, since N fertiliser~~ was only given to one of the two subplots in each block (Figure 1b). Fertilised subplots of the *RG1* site received 350 kg ha<sup>-1</sup> NS 27-4 fertiliser on 16 April (~~DOY106DOY 106~~), corresponding to 94.5 kg N ha<sup>-1</sup>. Site *RG2* was fertilised with 20-25 Mg ha<sup>-1</sup> acidified cattle slurry (pH 6) on 5 May (~~DOY125DOY 125~~), and again on 2 July (~~DOY183DOY 183~~), each time corresponding to 90-110 kg total N ha<sup>-1</sup>. ~~After~~On the day of the second slurry application, *RG2* further received 50 kg N ha<sup>-1</sup> as NS 27-4 fertiliser, which was applied by mistake to both fertilisation subplots. The *ARI* site received 100 kg N ha<sup>-1</sup> as liquid NPS 20-3-3 fertiliser on 21 May (~~DOY141DOY 141~~), while the *AR2* site received 110 kg N ha<sup>-1</sup> as NS 21-24 pelleted fertiliser on 30 April (~~DOY120DOY 120~~). The NS fertilisers contained equal amounts of ammonium (NH<sub>4</sub><sup>+</sup>) and nitrate (NO<sub>3</sub><sup>-</sup>), while N in the NPS fertiliser was mainly as NH<sub>4</sub><sup>+</sup>.

At the *RG1* site, the grass was cut in late August, while at the *RG2* site the grass was cut in late June and on 9 September (~~DOY252DOY 252~~). Potato harvest at the *ARI* site took place in mid-September (DOY 258), with interruptions due to heavy rainfall. At the *AR2* site, the potato harvest took place on 23 September (~~DOY266DOY 266~~).

## 2.4 Field campaigns

~~A~~Based on the patterns of N<sub>2</sub>O emissions observed by Petersen et al. (2012), a monitoring program was conducted during spring from 3 March (~~DOY63DOY 63~~) to 16 June (~~DOY169DOY 169~~), and during autumn from 3 September (~~DOY245DOY 245~~) to 10 November (~~DOY314DOY 314~~). Weekly measurement campaigns were conducted at each of the four sites insofar as field operations permitted. Thus, during spring there were 14, 12, 14 and 15 weekly campaigns at the *RG1*, *ARI*, *RG2*, and *AR2* sites, respectively. During autumn there were 10, 10, 7 and 10 weekly campaigns at the *RG1*, *ARI*, *RG2*, and *AR2* sites, respectively. Field trips included sampling at two sites, either *ARI* + *RG1* or *AR2* + *RG2*, and thus all four sites were visited during two field trips on consecutive days. Campaigns included registration of weather conditions and WT depth, soil sampling, soil gas sampling, and N<sub>2</sub>O flux measurements. With a few exceptions, (~~DOY 169 and 265 at site *RG1*, DOY 132 and 314 at site *AR2*~~), each campaign was initiated between 9:00 and 12:00; the order of sites visited in each trip alternated from week to week.

### 2.4.1 Climatic conditions

Air temperature, relative humidity and barometric pressure were logged at the weather station located at *RG1*. During field campaigns, the WT depth was first determined in each of the three piezometers using a Model 101 water level meter (Solinst; Georgetown, Canada). At *ARI* and *AR2*, WT depth in Block 3 was further recorded at 30-minute time resolution for a period during autumn using MaT Level2000 data loggers (MedgeTech; Warner, NH, USA). Soil temperatures at 5, 10 and 30 cm depth were measured in each block using a high precision thermometer (GMH3710, Omega Newport, Deckenpfronn, Germany), and in addition continuous measurements of soil temperature at 5 cm depth were collected in block 2 at each site using HOBO Pendant Temperature Data Loggers (Onset Computer Corp., Bourne, MA).

#### 2.4.2 Soil sampling

During all field campaigns, soil samples were collected separately from fertilised and unfertilised subplots by random sampling of six 20 mm-diameter cores to 50 cm depth- (two per block). Each core was split into 0-25 and 25-50 cm depth, and the six subsamples from each depth were pooled. The pooled samples were transported back to the laboratory in a cooling box and stored at -20°C for later analysis of mineral N and gravimetric water content.

On 23 April (DOY113), and again on 2 September (DOY245), undisturbed soil cores (50 mm diameter, 30 cm segments) were collected to 1 m depth within 1 m distance from the positions of flux measurements in Block 3 of sites *RG1* and *AR1* (cf. Figure 1b). A stainless steel corer (04.15 SA/SB liner sampler, Eijkelkamp, Giesbeek, Netherlands) equipped with a transparent plastic sleeve was used. The steel corer's lower end was capped with a 4 cm long cutting head, and hence sampling depths were 0 to 30 cm, 34 to 64 cm and 68 to 98 cm. The intact cores were capped and sealed, and transported in a cooling box to the laboratory, where they were stored at -20°C: until analysis (see section 2.4.5).

#### 2.4.3 Soil gas sampling

Soil gas samples were collected in 6 mL pre-evacuated Exetainers (Labco Ltd, Lampeter, UK) as described by Petersen (2014) and ~~shown~~ demonstrated in Figure S2. In brief, the diffusion probes were flushed via the inlet tube with 10 mL N<sub>2</sub> containing 50 µL L<sup>-1</sup> ethylene (AGA, Enköping, Sweden) as a tracer, ~~which displaced the gas in the diffusion cell, though with some mixing of sample and flushing gas.~~ A three-way valve, mounted on the outlet tube, was fitted with a 10 mL glass syringe and an Exetainer. The displaced gas was collected in the glass syringe, ~~and from where the glass syringe the 10 mL soil gas sample, now partly diluted by the flushing gas, was could be~~ transferred to the Exetainer. After gas sampling, the probe was flushed with 2 × 60 mL N<sub>2</sub> to remove ethylene, and the Luer Lock fittings were capped. Samples of the N<sub>2</sub>/ethylene gas mixture used for sample displacement were also transferred directly to Exetainers for gas chromatographic analysis ( $n = 3$ ) as reference for the calculation of dilution factors (Petersen, 2014). Sampling for soil gas was done in parallel with flux measurements, except when equipment had to be removed during periods with field operations. Due to damage of some probes during spring, it was decided to discontinue soil gas sampling in the unfertilised grassland subplot ~~at site~~ *RG2-NF*, which had by mistake received fertiliser on DOY 183.

#### 2.4.4 Nitrous oxide flux measurements

Gas fluxes were measured with static chambers (60 cm × 60 cm × 40 cm) constructed from 4-mm white PVC, and equipped with a closed-cell rubber gasket (Emka Type 1011-34; Megatrade, Hvidovre, Denmark) as ~~seal~~ sealing during chamber deployment. Chambers were further equipped with a 12V fan (RS Components, Copenhagen, Denmark) for headspace mixing that was connected to an external battery (Yuasa Battery Inc.; Laureldale, PA), as well as a vent tube with outlet near the ground to minimise effects of wind (Conen and Smith, 1998; Hutchinson and Mosier, 1981). Also, chambers were equipped with an internal temperature sensor (Conrad Electronic SE; Hirschau, Germany), and a butyl rubber septum on top of each chamber for gas sampling. Handles attached to the top were used for straps fixing the

chamber firmly against the collar. Gas samples (10 mL) were taken with a syringe and hypodermic needle immediately after chamber deployment, and then 15, 30, 45 and 60 minutes after closure. Gas samples were transferred to 6 mL Exetainer vials, leaving a 4 mL overpressure.

#### 2.4.5 Soil analyses

Soil samples collected during the weekly campaigns were sieved (6 mm) and subsampled for determination of soil mineral N and gravimetric water content. Approximately 10 g field moist soil was mixed with 40 mL of 1 M potassium chloride (KCl) and shaken for 30 min, and then filtered through 1.6 µm glass microfibre filters. Concentrations of  $\text{NH}_4^+$  and  $\text{NO}_2^- + \text{NO}_3^-$  in filtered KCl extracts were determined by autoanalyser (Model 3; Bran+Luebbe GmbH, Norderstedt, Germany) using standard colorimetric methods (Keeney and Nelson, 1982). Gravimetric soil water content was determined after drying of soil samples at 80°C for 48 hours.

Additional soil characteristics were determined on the intact soil cores collected in April and September at *AR1* and *RG1*. Five cm sections were subsampled from selected depths and analysed for water content, pH, electrical conductivity (EC), total soil organic C and N, and  $\text{NO}_2^-$ . Soil pH and EC were measured with a Cyberscan PC300 (Eutech Instruments; Singapore) in a soil:water solution (1:2.5, w/v). Total soil organic C and total N were measured by high temperature combustion with subsequent gas analysis using a vario MAX cube CN analyser (Elementar Analysensysteme GmbH; Langenselbold, Germany). Soil  $\text{NO}_2^-$ -N concentrations were analysed in soil:water extracts (1:5, w/v) using a modified Griess-Ilosvay method (Keeney and Nelson, 1982). Total organic C and total N were further determined in bulk soil samples (0-25 cm and 25-50 cm depth) collected at *RG2* and *AR2* in the same weeks as sampling of intact cores took place at *AR1* and *RG1*.

The concentration of total reactive Fe (TRFe) at selected depth intervals was determined in the samples from both April and September samplings of intact soil cores. The analysis of TRFe was done using a dithionite-citrate extraction (Carter and Gregorich, 2007; Thamdrup et al., 1994) followed by  $\text{Fe}^{2+}$  analysis with the colorimetric ferrozine method, which ~~included~~includes hydroxylamine as reducing agent (Viollier et al., 2000). The extraction dissolves free (ferric) Fe oxides (except magnetite,  $\text{Fe}_3\text{O}_4$ ), as well as (ferrous) Fe in FeS, but not  $\text{FeS}_2$ .

The intact soil cores, from the September sampling ~~only~~, were further analysed for acid volatile sulfides (AVS) and chromium reducible sulfur (CRS) as indices of FeS and  $\text{FeS}_2$ , respectively. Quantification of AVS and CRS was based on passive distillation adapted from Ulrich et al. (1997) and Burton et al. (2008). Briefly, 0.5 g soil and a trap with 4 mL alkaline Zn-acetate solution (5%) was placed in 120 mL butyl-stoppered (and crimp-sealed) serum bottles, which were evacuated (1 kPa) and pressurised with  $\text{N}_2$  (150 kPa) in three ~~times~~cycles to remove  $\text{O}_2$ , eventually leaving the headspace with  $\text{N}_2$  at atmospheric pressure. Acid volatile sulfide (primarily FeS) was liberated and trapped as ZnS after injection of 12 mL anoxic 2 M HCl followed by sonication (0.5 h) and incubation (24 h) on a rotary shaker (20°C). Using the same approach with replicate soil samples, combined AVS and CRS (primarily elemental S and  $\text{FeS}_2$ ) was trapped after injection of 12 mL 1 M  $\text{Cr}^{2+}$  in 2 M HCl, prepared by reduction of  $\text{CrCl}_3$  (Røy et al., 2014). Trapped sulfide (ZnS) in the two traps was measured colorimetrically using diamine reagent (Cline, 1969), and CRS was then calculated by difference.



Finally, the total reduction capacity of the peat at depths of 27-30 cm, 61-64 cm and 95-98 cm was determined. In brief, a suspension (soil:solution, 1:25; w/v) of oven dried (105°C) sieved soil (<2 mm) and 25 mM cerium (IV) sulfate reagent, Ce(SO<sub>4</sub>)<sub>2</sub> in 5% sulfuric acid (H<sub>2</sub>SO<sub>4</sub>), was shaken horizontally for 24 h at 275 ~~rounds~~revolutions per minute (rpm). After centrifugation at 2,000 rpm, residual Ce(IV) was measured by end-point titration using a solution of 5 mM FeSO<sub>4</sub> in 5% H<sub>2</sub>SO<sub>4</sub>. The amount of reduced compounds was calculated and expressed as meq kg<sup>-1</sup>.

#### 2.4.6 Gas analyses

Nitrous oxide concentrations were analysed on an Agilent 7890 gas chromatograph (GC) combined with a CTC CombiPal auto-sampler (Agilent, Nærum, Denmark). The instrument had a 2 m back-flushed pre-column with Haysep P connected to a 2 m main column with Poropak Q. From the main column, gas entered an electron capture detector (ECD). The carrier was N<sub>2</sub> at a flow rate of 45 mL min<sup>-1</sup>, and Ar-CH<sub>4</sub> (95%/5%) at 40 mL min<sup>-1</sup> was used as make-up gas. Temperatures of the injection port, columns and ECD were 80, 80 and 325°C, respectively. Concentrations were quantified with reference to synthetic air and a calibration mixture containing 2013 nL L<sup>-1</sup> N<sub>2</sub>O. Soil profile N<sub>2</sub>O concentrations were frequently at several hundred μL L<sup>-1</sup>; linearity of the EC detector response was ascertained up to 1600 μL L<sup>-1</sup>, but the entire range was not included in analytical runs as a standard practice, and therefore the higher equivalent gas phase concentrations are relatively uncertain.

Ethylene concentrations in soil gas samples and flushing gas were analysed following a separate injection with an extended run time. All GC settings were as described above, except that run time was different, and gas from the main column was directed to a flame ionisation detector supplied with 45 mL min<sup>-1</sup> H<sub>2</sub>, 450 mL min<sup>-1</sup> air, and 20 mL min<sup>-1</sup> N<sub>2</sub>; the detector temperature was 200°C.

#### 2.5 Data processing and statistical analyses

Equivalent soil gas phase concentrations of N<sub>2</sub>O were calculated assuming full equilibrium (Petersen, 2014) according to Eq. 3:

$$c_s = c_m / \left[ 1 - \frac{(V + d_{out})a_m}{(V - d_{in})a_F} \right] \quad (3)$$

where  $c_s$  is the concentration of N<sub>2</sub>O in the diffusion cell and  $c_m$  the observed concentration (μL L<sup>-1</sup>);  $V$ ,  $d_{in}$  and  $d_{out}$  are the volumes of the diffusion cell, inlet tube and outlet tube, respectively (L);  $a_m$  is the concentration of the tracer ethylene in the gas sample analysed (μL L<sup>-1</sup>); and  $a_F$  is the concentration of ethylene in the flushing gas (μL L<sup>-1</sup>).

Nitrous oxide mixing ratios were converted to units of mass per volume using the ideal gas law and values of pressure and air temperature recorded by the weather station. Individual N<sub>2</sub>O fluxes were calculated in R (version 3.2.5, R Core Team, 2016) using the package HMR (Pedersen et al., 2010). This program analyses non-linear concentration-time series with a regression-based extension of the model of Hutchinson and Mosier (1981), and linear concentration-time series by linear regression (Pedersen et al., 2010). Statistical data ( $p$  value, 95% confidence limits) are provided by

Formatted: Indent: First line: 0.25"

HMR for both categories of fluxes. The choice to use a linear or non-linear flux model was made based on scatter plots and the statistical output.

The temporal dynamics of N<sub>2</sub>O fluxes were analysed [by season for individual site-crop combinations](#) using a generalised linear mixed model defined with the identity link function, the gamma distribution (see Jørgensen and Labouriau, 2012; McCullagh and Nelder, 1989), and Gaussian random components. The model contained a fixed effect representing the interaction between [crop](#), fertilisation and sampling day, and random effects representing site and sampling position. The model for daily N<sub>2</sub>O emission ~~described above~~ was used to estimate cumulative emissions by integrating the flux curves over time. Treatment effects were then analysed by specially designed linear contrasts as described in detail by Duan et al. (2017), who showed that models with untransformed responses (when using adequate distributions) allow simple statistical inference of the time-integrated N<sub>2</sub>O emissions. [WT levels at different time points were compared For each site-crop combination by permutation tests \(Good, 2005\) with 9999 permutations respecting the block structure.](#)

The dependence structure of variables that were potential drivers of N<sub>2</sub>O fluxes were studied using a class of multivariate models called “graphical models” (Whittaker, 1990, see also Labouriau et al., 2008a,b; and Lamandé et al., 2011 for applications in soil science). These models represent the dependence of variables using an undirected graph (not to be confounded with the word “graph” used to refer to a plot), which is a mathematical structure composed of *vertices*, represented by points, and *edges* connecting pairs of vertices, represented by lines connecting points, according to the convention explained below. In graphical models, the variables of interest are the vertices of the graph (represented as labelled points). Here the variables used were: soil temperature at 5 cm depth (Temp5); soil temperature at 30 cm depth (Temp30); NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> concentrations in the top soil (AmmoniumT and NitrateT); N<sub>2</sub>O concentration of the soil gas diffusion probe closest to, but above the WT, i.e., in the capillary fringe (N<sub>2</sub>OWT); and finally, the N<sub>2</sub>O flux (N<sub>2</sub>O-flux). The dependence structure of these variables was characterised by the conditional covariances between each pair of variables given the other variables. Those conditional covariances were simultaneously estimated using the available data according to a statistical model. The graph representation of the model is constructed by connecting the pairs of vertices (i.e., pairs of variables) by an edge when the conditional correlation of the two corresponding variables, given all the other variables, is different from zero. It is possible to show that two variables directly connected in the graph carry information on each other that is not already contained in the other variables (see Whittaker, 1990, Jørgensen and Labouriau, 2012). Moreover, the absence of an edge connecting two vertices indicates that (even a possible) association between the two corresponding variables can be entirely explained by the other variables. According to the general theory of graphical models, if two groups of variables, say A and B, are separated in the graph by a third group of variables, say C (i.e., every path connecting an element of A with an element of B necessarily contains an element of C), then A and B are conditionally uncorrelated given C (see Lauritzen, 1999). This property, called the separation principle, was used below to draw non-trivial conclusions on the interrelationship between N<sub>2</sub>O-flux related variables. The graphical models were inferred by finding the model that minimised the BIC (Bayesian information criterion, i.e., a penalised version of the likelihood function) as implemented in the R package gRapHD (Abreu et al., 2010). This inference procedure yields an optimal representation of the data in the sense that the probability of correct specification of the model, when using this penalisation, tends to one as the

305 number of observations increases (see Haughton, 1988). The confidence intervals for the conditional correlations were  
obtained by a non-parametric bootstrap procedure (Davidson and Hinkley, 1997) with 10,000 bootstrap samples.  
Separate analyses were conducted for each combination of site, crop and season ~~and crop, since different dependency~~  
~~patterns appear in those groups.~~

## 310 3 Results

### 3.1 Climatic conditions

In 2015, the annual mean air temperature in the area of this study was 8.7°C, and annual precipitation was 920 mm.  
This was slightly above the ten-year (2009-2018) average temperature of 8.3°C, and well above the ten-year average  
annual precipitation of 798 mm. During the spring monitoring period, the daily mean air temperature varied between 1  
315 and 15°C, with an increasing trend over the period, and total rainfall was 220 mm. During the autumn monitoring  
period, the daily mean air temperature declined from 15 to 5°C, and total rainfall was 148 mm; the most intense daily  
rain events during spring and autumn were 16.9 and 33.2 mm, respectively.

Soil temperature at 5 cm depth showed a clear diurnal pattern (Figure S3), but at all four sites the temperature at the  
time of chamber deployment was close to the daily mean temperature at this depth. Thus, across the four sites the  
320 average deviation ranged from 0.2 to 0.9°C, and the largest deviations on a single day were -2.0 and 2.1°C,  
respectively.

### 3.2 Soil characteristics

~~Soil~~Several soil characteristics were determined by analyses of intact cores collected in late April (DOY 113) 2015  
(Table 1). At all sites the soil was acidic, with pH ranging from 4.7 to 5.4. At the paired sites *ARI* and *RG1*, a weak  
325 decline in pH was indicated at 40-50 cm depth. Electrical conductivity at *ARI* and *RG1* sites ranged from 0.15 to 0.91  
mS cm<sup>-1</sup>, with no obvious trends in the data; the highest value (0.91 mS cm<sup>-1</sup>) occurred at site *ARI* at 93-98 cm in a  
layer dominated by sand underlying the peat.

The organic matter composition of soil profiles at the four sites varied. Total organic C concentrations at sites *ARI*  
and *RG1* were 34-43% in the upper 0-40 cm, but then dropped to only 0.3-0.6% at c. 1 m depth in the sand. The peat  
330 was amorphous and well-decomposed at 0-20 cm depth, while the underlying peat ~~was dominated by~~contained intact  
plant debris. At site *RG2*, the process of peat degradation was evident even at 0-50 cm depth, where TOC  
concentrations only just met the requirements for being defined as an organic soil; ~~i.e.,~~ the organic C content was below  
20 and 10% at 0-25 and 25-50 cm depth, respectively. Site *AR2* was characterised by a uniform peat layer (33-38%  
organic C) at 0-50 cm depth. Across all sites, the C:N ratios ranged between 14 and 26 in the organic soil layers.

335 Two iron sulfide fractions, as well as total reactive iron, were quantified. Acid volatile sulfide ranged from 1.7 to  
4.9  $\mu\text{g S g}^{-1}$  soil across the four sites and showed no clear relationship with soil depth. This was also the case for CRS,  
which ranged from 24 to 155  $\mu\text{g S g}^{-1}$  dry weight soil. Total reactive Fe (TRFe) concentrations in soil profiles from sites  
340 *ARI* and *RG1* ranged from 1.19 to 4.99  $\text{mg g}^{-1}$  dry weight soil at 0-50 cm depth, and hence concentrations of reactive Fe  
were up to 1500 times higher than concentrations of Fe in AVS (assuming this was  $\text{FeS}$ ), and 25-120 times higher than  
Fe in CRS (assuming this was  $\text{FeS}_2$ ). At sites *ARI* and *RG1*, TRFe declined below 20 cm depth and was close to zero in  
the sand below the peat layer (Table 1). The highest concentrations of TRFe at sites *RG1* (Figure 2b) and *ARI* (Figure  
2d) occurred at 20 cm depth on 23 April. (DOY 113). At site *ARI*, a sink for TRFe at 40-60 cm depth was indicated:  
There and, disregarding Depth 6 (93-98 cm), the concentration of TRFe at Depth 5 (47.5-52.5 cm) was significantly  
lower than concentrations at more shallow depths ( $p < 0.05$ ). Differences in TRFe concentration were only minor  
345 observed at site *RG1*, and any differences in the distribution of TRFe between seasons, were probably also minor (not  
tested). There was a strong correlation between TRFe and TOC across all sites ( $r = 0.88$ ,  $n = 16$ ).

The total reduction capacity was determined by a wet oxidation procedure using  $\text{Ce}(\text{SO}_4)_2$ . At both *ARI* and *RG1*,  
the total reductive capacity of the peat at 27-30 cm depth was outside the range of the analytical method at  $>11,500$  meq  
 $\text{kg}^{-1}$ . The reduction capacity dropped to around 1000 meq  $\text{kg}^{-1}$  at 60 to 65 cm depth with a declining organic matter  
350 content, and 50 to 50-100 meq  $\text{kg}^{-1}$  in the sandy layer at 100-93 cm depth.

### 3.3 Soil mineral N dynamics

Soil concentrations of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  at 0-25 and 25-50 cm depth were determined in connection with field campaigns  
(Tables S1-S4). Since subsamples were pooled for analysis, only a qualitative description of the effects of treatments  
355 and temporal dynamics is possible. The residence time for mineral N in the soil solution was generally appeared to be  
longer at *AR* compared to *RG* sites. At *AR* sites, there was an accumulation of mineral N (Table S2, S4) at both depth  
intervals during May, that also occurred before N fertilisation. Mineral N concentrations were apparently greater at *ARI*  
compared to *AR2*, and at site *AR2* only  $\text{NO}_3^-$  accumulated. Fertilisation increased Following fertilisation,  $\text{NH}_4^+$ -N and  
 $\text{NO}_3^-$ -N concentrations were 100-200  $\mu\text{g g}^{-1}$  dry weight soil at all sites except *RG2* (Table S3), where acidified cattle  
360 slurry was applied. Nitrate accumulated at all sites in the weeks after fertilisation was observed at  
all sites, and also there was evidence for some transport to 25-50 cm depth.

Nitrite-N concentrations were determined in soil profiles from the cores sampled at sites *RG1* and *ARI* on 23 April  
(DOY 113) and 2 September (DOY 245) 2015. Both fertilised and unfertilised subplots were represented, although at  
365 site *ARI* the *RG1*, fertilisation had not yet taken place at the time of sampling in April. There was variation at depth in  
the soil, which could not be explained by fertilisation one week earlier, but a two-sample *t* test did not find evidence for  
an effect on  $\text{NO}_2^-$  availability ( $p = 0.19$ ), and therefore the results from both *ARI* and *RG1* subplots are presented  
together. In April, the average concentration of  $\text{NO}_2^-$ -N at both sites was highest (c. 10  $\mu\text{g g}^{-1}$  dry weight soil) around 40  
cm depth and declined towards the surface and deeper layers (Figure 2a,c). However, due to heterogeneity of soil  
profiles, the differences between depths were not significant ( $p = 0.22$  for *RG1* and  $p = 0.06$  for *ARI*). A decline in  $\text{NO}_2^-$

Formatted: Font: Italic

-N concentration was indicated at 50 cm depth at site *ARI* ~~relative to site *RG1*, and also, where~~ a depletion of TRFe was indicated. However, there was also ~~a lower concentration of peatless organic matter~~ (cf. TOC in Table 1), which may account for this difference. In September, NO<sub>2</sub>-N concentrations were <1 µg g<sup>-1</sup> dry weight soil at both sites, while the ~~much higher~~ concentrations of TRFe were comparable to those in April.

### 3.4 Groundwater table dynamics

Across the four sites, WT changes ranged from 60 to 100 cm. During spring, WT depth at sites *RG1* and *ARI* varied between 17 and 81 cm, with a steady decline until the end of April (~~DOY120~~), and by DOY 119 the WT depth was significantly (all  $p$  values < 0.05) below that ~~was of all previous samplings except DOY 112~~ (Figures 3 and 4). Then followed ~~by a period with fluctuations frequent rainfall, where WT fluctuated (no significant changes) around 60-80 cm depth due to frequent rainfall (Figures 3 and 4).~~ During the first half of September (~~DOY246~~ DOY 246 to 259), rainfall caused the WT to rise from 80 to 40 cm depth ( $p < 0.05$ ; Figures 5 and 6). On two occasions (~~DOY248~~ DOY 248 and 260) the WT depth rose to 20 cm depth and only gradually declined during the following days (data not shown). From mid-September (DOY 258) ~~then there~~ followed a period with a gradual WT decline until early November (DOY 308), where upon the WT ~~rose showed an increasing trend~~ from 90 to 45 cm depth during a week with intense rainfall. At site *RG2*, (Figure 3) the WT ~~was initially declined ( $p < 0.05$ ) and then remained~~ mostly at 50-60 cm depth during spring, with a temporary rise to 30 cm depth ~~by on~~ 3 June (~~DOY139; see Figure 3~~; DOY 139;  $p < 0.05$ ). In the autumn, sampling campaigns ~~could not be initiated until DOY260 due to harvest were resumed on DOY 245~~. By this time, the WT was close to the surface following intense rainfall, but then declined ( $p < 0.05$ ) to 80-100 cm in the sandy subsoil (Figure 5). The WT ~~depth~~ at site *AR2* ~~was consistently declined initially ( $p < 0.01$ ) and then fluctuated~~ between 45 and 60 cm depth during spring except for a transient increase ( $p = 0.05$ ) to 35 cm depth in early June (Figure 4). During autumn, the WT rose ( $p = 0.05$ ) to the soil surface in September (~~DOY260~~ DOY 260), and then gradually withdrew ( $p = 0.05$ ) until early November (DOY 307) when rainfall caused a c. 40 cm increase (Figure 6), as also observed at sites *RG1* and *ARI*.

### 3.5 Soil N<sub>2</sub>O concentration profiles

~~The distribution and temporal dynamics of N<sub>2</sub>O in the soil profiles showed important contrasts between grassland and arable sites.~~ Equivalent gas phase concentrations of N<sub>2</sub>O in passive diffusion samplers were determined concurrently with gas sampling, and results are presented as contour plots (Figures 3-6; data in Table S5). Concentrations in many cases varied by several orders of magnitude between sites and sampling days, and between depths within individual profiles, and therefore a logarithmic grey scale ~~was is~~ used to show ~~trends gradients~~. The gaps in Figures 3-6 indicate periods, where diffusion probes could not be installed or were temporarily removed due to field operations.

Under the rotational grass at site *RG1*, soil N<sub>2</sub>O concentrations during spring were mostly between 0.1 and 3 µL L<sup>-1</sup> (Figure 3). A higher concentration (15 µL L<sup>-1</sup>) was observed at 40-80 cm depth in the fertilised subplot around ~~DOY139~~ DOY 139, but only ~~in at the lower position~~ (Block 3) of the field plot. At site *RG2*, the concentrations of N<sub>2</sub>O in the soil during spring were generally similar to those of *RG1*, although there were more values in the 1-10 µL L<sup>-1</sup> concentration range ~~(in the unfertilised plot (Figure 3, Table S5)).~~ However, on 3 June (~~DOY154~~ DOY 154) a significant

Formatted: Indent: First line: 0.25"

increase in much higher N<sub>2</sub>O concentration occurred was observed in the fertilised part of the plot with a maximum of 560  $\mu\text{L L}^{-1}$  at 100 cm depth (i.e., well below the WT). Soil N<sub>2</sub>O concentrations in the unfertilised plot were also increased/elevated around this time, but only to e-a maximum of 15  $\mu\text{L L}^{-1}$  and mainly near the soil surface.

The arable site *ARI*, with sampling positions located in a different field, but only 10-20 m from those of site *RG1*, showed had apparently very different soil N<sub>2</sub>O concentration dynamics during spring (Figure 4). There was a consistent accumulation of N<sub>2</sub>O at 50 and 100 cm depth where seasonal concentrations averaged 340 and 424  $\mu\text{L L}^{-1}$ , respectively. In contrast, at 5, 10 and 20 cm depth the average N<sub>2</sub>O concentrations were 10-30  $\mu\text{L L}^{-1}$ , and there was no clear response to fertilisation on DOY141+DOY 141 in terms of soil N<sub>2</sub>O accumulation. The soil N<sub>2</sub>O concentrations suggested that there was considerable within-site heterogeneity in soil conditions, as the highest concentrations were often observed in the unfertilised subplot. Between DOY75 and DOY100, the Maximum concentrations of N<sub>2</sub>O peaked at of nearly 1500  $\mu\text{L L}^{-1}$  were recorded at 50 cm depth and were 2-3 fold higher than on DOY 99, and a similar concentration at 100 cm depth on DOY 112, both in the unfertilised subplot. At site *AR2*, the highest soil N<sub>2</sub>O concentrations during early spring were consistently observed at 20 cm depth, but then seemed to gradually declining/decline to reach the background level of 0.3  $\mu\text{L L}^{-1}$  in mid-May (around DOY130+DOY 130). In the unfertilised field-plots/subplot, the N<sub>2</sub>O concentration then increased again-reached 272  $\mu\text{L L}^{-1}$  at 20 cm depth to reach 272  $\mu\text{L L}^{-1}$  following rainfall, and with WT rise to at 35 cm depth. With fertilisation, soil N<sub>2</sub>O concentrations were even higher at 10 cm depth and reached nearly 400  $\mu\text{L L}^{-1}$  in mid-June. The observed accumulation of N<sub>2</sub>O near the soil surface was accompanied by increasing N<sub>2</sub>O emissions during this period (Figure 4).

During autumn, N<sub>2</sub>O concentrations in the soil profile at the *RG1* and *RG2* sites varied between 0 and 12  $\mu\text{L L}^{-1}$ , with a tendency for higher concentrations at 10-20 cm depth (Figure 5). At site *RG1*, where both fertilised and unfertilised subplots could be sampled, this was apparently independent of fertilisation.

September was characterised by heavy rainfall (114 mm in total), and at site *ARI* a substantial rise in the WT from 80 to 40 cm depth was observed (Figure 6). Soil N<sub>2</sub>O concentrations showed a dual-pattern, with maxima at 10 and 100 cm depth through to DOY266+DOY 266 (end of September), and after this time soil N<sub>2</sub>O accumulation rapidly declined as the concurrently with WT withdrew/drawdown. Nitrous oxide concentrations equivalent to several hundred  $\mu\text{L L}^{-1}$  were measured even at 5 cm depth during this period. During late autumn, the N<sub>2</sub>O concentration at 0-50 cm depth varied between 0 and 20  $\mu\text{L L}^{-1}$ , whereas at 100 cm depth it remained high at 100-850  $\mu\text{L L}^{-1}$ . At site *AR2*, the groundwater level was higher than at *ARI* and reached/came close to the soil surface by mid-September (DOY 260). Soil N<sub>2</sub>O accumulated in both fertilised and unfertilised subplots following saturation of the soil, again with the highest concentrations apparently occurred at 20 cm depth. A secondary increase was observed near the soil surface at the last sampling on DOY314+DOY 314 in November, in response to a period with rainfall and a rapid WT rise.

### 3.6 Nitrous oxide emissions

The weekly sampling campaigns during spring and autumn showed much that with few exceptions N<sub>2</sub>O emissions, expressed as average daily rates, were significantly higher N<sub>2</sub>O emissions at arable compared to grassland sites independent of season and fertiliser N application. At site *RG1*, N<sub>2</sub>O emissions during (Table 2). One exception was

Commented [SOP1]: Statistik?

Commented [SOP2]: Statistik?



treatment *RG2-F* in spring ranged from 0 to 550  $\mu\text{g N}_2\text{O m}^{-2} \text{h}^{-1}$ , with no effect of fertiliser amendment (Figure 3). The grass in the fertilised subplot showed a clear response to fertilisation that indicated uptake of fertiliser N. At site *RG2*, where a peak in  $\text{N}_2\text{O}$  emissions occurred was indicated ( $p = 0.06$ ) on DOY 154, and the flux was still elevated at the next two samplings (Figure 3). This high flux coincided with the accumulation of  $\text{N}_2\text{O}$  in the soil profile described above, and it was the only time that an effect of fertilisation was observed. The grass in the fertilised subplot showed a clear visual response to fertilisation, which indicated that fertiliser N was effectively taken up by the sward.

At site *AR1*, the  $\text{N}_2\text{O}$  fluxes were generally much higher than at grassland sites during spring (Figure 4). Fluxes during early spring reached 2000-6000  $\mu\text{g N}_2\text{O m}^{-2} \text{h}^{-1}$  and were higher than in late spring where, as for site *RG1* (Figure 4). Since there was no effect of N fertilisation was observed. Hence (cf. Table 2), the higher emissions were associated with site differences probably derived from soil N pools and caused by factors other than fertilisation. The potato field at site *AR2* showed a different pattern, with  $\text{N}_2\text{O}$  fluxes remaining low during early spring, and for several weeks after fertilisation. The highest emissions occurred, independent of fertilisation, an increasing trend (not significant) was observed in June following a WT rise to in WT, which was at 35 cm depth on DOY 154.

In the autumn,  $\text{N}_2\text{O}$  fluxes from site *RG1* were consistently low (Figure 5). The first sampling at site *RG2* was on DOY 259 in mid-September, where a high flux of 3000  $\mu\text{g N}_2\text{O m}^{-2} \text{h}^{-1}$  was seen, which dropped to near zero within 1-2 weeks. Nitrous oxide emissions at site *AR1* were high during September at 4000-10,000  $\mu\text{g N}_2\text{O m}^{-2} \text{h}^{-1}$  independent of across the two N fertilisation treatments (Table S2), and subsequently declined ( $p < 0.05$ ) to near zero (Figure 6). High fluxes were observed on the first sampling day of this monitoring period, DOY 246, while DOY 246, at a time where WT depth was still at 40 to 80 cm depth. However, this followed. Instead the high  $\text{N}_2\text{O}$  fluxes may have been triggered by saturation of the top soil after 10 and 22 mm rainfall on the previous two days. Rainfall. Additional rainfall during the following days then was accompanied by a rise in WT. The subsequent decline in  $\text{N}_2\text{O}$  emissions at *AR* sites coincided with WT withdrawal draw-down and drainage of the top soil.

Cumulative  $\text{N}_2\text{O}$  emissions were calculated for the 99-105 days of monitoring period in spring, and for as well as the 47-69 d period in the autumn. Daily rates were surprisingly consistent in the two periods (Table 2), and therefore the following refers to cumulative emissions across both periods. At *RG* sites, the average  $\text{N}_2\text{O}$  flux from site *RG1*, the total emission was 4 kg  $\text{N}_2\text{O-N ha}^{-1}$  independent of fertilisation, whereas at *RG2* the fertilised grassland was significantly higher than from and unfertilised grass (7.3 vs. 2.0 subplots were different at 18 and 5 kg  $\text{N}_2\text{O-N ha}^{-1}$ ) during spring, respectively. At *AR* sites with potato, there was no significant effect of N fertilisation, but and the cumulative  $\text{N}_2\text{O}$  emissions of 15-17 were 18-20 kg  $\text{N}_2\text{O-N ha}^{-1}$  were at *AR2*, but much higher than from *RG* sites. In the autumn, the average cumulative emissions at the *RG* and *AR* sites were 2 and 15 at 44-52 kg  $\text{N}_2\text{O-N ha}^{-1}$ , respectively, at site *AR1*.

### 3.7 Interrelationships between driving variables of $\text{N}_2\text{O}$ production

Graphical models were used to study the dependence structure among selected soil variables and  $\text{N}_2\text{O}$  fluxes. Interestingly, at both *RG* sites, and in both spring (Figure 7a) and autumn (Figure 7b), and at *AR* sites in spring (Figure

Formatted: Indent: First line: 0.25"

7e), the only variable with a direct link to seasons, N<sub>2</sub>O flux was consistently depended on N<sub>2</sub>O<sub>WT</sub>, i.e., soil N<sub>2</sub>O concentration in the capillary fringe (N<sub>2</sub>O<sub>WT</sub>), indicating that N<sub>2</sub>O<sub>WT</sub> carried information on the N<sub>2</sub>O flux that could not be explained by indirect correlations between the other variables. Moreover, the variable Figure 7a,b and 8a,b). This was also the case for both AR sites in spring (Figure 7c,d). Where N<sub>2</sub>O<sub>WT</sub> separated N<sub>2</sub>O flux from the other variables in the graph which, this indicates according to the separation principle (an instance of the general theory of graphical models), indicates section 2.5) that information about this variable N<sub>2</sub>O<sub>WT</sub> rendered all the other variables uninformative with respect to N<sub>2</sub>O flux. For example, in the analysis of AR sites AR1 in spring (Figure 7e7b), the variables N<sub>2</sub>O flux and Temp5 were not directly connected, and therefore any correlation between Temp5 and N<sub>2</sub>O flux could be completely explained by other variables.

The only exception to this pattern was graphical model results were different for AR sites in the autumn (Figure 7d8c,d), where instead two other variables showed a significant relationship with N<sub>2</sub>O flux; one variable was Nitrate T, i.e., soil temperature at 30 cm depth and (site AR1 only) NO<sub>3</sub>-N concentration in the top soil, and the other variable was soil temperature at 30 cm depth, showed significant correlation with N<sub>2</sub>O flux ( $p < 0.05$ ). All other relationships variables were unrelated to N<sub>2</sub>O flux, or could be accounted for by other variables.

#### 4 Discussion

This study investigated seasonal dynamics of N<sub>2</sub>O emissions and soil conditions in an area, which has been designated as a hotspot for N<sub>2</sub>O emissions (Leppelt et al., 2014). Spring and autumn monitoring periods together covered between 152 and 174 d, and cumulative N<sub>2</sub>O emissions during these periods were in total 3-64-5 kg N<sub>2</sub>O-N ha<sup>-1</sup> for rotational grass if disregarding the two-week period after slurry application at RG2-F, and 49-2420-50 kg N<sub>2</sub>O-N ha<sup>-1</sup> for arable sites with a potato crop. These numbers, representing <6 month periods, thus confirmed previous results (Petersen et al., 2012) that annual N<sub>2</sub>O emissions from organic soil in this area are comparable to (RG), or clearly above (AR), the IPCC emission factors for drained organic soil of 8 and 13 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup> for nutrient rich grassland and cropland, respectively (IPCC, 2014). The area has been characterised as potentially acid sulfate soil (Madsen and Jensen, 1988), and a previous study showed groundwater sulfate concentrations in excess of 100 mg L<sup>-1</sup> (Petersen et al., 2012). We therefore hypothesised that NO<sub>3</sub><sup>-</sup> reduction coupled with FeS<sub>2</sub> oxidation could be a pathway of N<sub>2</sub>O formation in this acid organic soil,

Pyrite, measured as CRS, was quantified at selected depths (Table 1), and with). Soil bulk density of the peat varying varied between 0.15 and 0.3 g cm<sup>-3</sup> (data not shown), and the total amount of CRS at 0 to 50 cm depth would thus could therefore be estimated to 200-350 mmol FeS<sub>2</sub> m<sup>-2</sup>. The N<sub>2</sub>O emissions observed during spring and autumn monitoring periods constituted up to 145 mmol N m<sup>-2</sup> in total (site AR1), and it is thus theoretically possible that the process described by Eq. 2 contributed to emissions of N<sub>2</sub>O. However, the FeS<sub>2</sub> concentration (0.74-2.4 mmol kg<sup>-1</sup>) represented a minor part of the total reduction capacity (>11,500 meq kg<sup>-1</sup> at 27-30 cm depth). Also, the concentration average concentrations of total reactive Fe was were 25-120 times higher than that of FeS<sub>2</sub> (though less in terms of reduction equivalents). Reducing It is therefore likely that reducing agents other than FeS<sub>2</sub> were therefore likely

Formatted: Indent: First line: 0.2"

510 ~~to be more~~ important, a conclusion that was later supported by a laboratory study in which peat amended with FeS<sub>2</sub> did  
not show enhanced N<sub>2</sub>O production (Taghizadeh-Toosi et al., ~~submitted~~-2019). Other possible drivers of N<sub>2</sub>O  
emissions were therefore considered.

#### 4.1 Environmental drivers of N<sub>2</sub>O emissions

515 A limited number of potential drivers (rainfall, temperature, soil mineral N and soil concentrations of N<sub>2</sub>O) were  
monitored to help explain N<sub>2</sub>O emission dynamics. Soil N<sub>2</sub>O concentration profiles showed complex patterns where, for  
example, the highest concentrations were sometimes observed above, and sometimes below the WT depth at both RG  
(Figure 3) and AR sites (Figure 4). Fertilisation in spring was associated with higher concentrations of N<sub>2</sub>O below the  
WT depth at sites RG2 and AR2, which indicated downward transport of fertiliser N, but this was not reflected in  
elevated N<sub>2</sub>O emissions. The reason may be that in wet soil the time required to reach a steady state between N<sub>2</sub>O  
520 production and emissions from the soil surface can be significant and increases with distance (Jury et al., 1982). In  
accordance with this, Clough et al. (1999) observed a delay of 11 days before <sup>15</sup>N enriched N<sub>2</sub>O produced at 80 cm  
depth was released from the soil surface at a corresponding rate. Since the presence of air-filled porosity is critical for  
the exchange of gases between soil and the atmosphere (Jury et al., 1982), the soil N<sub>2</sub>O concentration closest to, but  
above the WT depth (N<sub>2</sub>O<sub>WT</sub>), was taken to represent "subsoil" processes stimulating N<sub>2</sub>O emissions.

525 The regulation of N<sub>2</sub>O emissions was investigated using a statistical method represented by graphical models.  
~~It~~Among the factors considered, the graphical models for individual site-crop combinations consistently identified N<sub>2</sub>O  
concentration in the capillary fringe as the strongest predictor of N<sub>2</sub>O emissions from both grassland and arable ~~soil~~sites  
in spring, (Figure 7a-d), and from grassland ~~soil~~sites in the autumn, (Figure 8a,b). The implication is that N  
transformations at depth in the soil, and not in the top soil, were the main source of N<sub>2</sub>O escaping to the atmosphere in  
530 these cases. In accordance with this, there was no ~~immediate~~apparent effect of N fertilisation on emissions of N<sub>2</sub>O  
~~independent, with emissions at RG2 after the application of land-use acidified cattle slurry as a notable exception.~~ Other  
studies also found a limited response to fertilisation (Maljanen et al., 2003; Regina et al., 2004), although Regina et al.  
(2004) later observed a peak in N<sub>2</sub>O emissions after rainfall. Goldberg et al. (2010) reported that N<sub>2</sub>O emissions from a  
minerotrophic fen were produced at 30-50 cm depth, in accordance with the observations presented here, where the  
535 highest concentrations of N<sub>2</sub>O were mostly observed at 20 or 50 cm depth (Table S5).

Peat decomposing in the capillary fringe during WT drawdown could have been the source of N for N<sub>2</sub>O  
production. It is well established that N<sub>2</sub>O emissions from organic soil may be enhanced by drainage (Martikainen et  
al., 1993; Taft et al., 2017), and the response will appear within days, as shown by Aerts and Ludwig (1997) in an  
incubation study with an oscillating WT. A stimulation of N<sub>2</sub>O emissions by WT drawdown was also observed by  
540 Goldberg et al. (2010) when simulating drought under field conditions, although a pulse of N<sub>2</sub>O also occurred after  
rewetting ~~in that study~~. In accordance with this, ~~rising WT and/or increasing soil wetness in effect of rewetting, rain  
events during~~ late spring and early autumn ~~consistently often~~ enhanced N<sub>2</sub>O emissions ~~at all sites in the present study.~~  
~~This was not necessarily a result of WT changes.~~ Despite 32 mm rainfall on ~~DOY244~~DOY 244 and 245, the WT depth  
at site AR1 was still at 40 to 80 cm ~~and could not account for the very high on DOY 246 (Figure 6), but~~ N<sub>2</sub>O emissions  
545 ~~observed on DOY246 (Figure 6) were high.~~ Well-degraded peat will release as little as 10% of its water to drainage

(Rezanezhad et al., 2016). ~~It~~, and it is therefore likely that ~~the~~ rain water was absorbed by peat above the WT and created conditions suitable for denitrification: close to the soil surface, which explains the significant correlation between N<sub>2</sub>O emissions and NO<sub>3</sub><sup>-</sup> (Figure 8c).

The increase in N<sub>2</sub>O emissions during WT cycles reported by Aerts and Ludwig (1997) was observed only with eutrophic peat, whereas a mesotrophic peat showed no effect of WT dynamics on N<sub>2</sub>O emissions, which were consistently low. A similar interaction between nutrient status and WT depth was observed in field studies comparing N<sub>2</sub>O emissions from minerotrophic and ombrotrophic boreal peatlands (Martikainen et al., 1993; Regina et al., 1996). In the present study, soil NH<sub>4</sub><sup>+</sup>-N and NO<sub>3</sub><sup>-</sup>-N concentrations at site *RG1* increased to 133 and 120 µg g<sup>-1</sup> dry weight soil upon fertilisation, respectively, but largely returned to the background level of around 5 and 10 µg g<sup>-1</sup> dry weight soil, respectively, within a week (Table S1). In contrast, at site *AR1* there was ~~significant~~ substantial accumulation of NH<sub>4</sub><sup>+</sup>-N and NO<sub>3</sub><sup>-</sup>-N even before fertilisation on ~~DOY141~~ DOY 141, and soil mineral N remained high for several weeks (Table S2). ~~This accumulation of soil mineral N around the time of potato crop establishment could~~ that may have stimulated N<sub>2</sub>O emissions in the arable soil. Grasslands on organic soil generally show lower emissions of N<sub>2</sub>O compared to arable organic soil (Eickenscheidt et al., 2015), presumably because plants compete successfully with microorganisms for available N. Schothorst (1977) estimated peat decomposition indirectly from the N-content in herbage yield of grassland and concluded that the soil supplied 96 kg N ha<sup>-1</sup> when the drainage depth was 25 cm, but 160 and 224 kg N ha<sup>-1</sup> with the drainage depth at 70 and 80 cm, respectively. Hence, plant uptake of N mineralised from soil organic matter above the WT likely ~~caused~~ contributed to the much lower N<sub>2</sub>O emissions from rotational grass in this study.

~~Nitrous oxide concentration profiles provided indirect information about soil mineral N dynamics.~~ At *RG* sites, soil N<sub>2</sub>O concentrations during spring were generally low and ~~thus~~ did not ~~provide clear~~ show evidence for microbial N transformations, which supports the conclusion above that plant uptake was a main sink for the N released during peat decomposition. ~~At site RG2 an~~ One exception was the accumulation of N<sub>2</sub>O ~~was seen at 4 m~~ 50-100 cm depth observed at site RG2, in late May (Figure 3), which could have been caused by leaching of mineral N from the acidified cattle slurry following extensive rain. In contrast, at *AR* sites ~~there was significant accumulation of~~ N<sub>2</sub>O accumulated in the soil throughout spring irrespective of fertilisation; at site *AR1* the highest concentrations occurred at 50 ~~to~~ and 100 cm depth, while at site *AR2* with a higher groundwater table the highest concentrations were at 20 cm depth, ~~in accordance with the higher groundwater table.~~ These observations indicated, It showed that N<sub>2</sub>O ~~was produced in the capillary fringe, consistent with~~ peat decomposition ~~as was~~ a source of mineral N, and a likely source of N<sub>2</sub>O above, and possibly also in below the saturated zone (see next section). ~~Following N fertilisation, the accumulation of N<sub>2</sub>O in the soil profile was mostly associated with precipitation and rising WT.~~

In the autumn, the graphical ~~model~~ models identified NO<sub>3</sub><sup>-</sup> in the top soil, (site AR1 only) and soil temperature at 30 cm depth, as ~~significant~~ predictors of N<sub>2</sub>O emissions at arable sites (Figure 7). The accumulation of NO<sub>3</sub><sup>-</sup> was much greater at site *AR1* compared to *AR2*, ~~suggesting differences in N mineralisation potentials, possibly because AR1 had better drainage of the top soil (e.g., WT at 80 vs. 40 cm depth on DOY 246; Figure 6).~~ It is not clear if the source of N was decomposing potato ~~erect~~ top residues or accelerated peat decomposition ~~following soil disturbance at harvest, or both.~~ Rainfall most likely triggered denitrification by rapidly increasing WT depth and soil water-filled pore space and raising WT depth, thereby impeding the ~~oxygen~~ O<sub>2</sub> supply to much of the soil profile (Barton et al., 2008). This

Formatted: Font: Italic


585 interpretation is supported by increasing N<sub>2</sub>O concentrations below, as well as above the WT depth depending on site  
and block, and in fertilised as well as unfertilised subplots (Figure 6). In an annual study, conducted in other parts of the  
Store Vildmose bog, Kandel et al. (2018) also measured high emissions of N<sub>2</sub>O from a potato crop, i.e., around 2000 µg  
N<sub>2</sub>O m<sup>-2</sup> h<sup>-1</sup> in October 2014 and 6000 µg N<sub>2</sub>O m<sup>-2</sup> h<sup>-1</sup> in June 2015, which coincided with NO<sub>3</sub><sup>-</sup> accumulation and  
rainfall. Precipitation was also high during September 2015, and ~~the rapid rise in WT toward the soil surface resulted~~  
~~in this was accompanied by~~ accumulation of N<sub>2</sub>O in the top soil at all sites. However, N<sub>2</sub>O concentrations reached only  
590 around 10 µL L<sup>-1</sup> at RG sites, as opposed to several hundred µL L<sup>-1</sup> at AR sites, confirming that soil mineral N  
availability was a limiting factor for N<sub>2</sub>O emissions.

#### 4.2 Pathways of N<sub>2</sub>O emissions

Bacterial nitrification, denitrification, and nitrifier-denitrification are all ~~potential-potentially significant~~ pathways  
of N<sub>2</sub>O formation (Braker and Conrad, 2011). The ~~significant relationship correlation~~ with NO<sub>3</sub><sup>-</sup> ~~in the top soil at AR~~  
~~sitesite ARI~~ in the autumn (Figure 78c) suggested that ~~here~~ denitrification activity ~~in the top soil~~ controlled N<sub>2</sub>O  
595 emissions ~~during this period. This was different in early spring, where, but mostly~~ soil mineral N concentrations were  
low, and ~~N<sub>2</sub>O accumulated near the WT depth. Here,~~ ammonia oxidation activity may ~~therefore have controlled~~ limited  
N<sub>2</sub>O emissions either directly, or indirectly *via* production of NO<sub>2</sub><sup>-</sup> or NO<sub>3</sub><sup>-</sup>. Ammonia oxidising bacteria (AOB) are  
scarce in acid peat despite the presence of nitrite oxidising bacteria (NOB) (Regina et al., 1996), and some studies  
indicate that ammonia oxidising archaea (AOA) predominate in both abundance and activity (Herrmann et al., 2012;  
600 Stopnišek et al., 2010). Stieglmeier et al. (2014) isolated an AOA from soil that emitted N<sub>2</sub>O at a rate corresponding to  
0.09% of the NO<sub>2</sub><sup>-</sup> produced independent of O<sub>2</sub> availability, but it is not known if this organism is present in acid  
organic soil, and at this time an indirect control of denitrification activity seems more plausible.

Stopnišek et al. (2010) found that AOA activity was not stimulated by an external source of NH<sub>4</sub><sup>+</sup> and concluded  
that the activity was associated with N released from decomposing soil organic matter. ~~Well-decomposed peat is~~  
605 ~~dominated by dead-end pores associated with plant cell remains, which are characterised by a slow exchange of solutes~~  
~~with active pore volumes (Hoag and Price, 1997), and hence ammonia oxidation in confined spaces could be important~~  
~~in organic soil.~~ The anaerobic conditions of saturated peat may have ~~been a limiting factor for~~ limited N mineralisation  
and ~~therefore hence~~ ammonia oxidation activity during early spring, a constraint which was alleviated as the WT  
declined and ~~oxygen entered deeper soil layers. Nitrite had accumulated at 20-50 cm depth in late April at both RG/~~  
610 ~~and ARI sites (Figure 2), which was consistent with peat decomposition and ammonia oxidation following WT~~  
~~drawdown. Total concentrations of NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> at 25-50 cm depth were significant (Tables S1 and S2), but well-~~  
~~decomposed peat is dominated by dead-end pores (Hoag and Price, 1997), and it is likely that ammonia oxidation to a~~  
~~large extent took place in such pores having a slow exchange of solutes with active pore volumes. The accumulation of~~  
~~NO<sub>2</sub><sup>-</sup> suggested there was an imbalance between ammonia oxidation and nitrite oxidation activity. O<sub>2</sub> entered deeper soil~~  
615 ~~layers.~~ Estop-Aragónés et al. (2012) found that oxic-anoxic interfaces in peat soil were located above the WT depth,  
and hence the capillary fringe ~~in this study~~ may have been still partly anoxic, ~~which can explain the correlation~~  
~~between N<sub>2</sub>O<sub>WT</sub> and N<sub>2</sub>O emissions in graphical models. In late April (DOY 113), NO<sub>2</sub><sup>-</sup> had accumulated at 20-50 cm~~  
~~depth at both RG/ and ARI sites (Figure 2), suggesting that there was an imbalance between ammonia oxidation and~~  
~~nitrite oxidation activity.~~ Oxygen affinity differs between nitrifiers, with AOA>AOB>NOB (Yin et al., 2018), and


620 ~~oxygen~~hence  $O_2$  limitation could ~~thus~~ have caused the accumulation of  $NO_2^-$ . In acid soil, this would result in product inhibition by  $HNO_2$ ; if there were no mechanism to remove  $NO_2^-$ , this would be especially true for *AR* sites, where mineral N accumulation was three to four times higher compared to *RG* sites (Tables S3-S6). Nitrifier-denitrification is ~~one~~a mechanism by which ammonia oxidisers can avoid  $HNO_2$  accumulation, and this process leads to  $N_2O$  formation (Braker and Conrad, 2011). Another potential sink for  $NO_2^-$  is chemodenitrification, an abiotic reaction in which  $NO_2^-$  reacts with  $Fe^{2+}$  to produce  $N_2O$  (Jones et al., 2015):




where in Eq. 34  $Fe(OH)_3$  is shown as anhydrous  $FeOOH$ . ~~Some~~A possible depletion of TRFe was indicated at 50 cm depth at site *ARI*, which coincided with a similar ~~depletion in pattern for~~  $NO_2^-$  (Figure 2). Nitrifier-denitrification and chemodenitrification are both sinks for  $NO_2^-$ , and therefore both pathways are potential sources of ~~the~~  $N_2O$  emissions ~~observed~~ during early spring.

~~The observation~~At *AR* sites there was often considerable accumulation of  $N_2O$  below the WT, which suggests ~~there was also an anaerobic pathway of  $N_2O$  formation. The fact~~ that TRFe concentrations were much higher than those of *AVS* or *CRS* (Table 1) makes it relevant to consider alternative reactions involving iron oxides/hydroxides, which have a potential to produce  $N_2O$ . One such recently described pathway is Feammox, a process whereby ammonia

635 oxidation coupled with ferric iron reduction can produce  $NO_2^-$  below pH 6.5 (Yang et al., 2012):



Nitrate can also be produced under these conditions (Yang et al., 2012; Guan et al., 2018):



A shuttle of  $Fe^{2+}$  between Feammox and chemodenitrification (Eq. 35 and Eq. 4) could ~~explain~~account for the accumulation of  $N_2O$  under anoxic conditions in the saturated zone, ~~where~~presumably ~~with~~ the availability of  $NH_4^+$  from peat mineralisation ~~would be~~a limiting factor. The confirmation of pathways will require more detailed investigations that should include molecular analyses targeting microbial communities in the soil profile.

## 5 Conclusion

645 ~~Nitrous oxide~~In this one-year study,  $N_2O$  emissions were ~~clearly~~consistently higher from arable sites compared to rotational grass. ~~This was independent of fertilisation~~There were strong seasonal dynamics in  $N_2O$  emissions, and ~~instead  $N_2O$  emissions could be associated with soil N mineralisation, rainfall patterns and temperature, as hypothesized~~we present evidence that different pathways were involved. Concentrations of pyrite were low compared to the total reduction capacity of the peat, and Fe was predominantly in forms other than pyrite. ~~While the~~The hypothesis, that  ~~$N_2O$  was produced by  $NO_3^-$  reduction coupled with  $FeS_2$  oxidation, could not be dismissed, it is likely that other processes were more~~ was an important. ~~There were strong seasonal dynamics in  $N_2O$  source of  $N_2O$  could therefore not be confirmed. Nitrous oxide emissions, and evidence that different pathways were involved. We propose~~



that oxidation of N mineralised from decomposing peat after WT drawdown in during spring were independent of fertilisation, since there was followed by mostly no effect of mineral N in the top soil. The significant effect of N<sub>2</sub>O concentration in the capillary fringe indicated that emissions during spring, and for grassland during the autumn, were associated with soil N mineralisation in this environment, as modified by rainfall patterns and WT dynamics. We propose that chemodenitrification (or nitrifier-denitrification) of NO<sub>2</sub><sup>-</sup> produced in the capillary fringe is a main source of N<sub>2</sub>O in acid organic soil during spring, whereas in the autumn, where NO<sub>3</sub><sup>-</sup> accumulated in arable soil after harvest, N<sub>2</sub>O emissions were associated with rising WT and heterotrophic denitrification as the can be a main pathway in arable soil as a result of NO<sub>3</sub><sup>-</sup> accumulation. Mitigating N<sub>2</sub>O emissions from acid organic soil is challenged by the complexity of underlying processes. However, reducing mineral N accumulation by ensuring a vegetation cover outside the main cropping season, and stabilising the WT depth by effective drainage, are potential mitigation strategies.

*Author contributions.* ATT, LEL, TJ and SOP designed the study. ATT, LEL, VE and SOP carried out sampling and analyses. ATT, RL and SOP were responsible for data analyses. ATT and SOP prepared the manuscript with contributions from all co-authors.

*Acknowledgements.* This study received financial support from the Danish Research Council for the project “Sources of N<sub>2</sub>O in arable organic soil as revealed by N<sub>2</sub>O isotopomers” (DFF – 4005-00448). We would like to thank the dedicated staff involved in field campaigns, including Bodil Stensgaard, Søren Erik Nissen, Sandhya Karki, Kim Johansen, Karin Dyrberg, Holger Bak and Stig T. Rasmussen. We would also like to acknowledge the support of three farmers hosting the field sites: Poul-Erik Birkbak, Rasmus Christensen and Jørn Christiansen.

## References

- Abreu, G.C.G., Edwards, D., and Labouriau, R.: High-dimensional graphical model search with the gRapHD R package. *J. Stat. Softw.*, 37, 1-18, doi: 10.18637/jss.v037.i01., 2010
- Aerts, R., and Ludwig, F.: Water-table changes and nutritional status affect trace gas emissions from laboratory columns of peatland soils. *Soil Biol. Biochem.*, 29, 1691-1698, doi: 10.1016/S0038-0717(97)00074-6, 1997
- Barton, L., Kiese, A., Gatter, D., Butterbach-Bahl, K., Buck, R., Hinz, C., and Murphy, D. V.: Nitrous oxide emissions from a cropped soil in a semi-arid climate. *Glob. Change Biol.*, 14, 177-192, doi: 10.1111/j.1365-2486.2007.01474.x, 2008.
- Braker, G., and Conrad, R.: Diversity, structure, and size of N<sub>2</sub>O-producing microbial communities in soils—what matters for their functioning? *Adv. Appl. Microbiol.*, 75, 33-70, doi: 10.1016/B978-0-12-387046-9.00002-5, 2011.
- Burton, E.D., Sullivan, L.A., Bush, R.T., Johnston, S.G., and Keene, A.F.: A simple and inexpensive chromium-reducible sulfur method for acid-sulfate soils. *Appl. Geochem.*, 23, 2759-2766, doi: doi.org/10.1016/j.apgeochem.2008.07.007, 2008.

690 Carter, M.R., and Gregorich, E.G., (Eds.): Soil sampling and methods of analysis, Second edition, CRC Press, USA  
2007.

Cline, J.D.: Spectrophotometric determination of hydrogen sulfide in natural waters. *Limnol. Oceanogr.*, 14, 454-458,  
doi: 10.4319/lo.1969.14.3.0454, 1969.

695 Clough T.J., Jarvis S.C., Dixon E.R., Stevens R.J., Laughlin R.J., and Hatch D.J.: Carbon induced subsoil  
denitrification of <sup>15</sup>N-labelled nitrate in 1 m deep soil columns. *Soil Biol. Biochem.* 31, 31-41,  
doi.org/10.1016/S0038-0717(98)00097-2, 1999.

Conen, F., and Smith, K.A.: A re-examination of closed flux chamber methods for the measurement of trace gas  
emissions from soils to the atmosphere. *Eur. J. Soil Sci.*, 49, 701-707, doi: 10.1046/j.1365-  
2389.1998.4940701, 1998.

700 Davidson, A.C., Hinkley, D.V.: *Bootstrap Methods and their Application*, 1st ed. Cambridge University Press, New  
York, NY, 1997.

Duan, Y.F. Kong, X.-W., Schramm, A., Labouriau, R., Eriksen, J., and Petersen, S.O.: Microbial N transformations and  
N<sub>2</sub>O emission after simulated grassland cultivation: effects of the nitrification inhibitor 3,4-Dimethylpyrazole  
Phosphate (DMPP). *Appl. Environ. Microbiol.*, 83, e02019-16, doi: 10.1128/AEM.02019-16, 2017.

705 Eickenscheidt, T., Heinichen, J., and Drösler, M.: The greenhouse gas balance of a drained fen peatland is mainly  
controlled by land-use rather than soil organic carbon content. *Biogeosciences*, 12, 5161-5184, doi:  
https://doi.org/10.5194/bg-12-5161-2015, 2015.

Estop-Aragónés, C., Knorr, K.-H., and Blodau, C.: Controls on *in situ* oxygen and dissolved inorganic carbon dynamics  
in peats of a temperate fen. *J. Geophys. Res.*, 117, G02002, doi: 10.1029/2011JG001888, 2012.

710 Gillham, R.W.: The capillary fringe and its effect on water-table response. *J. Hydrol.*, 67, 307-324,  
doi.org/10.1016/0022-1694(84)90248-8, 1984.

Guan, Q.S., Cao, W.Z., Wang, G.J., Wu, F.F., Wang, C., Jiang, C., Tao, Y.R., and Gao, Y.: Nitrogen loss through  
anaerobic ammonium oxidation coupled with iron reduction in a mangrove wetland. *Eur. J. Soil Sci.*, 69, 732-  
741, doi: 10.1111/ejss.12552, 2018.

715 Goldberg, S.D., Knorr, K.-H., and Gebauer, G.: N<sub>2</sub>O concentration and isotope signature along profiles provide deeper  
insight into the fate of N<sub>2</sub>O in soils. *Isot. Environ. Health Stud.*, 44, 377-391, doi:  
dx.doi.org/10.1080/10256010802507433, 2008.

Goldberg, S.D., Knorr, K.-H., Blodau, C., Lischeid, G., and Gebauer, G.: Impact of altering the water table height of an  
acidic fen on N<sub>2</sub>O and NO fluxes and soil concentrations. *Glob. Change Biol.*, 16, 220-233, doi:  
10.1111/j.1365-2486.2009.02015.x, 2010.

720 Good, P.I.: *Permutation, Parametric and Bootstrap Tests of Hypotheses. Third Edition. Springer, New York, 2005.*

Haughton, D.M.A.: On the choice of a model to fit data from an exponential family. *Ann. Statist.*, 16, 342-335, doi:  
10.1214/aos/1176350709, 1988.

725 Herold, M.B., Baggs, E.M., and Daniell, T.J.: Fungal and bacterial denitrification are differently affected by long-term  
pH amendment and cultivation of arable soil. *Soil Biol. Biochem.*, 54, 25-35, doi:  
doi.org/10.1016/j.soilbio.2012.04.031, 2012.

- Herrmann, M., Hädrich, A., and Küsel, K.: Predominance of thaumarchaeal ammonia oxidizer abundance and transcriptional activity in an acidic fen. *Env. Microbiol.*, 14, 3013-3025, doi: 10.1111/j.1462-2920.2012.02882.x, 2012.
- 730 Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Jamsranjav, B., Fukuda, M. & Troxler, T.: Supplement to the 2006 guidelines for national greenhouse gas inventories: wetlands. Intergovernmental Panel on Climate Change (IPCC), Geneva, Switzerland, 354 pp, 2014.
- Hoag, R.S., and Price J.S.: The effects of matrix diffusion on solute transport and retardation in undisturbed peat in laboratory columns. *J Contam Hydrol* 28:193-205, 1997.
- 735 Hutchinson, G.L., and Mosier, A.R.: Improved soil cover method for field measurement of nitrous oxide fluxes. *Soil Sci. Soc. Am. J.*, 45, 311-316, doi: 10.2136/sssaj1981.03615995004500020017x, 1981.
- IPCC: 2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands, Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M. and Troxler, T.G. (eds). Published: IPCC, Switzerland, 2014.
- Jones, L.C., Peters, B., Pacheco, J.S.L., Casciotti, K.L., and Fendorf, S.: Stable isotopes and iron oxide mineral products as markers of chemodenitrification. *Environ. Sci. Technol.*, 49, 3444-3452, doi: 10.1021/es504862x, 2015.
- 740 Jury W.A., Letey J., and Collins T.: Analysis of chamber methods used for measuring nitrous oxide production in the field. *Soil Sci. Soc. Am. J.* 46, 250–256, doi:10.2136/sssaj1982.03615995004600020007x, 1982.
- Jørgensen, B., and Labouriau, R.: Exponential families and theoretical inference. Springer, Monografias de Matemática, Rio de Janeiro, Brazil, 2012.
- 745 Jørgensen, C.J., Jacobsen, O.S., Elberling, B., and Aamand, J.: Microbial oxidation of pyrite coupled to nitrate reduction in anoxic groundwater sediment. *Environ. Sci. Technol.* 43, 4851–4857, doi: 10.1021/es803417s, 2009.
- Kandel, T.P., Lærke, P.E., and Elsgaard, L.: Annual emissions of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from a temperate peat bog: Comparison of an undrained and four drained sites under permanent grass and arable crop rotations with cereals and potato. *Agric. Forest Meteorol.* 256-257, 470-481, doi: doi.org/10.1016/j.agrformet.2018.03.021, 2018.
- 750 Keeney, D.R., and Nelson, D.W.: Nitrogen–inorganic forms. In: A.L. Page, T.H. Miller and D.R. Keeney (Eds.), *Methods of Soil Analysis. Part 2. Agronomy Monographs*, 9. American Society of Agronomy and Soil Science Society of America, Madison, WI, pp. 643-692, 1982.
- 755 Kits, K. D., Jung, M.Y., Vierheilig, J., Pjevac, P., Sedlacek, C.J., Liu, S., Herbold, C., Stein, L.Y., Richter, A., Wissel, H., Brüggemann, N., Wagner, M., and Daims, H.: Low yield and abiotic origin of N<sub>2</sub>O formed by the complete nitrifier *Nitrospira inopinata*. *Nature Comm.* 10, 1836, doi: //doi.org/10.1038/s41467-019-09790-x, 2019.
- Kristensen, M. K.: Vildmosearbejdet. Det Danske Forlag, Copenhagen, Denmark, 219 pp. (in Danish), 1945.
- Labouriau, R., and Amorim, A.: Comment on ‘An Association Between the Kinship and Fertility of Human Couples’. *Science*, 322 (5908):1634, doi: 10.1126/science.1161907, 2008a.
- 760 Labouriau, R., and Amorim, A.: Human fertility increases with marital radius. *Genetics*, 178, 601-603, doi: 10.1534/genetics.107.072454 , 2008b.

Lamandé, M., Labouriau, R., Holmstrup, M., Torp, S. B., Heckrath, G., Iversen, B. V., and Jacobsen, O.H.: Density of macropores as related to soil and earthworm community parameters in cultivated grasslands. *Geoderma*, 162, 319-326, doi: doi.org/10.1016/j.geoderma.2011.03.004, 2011.

Lauritzen, S. L.: Causal Inference from Graphical Models. In: Complex Stochastic Systems, O. E. Barndorff-Nielsen, D. R. Cox and C. Klüppelberg (Editors), New York, 1999.

Leppelt, T., Dechow, R., Gebbert, S., Freibauer, A., Lohila, A., Augustin, J., Drösler, M., Fiedler, S., Glatzel, S., Höper, H., Järveoja, J., Lærke, P. E., Maljanen, M., Mander, Ü., Mäkiranta, P., Minkinen, K., Ojanen, P., Regina, K., and Strömberg, M.: Nitrous oxide emission budgets and land-use-driven hotspots for organic soils in Europe. *Biogeosci.*, 11, 6595-6612, doi: doi.org/10.5194/bg-11-6595-2014, 2014.

Liu, B., Frostegård, Å., and Bakken, L.R.: Impaired reduction of N<sub>2</sub>O to N<sub>2</sub> in acid soils is due to a posttranscriptional interference with the expression of nosZ. *MBio*, 5, e01383-14, doi: 10.1128/mBio.01383-14, 2014.

Madsen, H.B., and Jensen, N.H.: Potentially acid sulfate soils in relation to landforms and geology. *Catena*, 15, 137-145, doi: doi.org/10.1016/0341-8162(88)90025-2, 1988.

Maeda, K., Spor, A., Edel-Hermann, V., Heraud, C., Breuil, M. -C., Bizouard, F., Toyoda, S., Yoshida, N., Steinberg, C., and Philippot, L.: N<sub>2</sub>O production, a widespread trait in fungi. *Nature Sci. Rep.*, 5, 9697, doi: 10.1038/srep09697, 2015.

Maljanen, M., Liikanen, A., Silvola, J., and Martikainen, P.J.: Nitrous oxide emissions from boreal organic soil under different land-use. *Soil Biol. Biochem.*, 35, 1-12, doi: doi.org/10.1016/S0038-0717(03)00085-3, 2003.

Mander, Ü., Uuemaa, E., Kull, A., Kanal, A., Maddison, M., Soosaar, K., Salm, J.-O., Lesta, M., Hansen, R., Kuller, R., Harding, A., and Augustin, J.: Assessment of methane and nitrous oxide fluxes in rural landscapes. *Landscape Urban Plan.*, 98, 172-181, doi: doi.org/10.1016/j.landurbplan.2010.08.021, 2010.

Martikainen, P.J., Nykanen, H., Crill, P., and Silvola, J.: Effect of a lowered water-table on nitrous-oxide fluxes from northern peatlands. *Nature*, 366, 51-53, doi: 10.1038/366051a0, 1993.

McCullagh, P., and Nelder, J.A.: Generalized Linear Models, Second Edition. Chapman and Hall/CRC, London, 1989.

Mu, Z., Huang, A., Ni, J., and Xie, D.: Linking annual N<sub>2</sub>O emission in organic soils to mineral nitrogen input as estimated by heterotrophic respiration and soil C/N ratio. *PLOS ONE*, 9, e96572, doi: doi.org/10.1371/journal.pone.0096572, 2014.

Palomo, A., Dechesne, A., and Smets, B.F.: Genomic profiling of *Nitrospira* species reveals ecological success of comammox *Nitrospira*. *bioRxiv* doi: <http://dx.doi.org/10.1101/612226>, 2019.

Pedersen, A.R., Petersen, S.O., and Schelde, K.: A comprehensive approach to soil-atmosphere trace-gas flux estimation with static chambers. *Eur. J. Soil Sci.*, 61, 888-902, doi: 10.1111/j.1365-2389.2010.01291.x, 2010.

Petersen, S.O.: Diffusion probe for gas sampling in undisturbed soil. *Eur. J. Soil Sci.*, 65, 663-671, doi: 10.1111/ejss.12170, 2014.

Petersen, S.O., Hoffmann, C. C., Schäfer, C.-M., Blicher-Mathiesen, G., Elsgaard, L., Kristensen, K., Larsen, S.E., Torp, S.B., and Greve, M.H.: Annual emissions of CH<sub>4</sub> and N<sub>2</sub>O, and ecosystem respiration, from eight organic soils in Western Denmark managed by agriculture. *Biogeosci.*, 9, 403-422, doi: doi.org/10.5194/bg-9-403-2012, 2012.

R Core Team: R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Vienna, Austria. <https://www.R-project.org/>, 2016.

- Regina, K., Budiman, A., Greve, M.H., Grønlund, A., Kasimir, Å., Lehtonen, H., Petersen, S.O., Smith, P., and Wösten, H., 2015. GHG mitigation of agricultural peatlands requires coherent policies. *Clim. Policy*, 16, 522–541, doi: doi.org/10.1080/14693062.2015.1022854, 2016.
- 805 Regina, K., Nykanen, H., Silvola, J., and Martikainen, P.J.: Fluxes of nitrous oxide from boreal peatlands as affected by peatland type, water table level and nitrification capacity. *Biogeochem.*, 35, 401–418, doi: doi.org/10.1007/BF02183033, 1996.
- Regina, K., Syväsalö, E., Hannukkala, A., and Esala, M.: Fluxes of N<sub>2</sub>O from farmed peat soils in Finland. *Eur. J. Soil Sci.*, 55, 591–599, doi: doi.org/10.1111/j.1365-2389.2004.00622.x, 2004.
- 810 Rezanezhad, F., Price, J.S., Quinton, W.L., Lennartz, B., Milojevic, T., and Van Cappellen, P.: Structure of peat soils and implications for water storage, flow and solute transport: A review update for geochemists. *Chem. Geol.*, 429, 75–84, <http://dx.doi.org/10.1016/j.chemgeo.2016.03.010>, 2016.
- Røy, H., Weber, H.S., Tarpgaard, I.H., Ferdelman, T.G., and Jørgensen, B.B.: Determination of dissimilatory sulfate reduction rates in marine sediment via radioactive <sup>35</sup>S tracer. *Limnol. Oceanogr. Methods*, 12, 196–211, doi: 10.4319/lom.2014.12.196, 2014.
- 815 Schothorst, C.J.: Subsidence of low moor peat soils in the western Netherlands. *Geoderma*, 17, 265–291, doi: doi.org/10.1016/0016-7061(77)90089-1, 1977.
- Schäfer, C.-M., Elsgaard, L., Hoffmann, C.C. and Petersen, S.O.: Seasonal methane dynamics in three temperate grasslands on peat. *Plant Soil*, 357, 339–353, doi: doi.org/10.1007/s11104-012-1168-9, 2012.
- 820 Spott, O., Russow, R., and Stange, C.F.: Formation of hybrid N<sub>2</sub>O and hybrid N<sub>2</sub> due to codenitrification: First review of a barely considered process of microbially mediated N-nitrosation. *Soil Biol. Biochem.*, 43, 1995–2011, doi: doi.org/10.1016/j.soilbio.2011.06.014, 2011.
- Stieglmeier, M., Mooshammer, M., Kitzler, B., Wanek, W., Zechmeister-Boltenstern, S., Richter, A., and Schleper, C.: Aerobic nitrous oxide production through N-nitrosating hybrid formation in ammonia-oxidizing archaea. *ISME J.*, 8, 1135–1146, doi: 10.1038/ismej.2013.220, 2014.
- 825 Stopnišek, N., Gubry-Rangin, C., Höfferle, S., Nicol, G. W., Mandic-Mulec, I., and Prosser, J.I.: Thaumarchaeal ammonia oxidation in an acidic forest peat soil is not influenced by ammonium amendment. *Appl. Environ. Microb.*, 76, 7626–7634, doi: 10.1128/AEM.00595-10, 2010.
- Taft, H.E., Cross, P.A., Edwards-Jones, G., Moorhouse, E.R., and Jones, D.L.: Greenhouse gas emissions from intensively managed peat soils in an arable production system. *Agr. Ecosyst. Environ.*, 237, 162–172, doi: doi.org/10.1016/j.agee.2016.11.015, 2017.
- 830 Taghizadeh-Toosi, A., [Clough, T.](#), Petersen, S.O., Clough, T., and Elsgaard, L.: Nitrous oxide (N<sub>2</sub>O) turnover [dynamics](#) in agricultural peat soil [-in response to availability](#) role of nitrate, nitrite, and iron sulfides. [Submitted Geomicrobiol. J. doi: doi.org/10.1080/01490451.2019.1666192, 2019.](#)
- 835 Thamdrup, B., Fossing, H. and Jørgensen, B.B.: Manganese, iron, and sulfur cycling in a coastal marine sediment, Aarhus Bay, Denmark. *Geochim. Cosmochim. Acta*, 58, 5115–5129, doi: 10.1016/0016-7037(94)90298-4, 1994.
- Torrento C, Cama J, Urmeneta J, Otero N, and Solar A.: Denitrification of groundwater with pyrite and *Thiobacillus denitrificans*. *Chem. Geol.*, 278, 80–91, doi: dx.doi.org/10.1016/j.chemgeo.2010.09.003, 2010.

Field Code Changed

- 840 Toyoda, S., Yoshida, N., and Koba, K.: Isotopocule analysis of biologically produced nitrous oxide in various environments. *Mass Spectrometry Reviews* 36, 135-160, doi: doi.org/10.1002/mas.21459, 2017.
- Tubiello, F.N., Biancalani, R., Salvatore, M., Roissi, S., and Conchedda, G.: A worldwide assessment of greenhouse gas emissions from drained organic soils. *Sustainability*, 8, 371, 13 pp, doi: 10.3390/su8040371, 2016.
- 845 Ulrich, G.A., Krumholz, L.R., and Suflita, J.M.: A rapid and simple method for estimationg sulfate reduction activity and quantifying inorganic sulfides. *Appl. Environ. Microbiol.*, 63, 1627-1630, 1997.
- Van Cleemput, O. and Samater, A.H.: Nitrite in soils: Accumulation and role in the formation of gaseous N compounds. *Fert. Res.*, 45, 81-89, doi: doi.org/10.1007/BF00749884, 1996.
- Viollier, E., Inglett, P.W., Hunter, K., Roychoudhury, A.N., and Van Cappellen, P.: The ferrozine method revisited: Fe(II)/Fe(III) determination in natural waters. *Appl. Geochem.*, 15, 785-790, doi: doi.org/10.1016/S0883-2927(99)00097-9, 2000.
- 850 Whittaker, J.: *Graphical models in applied multivariate statistics*. John Wiley & Sons, Chichester, UK, 1990.
- Wrage-Mönnig, N., Horn, M.A., Well, R., Müller, C., Velthof, G., and Oenema, O.: The role of nitrifier denitrification in the production of nitrous oxide revisited. *Soil Biol. Biochem.* 123, A3-A16, doi:doi.org/10.1016/j.soilbio.2018.03.020, 2018.
- 855 Yang, W.H., Weber, K.A., and Silver, W.L.: Nitrogen loss from soil through anaerobic ammonium oxidation coupled to iron reduction. *Nat. Geosci.*, 5, 538-541, doi: 10.1038/ngeo1530, 2012.
- Yin, Z., Bi, X., and Xu, C.: Ammonia-oxidizing archaea (AOA) play with ammonia-oxidizing bacteria (AOB) in nitrogen removal from wastewater. *Archaea* Article ID 8429145, 9 pages, doi: doi.org/10.1155/2018/8429145, 2018.

860

**Table 1.** Selected characteristics of soil profiles at the four monitoring sites with rotational grass (*RG1*, *RG2*) and potato crop (*AR1*, *AR2*). All analyses were done in triplicate; results shown represent mean and standard error of two soil profiles (*n* = 2) for which a complete data set was available. Soils for analyses were collected in late April (DOY 113) except for AVS and CRS (early September). Abbreviations: EC, electrical conductivity; TOC, soil organic carbon; TRFe, total reactive iron; AVS, acid volatile sulfide; CRS, chromium reducible sulfur.

	Depth (cm)	pH	EC	TOC (g 100 g <sup>-1</sup> )	Total N (g 100 g <sup>-1</sup> )	C:N ratio	TRFe (mg Fe g <sup>-1</sup> )	AVS (μg S g <sup>-1</sup> )	CRS (μg S g <sup>-1</sup> )
<b><i>RG1</i></b>									
Depth 1	2.5-7.5	5.1 (0.2)	0.26 (0.10)	37.4 (0.2)	1.75 (0.00)	21.3	3.63 (0.11)	2.51 (0.86)	155 (62)
Depth 2	7.5-12.5	5.3 (0.1)	0.15 (0.02)	38.2 (0.2)	1.79 (0.01)	21.3	4.03 (0.44)	NA	NA
Depth 3	17.5-22.5	5.3 (0.5)	0.37 (0.18)	39.7 (0.3)	1.80 (0.04)	22.1	4.14 (0.32)	NA	NA
Depth 4	36-40	4.8 (0.1)	0.55 (0.02)	43.1 (2.7)	1.85 (0.03)	23.3	3.04 (0.26)	2.60 (0.87)	133 (64)
Depth 5	47.5-52.5	5.1 (0.3)	0.42 (0.13)	31.0 (15.6)	1.47 (0.64)	21.1	2.50 (0.55)	4.86 (1.07)	24 (17)
Depth 6	93-98	5.4 (0.0)	0.51 (0.06)	0.6 (0.3)	0.01 (0.01)	ND	0.14 (0.04)	NA	NA
<b><i>RG2</i></b>									
Depth 1	0-25	5.0	NA	19.8 (3.4)	1.34 (0.13)	14.8	2.29 (0.56)	NA	NA
Depth 2	25-50	5.1	NA	8.9 (3.0)	0.63 (0.23)	14.2	4.48 (NA)	1.71 (0.00)	33 (7.3)
<b><i>AR1</i></b>									
Depth 1	2.5-7.5	5.0 (0.1)	0.45 (0.04)	35.9 (0.1)	1.81 (0.02)	19.9	4.57 (0.09)	1.74 (0.02) <sup>§</sup>	141 (9)
Depth 2	7.5-12.5	5.2 (0.1)	0.42 (0.06)	34.2 (0.2)	1.76 (0.02)	19.4	4.66 (0.15)	NA	NA
Depth 3	17.5-22.5	5.2 (0.1)	0.34 (0.04)	41.0 (2.2)	1.93 (0.11)	21.3	4.99 (0.43)	NA	NA
Depth 4	36-40	4.7 (0.5)	0.37 (0.05)	41.1 (5.8)	1.84 (0.05)	22.4	3.23 (0.41)	2.17 (0.29)	49 (3)
Depth 5	47.5-52.5	4.7 (0.3)	0.48 (0.08)	5.9 (1.7)	0.37 (0.13)	16.3	1.19 (0.19)	1.98 (0.41)	137 (39)
Depth 6	93-98	5.4 (0.2)	0.91 (0.03)	0.3 (0.1)	0.00 (0.00)	ND	0.18 (0.02)	NA	NA
<b><i>AR2</i></b>									
Depth 1	0-25	5.1	NA	33.4 (1.2)	1.45 (0.03)	23.1	4.11 (0.03)	NA	NA
Depth 2	25-50	5.1	NA	38.4 (0.2)	1.46 (0.02)	26.2	3.78 (0.14)	1.65 (0.02)	45 (8)

ND – Not determined due to TOC and total N concentrations being at the limit of detection.

NA - Not analysed.

<sup>§</sup> Taghizadeh-Toosi et al. (2019) presented slightly different AVS and CRS results for *AR1* representing all the six profiles collected at this site.

Formatted Table

Deleted Cells

Deleted Cells

Deleted Cells

Deleted Cells

Deleted Cells

Deleted Cells

Deleted Cells

Formatted: Centered

**Table 2.** CumulativeDaily emissions of N<sub>2</sub>O (kg N<sub>2</sub>O ha<sup>-1</sup>) during the monitoring periods in spring (99-105 days) and autumn (47-69 days) monitoring period. Estimation for each season was performed using the trapezoidal approximation of the integral of the emission curve. Numbers in parentheses indicate 95% confidence intervals, and significant differences, (at 5% significance level), corrected for multiple testing by the single-step method, are indicated by asterisks. RG, rotational grass; AR, arable crop (potato); F, fertilised; NF, unfertilised letters.

	DOY	Cumulative N <sub>2</sub> O	RG-NF	Spring RG-F	AR-NF	Autumn		
Spring <sup>#</sup>	DOY		kg N <sub>2</sub> O-N ha <sup>-1</sup> d <sup>-1</sup>			g N <sub>2</sub> O-N ha <sup>-1</sup> d <sup>-1</sup>		
Rotational grass	RG1	RG-NF <sup>#</sup>	63-162 161	2.0-29 36	a	251-313 20	(16-25) a	
		RG-F	63-162 7.3-31	37	***a	15	(11-18) a	
	RG2	AR-NF	63-167 168	23	(17-28) 20.2)a	306	***50	(34-66) b
		AR-F	63-167 45.0-146	197	***bc	NS51	(39-63) b	
Autumn								
Arable	AR1	RG-NF	252-314 166	2.0-303 377	d	245-307 338	(239-437) d	
		RG-F	252-314 4.9-234	288	NScd	322	(224-421) d	
	AR2	AR-NF	246-314 168	43.6-102 120	***b	313	105	(70-141) bc
		AR-F	246-314 45.3-105	128	***b	NS127	(90-165) c	

\*\*\*,  $p < 0.001$ ; NS, not significant ( $p > 0.05$ )

<sup>#</sup>The monitoring periods (spring and autumn) were: DOY63-162 and DOY252-314 (RG1); DOY64-169 and DOY260-307 (RG2); DOY63-162 and DOY246-308 (AR1); DOY64-169 and DOY245-314 (AR2). <sup>#</sup> F, fertilised; NF, not fertilised.





## Figure captions

**Figure 1.** A. Location of sites *AR1* and *RG1* (both at 57°13'59.7"N, 9°50'40.3E), *RG2* (57°13'55.9"N, 9°52'20.2E) and *AR2* (57°13'7.6"N, 9°46'26.9E). B. Experimental design at each of the four sites, with three blocks centered around piezometers (●) and two subplots, one of which received N fertiliser at the rate of the surrounding field. Six collars for gas flux measurements (S1-S6) were distributed as indicated, and sets of 5 diffusion probes for soil gas sampling were installed near collars in selected positions (see text).

**Figure 2.** Nitrite-N (a, c) and total reactive iron, TRFe (b, d), in undisturbed soil cores were collected at sites *RG1* and *AR1* on 23 April (DOY113DOY 113; white symbols) and 2 September (DOY245DOY 245; grey symbols). Results shown are mean and standard error ( $n = 2$ ). At site *RG1* fertilisation had taken place one week earlier, but a two-sample *t* test did not find evidence for any effect on  $\text{NO}_2^-$  availability ( $p = 0.19$ ). The dotted lines indicate WT level on the two sampling dates.

**Figure 3.** The top panel shows rainfall, air temperature and management (F – fertilisation) at sites *RG1* (left panels) and *RG2* (right panels) during spring, 3 March (DOY63DOY 63) to 16 June (DOY169DOY 169). The middle section shows  $\text{N}_2\text{O}$  fluxes (black circles; mean  $\pm$  standard error,  $n = 3$ ) and contour plots of soil  $\text{N}_2\text{O}$  concentrations in fertilised subplots, and the lower section the corresponding results for unfertilised subplots. A logarithmic grey scale was used in order to show trends within both *RG* and *AR* treatments, and between depths. Soil gas sampling positions are indicated in the contour plots; numbers shown are  $\text{N}_2\text{O}$  concentrations ( $\mu\text{L L}^{-1}$ ). Green lines show the WT depth (which varied slightly between blocks). B2 and B3 refer to block number of diffusion probe positions.

**Figure 4.** The top panel shows rainfall, air temperature and management (T – tillage; F – fertilisation) at sites *AR1* (left panels) and *AR2* (right panels) during spring, 3 March (DOY63DOY 63) to 16 June (DOY169DOY 169). The middle section shows  $\text{N}_2\text{O}$  fluxes (black circles; mean  $\pm$  standard error,  $n = 3$ ) and contour plots of soil  $\text{N}_2\text{O}$  concentrations in fertilised subplots, and the lower section the corresponding results for unfertilised subplots. A logarithmic grey scale was used in order to show trends within both *RG* and *AR* treatments, and between depths. Soil gas sampling positions are indicated in the contour plots; numbers shown are  $\text{N}_2\text{O}$  concentrations ( $\mu\text{L L}^{-1}$ ). Gaps are indicated where soil gas sampling probes were installed late, or removed due to field operations. Green lines show the WT depth (which varied slightly between blocks). B2 and B3 refer to block number of diffusion probe positions.

**Figure 5.** The top panel shows rainfall, air temperature and management (H - harvest) at sites *RG1* (left panels) and *RG2* (right panels) during autumn, 3 September (DOY245DOY 245) to 10 November (DOY314DOY 314). The middle section shows  $\text{N}_2\text{O}$  fluxes (black circles; mean  $\pm$  standard error,  $n = 3$ ) and contour plots of soil  $\text{N}_2\text{O}$  concentrations in fertilised subplots, and the lower section the corresponding results for unfertilised subplots. A logarithmic grey scale was used in order to show trends within both *RG* and *AR* treatments, and between depths. Soil gas sampling positions are indicated in the contour plots; numbers shown are  $\text{N}_2\text{O}$  concentrations ( $\mu\text{L L}^{-1}$ ); the probes were absent in the unfertilised subplot after harvest. Green lines show the WT depth (which varied slightly between blocks). B2 and B3 refer to block number of diffusion probe positions.

**Figure 6.** The top panel shows rainfall, air temperature and management (H - harvest) at sites *AR1* (left panels) and *AR2* (right panels) during autumn, 3 September (DOY245DOY 245) to 10 November (DOY314DOY 314). The middle section shows N<sub>2</sub>O fluxes (black circles; mean  $\pm$  standard error,  $n = 3$ ) and contour plots of soil N<sub>2</sub>O concentrations in fertilised subplots, and the lower section the corresponding results for unfertilised subplots. A logarithmic grey scale was used in order to show trends within both *RG* and *AR* treatments, and between depths. Soil gas sampling positions are indicated in the contour plots; numbers shown are N<sub>2</sub>O concentrations ( $\mu\text{L L}^{-1}$ ). Green lines show the WT depth (which varied slightly between blocks). B2 and B3 refer to block number of diffusion probe positions.

**Figure 7.** Using Results of a graphical models, a statistical analysis was conducted for each site crop combination of crop (*RG*, *AR*) and season (in spring, autumn). a. *RG*, spring; b. *RG*, autumn; c. *AR*, spring; and d. *AR*, autumn. A. RG1-Spring; B. RG2-Spring; C. AR1-Spring; and D. AR2-Spring. The edges (“lines”) connecting vertices (“points”) indicate significant relationships conditional correlation between explanatory variables and the response variable (N<sub>2</sub>O flux). Statistical given the other variables. The statistical results for direct effects on N<sub>2</sub>O flux are: [1] 2.32 (0.12-9.14) (0.02-0.23,  $p = 0.014067$ ); [2] 0.7413 (0.06-3.05,  $p = 0.034$ ); [3] 0.78 (0.41-2.47,  $p = 0.0002$ ); [4] 1.34 (0.78-4.08) ( $p = 0.008$ ); [3] 0.18 (0.09-0.26,  $p = 0.001$ ); [4] 0.15 (0.04-0.23,  $p = 0.032$ ); and [5] 2.45 (1.10-9.90) (0.12-0.49,  $p = 0.0002$ ). Key to variables: AmmoniumT: NH<sub>4</sub><sup>+</sup>-N at 0-25 cm depth; NitrateT: NO<sub>3</sub><sup>-</sup>-N at 0-25 cm depth; N<sub>2</sub>O WT: equivalent soil gas phase concentration closest to, but above the water table depth; Temp5: soil temperature at 5 cm depth; Temp30: soil temperature at 30 cm depth.

**Figure 8.** Results of a graphical model for each site crop combination in the autumn. A. RG1-Autumn; B. RG2-Autumn; C. AR1-Autumn; and D. AR2-Autumn. The edges (“lines”) connecting vertices (“points”) indicate significant conditional correlation between the variables given the other variables. Statistical results for effects on N<sub>2</sub>O flux are: [1] 0.27 (0.19-0.38,  $p = 0.002$ ); [2] 0.16 (0.007-0.28,  $p = 0.049$ ); [3] 0.19 (0.09-0.28,  $p = 0.021$ ) [4] 0.15 (0.06-0.21,  $p = 0.032$ ); and [5] 0.29 (0.17-0.35,  $p = 0.005$ ). Key to variables: AmmoniumT: NH<sub>4</sub><sup>+</sup>-N at 0-25 cm depth; NitrateT: NO<sub>3</sub><sup>-</sup>-N at 0-25 cm depth; N<sub>2</sub>O WT: equivalent soil gas phase concentration closest to, but above the water table depth; Temp5: soil temperature at 5 cm depth; Temp30: soil temperature at 30 cm depth.

Formatted: Font: Not Bold


Formatted: Space Before: 0 pt, After: 8 pt

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic





**Figure 1**





Figure 2.







Figure 32.

Formatted: Left





**Figure 3**




Figure 4

Commented [SOP3]:  
AR2: 71 d vlærði mangler (21 cm)




Figure 5










Figure 6





Commented [SOP4]: Udsiftes

Figure 7



**Figure 8**

