



## Characterizing photosymbiosis in modern planktonic foraminifera

Haruka Takagi<sup>1,2</sup>, Katsunori Kimoto<sup>3</sup>, Tetsuichi Fujiki<sup>3</sup>, Hiroaki Saito<sup>1</sup>, Christiane Schmidt<sup>4</sup>, Michal Kucera<sup>4</sup>, Kazuyoshi Moriya<sup>5</sup>

<sup>1</sup>Atmosphere and Ocean Research Institute, The University of Tokyo, Japan

5 <sup>2</sup>Graduate School of Creative Science and Engineering, Waseda University, Japan

<sup>3</sup>Japan Agency for Earth-Marine Science and Technology, Japan

<sup>4</sup>MARUM - Center for Marine Environmental Sciences and Faculty of Geosciences, University of Bremen, Germany

<sup>5</sup>Department of Earth Sciences, School of Education, Waseda University, Japan

*Correspondence to:* Haruka Takagi ([htakagi@aori.u-tokyo.ac.jp](mailto:htakagi@aori.u-tokyo.ac.jp), [harurah-t@fuji.waseda.jp](mailto:harurah-t@fuji.waseda.jp))

10 **Abstract.** Photosymbiosis has played a key role in the diversification of foraminifera and their carbonate production through geologic history. However, identification of photosymbiosis in extinct taxa remains challenging and even among the extant species the occurrence and functional relevance of photosymbiosis remains poorly constrained. Here, we investigate photosymbiosis in living planktonic foraminifera by measuring active chlorophyll fluorescence with fast repetition rate fluorometry. This method provides unequivocal evidence for the presence of photosynthetic capacity in individual foraminifera and it allows us to characterize multiple features of symbiont photosynthesis including chlorophyll *a* (Chl *a*) content, potential photosynthetic activity ( $F_v/F_m$ ), and light absorption efficiency ( $\sigma_{PSII}$ ). To obtain robust evidence for the occurrence and importance of photosymbiosis in modern planktonic foraminifera, we conducted measurements on 1266 individuals from 30 species of the families Globigerinidae, Hastigerinidae, Globorotaliidae, and Candeinidae. Among the studied species, 19 were recognized as symbiotic and 11 as non-symbiotic. Of these, six species were newly confirmed as symbiotic and five as non-symbiotic. Photosymbiotic species have been identified in all families except the Hastigerinidae. A significant positive correlation between test size and Chl *a* content, found in 16 species, is interpreted as symbiont growth scaled to the growth of the host, consistent with persistent possession of symbionts through the lifetime of the foraminifera. The remaining three symbiont-bearing species did not show such a relationship, and their  $F_v/F_m$  values were comparatively low, indicating that their symbionts do not grow once acquired from the environment. The objectively quantified photosymbiotic characteristics have been used to design a metric of photosymbiosis, which allows the studied species to be classified along a gradient of photosynthetic activity, providing a framework for future ecological and physiological investigations of planktonic foraminifera.

### 1 Introduction

Planktonic foraminifera are unicellular marine zooplankton with calcite tests. Since they are geographically widespread and abundant, and can be preserved in seafloor sediments as microfossils, foraminifera are one of the most important archives of surface ocean conditions in the past. They have been used to investigate pelagic marine biodiversity dynamics from middle



Mesozoic to the present (Bolli et al., 1985; Norris, 1991; Boudagher-Fadel et al., 1997; Hull, 2017; Yasuhara et al., 2017). Recent studies of macroevolutionary dynamics of planktonic foraminifera emphasized the importance of species ecology including photosymbiosis (endosymbiosis with autotrophic algae) as a key player determining temporal and spatial patterns of species diversity (Ezard et al., 2011; Fenton et al., 2016). However, identifying photosymbiosis in extinct species is difficult and requires indirect evidence such as size-dependent stable isotopic trends (e.g., Pearson et al., 1993; Norris, 1996). These indirect methods must be benchmarked by observations from living foraminifera, where the presence of symbionts can be determined directly. A knowledge on the prevalence, diversity, and phylogenetic position of photosymbiosis is also required to elucidate ecological and evolutionary strategies of the involved clades and to characterize key features of foraminiferal test geochemistry like  $\delta^{13}\text{C}$  and  $\delta^{11}\text{B}$  (e.g., Spero and DeNiro, 1987; Hönisch et al., 2003; Henehan et al., 2013; Ezard et al., 2015).

Photosymbiosis in modern planktonic foraminifera has been empirically identified based on microscopic observations of intracellular algae (Lee et al., 1965; Anderson and Bé, 1976; Gastrich, 1987), and molecular confirmation of algal DNA extracted from a single foraminifer cell (Gast and Caron, 1996; Gast et al., 2000; Shaked and de Vargas, 2006; Bird et al., 2017, 2018). As a result, among the ~ 50 species of modern planktonic foraminifera, thirteen have so far been reported to be photosymbiotic (*Orbulina universa*, *Globigerinoides sacculifer*, *Globigerinoides conglobatus*, *Globigerinoides ruber*, *Globigerinella siphonifera*, *Turborotalita humilis*, *Globigerina bulloides*, *Neogloboquadrina dutertrei*, *Pulleniatina obliquiloculata*, *Globorotalia inflata*, *Globorotalia menardii*, *Candeina nitida*, and *Globigerinita glutinata*), and six to be symbiont-barren (*Hastigerina pelagica*, *Globigerina bulloides*, *Globorotalia truncatulinoides*, *Globorotalia hirsta*, *Neogloboquadrina incompta*, and *Neogloboquadrina pachyderma*) (Table 1). The remaining 30 species have not been systematically examined for the presence of symbionts. In a strict sense, in all previous studies on a photosymbiotic association, the authors could not differentiate whether the intracellular algae they identified were symbionts or just captured preys to be digested. Although observations of features like mitosis (cell-division) of the intracellular algal cells are strong evidence that these were alive within the foraminifera, the presence of intracellular algae alone does not guarantee that they act as photosymbionts. Many foraminiferal species ingest phytoplankton prey (Anderson et al., 1979), which makes it difficult to differentiate symbionts or prey, especially by DNA analysis. Since many species of planktonic foraminifera do not survive well in culture, it is hard to conduct behavioral or physiological experiments to confirm their symbiosis. These limitations have hindered the progress of studies of photosymbiosis targeting various species of planktonic foraminifera.

One solution to identify functional photosymbiosis is to detect a physiological signature of photosynthesis within the cell. This has been done by measurements of oxygen production with microelectrodes (Jørgensen et al., 1985; Rink et al., 1998; Lombard et al., 2009) or a determination of photosynthetic carbon fixation by measurements of  $^{14}\text{C}$ -tracer (Spero and Parker, 1985; Gastrich and Bartha, 1988). These studies were limited to established symbiotic species that are easy to culture (e.g., *O. universa*, *G. sacculifer*, and *G. siphonifera*). For the other species, especially non-spinose species (e.g., *N. dutertrei*, *P. obliquiloculata*, and *G. glutinata*), the physiological characteristics of their photosymbiosis have never been described. Therefore, our knowledge of modern photosymbiosis has been exclusively obtained from a small number of spinose symbiotic species. A powerful alternative to directly and unambiguously determine the presence of active photosynthesis in the



foraminifera is given by measurement of fluorescence induced by light capture in the photosystem II of the algal chlorophyll. These methods have been used in benthic symbiont-bearing foraminifera (e.g., Uthicke, 2006; Schmidt et al., 2011; Ziegler and Uthicke, 2011) and recently successfully adapted for application on single specimens of living planktonic foraminifera (Fujiki et al., 2014; Takagi et al., 2016, 2018). An active chlorophyll fluorometry performs non-destructive and non-invasive measurements of algal physiology based on real-time variable fluorescence profiles (Kolber and Falkowski, 1993), allowing us to quantify chlorophyll *a* content of a specimen, the health of its symbionts and their light-level adaptation (Fujiki et al., 2014; Takagi et al., 2018). The measurements can be performed almost immediately after collection, with minimal manipulations, thus minimizing damage to the foraminifera and circumventing culturing stress-induced artifacts. This approach can make a breakthrough in the study of photosymbiosis, not just because of its versatility but its potential to provide key quantitative attributes of the photosymbiosis.

Symbiotic relationships in foraminifera have been previously categorized as either obligate or facultative (Hemleben et al., 1989). Obligate photosymbiosis is essential for the host, making it functionally mixotrophic, an adaptive strategy to live in oligotrophic and well-lit parts of the ocean (Hallock, 1981; Lee, 2006). In facultative symbiosis, the foraminifera is not dependent upon it for survival and as a result, symbiotic algae in facultative symbiosis will be only found in some specimens of the host species. Facultative associations generally do not involve extensive metabolic adaptation of the host and can thus enhance the flexibility of nutritional sources with minimal energetic investment (Stoecker et al., 2009). In planktonic foraminifera, species always found with intact intracellular algae has been regarded as obligate symbiotic species, whereas species sometimes found with but sometimes without them has been termed as facultative symbiotic species (Hemleben et al., 1989). However, most of our knowledge of foraminiferal photosymbiosis is based on indirect evidence, insufficient to categorize planktonic foraminiferal photosymbiosis as either obligate or facultative. Rather, the persistence and functional relevance of the symbiotic relationship through a foraminiferal lifetime should be determined anew, using direct measurements, allowing us to correctly understand the function of each specific photosymbiotic relationship.

Here, we present the results of active chlorophyll fluorometry of 30 species of modern planktonic foraminifera obtained from 1266 individuals. The main purpose of this study is (1) to provide information on the biomass of symbionts (indicated by chlorophyll *a* content), (2) to qualify the functionality/fitness of symbionts (indicated by photophysiology), (3) to characterize the photosymbiotic features, and (4) to propose a new framework to characterize the photosynthetic activity of modern planktonic foraminifera.

## 2 Material and Methods

### 2.1 Sampling and identification of morphological species

Planktonic foraminifera were collected in central and western Pacific Ocean and north-eastern Atlantic Ocean (Fig. 1). We have sampled across much of the northern hemisphere tropical-subtropical gradient in both Pacific and Atlantic, to get the endemic species and to replicate for the others. Samples from the Pacific Ocean were taken onboard during RV Mirai cruises



MR13-04 and MR14-02, RV Kaiyo cruise KY14-09, RV Shinsei-maru cruise KS-16-9, and RV Hakuho-maru cruises KH-16-7 and KH-17-4 (Fig. 1a). The samples were collected either by vertical stratified towing (closing ring net or VMPS with 100- $\mu$ m mesh) or by pumped seawater (sampling depth, ca. 5 m). The pumped seawater was continuously opened to a 100- $\mu$ m-mesh net settled within a water tank to collect specimens as gently as possible. Some specimens were additionally collected from Tsugaru Strait, Sagami Bay, and off Sesoko Island by surface towing and vertical towing with a 100- $\mu$ m-mesh net to increase the taxonomic range of our analysis. Samples from the Atlantic were taken onboard during RV Meteor cruise M140 (Fig. 1b). A multi-closing net system (Multi-Plankton-Sampler) with 100- $\mu$ m mesh was used for stratified sampling of the water column. Samples from pumped seawater (sampling depth, ca. 8 m) were also collected in the same way as to the Pacific sampling.

Collected specimens were isolated immediately after collection with either brush or Pasteur pipets into Petri-dishes filled with 0.22- $\mu$ m-filtered or 0.45- $\mu$ m-filtered seawater, and rinsed several times. Specimens were identified to morphospecies level under a stereoscopic microscope, and the maximum test length (test size) were measured. We identified 30 morphospecies from four families (Globigerinidae, Hastigerinidae, Globorotaliidae, and Candeinidae) (Fig. 2). *Sphaeroidinella dehiscens* was identified only after it thickened its test during the adult stage under culture, though the data we used here is from the very first measurement after collection before the identification. We differentiated *G. ruber* white variety and pink variety based on the pigmentation in earlier whorls of the tests. *Globigerinella siphonifera* was divided into two morphotypes (Type I and Type II) based on the criteria described in Faber et al. (1988) and Huber et al. (1997). From among the isolated individuals, viable specimens were selected for analysis with the following criteria; (1) penultimate chamber was filled with cytoplasm and (2) the specimen was sticky when touched with a brush or the rhizopods were observed under a microscope. Screening for the presence of photosymbiosis was conducted on as many species and specimens as possible, regardless of locality and sampling depth. Photophysiological measurements were carried out only on specimens collected from the upper 100 m of the water column (corresponding approximately to the photic zone). The specimens were kept individually in a well of a culture dish filled with filtered seawater until the measurement. The duration between the collection and the measurement was no longer than 12 hrs. During this time, most spinose species recovered their spines.

## 2.2 Fast repetition rate fluorometry measurements and photophysiological parameters

Fast repetition rate (FRR) fluorometry, a kind of active fluorometry, can obtain photophysiological information of host-algal symbiotic consortia using various parameters of photosystem II (PSII) (Fig. 3). FRR fluorescence transients were measured either using an FRR fluorometer DF-03 or DF-14 (Kimoto Electric Co., Ltd.) (Table S1). FRR fluorometers generate a series of blue flashlets of an excitation light intensity of 30 mmol quanta  $\text{m}^{-2} \text{s}^{-1}$  with a wavelength of 470 nm with a 25 nm-bandwidth (DF-03) or a wavelength of 450 nm with a 10 nm-bandwidth (DF-14). Saturation protocols were consisting of 50 flashlets of 2  $\mu$ s duration at 4  $\mu$ s intervals (DF-03) or 100 flashlets of 1  $\mu$ s duration at 2  $\mu$ s intervals (DF-14). A fluorescence induction curve based on the biophysical model of Kolber et al. (1998) was numerically fitted to transients of chlorophyll fluorescence to derive PSII parameters. The parameters include minimum fluorescence ( $F_0$ ), maximum fluorescence ( $F_m$ ), variable



fluorescence [ $F_v (=F_m - F_0)$ ], maximum photochemical efficiency indicating photosynthetic activity ( $F_v/F_m$ ), and functional absorption cross-section of PSII indicating light absorption efficiency ( $\sigma_{PSII}$ ) (Fig. 3). Before measurements, specimens were confirmed that no visible contamination of algae or particles were present at the test surface and spines under a stereoscopic microscope. After 10-minutes dark adaptation, a specimen was transferred into a quartz glass cuvette with filtered seawater  
135 for the measurement.

### 2.3 Assessment of symbiont possession and parameters characterizing photosymbiosis

When chlorophyll fluorescence ( $F$ ) was detected from an individual foraminifera, the status of chlorophyll was categorized based on the detection of variable fluorescence ( $F_v$ ).  $F_v$  represents fluorescence transients during the saturation process of the reaction centers of PSII. It is detected only when the PSII captures photons and passes the product further through the chain of photosynthetic reactions; i.e., when actively photosynthesizing organisms are present in the specimen. When  $F_v$  was not  
140 not detected but  $F$  value was significantly higher than the background level of the fluorometer, chlorophyll was regarded to be present but non-functional, signifying remnants of phytoplankton prey. If no  $F$  was detected, the specimen had no chlorophyll (Figs. 3 and 4).

When functional chlorophyll was detected in a specimen, then the maximum fluorescence ( $F_m$ ) value was used to estimate  
145 chlorophyll  $a$  (Chl  $a$ ) content of the specimen based on a linear relationship between  $F_m$  and Chl  $a$  (cf. Fujiki et al., 2014; Takagi et al., 2016). Calibration line was established for each FRR fluorometer. A relationship between the Chl  $a$  content, an indicator of symbiont biomass, and the foraminiferal test size was then analyzed. As an indicator of symbiont density of an individual, Chl  $a$  content per protein biomass (Chl  $a$ /biomass) was also calculated. The protein biomass was estimated based on species-specific relationships with test size (exponential equation) proposed by Movellan (2013). For species whose test  
150 size-biomass relationship was not presented in his study, the protein biomass was estimated based on the relationship established by morphologically similar species (Table S2). As indicators of photosynthetic vitality and light absorption efficiency of symbionts, photophysiological parameters  $F_v/F_m$  and  $\sigma_{PSII}$  were used, respectively.

### 2.4 Statistical analysis

To compare the differences in the parameters (Chl  $a$ /biomass,  $F_v/F_m$ , and  $\sigma_{PSII}$ ) among species, statistical tests for comparison  
155 of differences in medians (Kruskal-Wallis test and post-hoc Steel-Dwass test for multiple comparison) were conducted. Species with less than 20 specimens were not tested due to small sample size. Principal component analysis (PCA) was performed to characterize photosymbiotic features of the studied species, based on the four diagnostic variables of photosymbiosis obtained in this study; (1) ratio of symbiont-bearing individuals, (2) correlation coefficient between test size and Chl  $a$  content, (3) Chl  $a$  content relative to the protein biomass (Chl  $a$ /biomass), and (4)  $F_v/F_m$  value. Species medians were used for the variables  
160 Chl  $a$ /biomass and  $F_v/F_m$  as representative values. In terms of the correlation coefficient of test size-Chl  $a$  relationship, negative values were considered as zero. K-means clustering was also performed to categorize photosymbiosis and to visualize the results of the PCA. All the statistical analyses were performed using R (R version 3.3.1, R Core Team, 2016).



### 3 Results

#### 3.1 Possession of symbionts

165 The results of the measurements on all 1266 specimens are shown in Table S1, including sampling locality, date, and the measured parameters. The incidence of each type of chlorophyll (functional, non-functional, and no chlorophyll) is summarized in Figure 5. Chlorophyll fluorescence, either functional or non-functional, was detected in 27 out of 30 species. The species *G. adamsi*, *N. incompta*, and *N. pachyderma* never showed any evidence for the presence of chlorophyll. Specimens of *G. scitula*, *G. crassaformis*, *G. truncatulinoides*, *H. pelagica*, *H. digitata*, *G. bulloides*, *T. quinqueloba*, and *T.*  
170 *fleisheri* never possessed functional chlorophyll, although many of them contained non-functional chlorophyll. Nineteen species contained functional chlorophyll, and can be considered symbiont-bearing: *O. universa*, *S. dehiscens*, *G. sacculifer*, *G. conglobatus*, *G. ruber* (white), *G. ruber* (pink), *G. rubescens*, *G. tenella*, *G. calida*, *G. siphonifera* Type I, *G. siphonifera* Type II, *T. humilis*, *P. obliquiloculata*, *N. dutertrei*, *G. inflata*, *G. menardii*, *C. nitida*, *G. glutinata* and *G. uvula*. Among these species, the percentage of symbiont-bearing individuals varied from 100 % (*S. dehiscens* and *G. conglobatus*) to 58 % (*G.*  
175 *calida*). Although the examined specimens included individuals collected at all depths, the percentages of non-functional or no-chlorophyll individuals were similar when removing the specimens collected below 100 m (Fig. S1). The incidence of symbiotic individuals was not significantly different between Pacific and Atlantic ( $p \gg 0.05$ , Fisher's exact test for species with more than 15 individuals in each basin, see Fig. S2). Moreover, the ontogenetic (size) trend in possession of symbionts was not apparent (Fig. 6).

180 *Globoturborotalita rubescens* and *G. tenella* have never been reported to possess symbionts, but we observed ovoid reddish brown symbionts along with their spines just as they are usually seen in *O. universa*, *G. ruber*, *G. conglobatus*, and *G. sacculifer* (Fig. S3). Therefore, although genetic information and detailed microscopic evidence are needed in the future, we categorize them here as dinoflagellate-bearing species. The remaining symbiont-bearing species that have never been reported before were *G. calida* and *G. uvula*. Symbionts of these species are treated here as uncharacterized. As precaution, the  
185 convincing symbiont-bearing species whose symbionts have not yet been identified by DNA analysis are treated as uncharacterized as well, *T. humilis*, *P. obliquiloculata*, *G. inflata*, *G. menardii*, *C. nitida*, and *G. glutinata* (Table 1).

#### 3.2 Test size–Chl *a* content relationship, and Chl *a*/protein biomass

Out of the 19 species which had functional chlorophyll (symbiont-bearing species), 16 species showed a statistically significant positive correlation between test size and Chl *a* content ( $p < 0.05$ , Fig. 6), with Chl *a* content being a power function of test  
190 size. The powers (scaling exponents) of the fitted functions varied from 1.33 (*G. tenella*) to 3.71 (*G. calida*) (Table 2). For the remaining three species, *T. humilis*, *P. obliquiloculata*, and *G. inflata*, their test size–Chl *a* relationships showed no significant correlation (Fig. 6).

The ratio of Chl *a* to protein biomass per individual showed clear differences among species (Fig. 7). *Globigerinoides conglobatus*, *G. sacculifer*, and *O. universa* showed significantly higher Chl *a*/biomass values (species median values were



195 4.8, 4.8, and 4.6 ng  $\mu\text{g}^{-1}$ , respectively), and *P. obliquiloculata* showed the lowest (median value 0.1 ng  $\mu\text{g}^{-1}$ ). Spinose species tended to show higher Chl *a*/biomass values than non-spinose species.

### 3.3 Photophysiological state

Overall,  $F_v/F_m$  values tended to be high in dinoflagellate-bearing species (species median values 0.46–0.53) (Fig. 8a). Amongst all 19 symbiont-bearing species,  $F_v/F_m$  value was highest in *S. dehiscens* (0.53), and lowest in *G. inflata* (0.33).  
200 Species to species comparison showed that *P. obliquiloculata* alone showed significantly lower  $F_v/F_m$  values ( $p \ll 0.01$ ).

On the other hand,  $\sigma_{\text{PSII}}$  was clearly low in dinoflagellate-bearing species (median values  $374\text{--}606 \times 10^{-20} \text{ m}^2 \text{ quanta}^{-1}$ ) and high in pelagophyte-bearing species (median values  $618\text{--}749 \times 10^{-20} \text{ m}^2 \text{ quanta}^{-1}$ ) (Fig. 8b). The highest and lowest  $\sigma_{\text{PSII}}$  in median were recorded in *N. dutertrei* ( $749 \times 10^{-20} \text{ m}^2 \text{ quanta}^{-1}$ ) and *C. nitida* ( $347 \times 10^{-20} \text{ m}^2 \text{ quanta}^{-1}$ ), respectively. Based on the statistical testing of the species to species difference in medians, *N. dutertrei* and *G. siphonifera* Type II (pelagophyte-  
205 bearing) showed no difference ( $p = 0.79$ ), and associated with the highest  $\sigma_{\text{PSII}}$ . *Globigerinoides ruber* (pink) alone showed significantly lower  $\sigma_{\text{PSII}}$  than the other dinoflagellate-bearing species ( $p \ll 0.01$ ), and the value was comparable to that of *C. nitida* ( $p = 1.0$ ).

### 3.4 Principal component analysis and clustering

The first principal component (PC1) alone accounted for 84.2 % of the total variance, and the second principal component (PC2) for 10.2 % (Fig. 9). In the PC1 score, the loading coefficient was positive for all variables related to photosymbiosis used in the analysis (0.96 for the ratio of symbiont-bearing individuals, 0.91 for the positive correlation coefficient of test size-Chl *a* content relationship, 0.96 for the  $F_v/F_m$  median value, and 0.82 for the Chl *a* content relative to protein biomass). Considering the high contribution to explaining the total variance, and the positive loading for the four variables, the PC1 score well represented photosymbiotic characteristics among foraminiferal species. In fact, the cluster analysis revealed that four  
215 clusters of species were separated along the PC1 score. The lowest PC1 score (−2.2) was recorded by non-symbiotic species (Cluster 4). The distribution of species along the PC1 score was relatively wide for the Cluster 2 and 3 (0.7–2.2, −0.6–0.2, respectively), whereas almost the same for the Cluster 1 with the highest score (2.3–2.5). The Cluster 1 and 2 consisted of the species with significant positive correlations between test size and Chl *a* content. The Cluster 1 was separated from Cluster 2 primarily due to their distinctly high PC2 score. The PC2 was characterized by Chl *a* content per protein biomass (Chl  
220 *a*/biomass) which exclusively had positive loading (0.57). Three species in the Cluster 1, *G. conglobatus*, *G. sacculifer*, and *O. universa*, were revealed to have significantly high Chl *a*/biomass as represented in Figure 7. The Cluster 2 consisted of 13 species which showed the widest distribution along with both PC1 and PC2 axes. Among the Cluster 2, the non-spinose species tended to show lower PC1 and PC2 scores compared to the spinose species. The Cluster 3 consisted of three species, *T. humilis*, *P. obliquiloculata*, and *G. inflata*. They were the species that possessed symbionts in most cases but without significant positive correlation in the test size-Chl *a* relationship. Overall, the clusters and the PC1 score depicted a clear tendency of  
225 photosymbiosis related features of the species.



## 4 Discussion

### 4.1 Characteristics and a new framework of planktonic foraminiferal photosymbiosis

230 Based on the result of the PCA and cluster analyses, 30 foraminiferal species were characterized and categorized into four groups (Fig. 9). The Cluster 4 is a group of non-symbiotic species. Of the 11 species in this group, six species were tested on their photosymbiosis for the first time, and revealed to be non-symbiotic: *G. adamsi*, *T. quinqueloba*, *H. digitata*, *G. scitula*, *G. crassaformis*, and *T. fleischeri*. An interesting feature of this group was that many species possessed non-functional chlorophyll (Fig. 5). For example, all the specimens in *G. scitula* and *G. crassaformis* had a certain amount of chlorophyll  
235 inside, but it was always non-functional and likely derived from prey. The occurrence of fresh (fluorescent) chlorophyll in these species was surprising, considering that most of these specimens were collected from a water depth below 300m (Table S1) where the chlorophyll concentration is low. They might incorporate sinking aggregates of phytoplankton remains as food (e.g., Anderson et al., 1979; Spindler et al., 1984), and chlorophyll or chloroplast itself might have remained undigested, resulting in non-functionality of chlorophyll. *Hastigerina pelagica* are known to show vertical depth segregation among the  
240 genotype (Weiner et al., 2012). It has been speculated that such segregation might be related to their possession of symbionts (e.g., Huber et al., 1997; Seears et al., 2012). Though our study did not identify their genotype, we revealed that this species never possessed symbionts even they were collected from shallower water depth (< 100m). The recent study showed that *G. bulloides* type IId possessed cyanobacterial symbionts (Bird et al., 2017). By using our fluorescence technique, chlorophyll fluorescence of cyanobacteria should also be detectable although the most effective wavelength of the fluorescence is slightly  
245 different. In fact, two specimens of this species showed possession of chlorophyll, yet they were non-functional (Table S1). This might indicate that possession of cyanobacterial symbionts may be a genotype-dependent, or regional or seasonal specific phenomenon.

On the other hand, five species were newly confirmed as symbiotic in this study; *S. dehiscens*, *G. rubescens*, *G. tenella*, *G. calida*, and *G. uvula*. All species in the Cluster 1 and 2 including the above five species showed relatively high rates of  
250 possession of symbionts, and exclusively showed significant positive correlations between the test size and Chl *a* content (Figs. 6 and 10). It was previously revealed that *G. sacculifer* and *G. siphonifera* Type II showed positive correlations between test size and Chl *a* content (Takagi et al., 2016). Similarly, *O. universa* has been demonstrated to have a positive relationship between test size and symbiont number itself (Spero and Parker, 1985). The capability of cell divisions of symbionts cannot be directly clarified from our active fluorescence-based study, but the significant positive correlation can be a strong indication  
255 for the growth of the symbiont population inside the host foraminifera. Hence, in addition to the high percentage of symbiont-bearing individuals in a species, such strong positive correlation should be a specific diagnostic of a persistent relationship of photosymbiosis through their lifetime. Moreover, species in the Cluster 1 should have the potential to perform more effective



photosynthesis due to the higher content of Chl *a* per protein biomass (Fig. 7). It may imply more phototrophic nature of these species.

260 The Cluster 1 and 2 included well-studied symbiotic species like *O. universa*, *G. ruber*, *G. sacculifer*, and *G. siphonifera* previously reported to be in “obligate” symbiosis (Hemleben et al., 1989). Amongst “facultative” symbiotic species inferred in previous studies, *N. dutertrei*, *G. menardii*, *C. nitida* and *G. glutinata* were revealed to have the persistent symbiotic relationships based on our test size-Chl *a* correlation analysis. In this study, not only so-far called “facultative” symbiotic species, but also most of the species were sometimes found without symbionts (Fig. 5). It was repeatedly observed that *G.*  
265 *sacculifer* and *G. siphonifera* digest their symbionts prior to gametogenesis (e.g., Bé et al., 1983; Faber et al., 1988; Takagi et al., 2016). Thus, symbiont-barren individuals could be present in the adult stage. However, the size of such symbiont-barren specimens recognized in this study was not necessarily large (Fig. 6). We speculate that they were in an unhealthy condition and going to die. In any case, the percentage of symbiont-barren individuals was small. We think the presence of symbiont-barren specimens in symbiont-bearing species, unless it is dominant, is not critical to describe the nature of photosymbiosis (i.e., conventional categorization of obligate or facultative symbiosis). Rather, the presence of such symbiont-barren  
270 individuals in these groups has led to the confusion in earlier works who placed some of these species into the category “facultative”. Nevertheless, the ratio of symbiont-bearing individuals may overall reflect the ecological differences among species like the persistence of symbiosis or the dependence on phototrophy that can represent a “strength” of photosymbiosis.

The Cluster 3 had intermediate features between persistent symbiosis (Cluster 1 and 2) and non-symbiosis (Cluster 4).  
275 Species do possess symbionts and can be called symbiotic species, but the significant correlation in test size-Chl *a* relationship which was common in the Cluster 1 and 2 was absent (Figs. 5 and 6). Contrary to the interpretation of the significant positive correlation in test size-Chl *a*, the absence of such a relationship indicates that the algae in the examined species were not able to increase their biomass as the host grew. It can be said the algae could not persistently reside in their host, in other words, the symbiosis is transient. *Pulleniatina obliquiloculata* and *G. inflata* are non-spinose species whose eating habits are reported  
280 to be primarily herbivorous (Anderson et al., 1979; Spindler et al., 1984). Therefore, the lack of the relationship with test size and Chl *a* content may reflect the residence time of algal prey. We speculate that all individuals that are capable to host symbionts can be temporarily symbiotic as long as the foraminifera incorporates a certain algal species. When a foraminifera maintains the algae for some time keeping them undigested and keeping their photosynthetic capability to provide photosynthates, the algae can serve as symbionts. Regardless of the role of the algae, i.e., symbionts or preys, when the algae  
285 are all digested, the foraminifera becomes temporarily chlorophyll-barren. If these are the case, possession of symbionts should be temporal, resulting in no size correlation in the Chl *a* content. If the symbionts do not increase inside the host, the Chl *a* content of a specimen is regulated by the incorporation frequency/rate of algal cells and their residence time inside the host (i.e., a balance between incorporation and digestion). They could potentially serve as symbionts if the residence time is long. This is similar to what is known of the behavior of benthic species with kleptoplasts; these are actively harvested and are  
290 functional, but wear off with time and have to be replenished (e.g., Bernhard and Bowser, 1999; Pillet et al., 2011). Digestion experiments for these species, therefore, an interesting subject in the future, and the FRR fluorometry can also serve such



culturing studies. Together with *P. obliquiloculata* and *G. inflata*, *T. humilis* which was previously inferred as “obligate” symbiotic species (Hemleben et al., 1989) fall into the Cluster 3 representing such transient symbiosis. However, caution should be paid for the narrow size range of *T. humilis* (97–168  $\mu\text{m}$ ) (Fig. 6). In addition, the specimens we analyzed were mostly with 13–15 chambers, probably in their adult stage. In this respect, since a sufficient size range of specimens with a variety of ontogenetic stages were not covered, it is difficult to strongly conclude that symbiosis in *T. humilis* is not persistent. Considering the  $F_v/F_m$  value of this species (0.51 in median), it was clearly higher compared to the other two species in the Cluster 3, *P. obliquiloculata* and *G. inflata* (0.36 and 0.33, respectively). Besides, the possession of symbionts of this species was 89 %, and higher than the other two as well (66 % for *P. obliquiloculata* and 69 % for *G. inflata*). We therefore interpret that *T. humilis* has established more persistent symbiosis compared to *P. obliquiloculata* and *G. inflata*.

Here, considering the above characterization of photosymbiosis, we propose a new framework of planktonic foraminiferal photosymbiosis (Fig. 11). As suggested in Stoecker et al. (2009), we think photosymbiosis can be regarded as a spectrum from absolute non-symbiosis (heterotrophy) to more robust symbiosis (higher extent of acquired phototrophy) which ends with a permanent plastid endosymbiosis seen in utter autotrophs. Each foraminiferal species can be located somewhere in-between phototrophy and heterotrophy. Since the PC1 score well represents the photosymbiotic characteristics, it is suitable as a quantitative indicator of the level of photosymbiosis in species which harbor active chlorophyll. Therefore, we aligned the species along with the PC1 score scale in the conceptual diagram (Fig. 11). An important point here is that this spectrum allowed us to gain an overview of the relative strength of photosymbiosis among species and across various families of planktonic foraminifera. The relative ordination may be amended by further exploration in the future, but we believe our thorough investigation could shed light on the species-specific difference in the nature of photosymbiosis in planktonic foraminifera. This would be a solid basis to help us to think about evolutionary aspects of photosymbiosis, its role in the earth system history, and possible effects on test geochemistry.

#### 4.2 Size scaling of Chl *a* content in symbiotic foraminifera

The significant positive correlation between test size and Chl *a* content (Figs. 6 and 10) is not just showing the growing nature of symbionts, but also has information on a kind of quantitative relationship in the host and symbionts based on their scaling exponent (Table 2). In theory, the size scaling exponent of 3 means that the dependent variable increases proportionally to the volume development. If the test shape is flatter, as is the case of *G. menardii*, the exponent should be smaller and approaching 2. Alternatively, when the test volume does not reflect the cytoplasm volume (the growth rate of cytoplasm is less than that of the test volume) like adult spherical specimens of *O. universa*, the scaling exponent results in relatively small values. The fact that all 16 species showed scaling exponent in the range of 2 to 3 (95 % confidence intervals overlap with this range, Table 2) indicates that the Chl *a* content, indirectly reflecting the symbiont biomass, increased in nearly proportional to the host’s test volume increased. This kind of size scaling theory across different species of planktonic foraminifera suggests a robust relationship between the host and symbionts.



The other notable point in the test size-Chl *a* relationship is that the spinose species commonly had higher contents of Chl  
325 *a* compared to the non-spinose species (Fig. 10). For example, when the test size was ca. 300  $\mu\text{m}$ , the macroperforate spinose  
group had almost five times higher Chl *a* than the microperforate non-spinose group, and 10 times higher than the  
macroperforate non-spinose group. The light-dark rhythm of symbiont deployment along the spines was commonly observed  
in *Globigerinoides*, *Orbulina* and *Globigerinella* species (Anderson and Bé, 1976; Bé et al., 1977; Hemleben and Spindler,  
1983; Takagi et al., 2016). Considering this phenomenon, the presence of spines could be seen as a character that facilitates  
330 symbiosis or at least allows the harboring of a larger symbiont population. For example, efficient lighting for each symbiont  
cell and maximizing total photosynthetic rates can be achieved due to the spherical distribution of symbionts along the radiating  
spines. This may contribute to the higher Chl *a* content in spinose species.

Moreover, the distribution of the plots in Figure 10 showed three clear clusters corresponding to each morphogroups,  
macroperforate spinose, macroperforate non-spinose, and microperforate non-spinose. It is also an interesting feature firstly  
335 revealed in this study. The largest morphological difference among these three groups is the test wall architecture including  
the density of pores, the presence of spines, and smoothness of test surface. Hence, these features might be related to the light  
penetration through the test wall. When the symbionts are sequestered within the test as seen in the non-spinose species, the  
test wall architecture is expected to affect the efficiency of photosynthesis. It may also be involved with lower Chl *a*/biomass  
observed in non-spinose species. If such microenvironmental condition surrounding the intracellular symbionts are measurable  
340 or theoretically modeled, it will enhance our understanding of the differences and the controlling factor of symbiont density.

### 4.3 Photophysiology and host-symbiont partnerships

When species are grouped based on the symbiont type they possess, dinoflagellate (*O. universa*, *G. sacculifer*, *G. conglobatus*,  
*S. dehiscens*, *G. ruber*, *G. tenella*, and *G. rubescens*). or pelagophyte (*G. siphonifera* Type II and *N. dutertrei*) (Table 1),  
photophysiological parameters were all significantly different between these groups. Chl *a*/biomass and  $F_v/F_m$  values were  
345 higher for dinoflagellate-bearing species ( $p \ll 0.01$  and  $p = 0.012$ , respectively, Figs. 7 and 8a), and  $\sigma_{\text{PSII}}$  values were higher  
for pelagophyte-bearing species ( $p \ll 0.01$ , Fig. 8b). As far as the species whose symbionts are known are compared, it seems  
that the symbiont photophysiology was overall related to the type of symbiont rather than the host size or the host  
morphological groups. In fact, we previously published experimental results on photophysiology of cultured *G. sacculifer*  
(dinoflagellate-bearing) and *G. siphonifera* Type II (pelagophyte-bearing), and reported lower  $F_v/F_m$  and higher  $\sigma_{\text{PSII}}$  in *G.*  
350 *siphonifera* Type II than in *G. sacculifer* (Takagi et al., 2016). In this study, what we observed was the same tendency of  
photophysiology corresponding to the type of symbionts they have, regardless of the phylogenetic position the host belongs  
to. The higher  $\sigma_{\text{PSII}}$  indicates a higher acclimation potential to a low-light environment. This observation is consistent with the  
living depth of the involved species. In general, dinoflagellate-bearing species like *G. ruber* and *G. sacculifer* prefer shallower  
habitat, and pelagophyte-bearing species like *N. dutertrei* and *G. siphonifera* Type II prefer relatively deeper water (Rebotim  
355 et al., 2017). Moreover, when *G. siphonifera* Type I and Type II were compared, the Type I having haptophyte symbionts  
showed significantly lower  $\sigma_{\text{PSII}}$  than the Type II (Fig. 8b). The previous report on the difference in pigment content of these



types also implied deeper habitat in *G. siphonifera* Type II (Bijma et al., 1998). The  $\sigma_{PSII}$  difference revealed in this study supports their arguments. Moreover, even in the time before the type difference of this species was recognized, *G. siphonifera* was often reported to have a bimodal vertical distribution (Tolderlund and Bé, 1971). It possibly reflected the difference of the light preference of their associating symbionts. The current knowledge on  $\sigma_{PSII}$  in foraminifera is still limited, but the observed consistency to their known depth preferences indicates that the symbiont acclimation potential may be one of the factors controlling the habitat selection of the host species.

The dinoflagellate-bearing species, *G. ruber* (pink) showed high  $F_v/F_m$  with relatively small variation, and interestingly, it was significantly higher than that of *G. ruber* (white) (Fig. 8a). In general,  $F_v/F_m$  values vary depending on the nutrient availability (Kolber et al., 1988; Parkhill et al., 2001); i.e., the higher  $F_v/F_m$  may be achieved by the higher nutrient supply to the symbionts. A recent study showed that the inorganic nutrients in the ambient seawater do not affect the  $F_v/F_m$  of *G. sacculifer*, suggesting that it is the internal supply of nutrients from the host to symbionts that can influence on the  $F_v/F_m$  (Takagi et al., 2018). In this context, it can be assumed that among the species having the same symbionts, the higher  $F_v/F_m$  possibly reflects the higher level of host-symbiont interaction. If it is the case, among the species used for the statistical analysis, it can be said that the strongest symbiotic relationship has been established in *G. ruber* (pink). In fact, however, the interspecific comparison may not be suitable because the other environmental factors which might affect the physiology of the host-symbiont consortia, such as seawater temperature, salinity, light intensity, and prey abundance, were not considered in this study. *Globigerinoides ruber* (pink) was collected only from the Atlantic cruise, whereas *G. ruber* (white) was collected from various oceanic realms (Table S1). It may also be involved with relatively constrained  $F_v/F_m$  values in *G. ruber* (pink) and contrastingly large variability in *G. ruber* (white). In order to discuss more detail on interspecific photophysiological differences, comparison of the photophysiological parameters for specimens cultured under controlled condition, or the compilation of individual data collected from a similar environmental condition is needed.

## 5 Conclusion and future perspectives

The present study was aimed to extend our understanding of photosymbiosis in modern planktonic foraminifera. A thorough investigation of 30 foraminiferal species was performed using FRR fluorometry. Eleven species showed no signal of photosynthesis, and were confirmed to be non-symbiotic. Nineteen species, in contrast, showed the functionality of photosynthesis which is convincing evidence of photosymbiosis. Of these species, we found significant positive correlations in test size-Chl *a* content relationship in 16 species, which were regarded to show persistent symbiotic relationships. Especially, dinoflagellate-bearing *G. sacculifer*, *G. globatus*, and *O. universa* had higher Chl *a* density, probably reflecting the higher potential of photosynthesis. The rest of three species, *T. humilis*, *P. obliquiloculata*, and *G. inflata* showed no significant size scaling relationship in Chl *a* content. Moreover, their  $F_v/F_m$  values and the symbiont possession rates were comparatively low. Based on a PCA using the four features relating to photosymbiosis, we ranked 30 species along with an integrated scale (the PC1 score scale). Finally, we proposed a new framework of photosymbiosis in planktonic foraminifera as a continuous spectrum implying a strength of photosymbiosis. In the context of nutrition, this concept represents a varying degree of



390 mixotrophy which is commonly seen in marine planktonic organisms (Stoecker et al., 2017). Besides, an interesting finding in  
this study was that the photophysiology was basically determined by the type of the symbiont they have, regardless of the  
phylogenetic position of the host or their test morphology. The physiological parameter, especially  $\sigma_{PSII}$ , seemed to correspond  
to the overall depth habitat of the host foraminifera. It might imply that the habitat of the host foraminifera is partly governed  
by the symbiont type. However, what is missing in our study is the taxonomy of the symbionts. Combining the information of  
395 FRR fluorometry, DNA, as well as microscopic evidence on their ultrastructure will provide a more comprehensive  
understanding of photosymbiosis in planktonic foraminifera.

*Data availability.* All data obtained in this study is in Supplementary materials.

400 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We acknowledge the Captains and the shipboard members of RV Mirai, RV Kaiyo, RV Shinsei-maru, RV  
Hakuho-maru, and RV Meteor. We would like to thank the staff at the Misaki Marine Biological Station (The University of  
Tokyo), the staff at the Sesoko Station, Tropical Biosphere Research Center (University of the Ryukyus), Tomohiko Kikuchi  
405 and Shinji Shimode (Yokohama National University) for helping the sampling. The graphs were drawn with Generic Mapping  
Tools (Wessel and Smith, 1998). This study was financially supported by JSPS KAKENHI Grant No. 24121003 (to H. Saito),  
25740014 and 16K00532 (to T. Fujiki), 13J05477, 16H06738, and 17J05887 (to H. Takagi).

## References

- Anderson, O. R., and Bé, A. W. H.: The ultrastructure of a planktonic foraminifer, *Globigerinoides sacculifer* (Brady), and its  
410 symbiotic dinoflagellates, *Journal of Foraminiferal Research*, 6, 1–21, 1976.
- Anderson, O. R., Spindler, M., Bé, A. W. H., and Hemleben, C.: Trophic activity of planktonic foraminifera, *Journal of Marine  
Biological Association of the UK*, 59, 791–799, 1979.
- Bé, A. W. H., Hemleben, C., Anderson, O. R., Spindler, M., Hacunda, J., and Tuntivate-Choy, S.: Laboratory and field  
observations of living planktonic foraminifera, *Micropaleontology*, 23, 155–179, 1977.
- 415 Bé, A. W. H., Anderson, O. R., Faber, W. W. Jr., and Caron, D. A.: Sequence of morphological and cytoplasmic changes  
during gametogenesis in the planktonic foraminifer *Globigerinoides sacculifer* (Brady), *Micropaleontology*, 29, 310–  
325, 1983.
- Bernhard, J. M., and Bowser, S. S.: Benthic foraminifera of dysoxic sediments: chloroplast sequestration and functional  
morphology, *Earth-Science Reviews*, 46, 149–165, 1999.



- 420 Bijma, J., Hemleben, C., Huber, B. T., Erlenkeuser, H., and Kroon, D.: Experimental determination of the ontogenetic stable isotope variability in two morphotypes of *Globigerinella siphonifera* (d'Orbigny), *Marine Micropaleontology*, 35, 141–160, 1998.
- Bird, C., Darling, K. F., Russell, A. D., Davis, C. V., Fehrenbacher, J., Free, A., Wyman, M., and Ngwenya, B. T.: Cyanobacterial endobionts within a major marine planktonic calcifier (*Globigerina bulloides*, Foraminifera) revealed  
425 by 16S rRNA metabarcoding, *Biogeosciences*, 14, 901–920, 2017.
- Bird, C., Darling, K. F., Russell, A. D., Fehrenbacher, J., Davis, C. V., Free, A., and Ngwenya, B. T.: 16S rRNA gene metabarcoding and TEM reveals different ecological strategies within the genus *Neogloboquadrina* (planktonic foraminifer), *Plos ONE*, 13, e0191653, 2018.
- Bolli, H. M., Saunders, J. B., and Perch-Nielsen, K.: *Plankton Stratigraphy*, volume 1, Cambridge University Press, Cambridge,  
430 1985.
- Boudagher-Fadel, M. K., Banner, F. T., and Whittaker, J. E.: *The Early Evolutionary History of the Planktonic Foraminifera*, Springer, Netherlands, 1997.
- Ezard, T. H. G., Aze, T., Pearson, P. N., and Purvis, A.: Interplay between changing climate and species' ecology drives macroevolutionary dynamics, *Science*, 332, 349–351, 2011.
- 435 Ezard, T., Edgar, K. M., and Hull, P. M.: Environmental and biological controls on size-specific  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  in recent planktonic foraminifera, *Paleoceanography*, 30, 151–173, 2015.
- Faber, W. W. Jr., Anderson, O. R., Lindsey, J. L., and Caron, D. A.: Algal–foraminiferal symbiosis in the planktonic foraminifer *Globigerinella aequilateralis*. I. Occurrence and stability of two mutually exclusive chrysophyte endosymbionts and their ultrastructure, *Journal of Foraminiferal Research*, 18, 334–343, 1988.
- 440 Faber, W. W. Jr., Anderson, O. R., and Caron, D. A.: Algal–foraminiferal symbiosis in the planktonic foraminifer *Globigerinella aequilateralis*. II. Effects of two symbiont species on foraminiferal growth and longevity, *Journal of Foraminiferal Research*, 19, 185–193, 1989.
- Fenton, I. S., Pearson P. N., Dunkley Jones T., and Purvis, A.: Environmental predictors of diversity in recent planktonic foraminifera as recorded in marine sediments, *PLoS ONE*, 11, e0165522, 2016.
- 445 Fujiki, T., Takagi, H., Kimoto, K., Kurasawa, A., Yuasa T., and Mino, Y.: Assessment of algal photosynthesis in planktic foraminifers by fast repetition rate fluorometry, *Journal of Plankton Research*, 36, 1403–1407, 2014.
- Gast, R. J., and Caron, D. A.: Molecular phylogeny of symbiotic dinoflagellates from Foraminifera and Radiolaria, *Molecular Biology and Evolution*, 13, 1192–1197, 1996.
- Gast, R. J., McDonnell, T. A., and Caron, D. A.: srDNA-based taxonomic affinities of algal symbionts from a planktonic  
450 foraminifera and a solitary radiolarian, *Journal of Phycology*, 36, 172–177, 2000.
- Gastrich, M. D.: Ultrastructure of a new intracellular symbiotic alga found within planktonic foraminifera, *Journal of Phycology*, 23, 623–632, 1987.



- Gastrich, M. D., and Bartha, R.: Primary productivity in the planktonic foraminifer *Globigerinoides ruber* (d'Orbigny), *Journal of Foraminiferal Research*, 18, 137–142, 1988.
- 455 Hallock, P.: Algal symbiosis: a mathematical analysis, *Marine Biology*, 62, 249–255, 1981.
- Hemleben, C., and Spindler, M.: Recent advances in research on living planktonic foraminifera, *Utrecht Micropaleontological Bulletins*, 30, 141–170, 1983.
- Hemleben, C., Spindler, M., and Anderson, O. R.: *Modern planktonic foraminifera*, Springer-Verlag, New York, 1989.
- 460 Henehan, M. J., Rae, J. W. B., Foster, G. L., Erez, J., Prentice, K. C., Kucera, M., Bostock, H. C., Martínez-Botí, M. A., Milton, J. A., Wilson, P. A., Marshall, B. J., and Elliott, T.: Calibration of the boron isotope proxy in the planktonic foraminifera *Globigerinoides ruber* for use in palaeo-CO<sub>2</sub> reconstruction, *Earth and Planetary Science Letters*, 364, 111–122, 2013.
- Hönisch, B., Bijma, J., Russell, A. D., Spero, H. J., Palmer, M. R., Zeebe, R. E., and Eisenhauer, A.: The influence of symbiont photosynthesis on the boron isotopic composition of foraminifera shells, *Marine Micropaleontology*, 49, 87–96, 2003.
- 465 Huber, B. T., Bijma, J., and Darling, K.: Cryptic speciation in the living planktonic foraminifer *Globigerinella siphonifera* (d'Orbigny), *Paleobiology*, 23, 33–62, 1997.
- Hull, P. M.: Emergence of modern marine ecosystems, *Current Biology*, 27, R466–R469, 2017.
- Jørgensen, B. B., Erez, J., Revsbech, N. P., and Cohen, Y.: Symbiotic photosynthesis in a planktonic foraminiferan *Globigerinoides sacculifer* (Brady), studied with microelectrodes, *Limnology and Oceanography*, 30, 1253–1267, 1985.
- 470 Kolber, Z. S., Zehr, J., and Falkowski, P. G.: Effects of growth irradiance and nitrogen limitation on photosynthetic energy conversion in Photosystem II, *Plant Physiology*, 88, 923–929, 1988.
- Kolber, Z. S., and Falkowski, P. G.: Use of active fluorescence to estimate phytoplankton photosynthesis in situ, *Limnology and Oceanography*, 38, 1646–1665, 1993.
- Kolber, Z. S., Prášil, O., and Falkowski, P. G.: Measurements of variable chlorophyll fluorescence using fast repetition rate techniques: defining methodology and experimental protocols, *Biochimica et Biophysica Acta*, 1367, 88–106, 1998.
- 475 Lee, J. J.: Algal symbiosis in larger foraminifera, *Symbiosis*, 42, 63–75, 2006.
- Lee, J. J., Freudenthal, H. D., Kossov, V., and Bé, A. W. H.: Cytological observations on two planktonic foraminifera, *Globigerina bulloides* and *Globigerinoides ruber*, *The Journal of Protozoology*, 12, 531–542, 1965.
- Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., Zweng, M. M., Paver, C. R., Reagan, J. R., Johnson, D. R., Hamilton, M., and Seidov, D.: *World Ocean Atlas 2013, Volume 1: Temperature*, in: *NOAA Atlas NESDIS 73*, edited by: Levitus, S. and Mishonov, A., 40, 2013.
- 480 Lombard, F., Erez, J., Michel, E., and Labeyrie, L.: Temperature effect on respiration and photosynthesis of the symbiont-bearing planktonic foraminifera *Globigerinoides ruber*, *Orbulina universa*, and *Globigerinella siphonifera*, *Limnology and Oceanography*, 54, 210–218, 2009.
- Movellan, A.: *La biomasse des foraminifères planctoniques actuels et son impact sur la pompe biologique de carbone*. PhD Thesis, University of Angers, 2013.
- 485 Norris, R. D.: Biased extinction and evolutionary trends, *Paleobiology*, 17, 388–399, 1991.



- Norris, R. D.: Symbiosis as an evolutionary innovation in the radiation of Paleocene planktonic foraminifera, *Paleobiology*, 22, 461–480, 1996.
- 490 Parkhill, J.-P., Maillet, G., and Cullen, J. J.: Fluorescence-based maximal quantum yield for PSII as a diagnostic of nutrient stress, *Journal of Phycology*, 37, 517–529, 2001.
- Pearson, P. N., Shackleton, N. J., and Hall, M. A.: Stable isotope paleoecology of middle Eocene planktonic foraminifera and multi-species isotope stratigraphy, DSDP Site 523, South Atlantic, *Journal of Foraminiferal Research*, 23, 123–140, 1993.
- 495 Pillet, L., de Vargas, C., and Pawlowski, J.: Molecular identification of sequestered diatom chloroplasts and kleptoplastidy in foraminifera, *Protist*, 162, 394–404, 2011.
- R Core Team: R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, 2016.
- 500 Rebotim, A., Voelker, A. H. L., Jonkers, L., Waniek, J. J., Meggers, H., Schiebel, R., Fraile, I., Schulz, M., and Kucera, M.: Factors controlling the depth habitat of planktonic foraminifera in the subtropical eastern North Atlantic, *Biogeosciences*, 14, 827–859, 2017.
- Rink, S., Kühl, M., Bijma, J., and Spero, H. J.: Microsensor studies of photosynthesis and respiration in the symbiotic foraminifer *Orbulina universa*, *Marine Biology*, 131, 583–595, 1998.
- Schiebel, R., and Hemleben, C.: *Planktic Foraminifers in the Modern Ocean*, Springer-Verlag, Berlin Heidelberg, 2017.
- 505 Schmidt, C., Heinz, P., Kucera, M., and Uthicke, S.: Temperature-induced stress leads to bleaching in larger benthic foraminifera hosting endosymbiotic diatoms, *Limnology and Oceanography*, 56, 1587–1602, 2011.
- Seears, H. A., Darling, K. F., and Wade, C. M.: Ecological partitioning and diversity in tropical planktonic foraminifera, *BMC Evolutionary Biology*, 12:54, 2012.
- Shaked, Y., and de Vargas, C.: Pelagic photosymbiosis: rDNA assessment of diversity and evolution of dinoflagellate symbionts and planktonic foraminiferal hosts, *Marine Ecology Progress Series*, 325, 59–71, 2006.
- 510 Spero, H. J., and DeNiro, M. J.: The influence of symbiont photosynthesis on the  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  values of planktonic foraminiferal shell calcite, *Symbiosis*, 4, 213–228, 1987.
- Spero, H. J., and Parker, S. L.: Photosynthesis in the symbiotic planktonic foraminifer *Orbulina universa*, and its potential contribution to oceanic primary productivity, *Journal of Foraminiferal Research*, 15, 273–281, 1985.
- 515 Spindler M., and Hemleben, C.: Symbionts in planktonic Foraminifera (Protozoa), in: *Endocytobiology, endosymbiosis and cell biology*, edited by: Schwemmler, W. and Schenk, H. E. A., Walter de Gruyter & Co, Berlin, 133–140, 1980.
- Spindler M., Hemleben, C., Salomons, J. B., and Smit, L. P.: Feeding behavior of some planktonic foraminifers in laboratory cultures, *Journal of Foraminiferal Research*, 14, 237–249, 1984.
- Stoecker, D. K., Johnson, M. D., de Vargas, C., and Not, F.: Acquired phototrophy in aquatic protists, *Aquatic Microbial Ecology*, 57, 279–310, 2009.



- 520 Stoecker, D. K., Hansen, P. J., Caron, D. A., and Mitra, A.: Mixotrophy in the marine plankton, *Annual Reviews of Marine Science*, 9, 311–335, 2017.
- Takagi, H., Kimoto, K., Fujiki, T., Kurasawa, A., Moriya, K., and Hirano, H.: Ontogenetic dynamics of photosymbiosis in cultured planktic foraminifers revealed by fast repetition rate fluorometry, *Marine Micropaleontology*, 122, 44–52, 2016.
- 525 Takagi, H., Kimoto, K., Fujiki, T., and Moriya, K.: Effect of nutritional condition on photosymbiotic consortium of cultured *Globigerinoides sacculifer* (Rhizaria, Foraminifera), *Symbiosis*, 76, 25–39, 2018.
- Tolderlund, D. S., and Bé, A. W. H.: Seasonal distribution of planktonic Foraminifera in the western North Atlantic, *Micropaleontology*, 17, 297–329, 1971.
- Uthicke, S.: Photosynthetic efficiency and rapid light curves of sediment-biofilms along a water quality gradient in the Great  
530 Barrier Reef, Australia, *Marine Ecology Progress Series*, 322, 61–73, 2006.
- Yasuhara, M., Tittensor, D. P., Hillebrand, H., and Worm, B.: Combining marine macroecology and palaeoecology in understanding biodiversity: microfossils as a model, *Biological Reviews*, 92, 199–215, 2017.
- Weiner, A., Aurahs, R., Kurasawa, A., Kitazato, H., and Kucera, M.: Vertical niche partitioning between cryptic sibling species of a cosmopolitan marine planktonic protist, *Molecular Ecology*, 21, 4063–4073, 2012.
- 535 Wessel, P., and Smith, W. H. F.: New, improved version of generic mapping tools released, *Earth & Space Sciences News*, *Transactions of the American Geophysical Union*, 79, doi:10.1029/98EO00426, 1998.
- Ziegler, M., and Uthicke, S.: Photosynthetic plasticity of endosymbionts in larger benthic coral reef foraminifera, *Journal of Experimental Marine Biology and Ecology*, 407, 70–80, 2011.

540



545

**Table 1.** Summary of species symbiotic ecology. <sup>1</sup>Spindler and Hemleben (1980); <sup>2</sup>Taylor (1982); <sup>3</sup>Hemleben and Spindler (1983); <sup>4</sup>Gastrich (1987); <sup>5</sup>Faber et al. (1989); <sup>6</sup>Hemleben et al. (1989); <sup>7</sup>Gast and Caron (1996); <sup>8</sup>Huber et al. (1997); <sup>9</sup>Shaked and de Vargas (2006); <sup>10</sup>Gast et al. (2000); <sup>11</sup>Fujiki et al. (2014); <sup>12</sup>Bird et al. (2017); <sup>13</sup>Schiebel and Hemleben (2017); <sup>14</sup>Bird et al. (2018). \*Based on microscopic observations of living specimens in this study. Comparison with the other dinoflagellate-bearing species revealed almost identical features of the symbionts (e.g., cell size, shape, color, see Figure S3).

Species	Previous studies		This study							Remarks
	Algal type	Obligate / facultative / none	Ratio of symbiotic individuals	Test size-Chl <i>a</i> correlation	$F_v/F_m$	$\sigma_{PSII} (\times 10^{-20} \text{ m}^2 \text{ quanta}^{-1})$	Chl <i>a</i> /biomass (ng $\mu\text{g}^{-1}$ )	Cluster		
	Microscopy-based	Molecular-based								
<b><i>Orbulina universa</i></b>										
Dinoflagellate <sup>1,4</sup>	<i>Pelagodinium béii</i> (Dinoflagellate) <sup>7,9</sup>	Obligate <sup>6</sup>	0.95	Positive	0.50	448	4.65	1		
<b><i>Sphaeroidinella dehisces</i></b>										
Not reported	Not reported	Not reported	1.00	Positive	0.53	606	2.36	2	Presence of dinoflagellate symbionts inferred <sup>13</sup>	
<b><i>Globigerinoides sacculifer</i></b>										
Dinoflagellate <sup>1,4</sup>	<i>Pelagodinium béii</i> (Dinoflagellate) <sup>7,9</sup>	Obligate <sup>6</sup>	0.96	Positive	0.51	453	4.78	1		
<b><i>Globigerinoides conglobatus</i></b>										
Dinoflagellate <sup>1,4</sup>	<i>Pelagodinium béii</i> (Dinoflagellate) <sup>7</sup>	Obligate <sup>6</sup>	1.00	Positive	0.50	449	4.80	1		
<b><i>Globigerinoides ruber</i></b>										
Dinoflagellate <sup>1,4</sup>	<i>Pelagodinium béii</i> (Dinoflagellate) <sup>7,9</sup>	Obligate <sup>6</sup>	white	0.98	Positive	0.49	469	3.09	2	
			pink	0.91	Positive	0.52	374	2.28	2	
<b><i>Globoturborotalita rubescens</i></b>										
Not reported	Not reported	Not reported	0.79	Positive	0.46	388	1.13	2	Probably dinoflagellate-bearing*	
<b><i>Globoturborotalita tenella</i></b>										
Not reported	Not reported	Not reported	0.77	Positive	0.51	421	2.12	2	Probably dinoflagellate-bearing*	
<b><i>Globigerinella calida</i></b>										
Not reported	Not reported	Not reported	0.58	Positive	0.44	492	1.33	2		
<b><i>Globigerinella siphonifera</i></b>										
Haptophyte <sup>1</sup> (Prymnesiophyte <sup>2</sup> ) / two different chrysothycophyte <sup>4,5,8</sup>	Type I Unclassified Haptophyceae <sup>10</sup>	Obligate <sup>6</sup> / facultative <sup>5</sup>	0.81	Positive	0.47	515	2.56	2	Extracellular commensal algae reported <sup>8</sup>	
	Type II <i>Pelagomonas calceolata</i> (Pelagophyte) <sup>11</sup>	Obligate <sup>6</sup> / facultative <sup>5</sup>	0.95	Positive	0.49	689	2.42	2	Extracellular commensal algae absent <sup>8</sup>	

***Globigerinella adamsi***

Not reported	Not reported	Not reported	0.00	–	–	–	–	4
--------------	--------------	--------------	------	---	---	---	---	---

***Globigerina bulloides***

Barren <sup>3,4,6</sup> / synechococcus <sup>12</sup>	Synechococcus <sup>12</sup>	None <sup>3,4,6</sup>	0.00	–	–	–	–	4
--	-----------------------------	-----------------------	------	---	---	---	---	---

***Turborotalita quinqueloba***

Not reported	Not reported	Not reported	0.00	–	–	–	–	4
--------------	--------------	--------------	------	---	---	---	---	---

***Turborotalita humilis***

Dinoflagellate <sup>1</sup> / haptophyte <sup>3</sup> / chrysophyte <sup>4</sup>	Not reported	Obligate <sup>6</sup>	0.89	Not significant	0.51	710	0.58	3
--	--------------	-----------------------	------	--------------------	------	-----	------	---

***Hastigerina pelagica***

Barren <sup>1,3</sup>	Not reported	None <sup>3,4,6</sup>	0.00	–	–	–	–	4
-----------------------	--------------	-----------------------	------	---	---	---	---	---

***Hastigerinella digitata***

Not reported	Not reported	Not reported	0.00	–	–	–	–	4
--------------	--------------	--------------	------	---	---	---	---	---

***Neogloboquadrina incompta***

Not reported	Barren <sup>14</sup>	None <sup>14</sup>	0.00	–	–	–	–	4
--------------	----------------------	--------------------	------	---	---	---	---	---

***Neogloboquadrina pachyderm***

Not reported	Not reported	None <sup>6</sup>	0.00	–	–	–	–	4
--------------	--------------	-------------------	------	---	---	---	---	---

***Neogloboquadrina dutertrei***

Barren <sup>3</sup> / chrysophyte <sup>4</sup> / pelagophyte <sup>14</sup>	Pelagophyte <sup>14</sup>	Facultative <sup>6</sup>	0.94	Positive	0.48	749	0.60	2
--	---------------------------	--------------------------	------	----------	------	-----	------	---

***Pulleniatina obliquiloculata***

Prymnesiophyte <sup>2</sup> / Barren <sup>3</sup> / chrysophyte <sup>4</sup>	Not reported	Facultative <sup>6</sup>	0.66	Not significant	0.36	518	0.07	3
--	--------------	--------------------------	------	--------------------	------	-----	------	---

***Globorotalia inflata***

Barren <sup>3</sup> / chrysophyte <sup>4</sup>	Not reported	Facultative <sup>6</sup>	0.69	Not significant	0.33	544	0.19	3
--	--------------	--------------------------	------	--------------------	------	-----	------	---

***Globorotalia menardii***

Prymnesiophyte <sup>2</sup> / Barren <sup>3</sup> / chrysophyte <sup>4</sup>	Not reported	Facultative <sup>6</sup>	0.87	Positive	0.50	498	0.58	2
---	--------------	--------------------------	------	----------	------	-----	------	---

***Globorotalia scitula***

Not reported	Not reported	Not reported	0.00	–	–	–	–	4
--------------	--------------	--------------	------	---	---	---	---	---

***Globorotalia crassaformis***

Not reported	Not reported	Not reported	0.00	–	–	–	–	4
--------------	--------------	--------------	------	---	---	---	---	---



***Globorotalia truncatulinoides***

Barren <sup>1,3,4</sup>	Not reported	None <sup>3,4,6</sup>	0.00	–	–	–	–	4
-------------------------	--------------	-----------------------	------	---	---	---	---	---

***Candeina nitida***

Chrysophyte <sup>4</sup>	Not reported	Facultative <sup>6</sup>	0.88	Positive	0.49	347	1.48	2
--------------------------	--------------	--------------------------	------	----------	------	-----	------	---

***Globigerinita glutinata***

Barren <sup>3</sup> / chrysophyte <sup>4</sup>	Not reported	Facultative <sup>6</sup>	0.68	Positive	0.50	632	0.71	2
--	--------------	--------------------------	------	----------	------	-----	------	---

***Globigerinita uvula***

Not reported	Not reported	Not reported	0.79	Positive	0.35	618	0.47	2
--------------	--------------	--------------	------	----------	------	-----	------	---

***Tenuitella fleisheri***

Not reported	Not reported	Not reported	0.00	–	–	–	–	4
--------------	--------------	--------------	------	---	---	---	---	---

---



550

**Table 2.** Scaling exponents (slopes in Figs. 6 and 10) for relationships between test size and Chl *a* content. Reduced major axis regression was used to estimate the scaling exponents after logarithmic transformation of the two variables. CI; confidence interval. When the correlation was not significant, the values are not shown. N; the number of specimens used for the analysis.


Species / morphogroup	N	Scaling exponent		
		Best estimate	2.5 % CI	97.5 % CI
<i>Orbulina universa</i>	75	1.90	1.60	2.26
<i>Sphaeroidinella dehiscentis</i>	7	2.91	1.92	4.43
<i>Globigerinoides sacculifer</i>	94	3.10	2.66	3.60
<i>Globigerinoides conglobatus</i>	18	1.83	1.25	2.68
<i>Globigerinoides ruber</i> (white)	49	2.36	1.90	2.93
<i>Globigerinoides ruber</i> (pink)	40	2.62	2.24	3.07
<i>Globoturbotalita rubescens</i>	15	1.59	1.09	2.30
<i>Globoturbotalita tenella</i>	10	1.33	0.87	2.04
<i>Globigerinella calida</i>	11	3.71	2.10	6.56
<i>Globigerinella siphonifera</i> Type I	61	3.57	2.85	4.47
<i>Globigerinella siphonifera</i> Type II	53	2.89	2.28	3.66
<i>Turbotalita humilis</i>	17	–	–	–
<i>Neogloboquadrina dutertrei</i>	91	3.16	2.79	3.59
<i>Pulleniatina obliquiloculata</i>	45	–	–	–
<i>Globorotalia inflata</i>	9	–	–	–
<i>Globorotalia menardii</i>	144	1.84	1.63	2.08
<i>Candeina nitida</i>	32	3.20	2.37	4.31
<i>Globigerinita glutinata</i>	69	2.43	1.97	2.99
<i>Globigerinita uvula</i>	11	2.66	1.71	4.12
Macroperforate spinose with dinoflagellate	308	2.52	2.37	2.67
Macroperforate spinose with non-dinoflagellate	125	3.06	2.64	3.55
Macroperforate non-spinose	235	2.17	1.95	2.41
Microperforate non-spinose	112	2.61	2.30	2.95



560

565

570




**Figure 1.** Maps showing the cruise tracks (lines) and the sampling points (circles). (a) Central and western Pacific area, and (b) northeastern Atlantic area. For detail sampling information, see Table S1. Annual sea surface temperature (SST) data was from World Ocean Atlas 2013 (Locarnini et al., 2013).

575



580




**Figure 2.** Photomicrographs of representative individuals for species analyzed. (1) *Orbulina universa*, (2) *Sphaeroidinella dehiscentis*, (3) *Globigerinoides sacculifer*, (4) *Globigerinoides conglobatus*, (5) *Globigerinoides ruber* (white), (6) *Globigerinoides ruber* (pink), (7) *Globoturborotalita rubescens*, (8) *Globoturborotalita tenella*, (9) *Globigerinella calida*, (10) *Globigerinella siphonifera* Type I, (11) *Globigerinella siphonifera* Type II, (12) *Globigerinella adamsi*, (13) *Globigerina bulloides*, (14) *Turborotalita quinqueloba*, (15) *Turborotalita humilis*, (16) *Hastigerina pelagica*, (17) *Hastigerinella digitata*, (18) *Neogloboquadrina incompta*, (19) *Neogloboquadrina pachyderma*, (20) *Neogloboquadrina dutertrei*, (21) *Pulleniatina obliquiloculata*, (22) *Globorotalia inflata*, (23) *Globorotalia menardii*, (24) *Globorotalia scitula*, (25) *Globorotalia crassaformis*, (26) *Globorotalia truncatulinoidea*, (27) *Candeina nitida*, (28) *Globigerinita glutinata*, (29) *Globigerinita uvula*, (30) *Tenuitella fleisheri*. Scale bars are 200  $\mu\text{m}$ .



595

600




Term	Definition	Indication
$F$	Chlorophyll fluorescence (arbitrary unit)	Presence of chlorophyll
$F_m$	Maximum fluorescence (arbitrary unit)	Chl a content
$F_0$	Minimum fluorescence (arbitrary unit)	–
$F_v$	Variable fluorescence $F_m - F_0$ (arbitrary unit)	Presence of functional chlorophyll indicating the presence of symbionts
$F_v/F_m$	Potential photochemical efficiency (dimensionless)	Photosynthetic activity indicating vitality of symbionts
$\sigma_{PSII}$	Functional absorption cross section of PSII photochemistry ( $\times 10^{-20} \text{ m}^{-2} \text{ quanta}$ )	Light absorption efficiency

**Figure 3.** Schematic diagram of fluorescence induction curves by fast repetition rate fluorometry and their interpretation. (a) Profile of a symbiotic individual. (b) Profile of a non-functional chlorophyll-bearing individual. (c) Profile of a non-symbiotic individual. Photosystem II parameters used in this study are also listed. All parameters are obtained in dark-adapted states.

610



615




620

625

**Figure 4.** Workflow of this study and four indices used to characterize photosymbiosis. Firstly, individual specimens were identified to morphospecies level, measured for the test size, and analyzed with active fluorometry to check the functionality of chlorophyll. Based on the fluorescence results, intracellular chlorophyll types (status) were categorized into three groups; functional chlorophyll, non-functional chlorophyll, and no chlorophyll. When chlorophyll was functional, the content of Chl *a* per individual and the photophysiological parameters were analyzed. Finally, four indices in bold (symbiont possession rate, test size-Chl *a* relationship, Chl *a*/biomass, and  $F_v/F_m$ ) were derived and used for characterization of photosymbiosis (see text for detail).


630



635

**Figure 5.** Summary of categorization of intracellular chlorophyll. The functionality of chlorophyll indicates the presence of symbionts. Numbers of specimens for three categories are represented in parentheses (functional chlorophyll / non-functional chlorophyll / no chlorophyll). The percentage of functional chlorophyll are essentially the same as the symbiont possession rate used as a variable to characterize photosymbiosis (see text for detail).

640




645

**Figure 6.** Relationships between test size and Chl *a* content for each species. Lines represent reduced major axis regression ( $y$ ;  $\log(\text{Chl } a)$ ,  $x$ ;  $\log(\text{test size})$ ). Specimens with no chlorophyll and non-functional chlorophyll (NC) are plotted at the bottom of each panel to show their test size information (these data are not used for the regression analysis). R, Pearson's correlation coefficient; p, p-value; N, number of specimens with functional chlorophyll (i.e., with symbionts).



650

655




660


665

670

**Figure 7.** Ratios of Chl *a* content (ng foraminifer<sup>-1</sup>) to protein biomass (µg foraminifer<sup>-1</sup>) of 19 symbiont-bearing species. Dots represent individual data sampled from the upper 100 m water depth. Box plots represent first and third quartiles as hinges, and midlines as medians with notch representing 95 % confidence interval of the medians. Means are also represented with open diamonds. Whiskers are extended up to 1.5 times interquartile ranges from the end of each box to the furthest datum within that distance. Violin plots show the distributions as Kernel density estimation. Numbers at either end of the panel are the sample size for each species. Species with more than 20 specimens were used for statistical testing (Kruskal-Wallis test for comparison of differences in medians, and post-hoc Steel-Dwass test for multiple comparison,  $p < 0.05$ ). Species with the same letter were not significantly different. Color symbols represent the difference of symbiotic algae (see Table 1). Note that the data are represented on a logarithmic scale.



**Figure 8.** Photophysiological parameters of 19 symbiont-bearing species. (a)  $F_v/F_m$ , and (b)  $\sigma_{PSII}$ . Dots represent individual data sampled from the upper 100 m water depth. Box plots represent first and third quartiles as hinges, and midlines as medians with notch representing 95 % confidence interval of the medians. Means are also represented with open diamonds. Whiskers are extended up to 1.5 times interquartile ranges from the end of each box to the furthest datum within that distance. Violin plots show the distributions as Kernel density estimation. Numbers at either end of the panels are the sample size for each species. Species with more than 20 specimens were used for statistical testing (Kruskal-Wallis test for comparison of differences in medians, and post-hoc Steel-Dwass test for multiple comparison,  $p < 0.05$ ). Species sharing the same letter were not significantly different. Color symbols represent the difference of symbiotic algae (see Table 1).




**Figure 9.** Results of cluster analysis and principal component analysis. (a) Cluster dendrogram obtained using Ward's method. (b) Biplot of principal component analysis. The colors of the symbols correspond to the four clusters. Vectors indicate the direction and strength of each variable to the overall distribution. The first axis explains 84.2 % of the variation, and the second axis 10.2 %. *Chl a*/biomass; *Chl a* content per protein biomass estimated from test size of individuals, Size-Chl cor.; correlation coefficient of test size-*Chl a* content relationship as an indicator of the persistence of symbionts, Ratio of symb.; ratio of symbiotic individuals,  $F_v/F_m$ ; median  $F_v/F_m$  value.



690

695

700




705

**Figure 10.** Relationships between test size and Chl *a* content for four groups. The 16 species with significant test size-Chl *a* correlation were used. Lines represent reduced major axis regression ( $y$ ;  $\log(\text{Chl } a)$ ,  $x$ ;  $\log(\text{test size})$ ).  $R$ , Pearson's correlation coefficient;  $p$ ,  $p$ -value;  $N$ , number of specimens with functional chlorophyll (with symbionts).



710

715



720

725

730

**Figure 11.** A conceptual diagram of the spectrum of planktonic foraminiferal photosymbiosis along the trophic gradient between permanent endosymbiosis (right) resulting in permanently integrated plastid (not found in planktonic foraminifera) and heterotrophy (left). Foraminiferal species are ordinated on the basis of their PC1 score of the principal component analysis conducted in this study (Fig. 9). Foraminiferal photosymbiosis has been acquired regardless of their morphological features (i.e., spinose or non-spinose, macroperforate or microperforate).