

Saltwater reduces potential CO₂ and CH₄ production in peat soils from a coastal freshwater
forested wetland

Kevan J. Minick^{a*}, Bhaskar Mitra^b, Asko Noormets^b, John S. King^a

^a*Department of Forestry and Environmental Resources, North Carolina State University,
Raleigh, NC, 27695, USA*

^b*Department of Ecosystem Science and Management, Texas A&M University, College Station,
TX, 77843, USA*

**Corresponding author: email kjminick@ncsu.edu; phone (919) 630-3307; fax NA*

*Keywords: extracellular enzyme activity, sea-level rise, methanogenesis, microbial biomass
carbon, carbon isotopes, ghost forest*

Abstract A major concern for coastal freshwater wetland function and health are the effects of saltwater intrusion on greenhouse gas production from peat soils. Coastal freshwater forested wetlands are likely to experience increased hydroperiod with rising sea level, as well as saltwater intrusion. These potential changes to wetland hydrology may also alter forested wetland structure and lead to a transition from forest to shrub/marsh wetland ecosystems. Loss of forested wetlands is already evident by dying trees and dead standing trees (“ghost” forests) along the Atlantic Coast of the US, which will result in significant alterations to plant carbon (C) inputs, particularly that of coarse woody debris, to soils. We investigated the effects of salinity and wood C inputs on soils collected from a coastal freshwater forested wetland in North Carolina, USA, and incubated in the laboratory with either freshwater or saltwater (2.5 or 5.0 ppt) and with or without the additions of wood. Saltwater additions at 2.5 ppt and 5.0 ppt reduced CO₂ production by 41 and 37 %, respectively, compared to freshwater. Methane production was reduced by 98 % (wood-free incubations) and by 75-87 % (wood-amended incubations) in saltwater treatments compared to the freshwater plus wood treatment. Additions of wood also resulted in lower CH₄ production from the freshwater treatment and higher CH₄ production from saltwater treatments compared to wood-free incubations. The $\delta^{13}\text{CH}_4\text{-C}$ isotopic signature suggested that in wood-free incubations, CH₄ produced from the freshwater treatment originated primarily from the acetoclastic pathway, while CH₄ produced from the saltwater treatments originated primarily from the hydrogenotrophic pathway. These results suggest that saltwater intrusion into coastal freshwater forested wetlands will reduce CH₄ production, but long-term changes in C dynamics will likely depend on how changes in wetland vegetation and microbial function influence C cycling in peat soils.

1 Introduction

Sea-level rise (SLR) threatens coastal regions around the world. Significantly, the rate of SLR is not uniform around the globe, with the highest rate occurring along the Atlantic coast of North America between Cape Hatteras and Cape Cod due to factors including local currents, tides, and glacial isostatic rebound (Karegar et al., 2017; Sallenger et al., 2012). Along with economic and cultural impacts, health of coastal forested ecosystems are expected to be impacted by SLR (Langston et al., 2017; Kirwan and Gedan 2019). For instance, salinization of coastal freshwater wetlands will likely impact vegetation community dynamics and regeneration in low lying ($< 1\text{m}$) wetlands (Langston et al., 2017). Understanding how coastal wetland ecosystems respond to extreme events, long-term climate change and a rapidly rising sea is essential to developing the tools needed for sustainable management of natural resources, and the building of resilient communities and strong economies. Because it has more than $5,180\text{ km}^2$ of coastal ecosystems and urban areas below 1 m elevation, the state of North Carolina is highly vulnerable to climate change and SLR and therefore saltwater intrusion (Riggs and Ames, 2008, Titus and Richman, 2001).

As sea level changes, coastal plant communities move accordingly up and down the continental shelf. In recent geologic time, sea level has risen about 3 m over the past $\sim 2,500$ years from sea level reconstructions adjacent to our study site (Kemp et al., 2011). The rate of SLR has varied greatly over that time, with periods of stability and change, and a geologically unprecedented acceleration in recent decades. The current distribution of coastal forested wetlands reflects the hydrologic equilibrium of the recent past climate, but the widespread mortality of such forests suggests that the rate of SLR is in a time of rapid change at a rate

potentially faster than the forest's capacity to move upslope, resulting in widespread death of coastal freshwater forested wetlands (Kirwan and Gedan 2019). Furthermore, dying coastal forests will alter the quantity and quality of organic matter inputs to the soil as vegetation shifts occur, as well as introduce a large pulse of woody debris into soils. This has the potential to alter C cycling processes responsible for storage of C in peat soils or loss of C as CO₂ and CH₄ (Winfrey and Zeikus, 1977).

Wetlands store more than 25% of global terrestrial soil C in deep soil organic matter deposits due to their unique hydrology and biogeochemistry (Batjes, 1996; Bridgham et al., 2006). Carbon storage capacity is especially high in forested wetlands characterized by abundant woody biomass, forest floors of *Sphagnum* spp., and deep organic soils. Across the US Southeast, soil organic C (SOC) in soils increases with proximity to the coast and is greatest in coastal wetlands (Johnson and Kern, 2003). Carbon densities are even higher in the formations of organic soils (Histosols) that occur across the region, typically ranging from 687 to 940 t ha⁻¹, but can be as high as 1,447 t ha⁻¹ (Johnson and Kern, 2003). As noted, forested wetlands, which historically have contributed to terrestrial C sequestration, are in serious decline and processes leading to destabilization of accumulated soil C are not represented in broad-scale ecosystem and land-surface models. The extent of changes in soil C cycling processes attributable to altered hydroperiod, saltwater intrusion, and structural changes in vegetation in these ecosystems remains unclear.

Saltwater intrusion, a direct result of SLR, into freshwater wetlands alters soil C cycling processes (Ardón et al., 2016; Ardón et al., 2018), particularly that of methanogenesis (Baldwin et al., 2006; Chambers et al., 2011; Dang et al., 2018; Marton et al., 2012), and microbial activity (e.g., extracellular enzyme activity, Morrissey et al., 2014; Neubauer et al., 2013). Saltwater

contains high concentrations of ions, particularly sulfate (SO_4^{2-}), which support high rates of SO_4^{2-} reduction compared to freshwater wetlands (Weston et al., 2011). Sulfate acts as a terminal electron acceptor in anaerobic respiration of SOC, and SO_4^{2-} reducers will typically increase in abundance in response to saltwater intrusion and out-compete other anaerobic microorganisms, particularly methanogens, for C (Bridgham et al. 2013; Dang et al., 2019; Winfrey and Zeikus, 1977). The effect of SO_4^{2-} on soil C cycling and competitive interactions with other anaerobic microbial processes also appears dependent on the concentration of the ion (Chambers et al., 2011). Even within freshwater forested wetlands, hydrology and microtopography interact to influence the amount of SO_4^{2-} within soils experiencing different levels of saturation, and therefore rates of SO_4^{2-} reduction (Minick et al., 2019a). A majority of saltwater intrusion studies on soil C dynamics though have focused on tidal freshwater wetlands, whereas non-tidal freshwater wetlands have received relatively little attention, partially due to their more confined distribution across the landscape. Nonetheless, they occupy critical zones within the coastal wetland ecosystem distribution and will be influenced by SLR differently than that of tidal wetlands. Tidal wetlands may experience short-term pulses of saltwater with tidal movement of water, while SLR effects on saltwater intrusion into non-tidal freshwater wetlands may result in more long-term saltwater inundation. This difference in saltwater inundation period may influence rates of soil CO_2 , CH_4 production, and microbial activity (Neubauer et al., 2013); and therefore should be considered in light of the hydrologic properties of non-tidal wetlands.

Saltwater intrusion into freshwater systems may also influence the CH_4 production pathways (Dang et al., 2019; Weston et al., 2011), as a result of saltwater-induced shifts in methanogenic microbial communities (Baldwin et al., 2006; Chambers et al., 2011; Dang et al., 2019). Stable isotope analysis of CO_2 and CH_4 indicate that acetoclastic methanogenesis is the

major CH₄ producing pathway in freshwater wetlands (Angle et al., 2016), but the influence of saltwater on the pathway of CH₄ formation in non-tidal freshwater forested wetlands has rarely been studied, particularly through the lens of CO₂ and CH₄ stable C isotope analysis. As ¹³C isotopic analysis of CH₄ is non-destructive and is long-proven as a reliable indicator of the CH₄ production pathway (Whiticar et al., 1986), utilization of this analysis provides easily attainable information on the effects of freshwater compared to saltwater on CH₄ production dynamics in coastal wetland ecosystems experiencing SLR-induced changes in hydrology and vegetation.

Our goal in this study was to test whether saltwater additions alter the production of CO₂, CH₄, and microbial activity from organic soils of a non-tidal temperate freshwater forested wetland in coastal North Carolina, US, and whether effects differ in response to additions of wood. Although many studies have focused on salinity pulses in tidal freshwater wetlands, less attention has been given to the effects of sustained saltwater intrusion on soil C dynamics. We expect saltwater intrusion due to SLR will be more persistent in non-tidal wetlands. Therefore, we investigated the effects of sustained saltwater inundation, using a laboratory microcosm experiment, on greenhouse gas production and microbial activity (e.g., microbial biomass C and extracellular enzyme activity). Wood additions to microcosms were utilized to mimic the potential large pulses of wood to peat soils as forest dieback occurs along the aquatic-terrestrial fringes of the Atlantic Coast and these wetlands transition to shrub/marsh ecosystems (Kirwan and Gedan 2019); thereby providing a large and widespread pulse of coarse woody debris to wetland soils and potentially altering soil C cycling.

2 Methods

2.1 Field Site Description

The field site was located in the Alligator River National Wildlife Refuge (ARNWR) in Dare County, North Carolina (35°47'N, 75°54'W) (Figure 1). The ARNWR was established in 1984 and is characterized by a diverse assemblage of non-tidal pocosin wetland types (Allen et al., 2011). ARNWR has a network of roads and canals, but in general contains vast expanses of minimally disturbed forested- and shrub-wetlands. Thirteen plots were established in a 4 km² area in the middle of a bottomland hardwood forest surrounding a 35-meter eddy covariance flux tower (US-NC4 in the AmeriFlux database; Minick et al., 2019a). Of the 13 plots (7 m radius), four central plots were utilized for this study which have been more intensively measured for plant and soil properties and processes (Miao et al. 2013, Miao et al., 2017, Minick et al 2019a, 2019b, Mitra et al. 2019). Over-story plant species composition was predominantly composed of black gum (*Nyssa sylvatica*), swamp tupelo (*Nyssa biflora*), bald cypress (*Taxodium distichum*), with occasional red maple (*Acer rubrum*), sweet gum (*Liquidambar styraciflua*), white cedar (*Chamaecyparis thyoides*), and loblolly pine (*Pinus taeda*). The understory was predominantly fetterbush (*Lyonia lucida*), bitter gallberry (*Ilex albra*), red bay (*Persea borbonia*), and sweet bay (*Magnolia virginiana*). Mean air temperature and precipitation from climate records of an adjacent meteorological station (Manteo AP, NC, 35°55'N, 75°42'W, National Climatic Data Center) for the period of 2008 – 2018 was 17.0 ± 0.30 °C and 932 ± 38 mm, respectively. These wetlands are characterized by a hydroperiod that responds over short time scales and is driven primarily by variable precipitation patterns. Soils are classified as a Pungo series (very poorly drained dystic thermic typic Haplosaprist) with a deep, highly decomposed muck layer overlain by a shallow, less decomposed peat layer and underlain by highly reduced mineral sediments of

Pleistocene origin (Riggs, 1996). Soils from the surface of hummocks have a pH of 4.2 ± 0.1 , C concentration of 49 ± 1.3 %, and a $\delta^{13}\text{C}$ value of -29.1 ± 0.29 ‰ (Minick et al. 2019b). Ground elevation is below < 1 m above sea level. Sea-level rise models of coastal NC show that ARNWR will experience almost complete inundation by 2100, with attendant shifts in ecosystem composition (DOD, 2010).

2.2 Sample Collection

Soil samples were collected on February 6, 2018, from surface organic soils by removing seven $10 \times 10 \text{ cm}^{-2}$ monoliths from hummocks to the depth of the root mat (approximately 6.3 cm) using a saw and a $10 \times 10 \text{ cm}^{-2}$ PVC square. The seven soil samples were composited by plot and stored on ice for transport back to the laboratory. In the laboratory, roots and large organic matter were removed by hand and gently homogenized. Soils samples were then stored in the dark at 4°C for seven weeks before initiating the laboratory incubation.

Freshwater and saltwater for the experiment was collected from water bodies surrounding the ARNWR on March 7, 2018 (Figure 1). Freshwater was collected from Milltail Creek, which runs Northwest from the center of ARNWR to Alligator River and drains our forested wetland study site. Freshwater salt concentration was 0 ppt. Saltwater was collected from Roanoke Sound to the east of ARNWR and had a salt concentration of 19 ppt (Figure 1). Freshwater and saltwater were mixed together to get the desired salt concentration for the saltwater treatments (2.5 and 5.0 ppt). These concentrations of saltwater were chosen due to the salinity levels in the Croatan and Pamlico Sounds, which are adjacent to ARNWR (Figure 1). Salinity in these waters range from approximately 1 to 5 ppt (unpublished data). Prior to mixing, freshwater and

saltwater was filtered through a Whatman #2 filter (8 μm). Neither saltwater nor freshwater were sterile filtered, therefore microbial communities from each water source were mixed together and added to the incubations. This could influence the response of soil microbes to the various treatments, but also represents what would occur under future projections of SLR in this region and the resulting mixing of freshwater and saltwater within the wetland. Four water samples of each freshwater and saltwater mixture were sent to the NCSU Environmental and Agricultural Testing Service laboratory for analysis of total organic C (TOC), ammonium (NH_4^+), nitrate (NO_3^-), phosphate (PO_4^-), SO_4^- , calcium (Ca^{2+}), magnesium (Mg^{2+}), sodium (Na^+), potassium (K^+), and chlorine (Cl^-). Analysis of TOC was made using a TOC analyzer (Schimadzu Scientific Instruments, Durham, NC). Analysis of NH_4^+ , NO_3^- , and PO_4^- , was made using Lachat Quikchem 8500 flow injection analysis system (Lachat Instruments, Milwaukee, WI). Sulfate and Cl^- were measured on a Dionex ion chromatograph (Thermo Fisher Scientific, Waltham, MA). Finally, a Perkin Elmer 8000 inductively-coupled plasma-optical emission spectrometer (Perkin Elmer, Waltham, MA) was used to analyze water samples for Ca^{2+} , Mg^{2+} , Na^+ , K^+ , and Cl^- .

2.3 Incubation Setup

Incubation water treatments included: 1) soils incubated at 65 % water holding capacity (WHC) (Dry); 2) soils incubated at 100% WHC with freshwater (0 ppt); 3) soils incubated at 100% WHC with a saltwater concentration of 2.5 ppt (2.5 ppt); and 4) soils incubated at 100% WHC with a saltwater concentration of 5.0 ppt (5.0 ppt). A subsample of each fresh soil (soils stored at 4 °C) was dried at 105°C to constant mass to determine gravimetric soil water content.

208 Approximately 150 – 200 g fresh soil (20 – 25 g dry weight) collected from each plot was
209 weighed into 1 L canning jars. For water addition estimates, WHC was calculated by placing a
210 subsample of fresh soil (approximately 2 g fresh weight) in a funnel with a Whatman #1 filter
211 and saturating with deionized H₂O (dH₂O). The saturated sample was allowed to drain into a
212 conical flask for 2 h. After 2 h, the saturated soil was weighed, dried at 105 °C to constant mass,
213 and weighed again to determine WHC. It is important to note that the 100% WHC moisture
214 level resulted in soils being completely flooded (with either freshwater or saltwater) with water
215 covering the surface of the incubated soils, thereby allowing for the development of CH₄
216 producing conditions similar to that observed in the field for surface soils. After soil and water
217 additions, the remaining headspace was estimated for each individual incubation vessel
218 (approximately 750 mL) and used in the calculation of gas production rates. Following wood-
219 additions (see below), incubation vessels from each of the eight treatments were incubated in the
220 dark in the laboratory for 98 d at 20 – 23 °C.

221 Two sets of incubations were set up with the above mentioned water treatments. We
222 added ¹³C-depleted American sweetgum (*Liquidamber styraciflua*) wood to half the incubation
223 vessels (0.22 g wood per g soil) (wood-amended), while the other half were incubated without
224 wood (wood-free). Trees were grown at the Duke FACE site under elevated CO₂ concentrations
225 (200 ppm CO₂ above ambient) using natural gas derived CO₂ with a depleted ¹³C signature
226 compared to that of the atmosphere (Feng et al., 2010; Schlesinger et al., 2006). The site was
227 established in 1983 after clear cut and burn (Kim et al., 2016). Trees were grown under elevated
228 CO₂ from 1994 to 2010 at which point they were harvested (Kim et al., 2016). Cookies were
229 removed from harvested trees, dried to a constant moisture level and stored at -20 °C until use.
230 The bark layer was removed and the outer six tree rings of multiple cookies were removed with a

chisel. Wood was then finely ground in a Wiley Mill (Thomas Scientific, Swedesboro, NJ, USA) and analyzed for C content and ^{13}C signature on a Picarro G2201-i Isotopic CO_2/CH_4 Analyzer outfitted with a Costech combustion module for solid sample analysis (Picarro Inc., Sunnyvale, CA USA). For $\delta^{13}\text{C}$ analysis of solids (e.g., wood, microbial biomass extracts, soils), certified solid standards were used to develop a standard curve from the expected and measured $\delta^{13}\text{C}$ values ($R^2 > 0.999$). These standards included USGS 40 (L-glutamic acid) ($\delta^{13}\text{C} = -26.39$ ‰; USGS Reston Stable Isotope Laboratory, Reston, VA, USA), protein ($\delta^{13}\text{C} = -26.98$ ‰; Elemental Microanalysis Ltd, Okehampton, UK), urea ($\delta^{13}\text{C} = -48.63$ ‰; Elemental Microanalysis Ltd, Okehampton, UK), atropine ($\delta^{13}\text{C} = -18.96$ ‰; Costech Analytical Technologies, Inc, Valencia, CA, USA), and acetanilide ($\delta^{13}\text{C} = -28.10$ ‰; Costech Analytical Technologies, Inc, Valencia, CA, USA). For C concentration, atropine standards were weighed out over a range of C concentrations that encompassed the expected C concentrations of the unknown samples and within the measurement range of the instrument. A standard curve for C concentration was also developed from the expected and measured C concentration of the atropine standards ($R^2 > 0.99$). All unknown sample's C concentration and $\delta^{13}\text{C}$ value were adjusted using the linear equations derived from the appropriate standard curve. The $\delta^{13}\text{C}$ values were reported in parts per thousand (‰) relative to the Vienna Pee Dee Belemnite (VPDB) standard. Wood had a C content of 45.6 ± 0.21 % and $\delta^{13}\text{C}$ value of -40.7 ± 0.06 ‰, which was within the range of -42 to -39 ‰ measured on fresh pine needles and fine roots (Schlesinger et al., 2006), and more depleted in ^{13}C compared to that measured in hummock surface soils from our site (-29.1 ± 0.29 ‰; Minick et al. 2019b).

2.4 CO_2 and CH_4 Sample Collection and Analysis

Headspace gas samples were collected from incubation vessels 15 times over the course of the 98 d incubation (days 1, 4, 8, 11, 15, 19, 25, 29, 29, 47, 56, 63, 70, 84, 98). Incubation lids were loosened between measurements to allow for gas exchange with the ambient atmosphere. Four blank incubations (empty jars; no soil, water, or wood) were set up and treated in the exact same manner as incubations containing soil, water, and wood. Blanks were used to measure soil-free CO₂ and CH₄ concentrations in incubations, which were always well below the detection limit of the gas analyzer (described below). Prior to each measurement, incubation vessels were removed from the dark, sealed tightly, and flushed at 20 psi for three minutes with CO₂/CH₄ free zero air (Airgas, Radnor, PA, USA). Following flushing, incubation vessels were immediately placed back in the dark (2-6 h over the first 39 days and 12-18 h over the remainder of the incubation) before taking a gas sample for analysis. Approximately 300 mL of headspace gas was removed using a 50 mL gas-tight syringe and transferred to an evacuated 0.5 L Tedlar gas sampling bag (Restek, Bellefonte, PA, USA). Simultaneous analysis of CO₂ and CH₄ concentrations and $\delta^{13}\text{C}$ isotopic signature were conducted on a Picarro G2201-i Isotopic CO₂/CH₄ Analyzer (Picarro Inc., Sunnyvale, CA USA). For $\delta^{13}\text{C}$ analysis of gases (e.g., CO₂ and CH₄), certified gas standards were used to develop a standard curve from the expected and measured $\delta^{13}\text{C}$ values ($R^2 > 0.99$). The gas standards for ¹³CO₂ analysis included gas tanks containing: 1) 372 ppm CO₂ with a $\delta^{13}\text{C}$ value of -11.0 ± 0.25 ‰ (Airgas, Inc., Radnor, PA); 2) 420 ppm CO₂ with a $\delta^{13}\text{C}$ value of -10.3 ± 0.18 ‰ (Airgas, Inc., Radnor, PA); 3) 768 ppm CO₂ with a $\delta^{13}\text{C}$ value of -29.5 ± 0.14 ‰ (Airgas, Inc., Radnor, PA); and 4) 3000 ppm CO₂ with a $\delta^{13}\text{C}$ value of -34.4 ± 0.3 ‰ (Airgas, Inc., Radnor, PA). The gas standards for ¹³CH₄ analysis included gas tanks containing: 1) 1.75 ppm CH₄ with a $\delta^{13}\text{C}$ value of -43.2 ± 0.07 ‰ (Airgas,

Inc., Radnor, PA); 2) 2.00 ppm CH₄ with a $\delta^{13}\text{C}$ value of -42.7 ± 0.20 ‰ (Airgas, Inc., Radnor, PA); 3) 10.00 ppm CH₄ with a $\delta^{13}\text{C}$ value of -68.6 ± 1.00 ‰ (Airgas, Inc., Radnor, PA); and 4) 15.08 ppm CH₄ with a $\delta^{13}\text{C}$ value of -29.5 ± 0.14 ‰ (Airgas, Inc., Radnor, PA). For CO₂ and CH₄ concentration, a concentrated gas standard (gas mix containing 4043 ppm CO₂ and CH₄) (Airgas, Inc., Radnor, PA) was diluted with zero air gas, providing a range of CO₂ and CH₄ concentrations that encompassed the expected gas concentrations of the unknown samples. A standard curve for gas concentration was developed from the expected and measured gas concentration of the diluted gas standards ($R^2 > 0.99$). All unknown gas sample CO₂ and CH₄ concentrations and $\delta^{13}\text{C}$ values were adjusted using the linear equations derived from the appropriate standard curve. The $\delta^{13}\text{C}$ values were reported in parts per thousand (‰) relative to the Vienna Pee Dee Belemnite (VPDB) standard. Production rates of CO₂-C and CH₄-C were calculated as well as daily cumulative CO₂-C and CH₄-C production summed over the course of the 98 d incubation. Small subsamples (approximately 1.0 g dry weight) of soil were removed periodically from each incubation vessel for extracellular enzyme analysis (see below). Removal of soil was accounted for in subsequent calculations of gas production rates. Incubation vessel water levels (mass basis) were checked and adjusted three times per week using either freshwater or saltwater.

The proportion of wood-derived CO₂ at each sampling date was calculated using ¹³CO₂ data and the ¹³C of depleted wood (-40.07) in a two pool flux model (Fry 2006), with the depleted wood signature as one end-point and the ¹³CO₂ of wood-free incubations as the other endpoint.

$$\% \text{ C} = ((\delta^{13}\text{C}_{\text{CO}_2\text{wood} + \text{soil}} - \delta^{13}\text{C}_{\text{CO}_2\text{wood-free soil}}) / (\delta^{13}\text{C}_{\text{wood}} - \delta^{13}\text{C}_{\text{CO}_2\text{wood-free soil}})) * 100$$

Where $\delta^{13}\text{C}_{\text{CO}_2\text{wood+ soil}}$ is the $\delta^{13}\text{C}$ value of CO_2 produced from soils incubated with the addition of ^{13}C -depleted wood, $\delta^{13}\text{C}_{\text{wood-free soil}}$ is the $\delta^{13}\text{C}$ value of CO_2 produced from soils incubated without the addition of ^{13}C -depleted wood, and $\delta^{13}\text{C}_{\text{wood}}$ is the average $\delta^{13}\text{C}$ value of the ^{13}C -depleted wood. Total wood-derived CO_2 was calculated using cumulative CO_2 produced over the 98 d incubation and the average $^{13}\text{CO}_2$ across the whole incubation.

2.5 Soil Characteristics

Soil organic C concentration and $\delta^{13}\text{C}$ were analyzed on the four replicate soil samples prior to the start of the incubation (initial soil samples) and on soils from each of the thirty incubations following the 98 d incubation period. The initial C analysis was performed on samples removed prior to incubation. Soils were finely ground in a Wiley Mill (Thomas Scientific, Swedesboro, NJ, USA) prior to analysis on a Picarro G2201-i Isotopic CO_2/CH_4 Analyzer outfitted with a Costech combustion module for solid sample analysis (Picarro Inc., Sunnyvale, CA USA). Carbon concentration and ^{13}C calibration standards were the same as those described for the analysis of the ^{13}C -depleted wood.

Soil pH and redox potential ($E_h = \text{mV}$) were measured in each incubation within one hour following sampling of headspace gas. Soil pH was measured on the four replicate soil samples immediately prior to the start of the incubation with a glass electrode in a 1:2 mixture (by mass) of soil and distilled water (dH_2O). Soil redox potential ($E_h = \text{mV}$) was measured using a Martini ORP 57 ORP/ $^{\circ}\text{C}/^{\circ}\text{F}$ meter (Milwaukee Instruments, Inc., Rocky Mount, NC, USA) .

2.6 Microbial Biomass Carbon and $\delta^{13}\text{C}$ Isotopic Signature

Microbial biomass C (MBC) was estimated on soils collected from incubations on day 1 (after 24 hour post-treatment incubation) and day 98 (following the end of the incubation). The chloroform fumigation extraction (CFE) method was adapted from Vance et al. (1987) in order to estimate MBC and $\delta^{13}\text{C}$. Briefly, one subsample of soil (approximately 0.5 g dry weight each) was placed in a 50 mL beaker in a vacuum desiccator to be fumigated. Another subsample was placed into an extraction bottle for immediate extraction in 0.5 M K_2SO_4 by shaking for 1 hr and subsequently filtering through Whatman #2 filter paper to remove soil particles. The samples in the desiccator were fumigated with ethanol-free chloroform (CHCl_3) and incubated under vacuum for 3 d. After the 3 d fumigation, samples were extracted similar to that of non-fumigated samples. Filtered 0.5 M K_2SO_4 extracts were dried at 60 °C in a ventilated drying oven and then ground to a fine powder with mortar and pestle before analysis of C concentration and $\delta^{13}\text{C}$ on a Picarro G2201-i Isotopic CO_2/CH_4 Analyzer outfitted with a Costech combustion module for solid sample analysis (Picarro Inc., Sunnyvale, CA USA). Carbon concentration and ^{13}C calibration standards were the same as those described for the analysis of the ^{13}C -depleted wood. Microbial C biomass was determined using the following equation:

$$\text{MBC} = \text{EC} / k_{\text{EC}}$$

where the chloroform-labile pool (EC) is the difference between C in the fumigated and non-fumigated extracts, and k_{EC} (extractable portion of MBC after fumigation) is soil-specific and estimated as 0.45 (Joergensen, 1996).

The $\delta^{13}C$ of MBC was estimated as the $\delta^{13}C$ of the C extracted from the fumigated soil sample in excess of that extracted from the non-fumigated soil sample using the following equation:

$$\delta^{13}C_{MBC} (\text{‰}) = ((\delta^{13}C_f \times C_f) - (\delta^{13}C_{nf} \times C_{nf})) / (C_f - C_{nf})$$

where C_f and C_{nf} is the concentration (mg kg^{-1} soil) of C extracted from the fumigated and non-fumigated soil samples, respectively, and $\delta^{13}C_f$ and $\delta^{13}C_{nf}$ is the ^{13}C natural abundance (‰) of the fumigated and non-fumigated soil samples, respectively.

2.5 Extracellular Enzyme Analysis

The potential activity of five extracellular enzymes was quantified on soil samples and on days 1, 8, 35, and 98 of the soil incubation. The enzymes chosen for this experiment represent a range of compounds in which they degrade, including fast and slow cycling C compounds, as well as ones that target nitrogen, phosphorus, and sulfate containing compounds. The Enzyme Commission number (EC) is stated in parenthesis after each enzyme, which classifies them by the chemical reaction catalyzed by each enzyme. The specific enzymes measured were: β -glucosidase (BG; EC: 3.2.1.21), xylosidase (XYL; EC 3.2.1.37), peroxidase (PER; EC: 1.11.1.7), β -glucosaminidase (NAGase; EC: 3.2.1.30), alkaline phosphatase (AP; EC: 3.1.3.1), and

arylsulfatase (AS; EC: 3.1.6.1). Carbon-degrading enzymes BG, XYL, and PER degrade sugar, hemicellulose, and lignin, respectively, while the N-degrading enzyme, NAGase, degrades chitin. Enzymes AP and AS degrade phosphorus and SO_4^{2-} containing compounds, respectively. Substrates for all enzyme assays were dissolved in 50 mM, pH 5.0 acetate buffer solution for a final concentration of 5 mM substrate.

Hydrolytic enzymes (BG, XYL, NAGase, AP, and AS) were measured using techniques outlined in Sinsabaugh et al. (1993). Approximately 0.8 g dry weight of soil sample was suspended in 50 mL of a 50 mM, pH 5.0 acetate buffer solution and homogenized in a blender for 1 min. In a 2 mL centrifuge tube, a 0.9 mL aliquot of the soil-buffer suspension was combined with 0.9 mL of the appropriate 5 mM p-nitrophenyl substrate solution for a total of three analytical replicates. Additionally, duplicate background controls consisting of 0.9 mL aliquot of soil-buffer suspension plus 0.9 mL of acetate buffer were analyzed, as well as four substrate controls consisting of 0.9 mL substrate solution plus 0.9 mL buffer. The samples were agitated for 2-5 hr. Samples were then centrifuged at 8,160 g for 3 min. Supernatant (1.5 mL) was transferred to a 15 mL centrifuge tube containing 150 μL 1.0 M NaOH, followed by the addition of 8.35 mL dH_2O . The resulting mixture was vortexed and a subsample transferred to a cuvette and the optical density at 410 nm was measured on a spectrophotometer (Beckman Coulter DU 800 Spectrophotometer, Brea, CA, USA).

The oxidative enzyme (PER) was measured using techniques outlined in Sinsabaugh et al. (1992). PER is primarily involved in oxidation of phenolic compounds and depolymerization of lignin. The same general procedure for hydrolytic enzymes was followed utilizing a 5 mM L-3,4-Dihydroxyphenylalanine (L-DOPA) (Sigma-Aldrich Co. LLC, St. Louis, MO, USA) solution as the substrate plus the addition of 0.2 mL of 0.3% H_2O_2 to all sample replicates and substrate

controls. After set up of analytical replicates and substrate and background controls, the samples were agitated for 2-3 hr. Samples were then centrifuged at 8,160 g for 3 min. The resulting supernatant turns an intense indigo color. Supernatant (1.4 mL) was transferred directly to a cuvette and the optical density at 460 nm was measured on a spectrophotometer.

For all enzymes, the mean absorbance of two background controls and four substrate controls was subtracted from that of three analytical replicates and divided by the molar efficiency ($1.66/\mu\text{mol}$), length of incubation (h), and soil dry weight. Enzyme activity was expressed as μmol substrate converted per g dry soil mass per hour ($\mu\text{mol g}^{-1} \text{h}^{-1}$). Daily cumulative enzyme activity was calculated and summed over the course of the 98 d incubation.

2.6 Statistical Analysis

Water chemistry, cumulative CO_2 production, cumulative CH_4 production, cumulative enzyme activity, post-incubation SOC concentration and $\delta^{13}\text{C}$, and wood-derived and wood-associated SOC, CO_2 , and MBC were analyzed using a one-way ANOVA (PROC GLM package). Microbial biomass C, MBC ^{13}C , pH, Eh, $\delta^{13}\text{CO}_2$, and $\delta^{13}\text{CH}_4$ were analyzed using repeated-measures ANOVA (PROC MIXED package) with time (Time) as the repeated measure and the incubation treatment as the fixed effect. All data for wood-free and wood-amended soils were analyzed separately. Raw data were natural log-transformed where necessary to establish homogeneity of variance. If significant main effects or interactions were identified in the one-way or repeated-measures ANOVA ($P < 0.05$), then post-hoc comparison of least-squares means was performed. All statistical analyses were performed using SAS 9.4 software (SAS Institute, Cary, NC, USA).

3 Results

3.1 Water and Soil Properties

Freshwater had higher concentrations of TOC compared to the saltwater treatments (Table 1). Concentration of SO_4^{2-} , Cl^- , Na^+ , Ca^{2+} , Mg^{2+} , and K^+ were higher in saltwater treatments compared to freshwater and were approximately twice as high in the 5.0 ppt saltwater treatment compared to 2.5 ppt saltwater (Table 1).

Initial (pre-incubation) SOC concentration was $490 \pm 27 \text{ g kg}^{-1}$ with a $\delta^{13}\text{C}$ value of $-28.5 \pm 0.32 \text{ ‰}$. After 98 d of incubation, SOC concentration in wood-free incubations was lower in the 5.0 ppt saltwater treatment, although no difference in soil $\delta^{13}\text{C}$ was found between treatments (Table 2). For wood-amended incubations, post-incubation SOC concentration was lower in the 5.0 ppt saltwater treatment compared to the dry and freshwater treatment (Table 2). Overall, the $\delta^{13}\text{C}$ of wood-free ($-29.5 \pm 0.08 \text{ ‰}$) and wood-amended soils ($-30.5 \pm 0.12 \text{ ‰}$) after 98 days of incubation were significantly different ($F = 49.6$; $P < 0.0001$).

Soil pH was significantly lower in the saltwater treatments in both wood-free and wood-amended soils compared to the dry and freshwater treatments (Table 3; Figure 2A-B). After an initial drop of pH in saltwater treatments (wood-free and wood-amended) to between 3.2 and 3.4 pH, pH steadily climbed back up to between 3.8 and 4.2 pH (Figure 2A-B). In wood-free soils, differences in soil Eh between treatments was variable over time, with both the 5.0 ppt saltwater treatment and the freshwater treatment having the lowest redox potential at different time points throughout the incubation (Table 3; Figure 2C), but fell below -124 mV on average. In wood-

amended soils, Eh dropped quickly to between -200 and -400 mV over the first 30 days for saltwater incubated soils (Table 3; Figure 2D), before rising to between -100 to 0 mV for the rest of the incubation period. In freshwater incubated soils, Eh rose quickly back to between -50 to 50 mV by day 15 and remained at this level for the rest of the incubation period, while saltwater treatments had significantly lower Eh between days 8 and 25.

3.2 CO₂, CH₄, $\delta^{13}\text{CO}_2\text{-C}$, and $\delta^{13}\text{CH}_4\text{-C}$

In wood-free incubations, cumulative CO₂ production was not different between the dry and freshwater treatments, but was higher than that produced from saltwater treatments (Table 4; Figure 3A). Cumulative CO₂ produced from wood-amended soils was highest in the dry treatment compared to all other treatments (Table 4; Figure 3B). Wood-derived CO₂ (calculated as the difference between cumulative CO₂ produced from wood-amended and wood-free incubations) was highest in the dry treatment (Table 4; Figure 3C). This finding was also confirmed by calculating cumulative wood-derived C using the ¹³C two-pool mixing model, with the highest proportion found in the dry treatment ($54 \pm 4.6 \%$) compared to soils incubated with freshwater ($42 \pm 1.7 \%$), 2.5 ppt saltwater ($37 \pm 1.0 \%$), and 5.0 ppt saltwater ($38 \pm 1.5 \%$) ($F = 10.1$; $P = 0.001$).

Cumulative CH₄ production was highest in the freshwater treatment compared to the saltwater treatments in both wood-free and wood-amended incubations (Table 4; Figure 3D-E). The difference between cumulative CH₄ produced from wood-amended and wood-free incubations was lower (and exhibited a negative response to wood additions) in the freshwater

treatment compared to both saltwater treatments (Table 3; Figure 3F), which both had a slight positive response to wood additions.

The CO₂:CH₄ ratio, in wood-free incubations, was calculated only for soils incubated under saturated conditions with freshwater or saltwater. The CO₂:CH₄ ratio, in wood-free incubations, was highest in freshwater (6 ± 3.4), compared to the 2.5 ppt saltwater (136 ± 33.9) and 5.0 ppt saltwater (102 ± 30.3) ($F = 24.8$; $P = 0.0002$). The CO₂:CH₄ ratio, in wood-amended incubations, was highest in freshwater (9 ± 0.8), compared to the 2.5 ppt saltwater (53 ± 20.3) and 5.0 ppt saltwater (107 ± 37.7) ($F = 9.2$; $P = 0.007$).

The $\delta^{13}\text{CO}_2\text{-C}$ and wood-derived CO₂ (estimated by ^{13}C two-pool mixing model) exhibited a time by treatment interaction for both wood-free and wood-amended incubations (Table 3; Figure 4A-B). In general, $\delta^{13}\text{CO}_2\text{-C}$ in wood-free and wood-amended incubations was depleted in the dry treatment (and remained steady throughout the incubation period) compared to all other treatments, especially after day 15. The proportion of wood-derived CO₂ was initially higher in freshwater and saltwater treatments (after day 1) but gradually dropped over the course of the incubation, while the proportion of wood-derived CO₂ from the dry treatment dropped quickly after the first sampling date (day 1) and remained steady (approximately 50-60 %) for the remainder of the incubation period (Figure 4C).

The $\delta^{13}\text{CH}_4\text{-C}$ (Table 3; Figure 5) exhibited a treatment and time effect (Table 3; Figure 5A-B), but only for wood-free incubations. For wood-free incubations, average $^{13}\text{CH}_4\text{-C}$ across the course of the incubation was enriched in the freshwater treatment (-67.8 ± 2.4 ‰) compared to the 2.5 ppt (-80.1 ± 2.4 ‰) and 5.0 ppt (-82.3 ± 2.0 ‰) saltwater treatments (Figure 5C). No difference in the $\delta^{13}\text{CH}_4\text{-C}$ was found in wood-amended incubations (Figure 4b, d), which ranged from between -78 to -75 ‰ for all treatments.

3.3 Microbial Biomass Carbon and Extracellular Enzyme Activity

Initially, MBC was lowest in the dry treatment of wood-free incubations and in the 5 ppt treatment of wood-amended incubations (Table 3; Table 5). Following the 98 day incubation, MBC was highest in the dry treatment of wood-free incubations, with no differences between the other treatments. In wood-amended incubations, final MBC was also highest in the dry treatment compared to both saltwater treatments. Initial $\delta^{13}\text{C}$ of MBC did not differ between treatments in either the wood-free or wood amended soils (Table 3; Table 5). After the 98 day incubation, ^{13}C of MBC in the wood-free treatments was depleted in the freshwater treatment and enriched in the 5.0 ppt saltwater treatment. In wood-amended incubations, ^{13}C of MBC was depleted in the dry treatment and enriched in the freshwater and 5.0 ppt saltwater treatments. Furthermore, the proportion of wood-derived MBC (as estimated by ^{13}C mixing model calculations) was highest in the dry treatment (31 %) and the 2.5 ppt saltwater treatment (21%) compared to the freshwater treatment (4%) (Table 5).

In wood-free incubations, activity of BG and NAGase was higher, while PER was lower, in the dry treatment compared to the saltwater treatments (Table 4; Table 5). Activity of AS was higher in the dry and freshwater treatments compared to saltwater treatments, in both wood-free and wood-amended incubations. In wood-amended incubations, BG and NAGase were highest in the dry treatment compared to the saltwater treatments. In the freshwater treatment, wood addition reduced activity of BG and NAGase compared to wood-free incubations (Figure 6A-B), but enhanced PER activity (Figure 6C). Wood addition also reduced AS and P activity across all treatments compared to wood-free incubations (Figure 6D-E).

4 Discussion

As forests within the lower coastal plain physiographic region of the southeastern US continue to experience increasing stresses from SLR, changes in microbial C cycling processes should be expected. Our results, combined with other field and lab experiments, confirm that saltwater intrusion into coastal freshwater forested wetlands can result in reductions in CO₂ and CH₄ production (Ardón et al., 2016; Ardón et al., 2018), but this may be balanced by long- and short-term effects of saltwater intrusion on these C cycling processes (Weston et al., 2011), as well as changes in C inputs due to forest-to-marsh transition. Further, wood additions to these wetland soils may reduce CH₄ production under freshwater conditions compared to the absence wood additions (Figure 3C and 3F), but slightly enhance CH₄ production under saltwater conditions. Our results also demonstrate that substantial quantities of CH₄ can be produced from freshwater wetland soils with redox potential between -100 to 100 mV, which may be related to the specific pathway of CH₄ production (acetoclastic versus hydrogenotrophic) (Angle et al., 2016), and challenges the widespread assumption that methanogenesis only occurs at very low redox potentials. Changes in the water table depth at the ARNWR is driven primarily by precipitation patterns (Minick et al., 2019a), resulting in the influx of oxygenated waters. Periodic *in situ* measurements of redox potential at the ARNWR indicate that standing water is relatively aerated (Eh = 175 - 260 mV), while surface soils of hummocks when not submerged are more aerated (Eh = 320 mV) than submerged hollow surface soils (Eh = 100 to 150 mV) and deeper organic soils (20 – 40 cm depth; Eh = 50 to 90 mV) (unpublished data). Furthermore, our results indicate that additions of new C to soils as wood may result in short-term reductions in

redox potential as anaerobic processes are enhanced due to the added C substrate and terminal electron acceptors are quickly reduced. As SLR continues to rise over the next century, more persistent saltwater intrusion may occur as rising brackish waters mix with non-tidal freshwater systems having important implications for both above- and below-ground C cycling dynamics. Although our study only looked at these effects in a controlled laboratory experiment, these data provide a baseline understanding of potential changes in C cycling dynamics in these wetlands due to SLR.

Saltwater additions decreased CO₂ production compared to freshwater in the wood-free soils, although post-incubation MBC and extracellular enzyme activity (e.g., BG, NAGase, and AP) were not different between these treatments. This has been found in other pocosin wetland soils on the coast of North Carolina (Ardón et al. 2018). Variable effects of salinity (and/or SO₄²⁻ additions) have been found on soil respiration, with some studies showing an increase (Marton et al., 2012; Weston et al., 2011), a decrease (Lozanovska et al. 2016; Servais et al. 2019), or no change (Baldwin et al., 2006). Krauss et al. (2012) found that permanently flooded saltwater treatments (expected in non-tidal wetlands) in a simulated coastal swamp mesocosm reduced soil respiration, whereas saltwater pulses (expected in tidal wetlands) had a variable effect on soil respiration. Alternatively, CO₂ production was not reduced in the saltwater compared to freshwater treatments in wood-amended soils, while post-incubation MBC was lower in the saltwater compared to freshwater, which suggests a shift in microbial carbon use efficiency.

Methane production was higher in the freshwater compared to saltwater treatments in both wood-amended and wood-free incubations. Numerous others studies have found that saltwater reduces CH₄ fluxes compared to freshwater, both within the field and laboratory. Reduced CH₄ production from saltwater treated soils primarily results from the availability of

more energetically favorable terminal electron acceptors (primarily SO_4^{2-}), which leads to the competitive suppression of methanogenic microbial communities by SO_4^{2-} reducing communities (Bridgham et al., 2013; Chambers et al., 2011; Winfrey and Zeikus, 1977), as methanogens and SO_4^{2-} reducers compete for acetate and electrons (Le Mer and Roger, 2001). Dang et al. (2019) did find partial recovery over time of the methanogenic community following saltwater inundation to freshwater soil cores, but interestingly this community resembled that of microbes performing hydrogenotrophic methanogenesis and not acetoclastic methanogenesis. Activity of arylsulfatase was also lower in saltwater amended soils. This also indicates a functional change in the microbial community, as microbes in the saltwater treatment are utilizing the readily available SO_4^{2-} pool, while microbes in the freshwater and dry treatments are still actively producing SO_4^{2-} -liberating enzymes to support their metabolic activities. Findings by Baldwin et al. (2006) support the effects of saltwater on changing the microbial community structure as well, in which reductions in CH_4 production in NaCl treated freshwater sediments were accompanied by a reduction in archaeal (methanogens) microbial population, establishing a link between shifting microbial populations and changing CH_4 flux rates due to saltwater intrusion.

Changes in the CH_4 production due to saltwater additions appear to be related to the dominant CH_4 producing pathway. The $^{13}\text{CH}_4$ isotopic signature in wood-free freshwater incubated soils indicated that acetoclastic methanogenesis was the dominant CH_4 producing pathway, while hydrogenotrophic methanogenesis dominated in the saltwater treatments. Acetoclastic methanogenesis produces isotopically enriched CH_4 compared to that of the hydrogenotrophic methanogenesis (Chasar et al., 2000; Conrad et al. 2010; Krohn et al. 2017; Sugimoto and Wada, 1993; Whiticar et al., 1986; Whiticar 1999). The differences in C discrimination between the two pathways is greater for the hydrogenotrophic compared to the

acetoclastic pathway, resulting in more depleted (-110 to -60 ‰) and more enriched (-60 ‰ to -50 ‰) $^{13}\text{CH}_4$, respectively. This has been confirmed in field and laboratory experiments (Conrad et al. 2010; Krohn et al. 2017; Krzycki et al., 1987; Sugimoto and Wada, 1993; Whiticar et al., 1986; Whiticar, 1999). Baldwin et al. (2006) also found that saltwater additions promoted the hydrogenotrophic methanogenic pathway. Further, recent studies have found that saltwater additions to soils result in a shift in the relative abundance of hydrogenotrophic methanogens (Chambers et al. 2011; Dang et al 2019), supporting the idea that saltwater may alter not only the production of CH_4 but also the pathway of methane production.

Changes in freshwater and saltwater hydrology due to rising seas is leading to dramatic shifts in the dominant plant communities within the ARNWR and across the southeastern US (Connor et al., 1997; DOD, 2010; Langston et al., 2017; Kirwan and Gedan 2019). This has the potential to alter the soil C balance due to introduction of large amounts of coarse woody debris as trees die. In our laboratory experiment, additions of wood resulted in changes in both CO_2 and CH_4 production, but the direction of change depended on if soils were incubated with freshwater or saltwater. Wood additions increased CO_2 production compared to wood-free soils, except in the freshwater treatment. This was particularly evident in the dry treatment where wood additions increased CO_2 production by approximately 32 %. For the dry treatment, wood-amended soils had the highest MBC and NAGase activity as microbes were likely immobilizing more N to support metabolic activities in the presence of added C (Fisk et al., 2015; Minick et al., 2017). Higher respiration with wood additions in the saltwater treatments likely resulted from enhanced metabolic activity of SO_4^{2-} reducing microbes in the presence of an added C source. On the other hand, wood additions resulted in a decline in CH_4 production from the freshwater treatment, while slightly enhancing CH_4 production from the saltwater treatments. Wood

additions also resulted in much lower redox potential, particularly in the saltwater treatments, and coupled with $^{13}\text{CH}_4$ stable isotope composition may have driven the higher levels of CH_4 production (via hydrogenotrophic methanogenesis) in the wood plus saltwater treatments. The suppression of CH_4 production by wood additions in the freshwater treatment was somewhat surprising given the positive effects of C additions on CH_4 production recently found in freshwater sediments (West et al. 2012), but likely resulted from enhancement of other, more energetically favorable redox reactions with the addition of a C source (e.g., wood). Furthermore, wood additions to freshwater incubations resulted in a decrease in MBC and activity of BG and NAGase enzymes compared to wood-free incubations and an increase in PER activity. This suggests that the microbial communities have altered their functional capacity in response to wood additions when exposed to freshwater. The $\text{CO}_2\text{:CH}_4$ ratio further indicated that, in freshwater, CH_4 production was quite high in relation to CO_2 production. This ratio was significantly higher for saltwater treatments as CH_4 production dropped drastically compared to freshwater. In wood-free incubations, the $\text{CO}_2\text{:CH}_4$ trend between freshwater and saltwater treatments was parabolic, but was linear upward in wood-amended soils. This suggests that interactions between saltwater concentration and coarse woody debris (in the form of dead and dying trees; Kirwan and Gedan 2019) may be important to understand when determining effects of saltwater intrusion on greenhouse gas production in freshwater forested wetlands.

Findings from this study indicate that substantial changes in the greenhouse gas production and microbial activity are possible due to saltwater intrusion into freshwater wetland ecosystems but that the availability of C in the form of dead wood (as forests transition to marsh) may alter the magnitude of this effect. At ARNWR and similar coastal freshwater forested wetlands, saltwater intrusion may reduce both CO_2 and CH_4 emissions from soils to the

atmosphere. Sea-level rise will likely lead to dramatic and visually striking changes in vegetation, particularly transitioning forested wetlands into shrub or marsh wetlands (Kirwan and Gedan 2019), which has resulted in the widespread occurrence of “ghost” forests along the Atlantic coast (Kirwan and Gedan 2019). As forested wetlands are lost, dead trees could provide a significant source of C to already C-rich peat soils, with the potential to alter CO₂ and CH₄ production. The long-term effect of forest-to-marsh transition on ecosystem C storage will likely depend on the balance between dead wood inputs and effects of SLR and vegetation change on future C inputs and soil microbial C cycling processes. Future work should include investigation of these C cycling and microbial processes at the field-scale and expand to a wider range of non-tidal wetlands within the southeastern US region.

Author contribution

All authors contributed to the conception and design of the study. KM wrote the first draft of the manuscript. KM collected the samples from the field and performed laboratory analysis. All authors contributed to manuscript revision and approved the submitted version.

Competing Interest

The authors declare that they have no conflict of interest.

Acknowledgements

We thank numerous undergraduate researchers for their invaluable help collecting samples from the field and analyzing samples in the laboratory. We also thank two reviewers for their comments, which significantly improved the manuscript. Primary support was provided by USDA NIFA (Multi-agency A.5 Carbon Cycle Science Program) award 2014-67003-22068. Additional support was provided by DOE NICCR award 08-SC-NICCR-1072, the USDA Forest Service Eastern Forest Environmental Threat Assessment Center award 13-JV-11330110-081, and Carolinas Integrated Sciences and Assessments award 2013-0190/13-2322. The USFWS Alligator River National Wildlife Refuge provided helpful scientific discussions, the forested wetland research site, and valuable in-kind support.

References

- Allen, T., Wang, Y., Gore, B., Swords, J., and Newcomb, D.: Coastal Wetland mapping using time series SAR imagery and LiDAR: Alligator River National Wildlife Refuge, North Carolina, in: Proceedings Pecora 18 Symposium, Herndon, Virginia, November 14-17, 2011.
- Angle, J. C., Morin, T. H., Solden, L. M., Narrowe, A. B., Smith, G. J., Borton, M. A., Reyes-Sanchez, C., Daly, R. A., Mirfenderesgi, G., and Hoyt, D. W.: Methanogenesis in oxygenated soils is a substantial fraction of wetland methane emissions, *Nature communications*, 8, 1567, doi: 10.1038/s41467-017-01753-4, 2017.
- Ardón, M., Helton, A. M., and Bernhardt, E. S.: Drought and saltwater incursion synergistically reduce dissolved organic carbon export from coastal freshwater wetlands, *Biogeochemistry*, 127, 411-426, doi: 10.1007/s10533-016-0189-5, 2016.

665 Ardón, M., Helton, A. M., and Bernhardt, E. S.: Salinity effects on greenhouse gas emissions
666 from wetland soils are contingent upon hydrologic setting: a microcosm experiment,
667 Biogeochemistry, 1-16, <https://doi.org/10.1007/s10533-018-0486-2>, 2018.

668 Baldwin, D. S., Rees, G. N., Mitchell, A. M., Watson, G., and Williams, J.: The short-term
669 effects of salinization on anaerobic nutrient cycling and microbial community structure in
670 sediment from a freshwater wetland, Wetlands, 26, 455-464,
671 [https://doi.org/10.1672/0277-5212\(2006\)26\[455:TSEOSO\]2.0.CO;2](https://doi.org/10.1672/0277-5212(2006)26[455:TSEOSO]2.0.CO;2), 2006.

672 Batjes, N. H.: Total carbon and nitrogen in the soils of the world, Eur. J. Soil Sci., 47, 151-163,
673 https://doi.org/10.1111/ejss.12114_2, 1996.

674 Bridgham, S. D., Megonigal, J. P., Keller, J. K., Bliss, N. B., and Trettin, C.: The carbon balance
675 of North American wetlands, Wetlands, 26, 889-916, [https://doi.org/10.1672/0277-5212\(2006\)26\[889:TCBONA\]2.0.CO;2](https://doi.org/10.1672/0277-5212(2006)26[889:TCBONA]2.0.CO;2), 2006.

677 Bridgham, S. D., Cadillo-Quiroz, H., Keller, J. K. and Zhuang, Q.: Methane emissions from
678 wetlands: biogeochemical, microbial, and modeling perspectives from local to global
679 scales, Global Change Biol., 19, 1325-1346, <https://doi.org/10.1111/gcb.12131>, 2013.

680 Chambers, L. G., Reddy, K. R., and Osborne, T. Z.: Short-term response of carbon cycling to
681 salinity pulses in a freshwater wetland, Soil Sci. Soc. Am. J., 75, 2000-2007,
682 doi:10.2136/sssaj2011.0026, 2011.

683 Chambers, L. G., Guevara, R., Boyer, J. N., Troxler, T. G. and Davis, S. E.: Effects of salinity
684 and inundation on microbial community structure and function in a mangrove peat soil,
685 Wetlands, 36, 361-371, <https://doi.org/10.1007/s13157-016-0745-8>, 2016.

686 Chasar, L., Chanton, J., Glaser, P., and Siegel, D.: Methane concentration and stable isotope
687 distribution as evidence of rhizospheric processes: Comparison of a fen and bog in the

688 Glacial Lake Agassiz Peatland complex, *Annals of Botany*, 86, 655-663,
689 <https://doi.org/10.1006/anbo.2000.1172>, 2000.

690 Conner, W., McLeod, K. and McCarron, J.: Flooding and salinity effects on growth and survival
691 of four common forested wetland species, *Wetlands Ecol. Manage.*, 5, 99-109,
692 <https://doi.org/10.1023/A:1008251127131>, 1997.

693 Conrad, R., Klose, M., Claus, P., and Enrich-Prast, A.: Methanogenic pathway, ^{13}C isotope
694 fractionation, and archaeal community composition in the sediment of two clear-water
695 lakes of Amazonia, *Limnol. Oceanogr.*, 55, 689-
696 702, <https://doi.org/10.4319/lo.2010.55.2.0689>, 2010.

697 Craft, C., Clough, J., Ehman, J., Guo, H., Joye, S., Machmuller, M., Park, R., and Pennings, S.:
698 Effects of accelerated sea level rise on delivery of ecosystem services provided by tidal
699 marshes: a simulation of the Georgia (USA) Coast, *Frontiers in Ecology and the*
700 *Environment*, 7, 73, 2009.

701 Department of Defense (DOD): Responding to climate change, *Natural Selections*, 6, 2-4, 2010.

702 Feng, X., Xu, Y., Jaffé, R., Schlesinger, W. H., and Simpson, M. J.: Turnover rates of
703 hydrolysable aliphatic lipids in Duke Forest soils determined by compound specific ^{13}C
704 isotopic analysis, *Org. Geochem.*, 41, 573-579,
705 <https://doi.org/10.1016/j.orggeochem.2010.02.013>, 2010.

706 Fisk, M., Santangelo, S., and Minick, K.: Carbon mineralization is promoted by phosphorus and
707 reduced by nitrogen addition in the organic horizon of northern hardwood forests, *Soil*
708 *Biol. Biochem.*, 81, 212-218, <https://doi.org/10.1016/j.soilbio.2014.11.022>, 2015.

709 Fry, B. 2006. *Stable Isotope Ecology*. Springer, New York, NY.

710 Joergensen, R. G.: The fumigation-extraction method to estimate soil microbial biomass:
 711 calibration of the kEC value. *Soil Biol. Biochem.*, 28, 25-31,
 712 [https://doi.org/10.1016/0038-0717\(95\)00102-6](https://doi.org/10.1016/0038-0717(95)00102-6), 1996.

713 Johnson, M.G., and Kern, J.S.: Quantifying the organic carbon held in forested soils of the
 714 United States and Puerto Rico. Chapter 4, Kimble, JS (ed.), *The Potential of U.S. Forest*
 715 *Soils to Sequester Carbon and Mitigate the Greenhouse Effect*. CRC Press LLC, Boca
 716 Raton, FL, 2003.

717 Karegar, M. A., Dixon, T. H., Malservisi, R., Kusche, J., and Engelhart, S. E.: Nuisance flooding
 718 and relative sea-level rise: the importance of present-day land motion, *Scientific reports*,
 719 7, 11197, doi: 10.1038/s41598-017-11544-y, 2017.

720 Kim, D., Oren, R., and Qian, S. S.: Response to CO₂ enrichment of understory vegetation in the
 721 shade of forests, *Global Change Biol.*, 22, 944-956, <https://doi.org/10.1111/gcb.13126>,
 722 2016.

723 Kirwan, M.L., and Gedan, K.B.: Sea-level driven land conversion and the formation of ghost
 724 forests, *Nature Climate Change*, 9, 450-457, <https://doi.org/10.1038/s41558-019-0488-7>
 725 2019.

726 Krauss, K. W., Whitbeck, J. L., and Howard, R. J.: On the relative roles of hydrology, salinity,
 727 temperature, and root productivity in controlling soil respiration from coastal swamps
 728 (freshwater), *Plant Soil*, 358, 265-274, <https://doi.org/10.1007/s11104-012-1182-y>,
 729 2012.

730 Krohn, J., Lozanovska, I., Kuzyakov, Y., Parvin, S., Dorodnikov, M.: CH₄ and CO₂ production
 731 below two contrasting peatland micro-relief forms: An inhibitor and $\delta^{13}\text{C}$ study. *Science*

732 of The Total Environment, 586, 142-151, <https://doi.org/10.1016/j.scitotenv.2017.01.192>,
733 2017.

734 Krzycki, J. A., Kenealy, W. R., Deniro, M. J., and Zeikus, J. G.: Stable carbon isotope
735 fractionation by *Methanosarcina barkeri* during methanogenesis from acetate, methanol,
736 or carbon dioxide-hydrogen, Appl. Environ. Microbiol., 53, 2597-2599, 1987.

737 Langston, A. K., Kaplan, D. A., and Putz, F. E.: A casualty of climate change? Loss of
738 freshwater forest islands on Florida's Gulf Coast, Global Change Biol., 23, 5383-5397,
739 <https://doi.org/10.1111/gcb.13805>, 2017.

740 Le Mer, J., and Roger, P.: Production, oxidation, emission and consumption of methane by soils:
741 a review, Eur. J. Soil Biol., 37, 25-50, [https://doi.org/10.1016/S1164-5563\(01\)01067-6](https://doi.org/10.1016/S1164-5563(01)01067-6),
742 2001.

743 Lee, J. K., Park, R. A., and Mausel, P. W.: Application of geoprocessing and simulation
744 modeling to estimate impacts of sea level rise on the northeast coast of Florida,
745 Photogrammetric Engineering and Remote Sensing;(United States), 58, 1992.

746 Lozanovska, I., Kuzyakov, Y., Krohn, J., Parvin, S., and Dorodnikov, M.: Effects of nitrate and
747 sulfate on greenhouse gas emission potentials from microform-derived peats of a boreal
748 peatland: A ^{13}C tracer study, Soil Biol. Biochem., 100, 182-191,
749 <https://doi.org/10.1016/j.soilbio.2016.06.018>, 2016.

750 Marton, J. M., Herbert, E. R., and Craft, C. B.: Effects of salinity on denitrification and
751 greenhouse gas production from laboratory-incubated tidal forest soils, Wetlands, 32,
752 347-357, <https://doi.org/10.1007/s13157-012-0270-3>, 2012.

753 Miao, G., Noormets, A., Domec, J., Trettin, C.C., McNulty, S.G., Sun, G., and King, J.S.: The
 754 effect of water table fluctuation on soil respiration in a lower coastal plain forested wetland
 755 in the southeastern US, *Biogeosciences* 118, 1748-1762, doi:10.1002/2013JG002354, 2013.

756 Miao G, Noormets A, Domec J-C, Fuentes M, Trettin CC, Sun G, McNulty SG, King JS:
 757 Hydrology and microtopography control carbon dynamics in wetlands: implications in
 758 partitioning ecosystem respiration in a coastal plain forested wetland, *Agricultural and*
 759 *Forest Meteorology*, 247, 343-355, <https://doi.org/10.1016/j.agrformet.2017.08.022>,
 760 2017.

761 Mitra, B., Miao, G., Minick K.J., McNulty S., Sun G., Gavazzi, M., King J.S., and Noormets A.,
 762 Disentangling the effects of temperature, moisture and substrate availability on soil CO₂ efflux.
 763 *Journal of Geophysical Research: Biogeosciences* 124, <https://doi.org/10.1029/2019JG005148>,
 764 2019.

765 Minick, K. J., Kelley, A. M., Miao, G., Li, X., Noormets, A., Mitra, B., and King, J. S.:
 766 Microtopography alters hydrology, phenol oxidase activity and nutrient availability in
 767 organic soils of a coastal freshwater forested wetland, *Wetlands* 39, 263-273,
 768 <https://doi.org/10.1007/s13157-018-1107-5>, 2019a.

769 Minick, K. J., Mitra, B., Li, X., Noormets, A., and King, J. S.: Water table drawdown alters soil
 770 and microbial carbon pool size and isotope composition in coastal freshwater forested
 771 wetlands, *Frontiers in Forests and Global Change*, 2, 1-19,
 772 <https://doi.org/10.3389/ffgc.2019.00007>, 2019b.

773 Morrissey, E. M., Gillespie, J. L., Morina, J. C., and Franklin, R. B.: Salinity affects microbial
 774 activity and soil organic matter content in tidal wetlands, *Global Change Biol.*, 20, 1351-
 775 1362, <https://doi.org/10.1111/gcb.12431>, 2014.

776 Neubauer, S., Franklin, R., and Berrier, D.: Saltwater intrusion into tidal freshwater marshes
 777 alters the biogeochemical processing of organic carbon, *Biogeosciences*, 10, 8171-8183,
 778 <https://doi.org/10.5194/bg-10-8171-2013>, 2013.

779 Paerl, H. W., Crosswell, J. R., Van Dam, B., Hall, N. S., Rossignol, K. L., Osburn, C. L.,
 780 Hounshell, A. G., Sloup, R. S., and Harding, L. W.: Two decades of tropical cyclone
 781 impacts on North Carolina's estuarine carbon, nutrient and phytoplankton dynamics:
 782 implications for biogeochemical cycling and water quality in a stormier world,
 783 *Biogeochemistry*, 141, 307-332, <https://doi.org/10.1007/s10533-018-0438-x>, 2018.

784 Riggs, S. R.: Sediment evolution and habitat function of organic-rich muds within the Albemarle
 785 estuarine system, North Carolina, *Estuaries* 19, 169–185,
 786 <https://doi.org/10.2307/1352223>, 1996.

787 Riggs, S. R., and Ames, D. V.: Drowning the North Carolina coast: Sea-level rise and estuarine
 788 dynamics. North Carolina Sea Grant, Raleigh, NC, 2008.

789 Sallenger, A. H., Doran, K. S., and Howd, P. A.: Hotspot of accelerated sea-level rise on the
 790 Atlantic coast of North America, *Nature Climate Change*, 2, 884, doi:10.1038/nclimate1597,
 791 2012.

792 Schlesinger, W., Bernhardt, E., DeLucia, E., Ellsworth, D., Finzi, A., Hendrey, G., Hofmockel,
 793 K., Lichter, J., Matamala, R. and Moore, D.: The Duke Forest FACE experiment: CO₂
 794 enrichment of a loblolly pine forest, in: *Managed Ecosystems and CO₂*, Springer, 197-
 795 212, 2006.

796 Sinsabaugh, R., Antibus, R., Linkins, A., McClaugherty, C., Rayburn, L., Repert, D., and
 797 Weiland, T.: Wood decomposition over a first-order watershed: mass loss as a function of

798 lignocellulase activity, *Soil Biol. Biochem.*, 24, 743-749, <https://doi.org/10.1016/0038->
 799 0717(92)90248-V, 1992.

800 Sinsabaugh, R. L., Antibus, R., Linkins, A., McClaugherty, C., Rayburn, L., Repert, D., and
 801 Weiland, T.: Wood decomposition: nitrogen and phosphorus dynamics in relation to
 802 extracellular enzyme activity, *Ecology*, 74, 1586-1593, <https://doi.org/10.2307/1940086>,
 803 1993.

804 Sugimoto, A., and Wada, E.: Carbon isotopic composition of bacterial methane in a soil
 805 incubation experiment: Contributions of acetate and CO₂H₂, *Geochim. Cosmochim. Acta*,
 806 57, 4015-4027, [https://doi.org/10.1016/0016-7037\(93\)90350-6](https://doi.org/10.1016/0016-7037(93)90350-6), 1993.

807 Titus, J. G., and Richman, C.: Maps of lands vulnerable to sea level rise: modeled elevations along
 808 the US Atlantic and Gulf coasts, *Climate research*, 18, 205-228, doi:10.3354/cr01, 2001.

809 Vance, E. D., Brookes, P. C. and Jenkinson, D. S.: An extraction method for measuring soil
 810 microbial biomass C, *Soil Biol. Biochem.*, 19, 703-707, <https://doi.org/10.1016/0038->
 811 0717(87)90052-6, 1987.

812 West, W. E., Coloso, J. J., and Jones, S. E.: Effects of algal and terrestrial carbon on methane
 813 production rates and methanogen community structure in a temperate lake sediment,
 814 *Freshwat. Biol.*, 57, 949-955, <https://doi.org/10.1111/j.1365-2427.2012.02755.x>, 2012.

815 Weston, N. B., Vile, M. A., Neubauer, S. C., and Velinsky, D. J.: Accelerated microbial organic
 816 matter mineralization following salt-water intrusion into tidal freshwater marsh soils,
 817 *Biogeochemistry*, 102, 135-151, <https://doi.org/10.1007/s10533-010-9427-4>, 2011.

818 Whiticar, M. J., Faber, E., and Schoell, M.: Biogenic methane formation in marine and
 819 freshwater environments: CO₂ reduction vs. acetate fermentation—isotope evidence,

Geochim. Cosmochim. Acta, 50, 693-709, [https://doi.org/10.1016/0016-7037\(86\)90346-7](https://doi.org/10.1016/0016-7037(86)90346-7), 1986.

Whiticar, M. J.: Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane, Chem. Geol., 161, 291-314, [https://doi.org/10.1016/S0009-2541\(99\)00092-3](https://doi.org/10.1016/S0009-2541(99)00092-3), 1999.

Winfrey, M. R., and Zeikus, J. G.: Effect of sulfate on carbon and electron flow during microbial methanogenesis in freshwater sediments, Appl. Environ. Microbiol., 33, 275-281, 1977.

835 **Tables and Figures**

836

837 Table 1. Total organic C (TOC) and ion concentrations (mg L⁻¹) in freshwater (0 ppt), 2.5 ppt saltwater, and 5.0 ppt saltwater.

838 Standard errors of the mean are in parenthesis (n=4). Values with different superscript lowercase letters are significantly different (*P* <
839 0.05).

840

Treatment	TOC	SO ₄ ²⁻	Cl ⁻	Na ⁺	NH ₄ ⁺	NO ₃ ⁻	PO ₄ ³⁻	Ca ²⁺	Mg ²⁺	K ⁺
0 ppt	44 (0.3) ^a	1 (0.1) ^a	17 (0.2) ^a	8 (0.1) ^a	0.00 (0.000) ^a	0.00 (0.000) ^a	0.00 (0.000) ^a	1 (0.0) ^a	1 (0.0) ^a	0.2 (0.0) ^a
2.5 ppt	40 (0.7) ^b	162 (1.3) ^b	1391 (42.8) ^b	538 (19.2) ^b	0.06 (0.004) ^b	0.06 (0.000) ^a	0.01 (0.000) ^a	23 (0.3) ^b	64 (2.6) ^b	19 (0.3) ^b
5.0 ppt	38 (0.1) ^b	319 (6.5) ^c	2695 (22.6) ^c	1039 (15.9) ^c	0.07 (0.004) ^b	0.07 (0.004) ^a	0.01 (0.000) ^b	44 (1.0) ^c	125 (2.1) ^c	36 (0.4) ^c

841

842

843

844

845

846

847

848

849

850

851

852 Table 2. Post-incubation soil organic C (SOC) concentration (g kg^{-1}), SOC $\delta^{13}\text{C}$ (‰), and wood-derived SOC (%) (estimated from ^{13}C
853 two pool mixing model) for soil samples collected from the field and incubated for 98 d in the laboratory under dry conditions (Dry)
854 or fully saturated with freshwater (0 ppt) or saltwater (2.5 and 5.0 ppt) and with (+ Wood) or without addition of ^{13}C -depleted wood.
855 Standard errors of the mean are in parenthesis (n=4). Data from wood-free and wood-amended soils were analyzed separately. Values
856 followed by different superscript lowercase letters are significantly different between the four treatments of the wood-free or wood-
857 amended soils ($P < 0.05$).

858

Treatment	Post-SOC Concentration (g kg^{-1})	Post-SOC $\delta^{13}\text{C}$ (‰)	Wood-derived SOC (%)
Dry	495 (1.5) ^b	-29.5 (0.20) ^a	.
0 ppt	493 (3.3) ^b	-29.5 (0.18) ^a	.
2.5 ppt	488 (4.9) ^b	-29.5 (0.20) ^a	.
5.0 ppt	460 (8.6) ^a	-29.5 (0.16) ^a	.
Dry + Wood	491 (4.7) ^{ab}	-30.4 (0.30) ^a	8 (2.5)
0 ppt + Wood	502 (4.6) ^a	-30.7 (0.22) ^a	12 (0.4)
2.5 ppt + Wood	477 (4.9) ^{bc}	-30.6 (0.35) ^a	10 (1.4)
5.0 ppt + Wood	470 (4.6) ^c	-30.4 (0.14) ^a	10 (2.0)

859

860

861

862

863

864

865

866

867

868

869

870

871 Table 3. Results (F-values and significance) from the repeated measures ANOVA of pH, Eh, microbial biomass C (MBC), $\delta^{13}\text{C}$
872 isotopic signature of MBC, $\delta^{13}\text{CO}_2$, and $\delta^{13}\text{CH}_4$ measured in soils collected from a coastal freshwater forested wetland and incubated
873 in the laboratory for 98 d under fully saturated conditions with either freshwater or saltwater (2.5 ppt and 5.0 ppt). Data from wood-
874 free and wood-amended soils were analyzed separately.

875

Source	pH	Eh	MBC	MBC ^{13}C	$\delta^{13}\text{CO}_2$	$\delta^{13}\text{CH}_4$
Wood-Free						
Treatment	26.6***	4.5*	3.7*	3.2*	351.7***	60.5***
Time	4.4***	40.7***	40.9***	15.8**	24.2***	8.3***
Treatment x Treatment	1.22	3.7***	27.3***	3.3*	6.4***	1.1
Wood-Amended						
Treatment	29.0***	13.6***	39.9***	2.6	129.8***	0.3
Time	18.3***	30.1***	111.0***	3.7	34.8***	1.4
Treatment x Treatment	1.4	3.4***	24.2***	5.5**	8.3***	1.0

876 *P < 0.05, **P < 0.01, ***P < 0.0001

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891 Table 4. Results (F-values and significance) from the one-way ANOVA of cumulative gas production and extracellular enzyme
892 activity (BG: β -glucosidase; PER: peroxidase; NAGase: glucosaminidase; AP: alkaline phosphatase; and AS: arylsulfatase) from soils
893 collected from a coastal freshwater forested wetland and incubated in the laboratory for 98 d under dry conditions or fully saturated
894 conditions with either freshwater or saltwater (2.5 ppt and 5.0 ppt). Data from wood-free and wood-amended soils were analyzed
895 separately.

896

Source	CO ₂	CH ₄	BG	PER	NAGase	AP	AS
Wood-Free							
Treatment	20.4***	15.6***	7.2**	11.9**	9.5**	0.9	15.8**
Wood-Amended							
Treatment	13.3**	36.7***	16.6**	2.5	32.0***	2.3	31.2***

897 *P < 0.05, **P < 0.01, ***P < 0.0001

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915 Table 5. Initial (1 d) and final (98 d) microbial biomass C (MBC) (mg kg⁻¹), MBC δ¹³C (‰), wood-derived MBC (%) (estimated
 916 using ¹³C two pool mixing model), and cumulative extracellular enzyme activity (μmol g⁻¹) (BG: β-glucosidase; PER: peroxidase;
 917 NAGase: glucosaminidase; AP: alkaline phosphatase; and AS: arylsulfatase) for soils incubated under dry conditions (Dry) or
 918 saturated conditions with freshwater (0 ppt) or saltwater (2.5 and 5.0 ppt) and with (+ Wood) or without addition of ¹³C-depleted
 919 wood. Standard errors of the mean are in parenthesis (n=4). Values followed by different superscript lowercase letters are
 920 significantly different between the four treatments for the wood-free or wood-amended soils (*P* < 0.05).
 921

Treatment	Initial MBC Concentration (mg kg ⁻¹)	Final MBC Concentration (mg kg ⁻¹)	Initial MBC δ ¹³ C (‰)	Final MBC δ ¹³ C (‰)	Wood- derived MBC (%)	BG	PER	NAGase	AP	AS
Dry	2238 (400) ^c	4077 (387) ^a	-27.0 (0.43) ^a	-28.4 (0.28) ^{ab}	.	547 (37) ^a	176 (14) ^a	240 (20) ^a	7599 (1038) ^a	47 (2) ^a
0 ppt	3982 (196) ^{ab}	2657 (344) ^b	-27.3 (0.19) ^a	-28.9 (0.16) ^a	.	479 (18) ^{ab}	197 (38) ^a	194 (11) ^{ab}	6308 (517) ^a	47 (8) ^a
2.5 ppt	7334 (1177) ^a	2495 (195) ^b	-27.8 (0.51) ^a	-27.9 (0.03) ^{ab}	.	389 (33) ^b	412 (75) ^b	159 (9) ^b	6539 (183) ^a	19 (3) ^b
5.0 ppt	6483 (104) ^{ab}	2114 (135) ^b	-27.0 (0.30) ^a	-27.4 (0.15) ^b	.	379 (27) ^b	490 (30) ^b	154 (8) ^b	6387 (529) ^a	15 (2) ^b
Dry + Wood	4444 (579) ^a	5174 (249) ^a	-29.3 (0.40) ^a	-32.1 (0.44) ^a	31 (4.9) ^a	554 (37) ^a	243 (22) ^a	275 (17) ^a	7247 (887) ^a	40 (2) ^a
0 ppt + Wood	5376 (330) ^a	1832 (102) ^b	-29.8 (0.37) ^a	-29.4 (0.15) ^b	4 (1.1) ^b	349 (24) ^b	275 (44) ^a	153 (11) ^b	4965 (459) ^a	36 (3) ^a
2.5 ppt + Wood	5173 (405) ^a	748 (124) ^c	-30.1 (0.25) ^a	-30.4 (0.95) ^{ab}	21 (7.8) ^a	368 (12) ^b	365 (30) ^a	150 (6) ^b	5548 (653) ^a	14 (3) ^b
5.0 ppt + Wood	2123 (400) ^b	790 (87) ^c	-29.9 (0.43) ^a	-29.7 (0.37) ^b	18 (1.9) ^{ab}	369 (13) ^b	326 (38) ^a	150 (6) ^b	5893 (495) ^a	13 (2) ^b

922

Figure 1. Location of the Alligator River National Wildlife Refuge (ARNWR) within North America (left panel, indicated by box) and location of ARNWR within eastern North Carolina, USA and surrounding freshwater (Alligator River and Albermarle Sound) and saltwater (Pamlico Sound, Croatan Sound, and Roanoke Sound) bodies (right panel). Dotted arrows indicate the location of important surrounding water bodies. The star represents the approximate location of soil and freshwater (from Milltail Creek) sampling locations within the freshwater forested wetlands of ARNWR. The black circle represents the approximate location of saltwater sampling (at the Melvin Daniels Bridge, Roanoke Sound) from the Roanoke Sound. The saltwater was sampled approximately 20 miles east of the soil and freshwater samples.




Figure 2. pH for wood-free soils (A) and wood-amended soils (B) and redox potential for wood-free soils (C) and wood-amended soils (D) measured over the course of the 98 d laboratory incubation. Symbols represent mean with standard error (n=4). An asterisk ($P < 0.05$) indicates significant differences between treatment means at each time point.




Figure 3. Cumulative CO₂ production from wood-free soils (A), wood-amended soils (B), and the wood-associated CO₂ production (C); and cumulative CH₄ production for wood-free soils (D), wood amended soils (E), and the wood-associated CH₄ production (F). Panels C and F refer to the difference between wood-amended and wood-free soils. Bars represent mean with standard error (n=4). Bars with different uppercase letters are significantly different ($P < 0.05$).




Figure 4. The $\delta^{13}\text{CO}_2$ values measured over the course of the 98 d laboratory incubation for wood-free soils (A), wood-amended soils (B), and the proportion of wood-derived CO_2 (C). Bars represent mean with standard error (n=4). An asterisk ($P < 0.05$) indicates significant differences between treatment means at each time point.




Figure 5. The $\delta^{13}\text{CH}_4$ values measured over the course of the 98 d laboratory incubation for wood-free soils (A) and wood-amended soils (B) and the average $\delta^{13}\text{CH}_4$ across the entire incubation for wood-free soils (C) and wood-amended soils (D). Symbols or bars represent mean with standard error (n=4). Treatment means with different lowercase letters are significantly different within a sampling time point ($P < 0.05$).




Figure 6. Wood-associated (wood-amended – wood-free) enzyme activity (BG: β -glucosidase; PER: peroxidase; NAGase: glucosaminidase; AP: alkaline phosphatase; and AS: arylsulfatase). Bars represent mean with standard error (n=4). Treatment means with different upper letters are significantly different ($P < 0.05$).

