

Evidence for microbial iron reduction in the methanic sediments of the oligotrophic SE Mediterranean continental shelf

Hanni Vigderovich¹, Lewen Liang², Barak Herut³, Fengping Wang², Eyal Wurgaft^{1,4}, Maxim Rubin-Blum³ and Orit Sivan¹

¹The Department of Geological and Environmental Sciences, Ben-Gurion University of the Negev, Beer-Sheva, 8410501, Israel.

²School of Life Sciences and Biotechnology, Shanghai JiaoTong University, Shanghai, 200240, P.R.China.

³Israel Oceanographic and Limnological Research, Haifa, 31080, Israel.

⁴Currently: The Department of Marine Chemistry and Biochemistry, Woods-Hole Oceanographic Institution, Woods-Hole, USA

Correspondence to: Orit Sivan (oritsi@bgu.ac.il)

Abstract. Dissimilatory iron reduction is probably one of the oldest types of metabolisms that still participates in important biogeochemical cycles, such as the carbon and sulfur cycles. It is one of the more energetically favorable anaerobic microbial respiration processes and is usually coupled to the oxidation of organic matter. Traditionally this process is thought to be limited to the shallow part of the sedimentary column in most aquatic systems. However, iron reduction has also been observed in the methanic zone of many marine and freshwater sediments, well below its expected zone, occasionally accompanied by decreases in methane, suggesting a link between the iron and the methane cycles. Nevertheless, the mechanistic nature of this link (competition, redox or other) has yet to be established, and has not been studied in oligotrophic shallow marine sediments. In this study we present combined geochemical and molecular evidences for microbial iron reduction in the methanic zone of the oligotrophic Southern Eastern (SE) Mediterranean continental shelf. Geochemical pore-water profiles indicate iron reduction in two zones, the uppermost part of the sediment, and the deeper zone, located in the enhanced methane concentration layer. Results from a slurry incubation experiment indicate that the deep methanic iron reduction is microbially mediated. The sedimentary profiles of microbial abundance and qPCR of the *mcrA* gene, together with Spearman correlation between the microbial data and Fe(II) concentrations in the pore-water, suggest types of potential microorganisms that may be involved in the iron reduction via several potential pathways: H₂ or organic matter oxidation, an active sulfur cycle or iron driven anaerobic oxidation of methane. We suggest that intensive upward migration of methane in the sedimentary column and its oxidation by sulfate may fuel the microbial activity in the sulfate methane transition zone (SMTZ). The biomass, created by this microbial activity, can be used below by the iron reducers in the methanic zone of the sediments of the SE Mediterranean.



1 Introduction

Iron (Fe) is the fourth most abundant element in the Earth's crust. It appears as elemental Fe, Fe(II) and Fe(III), and has an important geobiological role in natural systems (e.g. Roden, 2006). Dissimilatory microbial iron reduction may be one of the first evolutionary metabolisms, and plays a key role in the reductive dissolution of Fe(III) minerals in the natural environment (Lovley and Phillips, 1986; Lovley


41 et al., 1987; Lovley and Phillips, 1988; Lovley, 1997; Weber et al., 2006) and in the mineralization of
42 organic matter in freshwater sediments (Roden and Wetzel, 2002). It also serves as a redox wheel that
43 drives the biogeochemical cycles of carbon, nitrogen, sulfur and phosphorous (Li et al., 2012 ; Slomp et
44 al., 2013; Sivan et al., 2014; Egger et al., 2016; Ettwig et al., 2016; Riedinger et al., 2017; März et al.,
45 2018).

46 Dissimilatory iron reduction is part of the anaerobic respiration cascade, in which different organic
47 substrates are used for energy by microorganisms and oxidized to dissolved inorganic carbon (DIC). This
48 is accomplished by reduction of electron acceptors, other than oxygen, according to their availability and
49 energy yield. Denitrification is the first respiratory process in anoxic sediments, followed by manganese
50 reduction, iron reduction and then sulfate reduction. Methane (CH₄) production (methanogenesis) by
51 archaeal methanogens is traditionally considered to be the terminal process of microbial organic matter
52 mineralization in anoxic environments, after the other electron acceptors have been exhausted (Froelich
53 et al., 1979). When the produced methane diffuses away from the methanic layer and meets an electron
54 acceptor it can be consumed by microbial oxidation (methanotrophy). In marine sediments anaerobic
55 oxidation of methane (AOM) coupled to sulfate reduction has been shown to occur (Iversen and
56 Jørgensen, 1985; Hoehler et al., 1994; Hinrichs et al., 1999; Boetius et al., 2000; Orphan et al., 2001;
57 Knittel and Boetius, 2009), and was found to consume up to 90 % of the methane that diffuses upwards
58 to the sulfate methane transition zone (SMTZ) (e.g. Neiwöhner et al., 1998; Valentine, 2002).

59 The classical process of dissimilatory iron reduction is coupled to the oxidation of organic matter
60 (organoclastic iron reduction) (Eq. 1, Lovley, 1991; Lovley et al., 1996). However, iron reduction can be
61 coupled to other processes as well, such as hydrogen (H₂) oxidation (hydrogenotrophic iron reduction)
62 (Eq. 1, Lovley, 1991). Additionally, Fe(III) can be reduced microbially (and also abiotically) by pyrite
63 oxidation (Eq. 2, Bottrell et al., 2000), leading to sulfur (S) intermediates, and followed by their
64 disproportionation to sulfate and sulfide via a "cryptic" sulfur cycle (e.g. Holmkvist et al., 2011).



67 Another recently discovered pathway of iron reduction is by AOM (Eq. 3).



69 This process in marine sediments was shown through incubation experiments with marine seeps
70 sediments (Beal et al., 2009; Sivan et al., 2014). It was also suggested to exist mainly through
71 geochemical profiles in deep sea sediments and their modeling (Sivan et al., 2007; März et al., 2008;
72 Riedinger et al., 2014), and in brackish coastal sediments (Slomp et al., 2013; Segarra et al., 2013; Egger
73 et al., 2014; Egger et al., 2016; Rooze et al., 2016; Egger et al., 2017). In freshwater environments, it
74 was suggested to occur in lakes (Crowe et al. 2011; Sivan et al., 2011; Norði et al., 2013), and in
75 denitrifying cultures from sewage, where it was performed by methanogens (Ettwig et al., 2016). Iron-
76 coupled AOM in natural lake sediments was indicated using isotope pore-water depth profiles (Sivan et
77 al., 2011), rate modeling based on these profiles (Adler et al., 2011), microbial profiles (Bar-Or et al.,

78 2015), and directly from a set of sediment slurry incubation experiments (Bar-Or et al. 2017). The few
79 microbial studies on iron-coupled AOM (mainly in cultures) showed either the involvement of
80 methanogenic/methanotrophic archaea (Scheller et al., 2016; Ettwig et al., 2016; Rotaru and Thamdrup,
81 2016; Cai et al., 2018; Yan et al., 2018) or a cooperation between methanotrophs and methanogens (Bar-
82 Or et al., 2017).

83 Whereas Fe(II) is highly soluble, Fe(III) which is the most abundant species of iron under natural
84 conditions, appears as low-solubility oxidized minerals. This makes iron usage a challenge to
85 microorganisms, which need to respire these iron-oxide minerals, thus rendering many of the iron-oxide
86 minerals effectively unavailable for reduction and leading to the dominance of sulfate reducing bacteria
87 beyond a certain depth. Therefore, it is not trivial to observe iron reduction below the upper iron reduction
88 depth, in the methanic zone, where iron-oxides are assumed to be of low reactivity. Moreover, this type
89 of iron reduction is occasionally accompanied by depletion in methane concentrations, suggesting a
90 possible link between the iron and the methane cycles. There are three potential mechanisms that can
91 link the cycles: 1) a competition between methanogens and iron-reducing bacteria over substrate, 2) a
92 metabolism switch of methanogens from methanogenesis to iron reduction, and 3) iron coupled AOM,
93 as mentioned above. Previous observations in other environments demonstrated the inhibition of
94 methanogenesis under iron-reducing conditions due to competition between methanogens and iron-
95 reducing bacteria for the common acetate and hydrogen substrates (Lovley and Phillips, 1986; Roden
96 and Wetzel, 1996; Conrad, 1999; Roden, 2003). Different methanogens can also utilize iron directly, by
97 reducing Fe(III). This was shown in pure cultures with the amorphous Fe(III) oxyhydroxide (Bond and
98 Lovley., 2002), in pure cultures close to natural sedimentary conditions (Sivan et al., 2016), in natural
99 lake sediments with different iron oxides (i.e. amorphous iron, goethite, hematite and magnetite) (Bar-or
100 et al., 2017), in anoxic ferruginous lake sediment enrichments (Bray et al., 2018), and in iron-rich clays
101 (Liu et al., 2011; Zhang et al., 2012; Zhang et al., 2013).

102 Despite the above studies, the nature of the link between the biogeochemical cycles of iron and methane
103 in the methanic zone of marine sediments, which creates suitable conditions for iron reduction, has not
104 yet been determined. Furthermore, this microbial iron reduction in methanic zones has not been shown
105 in the sediments of oligotrophic shallow marine environments. In this study we report the observation of
106 microbial iron reduction in the methanic depth of marine sediments from the oligotrophic SE
107 Mediterranean continental shelf. The microbial iron reduction is observed by using geochemical pore-
108 water profiles, qPCR profile of archaea, bacteria and the *mcrA* functional gene and 16S rRNA gene
109 sequencing profile at three different stations combined with a simple slurry incubation experiment from
110 the methanic zone. The slurries were amended with hematite and magnetite. Given their low reactivity
111 these are the expected Fe(III) minerals to survive the sulfide zone (Canfield, 1989; Poulton et al., 2004).
112 Furthermore, these minerals were found to be active in iron-coupled AOM in lake sediments (Bar-Or et
113 al., 2017). The profiles, the incubation experiment as well as the related microorganisms, are discussed
114 in terms of the possible links between the cycles of iron and methane.

115 **2 Methods**

116 **2.1 Study site**

117 The surface water in the Levantine Basin of the SE Mediterranean Sea, including Israel's continental
118 shelf, is an oligotrophic nutrient-poor marine system (Herut et al., 2000; Kress and Herut, 2001). The
119 continental shelf narrows from south to north and is composed of Pliocene-Quaternary Nile-derived
120 sediments. The sedimentation rate decreases with increasing distance from the Nile Delta and from the
121 shoreline (Nir, 1984; Sandler and Herut, 2000). Off the shore of Israel, the sedimentation rate is relatively
122 high at $\sim 0.1 \text{ cm y}^{-1}$ (Bareket et al., 2016). The bottom seawater along the continental shelf is well
123 oxygenated and sulfate concentrations at the water-sediment interface are $\sim 30 \text{ mmol L}^{-1}$ (Sela-Adler et
124 al., 2015). The central and eastern regions of the Levantine Basin have relatively low total organic carbon
125 (TOC) content ($\sim 0.1 - 1.4\%$; Almogi-Labin et al., 2009; Sela-Adler et al., 2015; Astrahan et al., 2017)
126 as compared to the Western Mediterranean Basin and offshore the Nile River delta ($1 - 2\%$). Along the
127 Egyptian coast, the TOC in surface sediments on the shelf reach maximum values of 1.5% (Aly Salem
128 et al., 2013). The finding of a free gas zone, which is located from few to tens meters below the seafloor
129 (i.e. gas front), in seismic profiles within the sediments of the continental shelf of Israel (Schattner et al.,
130 2012) led to the discovery of biogenic methane formation at some locations in the shallow sediments
131 (Sela-Adler et al., 2015).

132 **2.2 Sampling**

133 Seven sediment cores ($\sim 5 - 6 \text{ m}$ long) were collected using a Benthos 2175 piston corer, from the
134 undisturbed sediments of the SE Mediterranean continental shelf of Israel at water depths of $81 - 89 \text{ m}$
135 from three stations; SG-1, PC-3 and PC-5 (Fig. 1). The cores were sampled during cruises of R.V.
136 *Shikmona* between 2013 to 2017, and by the R.V. *Bat-Galim* on January 2017 (Table 1). The sediment
137 cores were sliced on board every $25 - 35 \text{ cm}$ within minutes upon retrieval from the seafloor. This area
138 was previously investigated for other purposes, such as the sulfate reduction in the SMTZ (Antler et al.,
139 2015; Wurgaft et al., 2019.), and the possibility of methanogenesis (Sela-Adler et al., 2015).

140 From each interval, a 2.5 mL of sediment sample was collected and inserted immediately into an anoxic
141 10 mL glass bottle filled with 5 mL NaOH 1.5 N for headspace measurements of methane concentration
142 (after Nüsslein et al, 2003). Approximately 3 mL of sediment was sampled every 50 cm for porosity. In
143 addition, another 2.5 mL sediment sample was taken from each segment of the cores and transferred into
144 a 20 mL glass bottle filled with NaCl saturated solution for H_2 concentrations measurements. Sediment
145 samples from each segment of the cores were centrifuged on board if possible or in the lab within a day
146 by Sorval centrifuge at 9500 RPM under $4 \text{ }^\circ\text{C}$ and Ar atmosphere in order to extract pore-water for
147 chemical analysis. The supernatant was filtered ($0.22 \text{ }\mu\text{m}$) and analyzed for Fe(II), sulfate, sulfide, DIC
148 and the stable carbon isotope composition of the DIC ($\delta^{13}\text{C}_{\text{DIC}}$). After the pore-water extraction, the
149 sediment was analyzed for the content of the different reactive iron minerals (Table 2). In addition,
150 sediment sub-sample from each segment of the January 2017 core from Station SG-1 was kept at $-20 \text{ }^\circ\text{C}$
151 for molecular analysis. Due to high water content and movement in the uppermost part of the sediments,
152 two $\sim 30 \text{ cm}$ sediment cores were also sub-sampled separately, using a 0.0625 m^2 box corer (Ocean

153 Instruments BX 700 AI) and Perspex tubes during the September 2015 and January 2017 cruises. The
154 short cores were stored at 4 °C, cut in the lab within 24 hours after their collection and their results are
155 presented for the top sediment (Fig. 2a – d).

156 **2.3 Slurry incubation experiment**

157 The experimental set-up (Table 2) consisted of 11 bottles with sediment from the methanic zone (265-
158 285 cm depth) from Station SG-1, where iron reduction was apparent from the pore-water profiles (Fig.
159 2d). Prior to the beginning of the experiment, sediment from the designated depth had been homogenized
160 in an anoxic bag under N₂ atmosphere. It was then transferred under anoxic conditions to a 250 mL glass
161 bottle with the addition of synthetic sea water without sulfate to reach a 1:1 sediment:water slurry ratio
162 for 3 months incubation period. After the incubation period the slurry was sub-divided anoxically to the
163 11 experiment bottles (60 mL each), and synthetic sea water was added for final sediment:water ratio of
164 1:3. The bottles were sealed with a crimped cap and were flushed with N₂ for 5 minutes, shaken
165 vigorously and flushed again, (repeated 3 times). Three experimental bottles were autoclaved twice to
166 serve as "killed" control for the experiment. The experimental bottles were amended with 1.6 g L⁻¹ of
167 hematite (Fe₂O₃) or 2.3 g L⁻¹ of magnetite (Fe₃O₄) to reach Fe(III) final concentration of 10 mmol L⁻¹.
168 The three killed bottles were amended with the iron oxides after they cooled down to room temperature.
169 H₂ was added to some treatments to test its potential as an electron donor. One mL of H₂ was injected by
170 gas tight syringe to the three killed bottles, to two bottles with the addition of hematite and to two bottles
171 with the addition of magnetite (to reach final concentration of ~4% of the head space volume). The
172 experimental bottles were sampled several times for dissolved Fe(II) concentrations during the 14 day
173 experiment period.

174 **2.4 Analytical methods**

175 **2.4.1 Pore-water analyses**

176 Methane concentrations in the pore-water were analyzed by Focus Gas Chromatograph (GC; Thermo)
177 equipped with FID detector with a detection limit of 50 µmol L⁻¹. To calculate the methane concentrations
178 the sediment porosity was considered. Porosity was determined by drying wet sediment samples at 60
179 °C until there was no weight loss (~48 h). It was calculated as the weight loss from the initial weight of
180 the samples. H₂ concentrations were analyzed in a Reducing Compound Photometer Gas Chromatograph
181 (RCP-GC; Peak Laboratories). Dissolved Fe(II) concentrations were measured using the ferrozine
182 method (Stookey, 1970) by a spectrophotometer at 562 nm wavelength with detection limit of 1 µmol L⁻¹.
183 Sulfide was measured using the Cline (1969) method by a spectrophotometer at 665 nm wavelength
184 with detection limit of 1 µmol L⁻¹. Total sulfur concentrations were measured in an inductively coupled
185 plasma atomic emission spectrometer (ICP-AES), Perkin Elmer Optima 3300, with an analytical error of
186 ±1% (average deviations from repeated measurements of a seawater standard). Since sulfide was not
187 detected in any of the sediment cores, the total sulfur concentration in each pore-water sample was
188 assumed to be the sulfate concentration of that sample. The δ¹³C_{DIC} values were measured on a DeltaV
189 Advantage Thermo© isotope-ratio mass-spectrometer (IRMS) at a precision of ±0.1 ‰. Results are
190 reported versus VPDB standard. Pore-water profiles of dissolved total sulfur, CH₄, δ¹³C_{DIC}, Fe(II) and
191 H₂ were produced during the study, and all of them are presented (Fig. 2). For each profile where

192 duplicate samples were taken the error bar is that of the average deviation of the mean of the duplicates,
193 in cases where only single samples were taken, it is the analytical error (if larger than the symbol).

194 **2.4.2 Sediment analysis**

195 Reactive Fe(III) in the sediments was measured according to Poulton and Canfield (2005) definition and
196 sequential extraction procedure. The different reactive iron minerals were separated to (1) carbonate-
197 associated Fe (Fe_{carb}) (i.e. siderite and ankerite); (2) easily reducible oxides (Fe_{ox1}) (i.e. ferrihydrite and
198 lepidocrocite); (3) reducible oxides (Fe_{ox2}) (i.e. hematite, goethite and akageneite) and (4) magnetite
199 (Fe_{mag}). Sediment samples were dried at 60°C, then, approximately 0.6 g dry sediment was inserted to a
200 centrifuge tube with 10 ml of a specific extractant at every stage under atmospheric conditions and
201 constant agitation (Table 3). The fluids were separated from the sediment by centrifugation and removed
202 from the tube with Pasteur pipette after every extraction stage. At the end of each extraction stage, the
203 extractant was transferred to a 15 mL falcon tube with 0.1 mL ascorbic acid and 0.1 mL ferrozine solution
204 to reduce all the Fe(III) to Fe(II) and fix it, then it was measured spectrophotometrically. The results
205 presented as "total reactive Fe(III)" are the sum of Fe_{ox1} , Fe_{ox2} and Fe_{mag} . The profile of pyrite (Fe_{py}) was
206 taken from Wurgaft et al., (2019).

207 **2.4.3 Quantitative PCR and 16S rRNA gene V4 amplicon pyrosequencing**

208 DNA was extracted from the sediment core of Station SG-1 from January 2017 using Power Soil DNA
209 Kit (MoBio Laboratories, Inc., Carlsbad, CA, USA) following manufacturer's instructions. Copy
210 numbers of selected genes were estimated with quantitative PCR (qPCR) as described previously (Niu
211 et al., 2017) using specific primers: Uni519f/Arc908R and bac341f/519r for archaeal and bacterial 16S
212 rRNA genes, respectively, and mlas/mcrA-rev for the *mcrA* gene, which encodes the α -subunit of methyl-
213 coenzyme M reductase. The amplification efficiency was 94.5%, 106.3% and 92.4% for the archaeal 16S
214 rRNA, bacterial 16S rRNA and the *mcrA* gene, respectively (the respective R^2 of the standard curve was
215 0.998, 0.998 and 0.995).

216 The V4 regions of bacterial and archaeal 16S rRNA genes were amplified using barcoded 515FB/806RB
217 primers (Walters et al., 2015) and Arch519/Arch806 primers (Song et al., 2013), respectively. PCR
218 mixture contained 6 – 10 ng total DNA, 5 μL 10 \times Ex Taq buffer, 4 μL 2.5 mmol L⁻¹ dNTP mix, 1 μL of
219 each primer, 0.25 μL Ex Taq polymerase (Ex-Taq; TaKaRa, Dalian, China) and 5 μL bovine serum
220 albumin (25 mg mL⁻¹) in a total volume of 50 μL . DNA was sequenced as 2 \times 150 bp reads using Illumina
221 MiSeq platform (Illumina, USA). Sequence quality assessments, chimera detection and down-stream
222 phylogenetic analyses were conducted in QIIME (Caporaso et al., 2010). Taxonomical assignments for
223 each OTU were performed in QIIME using the BLAST method and the SILVA128 reference database.
224 24056 to 132042 high quality sequences were obtained per sample, with the proportion of high-quality
225 sequence versus total sequence between 81.97 – 99.89%. Spearman correlation was performed using the
226 online calculator (<http://www.sthda.com/english/rsthd/correlation.php>) to test the relevance of
227 microbial abundance and communities with Fe(II) concentration along the depth of the sediment core
228 from 185 cm to the bottom 575 cm, which is the methanic zone of the sediment core according to the
229 geochemical profile (see the results below).

230 **3 Results**

231 **3.1 Geochemical profiles**

232 Geochemical pore-water profiles of several sediment cores from the three stations (SG-1, PC-3 and PC-
233 5 (Fig. 1, Table 1)) were produced in order to characterize the iron reduction process in the methanic
234 zone of the SE Mediterranean continental shelf and to identify its potential sources. The pore-water
235 profiles at Station SG-1 (Fig. 2a) show complete depletion of total sulfur at approximately 150 cm depth
236 in all cores. Sulfide concentrations were below the detection limit in all cores, indicating that the total
237 sulfur is mostly sulfate. The methane concentrations in the pore-water (Fig. 2b) show an increase with
238 depth immediately after the consumption of sulfate. The maximum methane concentration was
239 approximately 10 mmol L^{-1} at $\sim 140 \text{ cm}$ depth in June 2015. The other methane depth profiles show an
240 increase in the concentrations to approximately 2 mmol L^{-1} and then leveling off throughout the bottom
241 of the cores ($\sim 600 \text{ cm}$). Detected dissolved Fe(II) concentrations (Fig. 2d) were found in the upper iron
242 reduction zone (between $30 - 90 \text{ cm}$ depth), and a second peak was found in the deeper part of the
243 sediment, at the methanic zone (below 180 cm depth). Maximum dissolved Fe(II) concentrations reached
244 $84 \text{ } \mu\text{mol L}^{-1}$ in the upper iron reduction zone of the sediments and $65 \text{ } \mu\text{mol L}^{-1}$ in the methanic zone. The
245 $\delta^{13}\text{C}_{\text{DIC}}$ values (Fig. 2c) were the lowest ($-35 \text{ } \text{‰}$) as expected at the SMTZ depth, and the highest in the
246 methanic zone. H_2 concentrations (Fig. 2e) decreased to a minimum of $0.017 \text{ } \mu\text{mol L}^{-1}$ at 155 cm depth,
247 and then increased to a maximum of $0.147 \text{ } \mu\text{mol L}^{-1}$ at 485 cm depth.

248 Pore-water profiles from Station PC-3 (Fig. 2g – l) show similar patterns to Station SG-1 on all three
249 sampling dates, but with lower methane concentrations. Total sulfur (Fig. 2g) was completely depleted
250 within the upper 300 cm depth. Sulfide concentrations were below the detection limit at this station as
251 well. Methane profiles show an increase in methane concentration immediately after the consumption of
252 sulfate. The maximum methane concentration (Fig. 2h) reached 0.8 mmol L^{-1} at 450 cm depth in the
253 Aug-13 core. The dissolved Fe(II) profiles (Fig. 2j) show two peaks at this station as well, one in the
254 upper part of the sediment with maximum value of $32 \text{ } \mu\text{mol L}^{-1}$ at 177 cm depth, and another one with
255 maximum value of $64 \text{ } \mu\text{mol L}^{-1}$ at 390 cm depth at the methanic depth. The $\delta^{13}\text{C}_{\text{DIC}}$ values (Fig. 2i)
256 decreased from approximately $-10 \text{ } \text{‰}$ at the water-sediment interface to $-20 \text{ } \text{‰}$ at the SMTZ. Below that
257 zone there was an increase in $\delta^{13}\text{C}_{\text{DIC}}$ values to about $-5 \text{ } \text{‰}$ due to methanogenesis. H_2 concentrations
258 (Fig. 2k) remained around $2 \text{ } \mu\text{mol L}^{-1}$ along the core. The three deviating points that do not fit the clear
259 pattern are attributed to an analytical or sampling error.

260 Pore-water profiles from the core collected at Station PC-5 (Fig. S1) resemble the profiles of Station PC-
261 3. Total sulfur was depleted at approximately 300 cm , and methane concentrations increased below that
262 depth to 0.3 mmol L^{-1} . The Fe(II) profile shows two peaks in this core as well, one in the upper sediment
263 of $20 \text{ } \mu\text{M}$ at 150 cm depth and the second of $30 \text{ } \mu\text{mol L}^{-1}$ in the methanic zone. The $\delta^{13}\text{C}_{\text{DIC}}$ value
264 decreased from $-5 \text{ } \text{‰}$ at the water-sediment interface to $-25 \text{ } \text{‰}$ at the SMTZ, and below that depth
265 increased to $-17 \text{ } \text{‰}$ at the methanic zone.

266 In addition to the dissolved constituents' profiles, reactive iron minerals were extracted from the sediment
267 collected on September 2015, and operationally defined iron mineral fractions profiles from Stations SG-

268 1 and PC-3 were produced (Fig. 2f and l). In Station SG-1 there appears to be a slight variability in the
269 content of the minerals (Fig. 2f). The Fe_{carb} content in the upper part of the sediment was 0.22 dry wt%,
270 increased to ~0.45 dry wt % at 103 cm depth and then remained constant. The Fe_{ox1} content was 0.49 dry
271 wt % in the upper part of the sediment, peaked at 203 cm depth to 0.64 dry wt % and then decreased to
272 0.50 dry wt % at the bottom of the core. The Fe_{ox2} content was 2.15 dry wt % in the upper part of the
273 sediment, decreased to 1.03 dry wt % at 312 cm depth, and then it increased to 1.55 dry wt % at 427 cm
274 depth. Fe_{mag} content was 0.34 dry wt % in the upper part of the sediment, decreased to 0.32 dry wt % at
275 153 cm depth, increased to 0.35 at 253 cm depth, decreased to 0.23 dry wt % at 312 cm depth, and
276 increased again to 0.35 dry wt % at the bottom. A pyrite content profile from Station SG-1 was also
277 produced (Wurgaft et al., 2019) from the September 2015 cruise and shows two peaks; the uppermost of
278 1.10 wt % at 153 cm depth, and the lower one of 1.80 wt % at 312 cm depth. The total reactive Fe(III)
279 oxides profile showed a general decrease from 3.00 dry wt % at 13 cm depth to 2.27 dry wt % at 507 cm
280 depth, with two minimum peaks of 2.42 dry wt % at 103 cm and of 1.88 dry wt % at 312 cm.

281 In Station PC-3 there appeared to be smaller changes in the different iron mineral fractions with depth
282 (Fig. 2l). The Fe_{carb} content in the upper part of the sediment was 0.50 dry wt % and reached 0.69 dry wt
283 % in the deep sediment. The Fe_{ox1} content was approximately 1.00 dry wt % throughout the sediment
284 column. The Fe_{ox2} content was 0.78 dry wt % in the upper part of the sediment, increased to 0.89 dry wt
285 % at 167 cm depth and then decreased to 0.76 dry wt % at 495 cm depth. Fe_{mag} content was 0.83 dry wt
286 % in the upper part of the sediment, increased to 0.89 dry wt % at 167 cm, and then decreased again to
287 0.75 dry wt % at 495 cm depth. The total reactive Fe(III) oxides content varied between 2.10 dry wt %
288 (at 167 cm depth) and 1.76 dry wt % (at 137 cm depth).

289 **3.2 Abundance and diversity of bacteria and archaea**

290 The qPCR of bacterial and archaeal 16S rRNA genes from the SG-1 core (collected on January 2017)
291 revealed an abundance of bacterial genes between $1.46 - 9.45 \times 10^6$ copies per g wet sediment, while that
292 of archaea was between $8.15 \times 10^5 - 2.25 \times 10^7$ copies per g wet sediment (Fig. 3). The abundance of
293 bacteria and archaea decreased gradually in the top 95 cm, increased sharply at 125 cm depth within the
294 SMTZ, remained relatively stable with high abundance at 185 – 245 cm (the top layer of the methanic
295 zone), and then decreased. Notably, the abundance of both bacteria and archaea peaked within the
296 methanic zone at 245 cm in correspondence with Fe(II) concentration peak. However, it is not feasible
297 to compare the abundance of archaea and bacteria by this method due to bias caused by the PCR primers
298 used (Buongiorno et al., 2017). The abundance of the *mcrA* gene (Fig. 3) increased sharply from the
299 surface layer to the SMTZ, peaked at 155 cm and remained stable at 155 – 245 cm, indicative of active
300 anaerobic methane metabolism in the SMTZ and an active methanic zone. Spearman correlation test
301 (Table S2) shows that the abundance of the bacteria and archaea 16S rRNA genes and *mcrA* genes
302 correlated with Fe(II) concentration in the methanic zone, where *mcrA* gene correlated the most
303 significantly ($r = 0.5429$, p value = 0.04789).

304 Illumina-sequencing of the 16S rRNA gene revealed diverse bacterial and archaeal communities
305 throughout the SG-1 core (Fig. 4). Although no clear plateau was observed on species rarefaction curve

306 for the current sequencing depth (Fig. S2), Shannon diversity indices reached stable values, indicating
307 that those sequences well covered the diversity of bacterial and archaeal populations in the samples (Fig.
308 S3). Shannon index, based on 16S rRNA gene sequences, shows higher diversity in the top layers of the
309 sediment along with similar values through the core using the bacterial primers, while for sequences
310 using archaeal primers, the values varied in different layers (Table S1). The bacterial sequences were
311 affiliated with the following phyla: Planctomycetes (25.7%), Chloroflexi (23.2 %), Proteobacteria
312 (12.9%), Deinococcus-Thermus (9.9 %), Acidobacteria (3.5%), Aminicenantes (3.3 %), Spirochaetes
313 (2.3%), Deferribacteres (1.7%), Elusimicrobia (1.6%), Aerophobetes (1.6%), Nitrospirae (1.4%),
314 Firmicutes (1.4 %), Actinobacteria (1.4 %), TM6 (Dependentiae) (1.2%), Marinimicrobia (SAR406 clade)
315 (1.0%), and other taxa with less than 1% of the bacterial communities (Fig. 4a). Bathyarchaeota were the
316 predominant archaea in all the sediment layers, based on the high relative abundance of their 16S rRNA
317 gene sequences (91.0%). The remaining archaeal phyla comprised Euryarchaeota (3.2%),
318 Thaumarchaeota (2.4%), Lokiarchaeota (1.0%), and other phyla with less than 1% of the archaeal
319 communities (Fig. 4b). Spearman correlation analysis (Table S2) revealed that uncultured SBR1093 ($r =$
320 0.6176 , p value = 0.01859) from bacterial Candidate Phylum SBR1093, subgroup 26 of Acidobacteria ($r =$
321 0.5841 , p value = 0.02828), the uncultured bacterium from TK10 Class of Chloroflexi phylum ($r =$
322 0.5297 , p value = 0.0544) and uncultured Bathyarchaeota sp. (archaea) ($r = 0.5516$, p value = 0.04388)
323 correlated significantly with Fe(II) concentration.

324 **3.3 Incubation experiment**

325 Sediment from the observed deep iron reduction zone of Station SG-1 from January 2017 core was used
326 for a simple short-term (couple of weeks) slurry incubation experiment in order to characterize the iron
327 reduction process in the methanic zone. Hematite and magnetite, which were expected to survive the
328 sulfate zone, and were shown to be a source for AOM in lake sediments, were added to the slurries.
329 Indeed, the operationally defined iron mineral fractions profiles (Fig. 2f) confirm that hematite and
330 magnetite were abundant in the methanic zone in this core.

331 The results of the experiment are shown in figure 5. Dissolved Fe(II) concentrations show significant
332 increase from $11 \mu\text{mol L}^{-1}$ to approximately $90 \mu\text{mol L}^{-1}$ during the first three days in all the experimental
333 bottles, except for the killed bottles, implying that the reduction is microbially mediated. Another
334 observation was that the microorganisms were able to reduce both hematite and magnetite to the same
335 extent. In addition, no difference in the Fe(II) concentrations between bottles with and without the
336 addition of H_2 was observed.

337 **4 Discussion**

338 **4.1 General**

339 This study was performed in the SE Mediterranean (Fig. 1) above the area of a recently discovered gas
340 front (Schattner et al., 2012). The investigated methane was found in the shallow sediments ($\sim 1\text{-}5$ m
341 deep) and seems biogenic based on its low $\delta^{13}\text{C}_{\text{CH}_4}$ -values and high C1/C2 ratio (Sela-Adler et al., 2015).
342 Station SG-1 is located at the center of the gas front area, while Stations PC-3 and PC-5 are located at
343 the edges, and indeed methane related processes were more intensive at Station SG-1. The source of this

344 gas front is not certain, but it was speculated to be terrestrial organic matter (Schattner et al., 2012). Our
345 results suggest that there are two sources for methane in our shallow sediment: the first is from migration
346 of methane from this gas front area (Wurgaft et al., 2019), and the second is from *in-situ* methane
347 formation, where the relative contribution of each source is currently unknown. *In-situ* methanogenesis
348 in the shallow shelf sediments is evident by the geochemical profiles of $\delta^{13}\text{C}_{\text{DIC}}$ and $\delta^{13}\text{C}_{\text{CH}_4}$ (Sela-Adler
349 et al., 2015), by the microbial population abundance profile and by the functional *mcrA* gene profile
350 (Figs. 3 and 4, further discussed below). The TOC content in the methanic zone are $\sim 0.8\%$ at Station SG-
351 1 (Sela-Adler et al., 2015; Wurgaft et al., 2019) and $\sim 1\%$ at Station PC-3 (Sela-Adler et al., 2015), and
352 these levels are known to be able to support *in-situ* methanogenesis (Sivan et al., 2007).

353 The comparison between the sites show that methane reaches the highest concentrations at Station SG-1
354 (up to the saturation level (Sela-Adler et al., 2015)), specifically in the June 2015 profile (Fig. 2b). This
355 leads to intensive AOM by sulfate at the SMTZ, causing it to occur at shallower depth and to produce
356 lower $\delta^{13}\text{C}_{\text{DIC}}$ values than the other two stations. The relation between the upward fluxes of methane, the
357 SMTZ depth and the $\delta^{13}\text{C}_{\text{DIC}}$ values fit previous studies (e.g. Sivan et al., 2007). The higher methane
358 concentrations in the June 2015 profile is presumably due to intensive migration of methane from the
359 deeper sediments and/or more intensive methane production at the exact location of the core collected at
360 that time. The H_2 concentrations at Station SG-1 (Fig. 2e) were lower by two orders of magnitude than
361 the concentrations at Station PC-3 (Fig. 2k), perhaps due to more intensive hydrogen consuming
362 processes at Station SG-1 (i.e. sulfate reduction, methanogenesis, iron reduction (Conrad et al., 1986;
363 Lovley, 1991). Dissolved Fe(II) pore-water profiles (Figs. 2d and j) show some variability between the
364 cores within the same station, probably as a result of environmental variations.

365 Despite the pore-water profiles variability between the stations, they show a resemblance in their trends.
366 All geochemical pore-water and iron mineral fraction profiles suggest that the sediments in this area of
367 the SE Mediterranean shelf can be classified into three general depth-zones (Fig. 2): **zone 1** is the upper
368 part of the sediment, where the classical iron reduction occurs, probably coupled to organic matter
369 oxidation, with sulfate reduction below it; **zone 2** is the SMT depth, where methane starts to increase,
370 sulfate is completely depleted, and Fe(II) (Fig. 2d and j) is either present in low concentrations or absent
371 (probably due to the precipitation of iron-sulfide minerals). In addition, the $\delta^{13}\text{C}_{\text{DIC}}$ values are the lowest
372 in this zone, as expected from the intensive AOM process there, which uses the isotopically light carbon
373 of the methane as a carbon source with small fractionation (Whiticar, 1999; Holler et al., 2009); **zone 3**
374 is the methanic zone, where methane concentrations increased to the highest values in all stations, as did
375 the $\delta^{13}\text{C}_{\text{DIC}}$ since the carbon source for the methane comes mainly from CO_2 , leaving the residual DIC
376 heavier by about 60 ‰ (Whiticar, 1999; Conrad, 2005). At this zone, local maxima of Fe(II)
377 concentrations in the pore-water were found in all cores, indicating reduction of iron oxides. The slurry
378 experiment results show only a slight increase in Fe(II) concentrations in the killed bottles compared to
379 their significant increase in the non-killed bottles, inferring that the iron reduction in zone 3 is microbial
380 (Fig. 5).

381 **4.2 Potential methanic iron reduction pathways**

382 This observed intensive iron reduction in the methanic sediments is the first discovered in the SE
383 Mediterranean shelf. The phenomenon of iron reduction in the methanic depth has been observed before
384 in other marine provinces (Jørgensen et al., 2004; März et al., 2008; Slomp et al., 2013; Riedinger et al.,
385 2014; Treude et al., 2014; Oni et al., 2015; Egger et al., 2016). Yet, the type of link to the methane cycle
386 is not well understood. Usually, iron reduction is coupled to oxidation of organic matter (Lovley and
387 Phillips, 1988) and is performed by iron-reducing bacteria, which is probably the case in zone 1. It is
388 however questionable if this also stands for zone 3 and if not, what process is responsible for the iron
389 reduction at this depth and its relation to methane. The iron reduction in zone 3 can occur potentially via
390 four pathways: 1) oxidation of organic matter arriving from the SMTZ, where it is produced by the
391 microorganisms that live there and benefit from the upward migrating methane, 2) oxidation of the
392 methane itself, 3) H₂ oxidation or 4) oxidation of sulfur species through a cryptic cycle.

393 The oligotrophic nature of the water column in the studied area would suggest that intensive bacterial
394 iron reduction coupled with the oxidation of organic matter in zone 3 is less likely. Nevertheless, we
395 observe high methane concentrations in zone 3 in all three stations, where part of it is from upward
396 migration. This indicates that regardless of the surface water oligotrophic nature, the TOC substrate may
397 be enough to sustain all the microbial activity and to take part in the iron reduction process in the methanic
398 zone. This is possibly due to biomass production in the SMTZ (i.e. the microbial community including
399 ANMEs and sulfate reducing bacteria (Boetius et al., 2000)) and its rapid use in the methanic zone (so
400 the TOC content still seems low).

401 The importance of the methane flux as a carbon source that supports the deep microbial community at
402 zone 2 and 3 in the sediments of the SE Mediterranean can be illustrated by comparing the organic carbon
403 flux from the photic zone, with the flux of organic carbon that is oxidized by sulfate in the pore-water.
404 Using traps, Moutin and Raimbault (2002) estimated an export flux of $7.4 \pm 6.3 \text{ mgC m}^{-2} \text{ d}^{-1}$, which leaves
405 the photic zone. However, Wurgaft et al. (2019) estimated that the flux of DIC toward the SMTZ from
406 sulfate reduction is equivalent to $8 \pm 3 \text{ mgC m}^{-2} \text{ d}^{-1}$. Whereas the difference between the two fluxes is
407 statistically insignificant, it should be noted that the flux of organic material that survives aerobic
408 oxidation in the water column and the upper part of the sediment column, as well as anaerobic oxidation
409 by other electron acceptors with higher energy yield (Froelich et al., 1979; Emerson et al., 1980), is likely
410 to be substantially smaller than the flux measured by Moutin and Raimbault (2002). Therefore, it is
411 unlikely that export flux from the photic zone constitutes the sole source of carbon to the SMTZ. Wurgaft
412 et al. (2019) suggested that “external” methane, originates in deeper portions of the sediments, provides
413 an important source of carbon to the SMTZ in Station SG-1. Such fluxes of “external” methane are
414 common along continental margin sediments (e.g. Milkov and Sassen, 2002; Milkov, 2004; Zhang and
415 Lanoil, 2004; Paull et al., 2008; Fischer et al., 2013). Here, we suggest that this supply of methane, leads
416 to intensive sulfate-mediated AOM in the SMTZ, and that this intensive process and biomass may serve
417 as an additional substrate that “fuels” zone 3, activating the iron-oxides.

418 The recently discovered iron-coupled AOM process (Eq. 3) is the second potential process that can
419 involve iron-oxide reduction in the deep methanic zone (Sivan et al., 2011; Segarra et al., 2013; Slomp

420 et al., 2013; Riedinger et al., 2014; Egger et al., 2015; Rooze et al., 2016; Egger et al., 2017; Bar-Or et
421 al., 2017). Fe(III) as an electron acceptor for AOM provides a greater free energy yield than sulfate
422 (Zehnder and Brock, 1980), and its global importance was emphasized (Sivan et al., 2011; Segarra et al.,
423 2013; Sivan et al., 2014). Two of the main environmental conditions for iron-coupled AOM to occur are
424 high dissolved methane concentrations and abundant reducible iron oxides (Riedinger et al., 2005;
425 Riedinger et al., 2014; Egger et al., 2017). Thus, from our profiles it seems that AOM could be a valid
426 option, considering the high methane concentrations and the high sedimentation rates (0.1 cm y^{-1} (Bareket
427 et al., 2016)), which allow the iron oxides to survive the sulfidic zone and reach the methanic zone
428 (Riedinger et al., 2005; Riedinger et al., 2014; Egger et al., 2017). This can also be inferred from figure
429 6, where some association was observed between the dissolved Fe(II) concentrations and the methane
430 concentrations in zone 3. It seems that at high concentrations of Fe(II), methane concentrations are low
431 and vice versa. This could be a result of iron-coupled AOM that uses methane to reduce Fe(III)-oxides,
432 releasing dissolved Fe(II) to the pore-water. It can also suggest a type of competitive relationship between
433 methanogenesis and microbial iron reduction, or microbial population switching from methanogenesis
434 to iron reduction metabolism (e.g. Sivan et al., 2016). It should be noted that our experiment was not
435 designed to test AOM due to its short time scale of a few weeks, hence another long experiment with the
436 addition of the ^{13}C -labeled methane will enable us to shed more light on this association.

437 The third potential process that can be coupled to iron reduction in the methanic zone is H_2 oxidation. H_2
438 is an important intermediate in anoxic aquatic sediments. In this type of environment, it is produced
439 mainly by fermentation of organic matter (Chen et al., 2006), and can be involved in different microbial
440 processes where each process would need a certain amount of H_2 in order to occur (Lovley and Goodwin,
441 1988). The H_2 levels at Stations SG-1 and PC-3 (Fig. 2e and k) are relatively high in comparison to other
442 marine environments (Lilley et al., 1982; Novelli et al., 1987), suggesting that there is enough H_2 to
443 sustain the iron reduction process. The relatively high H_2 concentrations at these stations could be
444 explained by the dominance of H_2 production processes (i.e. fermentation (Chen et al., 2006)) compared
445 to H_2 consuming processes (i.e. sulfate reduction, methanogenesis, iron reduction (Conrad et al., 1986;
446 Lovley, 1991)). At Station PC-3, the H_2 concentrations (Fig. 2k) are constant in zone 3, this suggest that
447 in addition to being produced, H_2 is consumed as well. At Station SG-1 (Fig. 2e) there is a maximum
448 peak at zone 3, indicating that there is either more H_2 production or less H_2 consumption at this zone
449 compared to zone 2. This is reasonable considering the intensive microbial activity in zone 2. The
450 decrease in the H_2 concentrations below the peak suggests that H_2 consuming processes are intensive in
451 this zone. The H_2 involvement was tested by injecting 1 mL of this gas to the experimental bottles in the
452 methanic iron reduction process (Fig. 5). We observed that the increase of Fe(II) concentration was
453 similar in the bottles with H_2 addition compared to the bottles without H_2 . This could mean that either
454 there is enough H_2 in the sediments as it is, as implied by the H_2 pore-water profiles, or that at the
455 methanic depth H_2 is not involved in the iron reduction process.

456 The fourth potential way to reduce iron in zone 3 is by an active sulfur cycle. The pyrite profile supports
457 this possibility by showing two peaks, uppermost in zone 2 of ~1 wt% and the other in zone 3 of ~2 wt%
458 at about 300 cm depth (Fig. 2f). The peak at 300 cm depth indicates possible active sulfur cycle, even

459 though sulfate is already undetected at 200 cm. Thus, a possible scenario is that Fe(III) is reduced by
460 pyrite oxidation (Eq. 3) (Bottrell et al., 2000), which triggers the ‘cryptic’ sulfur cycle, as observed in
461 other marine sediments (Holmkvist et al., 2011; Brunner et al., 2016; Egger et al., 2016). In this cycle,
462 elemental sulfur, and eventually by disproportionation also sulfide and sulfate, are produced. The sulfide
463 reacts with iron-oxide and precipitates as FeS or as pyrite (FeS₂) (Holmkvist et al., 2011). The sulfate
464 can inhibit methanogenesis (Mountfort et al., 1980; Mountfort and Asher, 1981), which can result in the
465 enhancement of the iron reduction process due to competition for substrate with the methanogenesis
466 process. Another indication for an active sulfur cryptic cycle comes from the 16S rRNA sequencing
467 analysis (Fig. 4), which shows that Proteobacteria, a potential sulfur related bacteria phylum, is one of
468 the most abundant phyla in the sediments. Moreover, the increase in the abundance of Sva0485 order of
469 the deltaproteobacteria class, a known sulfate reducer (Tan et al., 2019), with depth supports an active
470 sulfur cycle in zone 3 as well.

471 **4.3 Potential microbial players**

472 Our data profiles and incubations indicate that the observed iron reduction in the methanic zone of the
473 SE Mediterranean shelf is performed by microbial activity. The microbial results show first that the
474 abundances of the bacteria and archaea (Fig. 4) are typical to oligotrophic marine sediments (e.g. South
475 China Sea that contains ~0.5 – 1 % TOC (Yu et al., 2018)). Second, even though potential bacterial iron
476 reducers, such as *Alicyclobacillus*, *Sulfobacillus*, *Desulfotomaculum* genera (Firmicutes), *Acidiphilium*
477 (Alphaproteobacteria), *Desulfobulbus*, *Desulfuromonas*, *Geobacter*, *Geothermobacter*,
478 *Anaeromyxobacter* (Deltaproteobacteria) and *Shewanella* (Gammaproteobacteria) (Weber et al., 2006)
479 comprise less than 0.1% of bacteria detected in the methanic zone (from 185 cm and below), it appears
480 that both the microbial abundance and the Fe(II) concentration peaked at this zone. Cultivation efforts
481 indicated that archaeal methanogens may also play a role in iron reduction within sediments (Sivan et al.,
482 2016). Moreover, the relative abundance of methane-metabolizing archaea was shown to correlate with
483 Fe(II) concentrations in Helgoland muds from the North Sea, where microbial abundance and the Fe(II)
484 concentrations peaked at the methanic zone (Oni et al., 2015), similarly to the results found in the SE
485 Mediterranean sediments. It is possible that methane-metabolizing archaea were involved in the iron
486 reduction in the SE Mediterranean sediments, as the highest *mcrA* gene copies per gram wet sediment
487 were detected in the SMTZ and in the top of the methanic zone (Fig. 3) where the Fe(II) concentrations
488 are high (Fig. 2d). Methanotrophs, such as ANMEs, were found to be involved in iron-coupled AOM in
489 marine and freshwater cultures (Scheller et al., 2016; McGlynn et al., 2015; Ettwig et al., 2016; Cai et
490 al., 2018). ANMEs were found here with relatively low frequencies (ANME1, below 1% in most
491 samples, circa 5% in the 185 cm layer), and their role in iron reduction within the SE Mediterranean
492 sediments remains to be tested.

493 In our study, Spearman correlation analysis at Station SG-1 (Table S2) revealed that bacterial phyla
494 SBR1093 (candidate Phylum), Acidobacteria and Chloroflexi, as well as archaeal Phylum
495 Bathyarchaeota showed significant positive correlation with a Fe(II) concentration in the methanic zone.
496 The Candidate Phylum SBR1093 was firstly identified in phosphate-removing activated sludge from a
497 sequencing batch reactor (Bond et al., 1995), and continuously detected in a short-chain fatty acid rich

498 environment such as wastewater treatment, and marine sediments (Wang et al., 2014). It was thought to
499 be capable of growing autotrophically, but the metabolic capabilities related to iron reduction remain
500 unclear. Strains of Acidobacteria and Chloroflexi phylum were found to be capable of iron reduction
501 (Kawaichi et al., 2013; Kulichevskaya et al., 2014). In addition, members of Acidobacteria were found
502 in iron-coupled AOM enrichment (Beal et al., 2009). The metabolic properties of Subgroup 26 from
503 Acidobacteria and TK10 Class of Chloroflexi are still not known. Bathyarchaeota are globally distributed
504 and account for a considerable fraction of the archaeal communities in the marine sediments, particularly,
505 in the Mediterranean Pleistocene sapropels (Coolen et al., 2002; Zhou et al., 2018). While Bathyarchaeota
506 have diverse metabolic capabilities (Lloyd et al., 2013; Meng et al., 2014; Evans et al., 2015; He et al.,
507 2016; Yu et al., 2018; Feng et al., 2019), their role in iron reduction warrants further studies, as suggested
508 from their high abundance here. Therefore, iron reduction and methane cycling within the deep methanic
509 zone may be facilitated by an interplay among bacterial and archaeal groups, whose physiology and
510 functions needs further investigation.

511 **5 Conclusions**

512 Our study used combined geochemical and microbial profiles together with a slurry incubation
513 experiment to show microbial iron reduction in methanic sediments, and the potential microbial
514 population performing this reduction. The Spearman analysis points out several potential microbial
515 players (both bacterial and archaeal) that correlate to the dissolved Fe(II) profiles (e.g. Bathyarchaeota,
516 Acidobacteria and Chloroflexi). Moreover, our study emphasizes that this iron reduction in the methanic
517 zone can occur even in sediments of oligotrophic seas such as the SE Mediterranean. We suggest that the
518 availability of iron minerals for reduction is linked to intensive upward fluxes of methane and high
519 sulfate-AOM rates that may produce available biomass or/and hydrogen, which fuel deeper microbial
520 processes. The deep iron reduction may also be linked to a cryptic sulfur cycle and iron-coupled AOM.

521 **5 Author contribution**

522 H.V and O.S designed research; B.H and O.S. were the PIs of the cruises; H.V, E.W and L.L performed
523 research and analyzed the data; H.V, O.S, B.H, F.W, M.RB and L.L synthesized the data and wrote the
524 paper.

525 The authors declare that they have no conflict of interest.

526 **6 Acknowledgments**

527 We thank the captain and crew of the R/V Shikmona and R/V Bat Galim from the Israel Oceanographic
528 and Limnological Research for all their help during field sampling. Many thanks to E. Eliani-Russak for
529 her technical assistance in the lab and to V. Boyko for her help with the reactive iron speciation procedure.
530 We also thank all of Prof. O. Sivan's lab members for their help. We would like to thank also to the
531 anonymous reviewers for their helpful and constructive comments. This study was supported by the joint
532 grant of Israel Science Foundation and the National Natural Science Foundation of China (ISF-NSFC)
533 [grant number 31661143022 (FW) and 2561/16 (OS)]. Funding was provided to H. Vigderovich by the
534 Mediterranean Sea Research Center of Israel.

535 **References**

- 536 Adler, M., Eckert, W. and Sivan, O.: Quantifying rates of methanogenesis and methanotrophy in Lake
537 Kinneret sediments (Israel) using pore-water profiles, *Limnol. Oceanogr.*, 56(4), 1525–1535,
538 doi:10.4319/lo.2011.56.4.1525, 2011.
- 539 Almogi-Labin, A., Herut, B., Sandler, A. and Gelman, F.: Rapid changes along the Israeli
540 Mediterranean coast following the damming of the Nile and their influence on the Israeli inner shelf.
541 *Geol. Surv. of Israel, Israel*, 32, 2009.
- 542 Antler, G., Turchyn, A. V., Herut, B. and Sivan, O.: A unique isotopic fingerprint of sulfate-driven
543 anaerobic oxidation of methane, *Geology*, 43(7), 1–4, doi:10.1130/G36688.1, 2015.
- 544 Astrahan, P., Silverman, J., Gertner, Y. and Herut, B.: Spatial distribution and sources of organic
545 matter and pollutants in the SE Mediterranean (Levantine basin) deep water sediments, *Mar. Pollut.*
546 *Bull.*, 116(1–2), 521–527, doi:10.1016/j.marpolbul.2017.01.006, 2017.
- 547 Bar-Or, I., Ben-Dov, E., Kushmaro, A., Eckert, W. and Sivan, O.: Methane-related changes in
548 prokaryotes along geochemical profiles in sediments of Lake Kinneret (Israel), *Biogeosciences*, 12,
549 2847–2860, doi:10.5194/bg-12-2847-2015, 2015.
- 550 Bar-Or, I., Elvert, M., Eckert, W., Kushmaro, A., Vigderovich, H., Zhu, Q., Ben-Dov, E. and Sivan, O.:
551 Iron-Coupled Anaerobic Oxidation of Methane Performed by a Mixed Bacterial-Archaeal Community
552 Based on Poorly Reactive Minerals, *Environ. Sci. Technol.*, 51, 12293–12301,
553 doi:10.1021/acs.est.7b03126, 2017.
- 554 Bareket, M. M., Bookman, R., Katsman, R., de Stigter, H. and Herut, B.: The role of transport
555 processes of particulate mercury in modifying marine anthropogenic secondary sources, the case of
556 Haifa bay, Israel, *Mar. Pollut. Bull.*, 105(1), 286–291, doi:10.1016/j.marpolbul.2016.02.014, 2016.
- 557 Beal, E. J., House, C. H. and Orphan, V. J.: Manganese-and Iron-Dependent Marine Methane
558 Oxidation, *Science*, 325(5937), 184–187, doi:10.1126/science.1169984, 2009.
- 559 Boetius, A., Ravensschlag, K., Schubert, C. J., Rickert, D., Widdel, F., Gieseke, A., Amann, R.,
560 Jürgensen, B. B., Witte, U. and Pfannkuche, O.: A marine microbial consortium apparently mediating
561 anaerobic oxidation of methane, *Lett. to Nat.*, 407(October), 623–626, 2000.
- 562 Bond, D. R. and Lovley, D. R.: Reduction of Fe (III) oxide by methanogens in the presence and
563 absence of extracellular quinones, *Environ. Microbiol.*, 4(2), 115–124, doi:10.1046/j.1462-
564 2920.2002.00279.x, 2002.
- 565 Bond, P. L., Hugenholtz, P., Keller, J. and Blackall, L. L.: Bacterial Community Structures of
566 Phosphate-Removing and Non-Phosphate-Removing Activated Sludges from Sequencing Batch
567 Reactors, *Appl. Environ. Microbiol.*, 61(5), 1910–1916, 1995.
- 568 Bottrell, S. H., Parkes, R. J., Cragg, B. a. and Raiswell, R.: Isotopic evidence for anoxic pyrite
569 oxidation and stimulation of bacterial sulphate reduction in marine sediments, *J. Geol. Soc. London.*,

570 157(4), 711–714, doi:10.1144/jgs.157.4.711, 2000.

571 Bray, M. S., Jieying, W., Reed, B. C., Kretz, C. B., Belli, K. M., Simister, R. L., Henny, C., Stewart, F.
572 J., DiChristina, T. J., Brandes, J. A., Fowle, D. A., Crowe, S. A. and Glass, J. B.: Shifting microbial
573 communities sustain multi-year iron reduction and methanogenesis in ferruginous sediment
574 incubations, *Geobiology*, doi:http://dx.doi.org/10.1101/087783., 2018.

575 Brunner, B., Arnold, G. L., Røy, H., Müller, I. A. and Jørgensen, B. B.: Off Limits : Sulfate below the
576 Sulfate-Methane Transition, *Front. earth Sci.*, 4(July), 1–16, doi:10.3389/feart.2016.00075, 2016.

577 Buongiorno, J., Turner, S., Webster, G., Asai, M., Shumaker, A. K., Roy, T., Weightman, A.,
578 Schippers, A. and Lloyd, K. G.: Interlaboratory quantification of Bacteria and Archaea in deeply buried
579 sediments of the Baltic Sea (IODP Expedition 347), *FEMS Microbiol. Ecol.*, (93), 1–16,
580 doi:10.1093/femsec/fix007, 2017.

581 Cai, C., Leu, A. O., Jianhua, G. X., Yuexing, G., Zhao, F. J. and Tyson, G. W.: A methanotrophic
582 archaeon couples anaerobic oxidation of methane to Fe (III) reduction, *ISME J.*, 8(13), 1929–1939,
583 doi:10.1038/s41396-018-0109-x, 2018.

584 Canfield, D. E.: Reactive iron in marine sediments, *Geochim. Cosmochim. Acta*, 53(3), 619–632,
585 doi:10.1016/0016-7037(89)90005-7, 1989.

586 Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer,
587 N., Peña, A. G., Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T., Knights, D., Koenig, J. E.,
588 Ley, R. E., Lozupone, C. A., McDonald, D., Muegge, B. D., Pirrung, M., Reeder, J., Sevinsky, J. R.,
589 Turnbaugh, P. J., Walters, W. A., Widmann, J., Yatsunencko, T., Zaneveld, J. and Knight, R.: QIIME
590 allows analysis of high-throughput com- munity sequencing data., *Nat. Methods*, (7), 335–336,
591 doi:10.1038/nmeth.f.303, 2010.

592 Chen, W. H., Chen, S. Y., Kumar Khanal, S. and Sung, S.: Kinetic study of biological hydrogen
593 production by anaerobic fermentation, *Int. J. Hydrogen Energy*, 31(15), 2170–2178,
594 doi:10.4304/jcp.6.4.740-746, 2006.

595 Cline, J. D.: Spectrophotometric determination of hydrogen sulfide in natural waters, *Limnol.*
596 *Oceanogr.*, 454–458, doi:https://doi.org/10.4319/lo.1969.14.3.0454, 1969.

597 Conrad, R.: Contribution of hydrogen to methane production and control of hydrogen concentrations in
598 methanogenic soils and sediments, *FEMS Microbiol. Ecol.*, 28(3), 193–202, 1999.

599 Conrad, R.: Quantification of methanogenic pathways using stable carbon isotopic signatures : a review
600 and a proposal, *Org. Geochem.*, 36, 739–752, doi:10.1016/j.orggeochem.2004.09.006, 2005.

601 Conrad, R., Schink, B. and Phelps, T. J.: Thermodynamics of H₂-consuming and H₂-producing
602 metabolic reactions in diverse methanogenic environments under in situ conditions, *FEMS Microbiol.*
603 *Ecol.*, 38, 353–360, 1986.

604 Coolen, M. J. L., Cypionka, H., Sass, A. M., Sass, H. and Overmann, J.: Ongoing modification of

605 Mediterranean pleistocene sapropels mediated by prokaryotes, *Science*, 296(June), 2407–2411,
606 doi:10.1126/science.1071893, 2002.

607 Crowe, S. A., Katsev, S., Leslie, K., Sturm, A., Magen, C., Nomosatryo, S., Pack, M. A., Kessler, J. D.,
608 Reeburgh, W. S., Robert S, J. A., Gonzalez, L., Douglas Haffner, G., Mucci, A., Sundby, B. and Fowle,
609 D.: The methane cycle in ferruginous Lake Matano, *Geobiology*, 9, 61–78, doi:10.1111/j.1472-
610 4669.2010.00257.x, 2011.

611 Egger, M., Rasigraf, O., Sapart, C. J., Jilbert, T., Jetten, M. S. M., Röckmann, T., Van Der Veen, C.,
612 Bânda, N., Kartal, B., Ettwig, K. F. and Slomp, C. P.: Iron-mediated anaerobic oxidation of methane in
613 brackish coastal sediments, *Environ. Sci. Technol.*, 49(1), 277–283, doi:10.1021/es503663z, 2014.

614 Egger, M., Kraal, P., Jilbert, T., Sulu-Gambari, F., Sapart, C. J., Röckmann, T. and Slomp, C. P.:
615 Anaerobic oxidation of methane alters diagenetic records of sulfur, iron and phosphorus in Black Sea
616 sediments, *Biogeosciences Discuss.*, (March), 1–39, doi:10.5194/bg-2016-64, 2016.

617 Egger, M., Hagens, M., Sapart, C. J., Dijkstra, N., van Helmond, N. A. G. M., Mogollón, J. M.,
618 Risgaard-Petersen, N., van der Veen, C., Kasten, S., Riedinger, N., Böttcher, M. E., Röckmann, T.,
619 Jørgensen, B. B. and Slomp, C. P.: Iron oxide reduction in methane-rich deep Baltic Sea sediments,
620 *Geochim. Cosmochim. Acta*, 207, 256–276, doi:10.1016/j.gca.2017.03.019, 2017.

621 Emerson, S., Jahnke, R., Bender, M., Froelich, P. and Klinkhammer, G.: Early diagenesis in sediments
622 from the eastern equatorial pacific, i. pore water nutrient and carbonate results, *Earth Planet. Sci. Lett.*,
623 49, doi:https://doi.org/10.1016/0012-821X(80)90150-8, 1980.

624 Ettwig, K. F., Zhu, B., Speth, D., Keltjens, J. T., Jetten, M. S. M. and Kartal, B.: Archaea catalyze iron-
625 dependent anaerobic oxidation of methane, *Proc. Natl. Acad. Sci.*, 113(45), 12792–12796,
626 doi:10.1073/pnas.1609534113, 2016.

627 Evans, P. N., Parks, D. H., Chadwick, G. L., Robbins, S. J., Orphan, V. J., Golding, S. D. and Tyson,
628 G. W.: Methane metabolism in the archaeal phylum Bathyarchaeota revealed by genome-centric
629 metagenomics, *Science*, 350(6259), 434–438, doi:10.1126/science.aac7745, 2015.

630 Feng, X., Wang, Y., Zubin, R. and Wang, F.: Core Metabolic Features and Hot Origin of
631 Bathyarchaeota, *Engineering*, doi:10.1016/j.eng.2019.01.011, 2019.

632 Fischer, D., Mogollón, J. M., Strasser, M., Pape, T., Bohrmann, G., Fekete, N., Spiess, V. and Kasten,
633 S.: Subduction zone earthquake as potential trigger of submarine hydrocarbon seepage, *Nat. Geosci.*,
634 6(8), 1–5, doi:10.1038/ngeo1886, 2013.

635 Froelich, P. N., Klinkhammer, G. P., Bender, M. L., Luedtke, N. A., Heath, G. R., Cullen, D., Dauphin,
636 P., Hammond, D., Hartman, B. and Maynard, V.: Early oxidation of organic matter in pelagic
637 sediments of the eastern equatorial Atlantic: Suboxic diagenesis, *Geochim. Cosmochim. Acta*, 43,
638 1075–1090, doi:https://doi.org/10.1016/0016-7037(79)90095-4, 1979.

639 He, Y., Li, M., Perumal, V., Feng, X., Fang, J., Xie, J., Sievert, S. M. and Wang, F.: Genomic and

640 enzymatic evidence for acetogenesis among multiple lineages of the archaeal phylum Bathyarchaeota
641 widespread in marine sediments, *Nat. Microbiol.*, (April), 1–9, doi:10.1038/nmicrobiol.2016.35, 2016.

642 Herut, B., Almogi-Labin, A., Jannink, N. and Gertman, I.: The seasonal dynamics of nutrient and
643 chlorophyll a concentrations on the SE Mediterranean shelf-slope, *Oceanol. Acta*, 23(7), 771–782,
644 doi:10.1016/S0399-1784(00)01118-X, 2000.

645 Hinrichs, K., Hayes, J. M. and Sylva, S. P.: Methane-consuming archaeobacteria in marine sediments,
646 *Lett. to Nat.*, 398(April), 802–805, 1999.

647 Hoehler, T. M., Alperin, M. J., Albert, D. B. and Martens, C. S.: Field and laboratory studies of
648 methane oxidation in an anoxic marine sediment: evidence for a methanogen-sulfate reducer
649 consortium, *Global Biogeochem. Cycles*, 8(4), 451–463, doi:https://doi.org/10.1029/94GB01800, 1994.

650 Holler, T., Wegener, G., Knittel, K., Boetius, A., Brunner, B., Kuypers, M. M. M. and Widdel, F.:
651 Substantial 13 C / 12 C and D / H fractionation during anaerobic oxidation of methane by marine
652 consortia enriched in vitro, *Environ. Microbiol. Reports.*, 1, 370–376, doi:10.1111/j.1758-
653 2229.2009.00074.x, 2009.

654 Holmkvist, L., Ferdelman, T. G. and Jørgensen, B. B.: A cryptic sulfur cycle driven by iron in the
655 methane zone of marine sediment (Aarhus Bay, Denmark), *Geochim. Cosmochim. Acta*, 75(12), 3581–
656 3599, doi:10.1016/j.gca.2011.03.033, 2011.

657 Iversen, N. and Jørgensen, B. B.: Anaerobic methane oxidation rates at the sulfate-methane transition
658 in marine sediments from Kattegat and Skagerrak (Denmark), *Limnol. Oceanogr.*, 1983(5), 944–955,
659 1985.

660 Jørgensen, B. B., Böttcher, M. E., Lüschen, H., Neretin, L. N. and Volkov, I. I.: Anaerobic methane
661 oxidation and a deep H₂S sink generate isotopically heavy sulfides in Black Sea sediments, *Geochim.*
662 *Cosmochim. Acta*, 68(9), 2095–2118, doi:10.1016/j.gca.2003.07.017, 2004.

663 Kawaichi, S., Ito, N., Kamikawa, R., Sugawara, T., Yoshida, T. and Sako, Y.: *Ardenticatena maritima*
664 gen. nov., sp. nov., a ferric iron- and nitrate-reducing bacterium of the phylum ‘Chloroflexi’ isolated
665 from an iron-rich coastal hydrothermal field, and description of *Ardenticatena classis nov.*, *Int. J. Syst.*
666 *Evol. Microbiol.*, (63), 2992–3002, doi:10.1099/ijs.0.046532-0, 2013.

667 Knittel, K. and Boetius, A.: Anaerobic Oxidation of Methane : Progress with an Unknown Process,
668 *Annu. Rev. Microbiol.*, 63, 311–334, doi:10.1146/annurev.micro.61.080706.093130, 2009.

669 Kress, N. and Herut, B.: Spatial and seasonal evolution of dissolved oxygen and nutrients in the
670 Southern Levantine Basin (Eastern Mediterranean Sea): chemical characterization of the water masses
671 and inferences on the N : P ratios, *Deep Sea Research Part I: Oceanographic Research Papers*, 48,
672 2347–2372, doi:10.1016/S0967-0637(01)00022-X, 2001.

673 Kulichevskaya, I. S., Suzina, N. E., Rijpstra, W. I. C., Dedysh, S. N. and Damste, J. S. S.: a facultative
674 anaerobe capable of dissimilatory iron reduction from subdivision 3 of the Acidobacteria, *Int. J. Syst.*

675 *Evol. Microbiol.*, (64), 2857–2864, doi:10.1099/ijms.0.066175-0, 2014.

676 Li, Y., Yu, S., Strong, J. and Wang, H.: Are the biogeochemical cycles of carbon , nitrogen , sulfur ,
677 and phosphorus driven by the “ Fe III – Fe II redox wheel ” in dynamic redox environments ?, *J. soils*
678 *sediments*, 12(5), 683–693, doi:10.1007/s11368-012-0507-z, 2012.

679 Lilley, M. D., Baross, J. A. and Gordon, L. I.: Dissolved hydrogen and methane in Saanich Inlet,
680 British Columbia, *Deep Sea Res. Part A, Oceanogr. Res. Pap.*, 29(12), 1471–1484, doi:10.1016/0198-
681 0149(82)90037-1, 1982.

682 Liu, D., Dong Hailiang, H., Bishop, M. E., Wang, H., Agrawal, A., Tritschler, S., Eberl, D. D. and Xie,
683 S.: Reduction of structural Fe(III) in nontronite by methanogen *Methanosarcina barkeri*, *Geochim.*
684 *Cosmochim. Acta*, 75(4), 1057–1071, doi:10.1016/j.gca.2010.11.009, 2011.

685 Lloyd, K. G., Schreiber, L., Petersen, D. G., Kjeldsen, K. U., Lever, M. A., Steen, A. D., Stepanauskas,
686 R., Richter, M., Kleindienst, S., Lenk, S., Schramm, A. and Jørgensen, B. B.: Predominant archaea in
687 marine sediments degrade detrital proteins, *Nature*, 496(7444), 215–218, doi:10.1038/nature12033,
688 2013.

689 Lovley, D. R. and Goodwin, S.: Hydrogen concentrations as an indicator of the predominant terminal
690 electron-accepting reactions in aquatic sediments, *Geochim. Cosmochim. Acta*, 52, 2993–3003,
691 doi:10.1016/0016-7037(88)90163-9, 1988.

692 Lovley, D.: Microbial Fe (III) reduction in subsurface environments, *FEMS Microbiol. Rev.*, 20, 305–
693 313, doi:10.1111/j.1574-6976.1997.tb00316.x/full, 1997.

694 Lovley, D. R.: Dissimilatory Fe(III) and Mn(IV) reduction., *Microbiol. Rev.*, 55(2), 259–287, 1991.

695 Lovley, D. R. and Phillips, E. J.: Novel mode of microbial energy metabolism: organic carbon
696 oxidation coupled to dissimilatory reduction of iron or manganese., *Appl. Environ. Microbiol.*, 54(6),
697 1472–1480, doi:10.1103/PhysRevLett.50.1998, 1988.

698 Lovley, D. R. and Phillips, E. J. P.: Organic matter mineralization with reduction of ferric iron in
699 anaerobic sediments., *Appl. Environ. Microbiol.*, 51(4), 683–689, doi:10.1080/01490458709385975,
700 1986.

701 Lovley, D. R., Stolz, J. F., Nord, G. L. and Phillips, E. J. P.: Anaerobic production of magnetite by a
702 dissimilatory iron-reducing microorganism, *Nature*, 330(6145), 252–254, doi:10.1038/330252a0, 1987.

703 Lovley, D. R., Coates, J. D., BluntHarris, E. L., Phillips, E. J. P. and Woodward, J. C.: Humic
704 substances as electron acceptors for microbial respiration, *Nature*, 382(6590), 445–448,
705 doi:10.1038/382445a0, 1996.

706 März, C., Hoffmann, J., Bleil, U., Lange, G. J. De and Kasten, S.: Diagenetic changes of magnetic and
707 geochemical signals by anaerobic methane oxidation in sediments of the Zambezi deep-sea fan (SW
708 Indian Ocean), *Mar. Geol.*, 255(3–4), 118–130, doi:10.1016/j.margeo.2008.05.013, 2008.

709 März, C., Riedinger, N., Sena, C. and Kasten, S.: Phosphorus dynamics around the sulphate-methane
710 transition in continental margin sediments : Authigenic apatite and Fe (II) phosphates, *Mar. Geol.*,
711 404(July), 84–96, doi:10.1016/j.margeo.2018.07.010, 2018.

712 McGlynn, S. E., Chadwick, G. L., Kempes, C. P. and Orphan, V. J.: Single cell activity reveals direct
713 electron transfer in methanotrophic consortia, *Nature*, 526(7574), 531–535,
714 doi:doi:10.1038/nature15512, 2015.

715 Meng, J., Xu, J., Qin, D., He, Y., Xiao, X. and Wang, F.: Genetic and functional properties of
716 uncultivated MCG archaea assessed by metagenome and gene expression analyses, *ISME J.*, 8(3), 650–
717 9, doi:10.1038/ismej.2013.174, 2014.

718 Milkov, A. V: Global estimates of hydrate-bound gas in marine sediments : how much is really out
719 there?, *Earth-Science Rev.*, 66, 183–197, doi:10.1016/j.earscirev.2003.11.002, 2004.

720 Milkov, A. V and Sassen, R.: Economic geology of offshore gas hydrate accumulations and provinces,
721 *Mar. Pet. Geol.*, 19, 1–11, doi:doi.org/10.1016/S0264-8172(01)00047-2, 2002.

722 Mountfort, D. O. and Asher, R. A.: Role of Sulfate Reduction Versus Methanogenesis in Terminal
723 Carbon Flow in Polluted Intertidal Sediment of Waimea Inlet, Nelson, New Zealand, *Appl. Environ.*
724 *Microbiol.*, 42(2), 252–258, doi:0099-2240/81/080252-07\$02.00/0, 1981.

725 Mountfort, D. O., Asher, R. a, Mays, E. L. and Tiedje, J. M.: Carbon and electron flow in mud and
726 sandflat intertidal sediments at delaware inlet, nelson, new zealand., *Appl. Environ. Microbiol.*, 39(4),
727 686–94, doi:0099-2240/80/04-0686/09\$02.00/0, 1980.

728 Moutin, T. and Raimbault, P.: Primary production , carbon export and nutrients availability in western
729 and eastern Mediterranean Sea in early summer 1996 (MINOS cruise), *J. Mar. Syst.*, 34, 273–288,
730 doi:doi.org/10.1016/S0924-7963(02)00062-3, 2002.

731 Neiwöhner, C., Hensen, C., Kasten, S., Zabel, M. and Schulz, H. D.: Deep sulfate reduction completely
732 mediated by anaerobic methane oxidation in sediments of the upwelling area off Namibia,
733 *Geochim.Cosmochim.Acta*, 62(3), 455–464, 1998.

734 Nir, Y.: Recent sediments of the Israel Mediterranean continental shelf and slope, University of
735 Gothenburg. <http://hdl.handle.net/2077/13267>, 1984.

736 Niu, M., Fan, X., Zhuang, G., Liang, Q. and Wang, F.: Methane-metabolizing microbial communities
737 in sediments of the Haima cold seep area , northwest slope of the South China Sea, *FEMS Microbiol.*
738 *Ecol.*, (March), 1–13, doi:10.1093/femsec/fix101, 2017.

739 Norði, K. Á., Thamdrup, B. and Schubert, C. J.: Anaerobic oxidation of methane in an iron-rich Danish
740 freshwater lake sediment, *Limnol. Oceanogr.*, 58(2), 546–554, doi:10.4319/lo.2013.58.2.0546, 2013.

741 Novelli, P. C., Scranton, M. I. and Michener, R. H.: Hydrogen Distributions in Marine Sediments
742 Hydrogen distributions, *Limnol. Oceanogr.*, 32(3), 565–576, doi:10.4319/lo.1987.32.3.0565, 1987.

743 Nüsslein, B., Eckert, W. and Conrad, R.: Stable isotope biogeochemistry of methane formation in
744 profundal sediments of Lake Kinneret (Israel), *Limnol. Oceanogr.*, 48(4), 1439–1446,
745 doi:10.4319/lo.2003.48.4.1439, 2003.

746 Oni, O., Miyatake, T., Kasten, S., Richter-Heitmann, T., Fischer, D., Wagenknecht, Laura Kulkarni, A.,
747 Blumers, M., Shylin, S. I., Ksenofontov, Vadim Costa, B. F. O., Klingelhöfer, G. and Friedrich, M. W.:
748 Distinct microbial populations are tightly linked to the profile of dissolved iron in the methanic
749 sediments of the Helgoland mud area , North Sea, *Front. Microbiol.*, 6(May), 1–15,
750 doi:10.3389/fmicb.2015.00365, 2015.

751 Orphan, V. J., House, C. H. and Hinrichs, K.: Methane-Consuming Archaea Revealed by Directly
752 Coupled Isotopic and Phylogenetic Analysis, *Science*, 293(July), 484–488,
753 doi:10.1126/science.1061338, 2001.

754 Paull, C. K., Normark, W. R., Ussler, W., Caress, D. W. and Keaten, R.: Association among active
755 seafloor deformation , mound formation , and gas hydrate growth and accumulation within the seafloor
756 of the Santa Monica Basin , offshore California, *Mar. Geol.*, 250, 258–275,
757 doi:10.1016/j.margeo.2008.01.011, 2008.

758 Poulton, S. W. and Canfield, D. E.: Development of a sequential extraction procedure for iron:
759 Implications for iron partitioning in continentally derived particulates, *Chem. Geol.*, 214(3–4), 209–
760 221, doi:10.1016/j.chemgeo.2004.09.003, 2005.

761 Poulton, S. W., Krom, M. D. and Raiswell, R.: A revised scheme for the reactivity of iron
762 (oxyhydr)oxide minerals towards dissolved sulfide, *Geochim. Cosmochim. Acta*, 68(18), 3703–3715,
763 doi:10.1016/j.gca.2004.03.012, 2004.

764 Riedinger, N., Pfeifer, K., Kasten, S., Garming, J. F. L., Vogt, C. and Hensen, C.: Diagenetic
765 alterations of magnetic signals by anaerobic oxidation of methane related to a change in sedimentation
766 rate, *Geochim. Cosmochim. Acta*, 69(16), 4117–4126, doi:10.1016/j.gca.2005.02.004, 2005.

767 Riedinger, N., Formolo, M. J., Lyons, T. W., Henkel, S., Beck, A. and Kasten, S.: An inorganic
768 geochemical argument for coupled anaerobic oxidation of methane and iron reduction in marine
769 sediments., *Geobiology*, 12(2), 172–81, 2014.

770 Riedinger, N., Brunner, B., Krastel, S., Arnold, G. L., Wehrmann, L. M., Formolo, M. J., Beck, A.,
771 Bates, S. M., Henkel, S., Kasten, S. and Lyons, T. W.: Sulfur Cycling in an Iron Oxide-Dominated ,
772 Dynamic Marine Depositional System : The Argentine Continental Margin, *Front. earth Sci.*, 5(May),
773 doi:10.3389/feart.2017.00033, 2017.

774 Roden, E. E.: Fe(III) Oxide Reactivity Toward Biological versus Chemical Reduction, *Environ. Sci.*
775 *Technol.*, 37(7), 1319–1324, doi:10.1021/es026038o, 2003.

776 Roden, E. E.: Geochemical and microbiological controls on dissimilatory iron reduction, *Comptes*
777 *Rendus - Geosci.*, 338(6–7), 456–467, doi:10.1016/j.crte.2006.04.009, 2006.

778 Roden, E. E. and Wetzel, R. G.: Organic carbon oxidation and suppression of methane production by
779 microbial Fe(III) oxide reduction in vegetated and unvegetated freshwater wetland sediments, *Limnol.*
780 *Oceanogr.*, 41(8), 1733–1748, doi:10.4319/lo.1996.41.8.1733, 1996.

781 Roden, E. E. and Wetzel, R. G.: Kinetics of microbial Fe (III) oxide reduction in freshwater wetland
782 sediments, *Limnol. Ocean.*, 47(1), 198–211, doi:10.4319/lo.2002.47.1.0198, 2002.

783 Rooze, J., Egger, M., Tsandev, I. and Slomp, C. P.: Iron-dependent anaerobic oxidation of methane in
784 coastal surface sediments: potential controls and impact, *Limnol. Oceanogr.*, (1),
785 doi:10.1002/lno.10275, 2016.

786 Rotaru, B. A. and Thamdrup, B.: A new diet for methane oxidizers, *Biogeochemistry*, 351(6274), 658–
787 660, doi:10.1126/science.aaf0741, 2016.

788 Salem, D. M. S. A., Khaled, A. and Nemr, A. El: Assessment of pesticides and polychlorinated
789 biphenyls (PCBs) in sediments of the Egyptian Mediterranean Coast, *Egypt. J. Aquat. Res.*, 39(3),
790 141–152, doi:10.1016/j.ejar.2013.11.001, 2013.

791 Sandler, A. and Herut, B.: Composition of clays along the continental shelf off Israel: Contribution of
792 the Nile versus local sources, *Mar. Geol.*, 167(3–4), 339–354, doi:10.1016/S0025-3227(00)00021-9,
793 2000.

794 Schattner, U., Lazar, M., Harari, D. and Waldmann, N.: Active gas migration systems offshore northern
795 Israel, first evidence from seafloor and subsurface data, *Cont. Shelf Res.*, 48, 167–172,
796 doi:10.1016/j.csr.2012.08.003, 2012.

797 Scheller, S., Yu, H., Chadwick, G. L., McGlynn, S. E. and Orphan, V. J.: Artificial electron acceptors
798 decouple archaeal methane oxidation from sulfate reduction, *Science*, 351(6274), 1754–1756,
799 doi:10.1126/science.aad7154, 2016.

800 Segarra, K. E. a, Comerford, C., Slaughter, J. and Joye, S. B.: Impact of electron acceptor availability
801 on the anaerobic oxidation of methane in coastal freshwater and brackish wetland sediments, *Geochim.*
802 *Cosmochim. Acta*, 115, 15–30, doi:10.1016/j.gca.2013.03.029, 2013.

803 Sela-Adler, M., Herut, B., Bar-Or, I., Antler, G., Eliani-Russak, E., Levy, E., Makovsky, Y. and Sivan,
804 O.: Geochemical evidence for biogenic methane production and consumption in the shallow sediments
805 of the SE Mediterranean shelf (Israel), *Cont. Shelf Res.*, 101, 117–124, doi:10.1016/j.csr.2015.04.001,
806 2015.

807 Sivan, O., Schrag, D. P. and Murray, R. W.: Rates of methanogenesis and methanotrophy in deep-sea
808 sediments, *Geobiology*, 5(2), 141–151, doi:10.1111/j.1472-4669.2007.00098.x, 2007.

809 Sivan, O., Adler, M., Pearson, A., Gelman, F., Bar-Or, I., John, S. G. and Eckert, W.: Geochemical
810 evidence for iron-mediated anaerobic oxidation of methane, *Limnol. Oceanogr.*, 56(4), 1536–1544,
811 doi:10.4319/lo.2011.56.4.1536, 2011.

812 Sivan, O., Antler, G., Turchyn, A. V., Marlow, J. J. and Orphan, V. J.: Iron oxides stimulate sulfate-

813 driven anaerobic methane oxidation in seeps, *Proc. Natl. Acad. Sci.*, 111, E4139–E4147,
814 doi:10.1073/pnas.1412269111, 2014.

815 Sivan, O., Shusta, S. and Valentine, D. L.: Methanogens rapidly transition from methane production to
816 iron reduction, *Geobiology*, 190–203, doi:10.1111/gbi.12172, 2016.

817 Slomp, C. P., Mort, H. P., Jilbert, T., Reed, D. C., Gustafsson, B. G. and Wolthers, M.: Coupled
818 Dynamics of Iron and Phosphorus in Sediments of an Oligotrophic Coastal Basin and the Impact of
819 Anaerobic Oxidation of Methane, *Plos One*, 8(4), doi:10.1371/journal.pone.0062386, 2013.

820 Song, Z., Wang, F., Zhi, X., Chen, J., Zhou, E., Liang, F., Xiao, X., Tang, S., Jiang, H., Zhang, C. L.,
821 Dong, H. and Li, W.: Bacterial and archaeal diversities in Yunnan and Tibetan hot springs , China,
822 *Environ. Microbiol.*, 15, 1160–1175, doi:10.1111/1462-2920.12025, 2013.

823 Stookey, L. L.: Ferrozine-a new spectrophotometric reagent for iron, *Anal. Chem.*, 42(7), 779–781,
824 doi:10.1021/ac60289a016, 1970.

825 Tan, S., Liu, J., Fang, Y., Hedlund, B. P., Lian, Z., Li, L. H. J., Li, L. H. W., Dong, H. J. H. and Shu,
826 W.: Insights into ecological role of a new deltaproteobacterial order Candidatus
827 *Acidulodesulfobacterales* by metagenomics and metatranscriptomics, *ISME J.*, doi:10.1038/s41396-
828 019-0415-y, 2019.

829 Treude, T., Krause, S., Maltby, J., Dale, A. W., Coffin, R. and Hamdan, L. J.: Sulfate reduction and
830 methane oxidation activity below the sulfate-methane transition zone in Alaskan Beaufort Sea
831 continental margin sediments: Implications for deep sulfur cycling, *Geochim. Cosmochim. Acta*, 144,
832 217–237, doi:10.1016/j.gca.2014.08.018, 2014.

833 Valentine, D. L.: Biogeochemistry and microbial ecology of methane oxidation in anoxic
834 environments: A review, *Antonie van Leeuwenhoek, Int. J. Gen. Mol. Microbiol.*, 81, 271–282,
835 doi:10.1023/A:1020587206351, 2002.

836 Walters, W., Hyde, E. R., Berg-lyons, D., Ackermann, G., Humphrey, G., Parada, A., Gilbert, J. A. and
837 Jansson, J. K.: Transcribed Spacer Marker Gene Primers for Microbial Community Surveys, *Am. Soc.*
838 *Microbiol.*, 1(1), 1–10, doi:10.1128/mSystems.00009-15, 2015.

839 Wang, Z., Guo, F., Liu, L. and Zhang, T.: Evidence of Carbon Fixation Pathway in a Bacterium from
840 Candidate Phylum SBR1093 Revealed with Genomic Analysis, *Plos One*, 9(10),
841 doi:10.1371/journal.pone.0109571, 2014.

842 Weber, K. A., Urrutia, M. M., Churchill, P. F., Kukkadapu, R. K. and Roden, E. E.: Anaerobic redox
843 cycling of iron by freshwater sediment microorganisms, *Environ. Microbiol.*, 8(1), 100–113,
844 doi:10.1111/j.1462-2920.2005.00873.x, 2006.

845 Whiticar, M. J.: Carbon and hydrogen isotope systematics of bacterial formation and oxidation of
846 methane, *Chem. Geol.*, 161(1–3), 291–314, doi:10.1016/S0009-2541(99)00092-3, 1999.

847 Wurgaft, E., Findlay, A. J., Vigderovich, H., Herut, B. and Sivan, O.: Sulfate reduction rates in the

848 sediments of the Mediterranean continental shelf inferred from combined dissolved inorganic carbon
849 and total alkalinity profiles, *Mar. Chem.*, 211, 64–74, doi:10.1016/j.marchem.2019.03.004, 2019.

850 Yan, Z., Joshi, P., Gorski, C. A. and Ferry, J. G.: A biochemical framework for anaerobic oxidation of
851 methane driven by Fe(III)-dependent respiration, *Nat. Commun.*, (iii), 1–9, doi:10.1038/s41467-018-
852 04097-9, 2018.

853 Yu, T., Wu, W., Liang, W., Alexander, M. and Hinrichs, K.: Growth of sedimentary Bathyarchaeota on
854 lignin as an energy source, *Proc. Natl. Acad. Sci.*, 115(23), 6022–6027, doi:10.1073/pnas.1718854115,
855 2018.

856 Zehnder, a. J. B. and Brock, T. D.: Anaerobic methane oxidation: Occurrence and ecology, *Appl.*
857 *Environ. Microbiol.*, 39(1), 194–204, <http://aem.asm.org/content/39/1/194>, 1980.

858 Zhang, C. L. and Lanoil, B.: Geomicrobiology and biogeochemistry of gas hydrates and cold seeps,
859 *Chemical Geol.*, 205, 187–194, doi:10.1016/j.chemgeo.2004.01.001, 2004.

860 Zhang, J., Dong, H., Liu, D., Fischer, T. B., Wang, S. and Huang, L.: Microbial reduction of Fe(III) in
861 illite–smectite minerals by methanogen *Methanosarcina mazei*, *Chem. Geol.*, 292–293, 35–44,
862 doi:10.1016/j.chemgeo.2011.11.003, 2012.

863 Zhang, J., Dong, H., Liu, D. and Agrawal, A.: Microbial reduction of Fe(III) in smectite minerals by
864 thermophilic methanogen *Methanothermobacter thermautotrophicus*, *Geochim. Cosmochim. Acta*, 106,
865 203–215, doi:10.1016/j.gca.2012.12.031, 2013.

866 Zhou, Z., Pan, J., Wang, F., Gu, J. and Li, M.: Bathyarchaeota : globally distributed metabolic
867 generalists in anoxic environments, *FEMS Microbiol. Rev.*, (May), 639–655,
868 doi:10.1093/femsre/fuy023, 2018.

869

870

871 **Table 1:** Cores sampling details: dates, water depths and locations.

Date	station	water depth (m)	Latitude	Longitude
August 14, 2013	PC-5	87	32°55.47'	34°54.01'
	PC-3	81	32°55.29'	34°54.14'
February 6, 2014	PC-3	82	32°55.30'	34°54.14'
January, 2015	PC-3	82	32°55.30'	34°54.14'
June 9, 2015	SG-1	89	32°57.87'	34°55.30'
September 17, 2015	SG-1	84	32°57.91'	34°55.27'
January 24, 2017	SG-1	85	32°57.51'	34°55.15'

872

873 **Table 2:** Experimental set-up of the slurry incubation experiment.

Treatment	Number of bottles
Hematite	2
Magnetite	2
Hematite + H₂	2
Magnetite + H₂	3
Killed + hematite + H₂	2
Killed + magnetite + H₂	1

874

875 **Table 3:** Summary of reactive iron extraction procedure (after Poulton and Canfield, 2005).

Extractant	Target compounds	Analyzed species	Formula	Shaking time (h)
Magnesium chloride	Ion-exchangeable Fe(II)	Adsorbed ferrous iron	Fe ²⁺	2
Sodium acetate	Iron carbonates	Siderite Ankerite	FeCO ₃ Ca(Fe ⁺² ,Mg ⁺² ,Mn ⁺²)(CO ₃) ₂	24
Hydroxylamine hydrochloride	"Easily reducible" Iron(hydr)oxides	Ferrihydrite Lepidocrocite	Fe ³⁺ ₂ O ₃ *0.5(H ₂ O) γ-FeOOH	48
Sodium dithionite	"Reducible" oxides	Goethite, Hematite, Akageneite	α-FeOOH Fe ₂ O ₃ β-FeOOH	2
Ammonium oxalate	Poorly crystalline	Magnetite	Fe ₃ O ₄	6

876

877 **Figures captions:**

Figure 1: A map of the study area with the location of the three stations that were sampled SG-1, PC-3 and PC-5 (after Wurgaft et al., 2019).

878 **Figure 2:** Geochemical pore-water profiles of: total S, CH₄, δ¹³C_{DIC}, dissolved Fe(II), H₂ and extractable Fe fractions from sediment cores collected at the two stations: SG-1 (a-f) and PC-3 (g-l) in the SE Mediterranean. The profiles are divided roughly into three zones according to the dominant processes: upper microbial iron and sulfate reduction, sulfate-methane transition zone (SMTZ), and the methanic zone at the deep part. The dashed line in the CH₄ graph at SG-1 station represents the CH₄ saturation value in the pore-water (Sela-Adler et al., 2015). The following extractable Fe fraction profiles of stations SG-1 (f) and PC-3 (l) are from the September 2015 and January 2015 cruise (respectively): Fe_{carb} (●), Fe_{ox1} (■) Fe_{ox2} (▲), Fe_{mag} (▼), Fe_{py} (◆) (Wurgaft et al., 2019) and total reactive iron (●). The error bars for CH₄ are presented where duplicate sediment samples were collected. The error bars for Fe(II), δ¹³C_{DIC} and H₂ are presented where measurements from the same sample were repeated at least twice. The analytical errors were too small to be displayed.

Figure 3: Sedimentary depth profiles of bacterial and archaeal 16S rRNA and *mcrA* functional genes of station SG-1 from January 2017, divided to three zones (as described in figure 2). Triplicates were produced from each sample with error bars smaller than the symbols displayed.

Figure 4: Phyla level classification of bacterial (a) and archaeal (b) diversity in the sediments of Station SG-1 from January 2017.

Figure 5: Dissolved Fe(II) results of the sediment slurry incubation experiment. The sediment was collected from Station SG-1 on January 2017 from sediment depth of 265-285 cm. The error bars were smaller than the symbols displayed.

Figure 6: The relationship between dissolved Fe(II) concentrations and methane concentrations in zone 3 of (a) Station SG-1 and (b) Station PC-3. An inverse association is observed between the two species, suggesting a relationship of competition or iron-coupled AOM.

879 **Figures:**

880 **Figure 1**

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905


906

907

908

909

910



911 **Figure 2**

912


913



914

915

916

917







919 **Figure 3**

920

921

922

923




924 **Figure 4**


925

926

927




928 **Figure 5**



929

930

931 **Figure 6**



932

933

934

935

936

937

938

939