

# Using Respiration Quotients to Track Changing Sources of Soil Respiration Seasonally and with Experimental Warming

Caitlin Hicks Pries<sup>1,2</sup>, Alon Angert<sup>3</sup>, Cristina Castanha<sup>2</sup>, Boaz Hilman<sup>3,4</sup>, and Margaret S. Torn<sup>2</sup>

<sup>1</sup>Department of Biological Sciences, Dartmouth College, Hanover, NH, 03784, United States of America

<sup>2</sup>Climate and Ecosystem Science Division, Earth and Environmental Science Area, Lawrence Berkeley National Laboratory, Berkeley, CA, 94720, United States of America

<sup>3</sup>The Institute of Earth Sciences, The Hebrew University of Jerusalem, Givat-Ram, Jerusalem 91904, Israel

<sup>4</sup>Currently at: Department of Biogeochemical Processes, Max-Planck Institute for Biogeochemistry, Jena, 07745, Germany

*Correspondence to:* Caitlin Hicks Pries (Caitlin.Pries@dartmouth.edu)

**Abstract.** Developing a more mechanistic understanding of soil respiration is hampered by the difficulty in determining the contribution of different organic substrates to respiration and in disentangling autotrophic versus heterotrophic and aerobic versus anaerobic processes. Here, we use a relatively novel tool for better understanding soil respiration: the apparent respiration quotient (ARQ). ARQ is the amount of CO<sub>2</sub> produced in the soil divided by the amount of O<sub>2</sub> consumed and it changes according to which organic substrates are being consumed and whether oxygen is being used as an electron acceptor. We investigated how the ARQ of soil gas varied seasonally, by soil depth, and by *in situ* experimental warming (+4°C) in a coniferous forest whole-soil-profile warming experiment over two years. We then compared the patterns in ARQ to those of soil δ<sup>13</sup>CO<sub>2</sub>. Our measurements showed strong seasonal variations in ARQ from ≈0.9 during the late spring and summer to ≈0.7 during the winter. This pattern likely reflected a shift from respiration being fueled by oxidized substrates like sugars and organic acids derived from root and root respiration during the growing season to more reduced substrates such as lipids and proteins derived from microbial necromass during the winter. This interpretation was supported by δ<sup>13</sup>CO<sub>2</sub> values, which were relatively lower, like lipids, in the winter and relatively higher, like sugars, in the summer. Furthermore, experimental warming significantly changed how both ARQ and δ<sup>13</sup>CO<sub>2</sub> responded to soil temperature. Wintertime ARQ and δ<sup>13</sup>CO<sub>2</sub> values were higher in heated than in control plots, probably due to the warming-driven increase in microbial activity that may have utilized oxidized carbon substrates, while growing season values were lower in heated plots. Experimental warming and phenology change the sources of soil respiration throughout the soil profile. The sensitivity of ARQ to these changes demonstrates its potential as a tool for disentangling the biological sources contributing to soil respiration.

## 29 1 Introduction

30 Despite making extensive measurements of soil respiration (Bond-Lamberty and Thomson, 2010), scientists lack methods to  
31 disentangle the processes underlying, and substrates contributing to, soil respiration, which hampers predictions of terrestrial  
32 carbon cycle responses to global change (Phillips et al., 2017). Mechanistic uncertainty surrounding soil respiration is partly  
33 responsible for the 1000 Pg spread in model predictions of end-of-century terrestrial carbon-climate feedbacks (Friedlingstein  
34 et al., 2013). Soil respiration is the flux of CO<sub>2</sub> from the soil surface to the atmosphere, which is dominated by autotrophic  
35 respiration from plant roots and heterotrophic respiration from soil microbes. Heterotrophic respiration, which has increased  
36 globally over the past three decades (Bond-Lamberty et al., 2018), is itself the sum of various processes using different sources  
37 of energy. For example, microbes consume different organic substrates depending on what molecules are accessible and  
38 whether the microbes are living in the rhizosphere or bulk soil, and microbes utilize different terminal electron acceptors  
39 depending on O<sub>2</sub> availability in the microsites in which they reside (Keiluweit et al., 2016; Liptzin et al., 2011). The electron  
40 donors (the organic substrates) and the electron acceptors used by soil microbes during respiration cannot be resolved by  
41 measuring the CO<sub>2</sub> flux alone. Previous studies have used measurements of δ<sup>13</sup>C to partition respiration into autotrophic and  
42 heterotrophic components (e.g., Dorrepaal et al., 2009), radiocarbon to partition respiration sources by age (e.g., Trumbore,  
43 2000), or both isotopes in combination to more finely separate respiration among sources (e.g., Hicks Pries et al., 2013;  
44 Hopkins et al., 2012). However, isotopes are not the only way to disentangle soil respiration's various components (Subke et  
45 al., 2006).

46  
47 Our ability to understand soil respiration is limited by measuring only one half of the respiration equation, the CO<sub>2</sub> produced.  
48 Simultaneously measuring the O<sub>2</sub> consumed can provide a more mechanistic understanding of the processes and substrates  
49 contributing to soil respiration (Phillips et al., 2017). The paired measurements of CO<sub>2</sub> and O<sub>2</sub> can be used to calculate a  
50 respiration quotient (RQ; Angert and Sherer, 2011). All organic matter has an oxidative ratio (1/RQ), which can be calculated  
51 based on an elemental analysis of its C, H, O, and N (Masiello et al., 2008). The oxidation state of carbon in carbohydrates is  
52 0 with a corresponding RQ of 1 based on its elemental structure. More reduced energy sources such as lipids have lower RQ  
53 values (≈0.73) and the RQ of proteins range from 0.67 to 1; more oxidized sources such as organic acids have RQ ranges from  
54 1 to 4 (Masiello et al., 2008; Table 1). The RQ of aerobic respiration therefore changes based on what substrates are being  
55 consumed (Dilly, 2001; Theenhaus et al., 1997). Anaerobic respiration increases RQ to values greater than one, as electron  
56 acceptors like Fe(III) and NO<sub>3</sub><sup>-</sup> replace O<sub>2</sub>. Thus, RQ can help differentiate between the electron donors (organic substrates)  
57 and terminal electron acceptors used during soil respiration. We will refer to the 'apparent' respiration quotient (ARQ) because  
58 not all ecosystem CO<sub>2</sub> or O<sub>2</sub> fluxes are due to respiratory processes (Angert and Sherer, 2011). For example, fluctuating redox  
59 conditions can lead to consumption of O<sub>2</sub> during metal oxidation and drive ARQ below the value of the most reduced organic  
60 matter (Angert et al., 2015).

61

62 Thus far, CO<sub>2</sub>:O<sub>2</sub> ratios have been primarily used to understand large scale earth system processes and only few studies have  
63 examined processes within ecosystems. This ratio in atmospheric samples has been used to estimate a) the magnitude of the  
64 terrestrial carbon sink, because carbon uptake by terrestrial ecosystems is balanced by O<sub>2</sub> production whereas ocean CO<sub>2</sub>  
65 uptake is decoupled from O<sub>2</sub>, (Keeling, 1988; Keeling et al., 1996; Randerson et al., 2006; Worrall et al., 2013) and b)  
66 anthropogenic impacts on the carbon cycle, based on the principle that burning of reduced fossil fuels results in a different  
67 oxidative ratio than does photosynthesis and subsequent respiration of carbohydrates (Keeling, 1988). The CO<sub>2</sub>:O<sub>2</sub> ratio of  
68 ecosystem-atmosphere exchanges is an essential quantity in these carbon cycle calculations. CO<sub>2</sub>:O<sub>2</sub> ratios have been estimated  
69 from measurements of net ecosystem exchange of CO<sub>2</sub> and O<sub>2</sub> (e.g., Seibt et al., 2004) and from elemental analysis of biomass  
70 (Hockaday William C. et al., 2015; e.g., Masiello et al., 2008), both of which are assumed to be similar over multiyear  
71 timescales. In early carbon sink calculations, the oxidative ratio of ecosystem fluxes was assumed to be 1.1 (ARQ=0.9) based  
72 on a single study of temperate soils (Severinghaus, 1995). However, the few subsequent studies examining the CO<sub>2</sub>:O<sub>2</sub> ratio  
73 of soil respiration fluxes have shown soil fluxes can deviate widely from that value.

74

75 Soil ARQ from incubations shift as a result of temperature changes, substrate additions, and soil management. For example,  
76 the ARQ of peat soils decreased from about 1.1 to about 0.6 when temperatures increased from 0 to 20°C, attributed to changing  
77 substrate use (Chapman and Thurlow, 1998). Glucose additions to German forest soils increased soil ARQ to 0.95-1.0 from a  
78 basal value around 0.7 (Dilly, 2001; Theenhaus et al., 1997). Soils under organic agriculture were found to have a greater ARQ  
79 (1.19) than the same soils under conventional agriculture (0.72; Theenhaus et al., 1997). Soil ARQ in mesocosms containing  
80 pine seedlings changed seasonally and when the pine seedlings were cut, indicating the ratio is responsive to changes in  
81 vegetation (Andersen and Scagel, 1997; Scagel and Andersen, 1997).

82

83 Lastly, in one of the only studies using *in situ* measurements, soil ARQ taken from gas wells across multiple forested  
84 ecosystems ranged widely from 0.14 to 1.23 indicating the influence of abiotic processes that consume O<sub>2</sub> (Angert et al. 2015).  
85 The wide range in soil ARQ values associated with different biochemical conditions indicates the ratio has the potential to  
86 provide insight into the substrates contributing to respiration as well as into abiotic O<sub>2</sub> consumption. Finer scale research is  
87 needed, however, to explore ARQ values in the same soils under different conditions to learn what these values indicate about  
88 the processes and substrates contributing to soil respired CO<sub>2</sub>.

89

90 Here we investigated how the ARQ of soil gas *in situ* varied seasonally, by soil depth, and by experimental warming in a  
91 whole-soil-profile warming experiment in a well-drained, oxygenated coniferous forest soil (Hicks Pries et al., 2017). We  
92 characterized soil ARQ at 30 and 90 cm depths in the winter and growing season over two years and compared the patterns in  
93 ARQ to monthly patterns in soil profile δ<sup>13</sup>C. We hypothesized that ARQ values would change seasonally and with warming  
94 reflecting the values of the organic carbon substrates being consumed by microbes. Like ARQ, the δ<sup>13</sup>C of soil CO<sub>2</sub> is

95 influenced by the use of different organic substrates since more reduced substrates tend to have lower  $\delta^{13}\text{C}$  values (Bowling et  
96 al., 2008). By comparing ARQ values to other indicators of respiration sources, such as  $\delta^{13}\text{C}$ , augmented by what we  
97 understand about plant allocation of carbon substrates belowground, we aim to advance the utility of ARQ as a tracer of  
98 respiration processes.

## 99 **2 Methods**

### 100 **2.1 Warming Experiment**

101 The whole soil profile warming experiment is located at the University of California Blodgett Forest Research Station, in the  
102 Sierra Nevada foothills near Georgetown, CA at 1370 m above sea level. Mean annual precipitation is 1774 mm with most of  
103 it occurring from November through April and mean annual temperature is about 12.5°C (Bird and Torn, 2006). The  
104 experiment is in a thinned 80-year-old stand of mixed conifers including ponderosa pine (*Pinus ponderosa*), sugar pine (*Pinus*  
105 *lambertiana*), incense cedar (*Calocedrus decurrens*), white fir (*Abies concolor*), and douglas fir (*Pseudotsuga menziesii*). The  
106 soils are Holland series: fine-loamy, mixed, superactive, mesic Ultic Haploxeralfs of granitic origin with thick, >5 cm O  
107 horizons, minimal carbonates (Rasmussen et al., 2005), and a pH that ranges from 5.6 to 6.5 (Hicks Pries et al., 2018). The  
108 warming treatment warmed the soil +4°C to 1 m depth while maintaining the natural temperature gradient with depth and  
109 temporal variations in soil temperature as described in Hicks Pries et al. (2017). Briefly, there were three pairs of control and  
110 heated 3 m diameter circular plots heated by 22 vertical resistance heater cables in metal conduit (BriskHeat, Ohio, USA) that  
111 surrounded them. To compensate for surface heat loss, two concentric rings of heater cable at 1 and 2 m in diameter were  
112 installed 5 cm below the soil surface in heated plots. Unheated cables were installed similarly in control plots. Heating  
113 throughout the plot volume was generally even, ranging from 3.5 to 4.5°C except at 5 cm depth where the heated plots were  
114 on average only  $2.4 \pm 1.2^\circ\text{C}$  warmer than the control due to a lack of aboveground heating. Soil moisture was slightly decreased  
115 in the warmed plots by an average of 1.5-3.5% volumetric water content (Hicks Pries et al., 2017).

### 116 **2.2 Sample Collection and Analysis**

117 Dataloggers (CR1000, Campbell Scientific, Utah, USA) continuously recorded soil temperature and moisture at 30 min  
118 intervals. Temperature was monitored at 5, 15, 30, 50, 75, and 100 cm depths at a radial distance of 0.75 m from the center of  
119 each plot. Temperature probes consisted of thermistors (Omega 44005) epoxied to PVC rods, placed inside thin-walled steel  
120 conduit. To monitor soil moisture, we used an enviroSCAN (Sentek, Australia) probe fitted with capacitance sensors at 10, 30,  
121 50, and 90 cm at a radial distance of 0.75 m from the center of each plot. We calibrated the soil moisture measurements by  
122 comparing the sensor values at each depth to the volumetric water content measured in nearby (within 0.5 m) soil cores that  
123 were sampled five times over two years.

124

125 Each of the six plots has a set of gas wells at 15, 30, 50, 75, and 90 cm. The gas wells were 6.35 mm diameter stainless steel  
126 tubes inserted into the soil at a 45° angle to the desired depth and topped with straight swage pipefittings (Swagelok Ohio,  
127 USA) with septa. For CO<sub>2</sub> and δ<sup>13</sup>C measurements, samples were collected from the wells with a syringe on a nearly monthly  
128 basis from March 2014 through June 2017 (32 months total) and always during morning hours. After clearing the headspace  
129 in each well, a 25 ml gas sample was transferred to an evacuated 20 ml septum-topped glass vial. For analysis, 5 ml samples  
130 were injected into the small sample isotope module of a cavity ring down spectrometer (CRDS, Picarro, Santa Clara,  
131 California) where they were diluted with ultra zero air (without CO<sub>2</sub>). A four-point calibration curve ranging from 2,000 to  
132 20,000 ppm (δ<sup>13</sup>C=-26.7‰) was used to calculate the CO<sub>2</sub> concentration from the CRDS data and to correct for mass  
133 dependency of the δ<sup>13</sup>C measurement.

134

135 In July 2015, February 2016, April 2016, August 2016, March 2017, and June 2017, we collected additional samples from the  
136 30 (except July 2015) and 90 cm gas wells into 13 ml flasks equipped with O-ring valves (LouwersHanique, Hapert,  
137 Netherlands) to simultaneously measure CO<sub>2</sub> and O<sub>2</sub> concentrations in order to calculate ARQ. The flasks were analyzed in  
138 the laboratory at the Hebrew University by a closed system (The Hampadah; Hilman and Angert, 2016). This fully automated  
139 system uses an infra-red gas analyzer (IRGA) for CO<sub>2</sub> measurement (LI 840A LI-COR; Lincoln, NE, USA) and a fuel-cell  
140 based analyzer (FC-10; Sable Systems International, Las Vegas, NV, USA) for measuring O<sub>2</sub>. The flasks were analyzed within  
141 2-3 weeks of collection.

142

143 In June 2017, we also ran a set of short (3 hour) incubations of root-free soil and of excised roots collected adjacent to the  
144 experimental plots. We collected four mineral soil cores with a 5 cm diameter hammer corer, separated the cores into 0-20 and  
145 20-40 cm depths, and removed roots >1 mm diameter. Roots were collected from four 25 cm x 25 cm x 25 cm soil pits. We  
146 rinsed roots with water to remove soil and blotted them dry before placing them into mason jars. The root-free soil was also  
147 placed into mason jars, and both sets of mason jars were flushed with ambient, outside air. After a three-hour incubation of the  
148 root samples and a 21-hour incubation of the soil samples, the headspace was sampled for CO<sub>2</sub> and O<sub>2</sub> and analyzed as  
149 described above. Incubations were run at room temperature, which was similar to the field temperature at the time of collection.

### 150 2.3 Sample Calculations and Statistics

151 To calculate ARQ, we used the following equation from Angert et al. (2015):

152 
$$ARQ = -0.76 \frac{\Delta CO_2}{\Delta O_2}$$

153 Where ARQ is the apparent respiratory quotient, ΔCO<sub>2</sub> (ppmv) is the difference between CO<sub>2</sub> concentrations in the soil pore  
154 space gas and ambient (i.e., 0.5 to 1m aboveground) samples, ΔO<sub>2</sub> (ppmv) is the difference of the soil pore space O<sub>2</sub>  
155 concentration and ambient O<sub>2</sub> concentration, and 0.76 is the ratio of CO<sub>2</sub> to O<sub>2</sub> diffusivity in air (Massman, 1998). The negative  
156 sign is for convenience so the ARQ value will typically be positive, because the difference in O<sub>2</sub> concentration is always

157 negative. For the jar incubations we used the same equation without the 0.76 factor. Ambient CO<sub>2</sub> concentrations were  
158 measured in the field at the time of sampling with the CRDS, while the ambient O<sub>2</sub> concentration was assumed to be 20.95%  
159 (Rumble, 2019). To relate the δ<sup>13</sup>C value of soil pore space CO<sub>2</sub> to the δ<sup>13</sup>C of CO<sub>2</sub> production, we corrected the pore-space  
160 δ<sup>13</sup>C value for diffusion since <sup>13</sup>C diffuses slower in air than <sup>12</sup>C and thus the measured value does not accurately represent the  
161 value of production. For the correction, we used the following equation from Bowling et al. (2015):

$$162 \quad \delta_{production} = \frac{C_s(\delta_s - 4.4) - C_a(\delta_a - 4.4)}{1.0044(C_s - C_a)}$$

163 Where C<sub>s</sub> is the soil pore space CO<sub>2</sub> concentration (ppmv), δ<sub>s</sub> (‰) is the isotopic composition of soil pore space CO<sub>2</sub> and C<sub>a</sub>  
164 and δ<sub>a</sub> are the CO<sub>2</sub> concentration and isotopic composition of ambient air, respectively. The ambient CO<sub>2</sub> concentrations and  
165 δ<sup>13</sup>C values needed for these corrections were measured in the field at the time of sampling with the CRDS.

166  
167 To investigate the effects of season, warming treatment, and soil depth on ARQ and δ<sup>13</sup>C, we ran multiple regressions in R (R  
168 Development Core Team, 2017). Because ARQ was not sampled from both depths on all dates, we ran separate regressions  
169 for each depth (30 and 90 cm) and then ran a regression that included a depth effect while dropping the first sampling date. In  
170 all regressions, treatment and sampling date (as a factor) were fixed effects. Following Zuur et al. (2009), we used a full model  
171 with all fixed effects and their interactions to optimize the random effects and autocorrelation structure based on AIC. For both  
172 versions, we used the individual gas well as a random effect and a temporal autocorrelation did not improve the model, nor  
173 did an autocorrelation function graph indicate one was needed. We chose the significant fixed effects by performing a series  
174 of pairwise model comparisons using AIC and the *F* test, dropping the least significant variables each time until only variables  
175 that improved the model fit remained. The p-values reported are those from the t-tests of the summary.lme function of best fit  
176 model. We report conditional R<sup>2</sup> values calculated using the rsquared command in the piecewiseSEM package.

177  
178 To investigate seasonal patterns in δ<sup>13</sup>CO<sub>2</sub>, we had more data in terms of both length of time and temporal density of sampling  
179 and were thus able to treat month as a continuous variable. We fit a sine function and tested models including the first and  
180 second harmonics of the month effect as well as linear fixed effects of depth, treatment, and a depth by treatment interaction.  
181 Graphical exploration indicated the sinusoidal pattern differed slightly by year, so we also added a year effect to the second  
182 harmonic of the month effect. As above, we used the full fixed effect model to test the best random and autocorrelation  
183 structure. Individual gas well depth was used as a random effect and a correlation structure did not improve the model.

184  
185 To test relationships between ARQ and δ<sup>13</sup>CO<sub>2</sub>, and both ARQ and δ<sup>13</sup>CO<sub>2</sub> individually versus soil temperature and volumetric  
186 water content, we ran mixed-model regressions with individual gas well as a random effect. For the soil climate relationships,  
187 we used AIC and the *F* test to investigate whether the warming treatment and its interaction with soil temperature or VWC  
188 were significant fixed effects. We tested the need for autocorrelation structures based off of AIC and none improved the  
189 models. For all models, we graphically checked the residuals for violations of normality and heterogeneity of variance. For

190  $\delta^{13}\text{CO}_2$  analyses, we dropped the 15 cm depths due to their unusually low  $\delta^{13}\text{C}$  value ( $<-32\text{‰}$ ) after correction (Eq. 1), which  
191 indicated potential intrusion of atmospheric air during sampling that led to an overcorrection. We used one-way ANOVA's to  
192 compare the ARQ of soil and root incubations and the ARQ of two soil depths we incubated. All statistics were performed in  
193 R v 3.4.1 and regressions were done using the lme function (R Development Core Team, 2017).

### 194 **3 Results**

195 Both ARQ and  $\delta^{13}\text{CO}_2$  had similar, strong seasonal patterns (Fig. 1a and 1b). ARQ values were higher during the growing  
196 season ( $0.89 \pm 0.01$ ,  $n=42$ ) and lower during the winter ( $0.70 \pm 0.02$ ,  $n=23$ ). In ARQ regression analyses for both depths, there  
197 was a significant effect of date ( $p<0.0001$ ,  $n=59$ ) with February 2016 and March 2017 differing significantly from July 2015  
198 (90 cm only), April 2016, August 2016, and June 2017. Similarly,  $\delta^{13}\text{C}$  was higher during the summer (June through October,  
199  $-27.97 \pm 0.06$ ,  $n=311$ ) and lower during the winter and spring (November through May,  $-29.01 \pm 0.04$ ,  $n=447$ ). While individual  
200 dates were not compared statistically for  $\delta^{13}\text{CO}_2$ , the vast improvement in model fit using month as a sine function instead of  
201 a linear function or factor ( $\Delta\text{AIC}=114$ ) is strong statistical evidence for a seasonal effect (Fig. 2b). ARQ and  $\delta^{13}\text{CO}_2$  were  
202 significantly related according to the mixed effect regression model (Fig. 1c,  $p<0.0001$ ,  $n=64$ ,  $R^2=0.20$ ). However, the patterns  
203 in ARQ and  $\delta^{13}\text{CO}_2$  did not match during April.

204

205 Both ARQ and  $\delta^{13}\text{CO}_2$  differed by warming treatment (Fig. 2) and by depth (Table 2). For the ARQ of 30 cm depths, there  
206 was a significant treatment-by-date interaction ( $p=0.051$ ,  $n=30$ ) whereby heated plots had greater ARQ values during the  
207 winter months (February 2016 and March 2017; Fig. 2a). For the ARQ of 90 cm depths, the best fit model did not include a  
208 significant treatment effect or treatment-by-date interaction (Fig. 2a,  $n=35$ ). For  $\delta^{13}\text{CO}_2$  across all depths, treatment was a  
209 significant effect ( $p=0.0065$ ,  $n=758$ ) with warmed soil on average having a slightly higher  $\delta^{13}\text{CO}_2$  ( $-28.33 \pm 0.05$ ) than the  
210 control soil ( $-28.83 \pm 0.06$ ; Fig 2b). The treatment-by-depth interaction was not significant for  $\delta^{13}\text{CO}_2$  and was not included in  
211 the best fit model. Looking at depth only (Table 2), ARQ at 30 cm was marginally significantly greater than ARQ at 90 cm by  
212 0.07 units ( $p=0.099$ ,  $n=59$ ), while  $\delta^{13}\text{CO}_2$  increased with depth from  $-28.98$  at 30 cm to  $-28.34$  at 90 cm ( $p=0.0089$ ,  $n=758$ ).

213

214 Both ARQ and  $\delta^{13}\text{CO}_2$  showed strong relationships with soil climate that were significantly affected by the warming treatment  
215 (Fig 3). We tested relationships with soil temperature and soil moisture individually because of the strong negative correlation  
216 between temperature and moisture in this Mediterranean climate (pearson's  $r=-0.76$  to  $-0.78$ ). ARQ increased significantly  
217 with increasing soil temperatures ( $p<0.0001$ ,  $n=65$ ,  $R^2=0.52$ ; Fig 3a) with values increasing faster in control plots than in  
218 warmed plots ( $p=0.0051$ ). ARQ decreased with increased soil moisture ( $p<0.0001$ ,  $n=60$  due to missing VWC values,  $R^2=0.24$ ;  
219 Fig. 3b), and the decrease was faster in the control than in the warmed plots.  $\delta^{13}\text{CO}_2$  became higher with increasing soil  
220 temperatures ( $p<0.0001$ ,  $n=375$ ,  $R^2=0.33$ ; Fig. 3c) with values again increasing faster in the control than in the warmed plots

221 ( $p=0.02$ ).  $\delta^{13}\text{CO}_2$  decreased with increased soil moisture ( $p<0.0001$ ,  $n=345$  due to missing VWC values,  $R^2=0.30$ ; Fig. 3d),  
222 and treatment did not have a significant effect.

223

224 Our incubations of roots ( $n=4$ ) and of root-free soil ( $n=4$  per depth increment) indicated that heterotrophic and autotrophic  
225 respiration had significantly different ARQ values, at least during the summer when we performed the incubations. Roots had  
226 a greater ARQ ( $0.87 \pm 0.03$ ) than did root-free soil ( $0.78 \pm 0.02$ ; one-way ANOVA,  $p=0.029$ ). Furthermore, ARQ of the soil  
227 incubations significantly declined with depth from  $0.82 \pm 0.01$  at 0-20 cm to  $0.74 \pm 0.02$  at 20-40 cm (one-way ANOVA,  
228  $p=0.0053$ ).

## 229 **4 Discussion**

230 There are many factors that can affect ARQ; however, our evidence indicates the strong seasonal patterns in ARQ and  $\delta^{13}\text{CO}_2$   
231 were likely driven by changes in the amount of root-derived organic substrates providing energy for heterotrophic microbial  
232 respiration and changes in the contributions of autotrophic root respiration. This interpretation is supported by previous soil  
233 ARQ studies, our incubations, and the scientific understanding of how plant carbon inputs change seasonally. The seasonal  
234 range in ARQ from  $\approx 0.9$  during the growing season to  $\approx 0.7$  during the winter may reflect a shift in the molecules fueling  
235 respiration from more oxidized substrates like sugars and organic acids derived from roots in the summer to more reduced  
236 substrates in the winter such as lipids and proteins derived from microbial necromass. Previous incubations found that glucose  
237 additions increased ARQ (Dilly, 2001; Theenhaus et al., 1997). Other studies attributed a decline in ARQ during the time  
238 course of incubation to the depletion of labile carbon sources (Angert et al., 2015; Severinghaus, 1995). Our short-term  
239 incubations demonstrated that root respiration has a greater ARQ than microbial respiration from root-free soils. During the  
240 growing season, root respiration and exudation increase, which should increase ARQ, as seen in our data. In Eastern U.S.  
241 deciduous forests, root exudation rates tend to be lower in the winter and spring than in the summer and fall (Abramoff and  
242 Finzi, 2016; Phillips et al., 2008). Mass-specific fine root respiration rates were greater during the growing season (up to 8  
243  $\text{nmol CO}_2 \text{ g}^{-1} \text{ s}^{-1}$ ) than in the winter ( $<1 \text{ nmol CO}_2 \text{ g}^{-1} \text{ s}^{-1}$ ) and total belowground carbon flux was greatest from May through  
244 October (Abramoff and Finzi, 2016). Though these root studies were not from the western United States, eddy covariance data  
245 from a coniferous forest near our study site found that primary production was greatest during the summer months from June  
246 through mid-September (Goldstein et al., 2000).

247

248 Beyond the results of our root and root-free soil incubations, there is additional evidence that root and rhizosphere respiration  
249 should have a greater ARQ than microbial-derived respiration. For example, respiration of root tips is driven by sugar content  
250 and has an RQ of 1.0 (Saglio and Pradet, 1980). Furthermore, recent metabolomic analysis of root exudates identified sugars,  
251 carboxylic acids, amino acids, and phenolics as the main metabolites (Zhalnina et al., 2018), most of which are relatively  
252 oxidized energy sources with relatively high respiratory quotients. Thus, we would expect greater ARQ values during the



253 summer due to higher root activity. When trees are dormant, the lack of fresh inputs from roots may lead to more recycling of  
254 organic carbon within microbial biomass, wherein proteins and lipids are the first and third largest constituents by weight,  
255 making up to 55% and from 10-35% of a typical bacterial cell's dry mass, respectively (Kleber and Reardon, 2017; Neidhardt,  
256 1987). Lipids and proteins tend to be reduced and have the lowest RQ values of common organic substrates, likely explaining  
257 the lower wintertime ARQ values in our soils.

258

259 The seasonal pattern in  $\delta^{13}\text{CO}_2$  reinforces our interpretation that changes in respiration carbon sources were driving changes  
260 in ARQ. Soil  $\delta^{13}\text{CO}_2$  was more enriched in the summer and became more depleted in the winter by up to 2‰. In a  
261 comprehensive review of carbon isotopes in terrestrial ecosystems, Bowling et al. (2008) showed that plant lipids tend to be  
262 more depleted in  $^{13}\text{C}$  while sugars and organic acids tend to be more enriched in  $^{13}\text{C}$  relative to bulk leaf  $\delta^{13}\text{C}$ . While these  
263 numbers are based on plant lipids, if we assume microbial lipids are similarly depleted relative to other organic compounds,  
264 an increase in microbial necromass as an organic matter source relative to root-derived sources during the winter would cause  
265 the observed fluctuation in  $\delta^{13}\text{CO}_2$ . Furthermore, a chemical fractionation of soil organic matter found that the water-soluble  
266 fraction, which includes sugars, was 3-4‰ more enriched than the acid-insoluble pool (Biasi et al., 2005). While the  
267 interpretation of respiration  $\delta^{13}\text{C}$  by itself in  $\text{C}_3$  ecosystems can be difficult due to the small ‰ differences among carbon  
268 sources (e.g., Bowling et al., 2015), the simultaneous use of ARQ and  $^{13}\text{CO}_2$  helps strengthen interpretations.

269

270 Seasonality encompasses changes to phenology and soil climate, among other factors. Both ARQ and  $\delta^{13}\text{C}$  had significant  
271 positive relationships with soil temperature. In addition to the importance of plant phenology described above, temperature  
272 could have direct effects on respiration sources. Specifically, warmer temperatures can increase root exudation rates (Yin et  
273 al., 2013) and the relative contribution of autotrophic-derived, if not directly autotrophic, respiration to total soil respiration.  
274 In two subarctic ecosystems, warming increased the proportion of ecosystem respiration derived from autotrophs (which, using  
275 natural abundance radiocarbon as a tracer, included heterotrophic respiration of root exudates) relative to heterotrophs (Hicks  
276 Pries et al., 2015). However, temperatures can affect ARQ through more than just changing the contributions of autotrophic  
277 sources. Lower temperatures increase the thermodynamic favorability of the oxidation of reduced carbon in compounds like  
278 lipids (LaRowe and Van Cappellen, 2011), which could also explain the decrease in ARQ values at lower temperatures. For  
279  $\delta^{13}\text{C}$ , it is likely that phenological changes to organic carbon sources were more important than temperature per se. Several  
280 soil incubation studies show that increases in temperature cause respired  $\delta^{13}\text{CO}_2$  to decrease by about 0.12–0.35‰ for each  
281 1°C rise in temperature—the opposite of the relationship we found (Andrews et al., 2000; Biasi et al., 2005; Hicks Pries et al.,  
282 2013). In these incubations, which were devoid of new organic carbon inputs, unlike in situ conditions, the shift was attributed  
283 to changes to the microbial community that affected carbon source preferences (Andrews et al., 2000; Biasi et al., 2005).  
284 Furthermore, in a Mediterranean climate, phloem sap from trees has been shown to become more enriched in  $^{13}\text{C}$  during the  
285 summer (Merchant et al., 2010), matching our pattern in soil  $\delta^{13}\text{CO}_2$ .

286

287 While ARQ and  $\delta^{13}\text{CO}_2$  increased with soil temperature, experimental warming slowed that rate of increase so that both ARQ  
288 and  $\delta^{13}\text{CO}_2$  values were generally greater in the control than in the heated treatment at the warmest soil temperature.  
289 Concurrently, during the colder months, experimental warming caused greater ARQ values (as at 30 cm depths in February  
290 2016 and March 2017) and slightly higher  $\delta^{13}\text{CO}_2$  relative to the controls. The increase in ARQ and  $\delta^{13}\text{CO}_2$  with experimental  
291 warming during the colder soil temperatures of winter indicates proportionately more respiration of relatively oxidized, labile  
292 organic substrates in the heated treatment. Perhaps enhanced root growth and exudation in the heated treatment (Yin et al.,  
293 2013) could result in the increased availability of labile organic substrates, but this increase occurred in winter when trees were  
294 less active and was not seen during the growing season. The increase in ARQ and  $\delta^{13}\text{CO}_2$  could also be the result of preferential  
295 decomposition of more highly oxidized, labile substrates by a more active microbial population during the winter.  
296 Experimental warming increased microbial activity at all soil depths; warming increased  $\text{CO}_2$  production by 34 to 37% overall  
297 with about 40% of the warming response occurring below 15 cm in the soil profile (Hicks Pries et al., 2017). A warming-  
298 induced increase in the consumption of labile substrates could lead to exhaustion of the labile pool and eventually smaller  
299 warming-induced SOC losses as seen at Harvard Forest (Melillo et al., 2002, 2017). In fact, the trend towards decreased ARQ  
300 and  $\delta^{13}\text{CO}_2$  values during the warmer soil temperatures of the growing season could be due to a depletion of the labile SOC  
301 pool during the winter. Another potential explanation for lower values during the growing season could be a reduction in the  
302 proportion of soil respiration derived from roots. In one warming study, root respiration was less sensitivity to warming relative  
303 to heterotrophic respiration (Hartley et al., 2007). The warming treatment dried the soil slightly at Blodgett (Hicks Pries et al.,  
304 2017), which could stress roots during California's essentially rainless growing season. Future measurements of  $\text{CO}_2$   
305 production, ARQ, and  $\delta^{13}\text{CO}_2$  in trenched and untilled plots could help distinguish these possibilities.

306

307 Soil temperature and soil moisture were so strongly negatively correlated due to our study site's Mediterranean climate that it  
308 is difficult to separate their effects. ARQ and  $\delta^{13}\text{CO}_2$  were negatively correlated with volumetric water content, which was  
309 greatest when soil temperatures were coldest. Volumetric water content has the potential to control ARQ in several ways. First,  
310 increased soil moisture reduces  $\text{O}_2$  availability, which could increase ARQ values  $>1$  as  $\text{CO}_2$  is produced without  $\text{O}_2$   
311 consumption. However, during our study the soil remained oxic (soil  $\text{O}_2$  averaged 20% and the minimum was 17.38%). The  
312 negative relationship between ARQ and soil moisture indicates that anaerobic respiration was not a driver, and we only  
313 measured one ARQ value greater than one (1.03) during our study. However, diffusion rates are lower with higher soil  
314 moisture, which could make detection of high ARQ values difficult if anoxic conditions occur within microaggregates. In  
315 anoxic microaggregates, iron (II) is produced anaerobically, which is subsequently oxidized to iron (III) as the aggregate dries  
316 and becomes aerobic, a process that consumes  $\text{O}_2$  without producing  $\text{CO}_2$ , resulting in low ARQ values that can be detected  
317 as drying soils increase diffusion (Angert et al., 2015). In our soils, which tend to contain relatively high amounts of iron  
318 oxides (Rasmussen et al., 2005), iron oxidation could explain the 15% of ARQ values that were less than the reduced organic  
319 matter value of 0.7. Lastly, since  $\text{CO}_2$  is more soluble in water than is  $\text{O}_2$ , more  $\text{CO}_2$  relative to  $\text{O}_2$  is expected to dissolve in  
320 soil water, which would reduce ARQ values at higher moisture contents. However, different dissolution rates and iron

321 oxidation do not fully explain our data as the wide variability in ARQ values (0.44 to 0.94) at high volumetric water contents  
322 (0.27 to 0.31) can be best explained by time of year (Fig. A1), which again points to phenology as the main driver; the greater  
323 ARQ values are from April and June while the lower values are from February and March. Furthermore, there was a stronger  
324 relationship between observed and predicted ARQ in the temperature model than in the soil moisture model.

325

326 Experimental warming affected the relationship between ARQ and soil moisture. ARQ was greater in the heated treatment  
327 when soil moisture was high (winter) and lower in the heated treatment when soil moisture was low (growing season). Soil  
328 water sampled from lysimeters had a greater concentration of dissolved organic carbon in the warming treatment than in the  
329 control (unpublished data), which could deliver oxidized substrates to microbes during the winter rainy season.

330

331 The reasons for  $\delta^{13}\text{CO}_2$  decreasing with increasing volumetric water content are not clear. Based on kinetics, we would expect  
332 that as more  $\text{CO}_2$  dissolves in water, the soil air should become enriched in  $^{13}\text{CO}_2$  because dissolution discriminates against  
333 the heavy isotope and increasingly so at lower temperatures (Zhang et al., 1995), but our data were not consistent with this  
334 explanation. Another possibility is that advective transport of atmospheric  $\text{CO}_2$  through the soil is more likely at lower soil  
335 moisture content. While intrusion of atmospheric  $\text{CO}_2$  would increase the  $\delta^{13}\text{C}$  of soil air, it reduces the effective diffusion  
336 fractionation to  $<4.4\%$ , leading to overcorrected, and thus unrealistically low  $\delta^{13}\text{C}$  values, of which we did have several.

337

338 Depth was the only parameter by which ARQ and  $\delta^{13}\text{CO}_2$  did not change in concert with one another. ARQ decreased with  
339 depth while  $\delta^{13}\text{CO}_2$  increased. The decrease in ARQ with depth, which was more dramatic in the root-free soil incubations  
340 than in soil air (difference of 0.08 versus 0.03), is likely due to decreased plant inputs with fewer fine roots and less root  
341 exudation at depth (Hicks Pries et al., 2018; Tückmantel et al., 2017). The enrichment of soil  $\delta^{13}\text{CO}_2$  likely reflects the near-  
342 universal enrichment of soil organic carbon with depth due to catabolic carboxylation reactions (as microbial byproducts and  
343 necromass become a larger proportion of soil organic matter; Ehleringer et al., 2000; Torn et al., 2002) or the Suess effect (the  
344 continuing depletion of atmospheric  $\text{CO}_2$  over time due to the burning of fossil fuels). In our soils, there was about a 2%  
345 enrichment in bulk soil organic  $\delta^{13}\text{C}$  with depth (Hicks Pries et al., 2018).

## 346 **5 Conclusion**

347 Here we have shown, for the first time, both annual patterns in soil ARQ and how ARQ is affected by experimental warming.  
348 We inferred that seasonal patterns in ARQ were likely due to changes in the dominant substrates providing the energy for soil  
349 respiration with root-derived sugars and organic acids being the dominant substrates during the growing season and microbial  
350 necromass being the dominant substrate during the winter. These inferences of organic substrates were supported by soil  
351  $\delta^{13}\text{CO}_2$  measurements, which showed clear patterns despite our study system containing only  $\text{C}_3$  plants. We caution that direct  
352 experimental evidence of how ARQ changes with sources is needed before our inferences of substrate use can be proven.

353 However, our data indicate ARQ measurements can help to disentangle the biological sources contributing to soil respiration  
354 and to understand how sources are shifting due to global change. This application of ARQ worked well in our soils, which  
355 were well-drained, oxygenated, and lacked carbonates. The interpretation of soil ARQ values becomes more complex if those  
356 conditions are not met (Angert et al., 2015). The autotrophic and heterotrophic source separation in our incubations indicates  
357 ARQ has the potential to be used to partition soil respiration in a similar manner to natural abundance  $\delta^{13}\text{C}$  (e.g., Dorrepaal et  
358 al., 2009; Hicks Pries et al., 2013). To enable further applications of ARQ, more characterization is needed of the controls of  
359 the ratio, including incubation studies of sterile and ‘live’ soils under aerobic and anaerobic conditions and co-located  
360 measurements of ARQ fluxes and the oxidative ratio of organic matter sources as in Masiello et al. (2008). Such future  
361 investigations will help determine whether ARQ deserves a prominent place alongside natural abundance isotopes in the  
362 ecosystem ecology and biogeochemistry toolkit.

### 363 **Code and Data Availability**

364 Data (doi:10.15485/1596312) are publicly available on ESS-DIVE (<http://ess-dive.lbl.gov/>). The R code used for the  
365 statistics and to generate the figures in this paper is available as a supplement.

### 366 **Author Contribution**

367 CHP, AA, and MST conceived of the study. Field measurements were conducted by CHP and CC. Lab analyses were  
368 conducted by CHP and BH. Statistical analyses were conducted by CHP. CHP wrote the manuscript with feedback from all  
369 authors.

### 370 **Acknowledgements**

371 This work was supported as part of the Terrestrial Ecosystem Science Program by the Director, Office of Science, Office of  
372 Biological and Environmental Research, of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. We  
373 would like to acknowledge Rachel Porras for her assistance running the isotopic samples, and Bryan Curtis and Biao Zhu for  
374 their contributions to setting up the warming experiment.

### 375 **References**

- 376 Abramoff, R. Z. and Finzi, A. C.: Seasonality and partitioning of root allocation to rhizosphere soils in a midlatitude forest,  
377 *Ecosphere*, 7(11), n/a-n/a, doi:10.1002/ecs2.1547, 2016.
- 378 Andersen, C. P. and Scagel, C. F.: Nutrient availability alters belowground respiration of ozone-exposed ponderosa pine, *Tree*  
379 *Physiology*, 17(6), 377–387, 1997.

380 Andrews, J. A., Matamala, R., Westover, K. M. and Schlesinger, W. H.: Temperature effects on the diversity of soil  
381 heterotrophs and the  $\delta^{13}\text{C}$  of soil-respired  $\text{CO}_2$ , *Soil Biology and Biochemistry*, 32(5), 699–706,  
382 2000.

383 Angert, A. and Sherer, Y.: Determining the relationship between tree-stem respiration and  $\text{CO}_2$  efflux by  $\delta^{18}\text{O}/\text{Ar}$   
384 measurements, *Rapid Communications in Mass Spectrometry*, 25(12), 1752–1756, doi:10.1002/rcm.5042, 2011.

385 Angert, A., Yakir, D., Rodeghiero, M., Preisler, Y., Davidson, E. A. and Weiner, T.: Using  $\text{O}_2$  to study the relationships  
386 between soil  $\text{CO}_2$  efflux and soil respiration, *Biogeosciences*, 12(7), 2089–2099, doi:10.5194/bg-12-2089-2015, 2015.

387 Biasi, C., Rusalimova, O., Meyer, H., Kaiser, C., Wanek, W., Barsukov, P., Junger, H. and Richter, A.: Temperature-dependent  
388 shift from labile to recalcitrant carbon sources of arctic heterotrophs, *Rapid Communications in Mass Spectrometry*, 19(11),  
389 1401–1408, 2005.

390 Bird, J. A. and Torn, M. S.: Fine roots vs. needles: a comparison of  $^{13}\text{C}$  and  $^{15}\text{N}$  dynamics in a ponderosa pine forest soil,  
391 *Biogeochemistry*, 79(3), 361–382, 2006.

392 Bond-Lamberty, B. and Thomson, A.: A global database of soil respiration data, *Biogeosciences*, 7(6), 1915–1926,  
393 doi:10.5194/bg-7-1915-2010, 2010.

394 Bond-Lamberty, B., Bailey, V. L., Chen, M., Gough, C. M. and Vargas, R.: Globally rising soil heterotrophic respiration over  
395 recent decades, *Nature*, 560(7716), 80, doi:10.1038/s41586-018-0358-x, 2018.

396 Bowling, D. R., Pataki, D. E. and Randerson, J. T.: Carbon isotopes in terrestrial ecosystem pools and  $\text{CO}_2$  fluxes, *New  
397 Phytologist*, 178(1), 24–40, doi:10.1111/j.1469-8137.2007.02342.x, 2008.

398 Bowling, D. R., Egan, J. E., Hall, S. J. and Risk, D. A.: Environmental forcing does not induce diel or synoptic variation in  
399 the carbon isotope content of forest soil respiration, *Biogeosciences*, 12(16), 5143–5160, 2015.

400 Chapman, S. J. and Thurlow, M.: Peat respiration at low temperatures, *Soil Biology and Biochemistry*, 30(8), 1013–1021,  
401 doi:10.1016/S0038-0717(98)00009-1, 1998.

402 Dilly, O.: Microbial respiratory quotient during basal metabolism and after glucose amendment in soils and litter, *Soil Biology  
403 and Biochemistry*, 33(1), 117–127, doi:10.1016/S0038-0717(00)00123-1, 2001.

404 Dorrepaal, E., Toet, S., van Logtestijn, R. S. P., Swart, E., van de Weg, M. J., Callaghan, T. V. and Aerts, R.: Carbon respiration  
405 from subsurface peat accelerated by climate warming in the subarctic, *Nature*, 460(7255), 616–619, doi:10.1038/nature08216,  
406 2009.

407 Ehleringer, J. R., Buchmann, N. and Flanagan, L. B.: Carbon isotope ratios in belowground carbon cycle processes, *Ecological  
408 Applications*, 10(2), 412–422, 2000.

409 Friedlingstein, P., Meinshausen, M., Arora, V. K., Jones, C. D., Anav, A., Liddicoat, S. K. and Knutti, R.: Uncertainties in  
410 CMIP5 Climate Projections due to Carbon Cycle Feedbacks, *J. Climate*, 27(2), 511–526, doi:10.1175/JCLI-D-12-00579.1,  
411 2013.

412 Goldstein, A. H., Hultman, N. E., Fracheboud, J. M., Bauer, M. R., Panek, J. A., Xu, M., Qi, Y., Guenther, A. B. and Baugh,  
413 W.: Effects of climate variability on the carbon dioxide, water, and sensible heat fluxes above a ponderosa pine plantation in

414 the Sierra Nevada (CA), *Agricultural and Forest Meteorology*, 101(2–3), 113–129, doi:10.1016/S0168-1923(99)00168-9,  
415 2000.

416 Hartley, I. P., Heinemeyer, A., Evans, S. P. and Ineson, P.: The effect of soil warming on bulk soil vs. rhizosphere respiration,  
417 *Global Change Biology*, 13(12), 2654–2667, doi:10.1111/j.1365-2486.2007.01454.x, 2007.

418 Hicks Pries, C. E., Schuur, E. A. G. and Crummer, K. G.: Thawing permafrost increases old soil and autotrophic respiration  
419 in tundra: Partitioning ecosystem respiration using  $\delta^{13}\text{C}$  and  $\Delta^{14}\text{C}$ , *Glob Change Biol*, 19(2), 649–661,  
420 doi:10.1111/gcb.12058, 2013.

421 Hicks Pries, C. E., Logtestijn, R. S., Schuur, E. A., Natali, S. M., Cornelissen, J. H., Aerts, R. and Dorrepaal, E.: Decadal  
422 warming causes a consistent and persistent shift from heterotrophic to autotrophic respiration in contrasting permafrost  
423 ecosystems, *Global change biology*, 21(12), 4508–4519, 2015.

424 Hicks Pries, C. E., Castanha, C., Porras, R. C. and Torn, M. S.: The whole-soil carbon flux in response to warming, *Science*,  
425 355(6332), 1420–1423, 2017.

426 Hicks Pries, C. E., Sulman, B. N., West, C., O’Neill, C., Poppleton, E., Porras, R. C., Castanha, C., Zhu, B., Wiedemeier, D.  
427 B. and Torn, M. S.: Root litter decomposition slows with soil depth, *Soil Biology and Biochemistry*, 125, 103–114,  
428 doi:10.1016/j.soilbio.2018.07.002, 2018.

429 Hilman, B. and Angert, A.: Measuring the ratio of  $\text{CO}_2$  efflux to  $\text{O}_2$  influx in tree stem respiration, edited by M. Ryan, *Tree*  
430 *Physiology*, tpw057, doi:10.1093/treephys/tpw057, 2016.

431 Hockaday William C., Gallagher Morgan E., Masiello Caroline A., Baldock Jeffrey A., Iversen Colleen M. and Norby Richard  
432 J.: Forest soil carbon oxidation state and oxidative ratio responses to elevated  $\text{CO}_2$ , *Journal of Geophysical Research:*  
433 *Biogeosciences*, 120(9), 1797–1811, doi:10.1002/2015JG003010, 2015.

434 Hopkins, F. M., Torn, M. S. and Trumbore, S. E.: Warming accelerates decomposition of decades-old carbon in forest soils,  
435 *PNAS*, 109(26), E1753–E1761, doi:10.1073/pnas.1120603109, 2012.

436 Keeling, R. F.: Measuring correlations between atmospheric oxygen and carbon dioxide mole fractions: A preliminary study  
437 in urban air, *J Atmos Chem*, 7(2), 153–176, doi:10.1007/BF00048044, 1988.

438 Keeling, R. F., Piper, S. C. and Heimann, M.: Global and hemispheric  $\text{CO}_2$  sinks deduced from changes in atmospheric  $\text{O}_2$   
439 concentration, *Nature*, 381(6579), 218–221, doi:10.1038/381218a0, 1996.

440 Keiluweit, M., Nico, P. S., Kleber, M. and Fendorf, S.: Are oxygen limitations under recognized regulators of organic carbon  
441 turnover in upland soils?, *Biogeochemistry*, 127(2–3), 157–171, doi:10.1007/s10533-015-0180-6, 2016.

442 Kleber, M. and Reardon, P.: Biopolymers and Macromolecules, in *Encyclopedia of Engineering Geology*, edited by P.  
443 Bobrowsky and B. Marker, pp. 1–5, Springer International Publishing, Cham., 2017.

444 LaRowe, D. E. and Van Cappellen, P.: Degradation of natural organic matter: A thermodynamic analysis, *Geochimica et*  
445 *Cosmochimica Acta*, 75(8), 2030–2042, doi:10.1016/j.gca.2011.01.020, 2011.

446 Liptzin, D., Silver, W. L. and Detto, M.: Temporal Dynamics in Soil Oxygen and Greenhouse Gases in Two Humid Tropical  
447 Forests, *Ecosystems*, 14(2), 171–182, doi:10.1007/s10021-010-9402-x, 2011.

448 Masiello, C. A., Gallagher, M. E., Randerson, J. T., Deco, R. M. and Chadwick, O. A.: Evaluating two experimental approaches  
449 for measuring ecosystem carbon oxidation state and oxidative ratio, *J. Geophys. Res.*, 113(G3), G03010,  
450 doi:10.1029/2007JG000534, 2008.

451 Massman, W. J.: A review of the molecular diffusivities of H<sub>2</sub>O, CO<sub>2</sub>, CH<sub>4</sub>, CO, O<sub>3</sub>, SO<sub>2</sub>, NH<sub>3</sub>, N<sub>2</sub>O, NO, and NO<sub>2</sub> in air,  
452 O<sub>2</sub> and N<sub>2</sub> near STP, *Atmospheric Environment*, 32(6), 1111–1127, 1998.

453 Melillo, J. M., Steudler, P. A., Aber, J. D., Newkirk, K., Lux, H., Bowles, F. P., Catricala, C., Magill, A., Ahrens, T. and  
454 Morrisseau, S.: Soil warming and carbon-cycle feedbacks to the climate system, *Science*, 298(5601), 2173–2176, 2002.

455 Melillo, J. M., Frey, S. D., DeAngelis, K. M., Werner, W. J., Bernard, M. J., Bowles, F. P., Pold, G., Knorr, M. A. and Grandy,  
456 A. S.: Long-term pattern and magnitude of soil carbon feedback to the climate system in a warming world, *Science*, 358(6359),  
457 101–105, doi:10.1126/science.aan2874, 2017.

458 Merchant, A., Tausz, M., Keitel, C. and Adams, M. A.: Relations of sugar composition and  $\delta^{13}\text{C}$  in phloem sap to growth and  
459 physiological performance of *Eucalyptus globulus* (Labill), *Plant, Cell & Environment*, 33(8), 1361–1368, doi:10.1111/j.1365-  
460 3040.2010.02143.x, 2010.

461 Neidhardt, F. C.: Chemical composition of *Escherichia coli*, *Escherichia coli* and *Salmonella typhimurium*: cellular and  
462 molecular biology, 3–6, 1987.

463 Phillips, C. L., Bond-Lamberty, B., Desai, A. R., Lavoie, M., Risk, D., Tang, J., Todd-Brown, K. and Vargas, R.: The value  
464 of soil respiration measurements for interpreting and modeling terrestrial carbon cycling, *Plant Soil*, 413(1–2), 1–25,  
465 doi:10.1007/s11104-016-3084-x, 2017.

466 Phillips, R. P., Ehlitz, Y., Bier, R. and Bernhardt, E. S.: New approach for capturing soluble root exudates in forest soils,  
467 *Functional Ecology*, 22(6), 990–999, doi:10.1111/j.1365-2435.2008.01495.x, 2008.

468 R Development Core Team: R: A language and environment for statistical computing, R Foundation for Statistical Computing,  
469 Vienna, Austria. [online] Available from: <http://www.R-project.org>, 2017.

470 Randerson, J. T., Masiello, C. A., Still, C. J., Rahn, T., Poorter, H. and Field, C. B.: Is carbon within the global terrestrial  
471 biosphere becoming more oxidized? Implications for trends in atmospheric O<sub>2</sub>, *Global Change Biology*, 12(2), 260–271,  
472 doi:10.1111/j.1365-2486.2006.01099.x, 2006.

473 Rasmussen, C., Torn, M. S. and Southard, R. J.: Mineral assemblage and aggregates control carbon dynamics in a California  
474 conifer forest, *Soil Science Society of America Journal*, 69(6), 1711–1721, 2005.

475 Rumble, J.: *CRC Handbook of Chemistry and Physics*, 100th Edition, 100th ed., CRC Press., 2019.

476 Saglio, P. H. and Pradet, A.: Soluble Sugars, Respiration, and Energy Charge during Aging of Excised Maize Root Tips, *Plant*  
477 *Physiology*, 66(3), 516–519, doi:10.1104/pp.66.3.516, 1980.

478 Scagel, C. F. and Andersen, C. P.: Seasonal changes in root and soil respiration of ozone-exposed ponderosa pine (*Pinus*  
479 *ponderosa*) grown in different substrates, *The New Phytologist*, 136(4), 627–643, 1997.

480 Seibt, U., Brand, W. A., Heimann, M., Lloyd, J., Severinghaus, J. P. and Wingate, L.: Observations of O<sub>2</sub>:CO<sub>2</sub> exchange ratios  
481 during ecosystem gas exchange, *Global Biogeochem. Cycles*, 18(4), GB4024, doi:10.1029/2004GB002242, 2004.

482 Severinghaus, J. P.: Studies of the terrestrial O<sub>2</sub> and carbon cycles in sand dune gases and in Biosphere 2, Ph. D. thesis,  
483 Columbia University [online] Available from: <http://ci.nii.ac.jp/naid/10014595764/>, 1995.

484 Theenhaus, A., Maraun, M. and Scheu, S.: Substrate-induced respiration in forest and arable soils measured by O<sub>2</sub>-  
485 microcompensation: moisture conditions and respiratory quotient, *Pedobiologia*, 41(5), 449–455, 1997.

486 Torn, M. S., Lapenis, A. G., Timofeev, A., Fischer, M. L., Babikov, B. V. and Harden, J. W.: Organic carbon and carbon  
487 isotopes in modern and 100-year-old-soil archives of the Russian steppe, *Global Change Biology*, 8(10), 941–953,  
488 doi:10.1046/j.1365-2486.2002.00477.x, 2002.

489 Trumbore, S.: Age of soil organic matter and soil respiration: radiocarbon constraints on belowground C dynamics, *Ecological*  
490 *Applications*, 10(2), 399–411, 2000.

491 Tückmantel, T., Leuschner, C., Preusser, S., Kandeler, E., Angst, G., Mueller, C. W. and Meier, I. C.: Root exudation patterns  
492 in a beech forest: Dependence on soil depth, root morphology, and environment, *Soil Biology and Biochemistry*, 107, 188–  
493 197, doi:10.1016/j.soilbio.2017.01.006, 2017.

494 Worrall, F., Clay, G. D., Masiello, C. A. and Mynheer, G.: Estimating the oxidative ratio of the global terrestrial biosphere  
495 carbon, *Biogeochemistry*, 115(1–3), 23–32, doi:10.1007/s10533-013-9877-6, 2013.

496 Yin, H., Li, Y., Xiao, J., Xu, Z., Cheng, X. and Liu, Q.: Enhanced root exudation stimulates soil nitrogen transformations in a  
497 subalpine coniferous forest under experimental warming, *Global Change Biology*, 19(7), 2158–2167, doi:10.1111/gcb.12161,  
498 2013.

499 Zhalnina, K., Louie, K. B., Hao, Z., Mansoori, N., da Rocha, U. N., Shi, S., Cho, H., Karaoz, U., Loqué, D., Bowen, B. P.,  
500 Firestone, M. K., Northen, T. R. and Brodie, E. L.: Dynamic root exudate chemistry and microbial substrate preferences drive  
501 patterns in rhizosphere microbial community assembly, *Nature Microbiology*, 3(4), 470–480, doi:10.1038/s41564-018-0129-  
502 3, 2018.

503 Zhang, J., Quay, P. D. and Wilbur, D. O.: Carbon isotope fractionation during gas-water exchange and dissolution of CO<sub>2</sub>,  
504 *Geochimica et Cosmochimica Acta*, 59(1), 107–114, doi:10.1016/0016-7037(95)91550-D, 1995.

505 Zuur, A., Ieno, E. N., Walker, N., Saveliev, A. A. and Smith, G. M.: *Mixed Effects Models and Extensions in Ecology with*  
506 *R*, 2009 edition., Springer, New York, NY., 2009.

507

508



<b>Molecule</b>	<b>RQ<sup>a</sup></b>	<b><math>\delta^{13}\text{C}</math> (relative to bulk leaf)<sup>b</sup></b>
Organic acids	1.4 (0.97-4.2)	+0.75
Sugars	1.0	+1.5-2
Phenolics	0.95 (0.92-1.3)	NA
Proteins	0.77 (0.67-1.01)	+1
Lignin	0.88 (0.88-0.94)	-3
Lipids	0.73 (0.68-0.80)	-4

510 <sup>a</sup>Data from Masiello et al. 2008

511 <sup>b</sup>Data from Bowling et al. 2008

512 **Table 1. Respiration quotients (RQ; the inverse of reported oxidative ratios, which are based on elemental analyses) and relative**  
 513 **isotopic enrichment of common molecules/substrates for respiration found in soils. The most common RQ value is listed followed by**  
 514 **the range of potential RQ values in parentheses. The apparent respiration quotient is based on the simultaneous measurement of**  
 515 **soil CO<sub>2</sub> and O<sub>2</sub>.**

516

517

518


519

520

<b>Depth</b>	<b><math>\delta^{13}\text{CO}_2</math></b>	<b>ARQ</b>
<b>(cm)</b>	<b>(‰)</b>	
30	-29.0 ± 0.09 (191)	0.84 ± 0.02 (30)
50	-28.6 ± 0.08 (190)	
70	-28.4 ± 0.07 (191)	
90	-28.3 ± 0.08 (186)	0.81 ± 0.02 (35)

521 **Table 2. The mean ± SE (number of samples) of corrected  $\delta^{13}\text{CO}_2$  and ARQ of soil pore space by depth averaged over all timepoints.**


522



523

524 **Figure 1.** Mean ( $\pm$  SE) apparent respiration quotient (ARQ,  $n=12$  except  $n=6$  for 07/2015; a) and corrected  $\delta^{13}\text{CO}_2$  ( $n=24$  per date;   
 525 b) in soil pore air averaged across all depths and treatments by sampling month. The relationship between ARQ and  $\delta^{13}\text{CO}_2$  values   
 526 over the months when they were sampled simultaneously (c). The line shows the fit of a linear regression ( $p<0.0001$ ,  $n=64$ ,  $R^2=0.20$ ).


527



528

529 **Figure 2. Apparent respiration quotient (ARQ) by sampling date for heated (black) and control (grey) treatments at 30 cm (circles)**  
 530 **and 90 cm (triangles) depths (n=3 per date and depth combination; a). ARQ differed significantly among treatments during the**  
 531 **winter at 30 cm. Corrected  $\delta^{13}\text{CO}_2$  for all depths (30, 50, 70, and 90 cm) and months sampled (b). The lines represent the predicted**  
 532 **fit of a sinusoidal regression (see text) for an average soil depth in control (grey) and heated (black) treatments (n=758).**

533



534  
535

536 **Figure 3. The relationships of apparent respiration quotient (ARQ) by soil temperature (a, n=65) and soil moisture (b, n=60) and**  
 537  **$\delta^{13}\text{CO}_2$  by soil temperature (n=565, c) and soil moisture (n=535, d). Gray and black points represent data from control and heated**  
 538 **gas wells, respectively. The lines show the fit of a mixed model regression between each variable where individual gas well was**  
 539 **treated as a random effect. Separate grey (control) and black (heated) lines indicate there was a significant effect of warming**  
 540 **treatment on the relationship between the response variable and soil temperature.**

541