

1 Dear Dr. De Kauwe:

2 Thank you for the kind consideration. We have carefully checked and revised the manuscript
3 entitled “Soil carbon release responses to long-term versus short-term climatic warming in an
4 arid ecosystem” (bg-2019-236) based on your and the reviewers’ suggestions and comments.
5 We also checked and corrected it carefully for other minor issues such as typos, format style
6 of the journal. The main changes in the newly revised manuscript have been highlighted using
7 red font (the clean version) or the track changes mode (a marked-up manuscript version at the
8 end of this letter). Please see our point-by-point responses as following.

9 **C:** the original comments; **R:** the responses to the comments.

10

11 **Response to Dr. De Kauwe:**

12 **C:** Two reviewers have now seen your revised manuscript. I'm happy to recommend
13 acceptance subject to a few minor suggested revisions from the reviewers.

14 **R:** Thank you for the kind consideration. The manuscript has been carefully revised and
15 corrected accordingly.

16

17 **Response to R1:**

18 **C:** My only commentary is for clarifying the language of the authors’ sections of new text.
19 Please see below for specific examples and suggestions for edits.

20 **R:** Thank you for the kind comments. Yes, we have revised the relevant sections as kindly
21 suggested, and carefully checked and corrected the entire text.

22

23 **Specific Technical Comments:**

24 **C:** Line 12-13: Edit out “such as the changes in” and replace with ‘by altering rates of’ for a
25 stronger opening sentence

26 **R:** This has been done (please see the red words in the new version; lines 12-13).

27

28 **C:** Line 16: Remove “about” for ~3°C

29 **R:** It has been done. Many thanks.

30

31 **C:** Line 17: Remove “about” for ~4°C

32 **R:** It has been done.

33

34 **C:** Line 30: precipitation pulses during the growing season

35 **R:** It has been done.

36

37 **C:** Line 37: of the earth’s

38 **R:** It has been done.

39

40 **C:** Line 62: during the growing season

41 **R:** It has been done.

42

43 **C:** Lines 94-101: A previously published warming experiment is referenced but no explicit
44 mention of those results are given. The new, long-term warming experiment is then stated to
45 possibly have an opposite result, but again, no explicit statement of what that might be. Could
46 the authors clarify this paragraph?

47 **R:** We added the relevant information to be clarified: “A previous study has indicated that the
48 short-term (two-year) warming (2°C) did not affect significantly respiration rate during the
49 growing season” and revised this paragraph accordingly (Lines 94-101 in the newest version).

50

51 **C:** Line 104: Remove “about” for ~3 and ~4°C, or since this is clarified in the Abstract and
52 then outlined in the Methods a few paragraphs later, this detail could be omitted here.

53 **R:** Yes, we agree to delete the information here since this is clarified in the Abstract and then
54 outlined in the Methods later. Many thanks.

55

56 **C:** Line 132: remove long-term as this sentence (line 131 above) started by defining this as the

57 longer warmed plots

58 **R:** Yes, we agree to revise this sentence, and it has been done.

59

60 **C:** Line 137: space needed between regimes (one-year)

61 **R:** It has been done.

62

63 **C:** Line 148: Can remove the sentence defining the control plots as they were already defined

64 in lines 138-139

65 **R:** It has been removed. Thank you.

66

67 Line 150: were installed with a “dummy” heater...

68 **R:** It has been done.

69

70 **C:** Line 274: Respiration significantly increased with...

71 **R:** It has been done.

72

73 **C:** Line 290: end sentence with: from each other ($P= 0.45$)

74 **R:** The phrase has been added.

75

76 **C:** Line 340-342: Awkwardly worded sentence. Rework to state that the positive feedback loop

77 could be weakened with length or intensity of warming

78 **R:** The phrase has been revised as kind suggested (Lines 340-341 in the new version).

79

80 **C:** Line 343: Edit out Actually, and start sentence with Total respiration (Rs)

81 **R:** This has been done.

82

83 **C:** Line 350: add in “and” before consequently the total Rs

84 **R:** Yes, the “and” has been added.

85

86 *C:* Line 378: may rapidly increase

87 *R:* This has been done.

88

89 *C:* Line 382-384: Edit out “at a higher level” as substrate limitation is just another limitation,
90 not necessarily higher (or more important) than moisture, or soil type for instance. Unless you
91 mean higher level as in greater respiration....Please clarify

92 *R:* We agree to edit out the phrase “at a higher level” as kindly suggested.

93

94 *C:* Line 385: high meaning increased respiration?

95 *R:* Yes, the “high” means “increased”, and it was changed accordingly.

96

97 *C:* Line 414: thus affecting its temperature sensitivity; SWC becomes the main factor...

98 *R:* This has been done.

99

100 *C:* Line 419: warming; and – split into 2 full sentences

101 *R:* This has been done.

102

103 *C:* Line 421: implicate that multiple factors together...

104 *R:* This has been done.

105 Many thanks for the constructive suggestions and comments.

106

107 **Response to R2:**

108 *C:* Consider including the hypotheses in the abstract.

109 *R:* The hypotheses could not be necessary in the abstract. However, we revised the expression
110 to be more specific and stronger as kind suggested (bellow).

111

112 *C:* Hypothesis 1 can be more specific. Currently, the hypothesis states that "soil moisture plants

113 a key factor controlling R_s ." This hypothesis would be stronger if the author's specified how
114 soil moisture is expected to control R_s .

115 **R:** Yes, for the hypotheses 1, we changed to "decreased soil moisture plays a key role in
116 reducing R_s ". This can be more specific and stronger (Lines 107-108).

117

118 **C:** I do not necessarily feel you need to include your hypotheses in the abstract that is up to
119 you, but I do think you could revise the hypotheses so that they are more insightful, I think the
120 reviewer makes a good point.

121 **R:** Yes, the hypotheses could not be necessary in the abstract. And we have revised this
122 expression for the hypotheses as kindly suggested.

123 Many thanks for the constructive suggestions and comments.

124

125 Please see a marked-up manuscript version below:

126

127

128 **Soil carbon release responses to long-term versus short-term climatic**
129 **warming in an arid ecosystem**

130

131 Hongying Yu^{1,2}, Zhenzhu Xu^{1,*}, Guangsheng Zhou^{1,3,*}, and Yaohui Shi^{1,3}

132 ¹State Key Laboratory of Vegetation and Environmental Change, Institute of Botany,
133 Chinese Academy of Sciences, Beijing 100093, China

134 ²University of Chinese Academy of Sciences, Beijing, 100049, China

135 ³Chinese Academy of Meteorological Sciences, China Meteorological Administration,
136 Beijing 100081, China

137 *Authors for correspondence

138

139 **Abstract.** Climate change severely impacts grassland carbon cycling by altering
140 rates of litter decomposition and soil respiration (R_s), especially in arid areas
141 steppes. However, little is known about the R_s responses to different warming
142 magnitudes and watering pulses *in situ* in desert steppes. To examine their effects
143 on R_s , we conducted long-term moderate warming (four-year, $\approx 3^\circ\text{C}$), and short-
144 term acute warming (one-year, $\approx 4^\circ\text{C}$), and watering field experiments in a desert
145 grassland of Northern China. While experimental warming significantly reduced
146 average R_s by 32.5% and 40.8% under long-term moderate and short-term acute
147 warming regimes, respectively, watering pulses (fully irrigated the soil to field
148 capacity) stimulated it substantially. This indicates that climatic warming
149 constrains soil carbon release, which is controlled mainly by decreased soil
150 moisture, consequently influencing soil carbon dynamics. Warming did not change
151 the exponential relationship between R_s and soil temperature, whereas the
152 relationship between R_s and soil moisture was better fitted to a sigmoid function.
153 The belowground biomass, soil nutrition, and microbial biomass were not
154 significantly affected by either long-term or short-term warming regimes,
155 respectively. The results of this study highlight the great dependence of soil carbon
156 emission on warming regimes of different durations and the important role of
157 precipitation pulses during the growing season in assessing the terrestrial
158 ecosystem carbon balance and cycle.

159

160 **Key words:** Long-term warming; Precipitation pulse; Soil carbon release;
161 Response sensitivity; Belowground characteristics; Desert grassland.

162

163 **1 Introduction**

164 The global carbon (C) cycle is a critical component of the earth's biogeochemical
165 processes and plays a major role in global warming, which is mainly exacerbated
166 by the elevated carbon dioxide (CO_2) concentration in the atmosphere (e.g.,
167 Falkowski et al., 2000; Carey et al., 2016; Ballantyne et al. 2017; Meyer et al.,

删除了: such as the changes

删除了: around

删除了: around

删除了: in

172 2018). Soil respiration (R_s), mainly including the respiration of live roots and
173 microorganisms, is a key component of the ecosystem C cycle as it releases *c.* 80
174 Pg of C from the pedosphere to the atmosphere annually (Boone et al., 1998; Karhu
175 et al., 2014; Liu et al., 2016; Ma et al., 2014; Schlesinger, 1977). The effects of
176 both soil moisture and temperature on R_s processes and the eco-physiological
177 mechanism are reported extensively; however, it is not well known how soil
178 moisture modulates the response of R_s to changes in the duration and intensity of
179 warming, particularly in arid and semiarid areas, where water and nutrients are
180 both severely limited (e.g., Dacal et al., 2019; Fa et al., 2018; Reynolds et al., 2015;
181 Ru et al., 2018).

182 The desert steppe of China is *c.* 88 million hm^2 , accounting for 22.6% of all
183 grasslands in China, and is located in both arid and semiarid areas. More than 50%
184 of the total area of the steppe is facing severe degradation in terms of the decline
185 of community productivity and soil nutrient depletion, primarily due to improper
186 land use, such as over-grazing and adverse climatic changes, including heat waves
187 and drought stresses (Bao et al., 2010; Kang et al., 2007). Global surface
188 temperature—mainly caused by the anthropogenic CO_2 increase—is expected to
189 increase from 2.6 to 4.8°C by the end of this century, accelerating land degradation
190 (IPCC 2014; 2019). Moreover, the desert steppe ecosystem with low vegetation
191 productivity is vulnerable to its harsh environmental conditions, such as scarce
192 precipitation and barren soil nutrition. For instance, water deficit and heat waves
193 during the growing season can markedly decrease plant cover and productivity in
194 this arid ecosystem (Hou et al., 2013; Luo et al., 2018; Maestre et al., 2012; Yu et
195 al., 2018).

196 Numerous studies have shown that soil temperature and moisture are the two
197 crucial factors that mainly control R_s ; however, it is not well known how soil
198 moisture status mediates the response of R_s to the changes in the duration and
199 intensity of climatic warming. Soil temperature is the primary factor driving
200 temporal R_s variations (e.g., Carey et al., 2016; Gaumont-Guay et al., 2006; Li et
201 al., 2008; Wan et al. 2005). Generally, R_s is significantly and positively correlated
202 with soil temperature when soil moisture is ample (Curiel et al., 2003; Jia et al.,
203 2006; Lin et al., 2011; Reynolds et al., 2015; Yan et al., 2013). In general, the
204 seasonal variations of R_s coincide with the seasonal patterns of soil temperature
205 (Keith et al., 1997; Lin et al., 2011; Wan et al., 2007). For instance, Lin et al. (2011)
206 reported that 63 to 83% of seasonal variations of R_s are dominantly controlled by
207 soil temperature. Diurnal R_s variations are highly associated with variations in soil
208 temperature (Drewitt et al., 2002; Jia et al., 2006; Song et al., 2015). Soil
209 respiration, according to previous studies, is expected to increase with soil water
210 content (SWC) (e.g., Chen et al., 2008; Song et al., 2015; Wan et al., 2007; Yan et
211 al., 2013). However, when the SWC exceeds the optimal point to reach saturated
212 levels, R_s decreases (Huxman et al., 2004; Kwon et al., 2019; Moyano et al., 2012;

213 Moyano et al., 2013; Wang et al., 2014; Yan et al., 2018). In a study conducted in
214 a tall grass prairie, water addition dramatically increased soil CO₂ efflux (Liu et al.,
215 2002). Liu et al. (2009) showed a significant R_s increase after a precipitation pulse
216 in a typical temperate steppe. Therefore, in arid and semiarid regions, where soil
217 water is limited, the SWC may control R_s , and regulate the warming effect (Chen
218 et al., 2008; Curiel et al., 2003; Shen et al., 2015). Furthermore, the effect of
219 watering pulses depends on the pulse size, antecedent soil moisture conditions, soil
220 texture and plant cover (Cable et al., 2008; Chen et al., 2008; Shen et al., 2015;
221 Hoover et al., 2016). For instance, the results by Huxman et al. (2004) showed that
222 different precipitation pulses have different effects on carbon fluxes in these arid
223 and semiarid regions; and Sponseller (2007) indicated that CO₂ efflux increases
224 with storm size in a Sonoran Desert ecosystem.

225 A previous study ~~has indicated that the~~ short-term (two-year) warming (2°C),
226 ~~did not affect significantly respiration rate during the growing season~~ (Liu et al.,
227 2016). However, there is limited information about the long-term (four-year)
228 warming effects on R_s and the underlying mechanism. In this current study, we
229 expect that the long-term (four-year) warming ~~may have more profound effects~~ R_s
230 relative to previous two-year short term, ~~and the underlying mechanism under~~
231 longer term warming condition, and the role of soil water status to R_s responses to
232 climatic warming are also ~~required to be explored further~~. Thus, in the present study,
233 we used a randomized block design with three treatments: control (no warming, no
234 watering), long-term moderate warming (four years extending from 2011 to 2014),
235 and short-term acute warming (one year in 2014). Moreover, a watering pulse
236 treatment (a full irrigation to reach field capacity) was also established. We present
237 the following hypotheses: (i) both long- and short-term climatic warming can
238 reduce soil CO₂ efflux, in which ~~decreased~~ soil moisture ~~plays a key role in~~
239 ~~reducing~~ R_s in the arid ecosystem; and (ii) the dynamics of R_s in the water-limited
240 ecosystem can be driven mainly by the combination of soil temperature and soil
241 moisture, and soil moisture can modulate the response of R_s to warming.

242

243 2 Methods and materials

244 2.1 Experimental site

245 The experiment was conducted in a desert steppe about 13.5 km from Bailingmiao
246 in Damao County (110°19'53.3"E, 41°38'38.3"N; 1409 m above sea level),
247 situated in Nei Mongol, Northern China. This area is characterized by a typical
248 continental climate. The mean annual temperature of this area was 4.3°C with a
249 minimum of -39.4°C and a maximum of 38.1°C from 1955 to 2014. The mean
250 annual precipitation is 256.4 mm and approximately 70% of the annual
251 precipitation is distributed in the growth season period occurring from June to
252 August (Supplementary Figure S1). According to Chinese classification, the soil
253 type is called "chestnut" (Calcic Kastanozems in the FAO soil classification) with

删除了: reported the effects of relatively

删除了:

删除了: on soil respiration

删除了: different effects on R_s (i.e.,

删除了: , even reverse

删除了:)

删除了: uncertain

删除了: , about 3°C

删除了: , about 4°C

删除了: plays a key factor controlling

264 a bulk density of 1.23 g·cm⁻³ and a pH of 7.4. The area has not been grazed since
265 1980; the dominant species is *Stipa tianschanica* var. *klemenzi*, accompanied by
266 *Cleistogenes squarrosa*, *Neopallasia pectinata*, *Erodium stephanianum* and
267 *Artemisia capillaris* (e.g., Hou et al., 2013; Ma et al., 2018).

268 2.2 Experimental design

270 The warming experiment used a randomized block design. The long-term moderate
271 warming plots were exposed to long-term warming from early June to late August
272 (the growing season) for four years (2011–2014), while short-term acute warming
273 was manipulated only during the growing season (June to August) in 2014. The
274 targeted increases in temperatures relative to ambient temperature (control) are
275 around 3°C and 4°C under the long-term moderate warming (four-year), and short-
276 term acute warming regimes (one-year), respectively. Watering pulse treatments
277 were conducted in August in 2014 and 2017. The control plots received no
278 additional treatments of either temperature or water (they were recognized as
279 warming or watering control treatments). All of the warmed plots were heated 24
280 h/day by infrared (IR) lamps (1.0 m long) (GHT220-800; Sanyuan Huahui Electric
281 Light Source Co. Ltd., Beijing, China) at 800 W during growing seasons in the
282 experimental years (2011–2014). The IR lamp heights above the ground were 1.5
283 m and 1.0 m in moderately and acutely warmed plots, respectively. This facility
284 can effectively mimic different climatic warming regimes in field *in situ*, as
285 previously reported (e.g., Hou et al., 2013; Ma et al., 2018; Yu et al., 2018). The
286 watering pulse plots were fully irrigated to field capacity to simulate a watering
287 pulse on August 19, 2014, and August 14, 2017. For the field warming facility, to
288 simulate the shading effects, the control plots were installed with a “dummy”
289 heater similar to those used for the warmed plots. There were a total of 15
290 experimental plots (2 m × 2 m) arranged in a 3 × 5 matrix with each treatment
291 randomly replicated once in each block across three experimental blocks; a 1 m
292 buffer for each adjacent plot was made.

293

294 2.3 Soil temperature and moisture

295 At the center of each plot, a thermocouple (HOBO S-TMB-M006; Onset Computer
296 Corporation, Bourne, MA, USA) was installed at a depth of 5 cm to measure the
297 soil temperature, and a humidity transducer (HOBO S-SMA-M005; Onset
298 Computer Corporation, Bourne, MA, USA) was installed at a depth of 0 to 20 cm
299 to monitor the soil moisture (v/v). Continuous half-hour measurements were
300 recorded by an automatic data logger (HOBO H21-002; Onset Computer
301 Corporation, Bourne, MA, USA).

302

303 2.4 Soil respiration

304 The soil respiration was measured with a Li-8100 soil CO₂ Flux System (LI-COR

删除了: that included three treatments: control (i.e., ambient temperature), long-term moderate warming, short-term acute warming

删除了:

删除了: The neither watering nor warming plots were made as the control plots.

删除了: were designed to

删除了: install a “dummy” heater

313 Inc., Lincoln, NE, USA) with the R_s chamber mounted on polyvinyl chloride (PVC)
314 collars. Fifteen PVC collars (10 cm inside diameter, 5 cm in height) were inserted
315 into the soil 2 to 3 cm below the surface. They were randomly placed into the soil
316 in each plot after clipping all plants growing in the collar placement areas. The
317 collars were initially placed a day before measurements were begun to minimize
318 the influence of soil surface disturbance and root injury on R_s (Bao et al., 2010;
319 Wan et al., 2005). Respirations for the control and all of the warmed plots were
320 measured from 6:00 a.m. to 6:00 p.m. on July 7 and 8 and August 18, 19, 20 and
321 21, 2014. The R_s for watering pulse treatment was measured after the water
322 additions on August 19, 2014, and August 14, 15, 16 and 17, 2017. To stabilize the
323 measurement, R_s was measured only on the selected typical days (i.e., mildly windy,
324 sunny days). The R_s in all plots was measured once every 2 h on that day and each
325 measurement cycle was finished within 30 min to minimize the effects of
326 environmental variables, such as temperature and light. Thus, a total of six
327 measurement cycles was completed each day. The soil water content (SWC, 0–20
328 cm soil depth) in watering plots was measured using the Field Scout TDR 300 Soil
329 Moisture Meter (Spectrum Technologies, Inc., Aurora, IL, USA).

330

331 **2.5 Belowground biomass and related soil characteristics**

332 Soil samples of 0 to 10 cm in depth were taken from each collar after the R_s
333 measurements and then passed through a 1 mm sieve to separate the roots. The
334 roots were washed and oven-dried at 70°C for 48 h to a constant weight and then
335 weighed. Subsamples of each soil sample were separated to determine the
336 gravimetric water content and soil chemical properties. Briefly, to determine the
337 soil organic C (SOC) content, we mixed a 0.5 g soil sample, 5 ml of concentrated
338 sulfuric acid (18.4 mol L⁻¹), and 5.0 ml of aqueous potassium dichromate (K₂Cr₂O₇)
339 (0.8 mol L⁻¹) in a 100 ml test tube, then heated them in a paraffin oil pan at 190°C,
340 keeping them boiling for 5 minutes. After cooling, the 3 drops of phenanthroline
341 indicator were added and then the sample was titrated with ferrous ammonium
342 sulphate (0.2 mol L⁻¹) until the color of the solution changed from brown to purple
343 to dark green (Nelson and Sommers, 1982; Chen et al., 2008; Edwards et al., 2013).
344 The soil ammonium-nitrogen (N) (NH₄⁺-N) concentration and the nitrate-N (NO₃⁻
345 -N) concentration were extracted with a potassium chloride (KCl) solution and
346 measured using a flow injection analyzer (SEAL Auto Analyzer 3; SEAL
347 Analytical, Inc., Mequon, WI, USA) (Liu et al. 2014). Soil samples (0–10 cm in
348 depth) from each collar were oven-dried at 105°C for at least 48 h and weighed to
349 determine the SWC. The soil microbial biomass C (MBC) and microbial biomass
350 N (MBN) were measured using the chloroform-fumigation extraction method and
351 calculated by subtracting extractable C and N contents in the unfumigated samples
352 from those in the fumigated samples (Liu et al., 2014; Rinnan et al., 2009). All
353 extracts were stored at 4°C until further testing commenced.

354

355 2.6 Statistical analysis

356 All statistical analyses were performed using IBM SPSS Statistics 21.0 (IBM,
357 Armonk, NY, USA). All the data were normal as tested by the Shapiro-Wilk
358 method. A one-way analysis of variation (ANOVA) with LSD multiple range tests
359 was conducted to test the statistical significance of the differences in the mean
360 values of the soil temperature, soil moisture, R_s , belowground biomass, SOC,
361 NH_4^+ -N and NO_3^- -N concentrations, and MBC and MBN concentrations at depths
362 of 0 to 10 cm among the different treatments. A linear regression analysis was also
363 used to test the relationship between the SWC and R_s . The relationship between R_s
364 and the soil temperature in each treatment was tested with an exponential function.

365 We used Q_{10} to express the temperature sensitivity of R_s and calculated it
366 according to the following equations:

367

$$368 R_s = ae^{bT_s} \quad (1)$$

369

$$369 Q_{10} = e^{10b} \quad (2)$$

370

371 Here, T_s is the soil temperature, a refers to the intercept of R_s when the soil
372 temperature is 0°C , and b is the temperature coefficient reflecting the temperature
373 sensitivity of R_s and is used to calculate Q_{10} (Lloyd and Taylor, 1994; Luo et al.,
374 2001; Shen et al., 2015).

375 The relationship between R_s and the SWC was further conducted to fit the
376 Gompertz function, a sigmoid function (Gompertz, 1825; Yin et al., 2003), which
377 could express that the linear increase is rapid followed by a leveling off:

378

$$379 R_s = a * e^{-b * (\exp(-k * \text{SWC}))} \quad (3)$$

380

381 Here, a is an asymptote; the SWC halfway point of $a/2$ equals $-\ln(\ln(2)/b)/c$. The
382 turning point of the maximum rate of R_s increase equals ak/e when the SWC equals
383 $\ln(b)/k$. Thus, from the sigmoid function curve, the thresholds of the changes in R_s
384 with increasing SWC can be obtained from the Gompertz function (Gompertz,
385 1825; Yin et al., 2003).

386 A non-linear regression model was used to fit the relationship of R_s with both
387 soil temperature and soil moisture (Savage et al., 2009):

388

$$389 R_s = (R_{\text{ref}} * Q_{10}^{(T_s-10)/10}) * \beta^{(\text{SWC}_{\text{OPT}} - \text{SWC})^2} \quad (4)$$

390

391

392 where T_s is the soil temperature at a soil depth of 5 cm, R_{ref} is R_s at 10°C and Q_{10}
393 is a unitless expression in R_s for each increase in 10°C . SWC is water content in 0

394 to 20 cm soil depth, SWC_{OPT} is the optimal water content and β is a parameter
395 modifying the shape of the quadratic fit.

396 Following the key factors selected by the stepwise regression method, a path
397 analysis was used to examine the primary components directly and indirectly
398 affecting R_s , by integrating both the stepwise linear regression module and Pearson
399 correlation analyses (Gefen et al., 2000). The statistical significances were set at P
400 < 0.05 for all tests, unless otherwise indicated.

401

402 3 Results

403 3.1 Warming effects on belowground characteristics

404 The soil temperatures at a soil depth of 5 cm in the warmed plots were much higher
405 than those in the control plots (Figure 1). During growing season, the mean soil
406 temperatures in the control, the moderately and acutely warmed plots were 21.9°C
407 (± 0.13 SE), 24.5°C (± 0.15), and 25.0°C (± 0.18), respectively. The moderately and
408 acutely warmed plots were respectively increased by 2.6°C ($P < 0.001$) and 3.1°C
409 ($P < 0.001$) compared to those in the control plots. The SWC in the moderately and
410 acutely warmed plots (0–20 cm soil profile, defined as ratios of water volume and
411 soil volume) were significantly reduced ($P < 0.001$) compared to those in the
412 control plots (Figure 1), indicating that warming led to marked declines in the SWC,
413 consequently enhancing drought stress. On August 18, 19, 20 and 21, which were
414 the dates that we measured R_s , the daily soil temperatures in the moderately and
415 acutely warmed plots were around 3°C and 4°C higher than those in the control
416 plots, respectively. All belowground variables (belowground biomass, soil N and
417 microbial characteristics) were not significantly altered by warming regimes at the
418 site of this experiment (Supplementary Table S1; $P > 0.05$). However, the organic
419 soil carbon content tended to decrease with long-term climatic warming.

420

421 3.2 Watering pulse effects on R_s

422 Soil respiration, significantly increased with SWC both linearly ($R^2 = 0.83$; $P <$
423 0.01) and quadratically ($R^2 = 0.88$; $P < 0.01$, Figure 2A). Moreover, the Gompertz
424 function was well fitted to their relationship ($R^2 = 0.87$; RMSE = 4.88) (Figure 2B).
425 From the Gompertz functional curve, the R_s asymptote value, as an estimated
426 maximum, was $3.76 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ when the optimal SWC was 22.85%. In the
427 watering plots, an exponential function was well fitted to the relationship between
428 soil respiration and the soil temperatures ($R^2 = 0.31$; $P < 0.01$), with a temperature
429 sensitivity (Q_{10}) of 1.69. However, the exponential function was not well fitted in
430 the control plots (Figure 3A).

431

432 3.3 Effects of warming regimes on R_s

433 Warming regimes resulted in marked declines in R_s . Whereas no difference in R_s
434 was observed in July, during August average R_s values were 1.57, 1.06, and 0.93

删除了: The R_s

436 $\mu\cdot\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ in the control, moderately warmed and acutely warmed plots,
437 respectively, indicating that warming regimes resulted in marked declines (Figure
438 4). Changes in R_s differed significantly between the control and both warmed plots
439 ($P < 0.01$), while the R_s in the two warmed plots did not significantly differ from
440 each other ($P = 0.45$). The relationships between the R_s and soil temperature of
441 each treatment were well fitted by the exponential equations ($P < 0.05$) (Figure
442 3B). The Q_{10} values were 1.88, 2.12 and 1.58 in the temperature controlled,
443 moderate and acute warming treatments, respectively (Figure 3B). It indicated that
444 R_s increases exponentially with temperature in watered plots but was lower and
445 insensitive to temperature in the control plots (Figure 3A); and that long-term
446 warming rather than temporary high temperature reduced R_s , despite having a
447 positive relationship with soil temperature (Figure 3B, 4).

448

449 **3.4 Interactive effects on R_s from soil temperature and soil water content**

450 Across all watering and warming treatments, generally, a high temperature led to
451 an increase in R_s under ample soil moisture, whereas R_s was limited under a soil
452 water deficit. As shown in Figure 5, A non-linear regression model (equation 4)
453 was well fitted to the relationship of R_s with both soil temperature and soil moisture
454 in the control plots ($R^2 = 0.40$, RMSE = 0.60). Based on the function $R_s =$
455 $(0.733*1.796^{(T_s-10)/10})*\beta^{(0.229-SWC)^2}$, the key parameters were obtained: R_{ref} , a R_s at
456 10°C , was $0.73 \mu\cdot\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$; Q_{10} , a unitless expression in R_s for each increase in
457 10°C , was 1.80; and β , a parameter modifying the shape of the quadratic fit, was
458 0.001 (Figure 5).

459

460 **3.5 Effects of multiple factors on R_s : a path analysis**

461 Based on a stepwise regression analysis of the relationships between the R_s and
462 multiple factors, four key factors were screened: soil temperature, soil moisture,
463 belowground biomass and SOC. Their effects on R_s were further determined by a
464 path analysis. The results showed that soil moisture and soil temperature were two
465 major direct factors controlling R_s (the two direct path coefficients were 0.72 and
466 0.55, respectively). SOC had the highest indirect effect on R_s (the indirect path
467 coefficient was 0.57). Soil moisture highly correlated with R_s ($R = 0.78$, $P < 0.01$;
468 Supplementary Table S2, Figure 6), indicating again that the soil water status may
469 impose the greatest effect on the carbon release from soil in the desert grassland.

470

471 **4. Discussion**

472 **4.1 Warming effects on R_s**

473 Previous studies have shown positive R_s responses to increased soil temperatures
474 below a critical high temperature (e.g., Carey et al., 2016; Drewitt et al., 2002;
475 Gaumont-Guay et al., 2006; Meyer et al., 2018). However, in the current study site,
476 the climatic warming finally reduced the average R_s by 32.5% and 40.8% under

477 long-term versus short-term climatic warming conditions in the desert dryland,
 478 respectively, which chiefly confirmed our first hypothesis. In a semiarid grassland
 479 on the Loess Plateau of China, the total R_s was also constrained substantially by a
 480 field manipulative experiment (Fang et al., 2018). This result may have been
 481 caused by the following factors. First, high temperatures may cause thermal stress
 482 on microbes and subsequently reduce microbial respiration (i.e., heterotrophic, R_h ,
 483 Chang et al., 2012; Dacal et al., 2019). For instance, in an alpine steppe on the
 484 Tibetan Plateau, microbial respiration was significantly reduced when the
 485 temperature rose to 30°C (Chang et al., 2012). Second, in the desert grassland,
 486 where water is often limited, the SWC becomes the primary factor affecting R_s
 487 (Supplementary Table S2; Figure 6), while warming can cause greater
 488 evapotranspiration, consequently lessening soil moisture (Figure 1), and finally
 489 reducing R_s (Munson et al., 2009; Wan et al., 2007; Yan et al., 2013). The decreases
 490 in average R_s with warming implicate that the positive feedback loop could be
 491 weakened with length or intensity of warming.
 492 Total respiration (R_s), [the sum of root (autotrophic, R_a) and R_h respiration—the
 493 former accounting for *c.* 22 % of the total R_s in the ecosystem, Liu et al., 2016]
 494 may acclimatize to warming within an appropriate range of temperature change at
 495 an ample soil moisture; however, it decreases with increasing temperatures above
 496 an optimum level. The mechanisms may include: within an appropriate range of
 497 temperature change at an ample soil moisture, climatic warming can enhance both
 498 plant root (Luo et al., 2001; Liu et al., 2016) and microbial activities (Tucker et al.,
 499 2014), leading to increases in both R_a and R_h , and consequently the R_s (Luo et al.,
 500 2001; Tucker et al., 2014; Xu et al., 2019). However, when warming continues or
 501 with increasing temperatures above an optimum level, the root growth can be
 502 constrained, directly reducing R_a (Carey et al., 2017; Liu et al., 2016; Luo et al.,
 503 2001; Wan et al., 2007); and the limitation to microbial activities may also occur
 504 (Tucker et al., 2013; Yu et al., 2018), decreasing the R_h (Bérard et al., 2011; Tucker
 505 et al., 2013; Bérard et al., 2015; Romero-Olivares et al., 2017). In addition,
 506 decreases in soil enzyme pools and its activity under warming may also contribute
 507 to a reduction in R_h (e.g., Alvarez et al., 2018). Further, R_s decreases with warming
 508 under water deficit (Moyano et al., 2013; Wang et al., 2014; and see below).
 509 Together, the declines in both root and microbial respirations finally reduce the R_s .
 510 Nevertheless, the drastic declines in R_s under both long-term and short-term
 511 climatic warming regimes in the desert dryland ecosystem may be driven by
 512 multiple factors, including the ecosystem type, time and soil features (Liu et al.,
 513 2016; Wan et al., 2007; Meyer et al., 2018; Thakur et al., 2019). It implies that the
 514 effects of multiple factors should be considered in assessing the carbon balance
 515 between ecosystem and atmosphere.

516

517 4.2 Interactive effect of soil water status and temperature

删除了: positive feedback

删除了: on climatic change may be weakened with warming time or under an acute warming condition

删除了: Actually,

设置了格式: 字体: 倾斜

设置了格式: 下标

删除了: the R_s

523 As stated above, in an arid ecosystem, soil water deficit is a primary factor
524 inhibiting soil carbon release (Supplementary Table S2; Figure 6; Liu et al., 2016;
525 Munson et al., 2009; Yan et al., 2013). Thus, R_s linearly increases with increasing
526 soil moisture. However, it could be leveled off or decreased when soil moisture
527 exceeds an optimal level for the soil carbon release (Huxman et al., 2004; Moyano
528 et al., 2013; Wang et al., 2014). Thus, the relationship between R_s and SWC may
529 be well fitted to the Gompertz functional curve model, a sigmoid function
530 (Gompertz, 1825; Yin et al., 2003), which can be confirmed by the present results
531 in the native arid desert ecosystem (Figure 2). The mechanisms mainly are: an
532 increase in SWC may rapidly increase microbial activities (Cable et al., 2008;
533 Meisner et al., 2015; Wu & Lee, 2011), and enhance root growth (Xu et al., 2014),
534 leading to a linear increase in R_s . However, when soil moisture reaches an ample
535 level, microbial activities may also reach a maximum where the limiting effects of
536 substrate occur (Skopp et al., 1990), finally maintaining a stable change in R_s .
537 Similar response to watering appears for root growth (Xu et al., 2014), and also
538 similarly leading to a stable change in R_s . Thus, R_s can be leveled off at an increased
539 and stable level. Moreover, the decrease in R_s at a saturated SWC level may be
540 ascribed to inhibitions of both root systems and microbial activities under the
541 anaerobic environment (Drew 1997; Huxman et al., 2004; Kwon et al., 2019;
542 Sánchez-Rodríguez et al. 2019; Yan et al., 2018). The model concerning the
543 relationship R_s with a broad range of SWC is helpful to assess and predict the
544 dynamics in soil carbon release in natural arid ecosystems.

545 As indicated by Tucker and Reed (2016), soil water deficit can shrink the R_s
546 itself and its response to temperature, suggesting the changes in R_s may be
547 determined simultaneously by both soil temperature and water status (Janssens et
548 al., 2001; Yan et al., 2013; Sierra et al., 2015). Moreover, in the present experiment,
549 the interactive effects of both factors were tested based on the relationship of R_s
550 with both soil temperature and soil moisture in a non-linear regression model
551 (Savage et al., 2009). The model utilized was well fitted but marginally so ($R^2 =$
552 0.40 , $RMSE = 0.596$; Figure 5), indicating that both the soil temperature and soil
553 water content coordinated the changes in R_s . However, this interaction may also be
554 affected simultaneously by other abiotic and biotic factors, such as soil nutrition
555 availability and soil microbe activity (e.g., Camenzind et al., 2018; Karhu et al.,
556 2014; Thakur et al., 2019; Zhang et al., 2014).

557

558 4.3 Key factors and the influence path

559 As noted above, R_s is affected by several abiotic and biotic factors. The current
560 results showed that soil moisture and soil temperature were two major direct
561 factors, and SOC only was an indirect factor controlling R_s (Supplementary Table
562 S2, Figure 6). Importantly, soil moisture, with both the highest direct path
563 coefficients (0.7) and correlation coefficient (0.8) for R_s , may become the most

删除了: rapidly

删除了: at a higher level

删除了: at a higher level

删除了: high

568 important factor affecting R_s in this desert steppe. These findings agree with the
569 previous results: improved soil water status had a significantly positive effect on
570 R_s (e.g., Chen et al., 2008; Liu et al., 2016; Xu et al., 2016). Furthermore, the soil
571 moisture condition can mediate the relationship between soil temperature and R_s ,
572 thus affecting its temperature sensitivity; SWC becomes the main factor
573 controlling R_s , especially in arid ecosystems, such as desert steppes, where the
574 available soil water is limited (Conant et al., 2000; Curiel et al., 2003; Fa et al.,
575 2018; Jassal et al., 2008; Roby et al., 2019). Thus, under both the long-term and
576 short-term climatic warming regimes, soil moisture could modulate the response
577 of R_s to warming. The changes in R_s might be driven by both soil temperature and
578 soil moisture as two key factors, and SOC as an indirect factor, thus mostly
579 confirming our second hypothesis. The findings again implicate that multiple
580 factors together coordinate R_s , and provide new insight into how to control soil
581 carbon release in arid ecosystems. The models on the R_s changes should consider
582 multiple-factor effects of soil carbon dynamics when assessing and predicting
583 carbon cycle, and its climate feedback.

584

585 4.4 Warming effects on the variables belowground

586 Elevated temperature has been shown to increase or decrease root productivity and
587 biomass, depending on experimental sites and vegetation types (Bai et al., 2010;
588 Fan et al., 2009; Litton and Giardina, 2008; Wan et al., 2004). The decreased
589 availability of soil nutrients apparently limits root growth, finally inducing root
590 mortality and weakening responses to the elevated temperature (Eissenstat et al.,
591 2000; Johnson et al., 2006; Wan et al., 2004; Zhang et al., 2014). In our experiment,
592 no significantly different changes occurred in either soil $\text{NH}_4^+\text{-N}$ or $\text{NO}_3^-\text{-N}$
593 concentrations among the three treatments (Supplementary Table S1), and these
594 might be linked to the non-significant response of belowground biomass to
595 increasing temperature. Microbial biomass and its activities in soil depend on the
596 root biomass, SWC and soil N conditions (Liu et al., 2014; Rinnan et al., 2007;
597 Zhang et al., 2008; Zhang et al., 2014). Warming regimes had no significant effects
598 on either MBC or MBN in the current study (Supplementary Table S1), which
599 might be due to the lack of any difference in the changes in basic soil nutrition
600 status, such as the N conditions, among the three warming treatments. This result
601 is consistent with that of Zhang et al. (2005) and Liu et al. (2015). Moreover, in
602 the present study, SOC concentrations were not significantly affected by climatic
603 warming (Supplementary Table S1), which is inconsistent with the findings of
604 previous studies (Jobbágy and Jackson, 2000; Prietzel et al., 2016). However, there
605 might be a decreasing trend evident with long-term warming. For instance,
606 Crowther et al. (2016) reported a loss of approximately 30 ± 30 Pg of C in the
607 upper soil horizons at 1°C warming in global soil C stocks and projected a loss of
608 203 ± 161 Pg of C under 1°C of warming over 35 years. The C losses from soil

删除了: the

删除了: it

删除了: key

删除了: ;

删除了: and t

删除了: the

删除了: may

616 moving into the atmosphere may result in positive feedback regarding global
617 warming (Bradford et al., 2016; Dacal et al., 2019; Jenkinson et al., 1991; Liu et
618 al., 2016). However, SOC exerted an indirect effect via a path analysis (Figure 6).
619 For this difference, therefore, more evidence needs to be provided to address the
620 issue (Xu et al., 2019).

621 In conclusion, we determined the responses of R_s to field experimental long-
622 term versus short-term climatic warming and watering pulses in a desert steppe
623 ecosystem. We found the following: i) both long- and short-term warming
624 significantly reduced R_s during the peak growth season; ii) soil moisture was the
625 main factor controlling R_s in desert grassland; iii) R_s was significantly and
626 exponentially increased with soil temperature, meanwhile soil moisture condition
627 can mediate the relationship between soil temperature and R_s , thus affecting its
628 temperature sensitivity; and iv) belowground biomass, soil nutrition variables and
629 soil microbial characteristics showed no significant changes after either long-term
630 or short-term climatic warming. These findings may be useful to assess and predict
631 dynamics of soil CO₂ fluxes, particularly the feedback of warming to climatic
632 change, and finally optimize C management work in arid and semiarid regions
633 under the changing climate. However, the patterns of the changes in soil C fluxes
634 and the underlying mechanism in response to climatic change are markedly
635 complicated at various spatial-temporal scales during growing season—from site
636 and regional to global scales, and from daily, seasonal and yearly to decade scales—
637 —and still need to be investigated further (e.g., Ballantyne et al., 2017; Dacal et al.,
638 2019; Meyer et al., 2018; Romero-Olivares et al., 2017).

639

640 **Data availability.** The final derived data presented in this study are available at
641 <https://doi.org/10.5281/zenodo.3546062> (Yu et al., 2019).

642

643 **Supplement.** The supplement related to this article is available online at:

644

645 **Author contributions.** ZX and GZ conceived and designed this study. HY, ZX and
646 YS conducted this experiment and analysed the data. All authors wrote and
647 proofread this manuscript.

648

649 **Competing interests.** The authors declare that they have no conflict of interest.

650

651 **Acknowledgements.** This research was jointly funded by National Natural
652 Science Foundation of China (31661143028, 41775108), and the Special Fund for
653 Meteorological Scientific Research in the Public Interest (GYHY201506001-3).
654 We greatly thank Feng Zhang, Yuhui Wang, Bingrui Jia, Hui Wang, Minzheng
655 Wang, He Song for their loyal help during the present study. The authors also
656 greatly appreciate Dr. De Kauwe and the reviewers for their constructive comments.

657

658 **References**

- 659 Alvarez, G., Shahzad, T., Andanson, L., Bahn, M., Wallenstein, M.D., and Fontaine, S.:
660 Catalytic power of enzymes decreases with temperature: New insights for understanding
661 soil C cycling and microbial ecology under warming. *Glob. Change Biol.*, 24, 4238–4250,
662 <https://doi.org/10.1111/gcb.14281>, 2018.
- 663 Bai, W., Wan, S., Niu, S., Liu, W., Chen, Q., Wang, Q., Zhang, W., Han, X., and Li, L.:
664 Increased temperature and precipitation interact to affect root production, mortality, and
665 turnover in a temperate steppe: implications for ecosystem C cycling. *Glob. Change Biol.*,
666 16, 1306–1316, <https://doi.org/10.1111/j.1365-2486.2009.02019.x>, 2010.
- 667 Ballantyne, A., Smith, W., Anderegg, W., Kauppi, P., Sarmiento, J., Tans, P., Shevliakova, E.,
668 Pan, Y., Poulter, B., Anav, A., and Friedlingstein, P.: Accelerating net terrestrial carbon
669 uptake during the warming hiatus due to reduced respiration. *Nature Clim. Change*, 7,
670 148–152, <https://doi.org/10.1038/nclimate3204>, 2017.
- 671 Bao, F., Zhou, G. S., Wang, F. Y., and Sui, X. H.: Partitioning soil respiration in a temperate
672 desert steppe in Inner Mongolia using exponential regression method. *Soil Biol. Biochem.*,
673 42, 2339–2341, <https://doi.org/10.1016/j.soilbio.2010.08.033>, 2010.
- 674 Bérard, A., Bouchet, T., Sévenier, G., Pablo, A.L., and Gros, R.: Resilience of soil microbial
675 communities impacted by severe drought and high temperature in the context of
676 Mediterranean heat waves. *Eur. J. Soil Biol.* 47, 333–342,
677 <https://doi.org/10.1016/j.ejsobi.2011.08.004>, 2011.
- 678 Bérard, A., Sassi, M.B., Kaisermann, A., and Renault, P.: Soil microbial community responses
679 to heat wave components: drought and high temperature. *Clim. Res.* 66, 243–264,
680 <https://doi.org/10.3354/cr01343>, 2015.
- 681 Boone, R. D., Nadelhoffer, K. J., Canary, J. D., and Kaye, J. P.: Roots exert a strong influence
682 on the temperature sensitivity of soil respiration. *Nature*, 396, 570–572,
683 <https://doi.org/10.1038/25119>, 1998.
- 684 Bradford, M. A., Wieder, W. R., Bonan, G. B., Fierer, N., Raymond, P. A., and Crowther, T.
685 W.: Managing uncertainty in soil carbon feedbacks to climate change. *Nature Clim.*
686 *Change*, 6, 751–758, <https://doi.org/10.1038/nclimate3071>, 2016.
- 687 Cable, J. M., Ogle, K., Williams, D. G., Weltzin, J. F., and Huxman, T. E.: Soil texture drives
688 responses of soil respiration to precipitation pulses in the Sonoran Desert: Implications
689 for climate change. *Ecosystems*, 11, 961–979, [https://doi.org/10.1007/s10021-008-9172-](https://doi.org/10.1007/s10021-008-9172-x)
690 [x](https://doi.org/10.1007/s10021-008-9172-x), 2008.
- 691 Camenzind, T., Hättenschwiler, S., Treseder, K. K., Lehmann, A., and Rillig, M. C.: Nutrient
692 limitation of soil microbial processes in tropical forests. *Ecol., Monogr.*, 88, 4–21,
693 <https://doi.org/10.1002/ecm.1279>, 2018.
- 694 Carey, J. C., Tang, J., Templer, P. H., Kroeger, K. D., Crowther, T. W., Burton, A. J., Dukes, J.
695 S., Emmett, B., Frey, S. D., Heskell, M. A., and Jiang, L.: Temperature response of soil
696 respiration largely unaltered with experimental warming. *P. Natl. Acad. Sci. USA*, 113,
697 13797–13802, <https://doi.org/10.1073/pnas.1605365113>, 2016.
- 698 Chang, X., Wang, S., Luo, C., Zhang, Z., Duan, J., Zhu, X., Lin, Q., and Xu, B.: Responses of
699 soil microbial respiration to thermal stress in alpine steppe on the Tibetan plateau. *Euro.*
700 *J. Soil Sci.*, 63, 325–331, <https://doi.org/10.1111/j.1365-2389.2012.01441.x>, 2012.
- 701 Chen, S. P., Lin, G. H., Huang, J. H., and He, M.: Responses of soil respiration to simulated
702 precipitation pulses in semiarid steppe under different grazing regimes. *J. Plant Ecol.*, 1,
703 237–246, <https://doi.org/10.1093/jpe/rtn020>, 2008.

704 Conant, R. T., Klopatek, J. M., and Klopatek, C. C.: Environmental factors controlling soil
705 respiration in three semiarid ecosystems. *Soil Sci. Soc. Am. J.*, 64(1), 383–390,
706 <https://doi.org/10.2136/sssaj2000.641383x>, 2000.

707 Crowther, T. W., Todd-Brown, K. E., Rowe, C. W., Wieder, W. R., Carey, J. C., Machmuller,
708 M. B., Snoek, B. L., Fang, S., Zhou, G., Allison, S. D., and Blair, J. M.: Quantifying
709 global soil carbon losses in response to warming. *Nature*, 540, 104–108,
710 <https://doi.org/10.1038/nature20150>, 2016.

711 Curiel, J. C., Janssens, I. A., Carrara, A., Meiresonne, L., and Ceulemans, R.: Interactive effects
712 of temperature and precipitation on soil respiration in a temperate maritime pine
713 forest. *Tree Physiol.*, 23, 1263–1270, <https://doi.org/10.1093/treephys/23.18.1263>, 2003.

714 Dacal, M., Bradford, M. A., Plaza, C., Maestre, F. T., and García-Palacios, P.: Soil microbial
715 respiration adapts to ambient temperature in global drylands. *Nat. Ecol. Evol.*, 3, 232–
716 238, <https://doi.org/10.1038/s41559-018-0770-5>, 2019.

717 Drew, M.C.: Oxygen deficiency and root metabolism: injury and acclimation under hypoxia
718 and anoxia. *Annu. Rev. Plant Biol.*, 48, 223–250,
719 <https://doi.org/10.1146/annurev.arplant.48.1.223>, 1997.

720 Drewitt, G. B., Black, T. A., Nestic, Z., Humphreys, E. R., Jork, E. M., Swanson, R., Ethier, G.
721 J., Griffis, T., and Morgenstern, K.: Measuring forest floor CO₂ fluxes in a Douglas-fir
722 forest. *Agric. For. Meteorol.*, 110, 299–317, [https://doi.org/10.1016/S0168-](https://doi.org/10.1016/S0168-1923(01)00294-5)
723 [1923\(01\)00294-5](https://doi.org/10.1016/S0168-1923(01)00294-5), 2002.

724 Edwards, K. A. and Jfferies, R. L.: Inter-annual and seasonal dynamics of soil microbial
725 biomass and nutrients in wet and dry low-Arctic sedge meadows. *Soil Biol. Biochem.*, 57,
726 83–90, <https://doi.org/10.1016/j.soilbio.2012.07.018>, 2013.

727 Eissenstat, D. M., Wells, C. E., Yanai, R. D., and Whitbeck, J. L.: Research view: Building roots
728 in a changing environment: Implications for root longevity. *New Phytol.*, 147, 33–42.
729 2000.

730 Fa, K., Zhang, Y., Lei, G., Wu, B., Qin, S., Liu, J., Feng, W., and Lai, Z.: Underestimation of
731 soil respiration in a desert ecosystem. *Catena*, 162, 23–28,
732 <https://doi.org/10.1016/j.catena.2017.11.019>, 2018.

733 Falkowski, P., Scholes, R. J., Boyle, E. E. A., Canadell, J., Canfield, D., Elser, J., Gruber, N.,
734 Hibbard, K., Högberg, P., Linder, S., and Mackenzie, F. T.: The global carbon cycle: a test
735 of our knowledge of earth as a system. *Science*, 290, 291–296,
736 <https://doi.org/10.1126/science.290.5490.291>, 2000.

737 Fan, J. W., Wang, K., Harris, W., Zhong, H. P., Hu, Z. M., Han, B., Zhang, W. Y., and Wang, J.
738 B.: Allocation of vegetation biomass across a climate-related gradient in the grasslands
739 of Inner Mongolia. *J. Arid Environ.*, 73, 521–528,
740 <https://doi.org/10.1016/j.jaridenv.2008.12.004>, 2009.

741 Fang, C., Li, F., Pei, J., Ren, J., Gong, Y., Yuan, Z., Ke, W., Zheng, Y., Bai, X., and Ye, J. S.:
742 Impacts of warming and nitrogen addition on soil autotrophic and heterotrophic
743 respiration in a semi-arid environment. *Agric. For. Meteorol.*, 248, 449–457,
744 <https://doi.org/10.1016/j.agrformet.2017.10.032>, 2018.

745 Gaumont-Guay, D., Black, T. A., Griffis, T. J., Barr, A. G., Jassal, R. S., and Nestic, Z.:
746 Interpreting the dependence of soil respiration on soil temperature and water content in a
747 boreal aspen stand. *Agric. For. Meteorol.*, 140, 220–235,
748 <https://doi.org/10.1016/j.agrformet.2006.08.003>, 2006.

749 Gefen, D., Straub, D., and Boudreau, M. C.: Structural equation modelling and regression:
750 Guidelines for research practice. *Communications of the Association for Information*

751 Systems, 4: 7. <http://doi.org/10.17705/1CAIS.00407>, 2000.

752 Gompertz, B.: On the nature of the function expressive of the law of human mortality, and on
753 a new mode of determining the value of life contingencies. *Philos. TR. Soc. London*, 115,
754 513–583, <https://doi.org/10.1098/rstl.1825.0026>, 1825.

755 Han, G. X., Zhou, G. S., Xu, Z. Z., Yang, Y., Liu, J. L., and Shi, K.Q.: Soil temperature and
756 biotic factors drive the seasonal variation of soil respiration in a maize (*Zea mays* L.)
757 agricultural ecosystem. *Plant Soil*, 291, 15–26, [https://doi.org/10.1007/s11104-006-9170-](https://doi.org/10.1007/s11104-006-9170-8)
758 [8](https://doi.org/10.1007/s11104-006-9170-8), 2006.

759 Hoover, D. L., Knapp, A. K., and Smith, M. D.: The immediate and prolonged effects of
760 climate extremes on soil respiration in a mesic grassland. *J. Geophys. Res.-Biogeosci.*,
761 121, 1034–1044, <http://dx.doi.org/10.1002/2015JG003256>, 2016.

762 Hou, Y. H., Zhou, G. S., Xu, Z. Z., Liu, T., and Zhang, X. S.: Interactive effects of warming
763 and increased precipitation on community structure and composition in an annual forb
764 dominated desert steppe. *PLoS one*, 8, e70114.
765 <http://dx.doi.org/10.1371/journal.pone.0070114>, 2013.

766 Huxman, T. E., Snyder, K. A., Tissue, D., Leffler, A. J., Ogle, K., Pockman, W. T., Sandquist,
767 D. R., Potts, D. L., and Schwinning, S.: Precipitation pulses and carbon fluxes in semiarid
768 and arid ecosystems. *Oecologia*, 141, 254–268, [http://dx.doi.org/10.1007/s00442-004-](http://dx.doi.org/10.1007/s00442-004-1682-4)
769 [1682-4](http://dx.doi.org/10.1007/s00442-004-1682-4), 2004.

770 IPCC. 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II
771 and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change
772 [Core Writing Team, Pachauri RK, Meyer LA (eds.)]. IPCC, Geneva, Switzerland, pp151.

773 IPCC. 2019. Climate Change and Land: an IPCC special report on climate change,
774 desertification, land degradation, sustainable land management, food security, and
775 greenhouse gas fluxes in terrestrial ecosystems. [Armeth A, Barbosa H, Benton T et al. (eds)]
776 IPCC, <https://www.ipcc.ch/report/srcl/>.

777 Janssens, I. A., Kowalski, A. S., and Ceulemans, R.: Forest floor CO₂ fluxes estimated by eddy
778 covariance and chamber-based model. *Agric. For. Meteorol.* 106: 61–69,
779 [http://dx.doi.org/10.1016/S0168-1923\(00\)00177-5](http://dx.doi.org/10.1016/S0168-1923(00)00177-5), 2001.

780 Jassal, R. S., Black, T. A., Novak, M. D., Gaumont-Guay, D., and Nescic, Z.: Effect of soil water
781 stress on soil respiration and its temperature sensitivity in an 18-year-old temperate
782 Douglas-fir stand. *Glob. Change Biol.*, 14, 1–14, [http://dx.doi.org/10.1111/j.1365-](http://dx.doi.org/10.1111/j.1365-2486.2008.01573.x)
783 [2486.2008.01573.x](http://dx.doi.org/10.1111/j.1365-2486.2008.01573.x), 2008.

784 Jenkinson, D. S., Adams, D. E., and Wild, A.: Model estimates of CO₂ emissions from soil in
785 response to global warming. *Nature*, 351, 304–306, <http://dx.doi.org/10.1038/351304a0>,
786 1991.

787 Jia, B., Zhou, G., Wang, Y., Wang, F., and Wang, X.: Effects of temperature and soil water-
788 content on soil respiration of grazed and ungrazed *Leymus chinensis* steppes, Inner
789 Mongolia. *J. Arid Environ.*, 67, 60–76, <http://dx.doi.org/10.1016/j.jaridenv.2006.02.002>,

790 2006.

791 Jobbágy, E. G. and Jackson, R. B.: The vertical distribution of soil organic carbon and its
792 relation to climate and vegetation. *Ecol. Appl.*, 10, 423–436,
793 [https://doi.org/10.1890/1051-0761\(2000\)010\[0423:TVDOSO\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0423:TVDOSO]2.0.CO;2), 2000.

794 Johnson, M. G., Rygiewicz, P. T., Tingey, D. T., and Phillips, D. L.: Elevated CO₂ and elevated
795 temperature have no effect on Douglas-fir fine-root dynamics in nitrogen-poor soil. *New*
796 *Phytol.*, 170, 345–356, <http://dx.doi.org/10.1111/j.1469-8137.2006.01658.x>, 2006.

797 Kang, L., Han, X. G., Zhang, Z. B., and Sun, O. J.: Grassland ecosystems in China: review of
798 current knowledge and research advancement. *Philos. T. R. Soc. B*, 362, 997–1008,
799 <http://dx.doi.org/10.1098/rstb.2007.2029>, 2007.

800 Karhu, K., Auffret, M. D., Dungait, J. A., Hopkins, D. W., Prosser, J. I., Singh, B. K., Subke,
801 J. A., Wookey, P. A., Agren, G. I., Sebastia, M. T., Gouriveau, F., Bergkvist, G., Meir, P.,
802 Nottingham, A. T., Salinas, N., and Hartley, I. P.: Temperature sensitivity of soil
803 respiration rates enhanced by microbial community response. *Nature*, 513, 81–84,
804 <http://dx.doi.org/10.1038/nature13604>, 2014.

805 Keith, H., Jacobsen, K. L., and Raison, R. J.: Effects of soil phosphorus availability,
806 temperature and moisture on soil respiration in *Eucalyptus pauciflora* forest. *Plant*
807 *Soil*, 190, 127–141, <https://doi.org/10.1023/A:1004279300622>, 1997.

808 Kwon, M. J., Natali, S. M., Hicks, C. E., Schuur, E. A., Steinhof, A., Crummer, K. G., Zimov,
809 N., Zimov, S. A., Heimann, M., Kolle, O., and Göckede, M.: Drainage enhances modern
810 soil carbon contribution but reduces old soil carbon contribution to ecosystem respiration
811 in tundra ecosystems. *Glob. Change Biol.*, 25, 1315–1325,
812 <https://doi.org/10.1111/gcb.14578>, 2019.

813 Li, H. J., Yan, J. X., Yue, X. F., and Wang, M. B.: Significance of soil temperature and moisture
814 for soil respiration in a Chinese mountain area. *Agric. For. Meteorol.*, 148, 490–503,
815 <http://dx.doi.org/10.1016/j.agrformet.2007.10.009>, 2008.

816 Lin, X. W., Zhang, Z. H., Wang, S. P., Hu, Y. G., Xu, G. P., Luo, C. Y., Chang, X. F., Duan, J.
817 C., Lin, Q. Y., Xu, B., Wang, Y. F., Zhao, X. Q., and Xie, Z. B.: Response of ecosystem
818 respiration to warming and grazing during the growing seasons in the alpine meadow on
819 the Tibetan plateau. *Agric. For. Meteorol.*, 151, 792–802,
820 <http://dx.doi.org/10.1016/j.agrformet.2011.01.009>, 2011.

821 Litton, C. M. and Giardina, C. P.: Below-ground carbon flux and partitioning: global patterns
822 and response to temperature. *Funct. Ecol.*, 22, 941–954, <http://dx.doi.org/10.1111/j.1365->
823 [2435.2008.01479.x](http://dx.doi.org/10.1111/j.1365-2435.2008.01479.x), 2008.

824 Liu, L. T., Hu, C. S., Yang, P. P., Ju, Z. Q., Olesen, J. E., and Tang, J. W.: Effects of experimental
825 warming and nitrogen addition on soil respiration and CH₄ fluxes from crop rotations of
826 winter wheat–soybean/fallow. *Agric. For. Meteorol.*, 207, 38–47,

827 <https://doi.org/10.1016/j.agrformet.2015.03.013>, 2015.

828 Liu, L., Wang, X., Lajeunesse, M. J., Miao, G., Piao, S., Wan, S., Wu, Y., Wang, Z., Yang, S.,
829 Li, P. and Deng, M.: A cross-biome synthesis of soil respiration and its determinants under
830 simulated precipitation changes. *Glob. Change Biol.*, 22, 1394–1405,
831 <http://dx.doi.org/10.1111/gcb.13156>, 2016a.

832 Liu, T., Xu, Z. Z., Hou, Y. H., and Zhou, G. S.: Effects of warming and changing precipitation
833 rates on soil respiration over two years in a desert steppe of northern China. *Plant Soil*,
834 400, 15–27, <http://dx.doi.org/10.1007/s11104-015-2705-0>, 2016b.

835 Liu, W. X., Jiang, L., Hu, S. J., Li, L. H., Liu, L. L., and Wan, S. Q.: Decoupling of soil
836 microbes and plants with increasing anthropogenic nitrogen inputs in a temperate steppe.
837 *Soil Biol. Biochem.*, 72, 116–122, <http://dx.doi.org/10.1016/j.soilbio.2014.01.022>, 2014.

838 Liu, W. X., Zhang, Z., and Wan, S. Q.: Predominant role of water in regulating soil and
839 microbial respiration and their responses to climate change in a semiarid grassland. *Glob.*
840 *Change Biol.*, 15, 184–195, <http://dx.doi.org/10.1111/j.1365-2486.2008.01728.x>, 2009.

841 Liu, X. Z., Wan, S. Q., Su, B., Hui, D. F., and Luo, Y. Q.: Response of soil CO₂ efflux to water
842 manipulation in a tallgrass prairie ecosystem. *Plant Soil*, 240, 213–223,
843 <http://dx.doi.org/10.1023/a:1015744126533>, 2002.

844 Lloyd, J. and Taylor, J. A.: On the temperature dependence of soil respiration. *Funct. Ecol.*, 8,
845 315–323, <http://dx.doi.org/10.2307/2389824>, 1994.

846 Luo, Y. Q., Wan, S. Q., Hui, D. F., and Wallace, L. L.: Acclimatization of soil respiration to
847 warming in a tall grass prairie. *Nature*, 413, 622–625,
848 <http://dx.doi.org/10.1038/35098065>, 2001.

849 Ma, Q., Yu, H., Liu, X., Xu, Z., Zhou, G. and Shi, Y.: Climatic warming shifts the soil nematode
850 community in a desert steppe. *Climatic Change*, 150, 243–258,
851 <https://doi.org/10.1007/s10584-018-2277-0>, 2018.

852 Ma, Y. C., Piao, S. L., Sun, Z. Z., Lin, X., Wang, T., Yue, C., and Yang, Y.: Stand ages regulate
853 the response of soil respiration to temperature in a *Larix principis-rupprechtii* plantation.
854 *Agric. For. Meteorol.*, 184, 179–187, <http://dx.doi.org/10.1016/j.agrformet.2013.10.008>,
855 2014.

856 Maestre, F. T., Salguero-Gómez, R. and Quero, J. L.: It is getting hotter in here: determining
857 and projecting the impacts of global environmental change on drylands. *Philos. T. R. Soc.*
858 *B.*, 367, 3062–3075, <http://dx.doi.org/10.1098/rstb.2011.0323>, 2012.

859 Martins, C. S. C., Macdonald, C. A., Anderson, I. C., and Singh, B. K.: Feedback responses of
860 soil greenhouse gas emissions to climate change are modulated by soil characteristics in
861 dryland ecosystems. *Soil Biol. Biochem.*, 100, 21–32,
862 <http://dx.doi.org/10.1016/j.soilbio.2016.05.007>, 2016.

863 Meisner, A., Rousk, J., and Bååth E.: Prolonged drought changes the bacterial growth response
864 to rewetting. *Soil Biol. Biochem.* 88, 314–322,
865 <https://doi.org/10.1016/j.soilbio.2015.06.002>, 2015.

866 Meyer, N., Welp, G., and Amelung, W.: The temperature sensitivity (Q_{10}) of soil respiration:
867 controlling factors and spatial prediction at regional scale based on environmental soil
868 classes. *Glob. Biogeochem. Cycle*, 32, 306–323,
869 <http://dx.doi.org/10.1002/2017GB005644>, 2018.

870 Moncrieff, J. B., and Fang, C.: A model for soil CO₂ production and transport 2: application to
871 a Florida *Pinus elliotte* plantation. *Agric. For. Meteorol.*, 95, 237–256,
872 [https://doi.org/10.1016/S0168-1923\(99\)00035-0](https://doi.org/10.1016/S0168-1923(99)00035-0), 1999.

873 Moyano, F. E., Manzoni, S., and Chenu, C.: Responses of soil heterotrophic respiration to
874 moisture availability: an exploration of processes and models. *Soil Biol. Biochem.*, 59,
875 72–85, <http://dx.doi.org/10.1016/j.soilbio.2013.01.002>, 2013.

876 Moyano, F. E., Vasilyeva, N., Bouckaert, L., Cook, F., Craine, J., Yuste, J. C., Don, A., Epron,
877 D., Formanek, P., Franzluebbers, A., Ilstedt, U., Kätterer, T., Orchard, V., Reichstein, M.,
878 Rey, A., Ruamps, L., Subke, J. A., Thomsen, I. K., and Chenu, C.: The moisture response
879 of soil heterotrophic respiration: interaction with soil properties. *Biogeosciences*, 8,
880 1173–1182, <http://dx.doi.org/10.5194/bg-9-1173-2012>, 2012.

881 Munson, S. M., Benton, T. J., Lauenroth, W. K., and Burke, I. C.: Soil carbon flux following
882 pulse precipitation events in the shortgrass steppe. *Ecol. Res.*, 25, 205–211,
883 <https://doi.org/10.1007/s11284-009-0651-0>, 2009.

884 Nelson, D. W. and Sommers, L. E.: Dry combustion method using medium temperature
885 resistance furnace. In: Page AL, Miller RH, Keeney DR (eds). *Methods of Soil Analysis*,
886 Part 2. Chemical and Microbial Properties. Madison, WI: American Society of Agronomy
887 and Soil Science Society of America, 539–579, 1982.

888 Prietzel, J., Zimmermann, L., Schubert, A., and Christophel, D.: Organic matter losses in
889 German Alps forest soils since the 1970s most likely caused by warming. *Nat. Geosci.*, 9,
890 543–548, <http://dx.doi.org/10.1038/ngeo2732>, 2016.

891 Reynolds, L. L., Johnson, B. R., Pfeifer-Meister, L., and Bridgham, S. D.: Soil respiration
892 response to climate change in Pacific Northwest prairies is mediated by a regional
893 Mediterranean climate gradient. *Glob. Change Biol.*, 21, 487–500,
894 <http://dx.doi.org/10.1111/gcb.12732>, 2015.

895 Rinnan, R., Michelsen, A., Bååth, E., and Jonasson, S.: Fifteen years of climate change
896 manipulations alter soil microbial communities in a subarctic heath ecosystem. *Glob.*
897 *Change Biol.*, 13, 28–39, <http://dx.doi.org/10.1111/j.1365-2486.2006.01263.x>, 2007.

898 Rinnan, R., Stark, S., and Tolvanen, A.: Responses of vegetation and soil microbial
899 communities to warming and simulated herbivory in a subarctic heath. *J. Ecol.*, 97, 788–

800, <http://dx.doi.org/10.1111/j.1365-2745.2009.01506.x>, 2009.

901 Roby, M. C., Scott, R. L., Barron-Gafford, G. A., Hamerlynck, E. P., Moore, D. J.:
 902 Environmental and vegetative controls on soil CO₂ efflux in three semiarid ecosystems.
 903 *Soil Syst.*, 3, 6, <https://doi.org/10.3390/soilsystems3010006>, 2019.

904 Romero-Olivares, A. L., Allison, S. D., and Treseder, K. K.: Soil microbes and their response
 905 to experimental warming over time: A meta-analysis of field studies. *Soil Biol. Biochem.*,
 906 107, 32–40, <http://dx.doi.org/10.1016/j.soilbio.2016.12.026>, 2017.

907 Ru, J., Zhou, Y., Hui, D., Zheng, M., and Wan, S.: Shifts of growing-season precipitation peaks
 908 decrease soil respiration in a semiarid grassland. *Glob. Change Biol.*, 24, 1001–1011,
 909 <http://dx.doi.org/10.1111/gcb.13941>, 2018.

910 Sánchez-Rodríguez, A.R., Nie, C., Hill, P.W., Chadwick, D.R., Jones, D.L.: Extreme flood
 911 events at higher temperatures exacerbate the loss of soil functionality and trace gas
 912 emissions in grassland. *Soil Biol. Biochem.*, 130, 227–236,
 913 <https://doi.org/10.1016/j.soilbio.2018.12.021>, 2019.

914 Savage, K., Davidson, E. A., Richardson, A. D., and Hollinger, D. Y.: Three scales of temporal
 915 resolution from automated soil respiration measurements. *Agric. For. Meteorol.*, 149,
 916 2012–202, <http://dx.doi.org/10.1016/j.agrformet.2009.07.008>, 2009.

917 Schlesinger, W. H.: Carbon balance in terrestrial detritus. *Annu. Rev. Ecol. Evol. Syst.*, 8, 51–
 918 81, <http://dx.doi.org/10.1146/annurev.es.08.110177.000411>, 1977.

919 Shen, Z. X., Li, Y. L., and Fu, G.: Response of soil respiration to short-term experimental
 920 warming and precipitation pulses over the growing season in an alpine meadow on the
 921 Northern Tibet. *Appl. Soil Ecol.*, 90, 35–40,
 922 <http://dx.doi.org/10.1016/j.apsoil.2015.01.015>, 2015.

923 Sierra C. A., Trumbore S. E., Davidson E. A., Vicca S., and Janssens I.: Sensitivity of
 924 decomposition rates of soil organic matter with respect to simultaneous changes in
 925 temperature and moisture. *J. Adv. Model. Earth Syst.*, 7, 335–356,
 926 <http://dx.doi.org/10.1002/2014MS000358>, 2015.

927 Skopp, J., Jawson, M.D. and Doran, J.W.: Steady-state aerobic microbial activity as a function
 928 of soil water content. *Soil Sci. Soc. Am. J.* 54, 1619–1625,
 929 <https://doi.org/10.2136/sssaj1990.03615995005400060018x>, 1990.

930 Song, W. M., Chen, S. P., Wu, B., Zhu, Y. J., Zhou, Y. D., Lu, Q., and Lin, G. H.: Simulated
 931 rain addition modifies diurnal patterns and temperature sensitivities of autotrophic and
 932 heterotrophic soil respiration in an arid desert ecosystem. *Soil Biol. Biochem.*, 82, 143–
 933 152, <http://dx.doi.org/10.1016/j.soilbio.2014.12.020>, 2015.

934 Sponseller, R. A.: Precipitation pulses and soil CO₂ flux in a Sonoran Desert ecosystem. *Glob.*
 935 *Change Biol.*, 13, 426–436, <https://doi.org/10.1111/j.1365-2486.2006.01307.x>, 2007.

936 Thakur, M. P., Del Real, I. M., Cesarz, S., Steinauer, K., Reich, P. B., Hobbie, S., Ciobanu, M.,
 937 Rich, R., Worm, K., and Eisenhauer, N.: Soil microbial, nematode, and enzymatic
 938 responses to elevated CO₂, N fertilization, warming, and reduced precipitation. *Soil Biol.*

939 Biochem., 135, 184–193, <http://dx.doi.org/10.1016/j.soilbio.2019.04.020>, 2019.

940 Tucker, C. L. and Reed, S. C.: Low soil moisture during hot periods drives apparent negative
 941 temperature sensitivity of soil respiration in a dryland ecosystem: a multi-model
 942 comparison. *Biogeochemistry*, 128: 155–169, [http://dx.doi.org/10.1007/s10533-016-](http://dx.doi.org/10.1007/s10533-016-0200-1)
 943 [0200-1](http://dx.doi.org/10.1007/s10533-016-0200-1), 2016.

944 Tucker, C.L., Bell, J., Pendall, E., and Ogle K.: Does declining carbon-use efficiency explain
 945 thermal acclimation of soil respiration with warming? *Glob. Chang Biol.*, 19, 252–263,
 946 <https://doi.org/10.1111/gcb.12036>, 2013.

947 Wan, S. Q., Hui, D. F., Wallace, L., and Luo, Y. Q.: Direct and indirect effects of experimental
 948 warming on ecosystem carbon processes in a tallgrass prairie. *Glob. Biogeochem. Cycle*,
 949 19, 1–13, <http://dx.doi.org/10.1029/2004GB002315>, 2005.

950 Wan, S. Q., Norby, R. J., Ledford, J., and Weltzin, J. F.: Responses of soil respiration to
 951 elevated CO₂, air warming, and changing soil water availability in a model old-field
 952 grassland. *Glob. Change Biol.*, 13, 2411–2424, [http://dx.doi.org/10.1111/j.1365-](http://dx.doi.org/10.1111/j.1365-2486.2007.01433.x)
 953 [2486.2007.01433.x](http://dx.doi.org/10.1111/j.1365-2486.2007.01433.x), 2007.

954 Wan, S. Q., Norby, R. J., Pregitzer, K. S., Ledford, J., and O'Neill, E. G.: CO₂ enrichment and
 955 warming of the atmosphere enhance both productivity and mortality of maple tree fine
 956 roots. *New Phytol.*, 162, 437–446, <http://dx.doi.org/10.1111/j.1469-8137.2004.01034.x>,
 957 2004.

958 Wang, Y., Hao, Y., Cui, X. Y., Zhao, H., Xu, C., Zhou, X., and Xu, Z.: Responses of soil
 959 respiration and its components to drought stress. *J. Soils Sedim.*, 14, 99–109,
 960 <http://dx.doi.org/10.1007/s11368-013-0799-7>, 2014.

961 Wu, H. J., and Lee, X.: 2011. Short-term effects of rain on soil respiration in two New England
 962 forests. *Plant Soil* 338, 329–342, <https://doi.org/10.1007/s11104-010-0548-2>, 2011.

963 Xu, W., Yuan, W., Cui, L., Ma, M., and Zhang, F.: Responses of soil organic carbon
 964 decomposition to warming depend on the natural warming gradient. *Geoderma*, 343, 10–
 965 18, <https://doi.org/10.1016/j.geoderma.2019.02.017>, 2019.

966 Xu, Z., Hou, Y., Zhang, L., Tao, L., and Zhou, G.: Ecosystem responses to warming and
 967 watering in typical and desert steppes. *Sci., Rep.*, 6, 34801,
 968 <http://dx.doi.org/10.1038/srep34801>, 2016.

969 Xu, Z., Shimizu, H., Ito, S., Yagasaki, Y., Zou, C., Zhou, G. and Zheng, Y.: Effects of elevated
 970 CO₂, warming and precipitation change on plant growth, photosynthesis and peroxidation
 971 in dominant species from North China grassland. *Planta*, 239, 421–435,
 972 <https://doi.org/10.1007/s00425-013-1987-9>, 2014.

973 Yan, M. F., Zhou, G. S., and Zhang, X. S.: Effects of irrigation on the soil CO₂ efflux from
 974 different poplar clone plantations in arid northwest China. *Plant Soil*, 375, 89–97,
 975 <http://dx.doi.org/10.1007/s11104-013-1944-1>, 2013.

976 Yan, Z. B., Bond-Lamberty, K. E., Todd-Brown, V. L., Bailey, S., Li, C., Liu, C. Q., Liu C.: A
 977 moisture function of soil heterotrophic respiration that incorporates microscale processes,
 978 *Nat. Commun.*, 9, 2562, <http://doi:10.1038/s41467-018-04971-6>, 2018.

979 Yin, X., Goudriaan, J. A. N., Lantinga, E. A., Vos, J. A. N., and Spiertz, H. J.: A flexible

980 Gompertz function of determinate growth. *Ann. Bot.*, 91, 361–371,
981 <http://aob.oupjournals.org/cgi/doi/10.1093/aob/mcg029>, 2003.

982 Yu, H. Y., Chen, Y. T., Xu, Z. Z., and Zhou, G. S.: Analysis of relationships among leaf
983 functional traits and economics spectrum of plant species in the desert steppe of Nei
984 Mongol. *Chin. J. Plant Ecol.*, 38, 1029–1040, doi: 10.3724/SP.J.1258.2014.00097, 2014.

985 Yu, H. Y., Xu, Z. Z., Zhou, G. S., and Shi, Y. H.: The data for the article entitled "Soil carbon
986 release responses to long-term versus short-term climatic warming in an arid ecosystem"
987 [Data set]. Zenodo, <https://doi.org/10.5281/zenodo.3546062>, 2019.

988 Yu, H., Ma, Q., Liu, X., Xu, Z., Zhou, G. and Shi, Y.: Short-and long-term warming alters soil
989 microbial community and relates to soil traits. *Appl. Soil Ecol.*, 131, 22–28,
990 <https://doi.org/10.1016/j.apsoil.2018.07.006>, 2018.

991 Zhang, C. P., Niu, D. C., Hall, S. J., Wen, H. Y., Li, X. D., Fu, H., Wan, C. G., and Elser, J. J.:
992 Effects of simulated nitrogen deposition on soil respiration components and their
993 temperature sensitivities in a semiarid grassland. *Soil Biol. Biochem.*, 75, 113–123,
994 <http://dx.doi.org/10.1016/j.soilbio.2014.04.013>, 2014.

995 Zhang, N. L., Wan, S. Q., Li, L. H., Bi, J., Zhao, M. M., and Ma, K. P.: Impacts of urea N
996 addition on soil microbial community in a semi-arid temperate steppe in northern China.
997 *Plant Soil*, 311, 19–28, <http://dx.doi.org/10.1007/s11104-008-9650-0>, 2008.

998 Zhang, W., Parker, K. M., Luo, Y., Wan, S., Wallace, L. L., and Hu, S.: Soil microbial responses
999 to experimental warming and clipping in a tallgrass prairie. *Glob. Change Biol.*, 11, 266–
1000 277, <http://dx.doi.org/10.1111/j.1365-2486.2005.00902.x>, 2005.

1001 **Figure legends**

1002 Figure 1. Effects of warming on the soil temperature and soil moisture during the
1003 growth peak in 2014 (Mean \pm SE). Mean daily values were presented (n = 120).
1004 The mean values with the same lowercase letters on the SE bars are not different
1005 at $P < 0.05$ according to LSD multiple range tests (P values and F ratios are shown
1006 inside).

1007 Figure 2. Relationship between R_s and soil water content based on a linear (blue
1008 line) and a quadratic (black line) functional model (A), and Gompertz functional
1009 model (B). Close and open circles denote the data in 2014 and 2017, respectively.
1010 The close red circles indicate data used for the linear R_s response to SWC at low
1011 levels. The one open triangle may be an outlier point due to some errors, but it does
1012 not notably affect the functional fitting when removing it (ref. Figure S2). Based
1013 on Gompertz functional curve, the R_s asymptote value, as an estimated maximum,
1014 is $3.76 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ when the optimal SWC is 22.85% [The red line denotes the
1015 initial R_s response to SWC; the blue line denotes $R_s = \text{constant}$ value of the
1016 maximum estimated by the asymptote value; and the intersection of the two lines
1017 represents a point (the blue arrow) at which R_s levelled off]. Note, we measured
1018 the R_s during 9:00-10:00 in these cloudless days with calm/gentle wind in order to
1019 maintain other environmental factors such as soil temperature and radiation to
1020 relatively stable and constant. The data were collected in the plots of watering
1021 treatments (n = 92)..

1022 Figure 3. The relationships between soil respiration and soil temperature under
1023 both watering (n = 23-25, A), and warming treatments (n=28-33, B) (Mean \pm SE).

1024 Figure 4. Effects of warming regimes on average soil respiration in 2014 (mean \pm
1025 SE), the mean values with the same lowercase letters on the SE bars are not
1026 different at $P < 0.05$ according to LSD multiple range tests (P values and F ratios
1027 are shown inside).


1028 Figure 5. An interactive relationship of soil respiration with both soil temperature
1029 (T_s) and soil water content (SWC) based on a nonlinear mixed model ($R_s =$
1030 $(0.733*1.796^{(T_s-10)/10})*\beta^{(0.229-SWC)^2}$). The data were used in control plots in the
1031 warming experiment. The optimal SWC of 0.229 was estimated by the Gompertz
1032 functional curve (see Figure 2B).

1033 Figure 6. A diagram of the effects of key environmental factors on soil respiration
1034 and their relationships. Blue double-headed arrows represent the relationships
1035 between the key environmental factors, data on the arrows are correlation
1036 coefficients. Black arrows represent the relationships between soil respiration and
1037 the key environmental factors, data on the arrows are correlation coefficients (bold)
1038 and direct path coefficients (italic), respectively. *, $P < 0.05$; **, $P < 0.01$, n = 12.
1039 For other details, see Supplementary Table S2.


1040

1041 Supplementary Figure S1. Long-term air temperature (A) and total annual


1042 precipitation (B) records from 1955 to 2014 in the experiment site in the desert
1043 steppe ecosystem, Damao Banner, Nei Mongol, China.
1044 Supplementary Figure S2. Relationship between R_s and soil water content based
1045 on a linear (black line) and a quadratic (dotted line) functional model (A), and
1046 Gompertz functional model (B). Close and open circles denote the data in 2014
1047 and 2017, respectively. The close red circles indicate data used for the initial R_s
1048 response to SWC. The functional fitting does not substantially affect despite a
1049 slight improvement with greater R^2 values when the outlier point was removed (ref.
1050 Figure 2). Note, we measured the R_s during 9:00-10:00 in the cloudless days with
1051 calm/gentle wind in order to maintain other environmental factors such as soil
1052 temperature and radiation to relatively stable and constant ($n = 91$).




1053 **Figure 1.** Effects of warming on the soil temperature and soil moisture during the growth
 1054 peak in 2014 (Mean \pm SE). Mean daily values were presented (n = 120). The mean values with
 1055 the same lowercase letters on the SE bars are not different at $P < 0.05$ according to LSD
 1056 multiple range tests (P values and F ratios are shown inside).




1057 **Figure 2.** Relationship between R_s and soil water content based on a linear (blue line) and a
 1058 quadratic (black line) functional model (A), and Gompertz functional model (B). Close and
 1059 open circles denote the data in 2014 and 2017, respectively. The close red circles indicate data
 1060 used for the linear R_s response to SWC at low levels. The one open triangle may be an outlier
 1061 point due to some errors, but it does not notably affect the functional fitting when removing it
 1062 (ref. Figure S2). Based on Gompertz functional curve, the R_s asymptote value, as an estimated
 1063 maximum, is $3.76 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ when the optimal SWC is 22.85% [The red line denotes the
 1064 initial R_s response to SWC; the blue line denotes $R_s =$ constant value of the maximum estimated
 1065 by the asymptote value; and the intersection of the two lines represents a point (the blue arrow)
 1066 at which R_s levelled off]. Note, we measured the R_s during 9:00-10:00 in these cloudless days
 1067 with calm/gentle wind in order to maintain other environmental factors such as soil
 1068 temperature and radiation to relatively stable and constant. The data were collected in the plots
 1069 of watering treatments ($n = 92$).




1070 **Figure 3.** The relationships between soil respiration and soil temperature under both watering
 1071 (n = 23-25, A), and warming treatments (n=28-33, B) (Mean ± SE).
 1072



1073 **Figure 4.** Effects of warming regimes on average soil respiration in 2014 (mean \pm SE), the
 1074 mean values with the same lowercase letters on the SE bars are not different at $P < 0.05$
 1075 according to LSD multiple range tests (P values and F ratios are shown inside).



1076 **Figure 5.** An interactive relationship of soil respiration with both soil temperature (Ts) and soil
 1077 water content (SWC) based on a nonlinear mixed model ($R_s = (0.733 * 1.796^{(Ts-10)/10}) * \beta^{(0.229-SWC)^2}$,
 1078 B). The data were used in control plots in the warming experiment. The optimal SWC of 0.229
 1079 was estimated by the Gompertz functional curve (see Figure 2B).
 1080



1081 **Figure 6.** A diagram of the effects of key environmental factors on soil respiration and their
 1082 relationships. Blue double-headed arrows represent the relationships between the key
 1083 environmental factors, data on the arrows are correlation coefficients. Black arrows represent
 1084 the relationships between soil respiration and the key environmental factors, data on the arrows
 1085 are correlation coefficients (**bold**) and direct path coefficients (*italic*), respectively. *, $P < 0.05$;
 1086 **, $P < 0.01$, $n = 12$. For other details, see Supplementary Table S2.
 1087